Title: Experimental evolution of gene essentiality in bacteria

Authors: Liang Bao[1, *], Zan Zhu[1], Ahmed Ismail[1], Bin Zhu[2], Vysakh Anandan[1], Marvin Whiteley[3], Todd Kitten[1], Ping Xu[1, *]

Affiliations:
1, Department of Oral and Craniofacial Molecular Biology, Philips Institute for Oral Health Research, School of Dentistry, Virginia Commonwealth University, Virginia, USA.
2, Massey Cancer Center, Virginia Commonwealth University, Virginia, USA.
3, School of Biological Sciences, Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Georgia, USA.

Correspondence:

pxu@vcu.edu (P.X.), baol@vcu.edu (L.B.)

Summary

Essential gene products carry out fundamental cellular activities in interaction with other components. However, the lack of essential gene mutants and appropriate methodologies to link essential gene functions with their partners poses significant challenges. Here, we have generated deletion mutants in 32 genes previously identified as essential, with 23 mutants showing extremely slow growth in the SK36 strain of *Streptococcus sanguinis*. Whole-genome sequencing of 243 independently evolved populations of these mutants has identified >1000 spontaneous suppressor mutations in experimental evolution, many of which are new gene and pathway relationships, such as F1Fo-ATPase/V1V1-ATPase/TrkA1-H1 gene pathway. Patterns of spontaneous mutations occurring in essential gene mutants differed from those found in wildtype. While gene duplications occurred rarely, substitutions, deletions, and insertions were prevalent in evolved populations. These essential gene deletion mutants and spontaneous mutations fixed in the mutant populations during evolution established a foundation for understanding gene essentiality and the interaction of essential genes in networks.

Keywords

Gene essentiality; Experimental evolution; Spontaneous mutations; Whole-genome sequencing; *Streptococcus*; F1Fo-ATPase/V1V1-ATPase/TrkA1-H1 gene pathway.

Introduction

All living organisms require essential physical properties and biochemical capacities encoded by "essential" genes to sustain basic cellular activities. Essential genes are indispensable for an organism to survive or reproduce offspring, and their essentiality is highly dependent on the

environment in which the organism lives. [1,2] *In vivo*, essential gene products carry out these tasks
by interacting with other components. Therefore, the essentiality of genes can also depend on the
genetic context provided by the additional genes present. [2-4] Relationships may be characterized
by negative epistasis when the essentiality of a gene is dependent on the absence or impairment
of another gene's function, such as synthetic lethality [3,5] or positive epistasis interactions, where
an individual with two genes mutated is fitter than a strain possessing a mutant version of only
one of the two genes. [3,6,7] In contrast to binary classification, it is increasingly acknowledged that
essential genes exhibit a quantitative spectrum, displaying a gradient of essentiality [2] and a
conditional nature. [8] For some essential genes, under optimal controlled conditions, deletion can
still result in viable, albeit severely compromised individuals. [9,10] The loss of essential genes
produces intense selective pressure for the fixation of genomic suppressor mutations in the
descendants of the original mutants during subsequent generations, including aneuploidy of
chromosomes in the yeast *Saccharomyces cerevisiae*. [9,10] Furthermore, essential genes can lose
their essentiality during long-term evolution and acquire novel functions. [11-13] For instance,
unlike the essential characteristics of the critical cell cycle regulator *CDK1* in yeasts and animals,
[14] plant PSTAIRE-type cyclin-dependent kinases, known as *CDKAs* (plant homologs of *CDK1*),
play a role in environmental responses that is independent of the cell cycle, [11] likely after the
acquisition of the plant-specific cyclin-dependent kinases, *CDKBs*. [15]

Identifying genes that interact with essential genes can produce many potential benefits, both
from a theoretical perspective, such as understanding the basic principle of cellular life, [16,17] and
from a practical standpoint, such as addressing various challenges in synthetic biology [18] and
combating drug resistance of infectious diseases. [17] Experimental evolution or laboratory
evolution imposes a natural selective pressure on a population to redirect the evolution towards
fitness-improving phenotypes in a controlled laboratory setting. [9,10,19-25] This methodology has
been applied to identify spontaneous mutations associated with fitness-improving phenotypes,
such as multicellularity, [19] novel metabolic capacity, [20] restoration mobility of flagellar
mobility, [26] and fitness of minimal cells. [21] Recently, experimental evolution using mutants
deleted of essential genes in the yeast *Saccharomyces cerevisiae* revealed a rapid growth
phenotype after passages. [9,10] However, aneuploidy of chromosomes was prevalent among
evolved populations deleted of essential genes, which hindered the identification of causal
suppressors. [9,10] Here, we have established a transformation system to obtain numerous essential
gene knockout mutants, including *obgE*, *pfkA*, and subunit members of F1Fo-ATPase that have
been so far identified as essential in most bacterial species, including all species of Streptococcus
examined (http://www.essentialgene.org/), and performed experimental evolution of mutants
deleted for essential genes using a *Streptococcus* bacterium as our model. After allowing for
short-term adaptation, we have identified >1000 spontaneous suppressor mutations in 243
evolved populations deleted of essential genes, most of which are substitutions, deletions, and
insertions. This feature enabled us to map virtually all mutations to distinct genomic segments of
individual ORFs or intergenic regions. For comparison, we also evaluated the twelve mutations
that arose in six evolved populations in the wild-type (WT) cells. The mutations that occurred in
essential gene mutants displayed several characteristics that distinguished them from those that
evolved in WT cells during short-term evolution.

Results
85

Isolating essential gene mutants and analysis of short-term evolution

The genome of *Streptococcus sanguinis* SK36 comprises over 2000 ORFs, [27] of which 218 have been experimentally validated as essential for the organism's survival when cells are grown on a rich medium—brain heart infusion (BHI) broth or plates—under microaerobic conditions. [1] To isolate viable essential-gene deletion mutants, we have modified our transformation procedure by minimizing the stress during the procedures (see Materials and Methods). Although we still employed homologous recombination (Figure 1A and Fig S1A), we replaced our conventional 1-hour aerobic transformation with an anaerobic incubation and a duration of up to 24 hours. [1] Additionally, we optimized the selection conditions by covering the transformants with a thin layer of agar medium and extending the selection period for the transformants to four to six days, all conducted under anaerobic conditions. (See Materials and Methods.)

Using this procedure, we observed an interesting phenomenon. Specifically, for 23 of the essential genes, the transformants produced two types of colonies: large and small (Figure 1B, Fig S1B, and Fig S2A). As expected, genotyping of the small colonies revealed a complete replacement of the corresponding ORF with a kanamycin resistance (*kan*) gene (Fig S1C). In contrast, genotyping of the large colonies showed that the mutants displayed a "double band" genotype due to gene duplication of the target gene prior to the replacement of one copy by the *kan* gene (Fig S1D, Supplemental Table 1). This finding suggests that *S. sanguinis* SK36 regularly undergoes gene duplications, particularly between two copies of a multi-copy gene or locus, such as the rRNA operons, or other repeated elements. Alternatively, some large colonies lacked a *kan* gene, but contained point mutations in different genes, with the majority of isolates acquiring mutations in translational machinery genes that may confer resistance to kanamycin, [28] such as *fusA, rpsL* or *rplF* (Supplemental Table 1). Among fifty-four large colonies genotyped, fifteen showed the double-band genotype and no mutations in *fusA, rpsL*, or *rplF*. The remaining thirty-nine did not contain the *kan* gene, but thirty-eight, or 97.4% contained mutations in *fusA, rpsL* or *rplF* (Supplemental Table 1). This observation confirmed that the deletion of an essential gene significantly reduced the growth of the corresponding mutant. Therefore, these observations indicate that by selecting the survival, small colonies in the selection medium can obtain the essential gene deletion mutants. In contrast, for nine essential genes, the deletion mutants exhibited robust growth (Fig S2B). These nine genes included *nrdE, nrdF, nrdH,* and *nrdI* which were previously reported as non-essential under anaerobic conditions but essential in the presence of O_2, [29] along with *rexB* for ATP-dependent nuclease subunit B, *rpsA* for 30S ribosomal protein S1, *pdf* for peptide deformylase, and *g6pD* for glucose-6-phosphate dehydrogenase (Fig S2B).

We, therefore, directed our attention to the twenty-three essential genes that, when deleted, led to severe growth defects in the resultant mutants. These twenty-three genes encoding components of diverse pathways, include the eight subunit genes encoding the F1Fo-ATPase (*uncE*, *uncB*, *uncF*, *uncH*, *uncA*, *uncG*, *uncD*, and *uncC*); *ftsX,* encoding a permease-like cell division protein; five *fab* genes (*fabH*, *fabK*, *fabG*, *fabF*, and *fabZ*) that are involved in Type-II fatty acid synthesis (FAS II); two ribosomal protein genes (*rplV* and *rpmH*); two GTPase genes (*obgE* and *yihA*); *yqeG,* encoding a YqeG-family phosphatase; *pfkA*, encoding 6-phosphofructokinase, involved in glycolysis; *pngM*, encoding phosphoglucomutase; *dpcK*, encoding diphospho-CoA kinase, involved in CoA synthesis; and *dnaK*, encoding a heat shock protein chaperone. These

genes are widely distributed and are essential in various genera and species
(http://www.essentialgene.org/). [30] In total, we have created 23 essential gene mutants, each with
an individual gene deletion, along with one mutant deleted for the entire *f1fo* region encoding the
eight F1Fo-ATPase subunits, which we have named Δ*f1fo* (Fig S1 and S2A.1-A.2). Although the
deletion mutants of the 23 genes were initially generated under anaerobic condition in this study,
we have confirmed the viability of mutants under both anaerobic condition (Fig S3A) and
microaerobic condition (Fig S3B) as utilized for identifying the essential genome for *S. sanguinis*
SK36 in the previous report. [1]
140
In order to study the evolution of gene essentiality, we have conducted passage experiments
using multiple independently evolved populations of the 24 essential gene mutants. To form a
population, we selected three to five colonies from the selection medium, combined them into a
single inoculum, and subjected them to passage in BHI by 1:20 serial transfer. To ensure a
comparable number of generations, we closely monitored the growth of mutants during each
passage. We proceeded to the following passage only when the cell cultures had reached an
OD_{600} within the range of 0.1-0.5. As a control, WT cells, which reached an OD_{600} of
approximately 1.0 within 24 hours of growth, or 2-10 times more cells compared to the mutants
in each passage, were passaged daily for a total of nine times. This method was applied to
maintain consistency throughout the experimental process (Figure 1C, Fig S3C, and
Supplemental Table 2). While all mutants exhibited significant growth defects compared to WT,
we observed variations in their growth phenotypes. Notably, mutants lacking *pfkA, fabK, fabG,*
fabF, and *fabZ* displayed the most severe growth defects. In contrast, the growth of others,
including another *fab* mutant, *fabH,* was less severe, as evidenced by the OD_{600} measurements of
the evolved mutants (Figure 1C and Fig S3C, Supplemental Table 2).
156
In our passage experiments, we observed that the mutants in passage 6 (P6, "P" for passage)
exhibited enhanced fitness compared to those in early passages. This improvement revealed itself
in multiple ways. First, the time required for cultures to reach an OD_{600} of 0.1-0.5 in P0
compared to P6 (Figure 1C, Supplemental Table 3) indicated faster growth at P6, reflecting
improved fitness. Second, growth curve experiments using colonies of transformants (Figure 1D,
Supplemental Table 4) and evolved P6 populations (Figure 1E, Supplemental Table 4)
demonstrated faster growth rates in P6 compared to P1 using Δ*f1fo*.
164
165 **Identification of suppressor loci in evolved populations deleted of essential genes and in**
166 **WT**
167
To identify suppressor mutations in the evolved populations deleted of essential genes, we
performed whole-genome sequencing at P6. Specifically, we sequenced between two and
twenty-five independently evolved populations for each of the twenty-four essential gene
mutants, resulting in a total of 243 populations (Figure 2A and Supplemental Table 5). As
controls, we also sequenced six populations of WT, which were generated by aliquoting into six
separate tubes and passaged nine times, as well as the original WT (Figure 2A and Supplemental
Table 5).
175
We first considered mutations related to duplications of genes other than the essential gene
targeted for mutagenesis. Of the 243 evolved mutant populations, we found that twelve (4.9%)

displayed gene duplications of various sizes, ranging from 10 kb to 310 kb. By comparison, none
of the six evolved WT populations in P9 nor the original parental WT contained a gene
duplication (Fig S4A). This result suggests that increasing the copy number of a specific gene(s)
within these duplicated regions plays a crucial role in improving the fitness of mutants lacking
these particular essential genes. For the twelve evolved mutant populations with gene
duplications, two were from $\Delta f1fo$ (#107 and #109) (Figure 2B and Fig S4B), one from $\Delta uncE$
(#2) (Fig S4C), one from $\Delta uncH$ (#7) (Figure 2C), three from $\Delta fabH$ (#4, #6 and #14) (Figure 2D
and Fig S4D and S4E), one from $\Delta fabF$ (#4) (Figure 2E), two from $\Delta fabZ$ (#1 and #3) (Figure 2F
and 2G), one from $\Delta obgE$ (#5) (Figure 2H) and one from $\Delta dpcK$ (#17) (Figure 2I). The
duplicated regions found in populations of $\Delta f1fo$, $\Delta uncE$, $\Delta uncH$, $\Delta fabH$ and $\Delta fabF$ mutants
encompassed the same ~103 kb region (coordinate: 22,738-125,834) that is flanked by two
directly repeated rRNA operons (Figure 2B-E). The above-duplicated region of ~103 kb contains
113 ORFs, including nine ORFs encoding the entire V1Vo-ATPase operon that is related to the
proton pump function of F1Fo-ATPase, [31] one ORF encoding acyl carrier protein and one ORF
encoding acyltransferase that is related to lipid biosynthesis. [32] The duplicated region found in
$\Delta fabZ$#1 spans approximately 143 kb (coordinate: 1,053,488-1,197,391). The 10.1-kb
duplication in $\Delta fabZ$#3 (coordinate: 1,093,824-1,103,958) is within the duplicated 143 kb in
$\Delta fabZ$#1. This 10.1-kb region contains ten ORFs, including one ORF annotated as acyl-acyl
carrier protein thioesterase, which catalyzes the terminal reaction of fatty acid biosynthesis (Jing,
et al., 2011). In the case of $\Delta obgE$#5, a 37-kb duplication is present (coordinate: 1,775,394-
1,812,805), encompassing thirty ORFs, including one ORF encoding der GTPase. In $\Delta dpcK$#17,
a ~310-kb duplication occurred (coordinate: 385,678-697,581), encompassing 305 ORFs. The
flanking sequences of the duplicated regions in $\Delta fabZ$#1, $\Delta fabZ$#3, $\Delta obgE$#5 and $\Delta dpcK$#17
possessed no notable repeated sequences.
202
Next, we assessed substitutions, deletions, and insertions of less than 10 kb in the evolved
populations. The genome of *S. sanguinis* can be divided into 4,283 segments—2,340 ORFs and
1,943 intergenic regions (Supplemental Table 6). [33] We defined a segment as mutated if any
mutation within that segment was present in at least 30% of the sequence reads (see Materials
and Methods section). We selected 30% as the threshold based on our finding that confirmed
suppressor mutations, such as *trkA1-H1*, ranged from 34.8% to 100% in the evolved $\Delta f1fo$
populations in P6 (Figure 5D and Supplemental Table 5). There were six mutations (two
substitution mutations and four deletion mutations from the six evolved populations), where two
to four ORFs were affected by each mutation. For example, the C to T (at coordinate 1,313,563)
mutation in $\Delta fabF$#10 caused amino substitutions of two overlapped ORFs, encoding two
hypothetical proteins (Supplemental Table 5). The remaining mutations fell within single
segments of ORFs or intergenic regions. In total, we detected 1,272 mutated segments
(Supplemental Table 5), with 1,260 from evolved populations lacking essential genes and twelve
from the evolved WT populations (Supplemental Table 5). Our analysis revealed that each of the
243 (100%) evolved mutant populations contained at least one mutated segment, with the
majority containing from 2 to 10 (Figure 2J and Supplemental Table 5-7). In contrast, among the
six evolved WT populations, four had at least one segment containing a mutation in at least 30%
of the sequence reads, while the remaining two, as well as the parental WT population, had none
(Figure 2J and Supplemental Table 5). Furthermore, when we set the threshold to 50%, 241
evolved mutant populations, or 99.2%, had mutated segments, whereas none of the six evolved

WT populations contained mutated segments (Figure 2K and Supplemental Table 8) ($\chi 2$=154.69, df=1, n=249 populations, p=2.2e-16).

We then analyzed the presence of mutated segments belonging to essential ORFs (in each case, excluding the ORF that was intentionally deleted in each mutant, as well as the nine conditionally essential genes). We found that suppressor mutations belonging to essential segments were present in at least one evolved population from each essential gene mutant (Figure 2L and Supplemental Table 9-10). In contrast, none of the twelve mutated segments from the evolved WT populations were found to be within an essential ORF (Figure 2L and Supplemental Table 9-10) ($\chi 2$=5.74, df=1, n=25 groups, p=0.017). To illustrate this further, we compared the frequency of mutated segments in P1 and P6 for two specific cases: mutations of *ftsZ* (an essential segment encoding a structural homolog of tubulin in prokaryotes) in the evolved *Δf1fo#107* and *uncH* (an essential segment, encoding delta subunit of F1Fo-ATPase) in the evolved *ΔuncC#1*. The *uncH* gene is located 3.9 kb upstream of *uncC*, both of which are within the *f1fo* operon. The evolved *ΔuncC#1* population of P1 did not have any *uncH* mutations using 30% as the cutoff, although an *uncH* mutation was present in 12% of the sequence reads (Fig S5A-C). In contrast, the *ΔuncC#1* population of P6 contained 86% *uncH* mutation (Fig S5D-F) The resultant mutation replaced a 207-bp segment that was 51 bp downstream of the *uncH* start codon with an 82-bp fragment, causing early truncation (Fig S5G). For the second case, the *ftsZ* gene is located approximately 126 kb upstream of the *f1fo* operon. In the evolved *Δf1fo#107* population in P1, we did not detect any mutations in the *ftsZ* segment using 30% abundance as cutoff, although 27% of the sequence reads contained a *ftsZ* mutation (Fig S5G-H). In comparison, the evolved population of P6 contained 100% mutated *ftsZ*—a 248 bp insertion located 1015 bp downstream of the start codon (Fig S5I-J), leading to the truncation of the FtsZ protein (Fig S5K). These findings demonstrate an increment of mutated essential segments in the essential gene deletion mutant due to selection during passage.

Multi-drug resistance across ESKAPE pathogens (i.e., *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter spp.*) has been found to be commonly associated with mutations in six genes— namely, *rrs, gyrA, gyrB, rpoB, rpsL* and *rsmG* (also known as *gidB*).[34] We therefore asked whether mutations in the typical set of multi-drug resistance-associated genes[34] could appear in the evolved populations deleted of essential genes. Apart from the 16S rRNA gene, which is present in four copies (*rrs1* to *rrs4*), *S. sanguinis* has one copy of each of these genes. Among the 243 evolved populations deleted of essential genes, mutations of multi-drug resistance genes were present in *rrs2*, *rpoB*, *gyrB,* and *rspL* in a variety of populations derived from different essential gene mutants, while no mutations were detected in *rrs1*, *rrs3*, *rrs4*, *gyrA* or *rsmG*. Moreover, no mutation was detected within this set of multi-drug resistance-associated genes in the evolved WT populations (Figure 2M and Supplemental Table 11). These results showed that missense mutations in four of the same genes that promote multi-drug resistance also improve the fitness of mutants deleted of these particular essential genes.

In summary, compared to the evolved WT populations, mutants deleted for essential genes exclusively evolve gene duplications of greater than 10 kb, mutations in essential genes, and mutations of genes associated with multi-drug resistance. In addition, once mutations are observed, they are more quickly fixed in the population (Figure 2K).

269
270 **Construction of a network using WT and essential gene deletion mutants and their**
271 **suppressors**
272
To explore the relationships of suppressor mutations from the evolved populations, we
established a network of 25 groups, comprising the 24 mutants deleted for essential genes, along
with SK36, and their respective evolved suppressors across 249 populations (243 populations
from mutants and six populations from SK36). In these 249 populations, 1,272 mutations
occurred in 592 distinct segments, with 503 segments belonging to ORFs and 89 belonging to
intergenic regions (Figure 3, Supplemental Table 5, and Supplemental Table 12-14). This
network also encompasses a total of 914 edges connecting these 617 nodes (Figure 3 and
Supplemental Table 12-14).
281
Some mutations discovered in the evolved WT also appeared in evolved mutant populations. The
twelve mutations occurred in ten distinctive segments, with 5 segments specific to WT and 5
shared with populations derived from essential gene mutants. Notably, an ORF J1C87_09360,
annotated as PTS mannose/fructose/sorbose transporter family subunit IID, exhibited mutations
in two of the six evolved WT populations. This result is consistent with a previous study in SK36,
where fitness improvement was associated with mutations in three ORFs of the same operon
(J1C87_09355, J1C87_09360, and J1C87_09365). [35] In the current study, mutations in the three
OFRs were also observed in mutant populations of Δ*uncH*, Δ*fabG*, Δ*fabK*, Δ*fabF*, Δ*fabZ, and*
Δ*pfkA* (Fig S6A-B and Supplemental Table 15). Interestingly, most of the evolved populations
that contain suppressor mutations in the three PTS mannose/fructose/sorbose transporter subunits
are also those that showed the most severe growth defects, such as Δ*fabG*, Δ*fabK*, Δ*fabF*, Δ*fabZ*,
and Δ*pfkA* (Figure 1D and Fig S3C). These results demonstrate that mutations in some segments
that apparently improved fitness in WT could also contribute to the fitness improvement of
essential gene mutants.
296
Mutations in some segments that were present in evolved populations deleted of one essential
gene also appeared in evolved populations of other mutants deleted of another essential gene. For
example, most populations of the mutant deleted for the entire *f1fo* region (Δ*f1fo*) and the eight
F1Fo subunits deletion mutants (Δ*uncE*, Δ*uncB*, Δ*uncF*, Δ*uncH*, Δ*uncA*, Δ*uncG*, Δ*uncD* and
Δ*uncC)* contain suppressor mutations in J1C87_08610 (*trkA1*) and J1C87_08615 (*trkH1*) (Figure
3). Mutations in many F1Fo subunits are shared among the evolved populations of various other
subunit deletion mutants (Figure 3 and Supplemental Table 5). Furthermore, many of the
suppressor mutations in type II fatty acid synthesis (FAS II) gene mutants, such as Δ*fabZ*, Δ*fabF*,
Δ*fabG*, Δ*fabK* and Δ*fabH* and those in the CoA biosynthesis mutant Δ*dpcK*, occur in the same
ORFs related to fatty acid biosynthesis, including essential genes like acetyl-CoA carboxylase
carboxyl transferase subunit alpha (J1C87_09410: *accA*) and acetyl-CoA carboxylase biotin
carboxylase subunit (J1C87_09420: *accC*) (Figure 3). Similarly, the suppressors identified in the
populations of two ribosomal protein mutants *(*Δ*rpmH* and Δ*rplV)* and two GTPase mutants
(Δ*yihA* and Δ*obgE*) include mutations in 50S ribosomal protein *rplB* (J1C87_00620) and RNA
biosynthesis genes, such as *rpoB* (J1C87_01040) (Figure 3 and Supplemental Table 5). These
findings indicate that mutated segments shared across multiple evolved mutants can confer
fitness benefits to mutants deleted of different essential genes.
314

We also observed mutual suppressors, where an evolved population deleted of the first essential
ORF could contain a mutated segment within a second essential ORF, while an evolved
population deleted of the second essential ORF could contain a mutated segment within the first
essential ORF. For example, in the *obgE* deletion mutant *(ΔobgE)*, a mutation in *dnaK* occurred
(Figure 3 and Supplemental Table 5); conversely, in *ΔdnaK*, a mutated segment of *obgE* was
detected (Figure 3 and Supplemental Table 5). Similarly, in the F1Fo subunit deletion mutant
ΔuncF, a mutation of *dpcK* was also present (Figure 3 and Supplemental Table 5). In *ΔdpcK*,
mutated segments were detected within *uncA* and *uncD* (Figure 3 and Supplemental Table 5).
Other mutual suppressor pairs include those between subunits of F1Fo (Figure 3 and
Supplemental Table 5) and between components of the FASII system (Figure 3 and
Supplemental Table 5). These results suggest the presence of positive epistasis interactions
between different essential gene(s), where strains containing mutations in both essential genes
are fitter than those containing mutations in either of the essential genes.
328
329 **Possible explanations for parallel suppressor evolution in mutants deleted for particular**
330 **essential genes**
331
Given the potential role of parallel evolution for fitness improvement,[36-38] we next sought to
identify the most commonly mutated segments for each essential gene mutant and those
clustered in a particular operon. Utilizing these highly occurring suppressors (Supplemental
Table 5 and Supplemental Table 16), along with candidate genes within duplicated regions
(Figure 2B-H), we aimed to elucidate the mechanisms by which they improved the growth of
mutants deleted for essential genes.
338
Δf1fo suppressors
F1Fo is proposed to function as an ATP-dependent proton pump in streptococci, including *S.*
mutans and *S. sanguinis.*[39] Mutations in *trkA1* or *trkH1* are present in all thirteen evolved
populations of *Δf1fo*, with twelve populations containing *trkH1* mutations and one containing a
trkA1 mutation (Figure 4, Supplemental Table 5 and Supplemental Table 16). Similarly,
mutations in *trkA1*, *trkH1*, or a one base pair (bp) insertion mutation located 38 bp upstream of
the *trkA1-H1* operon (P_trkA1-H1, "P" for promoter) are found in twenty-five out of twenty-six
(96.2%) of the evolved populations deleted of single F1Fo subunits (Figure 4, Supplemental
Table 5 and Supplemental Table 16). Among the twenty-three mutations in the ORFs of *trkA1-*
H1, sixteen populations have *trkH1* mutations, and eight have *trkA1* mutations. Given the
comparable length of *trkA1* (1350 bp) and *trkH1* (1440 bp), it appears that mutations in the *trkH1*
gene are more likely than mutations in *trkA1* to improve the fitness of mutants deleted for F_1F_o
subunits.
352
We also observed that gene duplications occurred in the same region across populations deleted
of all F1Fo subunits or single subunits, i.e., *Δf1fo#107* and *#109* (Figure 2B and Fig S4B),
ΔuncE#2 (Fig S4) and *ΔuncH#7* (Figure 2C). This region of approximately 103 kb (coordinates:
22,738-125,834) contains the *v1vo* operon encoding the V1Vo-ATPase, which has been
proposed to function as a proton pump [31] and may compensate for the loss of *f1fo*. It is interesting
to note that *ΔuncH#7* is the only evolved population listed above that lacks mutations in the
flanking upstream regions or the ORFs of *trkA1* or *trkH1* (Figure 2C). More interestingly, among
the evolved populations deleted of F_1F_o subunits (thirteen *Δf1fo* and twenty-six mutants deleted

for single *unc* subunit genes), Δ*uncE#2* and Δ*uncH#7* are the only two populations that contain
amino acid substitution mutations of V-type ATP synthase subunits (Supplemental Table 5 and
Supplemental Table 16).
364
365 Suppressor mutations appeared in other F1Fo subunits or intergenic regions of *f1fo* in many
366 evolved populations deleted of a single subunit (Figure 4). This result suggests that for the F_1F_o
367 complex, when one subunit is absent, other subunits cannot function properly and their presence
368 is detrimental to the mutant.
369
370 Given that V1Vo may also function as an ATP-dependent proton pump [31] and the Trk system
371 works as a symporter (K+/Na+ or K+/H+), [40] we hypothesize that *v1vo* can partially compensate
372 for the loss of *f1fo*, that TrkA1-H1 is the major contributor to acidification of the cytosol when
373 pumping in K+, and that the function of TrkA1-H1 must therefore be inhibited when F1Fo is not
374 present (Figure 6).
375
ΔftsEX suppressors
FtsEX is crucial in regulating divisome assembly and activation at the plasma membrane and cell
wall. [41] In the evolved Δ*ftsX* populations, mutations in the *ftsE* gene of the *ftsEX* operon and the
vicRK (vicK/walK and *vicR/walR)* two-component system (TCS) frequently occurred (Figure 4).
Mutations in *ftsE* were present in eight out of nineteen evolved Δ*ftsX* populations (42.1%); in one
of these populations, Δ*ftsX#13*, a deletion mutation at 51 bp of the 693 bp *ftsE* ORF leads to a
frameshift in the N-terminal region (Figure 4). This observation suggests that in the absence of
ftsX, *ftsE* does not function properly and is deleterious to the cell. Although *vicR* and *vicK*
compose a TCS, *vicK* is not essential, while *vicR* is essential. [1,42] Fifteen populations contained
vicK mutations (with one frameshift and one truncation), and one population contained a *vicR*
mutation (an amino acid substitution, M54V), collectively representing 89.5% of the evolved
populations (Figure 4, Supplemental Table 5 and Supplemental Table 16). It was reported that
FtsEX interacts with the cell wall component peptidoglycan hydrolase, PcsB, which is positively
regulated by the *vicRK/walRK* two-component regulatory system. [43] Therefore, we hypothesize
that when FtsEX is absent and PcsB cannot function properly, there is a need to inhibit the
activity of PcsB's upstream regulatory factor, VicRK.
392
fab operon suppressors
The *fabH, fabF, fabG, fabZ, and fabK* genes belong to a FASII operon containing thirteen ORFs
in total, with a MarR family transcriptional regulator gene (*fabT*) upstream and seven other genes.
In the evolved populations of Δ*fabZ*, Δ*fabF*, Δ*fabG*, Δ*fabK*, and Δ*fabH*, we observed a
significant enrichment of mutations in genes encoding components of the FASII system (Figure
3, Figure 4, Supplemental Table 5, and Supplemental Table 16). This result suggests that when
one component of the FASII system is deleted, this can trigger genome changes in other FASII
components, leading to fitness improvement.
401
There is a notable number of populations containing a duplication of approximately 103 kb
(coordinate: 22,738-125,834) in the Δ*fabH,* Δ*fabF,* and Δ*fabG* mutants, which includes one ORF
encoding an acyltransferase and one ORF encoding acyl carrier protein. Additionally, a
duplicated region of 10.1 kb (coordinate: 1,093,824-1,103,958) is shared by Δ*fabZ #1* and Δ*fabZ*

#3 and contains acyl-acyl carrier protein thioesterase. We hypothesize that these duplications
may enhance fatty acid synthesis when *fabK, fabZ, fabH, fabF, or fabG* is defective.
408
ΔdpcK suppressors
DPCK is the final enzyme in the four-step pathway for CoA biosynthesis. [44] In the evolved
Δ*dpcK* populations*,* mutations ≤100 bp upstream of *acpS* (encoding holo-ACP synthase) are
present in eleven of the sixteen Δ*dpcK* populations. Holo-ACP synthase uses CoA as a substrate
for fatty acid biosynthesis. [45] Among the seventeen populations, one population, Δ*dpcK#17,*
contained a gene duplication; we were not able to identify the gene(s) responsible for improved
growth due to the large size of the duplication (>300 kb, containing 305 ORFs; Figure 2I). We
hypothesize that holo-ACP synthase is a significant contributor to the use of CoA, and when
there is a shortage of CoA, fatty acid production needs to be reduced by lessening the expression
of *acpS* (Figure 4, Supplemental Table 5 and Supplemental Table 16).
419
ΔpfkA suppressors
pfkA encodes 6-phosphofructokinase (PFK), responsible for catalyzing the phosphorylation of
fructose-6-phosphate to fructose-1,6-bisphosphate, a committing step in glycolysis. [46] In the
evolved Δ*pfkA* populations, we observed an enrichment of mutations in *fruR, fruB,* and *glck*.
Among the ten Δ*pfkA* evolved populations (Figure 4, Supplemental Table 5, and Supplemental
Table 16), six populations contained mutations in *fruR*, five populations had mutations in *fruB*,
and nine populations had mutations in *glcK*. The *glcK* gene encodes an ROK domain-containing
glucokinase, which phosphorylates glucose to produce glucose-6-phosphate, a progenitor of
fructose-6-phosphate. [47] The *fruR* gene encodes a transcriptional repressor of the *fruBA* operon,
while *fruB* (also known as *pfkB*) has been demonstrated to function as a 1-phosphofructokinase
in multiple bacterial species, [48] including streptococci. [49] Interestingly, the nature of the mutations
differed according to the gene; three of the eleven *glcK* mutations were truncations, and four of
the seven *fruR* mutations were truncations or frameshift mutations, while none of the seven *fruB*
mutations were truncations or frameshifts (Supplemental Table 5 and Supplemental Table 16).
434
We hypothesize that in the absence of *pfkA*, accumulated fructose-6-phosphate becomes toxic, as
has been shown previously in other bacterial species. [48] Elimination of GlcK activity would
reduce fructose-6-phosphate production. We further hypothesize that FruB possesses low levels
of fructose-6-phosphate kinase activity, thus replacing PfkA, but only when the expression of
fruB is increased due to *fruR* mutation or the fructose-6-phosphate kinase activity of FruB is
increased due to alteration of its sequence. This mechanism would be similar to findings in *E.*
coli, where a mutation that increased *fruB* expression was found to suppress the phenotype of
Δ*pfkA* mutants. [50,51]
443
ΔobgE suppressors
ObgE belongs to the TRAFAC (Translation Factor Association) class of P-loop GTPases, which
is highly conserved and essential in bacteria. [52] The ObgE GTPase is known to play a wide range
of functions, with some of the most significant roles including sporulation initiation in *Bacillus*
subtilis, [53] aerial mycelium formation in *Streptomyces*, [54,55] bacterial persistence in *E. coli* and
Pseudomonas aeruginosa, [56] involvement in ribosome biogenesis, [57,58] participation in DNA
replication, [59-61] and contributions to chromosome segregation and cell division. [62]
451

Among the fourteen evolved Δ*obgE* populations, eight populations had substitution mutations
upstream of *rpoB* (seven were an A to T mutation 321 bp upstream of the start codon, and the
other was an A to G mutation 24 bp upstream), and two populations contained substitution
mutations within the *rpoB* ORF (one at 905 bp and the other at 1058 bp from the start of the
3567 bp ORF). In total, 71.4% of the populations had mutations within or upstream of *rpoB*
(Figure 3, Supplemental Table 5 and Supplemental Table 16)). The *rpoB* gene codes for the beta-
subunit of RNA polymerase. In one of these populations, Δ*obgE#5*, a 37 kb gene duplication was
observed (coordinate: 1,775,394-1,812,805), encompassing thirty ORFs, including one annotated
as Der GTPase. It was reported that overexpression of *der* or *obgE* can suppress growth defects
of a mutant lacking *rrmJ* that encodes the methyltransferase for U2552 of the 23S rRNA. [63] From
this, we hypothesize that the duplication may partially compensate for the loss of ObgE function
by increasing *der* copy numbers, and a reduction in RNA synthesis is a significant fitness
adaptation when ObgE is defective.
465
466 *WT and other mutant suppressors*
467 Suppressor mutations in the evolved populations of Δ*pngM*, Δ*yqeG*, Δ*dnaK*, Δ*yihA* Δ*rpmH*,
468 Δ*rplV*, and WT are less easily explained, and further study is required to identify possible
469 mechanisms (Figure 4 and S16).
470
471 **Testing the working model involving *f1fo, v1vo, trkA1-H1* and *trkA2-H2***
472
473 To test the hypotheses proposed above, we have chosen F1Fo-ATPase, an evolutionarily ancient
474 enzyme complex that is highly conserved across virtually all forms of cellular life [64] and has
475 been reported as essential for entire genus of the streptococcal species. [30]
476
We began by testing whether increasing copy numbers of *v1vo* contributes to the fitness
improvement of the mutant deleted for the entire F1Fo complex or single subunits. To explore
this, we examined the gene duplication of the *v1vo* region at the early stage, P1 in the four
populations that contain gene duplications at P6 (Figure 2B-C, Fig S4B-C, and Fig S7A). Among
these four populations, one had the gene duplication at P1, while the other three acquired the
duplication during subsequent passages (Figure 5A and Fig S7A). This observation indicates that
gene duplications of the *v1vo* region evolved in these populations after initial passages.
484
485 To further confirm the increased fitness of *v1vo* region duplications in populations lacking *f1fo*,
486 we subcultured five Δ*f1fo#4* populations from the P6 stage, which did not initially contain gene
487 duplications in the *v1vo* region. After nine additional passages (P15), we observed that two out
488 of the five evolved populations had *v1vo* region duplications in P15 (Fig S7B).
489
To ascertain the fitness improvement by increasing *v1vo* copy numbers in mutants lacking *f1fo*,
we tested whether *v1vo* could be deleted in Δ*f1fo*. We first deleted the entire *v1vo*, which consists
of nine subunits, to generate the Δ*v1vo* mutant in the WT background. This Δ*v1vo* mutant grew
indistinguishably from the wild type in a rich medium (data not shown), demonstrating that *v1vo*
can be deleted when *f1fo* is intact (Figure 5C). However, when we attempted to delete *v1vo* in
Δ*f1fo#4*, where *f1fo* was removed, it resulted in a double-band genotype in all five Δ*v1vo*-
DB/Δf1fo#4 ("DB" for double-band) lines tested (Figure 5C and Fig S8). Consistent with these
findings, Δ*uncE#2* and Δ*uncH#7* contain amino acid substitution mutations of V1Vo ATP

synthase subunit A and K in the *v1vo* operon, respectively, also contain *v1vo* region duplications
(Supplemental Table 5). This finding indicates that increased copy numbers of the *v1vo* genes
compensated for the loss of F1Fo function. Overall, these results suggest that increasing the copy
numbers of *v1vo* contributes to the fitness improvement of mutants lacking *f1fo*.
502
We then investigated the impact of inhibiting *trkA1-H1* on enhancing the fitness of *Δf1fo*.
Initially, we compared the occurrence of *trkA1-H1* mutations between P1 and P6 stages across
twelve independently evolved populations of *Δf1fo* (*#101* to *#112*). While mutations in the *trkA1*
or *trkH1* segments were not detected in P1 using a 30% threshold in nine of the twelve
populations, we observed *trkA1-H1* mutations in all populations, with the mutation frequency
increasing during passage in all six independently evolved populations (Figure 5D and Table
S17). These mutations resulted in at least one frameshift and various substitutions in the
translation of TrkA1 or TrkH1 in six populations, while in the remaining populations, they led to
a histidine insertion or various substitutions in the TrkH1 protein (Supplemental Table 17). This
data indicates the fixation of *trkA1-H1* mutations during passage.
513
To further investigate the impact of *trkA1-H1*, we compared the growth of *Δf1fo* in the presence
and absence of *trkA1-H1*. *S. sanguinis* SK36 has two Trk systems, *trkA1-H1* and *trkA2-H2*. The
growth of *trkH2*, or double-gene deletion mutants of *trkA1-H1* or *trkA2-H2,* were
indistinguishable from that of WT (Figure 5E, Fig S9A-F and Supplemental Table 18). However,
combined deletion of either *trkA1* or *trkH1* along *trkH2* resulted in growth defects (Figure 5E-G,
Fig S9G-H, and Supplemental Table 18), indicating that *trkA1-H1* and *trkA2-H2* are redundant
for growth. In order to determine the effect of mutating the two Trk systems on the fitness of
Δf1fo, we transformed the *Δf1fo* construct into the backgrounds of *ΔtrkA1-H1*or *ΔtrkA2-H2*, and
compared the growth of *Δf1fo/ ΔtrkA1-H1* mutant and that of *Δf1fo/ΔtrkA2-H2*. As the controls,
we also included *Δf1fo#4*, which contains a frameshift mutation of *ΔtrkH1* in the *Δf1fo*
background (Supplemental Table 5). It showed that while the growth of *Δf1fo#4* and
Δf1fo/ΔtrkA1-H1 were comparable to each other, the growth of *Δf1fo/ΔtrkA2-H2* was
significantly inhibited compared to that of *Δf1fo#4* and *Δf1fo/ΔtrkA1-H1*. (Figure 5F-G and
Supplemental Table18). These findings suggest that inhibiting *trkA1-H1*, but not *trkA2-H2*,
contributes to the fitness improvement of evolved mutants deleted for *f1fo*.
529
In lactic acid bacteria, F1Fo-ATPase is believed to function by pumping out protons to alkalinize
the cytosol or to build negative internal and positive external membrane potentials. [65] To test
whether the growth defects in the *S. sanguinis Δf1fo* mutant are due to impaired function in
alkalinizing the cytosol or building a membrane potential, we compared the growth of *Δf1fo#4* in
a rich medium supplemented with lactic acid (which increases membrane potential and cytosolic
proton concentration) or NaOH (which lowers membrane potential and cytosolic proton
concentration). We observed that *Δf1fo#4* is susceptible to lactic acid treatment compared to WT
(Figure 5H and Supplemental Table 19). Conversely, the growth defect of *Δf1fo#4* can be
partially rescued by NaOH within a certain range (Figure 5I and Supplemental Table 19). These
results suggest that *f1fo* in *S. sanguinis* functions by pumping out protons to alkalinize the
cytosol.
541
We therefore suggest the following model for the relationship of *f1fo* with *v1vo, trkA1-H1,* and
trkA2-H2 (Figure 6). For streptococci, which do not contain the tricarboxylic acid cycle or

electron transport chain, energy production is mainly through lactic acid fermentation. [66] The
lactate from fermentation creates highly acidic cytosol, which would be toxic if H+ were not
exported. F1Fo pumps out H+ from the cytosol using ATP, and V1Vo-ATPase supplements the
function of F1Fo when the F1Fo is inhibited/inactivated. Both TrkA1-H1 and TrkA2-H2 are
potassium transporters and functionally redundant for that purpose. TrkA1-H1 pumps in
potassium together with H+, acidifying the cytosol when H+ is not exported. In contrast, TrkA2-
H2 does not pump protons into the cytosol when pumping in potassium (likely instead pumping
in Na+). Therefore, the function of TrkA1-H1 must be inhibited if F1Fo is not functional.
552
553 **Conserved functional relationships of *f1fo*, *trkA1-H1*, *trkA2-H2* and *v1vo* in different**
554 **streptococci**
555
To generalize the *f1fo* working model and assess F1Fo's essentiality in other strains and species,
we conducted essentiality evaluations in three other streptococci: *S. mutans* UA159, *S. sanguinis*
SK405, and *S. sanguinis* SK1058. Genomic sequencing of *S. mutans* UA159, [67] *S. sanguinis*
SK405 [33,] and *S. sanguinis* SK1058 [33] showed that *f1fo* and *trkA1-H1* are present in all three
species, as in SK36, while *v1vo* is present in the *S. sanguinis* strains only. *TrkA2-H2* is present in
both *S. sanguinis* SK36 and SK405 but not in *S. sanguinis* SK1058 or *S. mutans* UA159 (Figure
7A).
563
To assess whether *v1vo* could serve as a partial backup in cases of F1Fo loss-of-function, we
attempted to delete *f1fo* in *S. mutans* UA159, which lacks *v1vo*. Despite multiple attempts and a
total of 64 independent genotyping analyses, none of the transformants were deleted for *f1fo*,
instead producing a double band indicative of gene duplication prior to allelic exchange (Fig
S10A-B). This result highlights that in *S. mutans* UA159, lacking *v1vo, f1fo* cannot be deleted in
our condition. This observation supports the evolutionarily conserved relationship between *f1fo*
and *v1vo* observed in other streptococcal species.
571
572 We then investigated the impact of deleting *trkA1-H1* in the other two strains: *S. sanguinis*
573 SK1058, which only possesses *trkA1-H1*, and *S. sanguinis* SK405, which has both *trkA1-H1* and
574 *trkA2-H2*. Our findings showed that *trkA1-H1* transformants exhibited severe growth defects in
575 the SK1058 background compared to the SK405 background (Fig S11A-C). This result
576 demonstrates the requirements of the *Trk* system for growth within the species of *S. sanguinis*.
577
Next, we investigated the possibility of deleting *f1fo* in *S. sanguinis* SK1058 and SK405, both of
which contain *v1vo* and *trkA1-H1*. We could generate *f1fo* whole-region deletion mutants (*Δf1fo*)
in both SK1058 and SK405 backgrounds (Figure 7B, 7C, Fig S12). Furthermore, the growth of
mutants with *f1fo* deletions in SK405 was notably better than in SK1058 (Figure 7B, 7D, and
Supplemental Table 20). To explore the potential emergence of *trkA1-H1* suppressor mutations
during evolution, we conducted passage experiments using independently evolved populations
with *f1fo* deletions in SK405 (Fig S12A: three populations) and in SK1058 (Fig S12B: two
populations). It was observed that, after six passages, *trkA1-H1* suppressor mutations appeared in
all evolved *Δf1fo* populations in SK405 and SK1058 (Figure 7E). At least one population of
SK405 or SK1058 background are frameshift or truncation mutations of *trkA1-H1* (Figure 7E).
This result underscores the evolutionarily conserved occurrence of *trkA1-H1* suppressor
mutations in evolved populations with *f1fo* deletions.

590
591 **Discussion**
592
While identifying genes that interact with essential genes or compensate for their loss provides a
great benefit, [4] this task remains challenging, primarily due to the unavailability of essential-gene
deletion mutants and systematic methodologies to connect the function of essential genes with
their partners. Traditional approaches, such as antisense RNA knockdowns [48] and CRISPR
interference, [68,69] while valuable, may exhibit limitations, including incomplete target gene
silencing and suboptimal design efficacy, [69] thereby potentially compromising the reliability of
essential gene functional analyses. Our study successfully established a system to generate viable
mutants with deletions in essential genes (Figure 1 and Fig S1-S2). We observed that small
colonies likely represent the deletion mutants, while large colonies often exhibit the "double-
band" genotype or kanamycin resistance due to mutations in translational machinery.
603
With these observations, we have generated essential gene deletion mutants, many of which are
also essential across multiple species (http://www.essentialgene.org/). These essential gene
mutants provide an invaluable resource for dissecting the functions of essential genes. Using
these twenty-four essential gene mutants and WT as controls, we identified over 1000
spontaneous suppressor mutations in passage experiments (Figure 2). Leveraging these essential
gene mutants and distinct mutated segments, we constructed a functional network (Figure 3).
Within this network, we made several noteworthy observations. 1) Mutated segments that
appeared in the evolved WT population can also appear in the evolved populations of mutants
with deletions of essential genes, such as mutations in the PTS mannose/fructose/sorbose
transporter family subunit. 2) Suppressor mutations in certain genes, such as *rpoB* and *rpsL,* that
are found in the evolved populations of one deletion mutant can also be present in the evolved
populations of other mutants (Supplemental Table 5). 3) Mutual essential gene suppressor
mutations were discovered, where mutations in the first essential gene were found within a
mutant population deleted for the second essential gene, and vice versa.
618
As expected, our findings support the notion that mutants with deletions of essential genes face
stronger selective pressures when compared to that of the WT. These mutants are more likely to
undergo the following significant evolutionary changes. 1) Gene duplications of greater than 10
kb: Our data revealed large-scale gene duplications exclusively were present in mutants deleted
of essential genes (Figure 2B-H). 2) Substitutions, insertions, and deletions of less than 10 kb:
Small-scale mutations, especially those of >50% of sequence reads, were more common in
mutants deleted for essential genes (Figure 2I). 3) Mutations in essential genes: Mutations in
other essential genes were observed more exclusively in mutants deleted for an essential gene
(Figure 2K). 4) Mutations in multi-drug resistance genes: mutations of multi-drug resistance-
associated genes, such as *rpoB* and *rpsL,* are present in several evolved populations deleted for
essential genes, suggesting a shared selective pressure resulting from drug treatment and deletion
of essential genes (Supplemental Table 5). The suppressor mutations identified in our evolved
populations deleted of essential genes may reflect the differential gene essentiality over
evolutionary time in the pan-genome [11] and reproducibly changing gene essentiality across
independently evolving populations. [13]
634

Unexpectedly, as discussed above, for every essential gene mutated, we identified at least one
population containing a suppressor mutation in another essential gene (Figure 2L). Notably, the
essential genes were inactivated by the suppressor mutation in some of the evolved populations,
such as the >200bp insertion mutation in *ftsZ* in the evolved population of *Δf1fo #107*, the >200
bp deletion mutation of *uncH* in *ΔuncC#1*, and the 1 bp deletion mutation of *ftsE* in *ΔftsX#13*
(Supplemental Table 5). In contrast, for other essential gene mutation suppressors, the mutations
were not within the open reading frame (ORF) but were found within 400 base pairs upstream of
the start codons, likely affecting expression rather than inactivating function. For instance, in the
evolved *ΔdpcK* populations, the suppressors with the highest mutation frequency were located
upstream of the holo-ACP synthase gene, *acpS* (Figure 4). Similarly, in the evolved *ΔobgE*
populations, the majority of mutation suppressors were found upstream of *rpoB* (Figure 4).
646
Essential genes represent ideal drug targets for controlling infectious diseases and cancers. In
traditional medical treatment, the focus has primarily been on targeting a single essential gene
product. However, it's crucial to consider the evolution of gene essentiality during drug treatment
and the development of drug resistance over short-term evolution. We have observed that in
multiple populations deleted of essential genes, mutations in multiple segments appeared within
six passages, including those belonging to multi-drug resistance-associated genes (Figure 2M).
Considering that some of the suppressor mutations were gene duplications that presumably
confer fitness benefits by increasing gene copy numbers, a promising approach is to target not
only the essential gene itself but also the backup gene. This strategy can help prevent or reduce
the development of drug resistance.
657
To understand the relationships between the structure and function of a protein, it is of great
interest to know which amino acids or domains of the proteins are sensitive to mutations. In this
study, we have observed numerous mutations of *trkA1* and *trkH1* in *Δf1fo* or *f1fo* single subunit
deletion mutants, *vicK* in *ΔftsX*, *fruR* and *glcK* in *ΔpfkA*, as well as *fusA*, *rpsL* and *rplF* in
kanamycin-resistant mutants lacking the *kan* gene (Supplemental Table 1-2). These mutations
represent fitness-related protein sequence changes that improve the growth of resultant mutants
in a particular selective condition.
665
In this work, we have demonstrated the efficacy of the methodology developed for generating
deletion mutants of numerous essential genes in *S. sanguinis* SK36. Furthermore, we have
highlighted the potency of experimental evolution [24,25] in unraveling the mechanisms of gene
essentiality, with a particular focus on *f1fo* in *S. sanguinis* and *S. mutans*. It would be interesting
to test the conservation or divergence of mechanisms governing gene essentiality of other genes,
as well as across different *Streptococcus* species and other genera. While extracting valuable
information by identifying spontaneous suppressor mutations in populations lacking essential
genes is possible, conducting experimental evolution requires caution. First and foremost, the
choice of an appropriate organism is crucial. Ideally, the selected organism should have a single
chromosome, a small genome size, and a rapid cell division rate. Streptococci represent an ideal
candidate in this regard. They possess a single circular genome with a relatively modest size,
typically ranging from 2.0 to 3.0 megabases. [70] This choice contrasts with a previous study that
employed budding yeast with multiple chromosomes [4] where most mutations in populations
lacking essential genes were related to changes in ploidy, which could hinder the identification
of causal suppressors. On the other hand, the identification of mechanisms of genetic suppression

requires caution. For the suppressor mutations of low frequency or these substitution mutations
where we were unable to determine whether the evolved suppressor mutations increase or
decrease the activity of the gene product, further evidence is needed to understand the
mechanisms underlying the evolution of gene essentiality.
685
In general, essential genes tend to be more conserved and evolutionarily ancient compared to
non-essential genes. [54] It is intriguing to consider when genes emerge within a biological system
over time. The suppressor mutations in essential gene mutants, particularly the loss-of-function
mutations, can provide valuable insights into dependency relationships. For instance, the absence
of *f1fo* in a system may hinder the emergence of the *trkA1-H1* system. In cases where essential
genes function as a complex, such as *f1fo* or *ftsEX*, our findings indicate that deletion of one
component may result in additional deleterious effects caused by the remaining subunit(s)
(Supplemental Table 5). This suggests that all subunits of the complex need to be integrated into
the system, at least for these cases. Moreover, understanding the timelines of gene emergence in
a system is also crucial for guiding decisions in synthetic biology, determining which genes, as
well as the order of genes, to be introduced into a system.
697
698 **Materials and Methods**
699
Strains Used and Growth Conditions:
In this study, we primarily utilized *Streptococcus sanguinis* strain SK36. [27] *S. sanguinis* SK405
and SK1058 strains were described previously. [33] *S. mutans* UA159 was obtained from Ann
Progulske-Fox (University of Florida). All bacterial cultures were routinely maintained in a BHI
medium at 37°C under anaerobic conditions, using a Coy anaerobic chamber, or stored at -80 °C
for long-term storage.
706
707 *Transformation and Mutant Isolation:*
708 To prepare competent cells, *S. sanguinis* was cultured in 1 mL of Todd-Hewitt + HS medium [10]
709 and incubated overnight in anaerobic conditions at 37°C. The following day, a 5 μL aliquot of
710 the overnight culture was added to 1 mL of fresh TS+HS medium and incubated for an additional
711 3 hours.
712
For creating knockout constructs, we employed 1 kb of homologous 3' and 5' end sequences
flanking the coding sequence (CDS) and resistance genes, *kan* or *Erm*[r], in overlapping extension
PCR. [27] For transformation, we mixed 50 ng to 500 ng of PCR product DNA with 200 ng of CSP
and 300 μL of competent cells. The reactions continued for 24 hours under anaerobic conditions
at 37°C. Subsequently, transformation reaction mixtures were plated on BHI-agar containing
appropriate antibiotics (Kanamycin: 500 μg/mL or Erythromycin: 10 μg/mL) for selection. Once
the plates had dried, the cells were overlaid with a thin layer of agar by adding 1 mL of cooled
BHI-agar with antibiotics. The plates were incubated at 37°C under anaerobic conditions to
select transformants for 4 to 5 days.
722
For genotyping, genomic DNA (gDNA) was isolated from 2 mL of saturated cultures. In brief,
cells were precipitated by centrifugation at 10,000 rpm for 10 minutes at room temperature. They
were then resuspended in 200 μL of resuspension buffer (20 mM EDTA, 200 mM Tris-HCl, and
2% Triton-X 100) and lysed with 200 μL of AL lysis buffer (Qiagen, 19075). DNA was

precipitated by adding 1 mL of 100% ethanol supplemented with 100 mM of sodium acetate.
Following washing and drying, the DNA was dissolved in 100 μL of water and used for PCR
and/or whole-genome sequencing.
730
To generate deletion mutants *Δf1fo* in *ΔtrkA1-H1* and *ΔtrkA2-H2*, the *ΔtrkA1-H1* and *ΔtrkA2-H2*
mutants were first generated using *Erm*[r] (Fig S9). After confirmation of deletion of *ΔtrkA1-H1*
and *ΔtrkA2-H2, Δf1fo* of *kan* resistance construct was subsequentially transformed into *ΔtrkA1-*
H1 or *ΔtrkA2-H2*. To generate deletion mutants of both *trkA1-H1* and *trkA2-H2*, *trkA1*, *trkH1*, or
ΔtrkA1-H1 of erythromycin resistance constructs were transformed into a *ΔtrkH2* mutant of kan
resistance background (Xu, et al., 2011). To isolate *Δf1fo* and *ΔtrkA1-H1* mutants in the SK405
and SK1058 strains, we employed the same PCR construct used for deleting *f1fo* and *trkA1-H1*
in SK36, as there is >90% identity at the flanking regions between SK36, SK1058, and SK405.
739
Strain Passage:
Antibiotic-resistant colonies from selection plates were transferred to 1 mL of BHI containing
antibiotics and grown for 2 to 8 days under anaerobic conditions at 37°C. The duration depended
on when the OD_{600} reached 0.1-0.5, which was referred to as P0 (see Figure 1C). A 300 μL
aliquot of the cell culture was then transferred to 3 mL of BHI containing antibiotics, and growth
continued until saturation, resulting in P1. P1 was mixed by pipetting five times with a P-1000
pipette, and 50 μL of the mixed P1 was transferred to 1 mL of BHI containing antibiotics.
Growth proceeded until the cell culture reached an OD_{600} of 0.1 to 0.5, producing P2. The latter
process was repeated sequentially from P3 to P6, where the volume of P2 to P5 was 1 mL and
the final passage, P6 was 3 mL. Cell cultures of P1 and P6 were stored at -80°C in glycerol at the
final concentration of 20%. DNA extractions and genome sequencing were conducted in P6
and/or P1 when indicated.
752
753   For passages involving WT, glycerol stocks from -80°C were transferred to 1 mL of BHI and
754   grown for 24 hours, resulting in T0 (see Fig S4A). A 50 μL aliquot of T0 cell culture was
755   aliquoted into six 1.5 mL tubes, each containing 1 mL of BHI, and grown for 24 hours, resulting
756   in T1. A 50 μL aliquot of T1 cell culture was inoculated into 1 mL of BHI and grown for 24
757   hours, leading to T2. This process was repeated for T3 to T9.
758
759   *Whole-Genome Sequencing and Variant Calling:*
760   Whole-genome sequencing was carried out by the Genome Core at Virginia Commonwealth
761   University or by SeqCenter (https://www.seqcenter.com/) using the shotgun method with 2x150
762   paired-end sequencing. Whole-genome sequences with an average coverage of ≥200 were used
763   for subsequent analysis.
764
Fastq files were aligned to the SK36 reference genome [16] using Geneious Prime software
(https://www.geneious.com/) after trimming via the BBDuk method. Variations in the genome
were exported from Geneious Prime. To identify the mutated segment, the frequencies
(percentages) of all mutations belonging to a certain segment were assessed and the greatest
value of percentages were assigned to that segment. For the deletions, the read coverages were
calculated by averaging the base coverages that were 100 bp from the flanking sequence. For
network analysis, the software Cytoscape was used.[71]
772

Contributions

L.B. and P.X. conceived the project. L.B., T.K. and P.X. designed the experiments. L.B. constructed all the essential gene mutants and performed all experimental evolution studies. L.B., T.K., and P.X. analyzed the data with the help of Z.Z., A.I., B.Z., V.A. and M.W.. The manuscript was written and revised by L.B., T.K., and P.X..

Acknowledgments

The authors would like to thank Dr. Mark Running at the Department of Biology, University of Louisville, Dr. Tomomichi Fujita at the Department of Biological Sciences, Hokkaido University, Dr. Tim van Opijnen at Broad Institute, MIT and Harvard, and Dr. Yan Bao at the School of Agriculture and Biology, Shanghai Jiao Tong University, for discussions regarding experimental design and critical reading of the manuscript. We also thank Dr. Junling Ren at the Philips Institute for Oral Health Research, Virginia Commonwealth University for help with network analysis and Ms. Ayana Wyche at Virginia Commonwealth University for assistance with the culturing of bacteria. We are grateful to Vladimir Lee and Kalyan Maliempati for genomic sequencing services. This work was funded by grants from the National Institutes of Health R01DE030121 (P.X. and T.K.) and the CCTR Endowment Fund (P.X.).

Declaration of generative AI in scientific writing

During the preparation of this work the authors used ChatGPT-3 in order to improve readability and language. After using it, the authors reviewed and edited the content as needed and took full responsibility for the content of the publication.

Declaration of interests

A patent application has been submitted by P.X. and L.B. and the Virginia Commonwealth University based on these results.

References

1. Xu, P., Ge, X., Chen, L., Wang, X., Dou, Y., Xu, J.Z., Patel, J.R., Stone, V., Trinh, M., Evans, K., et al. (2011). Genome-wide essential gene identification in *Streptococcus sanguinis*. Scientific reports *1*, 125. 10.1038/srep00125.
2. Rancati, G., Moffat, J., Typas, A., and Pavelka, N. (2018). Emerging and evolving concepts in gene essentiality. Nat Rev Genet *19*, 34-49. 10.1038/nrg.2017.74.
3. Costanzo, M., VanderSluis, B., Koch, E.N., Baryshnikova, A., Pons, C., Tan, G., Wang, W., Usaj, M., Hanchard, J., Lee, S.D., et al. (2016). A global genetic interaction network maps a wiring diagram of cellular function. Science *353*. 10.1126/science.aaf1420.
4. Jana, B., Liu, X., Dénéréaz, J., Park, H., Leshchiner, D., Liu, B., Gallay, C., Veening, J.W., and van Opijnen, T. (2023). CRISPRi-TnSeq: A genome-wide high-throughput tool for bacterial essential-nonessential genetic interaction mapping. bioRxiv. 10.1101/2023.05.31.543074.

5.   Nijman, S.M. (2011). Synthetic lethality: general principles, utility and detection using
genetic screens in human cells. FEBS Lett *585*, 1-6. 10.1016/j.febslet.2010.11.024.
6.   Leaver, M., Domínguez-Cuevas, P., Coxhead, J.M., Daniel, R.A., and Errington, J.
(2009). Life without a wall or division machine in *Bacillus subtilis*. Nature *457*, 849-853.
10.1038/nature07742.
7.   Mutschler, H., Gebhardt, M., Shoeman, R.L., and Meinhart, A. (2011). A novel
mechanism of programmed cell death in bacteria by toxin-antitoxin systems corrupts
peptidoglycan synthesis. PLoS Biol *9*, e1001033. 10.1371/journal.pbio.1001033.
8.   Larrimore, K.E., and Rancati, G. (2019). The conditional nature of gene essentiality in
Pseudomonas aeruginosa. Curr Opin Genet Dev *58-59*, 55-61.
10.1016/j.gde.2019.07.015.
9.   Liu, G., Yong, M.Y., Yurieva, M., Srinivasan, K.G., Liu, J., Lim, J.S., Poidinger, M.,
Wright, G.D., Zolezzi, F., Choi, H., et al. (2015). Gene Essentiality Is a Quantitative
Property Linked to Cellular Evolvability. Cell *163*, 1388-1399.
10.1016/j.cell.2015.10.069.
10.   Rancati, G., Pavelka, N., Fleharty, B., Noll, A., Trimble, R., Walton, K., Perera, A.,
Staehling-Hampton, K., Seidel, C.W., and Li, R. (2008). Aneuploidy underlies rapid
adaptive evolution of yeast cells deprived of a conserved cytokinesis motor. Cell *135*,
879-893. 10.1016/j.cell.2008.09.039.
11.   Bao, L., Inoue, N., Ishikawa, M., Gotoh, E., Teh, O.K., Higa, T., Morimoto, T., Ginanjar,
E.F., Harashima, H., Noda, N., et al. (2022). A PSTAIRE-type cyclin-dependent kinase
controls light responses in land plants. Sci Adv *8*, eabk2116. 10.1126/sciadv.abk2116.
12.   Rosconi, F., Rudmann, E., Li, J., Surujon, D., Anthony, J., Frank, M., Jones, D.S., Rock,
C., Rosch, J.W., Johnston, C.D., and van Opijnen, T. (2022). A bacterial pan-genome
makes gene essentiality strain-dependent and evolvable. Nat Microbiol *7*, 1580-1592.
10.1038/s41564-022-01208-7.
13.   Couce, A., Limdi, A., Magnan, M., Owen, S.V., Herren, C.M., Lenski, R.E., Tenaillon,
O., and Baym, M. (2024). Changing fitness effects of mutations through long-term
bacterial evolution. Science *383*, eadd1417. 10.1126/science.add1417.
14.   Santamaría, D., Barrière, C., Cerqueira, A., Hunt, S., Tardy, C., Newton, K., Cáceres, J.F.,
Dubus, P., Malumbres, M., and Barbacid, M. (2007). Cdk1 is sufficient to drive the
mammalian cell cycle. Nature *448*, 811-815. 10.1038/nature06046.
15.   Nowack, M.K., Harashima, H., Dissmeyer, N., Zhao, X.A., Bouyer, D., Weimer, A.K.,
De Winter, F., Yang, F., and Schnittger, A. (2012). Genetic framework of cyclin-
dependent kinase function in *Arabidopsis*. Dev. Cell *22*, 1030-1040.
10.1016/j.devcel.2012.02.015.
16.   Glass, J.I., Merryman, C., Wise, K.S., Hutchison, C.A., and Smith, H.O. (2017). Minimal
Cells-Real and Imagined. Cold Spring Harb Perspect Biol *9*.
10.1101/cshperspect.a023861.
17.   Juhas, M., Eberl, L., and Church, G.M. (2012). Essential genes as antimicrobial targets
and cornerstones of synthetic biology. Trends Biotechnol *30*, 601-607.
10.1016/j.tibtech.2012.08.002.
18.   Hutchison, C.A., Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J.,
Ellisman, M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L., et al. (2016). Design and
synthesis of a minimal bacterial genome. Science *351*, aad6253. 10.1126/science.aad6253.

19.   Ratcliff, W.C., Denison, R.F., Borrello, M., and Travisano, M. (2012). Experimental
865   evolution of multicellularity. Proc Natl Acad Sci U S A *109*, 1595-1600.
866   10.1073/pnas.1115323109.
867   20.   Blount, Z.D., Borland, C.Z., and Lenski, R.E. (2008). Historical contingency and the
868   evolution of a key innovation in an experimental population of *Escherichia coli*. Proc
869   Natl Acad Sci U S A *105*, 7899-7906. 10.1073/pnas.0803151105.
870   21.   Moger-Reischer, R.Z., Glass, J.I., Wise, K.S., Sun, L., Bittencourt, D.M.C., Lehmkuhl,
871   B.K., Schoolmaster, D.R., Lynch, M., and Lennon, J.T. (2023). Evolution of a minimal
872   cell. Nature *620*, 122-127. 10.1038/s41586-023-06288-x.
873   22.   Kawecki, T.J., Lenski, R.E., Ebert, D., Hollis, B., Olivieri, I., and Whitlock, M.C. (2012).
874   Experimental evolution. Trends Ecol Evol *27*, 547-560. 10.1016/j.tree.2012.06.001.
875   23.   Barrick, J.E., and Lenski, R.E. (2013). Genome dynamics during experimental evolution.
876   Nat Rev Genet *14*, 827-839. 10.1038/nrg3564.
877   24.   McDonald, M.J. (2019). Microbial Experimental Evolution - a proving ground for
878   evolutionary theory and a tool for discovery. EMBO Rep *20*, e46992.
879   10.15252/embr.201846992.
880   25.   Van den Bergh, B., Swings, T., Fauvart, M., and Michiels, J. (2018). Experimental
881   Design, Population Dynamics, and Diversity in Microbial Experimental Evolution.
882   Microbiol Mol Biol Rev *82*. 10.1128/MMBR.00008-18.
883   26.   Taylor, T.B., Mulley, G., Dills, A.H., Alsohim, A.S., McGuffin, L.J., Studholme, D.J.,
884   Silby, M.W., Brockhurst, M.A., Johnson, L.J., and Jackson, R.W. (2015). Evolutionary
885   resurrection of flagellar motility via rewiring of the nitrogen regulation system. Science
886   *347*, 1014-1017. 10.1126/science.1259145.
887   27.   Xu, P., Alves, J.M., Kitten, T., Brown, A., Chen, Z., Ozaki, L.S., Manque, P., Ge, X.,
888   Serrano, M.G., Puiu, D., et al. (2007). Genome of the opportunistic pathogen
889   *Streptococcus sanguinis*. J Bacteriol *189*, 3166-3175. 10.1128/JB.01808-06.
890   28.   Lázár, V., Pal Singh, G., Spohn, R., Nagy, I., Horváth, B., Hrtyan, M., Busa-Fekete, R.,
891   Bogos, B., Méhi, O., Csörgő, B., et al. (2013). Bacterial evolution of antibiotic
892   hypersensitivity. Mol Syst Biol *9*, 700. 10.1038/msb.2013.57.
893   29.   Rhodes, D.V., Crump, K.E., Makhlynets, O., Snyder, M., Ge, X., Xu, P., Stubbe, J., and
894   Kitten, T. (2014). Genetic characterization and role in virulence of the ribonucleotide
895   reductases of *Streptococcus sanguinis*. J Biol Chem *289*, 6273-6287.
896   10.1074/jbc.M113.533620.
897   30.   Luo, H., Lin, Y., Gao, F., Zhang, C.T., and Zhang, R. (2014). DEG 10, an update of the
898   database of essential genes that includes both protein-coding genes and noncoding
899   genomic elements. Nucleic Acids Res *42*, D574-580. 10.1093/nar/gkt1131.
900   31.   Nelson, N., Perzov, N., Cohen, A., Hagai, K., Padler, V., and Nelson, H. (2000). The
901   cellular biology of proton-motive force generation by V-ATPases. J Exp Biol *203*, 89-95.
902   10.1242/jeb.203.1.89.
903   32.   Röttig, A., and Steinbüchel, A. (2013). Acyltransferases in bacteria. Microbiol Mol Biol
904   Rev *77*, 277-321. 10.1128/MMBR.00010-13.
905   33.   Baker, S.P., Nulton, T.J., and Kitten, T. (2019). Genomic, Phenotypic, and Virulence
906   Analysis of *Streptococcus sanguinis* Oral and Infective-Endocarditis Isolates. Infect
907   Immun *87*. 10.1128/IAI.00703-18.

34. Ghosh, A., N, S., and Saha, S. (2020). Survey of drug resistance associated gene
mutations in Mycobacterium tuberculosis, ESKAPE and other bacterial species.
Scientific reports *10*, 8957. 10.1038/s41598-020-65766-8.
35. Zeng, L., Walker, A.R., Lee, K., Taylor, Z.A., and Burne, R.A. (2021). Spontaneous
Mutants of Streptococcus sanguinis with Defects in the Glucose-Phosphotransferase
System Show Enhanced Post-Exponential-Phase Fitness. J Bacteriol *203*, e0037521.
10.1128/JB.00375-21.
36. Chevin, L.M., Martin, G., and Lenormand, T. (2010). Fisher's model and the genomics of
adaptation: restricted pleiotropy, heterogenous mutation, and parallel evolution.
Evolution *64*, 3213-3231. 10.1111/j.1558-5646.2010.01058.x.
37. Wichman, H.A., Badgett, M.R., Scott, L.A., Boulianne, C.M., and Bull, J.J. (1999).
Different trajectories of parallel evolution during viral adaptation. Science *285*, 422-424.
10.1126/science.285.5426.422.
38. Stern, D.L. (2013). The genetic causes of convergent evolution. Nat Rev Genet *14*, 751-
764. 10.1038/nrg3483.
39. Sheng, J., and Marquis, R.E. (2006). Enhanced acid resistance of oral streptococci at
lethal pH values associated with acid-tolerant catabolism and with ATP synthase activity.
FEMS Microbiol Lett *262*, 93-98. 10.1111/j.1574-6968.2006.00374.x.
40. Corratgé-Faillie, C., Jabnoune, M., Zimmermann, S., Véry, A.A., Fizames, C., and
Sentenac, H. (2010). Potassium and sodium transport in non-animal cells: the
Trk/Ktr/HKT transporter family. Cell Mol Life Sci *67*, 2511-2532. 10.1007/s00018-010-
0317-7.
41. Du, S., Pichoff, S., and Lutkenhaus, J. (2016). FtsEX acts on FtsA to regulate divisome
assembly and activity. Proc Natl Acad Sci U S A *113*, E5052-5061.
10.1073/pnas.1606656113.
42. Moraes, J.J., Stipp, R.N., Harth-Chu, E.N., Camargo, T.M., Höfling, J.F., and Mattos-
Graner, R.O. (2014). Two-component system VicRK regulates functions associated with
establishment of *Streptococcus sanguinis* in biofilms. Infect Immun *82*, 4941-4951.
10.1128/IAI.01850-14.
43. Sham, L.T., Barendt, S.M., Kopecky, K.E., and Winkler, M.E. (2011). Essential PcsB
putative peptidoglycan hydrolase interacts with the essential FtsXSpn cell division
protein in *Streptococcus pneumoniae* D39. Proc Natl Acad Sci U S A *108*, E1061-1069.
10.1073/pnas.1108323108.
44. Gupta, A., Singh, P.K., Sharma, P., Kaur, P., Sharma, S., and Singh, T.P. (2020).
Structural and biochemical studies of phosphopantetheine adenylyltransferase from
Acinetobacter baumannii with dephospho-coenzyme A and coenzyme A. Int J Biol
Macromol *142*, 181-190. 10.1016/j.ijbiomac.2019.09.090.
45. Heath, R.J., and Rock, C.O. (1996). Inhibition of beta-ketoacyl-acyl carrier protein
synthase III (FabH) by acyl-acyl carrier protein in *Escherichia coli*. The Journal of
biological chemistry *271*, 10996-11000. 10.1074/jbc.271.18.10996.
46. Phong, W.Y., Lin, W., Rao, S.P., Dick, T., Alonso, S., and Pethe, K. (2013).
Characterization of phosphofructokinase activity in *Mycobacterium tuberculosis* reveals
that a functional glycolytic carbon flow is necessary to limit the accumulation of toxic
metabolic intermediates under hypoxia. PLoS One *8*, e56037.
10.1371/journal.pone.0056037.

47.   Skarlatos, P., and Dahl, M.K. (1998). The glucose kinase of *Bacillus subtilis*. J Bacteriol
180, 3222-3226. 10.1128/JB.180.12.3222-3226.1998.
48.   Boulanger, E.F., Sabag-Daigle, A., Thirugnanasambantham, P., Gopalan, V., and Ahmer,
B.M.M. (2021). Sugar-Phosphate Toxicities. Microbiol Mol Biol Rev *85*, e0012321.
10.1128/MMBR.00123-21.
49.   Zeng, L., and Burne, R.A. (2021). Molecular mechanisms controlling fructose-specific
memory and catabolite repression in lactose metabolism by *Streptococcus mutans*. Mol
Microbiol *115*, 70-83. 10.1111/mmi.14597.
50.   Morrissey, A.T., and Fraenkel, D.G. (1972). Suppressor of phosphofructokinase
mutations of *Escherichia coli*. J Bacteriol *112*, 183-187. 10.1128/jb.112.1.183-187.1972.
51.   Babul, J. (1978). Phosphofructokinases from Escherichia coli. Purification and
characterization of the nonallosteric isozyme. The Journal of biological chemistry *253*,
4350-4355.
52.   Deckers, B., Vercauteren, S., De Kock, V., Martin, C., Lazar, T., Herpels, P., Dewachter,
L., Verstraeten, N., Peeters, E., Ballet, S., et al. (2023). YbiB: a novel interactor of the
GTPase ObgE. Nucleic Acids Res *51*, 3420-3435. 10.1093/nar/gkad127.
53.   Vidwans, S.J., Ireton, K., and Grossman, A.D. (1995). Possible role for the essential
GTP-binding protein Obg in regulating the initiation of sporulation in *Bacillus subtilis*. J
Bacteriol *177*, 3308-3311. 10.1128/jb.177.11.3308-3311.1995.
54.   Okamoto, S., Itoh, M., and Ochi, K. (1997). Molecular cloning and characterization of
the obg gene of *Streptomyces griseus* in relation to the onset of morphological
differentiation. J Bacteriol *179*, 170-179. 10.1128/jb.179.1.170-179.1997.
55.   Okamoto, S., and Ochi, K. (1998). An essential GTP-binding protein functions as a
regulator for differentiation in *Streptomyces coelicolor*. Mol Microbiol *30*, 107-119.
10.1046/j.1365-2958.1998.01042.x.
56.   Verstraeten, N., Knapen, W.J., Kint, C.I., Liebens, V., Van den Bergh, B., Dewachter, L.,
Michiels, J.E., Fu, Q., David, C.C., Fierro, A.C., et al. (2015). Obg and Membrane
Depolarization Are Part of a Microbial Bet-Hedging Strategy that Leads to Antibiotic
Tolerance. Molecular cell *59*, 9-21. 10.1016/j.molcel.2015.05.011.
57.   Feng, B., Mandava, C.S., Guo, Q., Wang, J., Cao, W., Li, N., Zhang, Y., Wang, Z., Wu,
J., Sanyal, S., et al. (2014). Structural and functional insights into the mode of action of a
universally conserved Obg GTPase. PLoS Biol *12*, e1001866.
10.1371/journal.pbio.1001866.
58.   Murti, K.G., Webster, R.G., and Jones, I.M. (1988). Localization of RNA polymerases on
influenza viral ribonucleoproteins by immunogold labeling. Virology *164*, 562-566.
10.1016/0042-6822(88)90574-0.
59.   Zielke, R., Sikora, A., Dutkiewicz, R., Wegrzyn, G., and Czyż, A. (2003). Involvement of
the cgtA gene function in stimulation of DNA repair in *Escherichia coli* and *Vibrio
harveyi*. Microbiology (Reading) *149*, 1763-1770. 10.1099/mic.0.26292-0.
60.   Czyz, A., Zielke, R., Konopa, G., and Wegrzyn, G. (2001). A Vibrio harveyi insertional
mutant in the *cgtA* (*obg*, *yhbZ*) gene, whose homologues are present in diverse organisms
ranging from bacteria to humans and are essential genes in many bacterial species.
Microbiology (Reading) *147*, 183-191. 10.1099/00221287-147-1-183.
61.   Courcelle, J., Khodursky, A., Peter, B., Brown, P.O., and Hanawalt, P.C. (2001).
Comparative gene expression profiles following UV exposure in wild-type and SOS-
deficient *Escherichia coli*. Genetics *158*, 41-64. 10.1093/genetics/158.1.41.

62.   Foti, J.J., Persky, N.S., Ferullo, D.J., and Lovett, S.T. (2007). Chromosome segregation
control by *Escherichia coli* ObgE GTPase. Mol Microbiol *65*, 569-581. 10.1111/j.1365-
2958.2007.05811.x.
63.   Tan, J., Jakob, U., and Bardwell, J.C. (2002). Overexpression of two different GTPases
rescues a null mutation in a heat-induced rRNA methyltransferase. J Bacteriol *184*, 2692-
2698. 10.1128/JB.184.10.2692-2698.2002.
64.   Nirody, J.A., Budin, I., and Rangamani, P. (2020). ATP synthase: Evolution, energetics,
and membrane interactions. J Gen Physiol *152*. 10.1085/jgp.201912475.
65.   Lund, P., Tramonti, A., and De Biase, D. (2014). Coping with low pH: molecular
strategies in neutralophilic bacteria. FEMS Microbiol Rev *38*, 1091-1125. 10.1111/1574-
6976.12076.
66.   Wang, Y., Wu, J., Lv, M., Shao, Z., Hungwe, M., Wang, J., Bai, X., Xie, J., and Geng, W.
(2021). Metabolism Characteristics of Lactic Acid Bacteria and the Expanding
Applications in Food Industry. Front Bioeng Biotechnol *9*, 612285.
10.3389/fbioe.2021.612285.
67.   Ajdic, D., McShan, W.M., McLaughlin, R.E., Savic, G., Chang, J., Carson, M.B.,
Primeaux, C., Tian, R., Kenton, S., Jia, H., et al. (2002). Genome sequence of
Streptococcus mutans UA159, a cariogenic dental pathogen. Proc Natl Acad Sci U S A
99, 14434-14439. 10.1073/pnas.172501299.
68.   Peters, J.M., Colavin, A., Shi, H., Czarny, T.L., Larson, M.H., Wong, S., Hawkins, J.S.,
Lu, C.H.S., Koo, B.M., Marta, E., et al. (2016). A Comprehensive, CRISPR-based
Functional Analysis of Essential Genes in Bacteria. Cell *165*, 1493-1506.
10.1016/j.cell.2016.05.003.
69.   Shields, R.C., Walker, A.R., Maricic, N., Chakraborty, B., Underhill, S.A.M., and Burne,
R.A. (2020). Repurposing the *Streptococcus mutans* CRISPR-Cas9 System to Understand
Essential Gene Function. PLoS Pathog *16*, e1008344. 10.1371/journal.ppat.1008344.
70.   Yother, J., Trieu-Cuot, P., Klaenhammer, T.R., and De Vos, W.M. (2002). Genetics of
streptococci, lactococci, and enterococci: review of the sixth international conference. J
Bacteriol *184*, 6085-6092. 10.1128/JB.184.22.6085-6092.2002.
71.   Otasek, D., Morris, J.H., Bouças, J., Pico, A.R., and Demchak, B. (2019). Cytoscape
Automation: empowering workflow-based network analysis. Genome Biol *20*, 185.
10.1186/s13059-019-1758-4.

1031
1032   **Figures and legends**
1033
1034   **Figure 1. Evolution of mutants deleted for essential genes in passage experiments.**
1035
(A) Schematic representation of suppressor evolution when a mutant deleted for an essential
gene is passaged multiple times. Shown is a replacement of a bacterial essential gene with the
kan gene by homologous recombination through transformation. Secondary mutations conferring
a fitness advantage become enriched in the population, and subsequent tertiary or quaternary
mutations build upon the previously fixed mutations. Bold circles indicate cells, and colors of the
bold circle indicate the development of cell lineages. Other symbols are defined in the figure.
1042
(B) Two types of colonies, small and large, as demonstrated for *f1fo* deletion, appear on selection
plates after 5 days of growth. Only the small colonies represent true deletions, denoted as Δ*f1fo*

(red circle). The large colonies are "double-band mutants," containing both the replacement of
the original *f1fo* with a *kan* gene and a wild-type *f1fo* copy, or they are mutants that did not
contain a *kan* gene, but contained point mutations in other genes, denoted as Δ*f1fo-DB* (black
circle, 'DB' for double-band).
1049
(C) Schematic representation of passage experiments. Illustrates the evolution of suppressors and
the fixation of mutated segments in a population deleted of an essential gene over multiple
passages. Yellow arrows indicate passaging process. The inverted trapezoids indicate the tubes
containing media and cells. The size of trapezoids indicates volume of cell cultures, where the
small trapezoids indicate 1.2 mL culture and big trapezoids indicate 2.5 mL culture. The bold
circles indicate cells. Blue arrows in P0 and P1 indicate the changes in cell lineages over passage.
1056
1057   (D) Incubation time (days) required for P0 and P6 mutants to reach an OD_{600} of 0.1 to 0.5.
1058
1059   (E and F) Growth curves display the growth of five independently evolved Δ*f1fo* colonies in P0
1060   (E) and P6 (F) in comparison to that of WT and four Δ*f1fo-DB* colonies. The inserted red
1061   numbers represent the means of OD_{600} values after 48 hours of growth under anaerobic
1062   conditions.
1063
1064
1065   **Figure 2. Identification of suppressors in mutants deleted for essential genes.**
1066
1067   (A) Number of independently evolved populations of different mutants.
1068
1069   (B-I) Evolved populations containing large-scale duplications. Red arrow, site of original
1070   essential gene deletion; Y axis indicates number of sequence reads mapped to the
1071   reference sequence at the coordinates shown on X axis.
1072
1073   (J) Boxplot of mutated segments (≥30% abundance) in the independently evolved populations of
1074   mutants deleted for essential genes or WT. Open circles indicate outliers.
1075
1076   (K) Stacked proportions of populations containing mutated segments comprising >50% (blue) or
1077   30%-50% (red) of sequence reads.
1078
1079   (L) Stacked proportions of mutated segments belonging to intergenic regions, non-essential
1080   ORFs, or essential ORFs.
1081
1082   (M) Stacked proportions of evolved populations with/without mutations in the multi-drug
1083   resistance genes (DRGs: *rpsL*, *gyrB*, *rpoB* and *rrs2*) shown.
1084
1085
1086   **Figure 3. Networks of suppressors and essential gene mutants.**
1087
A network was built from the lists of suppressor mutants derived from each essential gene
mutant and WT. Yellow-highlighted labels indicate the names of the essential genes mutated and
the blue-highlighted label indicates the passaged WT. In total, there are twenty-five groups. Gene

names without any marks, mutated segments belonging to essential genes; gene names with an
asterisk (*) on the left, conditional essential genes; gene name with two asterisks on the left, non-
essential genes; P_ followed by gene names and an asterisk, promoters; In_ followed by gene
names and an asterisk and int_ followed by number, intergenic regions; _ followed by number,
gene locus.
1096
1097
1098   **Figure 4. The most frequent suppressors identified in mutants deleted of essential genes.**
1099   The percentage of mutated segments in evolved populations in *Δf1fo*, mutants deleted of single
1100   *unc* subunit genes, denoted as *Δf1fo.s*, the other single mutants indicated, and WT. The names
1101   with an asterisk (*) indicate the intergenic regions.
1102
1103
1104   **Figure 5. Increasing *v1vo* copy number and inhibition of *trkA1-H1* independently**
1105   **contribute to the fitness improvement of evolved *Δf1fo* populations.**
1106
1107   (A) Gene duplication of *v1vo* region does not occur in P1 population (upper) and appeared in P6
1108   (lower) population of *Δf1fo#109.* Yellow arrows indicate passaging process. The black box
1109   indicates the *f1fo* region
1110
1111   (B) Population of *Δf1fo#4* at P6 that does not contain gene duplication of *v1vo* region (upper).
1112   *v1vo* region duplication was observed in P15 (lower). (Two out of sixed evolved populations; see
1113   S7). Arrows indicate passaging process.
1114
1115   (C) *v1vo* region deleted in WT (upper). Duplication of *v1vo* region *Δf1fo#4, denoted as v1vo-DB/*
1116   *Δf1fo#4.* The black box indicates the *f1fo* region. The red box indicates *v1vo* region.
1117
1118   (D) Mutation frequency in *trkH1* increased with passage in P6 compared to P1 in population
1119   *Δf1f#104.*
1120
1121   (E) Growth of strains indicated. 0.1 OD_{600} of cells in a volume of 2 µl were spotted directly (first
1122   column) or diluted 20-fold (column 2), or 400-fold (column 3) and grown on BHI-agar for two
1123   days.
1124
1125   (F) Growth of the strains indicated. 0.1 OD_{600} of cells in a volume of 2 µl were spotted directly
1126   (first column) or diluted 10-fold (column 2), or 100-fold (column 3), or 1000-fold (column 4),
1127   and grown for four days.
1128
1129   (G) Quantitative growth measurements of WT, *ΔtrkA1-H1, ΔtrkA2-H2, Δf1fo* in *ΔtrkA1-H1* and
1130   *Δf1fo* in *ΔtrkA2-H2.* Culture is in BHI for 24 hr in anaerobic conditions. Data are shown as
1131   means ± SD from three replicate cultures. Different letters indicate statistically significant
1132   differences (P≤0.05), determined by one -way ANOVA with Tukey multiple comparisons test.
1133
1134   (H) Sensitivity of *Δf1fo#4* and WT to low pH (lactic acid: LA). The values are relative OD_{600} to
1135   that of WT grown in plain BHI. X-axis denotes the vol (µL) of 90% LA in 25 mL of BHI. Data
1136   are shown as means ± SD from three replicate cultures.

1137
1138 (I) Sensitivity of *Δf1fo#4* and WT to high pH (NaOH). The values are relative OD_{600} to that of
1139 WT grown in plain BHI. X-axis denotes the vol (μL) of 5 M NaOH in 25 mL of BHI. Data are
1140 shown as means ± SD from three replicate cultures.
1141
1142
1143 **Figure 6. Working model for F1Fo, V1Vo, TrkA1-H1 and TrkA2-H2.**
1144
For streptococci, which do not contain a complete tricarboxylic acid cycle (TCA: shaded) or
electron transport chain (ETC: shaded), energy production is mainly through lactic acid
fermentation. The lactate from fermentation creates acidic conditions in the cytosol, which
becomes toxic if H+ is not exported. F1Fo pumps out H+ from the cytosol using ATP, and V1Vo
supplements or replaces the function of F1Fo when the F1Fo is inhibited/inactivated. Both
TrkA1-H1 and TrkA2-H1 are potassium transporters, and functionally redundant. TrkA1-H1
pumps in potassium together with H+, acidifying the cytosol when H+ is not exported. In
contrast, TrkA2-H2 does not pump protons into the cytosol when pumping in potassium (likely
instead pumping in Na+). Therefore, the function of TrkA1-H1 must be inhibited if F1Fo is not
functional. Solid lines indicate presence of reactions. Dashed lines indicate absence of reactions.
1155
1156
1157 **Figure 7 Essentiality of *f1fo* in *S. mutans* UA159, *S. sanguinis* SK36, SK405 and SK1058.**
1158
1159 (A) Distribution of the genes indicated in *S. mutans* UA159, *S. sanguinis* SK36, SK405 and
1160 SK1058.
1161
1162 (B) Growth of *Δf1fo* transformants in *S. sanguinis* SK405 (upper) and SK1058 (lower) on same
1163 BHI agar medium for five days. Enlarged images of the transformants are shown. Red circles
1164 indicate small colonies, *Δf1fo*. Black circles indicate big colonies, *Δf1fo-DB*.
1165
1166 (C) Deletion of *f1fo* in SK405 or SK1058 shown by whole genome sequencing. Black box
1167 indicates the location of *f1fo* region.
1168
1169 (D) Quantitative measurements of growth in wild-type SK36, SK405 or SK1058, *ΔtrkA1-H1* or
1170 *Δf1fo* mutants in the backgrounds of SK36, SK405 or SK1058. Cells were grown in BHI for 24
1171 hours. Data are shown as means ± SD from three replicate cultures. Different letters indicate
1172 statistically significant differences (P≤0.05), determined by one -way ANOVA with Tukey
1173 multiple comparisons test.
1174
1175 (E) Growth of SK36, SK405 or SK1058, *ΔtrkA1-H1* mutant in the backgrounds of SK36, SK405
1176 or SK1058. 0.1 OD_{600} of cells in a volume of 2 μl were spotted directly (first column) or diluted
1177 10-fold (column 2), or 100-fold (column 3), or 1000-fold (column 4). Growth of Cells were
1178 grown on BHI-agar for two days.
1179
1180 (F) Growth of SK36, SK405 or SK1058, *Δf1fo* mutant in the backgrounds of SK36, SK405 or
1181 SK1058. Cells were diluted as in Fig. E and grown on BHI-agar for four days.
1182

1183 (G) Location of *trkA1-H1* suppressor mutations in SK405 and SK1058. Mutation of *trkA1-H1*
1184 appeared in all evolved *Δf1fo* populations, three in SK405 and two in SK1058. X-axis indicates
1185 the genome coordinates.
1186
1187
1188 **Supplemental Figures, Tables and legends**
1189
1190
1191 **Fig S1. Genotyping and complementation of mutants deleted of *f1fo*.**
1192 (A) Gene knockout strategy to delete entire region of eight *f1fo* subunits by homologous
1193 recombination. The yellow blocks indicate the eight F1Fo subunit genes in WT (upper). The blue
1194 block indicates the *kan* gene in Δ*f1fo* mutants (lower).
1195
1196 (B) Growth of transformants for five days in selection medium. Red circle indicates small
1197 colonies, Δ*f1fo*. Black circle indicates big colonies, Δ*f1fo-DB* ('DB' for double-band). This image
1198 is a duplicate of Figure 1B as a demonstration of the transformants in selection medium.
1199
1200 (C) Whole-genome sequencing to confirm the genotypes of the small colonies for Δ*f1fo* mutants.
1201 Black box indicates the location of deleted *f1fo* region.
1202
1203 (D) Whole-genome sequencing of the big colonies, Δ*f1fo*-DB. Black box indicates the location
1204 of *f1fo* region. Red arrows indicate the two directly repeated *ugpC* genes, encoding glycerol-3-
1205 phosphate ABC transporter.
1206
1207
1208 **Fig S2. Isolation of viable mutants deleted essential genes and genotyping by whole-genome**
1209 **sequencing.**
1210
(A-B) Colonies of transformants (left) and whole-genome sequencing of colonies (right). Red
circles indicate the colonies picked for whole-genome sequencing (left). For genome sequencing
(right), the blue graphic above the sequence coordinates (indicated by a red arrow in the top
panel of A.1) indicates the extent of coverage, the yellow bars indicate the mismatches and the
bracket indicates the aligned sequence reads. Bacterial colonies on selective agar plate for five
days. (A.1-A.2) Gene deletion mutants with severe growth defects. (B) Gene deletion mutants
with robust growth. Scale bar: 1 mm. The image of Δ*f1fo* transformation is a duplicate of Figure
1B to demonstrate the transformants in selection medium.
1219
1220
1221 **Fig S3. Growth of mutants deleted of essential genes, relative to that of WT.**
1222
1223 (A-B) Growth mutants deleted of 23 essential genes under anaerobic condition (A) and
1224 microaerobic condition (B). 0.1 OD_{600} of cells in a volume of 2 µl were spotted directly (first
1225 column) or diluted 20-fold (column 2), or 400-fold (column 3) and grown on BHI-agar for two
1226 days. Scale bar is 1 cm.
1227

1228 (C) Comparison of the growth of the twenty-four evolved mutants and that of WT. The values
1229 are relative OD_{600} after 24 hr of growth in BHI liquid medium for 24 hours.
1230
1231
1232 **Fig S4. Large scale gene duplication present in the evolved populations of the mutants**
1233 **deleted essential genes but not in WT populations.**
1234
1235 (A) Whole-genome sequencing of parental WT and the six evolved WT populations after nine
1236 passages. The yellow arrow indicates the passage procedure.
1237
1238 (B-E) Evolved populations that contain gene duplication of greater than 10 kb in Δ*uncE#3* (B),
1239 Δ*f1fo#107* (C), Δ*fabH#4* (D) *and* Δ*fabH#6* (E). Red arrows (B-E) indicate the location of the
1240 gene intentionally deleted. The insert image (C) is an enlarged segments of *uncE* region, shown
1241 as a black box.
1242
1243
1244 **Fig S5. Appearance of mutations in essential genes in the evolved populations of strains**
1245 **intentionally deleted for different essential genes.**
1246
1247 (A-F) Aligned sequence reads from Δ*uncC#1* of P1 population (A-C) and P6 population (D-F).
1248 Red arrows indicate the deleted *uncC* in the genome (A and D) in the Δ*uncC#1* strain. Red box
1249 indicates mutation of *uncH* in Δ*uncC#1* (A and D). Enlarged mutated *uncH* region of aligned
1250 sequence reads in Δ*uncC#1* of P1 population (B-C) and P6 population (E-F). The yellow arrow
1251 indicates the passage procedure.
1252
1253 (G) Sequencing of the mutated *uncH* gene in Δ*uncC#1* in P6. Red letters indicate a fragment (82
1254 bp) that was present in mutated *uncH* gene (upper). Asterisk (*) indicates the stop codon that was
1255 highlighted. Vertical bars indicate the coordinates of the *uncH* ORF (lower). The coordinates
1256 between two red bars indicate replaced region.
1257
1258 (H-K) Aligned sequence reads from Δ*f1fo#107* of P1 population (H-I) and P6 population (J-K).
1259 Red arrows indicate the deleted *f1fo* in the genome (H and J) in the Δ*f1fo#107*. Red box indicates
1260 mutation of *ftsZ* in Δ*f1fo#107* (H and J). Enlarged mutated *ftsZ* region of aligned sequence reads
1261 in Δ*f1fo#107* of P1 population (H) and P6 population (I). The yellow arrow indicates the passage
1262 procedure. Fig S5J is a duplicate of Fig S4B to show a genome alignment of Δ*f1fo#107* of P6.
1263
1264 (L) Sequencing of the mutated *ftsZ* in Δ*f1fo#107* in P6. Red letters indicate a fragment (248 bp)
1265 that was present in mutated Δ*f1fo#107* (upper). Asterisk (*) indicates the stop codon that was
1266 highlighted. Vertical bars indicated the coordinates of the *uncH* ORF (lower). The red bar
1267 indicates the coordinate where insertion occurred.
1268
1269
1270 **Fig S6. Populations with and without mutations in three PTS mannose/fructose/sorbose**
1271 **transporter subunits.**
1272

1273 (A-B) Number (A) and percentage (B) of populations that have mutations in any of three PTS
1274 mannose/fructose/sorbose transporter subunits, namely J1C87_09355, J1C87_09360, and
1275 J1C87_09365.
1276
1277
1278 **Fig S7. Appearance of *v1vo* region gene duplication during passage. (Related to Figure 5A**
1279 **and 5B)**
1280
1281 (A-B) Alignment of sequence reads in populations of *Δf1fo#107, Δf1fo#109, uncE#2* and *uncH#7*
1282 at P1 (A) and P6 (B). The black box indicates the deleted *f1fo* region and the red arrows indicate
1283 the deleted *uncE* or *uncH*.
1284
1285 (C) Alignment of sequence reads of the P6 population in *Δf1fo#4* (upper) and five derived
1286 populations from P15 (lower). Rectangles represent the deleted *f1fo* region. Asterisks (*) indicate
1287 the two populations with *v1vo* region gene duplications.
1288
1289
1290 **Fig S8. Gene duplication in *Δv1vo-DB/Δf1fo#4* populations (related to Figure 5C).**
1291
1292 Alignment of sequence reads of five additional *Δv1vo-DB/Δf1fo#4* populations. The black box
1293 indicates the deleted *f1fo*. The red box indicates the gene duplication of *v1vo* region when one
1294 copy of *v1vo* was deleted.
1295
1296
1297 **Fig S9. Alignment of sequence reads of *trk* mutants with and without *f1fo* deletion**.
1298
1299 (A-K) Alignment of sequence reads of *ΔtrkA1* (A), *ΔtrkH1* (B), *ΔtrkA1-H1* (C), *ΔtrkA2* (D),
1300 *ΔtrkH2* (E), *ΔtrkA2-H2* (F), *ΔtrkA1/ΔtrkH2* (G), *ΔtrkH1/ΔtrkH2* (H), *ΔtrkA1-H1/ΔtrkH2* (I),
1301 *Δf1fo/ΔtrkA1-H1* (J), and *Δf1fo/ΔtrkA2-H2* (K). Red arrows indicated intended gene deletion.
1302 The block box indicates the deleted *f1fo region*. (related to Figure 5F).
1303
1304
1305 **Fig S10. *f1fo* cannot be deleted in *S. mutans* UA159.**
1306
1307 (A) Gene knockout strategy to delete entire region of eight *f1fo* subunits by homologous
1308 recombination. The purple blocks indicate the eight F1Fo subunit genes in WT (upper). The blue
1309 block indicates the *kan* gene in Δ*f1fo* mutants (lower). The two arrows indicate the primer set for
1310 the PCR amplicon of the region in (B).
1311
1312 (B) Genotyping the presence of wild-type *f1fo* colonies from selection medium. P: positive
1313 control using DNA template from wild-type. N: negative control using water as a template. The
1314 size of amplicons is about 1300 bp.
1315
1316
1317 **Fig S11. Deletion *of trkA1-H1* in *S. sanguinis* SK1058 and S. sanguinis SK405 (related to**
1318 **Figure 7C).**

1319
1320   (A-B) Alignment of sequence reads of *ΔtrkA1-H1* populations in SK405 (A) and in SK1058 (B)
1321   backgrounds.
1322
1323
1324   **Fig S12. Deletion *of f1fo* in *S. sanguinis* SK1058 and SK405 (related to Figure 7C).**
1325   Alignment of sequence reads of three *Δf1fo* populations in SK405 backgrounds (A) and two in
1326   SK1058 backgrounds (B) (related to Figure 7C). Rectangles represent the deleted *f1fo* region.

Fig.1

A

Transformation   Homologous recombination   Development of cell lineage during serial passages

Knockout construct
Chromosome
Mutations on chromosome
Passage

B

Δf1fo
Δf1fo-DB
1 mm

C

Days required to reach OD$_{600}$ of 0.1 to 0.5

ΔfabF ΔfabZ ΔfabK ΔfabG ΔpfkA ΔpmgM ΔobgE Δf1fo ΔuncE ΔuncB ΔuncF ΔuncH ΔuncA ΔuncG ΔuncD ΔuncC ΔyqeG ΔdnaK ΔfabH ΔrpmH ΔrplV ΔyihA ΔdpcK ΔftsX

■ P0   ■ P6

D

OD$_{600}$

Δf1fo#101
Δf1fo#102
Δf1fo#103
Δf1fo#104
Δf1fo#105
Δf1fo-DB#1
Δf1fo-DB#2
Δf1fo-DB#3
Δf1fo-DB#4
WT.1
WT.2
WT.3

1 mm

0.162

hours

E

OD$_{600}$

Δf1fo#101
Δf1fo#102
Δf1fo#103
Δf1fo#104
Δf1fo#105
Δf1fo-DB#1
Δf1fo-DB#2
Δf1fo-DB#3
Δf1fo-DB#4
WT.1
WT.2
WT.3

1 mm

0.469

hours

Fig.2

Fig.3

Fig.4

Fig.5

A P1 Δf1fo#109
P6

B P6 Δf1fo#4
P15

C Δv1vo#10
Δv1vo-C/Δf1fo#4

D **Mutation frequency of *trkH1***

● *substitution*   - - - - *30%*
▲ *frameshift*   ······· *100%*

P1 Δf1fo#104

P6

1   100   200   300   400   479
(aa)

E
WT
ΔtrkA1-H1
ΔtrkA2-H2
ΔtrkH2
ΔtrkA1-H1/ΔtrkH2
ΔtrkA1/ΔtrkH2
ΔtrkH1/ΔtrkH2
Δf1fo#4
1 cm

F
WT
ΔtrkA1-H1
ΔtrkA2-H2
Δf1fo/ΔtrkA1-H1
Δf1fo/ΔtrkA2-H2
Δf1fo#4

G
OD 600
WT, ΔTrkA1-H1, ΔTrkA2-H2, Δf1fo/ΔTrkA1-H1, Δf1fo/ΔTrkA2-H2
a a a b c

H
Relative sensitivity to lactic acid
■ *WT* ■ *f1fo#4*
0 10 20 30 40
μL

I
Relative sensitivity to NaOH
■ *WT* ■ *f1fo#4*
0 25 50 100 200
μL

Fig.6   Working model for F1Fo, V1Vo, TrkA1-H1 and TrkA2-H2

Fig.7