Genetic variation in the species *Arabidopsis thaliana* reveals the existence of natural heat resilience factors for meiosis

Jiayi Zhao[1], Huiqi Fu[1], Zhengze Wang[2], Min Zhang[1], Yaoqiong Liang[1], Xueying Cui[1], Wenjing Pan[1], Ziming Ren[3], Zhihua Wu[4], Yujie Zhang[1], Xin Gui[1], Li Huo[1], Xiaoning Lei[5], Chong Wang[6], Arp Schnittger[7], Wojciech P. Pawlowski[8], Bing Liu[1,✉]

[1]Arameiosis Lab, South-Central Minzu University, Wuhan 430074;

[2]College of Life Science and Technology, Huazhong Agricultural University, Wuhan 430070;

[3]Department of Landscape Architecture, Zhejiang Sci-Tech University, Hangzhou 310018;

[4]College of Life Sciences, Zhejiang Normal University, Jinhua 321004;

[5]School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai 200025;

[6]Shanghai Key Laboratory of Plant Molecular Sciences, Development Center of Plant Germplasm Resources, College of Life Sciences, Shanghai Normal University, Shanghai 200234;

[7]Department of Developmental Biology, University of Hamburg, Hamburg 22609;

[8]School of Integrative Plant Science, Cornell University, Ithaca, NY 14853;

[✉]To whom correspondence should be addressed: B.L. (arameiosis@163.com).

ORCID

Arp Schnittger: 0000-0001-7067-0091

Wojciech P. Pawlowski: 0000-0003-0770-2083

Bing Liu: 0000-0002-8547-0970

Short title: Natural genetic variation controls heat tolerance of meiosis in Arabidopsis

One-sentence summary: *Arabidopsis thaliana* reveals accession-specific heat tolerance of meiosis that is controlled by multiple genomic loci derived from natural variations.

Total wordcount (Introduction, Materials and Methods, Results and Discussion): 5736

Discussion section: 1122 (19.6% of the total wordcount)

Numbers of figures: 9 (all in color)

Summary

- Heat interferes with multiple meiotic processes leading to genome instability and sterility in flowering plants, including many crops. Despite its importance for food security, the mechanisms underlying heat tolerance of meiosis are poorly understood.

- We analyzed different meiotic processes in the Arabidopsis (*Arabidopsis thaliana*) accessions Columbia (Col) and Landsberg *erecta* (L*er*), their F1 hybrids and F2 offspring under heat stress (37°C).

- At 37°C, Col exhibits significantly reduced formation of double-stand breaks (DSBs) and completely abolished homolog pairing, synapsis and crossover (CO) formation. Strikingly, L*er* and L*er*/Col hybrids are much less affected than Col. Interestingly, only 10% ~ 20% of F2 offspring exhibit the same heat tolerance of meiotic recombination as parents, indicating that heat resilience in L*er* is controlled by the interplay of several loci.

- This study reveals the existence of natural heat resilience factors for meiosis in Arabidopsis, which have the great potential to be exploited in breeding programs.

Keywords

Meiosis, heat stress, meiotic recombination, homolog pairing, crossover, genetic variation, *Arabidopsis thaliana*

Introduction

Meiosis is a specialized type of cell division, which, in plants, generates spores with halved chromosome number required for sexual reproduction and maintenance of ploidy consistency over generations. During prophase I, homologous chromosomes (homologs) undergo recombination, resulting in reciprocal exchange of DNA fragments and thereby promoting genetic diversity. Meiotic recombination is initiated by the programmed formation of DNA double-strand breaks (DSBs) catalyzed by the conserved topoisomerase VI-like protein SPO11 together with other components of the DSB formation machinery (Vrielynck *et al.*, 2021; Nore *et al.*, 2022). The recombinases RAD51 and DMC1 catalyze DSB repair via homologous recombination (Kurzbauer *et al.*, 2012; Da Ines *et al.*, 2013; Chen, H *et al.*, 2021). In plants, deficiencies in DSB repair result in impaired crossover (CO) formation and/or cause chromosome fragmentation (Kurzbauer *et al.*, 2012; Da Ines *et al.*, 2013; Su *et al.*, 2017; Chen, H *et al.*, 2021). Although a large

number of DSBs is generated along chromosomes, only a small proportion of them are repaired into COs. In Arabidopsis as in many other species, most COs are catalyzed by the ZMM proteins resulting in type-I COs that are placed with a distance to each other (CO interference) (Ziolkowski, 2023). The formation of the remaining approximately 15% of COs in plants involves structure-selective endonucleases such as MUS81 giving rise to type-II COs, which are typically not subject to CO interference (Wang & Copenhaver, 2018). The process of COs formation can be visualized by the linking of homologs giving rise to bivalents connected by chiasmata. In the wild-type, all homologs become eventually connected through at least one CO (CO assurance) (Jones & Franklin, 2006), which is required for equal distribution of homologs after meiosis I.

During meiotic recombination, chromatids are organized into loops, the basal regions of which are anchored to an axis through the interactions between axis core proteins (e.g. ASY3 and ASY4 in Arabidopsis) and the axis-associated HORMA domain protein ASY1 (Armstrong *et al.*, 2002; Ferdous *et al.*, 2012; Chambon *et al.*, 2018; West *et al.*, 2019). The assembly of the chromosome axis is required for normal DSB formation and repair, homolog pairing and synapsis, and recombination (Ferdous *et al.*, 2012; Lambing, Christophe *et al.*, 2020; Lambing, C. *et al.*, 2020; Yang *et al.*, 2022).

The separation of homologs and sister chromatids is facilitated by spindles. Cohesin, a ring-shaped protein complex, assures the protection of the sister-chromatids union. The cohesion on chromosome arms is released during prophase and particularly during anaphase I allowing the separation of homologs (Yang *et al.*, 2019). The centromeric cohesion is removed in anaphase II paving the road for the segregation of sister chromatids (Onn *et al.*, 2008). The function and stability of centromeres are crucial for maintenance of cohesion and thus faithful chromosome segregation (McKinley & Cheeseman, 2016; Chen, Y. *et al.*, 2021).

Several studies have shown that meiosis in plants is sensitive to temperature variations (Bomblies *et al.*, 2015; Modliszewski & Copenhaver, 2017; Liu *et al.*, 2019; De Jaeger-Braet *et al.*, 2022; De Jaeger-Braet & Schnittger, 2024). Either elevated or reduced temperature can interfere with cytoskeleton organization and/or meiotic cell cycle transition resulting in defective cytokinesis and unreduced spores (Tang *et al.*, 2011; De Storme *et al.*, 2012; Liu *et al.*, 2018; De Storme & Geelen, 2020; Zhou *et al.*, 2022). In Arabidopsis, mildly decreased or increased temperature within the spectrum that does not reduce fertility (8-28°C) promotes type-I CO formation (Lloyd *et al.*, 2018; Modliszewski *et al.*, 2018). Higher temperatures (30-34°C), however, attenuate homolog synapsis and alter CO rate and distribution, affecting meiosis progression (De Storme & Geelen, 2020; De Jaeger-Braet *et al.*, 2022). In addition, increased

temperatures perturb stability of centromeres and induce unbalanced chromosome segregation in Arabidopsis

(Ahmadli *et al.*, 2023; Jin *et al.*, 2023; Crhak Khaitova *et al.*, 2024). At extremely high temperatures (36-38°C), homolog

pairing and synapsis and CO formation in the Arabidopsis accession Columbia (Col) is completely abolished, which

results in disrupted chromosome segregation and sterility (Lei *et al.*, 2020; Ning *et al.*, 2021; Fu *et al.*, 2022). Despite

the significance of heat stress on genetic makeup and genome stability in plants, the molecular mechanisms controlling

heat tolerance of meiosis remain poorly understood.

Interestingly, previous studies have shown that there are naturally existing genetic differences with respect to meiotic

recombination and stress tolerance in plants (Schulz *et al.*, 2015; Sidhu *et al.*, 2015; Ziolkowski *et al.*, 2017; Taghavi

Namin *et al.*, 2018; Lardon *et al.*, 2020; Chen, Ying *et al.*, 2021; Zhu *et al.*, 2021). Here, we report that the *Arabidopsis*

thaliana accessions Col and Landsberg *erecta* (L*er*) have striking differences in heat tolerance of meiotic recombination.

Our data reveal that meiotic chromosome stability in Col/L*er* F2 progeny is more sensitive to heat than in either inbreds

or their hybrids indicating that several loci contribute to the natural heat resistance.

Material and Methods

Plant materials and growth conditions

Arabidopsis thaliana (L.) accessions Columbia (Col-0) and Landsberg *erecta* (L*er*), the *atm-2* (Kurzbauer *et al.*, 2021),

atm-5 (Yao *et al.*, 2020), *er-105*, *mpk6*, *er-105 mpk6* (Cai *et al.*, 2017), *dmc1* (Couteau et al., 1999), *spo11-1-3* (Xue *et*

al., 2018), *msh4* (Higgins *et al.*, 2004) and *zyp1a/b* (Yang *et al.*, 2022) mutants were used in this study. Col/L*er* hybrids

were obtained by crossing using L*er* as pollen donor. The bi-allelic mutant $ATM^{atm-2/atm-5}$ was obtained by crossing the

heterozygous *atm-2* and *atm-5* mutants. Genotyping was performed using the primers listed in Table S1. Arabidopsis

were cultivated in a growth chamber with a 16 h day/8 h night and 20°C condition. For heat treatment, young flowering

plants were treated by 37°C for 24 h in a humid chamber with a 16 h day/8 h night condition. To avoid potential impact

of circadian clock, heat treatment was applied starting from 8:00-10:00 am.

Cytological analysis of meiocytes

After heat treatment, plants were transferred back to 20°C for 6-12 h recovery, and tetrad stage meiocytes were

analyzed (Lei *et al.*, 2020). A drop of 4.5% (w/v) lactopropionic orcein staining buffer was added to the surface of a tetrad-staged flower bud (floral bud stage 9, 0.3-0.4 mm) (Armstrong & Jones, 2003) followed by squashing to release meiocytes.

Preparation of chromosome spreads

Inflorescences were isolated and fixed in cold Carnoy's fixative for at least 24 h. Flower buds at stages beyond tetrads were removed, and the remaining buds were washed twice in distilled water and once in citrate buffer (10 mM, pH = 4.5), followed by incubation in a digestion enzyme mixture (0.3% pectolyase and 0.3% cellulase in citrate buffer) at 37°C for 3 h. Digested flower buds were subsequently washed once in distilled water, which thereafter were macerated in distilled water on a glass slide. Two rounds of 60% acetic acid were added to the slide, which was dried on a hotplate at 45°C. The slide was flooded with cold Carnoy's fixative and then was air dried. 4',6-diamidino-2-phenylindole (DAPI) was diluted to 5 μg/mL in Vectashield antifade mounting media (Vector Laboratories).

Immunolocalization of α-tubulin and meiotic recombination proteins

To perform immunolocalization of α-tubulin, inflorescences were fixed by 4% (w/v) paraformaldehyde under vacuum for 30 min, followed by twice washes in distilled water and once in citrate buffer (10 mM, pH = 6.0). Meiosis-staged flower buds were digested in an enzyme mixture consisting of 0.3% (w/v) cellulase (Sigma), 0.3% (w/v) pectolyase (Sigma) and 0.3% (w/v) cytohelicase (Sigma) in a humid chamber at 37°C for 3 h. After the digestion, anthers were dissected, squashed, and fixed on a slide by freezing in liquid nitrogen. Released cells on the slide were immobilized with a thin layer of gel containing 1% (w/v) gelatin, 1% (w/v) agarose and 2.5% (w/v) sucrose. After rinsing with phosphate buffered saline with 1% (v/v) Triton-X 100 for 30 min, the α-tubulin antibody diluted in phosphate buffered saline with 0.1% (v/v) Triton X-100 and 4.5 g/L bovine serum albumin was added and the slide was incubated overnight at 4°C. The slide washed three times with phosphate buffered saline with 0.02% tween-20 was incubated with a secondary antibody at 37°C for 1 h in the dark. Immunolocalization of meiotic recombination proteins was performed as previously described (Chelysheva *et al.*, 2010; Wang *et al.*, 2014), and the antibodies used in this study have been reported previously (Lei *et al.*, 2020; Ning *et al.*, 2021; Zhao *et al.*, 2023).

Quantification of fluorescent foci, fluorescence intensity and chromosome fragments

Fluorescent images were taken by an Olympus IX83 inverted fluorescence microscope with a X-Cite lamp and a Prime

BSI camera, and Z-stacks were processed using ImageJ. The fluorescent foci merged onto DAPI-stained chromosomes

were counted. For the pixel intensity plot, the pixel brightness through a region of interest was measured using ImageJ

and plotted against the X dimension. To score the number of chromosome fragments, chromosome bodies in

meiocytes at metaphase I, anaphase I and metaphase II that were obviously brighter than organelles were counted

followed by a subtraction of expected chromosome number (5 at metaphase I, 10 at anaphase I or metaphase II).

Statistical analysis

Significance was calculated using unpaired *t*-tests (Fig. 1b; Fig. 2c, d, h and j; Fig. 4b, d, and h; Fig. 6d and e; Fig. 7d and

g; Fig. 9b and d) or Chi-squared tests (Fig. 1f; Fig. 2f; Fig. 3j, l and t; Fig. 4f, j, l and n; Fig. 7c; Fig. 8c and h; Fig. 9g, h, k

and l) with the software GraphPad Prism. The significance level was set as $P < 0.05$. To calculate the frequency of

individuals or meiocytes showing either altered chromosome morphology in interkinesis (Fig. 6c, d and e), untimely

sister-chromatids segregation, or number of chromosome fragments (Fig. 7c, d and g), the individuals, in which a

meiocyte with the corresponding defect was observed, and at least three meiocytes at the corresponding stage were

found, were used for calculation. The numbers of analyzed meiocytes or individuals are shown in the figures or main

text. Correlation between different meiosis phenotypes was determined based on Spearman's rank correlation tests.

Coefficients were calculated using Corrplot package on the R statistical platform, and Ggpubr package was used for

visualization.

Results

Landsberg *erecta* exhibits higher heat tolerance of meiosis than Columbia

The *Arabidopsis thaliana* accessions Columbia (Col) and Landsberg *erecta* (L*er*), two frequently used varieties, exhibit

differences in their meiotic recombination rates as well as response to stresses (Schiavon *et al.*, 2007; Lasky *et al.*,

2014; Schulz *et al.*, 2015; Ziolkowski *et al.*, 2017; Choi *et al.*, 2019; Chen, Ying *et al.*, 2021). To test whether heat

tolerance of meiosis also varies, we compared the effect of extreme heat stress, i.e., 37°C, versus the control condition,

i.e., 20°C, between these two accessions. To this end, we analyzed tetrad formation at the end of meiosis by orcein

staining. At 20°C, both accessions generated typical tetrads that harbored four spores with equally-sized nuclei (Fig.

1a and b). Exposure to the extreme temperature resulted in 85.7% tetrads in Col that showed irregular configurations

with abnormal numbers and/or sizes of spores and nuclei (Fig. 1a and b). In striking contrast, only 8.8% abnormal

tetrads were found in L*er* (Fig. 1a and b), suggesting that meiosis in L*er* is more resilient to heat stress than Col.

Heat-induced formation of abnormal tetrads could result from defective meiotic recombination that leads to

univalents and consequently, chromosome mis-segregation. Alternatively, defects in microtubule-mediated

chromosome distribution and/or cytokinesis could be the underlying cause (Lei *et al.*, 2020). To determine which

meiotic process in Col is more affected by heat stress, we analyzed the spindle at metaphase I and II as well as

phragmoplast at telophase II (Liu *et al.*, 2017; Liu *et al.*, 2018; Lei *et al.*, 2020). At 20°C, the typical barrel-shaped

spindles and straight phragmoplasts were observed in both accessions. (Fig. S1, Fig. 1c). At 37°C, we did not observe

obvious defects in microtubule organization in prophase I meiocytes in either accession (Fig. S1) while Col and L*er*

showed similar rates of irregularities in morphology of spindles and phragmoplasts in meiosis I and/or II (Fig. 1d,

normal; e, abnormal; and f) that likely contributed to an unequal segregation of chromosomes (Fig. 1g). The higher

heat tolerance of meiosis in L*er* versus Col is thus likely not caused by the difference in microtubule organization, but

possibly occurs prior to metaphase I, i.e., in meiotic recombination.

Figure 1. L*er* exhibits a higher heat tolerance of meiosis than Col. (a) Representative images of tetrads with normal or abnormal

configurations. (b) Graph showing the frequencies of abnormal tetrads in Col and L*er* at 20°C or 37°C. Stars indicate significance

levels based on unpaired *t* tests. (c and d) Representative images of immunolocalization of α-tubulin in metaphase I, anaphase I,

metaphase II and telophase II meiocytes depicting normal microtubule configurations at 20°C (c) or 37°C (d). (e) Representative

images of immunolocalization of α-tubulin in metaphase I, anaphase I, metaphase II and telophase II meiocytes displaying altered

configurations at 37°C. (f) Graph showing the rates of spindle and phragmoplast with normal or abnormal configurations in Col and L*er* at 20°C or 37°C. Stars indicate significance levels based on χ^2-tests. (g) Representative images of immunolocalization of α-tubulin in tetrads showing normal or abnormal configurations at 37°C. White, DAPI; green, α-tubulin. Frequencies represent the average rates of the corresponding phenotypes; n indicates the number of the analyzed plants or meiocytes; ***, $P < 0.001$; *, $P < 0.05$; ns, $P > 0.05$; scale bars, 10 μm.

DSB and CO formation in L*er* are largely resistant to heat

To assess the possible resilience of meiotic recombination towards heat in L*er*, we analyzed chromosomes by spreading and scored bivalent formation in Col and L*er* at 20°C and 37°C. At 20°C, five bivalents were present in diakinesis in both accessions (Fig. 2a and c). While bivalent formation in Col was completely impaired at 37°C, this was not the case in L*er* (Fig. 2a and c). To examine then the effect of heat on CO formation, immunolocalization of the type-I CO marker protein HEI10 was performed and the number of HEI10 foci on chromosomes in diakinesis was quantified. At 20°C, as previously reported, Col showed a higher number of HEI10 foci than L*er* with 11.0 foci versus 9.4, respectively (Fig. 2b and d) (Ziolkowski *et al.*, 2017). At 37°C, no HEI10 foci were detected in Col whereas L*er* showed an average of 10.5 HEI10 foci per meiocyte, being not significantly different from the number at 20°C (Fig. 2b and d). Moreover, all analyzed anaphase I meiocytes in heat-stressed Col showed mis-segregation of homologs while only 41.5% unequal homolog segregation occurred in L*er* under heat stress (Fig. 2e and f). Based on these data, we conclude that heat stress does not reduce CO formation in L*er*. Since heat stress has earlier been found to partially suppress CO formation by reducing DNA double-strand break (DSB) formation (Ning *et al.*, 2021), we next analyzed the localization of the DSB repair recombinases RAD51 and DMC1, two widely used DSB markers (Kurzbauer *et al.*, 2012; Kurzbauer *et al.*, 2021). Consistent with previous reports, both the numbers of RAD51 and DMC1 foci on zygotene chromosomes in Col were significantly reduced after heat treatment (Fig. 2g-j) (Ning *et al.*, 2021; Zhao *et al.*, 2023). In L*er*, however, the focus numbers of both proteins were not significantly altered by heat (Fig. 2g-j), suggesting that heat stress does not reduce DSB formation in L*er*.

Figure 2. Heat stress does not reduce CO and DSB formation in L*er*. (a) Representative images of diakinesis meiocytes showing five,

four, one or no bivalents. The red and yellow arrows indicate univalents and bivalent, respectively. (b) Immunolocalization of HEI10

in diakinesis meiocytes of Col and L*er* at 20°C and 37°C. (c and d) Graphs showing the average numbers of bivalent (c) and HEI10

foci (d) per meiocyte in Col and L*er* at 20°C and 37°C. Stars indicate significance levels based on unpaired *t* tests. (e) Representative images of anaphase I meiocytes showing balanced and unequal homolog separation. (f) Graph showing the fraction of anaphase I meiocytes with equal or unequal homolog separation in Col and L*er* at 20°C and 37°C. Stars provide significance levels based on χ^2-tests. (g and i) Immunolocalization of RAD51 (g) and DMC1 (i) on zygotene chromosomes of Col and L*er* at 20°C and 37°C. (h and j) Graphs showing the average numbers of RAD51 (h) and DMC1 (j) foci per meiocyte in Col and L*er* at 20°C and 37°C. Stars indicate significance levels based on unpaired *t* tests; frequencies indicate the average number or fraction of the corresponding phenotypes; n indicates the number of the analyzed cells; ***, $P < 0.001$; **, $P < 0.01$; ns, $P > 0.05$; scale bars, 10 μm.

Homolog synapsis in L*er* is more tolerant to heat stress than in Col

Disrupted homolog pairing is the main cause of impaired CO formation in Col under heat stress (Ning *et al.*, 2021). Therefore, we compared heat tolerance of homolog pairing in Col and L*er*. We classified pachytene meiocytes into three types based on juxtaposition status of homologs: full-pairing, which indicates meiocytes exhibiting fully juxtaposed homologs (Fig. 3a, b and e); partial-pairing, which indicates meiocytes showing both juxtaposed and unpaired regions between homologs (Fig. 3f-i); and no pairing that represents impairment of homolog juxtaposition (Fig. 3c and d). As previously reported, homolog pairing and synapsis in Col were fully impaired at 37°C (Fig. 3a, c and j) (Ning *et al.*, 2021). In L*er*, however, 63.1% of meiocytes exhibited full pairing (Fig. 3e and j), and 35.4% of meiocytes showed partial pairing (Fig. 3f-i, red arrows; j), with only 1.5% meiocytes displaying no pairing (Fig. 3d and j). Next, we compared the impact of heat on chromosome morphology and the assembly of the chromosome axis and synaptonemal complex (SC) by analyzing the localization of ASY4 and ZYP1 on zygotene and pachytene chromosomes, respectively. At 20°C, ASY4 were loaded on the whole DAPI-stained chromosome regions showing a linear configuration (Fig. 3k and l). Heat induced patchy ASY4 signals in both Col and L*er* (Fig. 3k), but the fraction of this phenotype in Col was significantly higher than that in L*er* (Fig. 3l, $P < 0.001$). Immunolocalization of ZYP1 showed that the SC assembly in Col was disrupted by heat, which, however, occurred normally or was only partially perturbed in most pachytene meiocytes in L*er* (Fig. 3m-s, and t, ~95.6%). These data indicate that homolog pairing and synapsis in L*er* are less impacted by heat than in Col.

Figure 3. Homolog pairing and synapsis in L*er* are less impacted by heat than those in Col. (a-i) Pachytene chromosomes in Col (a and c) and L*er* (b, d-i) at 20°C (a and b) and 37°C (c-i). The red arrows indicate unpaired chromosome regions. (j) Graph showing the rates of pairing status of pachytene chromosomes in Col and L*er* at 20°C and 37°C. Stars indicate significance levels based on χ^2-tests. (k) Representative images of immunolocalization of ASY4 on chromosomes showing linear or patchy configurations. White, DAPI; green, ASY4. (l) Graph showing the rates of ASY4 configurations in Col and L*er* at 20°C and 37°C. Stars indicate significance levels based on χ^2-tests. (m-s) Immunolocalization of ZYP1 on pachytene chromosomes in Col (m and o) or L*er* (n, p-s) at 20°C (m and n) and 37°C (o-s). White, DAPI; red, ZYP1. (t) Graph showing the rates of ZYP1 configurations on pachytene chromosomes in Col and L*er* at 20°C and 37°C. Stars indicate significance levels based on χ^2-tests; frequencies indicate the rates of the corresponding phenotypes; n indicates the number of the analyzed cells; ***, $P < 0.001$; **, $P < 0.01$; scale bars, 10 μm.

Col/L*er* hybrids exhibit similar heat tolerance of meiotic recombination as L*er*

One of the main developmental differences between Col and L*er* is floral architecture (Fig. S2), which, in Arabidopsis,

is regulated by the receptor-like protein kinase ERECTA (ER) that acts upstream of the mitogen-activated protein kinase

cascade module MKK4/5-MPK3/6 (Meng *et al.*, 2012; Cai *et al.*, 2017). In addition, ER potentially plays a role in varied

heat responses between Col and L*er* (Zhang *et al.*, 2017; Lyu *et al.*, 2020). We thus examined whether the distinct heat

tolerance of meiotic recombination between Col and L*er* is owing to a loss of *ER*. To this end, we analyzed bivalent

formation, homolog separation, DMC1 and ASY4 localization, pachytene chromosome morphology as well as the SC

assembly in Col plants defective for ER and/or MPK6. Both the single *er* and *mpk6* and the double *er mpk6* mutants

displayed the same heat tolerance of meiotic recombination as Col (Fig. 4d, f, h, j and n).

To further explore the impact of the general genetic background on heat tolerance of meiosis, we analyzed

chromosome behavior in heat-stressed Col/L*er* hybrids, which exhibited the same flower architecture as Col (Fig. S2).

These plants produced a lower level of abnormal tetrads (~31.0%) than Col (Fig. 4a and b; Fig. 1b; $P < 0.001$).

Interestingly, bivalent formation in Col/L*er* hybrids was not reduced under heat stress (Fig. 4c and d), supporting the

conclusion that heat tolerance of meiotic recombination in L*er* is not conferred by the flower architecture. In addition,

balanced homolog segregation at anaphase I in heat-stressed Col/L*er* hybrids occurred at a similar frequency as in L*er*

(Fig. 4e and f; Fig. 2f; $P > 0.05$). Moreover, at 37°C, zygotene chromosomes in Col/L*er* hybrids showed an average

number of DMC1 foci (~116.0) significantly higher than those observed for Col (~74.1; $P < 0.001$) and L*er* (~101.6; $P <$

0.05) (Fig. 4g and h; Fig. 2j). Analysis of pachytene chromosome configurations and ASY4 localization revealed no

significant difference in heat tolerance between Col/L*er* hybrids and L*er* (Fig. 4i-l; Fig. 3j and l; $P > 0.05$). Furthermore,

most pachytene meiocytes (~74.1%) in Col/L*er* hybrids displayed a partial linear localization pattern of ZYP1, the extent

of which was higher than that in L*er* (Fig. 4m and n; Fig. 3t; $P < 0.001$); and the meiocytes showing linear or patchy

ZYP1 localization occupied 1.8% and 24.1%, respectively (Fig. 4m and n). Overall, these findings revealed similar heat

tolerance of meiotic recombination in Col/L*er* hybrids and in L*er*.

Figure 4. Col/L*er* hybrids exhibit similar heat tolerance of meiotic recombination as L*er*. (a) Representative images of normal and

abnormal tetrads. (b) Graph showing the rates of abnormal tetrad in Col/L*er* hybrids at 20°C and 37°C. (c) Representative images

of metaphase I chromosomes with different numbers of bivalents. The red and yellow arrows indicate univalents and bivalents, respectively. (d) Graph showing the average numbers of bivalent per diakinesis and/or metaphase I meiocyte in Col/Ler hybrids, the single *er-105* and *mpk6*, and the double *er-105 mpk6* mutants at 20°C and 37°C. (e) Representative images of anaphase I chromosomes showing balanced or unbalanced homolog separation. (f) Graph showing the frequencies of anaphase I meiocytes with balanced or unbalanced homolog separation in Col/Ler hybrids, the single *er-105* and *mpk6*, and the double *er-105 mpk6* mutants at 20°C and 37°C. (g) Immunolocalization of DMC1 on zygotene chromosomes in Col/Ler hybrids and *er-105* at 20°C and 37°C. White, DAPI; red, DMC1. (h) Graph showing the average numbers of DMC1 foci per meiocyte in Col/Ler hybrids and *er-105* at 20°C and 37°C. (i) Representative images of pachytene chromosomes showing full-pairing, partial-pairing, and no pairing. (j) Graph showing the rates of pachytene meiocytes exhibiting full-pairing, partial-pairing and no pairing in Col/Ler hybrids, the single *er-105* and *mpk6* and the double *er-105 mpk6* mutants at 20°C and 37°C. (k) Representative images of immunolocalization of ASY4 on zygotene chromosomes showing different configurations. White, DAPI; green, ASY4. (l) Graph showing the frequencies of ASY4 configurations in Col/Ler hybrids at 20°C and 37°C. (m) Representative images of immunolocalization of ZYP1 on pachytene chromosomes showing different configurations. White, DAPI; red, ZYP1. (n) Graph showing the rates of ZYP1 configurations on pachytene chromosomes in Col/Ler hybrids and the single *er-105* mutant at 20°C and 37°C. In (b), (d) and (h), stars indicate significance levels based on unpaired *t* tests; in (f), (j), (l) and (n), stars indicate significance levels based on χ^2-tests; ***, $P < 0.001$; **, $P < 0.01$; ns, $P > 0.05$; numbers or frequencies indicate the quantities or fractions of the corresponding phenotypes; n indicates the number of the analyzed individuals or cells. Scale bars, 10 μm.

Heat tolerance of pairing and CO formation is largely lost in the progeny of Col/Ler hybrids

To further analyze the genetic underpinnings of heat tolerance of meiotic recombination, we examined homolog pairing, synapsis and bivalent formation in the progeny of Col/Ler hybrids under heat stress. Unexpectedly, in a population of 1213 pachytene meiocytes from 132 Col/Ler F2 individuals, only 11.1% of meiocytes showed full-pairing of homologs. Partial-pairing and no pairing of homologs occurred 36.7% and 52.2%, respectively (Fig. 5a and b). Furthermore, only 19.8% of analyzed meiocytes (n = 3391) had five bivalents whereas 45.8% of meiocytes showed ten univalents, with the remaining 35% having either four, three, two or one bivalents occurring at roughly similar levels (Fig. 5a and c). Accordingly, 48.4% of F2 individuals (n = 155) showed an average of at most one bivalent per meiocyte, while 18.7% of F2 individuals showed an average of more than four bivalents per meiocyte (Fig. 5d). These data indicated that, in the context of bivalent formation, around half of the Col/Ler F2 individuals behaved like Col, while only about 20.0% individuals displayed a Ler-like resistance pattern under heat stress.

We analyzed the correlation of the impact of heat on pairing, bivalent formation, and homolog segregation in the heat-

stressed Col/L*er* F2 population by performing Spearman's rank correlation tests using the frequencies of meiocytes

showing different phenotypes (Fig. 5e). As expected, a strong and positive correlation was found between the numbers

of bivalents and homolog pairing (Fig. 5e), with the presence of two, three or four bivalents per meiocyte being

positively correlated with the partial-pairing phenotype (Fig. 5e). This correlation offered a proof that our assay system

worked. Moreover, positive correlations were found between balanced homolog segregation and bivalent formation

or successful pairing (Fig. 5e). Similar correlation patterns occurred in heat-stressed inbreds and F1 hybrids (Fig. S3).

These findings suggested that DSB formation and/or early stages of DSB repair-mediated homolog pairing are possibly

the primary targets of heat.

Figure 5. Chromosome pairing and bivalent formation in Col/L*er* F2 population under heat stress. (a) Representative images of

chromosomes in pachytene and diakinesis showing different phenotypes of homolog pairing and bivalent formation in Col/L*er* F2

population under heat stress. The red, yellow and green arrows indicate unpaired or unsynapsed regions, univalents and bivalents,

respectively. Scale bars, 10 μm. (b) Graph showing the fraction of pachytene meiocytes showing different pairing status in the

heat-stressed Col/L*er* F2 population. (c) Graph showing the fraction of diakinesis meiocytes showing different numbers of bivalents

in the heat-stressed Col/L*er* F2 population. (d) Graph showing the fraction of individuals in heat-stressed Col/L*er* F2 population

showing different average numbers of bivalents. The frequencies indicate the rates of the corresponding phenotypes; n indicates

the numbers of meiocytes or individuals; N indicates the average numbers of bivalent per meiocyte. (e) Graph showing correlation

between pairing, average number of bivalents per meiocyte, and homolog segregation. Spearman correlation analysis was

performed; numbers indicate the correlation coefficients between the corresponding phenotypes; significance levels of

correlation were determined based on *t*-tests; ***, $P < 0.001$; **, $P < 0.01$; ns, $P > 0.05$; the significance levels in the cells without

indication of significance values were not calculated.

Heat stress reveals different impacts on chromosome morphology in interkinesis between Col and L*er* inbreds and

their F2 offspring

At the end of meiosis I, homologs temporally decondense at interkinesis and then recondense at metaphase II (Fig.

S4). We found that heat induced a less-condensed or non-condensed chromosome configuration in interkinesis, which

also varied between Col and L*er* (Fig. 6a and c). The chromosomes exhibiting either kind of alterations in condensation

showed a lower DAPI fluorescence intensity compared with the ones displaying normal condensation (Fig. 6a and b).

At 37°C, 25.0% Col plants produced meiocytes with a less-condensed chromosome configuration, while, this

phenotype was not observed in any analyzed L*er* plant individuals (Fig. 6c). Meanwhile, meiocytes showing a non-

condensed chromosome configuration were observed in both Col and L*er* under heat stress (Fig. 6c). Both kinds of

abnormalities were observed in Col/L*er* hybrids and F2 population (Fig. 6c). However, in the F2 offspring that displayed

altered chromosome morphology, the fraction of meiocytes displaying non-condensed chromosomes was similar to

that in Col but higher than those in L*er* and Col/L*er* hybrids (Fig. 6d), while the fraction of meiocytes exhibiting the

less-condensed phenotype was higher than those in inbreds and hybrids (Fig. 6e).

Interestingly, positive correlations were found between the less-condensed phenotype and impaired pairing or

formation of only a single bivalent per meiocyte (Fig. 6f and g). In addition, normal morphology of interkinesis

chromosomes was highly correlated with the presence of five bivalents per meiocyte (Fig. 6g). The non-condensed

phenotype was positively associated with compromised pairing and reduced bivalent formation (Fig. 6f and g). Taken

together, these findings implied that the loci controlling heat tolerance of homolog pairing and/or CO formation may

have a potential role in regulating chromosome morphology in interkinesis under heat stress.

Figure 6. The impact of heat on chromosome morphology in interkinesis. (a) Representative images of different chromosome

morphologies in interkinesis. Scale bar, 10 μm. (b) Average pixel intensity quantified in interkinesis meiocytes showing either kind

of chromosome configuration of a section going through the middle of chromosomes (yellow lines). (c) Graph showing the rates

of individuals with less-condensed or non-condensed phenotypes in Col, Ler, Col/Ler hybrids and Col/Ler F2 at 20°C or 37°C. n

indicates the numbers of plant individuals. (d and e) Graphs showing the rates of meiocytes showing non-condensed (d) or less-

condensed (e) phenotypes in Col, Ler, Col/Ler hybrids and Col/Ler F2 at 20°C or 37°C. Stars indicate significance levels based on

unpaired *t* tests; n indicates the numbers of plant individuals; frequencies indicate the rates of the corresponding phenotypes. (f)

and (g) Graphs showing the correlation between pairing (f) and average number of bivalents per meiocyte (g) with the less-

condensed or non-condensed phenotypes. Spearman correlation analysis was performed; numbers indicate the correlation

coefficients between the corresponding phenotypes; stars indicate significance levels based on unpaired *t* tests; ***, $P < 0.001$;

**, $P < 0.01$; *, $P < 0.05$; ns, $P > 0.05$; the significance levels in the cells without indication of significance values were not calculated.

Heat stress induces high levels of chromosome instabilities in the Col/Ler F2 population

Heat did not significantly increase the fraction of individuals exhibiting premature segregation of sister chromatids in

Col and Ler (Fig. 7a-d). In the Col/Ler F2 population, however, 56.7% of individuals (n = 164) showed ectopic

segregation of sister chromatids from diakinesis to metaphase II (Fig. 7b and c), suggesting that sister-chromatid

cohesion was perturbed. In these defect-harboring F2 plants, the fraction of meiocytes showing untimely segregation

of sister chromatids was significantly higher than in parental genotypes (Fig. 7d). While we did not observe obvious

defects in chromosome integrity in Col and Ler under heat stress, 27.0% of individuals (n = 126) in the heat-stressed

Col/Ler F2 population produced chromosome fragments at metaphase I, anaphase I or metaphase II (Fig. 7e-g),

suggesting that DSB repair was defective in those F2 individuals under heat stress. Moreover, the average number of

fragments per meiocyte in those F2 individuals was higher than in the inbreds and hybrids (Fig. 7g). These data

suggested that heat tolerance of chromosome stability maintenance is affected in the Col/Ler F2 population.

Figure 7. The Col/L*er* F2 population shows a higher level of chromosome instability under heat stress. (a) and (b) Representative images of diakinesis, metaphase I, anaphase I, interkinesis and metaphase II meiocytes showing normal (a) or pre-mature segregation of sister chromatids (b) under heat stress. The yellow arrows indicate separated sister chromatids. (c) Graph showing the rates of individuals that show abnormal segregation of sister chromatids in Col, L*er*, Col/L*er* hybrids and Col/L*er* F2 at 20°C or 37°C. The indicated significance levels were determined based on χ^2-tests. (d) Graph showing the rates of meiocytes exhibiting premature separation of sister chromatids in Col, L*er*, Col/L*er* hybrids and Col/L*er* F2 at 20°C or 37°C. (e and f) Representative images of metaphase I, anaphase I and metaphase II chromosomes with normal (e) or interfered integrity (f) under heat stress. The red arrows indicate chromosome fragments. (g) Graph showing the average numbers of chromosome fragment per meiocyte in Col, L*er*, Col/L*er* hybrids and Col/L*er* F2 at 20°C or 37°C. Stars indicate significance levels based on unpaired *t* tests; frequencies indicate the rates of the corresponding phenotypes; n indicates the number of individuals; ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; ns, $P > 0.05$; scale bars, 10 μm.

Homolog pairing and synapsis facilitate timely sister-chromatids separation under heat stress

We then related homolog pairing and bivalent formation with the phenotype of premature sister-chromatids separation in our heat-stressed Col/L*er* F2 population. A positive but weak correlation was detected between the fraction of meiocytes showing cohesion defect with the fraction of meiocytes showing partial pairing, or the diakinesis meiocytes producing an average of one bivalent (Fig. 8a and b). Notably, among the meiocytes showing ectopic sister-chromatid separation, the meiocytes in metaphase I, anaphase I or metaphase II stages occupied 8.7%, 35.8% and 59.0%, respectively (Fig. 8c). This implied that heat stress primarily destabilizes centromeric cohesion in late or after meiosis I. Moreover, we found a strong and positive correlation between the fraction of meiocytes exhibiting chromosome fragmentations with meiocytes in pachytene undergoing normal homolog pairing (Fig. 8d) and with meiocytes in diakinesis producing five bivalents (Fig. 8e).

To further investigate the impact of homolog pairing and/or synapsis and CO formation on sister-chromatids cohesion under heat stress, we analyzed a *spo11-1* mutant (Col background), in which DSB formation is largely abolished and homolog pairing and recombination are impaired (Xue *et al.*, 2018). Untimely separation of sister chromatids was observed in *spo11-1* at both temperatures, but there was no significant difference (Fig. 8f-h, $P > 0.05$). Next, we analyzed the *dmc1* (Col background), in which DSBs are formed but homolog pairing and CO formation do not occur (Sanchez-Moran *et al.*, 2007), and the double *zyp1a/b* mutant (Col background), in which homolog pairing occurs but

synapsis is impaired (Capilla-Pérez *et al.*, 2021; France *et al.*, 2021; Yang *et al.*, 2022). Both *dmc1* and *zyp1a/b* exhibited

premature sister-chromatids segregation at 20°C, and, the fraction of meiocytes showing the cohesion defect in both

mutants were significantly increased under heat stress (Fig. 8f-h; *P* < 0.01 for *dmc1* and *P* < 0.001 for *zyp1a/b*).

Interestingly, the defect-harboring meiocytes in *dmc1* occurred by 66.7% at and/or prior to anaphase I, while, in

zyp1a/b, the irregularity was found by 79.2% at metaphase II (Fig. 8i). These findings implied that DMC1- and ZYP1-

dependent homolog pairing and/or synapsis play a role in facilitating timely separation of sister chromatids under heat

stress, yet possibly via different mechanisms. To address if the distinct distribution of differently-staged meiocytes

showing ectopic sister-chromatids separation between *dmc1* and *zyp1a/b* was owing to the difference in CO formation,

we examined a *msh4* mutant (Col background), which undergoes normal homolog synapsis but loses approximately

85% COs (Higgins *et al.*, 2004). No significant difference in the rates of meiocytes with cohesion defect was detected

at 20°C and 37°C (Fig. 8h, *P* > 0.05).

Figure 8. Homolog pairing and synapsis facilitate timely sister-chromatids separation under heat stress. (a and b) Graphs showing the correlation between the rates of meiocytes showing different pairing status (a) or bivalent formation (b) with the frequency of meiocytes with cohesion defect in the Col/L*er* F2 population under heat stress. (c) Graph showing the rates of metaphase I, anaphase I and metaphase II meiocytes with cohesion defect in the Col/L*er* F2 population under heat stress. (d and e) Graphs showing the correlation between the frequencies of meiocytes showing different pairing status (d) or bivalent formation (e) with the rate of meiocytes showing chromosome fragmentation in the Col/L*er* F2 population under heat stress. Spearman correlation analysis was performed. Numbers indicate the correlation coefficients between the corresponding phenotypes; significance of correlation was calculated using *t*-tests; the significance levels in the cells without indication of significance values were not calculated. (f and g) Representative images of diakinesis, metaphase I, anaphase I, interkinesis and metaphase II meiocytes showing normal (f) or untimely sister-chromatids separation (g). (h) Graph showing the rates of meiocytes with normal or untimely sister-chromatids separation in *spo11-1*, *dmc1*, *zyp1a/b* and *msh4* at 20°C and 37°C. (i) Graph showing the rates of different meiosis-staged meiocytes exhibiting untimely sister-chromatids separation in *dmc1* and *zyp1a/b* at 20°C and 37°C. For (c) and (h), χ^2-tests were performed; frequencies indicate the rates of the corresponding phenotypes; n indicates the number of analyzed cells; ***, $P < 0.001$; **, $P < 0.01$; ns, $P > 0.05$; scale bar, 10 μm.

Chromosome integrity in both Col and L*er* under heat stress relies on functional ATM

We previously reported that ATM-controlled DSB repair is required for maintenance of meiotic chromosome integrity in Col under heat stress (Zhao *et al.*, 2023). Thus, we decided to examine whether the role of ATM in protecting chromosome stability under heat stress varies between Col and L*er*. To this end, we took advantage of an *atm* mutant allele, *atm-5*, which is in L*er* background and shows defects in DSB repair (Yao *et al.*, 2020). We found that, as the case in *atm-2*, an allele in Col background (Zhao *et al.*, 2023), chromosome fragments in *atm-5* was significantly increased after heat treatment (Fig. 9a and b; $P < 0.0001$). A bi-allelic *atm* mutant $ATM^{atm-2/atm-5}$, which harbored the *atm-2* and *atm-5* alleles on either homolog, was generated, and behaved similarly as *atm-2* and *atm-5* under heat stress (Fig. 9c and d). We also examined the impact of heat on homolog pairing in *atm-5*. At 20°C, most pachytene meiocytes showed normal pairing and synapsis of homologs, and 16.9% of meiocytes showed a partial pairing phenotype (Fig. 9e and g). At 37°C, most pachytene meiocytes in *atm-5* showed a partial pairing phenotype, with only 3.9% cells showing an impaired homolog pairing (Fig. 9e and g), similar to the case in heat-stressed L*er* (Fig. 3j). Immunolocalization revealed an increased level of patchy ZYP1 configuration in *atm-5* compared with that in L*er* (Fig. 9f and h; Fig. 3t), which was likely caused by the disrupted chromosome integrity due to ATM dysfunction (Zhao *et al.*, 2023). Similar alterations in

homolog pairing and SC formation were observed in $ATM^{atm-2/atm-5}$ at either temperature (Fig. 9i-l). These data suggest that there is no accession-specific difference in dependency of ATM-mediated DSB repair for meiotic chromosome integrity under heat stress.

Figure 9. Chromosome integrity under heat stress in both Col and L*er* relies on ATM. (a and c), Metaphase I, anaphase I, metaphase II and telophase II chromosomes in *atm-5* (a) and $ATM^{atm-2/atm-5}$ (c) at 20°C and 37°C. (b and d), Graphs showing the average number of chromosome fragment in *atm-5* (b) and $ATM^{atm-2/atm-5}$ (d) at 20°C and 37°C. Significance levels are shown based on unpaired *t* tests. (e and i), Representative images of pachytene chromosomes showing fully-paired, partially-paired, or no pairing in *atm-5* (e) and $ATM^{atm-2/atm-5}$ (i) at 20°C and 37°C. (f and j), Representative images of immunolocalization of ZYP1 on pachytene chromosomes showing linear, partial linear, or patchy phenotypes in *atm-5* (f) and $ATM^{atm-2/atm-5}$ (j) at 20°C and 37°C. The blue arrow indicates

linear ZYP1 signals. (g and k), Graphs showing the rates of different pairing status of pachytene chromosomes in *atm-5* (g) and

ATM^atm-2/atm-5 (k) at 20°C and 37°C. Significance levels are shown based on χ^2-tests. (h and l), Graphs showing the rates of different

ZYP1 configurations on pachytene chromosomes in *atm-5* (h) and *ATM^atm-2/atm-5* (l) at 20°C and 37°C. Significance levels are shown

based on χ^2-tests; frequencies or numbers indicate the rate or quantity of the corresponding phenotypes; n indicates the number

of analyzed cells; ***, *P* < 0.001; scale bars, 10 μm.

Discussion

Our previous study demonstrated that heat stress reduces DSB formation and disrupts pairing as well as synapsis of

homologs in Col (Ning *et al.*, 2021; Fu *et al.*, 2022). In this study, we report that Col and L*er* have a striking difference

in heat tolerance of meiotic recombination. Specifically, heat stress does not impair SC formation and does not reduce

the numbers of RAD51 or DMC1 foci in L*er* (Fig. 2 and 3). Based on our previous studies, the distinct heat tolerance of

DSB formation between Col and L*er* is likely due to a difference in the activity and/or localization of the recombination

proteins, and not in transcription levels (Ning *et al.*, 2021; Zhao *et al.*, 2023). Since the number of bivalents is strongly

correlated with homolog pairing status (Fig. 5), it is likely that heat suppresses meiotic recombination predominantly

by interfering with DSB formation and/or the RAD51- and DMC1-mediated homolog pairing. In support, DMC1 has

been found to play a role in maintaining CO formation at high temperature in wheat (Draeger *et al.*, 2019; Draeger *et*

al., 2023). Nevertheless, since there is still a low level of DSBs in heat-stressed Col (Fig. 2), and some meiocytes in L*er*

display compromised homolog pairing under heat stress (Fig. 3), we propose that heat can also directly attenuate

homolog pairing apart from reducing DSB formation (Grelon *et al.*, 2001; Ning *et al.*, 2021; Fu *et al.*, 2022; Zhao *et al.*,

2023). Based on the analysis of Col/L*er* hybrids, we conclude that the heat tolerance of meiotic recombination in L*er*

is a dominant trait (Fig. 4). 10% ~ 20% of F2 offspring behaving as L*er* suggests that the dominance relies on the

presence of several alleles. While all alleles are present in Col/L*er* hybrids, the linkage of these alleles is broken in the

meiosis of these hybrids leading to a differentiated pattern of heat tolerance in the F2 population (Fig. 5).

Heat induces an increased fraction of interkinesis meiocytes displaying a non-condensed chromosome configuration

in Col, which does not occur in L*er* and the Col/L*er* F1 hybrids, but occurs again, and, at the same level, in the F2

population (Fig. 6d). Meanwhile, heat increases the rate of interkinesis meiocytes exhibiting a less-condensed

phenotype in some F2 individuals but not in inbreds or hybrids, which suggests a higher heat sensitivity in F2 offspring

(Fig. 6e). The heat-induced abnormalities in interkinesis phenotypically mimic slowed meiosis progression or a failure

of entering into meiosis II, which implies that meiosis progression and/or cell cycle transition may be perturbed possibly due to alterations in expression and/or function of *TAM/CYCA1;2* (*TAM*) or *OMISSION OF SECOND DIVISION 1* (*OSD1*), or other meiotic cell cycle regulators (Cromer *et al.*, 2012; Prusicki *et al.*, 2019; Ning *et al.*, 2021; Schindfessel *et al.*, 2022). Furthermore, correlation analysis suggests that meiotic recombination or its related regulators may impact meiotic cell cycle progression and/or chromosome morphology after anaphase I beyond its impact on prophase I progression under heat stress (Fig. 6) (De Jaeger-Braet *et al.*, 2022).

Incorrect sister-chromatids separation in the F2 offspring (Fig. 7) implies that heat perturbs the function of the cohesion complex. It has recently been reported that high temperatures reduce the amounts of CENH3 in Arabidopsis (Crhak Khaitova *et al.*, 2024), which may lead to interfered centromere function and cohesion instability. Notably, we found that most meiocytes showing cohesion defects are in anaphase I or metaphase II (Fig. 8), suggesting that heat primarily perturbs centromeric cohesion during anaphase I and interkinesis. In Arabidopsis, SHUGOSHIN 1 (SGO1) and PROTEIN PHOSPHATASE 2A (PP2A), and the Arabidopsis homolog of securin PATRONUS (PANS1) protect meiotic centromeric cohesion at anaphase I and diakinesis, respectively (Cromer *et al.*, 2013; Zamariola *et al.*, 2014; Yuan *et al.*, 2018; Cromer *et al.*, 2019). In addition, PP2A is co-expressed with genes involved in response to abiotic stresses including heat in multiple developmental stages in Arabidopsis (Khan *et al.*, 2020). It is thus possible that heat stress alters the expression and/or function of these cohesion regulators. In multiple species, bivalent formation is required for correct orientation of sister kinetochores and thus balanced chromosome segregation in meiosis I (Hunt *et al.*, 1995; Rebollo & Arana, 1995; Khodjakov *et al.*, 1997; Chan & Cande, 1998; Yamamoto & Hiraoka, 2003; Xue *et al.*, 2019). Our correlation analysis suggests that homolog pairing and/or synapsis but possibly not chiasmata formation may directly or indirectly impact timely sister-chromatids separation under heat stress (Fig. 8). However, heat-stressed *spo11-1* does not show a higher level of defective sister-chromatids separation (Fig. 8), suggesting that the effect of heat stress on homolog pairing-mediated cohesion stability may rely on DSB formation, and, the increased aneuploid nuclei due to chromosome mis-segregation observed in heat-stressed *spo11-2* are not owing to premature sister-chromatids separation (Crhak Khaitova *et al.*, 2024).

We observed an increase of chromosome fragments in some Col/L*er* F2 individuals but not in either inbreds or hybrids under heat stress (Fig. 7), suggesting that DSB repair is perturbed in those fragment-harboring F2 plants. This phenotype, together with the increased fraction of meiocytes showing premature sister-chromatid separation, imply that transgressive effects become visible in the F2 population showing defects exceeding that observed in parents.

ATM-mediated DSB repair is required for meiotic chromosome integrity in both Col and L*er* under heat stress (Fig. 9)

(Zhao *et al.*, 2023), dysfunction of which might be one of the mechanisms that underpin the heat-interfered

chromosome integrity in those F2 individuals. The upstream factor and/or signaling pathway that regulates ATM

expression or activity in DSB repair in response to heat stress require further investigation. Our data suggest that there

is no accession-specific difference in the dependency of ATM-mediated DSB repair for maintenance of meiotic

chromosome integrity under heat stress. Alternatively, ATM-mediated mechanism acts in an epistatic manner to

protect meiotic chromosome stability under heat stress. Interestingly, the fragmentation phenotype is strongly and

positively correlated with successful homolog pairing and bivalent formation (Fig. 8d and e). We speculate that these

associations are possibly a subsequent effect of heat stress on DSB formation and repair, the heat tolerance of which

are regulated via different pathways. Specifically, in the individuals that show normal homolog pairing and CO

formation, DSB formation likely occurs normally, however, DSB repair may be perturbed, which consequently leads to

lesions in chromosome integrity. On the contrary, the individuals with impaired homolog pairing and CO formation

likely have a significantly reduced DSB formation, which compensates for the perturbation of DSB repair, and thus

show relatively undisturbed chromosome integrity. Taken together, our study reveals that natural variation in multiple

genomic loci controls heat tolerance of meiosis in Arabidopsis, with DSB formation and homolog pairing possibly being

the predominant targets, which facilitate meiosis progression and genome stability after meiotic recombination at

high temperatures.

Acknowledgement

The authors thank the National Natural Science Foundation of China (32000245), Hubei Provincial Natural Science

Foundation of China (2024AFB695), the Fundamental Research Funds for the Central Universities of South-Central

Minzu University (CZZ24011), Knowledge Innovation Program of Wuhan-Shuguang Project (2022020801020410) and

general funds of University of Hamburg (to A.S.) for the financial support on this study. They thank Pingli Lu (Henan

University) for sharing the *atm-5* mutants. They thank Chao Yang (HZAU) for providing the *zyp1a/b* mutant. They thank

Yuan Qin (FAFU) for sharing the *er-105*, *mpk6* and the *er-105 mpk6* mutants.

Conflicts of interest

None declared.

Data availability

All data that support the conclusions drawn in our study are fully documented in this article and Supporting Information (including Fig. S1–S4; Table S1).

Author contributions

J.Z. and H.F. performed most cytological experiments, genotyped plants, analyzed data, and edited manuscript; Z.W., Z.R., Z.W. and X.L. performed statistical analysis; M.Z., Y.L., X.C., W.P., Y.Z., X.G. and L.H. analyzed chromosome spreads; C.W., A.S. and W.P. analyzed data and edited manuscript; B.L. conceived the project, designed experiments, analyzed data, wrote, and edited the manuscript.

Accession numbers

Accession numbers of genes studied in this work are: *SPO11-1* (AT3G13170), *RAD51* (AT5G20850), *DMC1* (AT3G22880), *ASY4* (AT2G33793), *ZYP1A* (AT1G22260), *ZYP1B* (AT1G22275), *HEI10* (AT1G53490), *ERECTA* (AT2G26330), *MPK6* (AT2G43790), *ATM* (AT3G48190) and *MSH4* (AT4G17380).

References

Ahmadli U, Kalidass M, Khaitova LC, Fuchs J, Cuacos M, Demidov D, Zuo S, Pecinkova J, Mascher M, Ingouff M, et al. 2023. High temperature increases centromere-mediated genome elimination frequency and enhances haploid induction in Arabidopsis. *Plant Communications* **4**(3): 100507.

Armstrong SJ, Caryl AP, Jones GH, Franklin FC. 2002. Asy1, a protein required for meiotic chromosome synapsis, localizes to axis-associated chromatin in Arabidopsis and Brassica. *Journal of Cell Science* **115**(Pt 18): 3645-3655.

Armstrong SJ, Jones GH. 2003. Meiotic cytology and chromosome behaviour in wild‐type Arabidopsis thaliana. *Journal of Experimental Botany* **54**(380): 1-10.

Bomblies K, Higgins JD, Yant L. 2015. Meiosis evolves: adaptation to external and internal environments. *New Phytologist* **208**(2): 306-323.

Cai H, Zhao L, Wang L, Zhang M, Su Z, Cheng Y, Zhao H, Qin Y. 2017. ERECTA signaling controls Arabidopsis inflorescence architecture through chromatin-mediated activation of PRE1 expression. *New Phytologist* **214**(4): 1579-1596.

Capilla-Pérez L, Durand S, Hurel A, Lian Q, Chambon A, Taochy C, Solier V, Grelon M, Mercier R. 2021. The synaptonemal complex imposes crossover interference and heterochiasmy in Arabidopsis. *Proc Natl Acad Sci U S A* **118**(12): e2023613118.

Chambon A, West A, Vezon D, Horlow C, De Muyt A, Chelysheva L, Ronceret A, Darbyshire A, Osman K, Heckmann S, et al. 2018. Identification of ASYNAPTIC4, a component of the meiotic chromosome axis. *Plant Physiology* **178**(1): 233-246.

Chan A, Cande WZ. 1998. Maize meiotic spindles assemble around chromatin and do not require paired chromosomes. *Journal of Cell Science* **111**(23): 3507-3515.

Chelysheva L, Grandont L, Vrielynck N, Guin Sl, Mercier R, Grelon M. 2010. An easy protocol for studying chromatin and recombination protein dynamics during Arabidopsis thaliana meiosis: immunodetection of cohesins, histones and MLH1. *Cytogenetic and Genome Research* **129**(1/3): 143-153.

Chen H, He C, Wang C, Wang X, Ruan F, Yan J, Yin P, Wang Y, Yan S. 2021. RAD51 supports DMC1 by inhibiting the SMC5/6 complex during meiosis. *Plant Cell* **33**(8): 2869-2882.

Chen Y, Dubois M, Vermeersch M, Inzé D, Vanhaeren H. 2021. Distinct cellular strategies determine sensitivity to mild drought of Arabidopsis natural accessions. *Plant Physiology* **186**(2): 1171-1185.

Chen Y, Zhang Q, Teng Z, Liu H. 2021. Centromeric transcription maintains centromeric cohesion in human cells. *Journal of Cell Biology* **220**(7).

Choi WG, Barker RJ, Kim SH, Swanson SJ, Gilroy S. 2019. Variation in the transcriptome of different ecotypes of Arabidopsis thaliana reveals signatures of oxidative stress in plant responses to spaceflight. *American Journal of Botany* **106**(1): 123-136.

Couteau F, Belzile F, Horlow C, Grandjean O, Vezon D, Doutriaux MP. 1999. Random chromosome segregation without meiotic arrest in both male and female meiocytes of a dmc1 mutant of Arabidopsis. *Plant Cell* **11**(9): 1623-1634.

Crhak Khaitova L, Mikulkova P, Pecinkova J, Kalidass M, Heckmann S, Lermontova I, Riha K. 2024. Heat stress impairs centromere structure and segregation of meiotic chromosomes in Arabidopsis. *eLife* **12**.

Cromer L, Heyman J, Touati S, Harashima H, Araou E, Girard C, Horlow C, Wassmann K, Schnittger A, De Veylder L, et al. 2012. OSD1 promotes meiotic progression via APC/C inhibition and forms a regulatory network with TDM and CYCA1;2/TAM. *Plos Genetics* **8**(7): e1002865.

Cromer L, Jolivet S, Horlow C, Chelysheva L, Heyman J, De Jaeger G, Koncz C, De Veylder L, Mercier R. 2013. Centromeric cohesion is protected twice at meiosis, by SHUGOSHINs at anaphase I and by PATRONUS at interkinesis. *Current Biology* **23**(21): 2090-2099.

Cromer L, Jolivet S, Singh DK, Berthier F, De Winne N, De Jaeger G, Komaki S, Prusicki MA, Schnittger A, Guérois R, et al. 2019. Patronus is the elusive plant securin, preventing chromosome separation by antagonizing separase. *Proc Natl Acad Sci U S A* **116**(32): 16018-16027.

Da Ines O, Degroote F, Goubely C, Amiard S, Gallego ME, White CI. 2013. Meiotic recombination in Arabidopsis is catalysed by DMC1, with RAD51 playing a supporting role. *Plos Genetics* **9**(9): e1003787.

De Jaeger-Braet J, Krause L, Buchholz A, Schnittger A. 2022. Heat stress reveals a specialized variant of the pachytene checkpoint in meiosis of Arabidopsis thaliana. *Plant Cell* **34**(1): 433-454.

De Jaeger-Braet J, Schnittger A. 2024. Heating up meiosis – chromosome recombination and segregation under high temperatures. *Current Opinion in Plant Biology* **80**: 102548.

De Storme N, Copenhaver GP, Geelen D. 2012. Production of diploid male gametes in Arabidopsis by cold-induced destabilization of postmeiotic radial microtubule arrays. *Plant Physiology* **160**(4): 1808-1826.

De Storme N, Geelen D. 2020. High temperatures alter cross-over distribution and induce male meiotic restitution in Arabidopsis thaliana. *Communications Biology* **3**(1): 187.

Draeger T, A CM, Alabdullah AK, Pendle A, Rey MD, Shaw P, Moore G. 2019. Dmc1 is a candidate for temperature tolerance during wheat meiosis. *Theoretical and Applied Genetics*.

Draeger TN, Rey MD, Hayta S, Smedley M, Martin AC, Moore G. 2023. DMC1 stabilizes crossovers at high and low temperatures during wheat meiosis. *Frontiers in Plant Science* **14**: 1208285.

Ferdous M, Higgins JD, Osman K, Lambing C, Roitinger E, Mechtler K, Armstrong SJ, Perry R, Pradillo M, Cuñado N, et al. 2012.

Inter-homolog crossing-over and synapsis in Arabidopsis meiosis are dependent on the chromosome axis protein AtASY3.
Plos Genetics **8**(2): e1002507-e1002507.

France MG, Enderle J, Röhrig S, Puchta H, Franklin FCH, Higgins JD. 2021. ZYP1 is required for obligate cross-over formation and
cross-over interference in Arabidopsis. *Proc Natl Acad Sci U S A* **118**(14).

Fu H, Zhao J, Ren Z, Yang K, Wang C, Zhang X, Elesawi IE, Zhang X, Xia J, Chen C, et al. 2022. Interfered chromosome pairing at
high temperature promotes meiotic instability in autotetraploid Arabidopsis. *Plant Physiology* **188**(2): 1210-1228.

Grelon M, Vezon D, Gendrot G, Pelletier G. 2001. AtSPO11-1 is necessary for efficient meiotic recombination in plants. *Embo j*
20(3): 589-600.

Higgins JD, Armstrong SJ, Franklin FCH, Jones GH. 2004. The Arabidopsis MutS homolog AtMSH4 functions at an early step in
recombination: evidence for two classes of recombination in Arabidopsis. *Genes & Development* **18**(20): 2557-2570.

Hunt P, LeMaire R, Embury P, Sheean L, Mroz K. 1995. Analysis of chromosome behavior in intact mammalian oocytes: monitoring
the segregation of a univalent chromosome during female meiosis. *Hum Mol Genet* **4**(11): 2007-2012.

Jin C, Sun L, Trinh HK, Danny G. 2023. Heat stress promotes haploid formation during CENH3-mediated genome elimination in
Arabidopsis. *Plant Reproduction* **36**(2): 147-155.

Jones GH, Franklin FC. 2006. Meiotic crossing-over: obligation and interference. *Cell* **126**(2): 246-248.

Khan ZH, Agarwal S, Rai A, Memaya MB, Mehrotra S, Mehrotra R. 2020. Co-expression network analysis of protein phosphatase
2A (PP2A) genes with stress-responsive genes in Arabidopsis thaliana reveals 13 key regulators. *Scientific Reports* **10**(1):
21480.

Khodjakov A, Cole RW, McEwen BF, Buttle KF, Rieder CL. 1997. Chromosome fragments possessing only one kinetochore can
congress to the spindle equator. *Journal of Cell Biology* **136**(2): 229-240.

Kurzbauer M-T, Janisiw MP, Paulin LF, Prusén Mota I, Tomanov K, Krsicka O, Haeseler Av, Schubert V, Schlögelhofer P. 2021. ATM
controls meiotic DNA double-strand break formation and recombination and affects synaptonemal complex organization
in plants. *Plant Cell*.

Kurzbauer MT, Uanschou C, Chen D, Schlogelhofer P. 2012. The recombinases DMC1 and RAD51 are functionally and spatially
separated during meiosis in Arabidopsis. *Plant Cell* **24**(5): 2058-2070.

Lambing C, Kuo PC, Tock AJ, Topp SD, Henderson IR. 2020. ASY1 acts as a dosage-dependent antagonist of telomere-led
recombination and mediates crossover interference in Arabidopsis. *Proc Natl Acad Sci U S A* **117**(24): 13647-13658.

Lambing C, Tock AJ, Topp SD, Choi K, Kuo PC, Zhao X, Osman K, Higgins JD, Franklin FCH, Henderson IR. 2020. Interacting genomic
landscapes of REC8-cohesin, chromatin, and meiotic recombination in Arabidopsis. *Plant Cell* **32**(4): 1218-1239.

Lardon R, Wijnker E, Keurentjes J, Geelen D. 2020. The genetic framework of shoot regeneration in Arabidopsis comprises master
regulators and conditional fine-tuning factors. *Communications Biology* **3**(1): 549.

Lasky JR, Des Marais DL, Lowry DB, Povolotskaya I, McKay JK, Richards JH, Keitt TH, Juenger TE. 2014. Natural variation in abiotic
stress responsive gene expression and local adaptation to climate in Arabidopsis thaliana. *Molecular Biology and*
Evolution **31**(9): 2283-2296.

Lei X, Ning Y, Eid Elesawi I, Yang K, Chen C, Wang C, Liu B. 2020. Heat stress interferes with chromosome segregation and
cytokinesis during male meiosis in Arabidopsis thaliana. *Plant Signaling & Behavior* **15**(5): 1746985.

Liu B, De Storme N, Geelen D. 2017. Gibberellin induces diploid pollen formation by interfering with meiotic cytokinesis. *Plant*
Physiology **173**(1): 338-353.

Liu B, De Storme N, Geelen D. 2018. Cold-induced male meiotic restitution in Arabidopsis thaliana is not mediated by GA-DELLA
signaling. *Frontiers in Plant Science* **9**: 91.

Liu B, Mo WJ, Zhang D, De Storme N, Geelen D. 2019. Cold influences male reproductive development in plants: a hazard to
fertility, but a window for evolution. *Plant Cell Physiology* **60**(1): 7-18.

Lloyd A, Morgan C, H. Franklin FC, Bomblies K. 2018. Plasticity of meiotic recombination rates in response to temperature in
Arabidopsis. *Genetics* **208**(4): 1409-1420.

Lyu YS, Shao YJ, Yang ZT, Liu JX. 2020. Quantitative proteomic analysis of ER stress response reveals both common and specific
features in two contrasting ecotypes of Arabidopsis thaliana. *Int J Mol Sci* **21**(24).

McKinley KL, Cheeseman IM. 2016. The molecular basis for centromere identity and function. *Nature Reviews Molecular Cell*
Biology **17**(1): 16-29.

Meng X, Wang H, He Y, Liu Y, Walker JC, Torii KU, Zhang S. 2012. A MAPK cascade downstream of ERECTA receptor-like protein
kinase regulates Arabidopsis inflorescence architecture by promoting localized cell proliferation. *Plant Cell* **24**(12): 4948-
4960.

Modliszewski JL, Copenhaver GP. 2017. Meiotic recombination gets stressed out: CO frequency is plastic under pressure. *Current*
Opinion in Plant Biology **36**: 95-102.

Modliszewski JL, Wang H, Albright AR, Lewis SM, Bennett AR, Huang J, Ma H, Wang Y, Copenhaver GP. 2018. Elevated
temperature increases meiotic crossover frequency via the interfering (Type I) pathway in Arabidopsis thaliana. *Plos*
Genetics **14**(5): e1007384.

Ning Y, Liu Q, Wang C, Qin E, Wu Z, Wang M, Yang K, Elesawi IE, Chen C, Liu H, et al. 2021. Heat stress interferes with formation
of double-strand breaks and homolog synapsis. *Plant Physiology* **185**(4): 1783-1797.

Nore A, Juarez-Martinez AB, Clément J, Brun C, Diagouraga B, Laroussi H, Grey C, Bourbon HM, Kadlec J, Robert T, et al. 2022.
TOPOVIBL-REC114 interaction regulates meiotic DNA double-strand breaks. *Nature Communications* **13**(1): 7048.

Onn I, Heidinger-Pauli JM, Guacci V, Unal E, Koshland DE. 2008. Sister chromatid cohesion: a simple concept with a complex
reality. *Annu Rev Cell Dev Biol* **24**: 105-129.

Prusicki MA, Keizer EM, van Rosmalen RP, Komaki S, Seifert F, Müller K, Wijnker E, Fleck C, Schnittger A. 2019. Live cell imaging
of meiosis in Arabidopsis thaliana. *eLife* **8**: e42834.

Rebollo E, Arana P. 1995. A comparative study of orientation at behavior of univalent in living grasshopper spermatocytes.
Chromosoma **104**(1): 56-67.

Sanchez-Moran E, Santos JL, Jones GH, Franklin FC. 2007. ASY1 mediates AtDMC1-dependent interhomolog recombination during
meiosis in Arabidopsis. *Genes & Development* **21**(17): 2220-2233.

Schiavon M, Zhang L, Abdel-Ghany SE, Pilon M, Malagoli M, Pilon-Smits EAH. 2007. Variation in copper tolerance in Arabidopsis
thaliana accessions Columbia, Landsberg erecta and Wassilewskija. *Physiologia Plantarum* **129**(2): 342-350.

Schindfessel C, Jin C, Geelen D. 2022. Heat induced male meiotic restitution and sexual polyploidisation through natural variation
of the Arabidopsis cyclin TAM/CYCA1;2. *BioRxiv*: 2022.2008.2011.503651.

Schulz E, Tohge T, Zuther E, Fernie AR, Hincha DK. 2015. Natural variation in flavonol and anthocyanin metabolism during cold
acclimation in Arabidopsis thaliana accessions. *Plant, Cell and Environment* **38**(8): 1658-1672.

Sidhu GK, Fang C, Olson MA, Falque M, Martin OC, Pawlowski WP. 2015. Recombination patterns in maize reveal limits to
crossover homeostasis. *Proc Natl Acad Sci U S A* **112**(52): 15982-15987.

Su H, Cheng Z, Huang J, Lin J, Copenhaver GP, Ma H, Wang Y. 2017. Arabidopsis RAD51, RAD51C and XRCC3 proteins form a
complex and facilitate RAD51 localization on chromosomes for meiotic recombination. *Plos Genetics* **13**(5): e1006827.

Taghavi Namin S, Esmaeilzadeh M, Najafi M, Brown TB, Borevitz JO. 2018. Deep phenotyping: deep learning for temporal
phenotype/genotype classification. *Plant Methods* **14**(1): 66.

Tang Z, Zhang L, Yang D, Zhao C, Zheng Y. 2011. Cold stress contributes to aberrant cytokinesis during male meiosis I in a wheat
thermosensitive genic male sterile line. *Plant Cell and Environment* **34**(3): 389-405.

Vrielynck N, Schneider K, Rodriguez M, Sims J, Chambon A, Hurel A, De Muyt A, Ronceret A, Krsicka O, Mézard C, et al. 2021.
Conservation and divergence of meiotic DNA double strand break forming mechanisms in Arabidopsis thaliana. *Nucleic*
acids research **49**(17): 9821-9835.

Wang Y, Cheng Z, Lu P, Timofejeva L, Ma H. 2014. Molecular cell biology of male meiotic chromosomes and isolation of male
meiocytes in Arabidopsis thaliana. *Methods Mol Biol* **1110**: 217-230.

Wang Y, Copenhaver GP. 2018. Meiotic recombination: mixing it up in plants. *Annual Review of Plant Biology* **69**: 577-609.

West AM, Rosenberg SC, Ur SN, Lehmer MK, Ye Q, Hagemann G, Caballero I, Usón I, MacQueen AJ, Herzog F, et al. 2019. A
conserved filamentous assembly underlies the structure of the meiotic chromosome axis. *eLife* **8**.
Xue M, Wang J, Jiang L, Wang M, Wolfe S, Pawlowski WP, Wang Y, He Y. 2018. The number of meiotic double-strand breaks
influences crossover distribution in Arabidopsis. *Plant Cell* **30**(10): 2628-2638.
Xue Z, Liu C, Shi W, Miao Y, Shen Y, Tang D, Li Y, You A, Xu Y, Chong K, et al. 2019. OsMTOPVIB is required for meiotic bipolar
spindle assembly. *Proc Natl Acad Sci U S A* **116**(32): 15967-15972.
Yamamoto A, Hiraoka Y. 2003. Monopolar spindle attachment of sister chromatids is ensured by two distinct mechanisms at the
first meiotic division in fission yeast. *Embo j* **22**(9): 2284-2296.
Yang C, Hamamura Y, Sofroni K, Böwer F, Stolze SC, Nakagami H, Schnittger A. 2019. SWITCH 1/DYAD is a WINGS APART-LIKE
antagonist that maintains sister chromatid cohesion in meiosis. *Nature Communications* **10**(1): 1755-1755.
Yang C, Sofroni K, Hamamura Y, Hu B, Elbasi HT, Balboni M, Chu L, Stang D, Heese M, Schnittger A. 2022. ZYP1-mediated
recruitment of PCH2 to the synaptonemal complex remodels the chromosome axis leading to crossover restriction.
Nucleic acids research **50**(22): 12924-12937.
Yao Y, Li X, Chen W, Liu H, Mi L, Ren D, Mo A, Lu P. 2020. ATM promotes RAD51-mediated meiotic DSB repair by inter-sister-
chromatid recombination in Arabidopsis. *Frontiers in Plant Science* **11**: 839.
Yuan G, Ahootapeh BH, Komaki S, Schnittger A, Lillo C, De Storme N, Geelen D. 2018. PROTEIN PHOSHATASE 2A B'α and β
maintain centromeric sister chromatid cohesion during meiosis in Arabidopsis. *Plant Physiology* **178**(1): 317-328.
Zamariola L, De Storme N, Vannerum K, Vandepoele K, Armstrong SJ, Franklin FC, Geelen D. 2014. SHUGOSHINs and PATRONUS
protect meiotic centromere cohesion in Arabidopsis thaliana. *Plant Journal* **77**(5): 782-794.
Zhang SS, Yang H, Ding L, Song ZT, Ma H, Chang F, Liu JX. 2017. Tissue-specific transcriptomics reveals an important role of the
unfolded protein response in maintaining fertility upon heat stress in Arabidopsis. *Plant Cell* **29**(5): 1007-1023.
Zhao J, Gui X, Ren Z, Fu H, Yang C, Wang W, Liu Q, Zhang M, Wang C, Schnittger A, et al. 2023. ATM-mediated double-strand
break repair is required for meiotic genome stability at high temperature. *Plant Journal* **114**(2): 403-423.
Zhou Q, Cheng X, Kong B, Zhao Y, Li Z, Sang Y, Wu J, Zhang P. 2022. Heat shock-induced failure of meiosis I to meiosis II transition
leads to 2n pollen formation in a woody plant. *Plant Physiology* **189**(4): 2110-2127.
Zhu L, Fernández-Jiménez N, Szymanska-Lejman M, Pelé A, Underwood CJ, Serra H, Lambing C, Dluzewska J, Bieluszewski T,
Pradillo M, et al. 2021. Natural variation identifies SNI1, the SMC5/6 component, as a modifier of meiotic crossover in
Arabidopsis. *Proc Natl Acad Sci U S A* **118**(33).
Ziolkowski PA. 2023. Why do plants need the ZMM crossover pathway? A snapshot of meiotic recombination from the perspective
of interhomolog polymorphism. *Plant Reproduction* **36**(1): 43-54.
Ziolkowski PA, Underwood CJ, Lambing C, Martinez-Garcia M, Lawrence EJ, Ziolkowska L, Griffin C, Choi K, Franklin FC,
Martienssen RA, et al. 2017. Natural variation and dosage of the HEI10 meiotic E3 ligase control Arabidopsis crossover
recombination. *Genes & Development* **31**(3): 306-317.

Supporting Information

Figure S1. Heat stress does not affect microtubule organization during prophase I in *Arabidopsis thaliana.*

Figure S2. Flower architectures in Col, L*er* and Col/L*er* hybrids.

Figure S3. Correlation analysis of synapsis, bivalent formation, and segregation of homologs at anaphase I in Col, L*er*

and Col/L*er* hybrids.

Fig S4. Representative images of meiotic chromosomes from anaphase I to metaphase II in wild-type Arabidopsis.

Table S1. Primers used in this study.

786