Title: Social odors drive hippocampal CA2 place cell responses to social stimuli

Running title: CA2 place cells respond to social odors

Emma Robson [a b 1], Margaret M. Donahue [a c 1], Alexandra J. Mably [a b], Peyton G. Demetrovich [a c], Lauren T. Hewitt [a c], Laura Lee Colgin [a b c]

[a] Center for Learning and Memory, The University of Texas at Austin, Austin, TX 78712

[b] Department of Neuroscience, The University of Texas at Austin, Austin, TX 78712

[c] Institute for Neuroscience, The University of Texas at Austin, Austin, TX 78712

[1]These authors contributed equally.

Corresponding author: Laura Lee Colgin, The Center for Learning and Memory, The University of Texas at Austin, 1 University Station Stop C7000, Austin, TX 78712, USA. Email:

colgin@mail.clm.utexas.edu

Abstract

Hippocampal region CA2 is essential for social memory processing. Interaction with social stimuli induces changes in CA2 place cell firing during active exploration and sharp wave-ripples during rest following a social interaction. However, it is unknown whether these changes in firing patterns are caused by integration of multimodal social stimuli or by a specific sensory modality associated with a social interaction. Rodents rely heavily on chemosensory cues in the form of olfactory signals for social recognition processes. To determine the extent to which social olfactory signals contribute to CA2 place cell responses to social stimuli, we recorded CA2 place cells in rats freely exploring environments containing stimuli that included or lacked olfactory content. We found that CA2 place cell firing patterns significantly changed only when social odors were prominent. Also, place cells that increased their firing in the presence of social odors alone preferentially increased their firing during subsequent sharp wave-ripples. Our results suggest that social olfactory cues are essential for changing CA2 place cell firing patterns during and after social interactions. These results support prior work suggesting CA2 performs social functions and shed light on processes underlying CA2 responses to social stimuli.

Keywords

hippocampus, place cells, CA2, social cognition, social memory, social odor

1. Introduction

Social recognition memory, or the ability to recognize and remember conspecifics, facilitates social interactions that are essential for animals to survive. Rats can discriminate between familiar conspecifics (Husted and McKenna, 1966; Thor and Holloway, 1982) based on their distinct characteristics and olfactory signatures (Gheusi et al., 1997; Popik et al., 1991; Sawyer et al., 1984). Many recent studies have implicated hippocampal area CA2 in social memory (for a review, see Oliva, 2022). Lesioning or inactivating CA2 pyramidal neurons leads to impaired social recognition memory in mice (Hitti and Siegelbaum, 2014; Stevenson and Caldwell, 2014) and optogenetic silencing of CA2 activity shows that CA2 is crucial for encoding, consolidating, and recalling social memories (Meira et al., 2018). Moreover, CA2 neurons are uniquely enriched in receptors selective for a variety of social neuropeptides, including vasopressin and oxytocin (Lee et al., 2008; Lin et al., 2018; Pagani et al., 2015; Raam et al., 2017; Wersinger et al., 2002, 2008).

All subregions of the hippocampus contain place cells, neurons that are selectively activated in a particular location in space (their "place fields") (O'Keefe, 1976; O'Keefe and Dostrovsky, 1971). Place cells alter their firing in response to significant environmental changes in a process called "remapping" (for a review, see Colgin et al., 2008). During "global" remapping, a place cell's place field may appear, disappear, or change location. During "rate" remapping, a place cell's place field location does not move, but its in-field firing rate changes. Neurophysiological recordings of place cells have shown that place fields in hippocampal subregions CA1 and CA3 remain relatively stable in unchanging environments (Muller et al., 1987; Thompson and Best,

1990). In contrast, CA2 place cells show gradual changes in firing patterns over time (Alexander

et al., 2016; Mankin et al., 2015). CA2 place cells also appear to be highly sensitive to small

updates to familiar environments (Wintzer et al., 2014). Previous work has shown that a

significant proportion of CA2 place cells globally remap when a familiar conspecific is presented

(Alexander et al., 2016). This remapping in response to social stimuli may support the ability of

CA2 to encode social memories (Hitti and Siegelbaum, 2014). However, it remains unclear

whether CA2 place field changes that occur in response to the presentation of a conspecific rat

are caused by integration of multimodal social stimuli or primarily due to a specific sensory

modality associated with the social interaction.

Previous work has also shown that CA2 firing patterns during sleep are altered following a

social experience (Oliva et al., 2020). During awake rest or sleep, hippocampal place cells fire

during distinctive events in the hippocampal local field potential (LFP) known as sharp wave-

ripples. Place cells that were active during exploratory behaviors later reactivate during sharp

wave-ripples, and reactivated place cell ensembles are believed to support memory

consolidation (Ego-Stengel and Wilson, 2010; Ramadan et al., 2009; Wilson and McNaughton,

1994). CA2 place cells that represent a conspecific have been shown to reactivate during sharp

wave-ripples following a social experience (Oliva et al., 2020). Further, disrupting CA2 sharp

wave-ripples impairs social recognition memory (Oliva et al., 2020). Therefore, the reactivation

of CA2 place cells that encode a social experience may promote consolidation of social

memories (Oliva et al., 2023).

In this study, we recorded CA2 place cells in dorsal hippocampus of rats during free exploration

of a two-dimensional spatial environment and during subsequent rest periods. We compared

changes in CA2 place cell firing patterns across sessions in which different social or non-social

stimuli spanning various sensory modalities were presented. Significant global remapping was

observed in CA2 place cells when social odors were presented in the absence of a rat and

when a familiar rat was presented in a soiled home cage containing social odors (as in our prior

study; Alexander et al., 2016). However, no significant changes in CA2 place cell firing rate

maps were observed when a familiar rat was presented in a clean and relatively odorless cage

or when a non-social odor was presented. These results suggest that olfactory cues are the key

sensory component of social experiences that drives CA2 place cell remapping. Furthermore,

CA2 place cells that increased their firing rates in response to social odors preferentially

increased their firing rates in sharp wave-ripples during subsequent rest periods. These results

improve our understanding of how CA2 place cells code information related to social

interactions.

2. Materials and Methods

2.1.   Subjects

Six wild-type male Sprague-Dawley rats (Inotiv, USA) were used for this study. Rats were

between the ages of 4 and 10 months at the time of surgery. Before surgery, rats were double or

triple housed and were pre-trained to freely explore an open field enclosure. After surgery, rats

were singly housed in custom-built acrylic cages (40 cm x 40 cm x 40 cm) containing

enrichment material (wooden blocks, paper towel rolls, etc.) and maintained on a reverse light

cycle (light: 8 p.m. – 8 a.m.). Rats were housed next to their former cage mates after recovering

from surgery and throughout behavioral testing. Rats recovered from surgery for at least one

week before behavioral training resumed. All behavioral experiments were performed during the

dark cycle. In order to encourage spatial exploration, four of the six rats were placed on a food

deprivation regimen. All rats maintained ~98% of their free-feeding body weight throughout the

experiment. All experiments were conducted according to the guidelines of the United States

National Institutes of Health Guide for the Care and Use of Laboratory Animals and under a

protocol approved by the University of Texas at Austin Institutional Animal Care and Use

Committee.

2.2.   Surgery and tetrode positioning

Drives with 14 independently movable tetrodes were implanted in five of the rats. A drive with 21

independently movable tetrodes was implanted in one of the rats. Drives were implanted above

a relatively lateral part of right dorsal hippocampus (anterior-posterior -3.8 mm from bregma,

medial-lateral -3.0 mm from bregma). Because the diameter of the bundle of cannula

surrounding tetrodes was wider than the mediolateral extent of CA2, this placement ensured

that at least some of the tetrodes would hit CA2. To stabilize the recording drives, eleven bone

screws were affixed to the skull and covered in dental acrylic. Two of the screws were

connected to the recording drive ground. Prior to surgery, tetrodes were built from 17 μm

polyimide-coated platinum-iridium (90/10%) wire (California Fine Wire, Grover Beach, California,

USA). The tips of tetrodes designated for single-unit recording were plated with platinum to

reduce single-channel impedances to ~150 to 300 kOhms. All tetrodes were lowered ~1 mm on

the day of surgery. Thereafter, tetrodes were slowly lowered to the hippocampal pyramidal cell

body layer over the course of several weeks.

One tetrode was designated for use as a reference for differential recording. All four wires of this

tetrode were connected to a single channel on the electrode interface board. The differential

recording reference tetrode was placed in an electrically quiet area approximately 1 mm above

the hippocampus and adjusted as needed to ensure quiescence. The reference signal was

duplicated using an MDR-50 breakout board (Neuralynx, Bozeman, MT, USA) and recorded

continuously to ensure that unit activity or volume conducted signals of interest were not

detected. Another tetrode was placed in the apical dendritic layer of CA1 to monitor LFPs and

guide placement of the other tetrodes using estimated depth and electrophysiological hallmarks

of the hippocampus (for example, sharp wave-ripples).

2.3.   Data acquisition

Data were acquired using a Digital Lynx system and Cheetah recording software (Neuralynx,

Bozeman, MT, USA). The recording setup has been described in detail previously (Hsiao et al.,

2016; Zheng et al., 2016). Briefly, LFP signals from one randomly chosen channel within each

tetrode were continuously recorded at a 2000 Hz sampling rate and filtered in the 0.1–500 Hz

band. Input amplitude ranges were adjusted before each recording session to maximize

resolution without signal saturation. Input ranges for LFPs generally fell within ±2,000 to

±3,000 µV. To detect unit activity, all four channels within each tetrode were bandpass filtered

from 600 to 6,000 Hz. Spikes were detected when the filtered continuous signal on one or more

of the channels exceeded a threshold set daily by the experimenter, which ranged from 55–

65 µV. Detected events were acquired with a 32,000 Hz sampling rate for 1 ms. For both LFPs

and unit activity, signals were recorded differentially against a dedicated reference channel (see

"Surgery and tetrode positioning" section above).

Videos of rats' behavior were recorded through the Neuralynx system with a resolution of 720 ×

480 pixels and a frame rate of 29.97 frames/s. Rat position and head direction were tracked via

an array of red and green or red and blue light-emitting diodes (LEDs) on a HS-54 or HS-27

headstage (Neuralynx, Bozeman, MT, USA), respectively.

2.4.   Behavioral task

Rats were familiarized to an open field arena (1 m x 1 m with 0.5 m wall height) for a minimum

of three days before recording started. Rats freely explored the open field arena for four 20-

minute sessions per day, with 10-minute rest sessions preceding and following each active

exploration session. During active exploration, small pieces of sweetened cereal or cookies

were randomly scattered to encourage rats to explore the entirety of the arena. Rats had to

cover at least 60% of the arena across each of the four sessions for a day to be included for

further analysis. During each rest session, rats rested in a towel-lined, elevated flowerpot

outside of the arena. A plexiglass standard rat housing cage was placed in one corner of the

arena for all recording sessions. In the first and fourth sessions (A and A'), this cage contained

only clean bedding. In the middle two sessions (B and B') of the different experimental

conditions, the cage contained various types of social stimuli (Figure 1). In the Social Odor

condition, the soiled bedding from the former home cage of familiar rats (or a familiar rat) was

used. In the Visual + Odor condition, a familiar rat was presented in its home cage containing

soiled bedding. In the Visual condition, a familiar rat was placed in the stimulus cage with clean

bedding and a filter-top lid to minimize social odors. In the Mirror condition, a familiar rat was

placed in the stimulus cage with clean bedding and a filter-top lid, but the cage was lined with a

one-way mirror attachment. This one-way mirror attachment prevented the stimulus rat from

seeing the implanted rat, limiting visually driven reciprocal interactions between the two rats. In

the Non-social Odor condition, three drops of hexyl acetate (Sigma-Aldrich Cat# 108154), which

has a fruity smell, were applied to a wooden block which was placed underneath clean bedding

in the stimulus cage. In the Control condition, a cage containing only clean bedding was

presented in all four sessions.

2.5.   Histology and tetrode localization

Following recording, rats were perfused with 4% paraformaldehyde solution in phosphate-

buffered saline. Brains were cut coronally in 30 μm sections using a cryostat. For two rats, brain

sections that contained tetrode tracks were immunostained with a monoclonal antibody specific

for the CA2-expressed protein, STEP (Cell Signaling, 4817, 1:1000 dilution, raised in mouse).

Sections were then incubated in an anti-mouse secondary antibody and developed with 3,3-

diaminobenzidine (brown) (as in Alexander et al., 2016). Sections were then counter-stained

with cresyl-violet (purple/blue) to indicate cell bodies (Supplementary Figure 1). Brain sections

that did not contain tetrode tracks were stained with cresyl violet only. In order to locate CA2 in

the sections with tetrode tracks, we examined STEP staining in the cell body layer and identified

the portion of the hippocampus where the cell body layer increases in size to denote the

boundaries of CA2 (as in Lu et al., 2015). For four rats, brains were immunostained for the CA2

marker Purkinje Cell Protein 4 (PCP4) (Figure 2). For this immunostaining protocol, sections

were washed three times for 15 minutes in tris-buffered saline (TBS) followed by a 10-minute

water wash. Sections were then permeabilized and washed for 15 minutes in TBS containing

0.3% Triton-X followed by three 10-minute washes in TBS. Sections were blocked for 30

minutes in 10% normal goat serum in TBS. Sections were incubated overnight with rabbit anti-

PCP4 (1:200, Sigma-Aldrich Cat# HPA005792) diluted in TBS containing 0.05% Tween. The

next day, sections were washed twice for 10 minutes in TBS and incubated overnight with

secondary fluorescent antibody (in one rat: Alexa Flour™-488 anti-rabbit, Thermo Fisher

Scientific; in three rats: Alexa Flour™-555 anti-rabbit, Thermo Fisher Scientific). All washes and

incubations were performed at room temperature. Slides were mounted using DAPI

Fluoromount-G (Fisher Scientific). Tetrode recording sites were identified by comparing

locations across adjacent sections.

2.6.   Spike sorting and unit selection

Spike sorting was performed manually using graphical cluster-cutting software (MClust, A.D.

Redish, University of Minnesota, Minneapolis, Minnesota) run in MATLAB (Mathworks). Spikes

were sorted using two-dimensional representations of waveform properties (i.e., energy, peak,

and peak-to-valley difference) from four channels. A single unit was accepted for further analysis

if the associated cluster was well isolated from, and did not share spikes with, other clusters on

the same tetrode. Units were also required to have a minimum 1 ms refractory period. Units with

mean firing rates above 5 Hz were considered putative interneurons and excluded from further

analysis. In order to be considered active in the arena, a unit had to reach a peak firing rate of at

least 1 Hz. In order to be included in the sharp wave-ripple firing analysis, a unit had to have

valid clusters in both the active exploration and rest sessions. CA2 cell yields for each condition

are reported in Tables 1 and 2.

2.7.   Place cell remapping analyses

Methods used to create firing rate maps for each single unit were based on methods used in our

prior study (Alexander et al., 2016; example rate maps for the present study shown in Figure 3).

First, the arena was divided into 4 cm^2 bins. The number of spikes that occurred within each bin

was divided by the time spent in that bin to determine the firing rate. Only spikes that occurred

while the rat was traveling 5 cm/s or faster were included. The rate map was smoothed with a

two-dimensional Gaussian kernel (standard deviation = 6 cm). To determine if a place cell

remapped during presentation of a social stimulus, a Pearson correlation coefficient *R* was

calculated for each unit between pairs of rate maps from control and social stimuli sessions (i.e.,

A-B, B-B', B'-A', A-B', and A-A' session pairs, see Figure 1). To determine if spatial correlation

coefficients differed across session pairs and conditions (i.e., Social Odor, Visual + Odor, Visual,

Mirror, Non-social Odor, Control), we used a generalized linear mixed model statistical analysis

(IBM SPSS Statistics, version 29.0.2.0). Condition and session pair were fixed factors, session

pairs were repeated measures within rats, and multiple place cells were nested within rats. A

condition by session pair interaction effect was also included in the model to determine whether

differences between session pairs varied across conditions. When a significant effect was

observed, post-hoc pairwise comparisons were performed with Bonferroni correction for multiple

comparisons. The estimated mean spatial correlations for each condition and session pair from

the generalized linear mixed model are shown in Figure 4A. The estimated mean spatial

correlations for each condition, collapsed across session pairs, are shown in Figure 4B.

Individual spatial correlation values for each cell are shown for each session pair and each

condition in Figure 4C-G.

To quantify potential rate remapping across the various conditions, we computed a rate overlap

measure across a pair of sessions for each cell. The rate overlap was defined as the ratio of the

mean in-field firing rate of a place cell in each session in a session pair, with the smaller of the

two firing rates in the numerator. To determine if rate overlap values differed across session

pairs and conditions (i.e., Social Odor, Visual + Odor, Visual, Mirror, Non-social Odor, Control),

we used a generalized linear mixed model statistical analysis (IBM SPSS Statistics, version

29.0.2.0). Condition and session pair were fixed factors, session pairs were repeated measures

within rats, and multiple place cells were nested within rats. A condition by session pair

interaction effect was also included in the model to determine whether differences between

session pairs varied across conditions. The estimated mean rate overlap values for each

condition and session pair from the generalized linear mixed model are shown in

Supplementary Figure 2A. The estimated mean rate overlap values for each condition,

collapsed across session pairs, are shown in Supplementary Figure 2B. Individual rate overlap

values for each cell are shown for each session pair and each condition in Supplementary

Figure 2C-G.

To determine if positions of place fields coherently shifted during remapping, we first identified

cells that remapped and sorted them into "turn on", "turn off", and "field shift" categories

(Supplementary Figure 3). Because place cells were classified as active if they reached a peak

firing rate of at least 1 Hz, "turn on" cells were defined as cells that had a peak firing rate less

than 1 Hz in Session A and greater than 1 Hz in Session B. We then calculated the distance

between the position of the peak firing rate of the cell in Session B and the stimulus cage for

"turn on" cells. "Turn off" cells were defined as cells that had a peak firing rate above 1 Hz in

Session A and below 1 Hz in Session B. The distance between the position of the peak firing

rate of the cell in Session A and the stimulus cage was calculated for "turn off" cells. Cells were

identified as "field shift" cells if they reached a peak firing rate above 1 Hz in Sessions A and B

but showed unstable place field locations. We assessed place field stability as follows, using

criteria defined in a prior study (Widloski and Foster 2022). Cell IDs were randomly shuffled

1000 times within each condition and rat. Spatial correlation coefficients were then calculated

for a given unit from session A and each shuffled rate map from session B. A place field was

considered stable between sessions A and B if the spatial correlation coefficient exceeded the

95th percentile of the shuffled distribution. For "field shift" cells, we calculated the distances

between the stimulus cage and the positions of the peak firing rates of the cell in Sessions A

and B. To determine if a cell's place field moved closer to the stimulus cage when stimuli were

presented, we calculated the difference between the distance estimates from Sessions A and B

for "field shift" cells. We used the Shapiro-Wilk test of normality to determine if the distribution of

distances was skewed within each condition (IBM SPSS Statistics, version 29.0.2.0). We

additionally used an independent samples Kruskal-Wallis test to assess the extent to which

distributions differed across conditions (IBM SPSS Statistics, version 29.0.2.0).

To quantify how selective a unit was for session A vs session B, we calculated a selectivity index

for each unit, as in Hwaun and Colgin (2019). The selectivity index was defined as $(\mu_B -$

$\mu_A)/(\mu_B + \mu_A)$, where μ_A was the mean firing rate in session A and μ_B was the mean firing rate in

session B. A value of -1 would indicate that a unit was exclusively active in session A, while a

value of 1 would indicate that a unit was exclusively active in session B.

2.8.   Place cell properties

Spatial information was calculated as previously described (Skaggs et al., 1996) using the

following formula:

260
$$\text{Spatial information} = \sum_i P_i \frac{\lambda_i}{\lambda} log2 \frac{\lambda_i}{\lambda},$$

where i = the spatial bin index, P_i = the probability of the rat being in the ith bin, λ_i = the mean

firing rate in the ith bin, and λ = the mean firing rate of the cell.

Place fields were separately identified for each cell in each session. Potential place fields were

identified as contiguous bins of the smoothed rate map (see "*Place cell remapping analyses*"

section) that exceeded 50% of the peak firing rate in that session. In order to be included for

further analysis, the peak firing rate within a place field had to exceed 1 Hz, and the size of the

place field had to exceed 10 cm^2. To test if place field properties differed across conditions, we

used a generalized linear mixed model statistical analysis (IBM SPSS Statistics, version

29.0.2.0). Multiple place cells were nested within rats, and sessions were repeated measures. A

condition by session interaction effect was included in the model to determine whether

differences between session pairs varied across conditions.

2.9. Exploration time

To investigate exploration of the stimulus cage, the arena was divided into 4 cm^2 bins, as in a

prior study (Zhu et al., 2023). The time spent in each bin during the first two minutes of every

session was determined for each day. These exploration maps were then smoothed with a two-

dimensional Gaussian kernel (standard deviation = 4 cm). Maps were averaged within and then

across rats and plotted as a heat map for each condition (Figure 5A). The time spent within 12

cm of the cage was then calculated. A generalized linear mixed model analysis (IBM SPSS

Statistics, V 29.0.2.0) was used to compare time spent investigating the stimulus cage across

different sessions and conditions, with conditions and sessions as fixed factors. Conditions were

repeated on different days within rats, with different days included as random factors within rats.

Sessions were repeated measures within conditions. Post-hoc tests were performed to compare

exploration times across conditions for the session of interest (i.e., Session B), using the

Bonferroni correction for multiple comparisons.

2.10.   Sharp wave-ripple detection and analysis

Sharp wave-ripples were detected from recordings obtained while rats rested in an elevated,

towel-lined flowerpot outside of the arena. Detection and analysis methods were similar to

previously published methods (Hwaun and Colgin, 2019). The LFP recorded on all tetrodes that

had CA2 place cells was band-pass filtered between 150 and 250 Hz. A Hilbert transform was

performed on the filtered LFP, and the absolute value of this signal was smoothed with a

Gaussian kernel (standard deviation = 25 ms). Potential sharp wave-ripple events were

detected when the signal exceeded at least 5 standard deviations above the mean and bounded

by crossings of the mean. Overlapping events were combined across tetrodes, so the time

interval of identified events could extend beyond sharp wave-ripples detected on a single

tetrode (as in Hwaun and Colgin, 2019; Karlsson and Frank, 2009). Potential sharp wave-ripple

events were kept for further analysis if they were between 50 and 500 ms in duration.

We then obtained the firing rate for each CA2 place cell around detected sharp wave-ripple

events by constructing a spike raster with a bin size of 1 ms and smoothing the raster with a

Gaussian kernel (standard deviation = 5 ms). To account for baseline firing rate differences

among individual cells, the firing rate was normalized by the average firing rate 400-100 ms

before ripple onset (Hwaun and Colgin, 2019). The peak normalized firing rate was obtained for

each unit by taking the maximum normalized firing rate after sharp wave-ripple onset. Because

a broad range of peak normalized firing rates was observed, we rescaled the distribution using a

$\log_{10}$ transformation of the peak normalized firing rates. To determine the extent to which sharp

wave-ripple-associated firing rates of individual cells were altered by presentation of a social

stimulus, we computed the difference between the peak normalized firing rates during the rest

periods after sessions A and B (i.e., peak normalized firing rate after session B - peak

normalized firing rate after session A) for the Control and Social Odor conditions. A positive

value indicates that a cell had higher sharp wave-ripple-associated firing in the rest session

following session B, and a negative value indicates that a cell had higher sharp wave-ripple-

associated firing in the rest session following session A. We used multiple linear regression

(IBM SPSS Statistics, V29.0.2.0) to assess the effects of session selectivity (as measured using

the selectivity index, see *Place cell remapping analyses* section) and experimental condition on

the difference in sharp wave-ripple-associated firing between rest sessions (as in Hwaun and

Colgin, 2019). The regression analysis was only performed for conditions that had more than

100 active cells during both run and rest sessions (Control condition: n = 130 cells from 5 rats,

Social Odor condition: n = 132 cells from 5 rats).

2.11. Data and Code Availability

MATLAB (Mathworks) scripts were custom written for the analyses in this study, based on

algorithms that have been used in prior studies, as described above. Scripts and data are

available upon request.

3. Results

In our prior study of CA2 place cells (Alexander et al., 2016), we compared CA2 place cell

responses to a rat in a soiled home cage and responses to the presentation of an object that

resembled a rat, namely a stuffed toy rat. However, the stuffed toy lacked some visual

components that occur during interactions with a live rat (e.g., motion content). Also, the toy rat

was presented in a clean cage and thus lacked the olfactory components of a social experience.

The major goal of the present study was to determine the extent to which different sensory

modalities associated with a social experience drive CA2 place cell remapping to social stimuli.

We employed an open field exploration behavior paradigm (see "*Behavioral task"* section of

Materials and Methods) in which different types of stimuli were presented across sessions for

various experimental conditions (i.e., Social Odor, Visual + Odor, Visual, Mirror, Non-social

Odor) and compared to a Control condition in which no stimuli were presented (Figure 1).

3.1. A significant proportion of CA2 place cells globally remapped to social odor stimuli

The olfactory system is highly important for social recognition and social behaviors in rodents

(Oettl and Kelsch, 2018). Therefore, we hypothesized that olfactory cues would be a key

sensory component of CA2 place cell responses to social stimuli. In the Social Odor condition,

rats explored an environment in which social odors were presented, but another rat was not

present. This condition isolated the olfactory content of a social interaction and preserved the

ethological relevance of the presented stimuli. Example CA2 place cell firing rate maps during

presentation of social odors (Figure 3B) show that a subset of cells changed the location of their

firing fields, or globally remapped, when social odors were presented. To quantify global

remapping in CA2 place cells, we compared firing rate map changes in cells recorded in the

Social Odor condition (145 cells in 6 rats, Table 1) to firing rate map changes in cells recorded in

the other conditions (see Figure 3 for example rate maps for all conditions and Table 1 for cell

yields). We calculated the spatial correlation between rate maps from pairs of sessions for each

cell (Figure 4). We found that spatial correlation values for the Social Odor condition were

significantly lower than the Control, Visual, Mirror, and Non-social Odor conditions (generalized

linear mixed model, no significant interaction between session pair and condition, $F_{(20, 3250)}$ =

0.843, $p = 0.662$ (Figure 4A); significant main effect of condition, $F_{(5, 3250)}$ = 12.7, $p < 0.001$

(Figure 4B); significant differences in post-hoc tests for Social Odor vs. Control ($t_{(3250)}$ = 6.3, p

< 0.001), Social Odor vs. Visual ($t_{(3250)}$ = 4.0, p = 0.001), Social Odor vs. Mirror ($t_{(3250)}$ = 3.6,

p = 0.003), and Social Odor vs. Non-social Odor ($t_{(3250)}$ = 2.8, p = 0.047)). In agreement with

our prior findings, CA2 place cells also showed significant global remapping when a familiar

conspecific rat was presented in a soiled home cage containing social odors (Visual + Odor

condition, Figures 3C and 4, significant differences in post-hoc tests for Visual + Odor vs.

Control (t(3250) = 6.4, p < 0.001), Visual + Odor vs. Visual (t(3250) = 4.5, p < 0.001), Visual +

Odor vs. Mirror (t(3250) = 4.1, p = 0.001), and Visual + Odor vs. Non-social Odor (t(3250) = 3.5,

p = 0.004)). In contrast, significant global remapping was not observed when a stimulus rat was

presented in a clean cage with a filter top, a condition that maintained all visual components of

social interactions and minimized social odors (Visual condition, Figures 3D and 4, no significant

differences in post-hoc test for Visual vs. Control (t(3250) = 1.8, p = 0.411)). Similarly, CA2 place

cell firing rate maps did not change significantly when a stimulus rat was presented in a clean

cage with a filter top and a one-way mirror to prevent visually mediated reciprocal interactions

between rats (Mirror condition, Figures 3E and 4, no significant differences in post-hoc test for

Mirror vs. Control (t(3250) = 2.3, p = 0.166)). To test whether remapping of CA2 place cell

responses was specific to social odors, we presented a non-social odor in the stimulus cage.

We found that presentation of a non-social odor did not significantly change CA2 place cell firing

rate maps (Non-social Odor condition, Figures 3F and 4, no significant differences in post-hoc

test for Non-social Odor vs. Control (t(3250) = 1.4, p = 0.770)) (see Table 3 for all pairwise

comparisons). Together, these results show that a subset of CA2 place cells globally remapped

when social odors were presented but not during interactions with another rat when social odors

were minimized.

While non-odorant social conditions and non-social odors were not found to induce global

remapping in CA2 place cells, it remains possible that these conditions induced rate remapping

in CA2. To test this possibility, we compared rate overlap values across conditions

(Supplementary Figure 2). Consistent with previous work (Alexander et al., 2016), we found that

none of the social conditions induced significant rate remapping in CA2 place cells. Further, the

presentation of a non-social odor did not induce significant rate remapping in CA2 place cells

(generalized linear mixed model, no significant interaction between session pair and condition,

$F(20,3213) = 0.7$, $p = 0.803$; no main effect of condition, $F(5,3213) = 2.0$, $p = 0.076$)).

3.2.   No coherent movement of place fields during remapping

CA1 place fields have been reported to shift with salient stimuli (Fenton et al., 2000; O'Keefe

and Conway, 1978) or move toward locations of salient rewarding stimuli (Breese et al., 1989;

Fenton et al., 2000). However, our previous work showed that fields of CA2 place cells in rats

did not move closer to a social stimulus during social remapping (Alexander et al., 2016).

Conversely, recordings of CA2 place cells in mice showed that a subpopulation of CA2 place

cells remapped when a conspecific mouse was moved to a new location of the arena, and this

subpopulation of "social-remapping" cells had fields significantly closer to the stimulus mouse

than fields of non-remapping ("social-invariant") cells (Oliva et al. 2020). To test if CA2 place

cells moved coherently towards or away from the stimulus cage in our experiment, we first

identified place cells that remapped between sessions A and B. These cells were identified as

cells that either gained a place field ("turn on" cells), lost their place field ("turn off" cells), or

shifted their place field ("field shift" cells). For "turn on" cells, we calculated the distance between

the cell's peak firing rate position in the session in which stimuli were presented (Session B) and

the position of the stimulus cage (Supplementary Figure 3A). Similarly, we calculated the

distance between the peak firing rate position and the position of the stimulus cage in Session A

for the "turn off" cells (Supplementary Figure 3B). If place cells preferentially gained or lost fields

close to the stimulus cage, we would have expected skewed distributions towards low values,

which we did not observe. However, definitive conclusions cannot be drawn from this

observation given the low number of cells that met the criteria for classification as "turn on" or

"turn off" cells.

To determine whether CA2 place cells preferentially shifted their place fields closer to the

stimulus cage, we estimated the distance between the stimulus cage and cells' peak firing rate

positions in Sessions A and B and then calculated the difference between these two distance

estimates (Supplementary Figure 3C). If place fields shifted closer to the cage, we would have

expected values to skew negatively. However, differences between place field distances from

the cage between Sessions A and B were normally distributed for all conditions (Shapiro-Wilk

test of normality, Control: $W(20) = 1.0$, $p = 0.717$; Social Odor: $W(30) = 1.0$, $p = 0.339$; Visual +

Odor: $W(16) = 0.9$, $p = 0.135$; Visual: $W(15) = 0.9$, $p = 0.142$; Mirror: $W(18) = 0.9$, $p = 0.346$;

Non-social Odor: $W(4) = 0.880$; $p = 0.340$). Further, the distributions of distance shifts did not

differ significantly across conditions (Kruskal-Wallis test, $H(5) = 6.3$, $p = 0.278$). Therefore,

consistent with our previous findings (Alexander et al., 2016), no coherent movement of place

fields toward or away from the stimulus cage was observed. However, our experimental design

did not allow us to identify "social-remapping" and "social-invariant" place cells as was done in a

previous study (Oliva et al. 2020). That is, social-remapping cells in the previous study in mice

were defined as place cells that displayed place fields that shifted in a manner that preserved a

fixed distance from a stimulus mouse, whereas place fields of "social-invariant" cells maintained

a fixed location in the environment. In the present experiment, only one stimulus was presented

in each session, and the location where it was presented did not change across sessions. It is

possible that the present sample of cells contained a mixture of social-remapping and social-

invariant place cells and that potential shifts in field relative to the social stimulus were masked

as a result. Alternatively, the lack of a preferential shift in place fields near the social stimulus

may be due to the nature of odor stimuli, as airborne odorants diffuse across widespread

locations within an environment. This may allow a rat to perceive odors regardless of their exact

location in the arena. It is also possible that the introduction of social odors to an environment

was sufficient to change its context as a whole. This generalizability may help rats to associate

conspecifics with larger areas of an environment.

3.3. CA2 place cell properties were not modulated by social stimuli

In addition to remapping, we examined general CA2 place cell properties during presentation of

social and non-social stimuli. We found that spatial information was unaffected by the inclusion

of social stimuli or a non-social odor in an environment (Supplementary Figure 4A; generalized

linear mixed model, no significant interaction between session and condition ($F_{(23,2596)}$ = 1.0,

p = 0.369)). We also saw no significant changes in peak firing rates (Supplementary Figure 4B;

generalized linear mixed model, no significant interaction between session and condition

($F_{(23,2413)}$ = 1.0, p = 0.478)), mean in-field firing rates (Supplementary Figure 4C; generalized

linear mixed model, no significant interaction between session and condition ($F_{(23,2413)}$ = 1.0,

p = 0.477)), or the number of place fields per cell (Supplementary Figure 4D; generalized linear

mixed model, no significant interaction between session and condition ($F_{(23, 2596)}$ = 0.9, p =

0.583)). These results suggest that CA2 place cells represented spatial locations similarly in

social and non-social conditions.

3.4. Rats explored all modalities of social stimuli

It is possible that the lack of remapping to non-olfactory social stimuli may be due to inadequate

exploration of these stimuli. To test this possibility, we examined rats' stimuli exploration time

during the first 2 minutes of each exploration session for each experimental condition (Figure 5).

We found that stimulus cage exploration time during session B of most social conditions (i.e.,

Social Odor, Visual, and Mirror) significantly increased compared to session B of the Control

condition (generalized linear mixed model, interaction between session and condition ($F_{(15,284)}$

= 2.3, p = 0.004); post-hoc tests, session B: significant difference between Control vs. Social

Odor ($t_{(284)}$ = 3.0, p = 0.037); Control vs. Visual ($t_{(284)}$ = 3.5, p = 0.009); and Control vs. Mirror

($t_{(284)}$ = 3.4, p = 0.010); no significant difference between Control vs. Visual + Odor ($t_{(284)}$ =

2.2, p = 0.325); no significant difference between Control vs. Non-social Odor ($t_{(284)}$ = 1.2, p >

0.999)). It is important to note that stimulus cage exploration times significantly increased in the

social conditions in which CA2 place cells did not show significant remapping (i.e., Visual and

Mirror conditions), suggesting that the lack of place cell remapping in these conditions was not

due to reduced exploration of social stimuli. Note also that exploration times did not significantly

differ between Visual + Odor and Control conditions, although significant remapping was

observed in the Visual + Odor condition. The insignificant difference in exploration times may be

due to the lower number of rats in the Visual + Odor condition (i.e., 4 rats in Visual + Odor

condition compared to 5-6 rats in other social conditions, see Table 1). Similarly, exploration

times also did not differ between Non-social Odor and Control conditions, perhaps due to the

lower number of rats in the Non-social Odor condition (i.e., only 3 rats, see Table 1).

Alternatively, the lack of ethological relevance of the hexyl acetate odor may have contributed to

a lack of increased exploration of the stimulus cage during presentation of the Non-social Odor.

3.5.   Place cells that responded to social odors were preferentially recruited into sharp

wave-ripples

CA2 place cells that encode a social experience with a novel conspecific have been shown to

reactivate during sharp wave-ripples in mice (Oliva et al., 2020). Therefore, we aimed to

determine whether presentation of social odors alone was sufficient to drive preferential sharp

wave-ripple-associated reactivation of CA2 place cells that responded to social odors. We used

a selectivity index (see Materials and Methods) to classify cells according to their firing

preferences during the Control and Social Odor conditions. The selectivity index defined the

extent to which CA2 place cells fired during session A (i.e., the session in which an empty cage

was presented) compared to session B (i.e., the session in which a cage containing social odors

was presented in the Social Odor condition). We then estimated the normalized average peak

firing rate of each cell during sharp wave-ripples in rest sessions after session A and session B

(see Materials and Methods). Next, we used the difference between a cell's peak normalized

firing rate during the rest session after session B and the rest session after session A as a

measure of the extent to which a cell changed its sharp wave-ripple-associated firing after

presentation of social odors. CA2 place cells show unstable firing patterns in unchanged

environments over time (Mankin et al., 2015). Thus, we performed the same analysis for CA2

place cells recorded in the Control condition to ensure that changes in sharp wave-ripple-

associated firing were not explained by the effect of time. We found that a cell's selectivity for

social stimuli in active exploratory sessions correlated with sharp wave-ripple-associated peak

firing rate changes in the Social Odor condition but not the Control condition (Figure 6; multiple

linear regression, $F(3,243) = 2.661$, $p = 0.049$; interaction between selectivity index and

condition, $t = 2.448$, $p = 0.015$, Pearson correlation between selectivity index and firing rate

changes for the Control condition $R = -0.102$, $p = 0.266$, Pearson correlation between selectivity

index and firing rate changes for the Social Odor condition $R = 0.227$, $p = 0.010$). Specifically,

cells that increased their firing rates during exploration of a social odor preferentially increased

their sharp wave-ripple-associated firing rates during rest following social exploration. These

results suggest that the presentation of a social odor alone is sufficient to preferentially

reactivate CA2 place cells that code social stimuli.

4. Discussion

An increasing number of lesion, genetic, optogenetic, and pharmacogenetic manipulation

studies have implicated CA2 in social memory processing (Hitti and Siegelbaum, 2014; Meira et

al., 2018; Oliva et al., 2020; Smith et al., 2016; Stevenson and Caldwell, 2014). In this study, we

expanded on prior results showing that CA2 place cells remap during a social interaction

(Alexander et al., 2016) by investigating how various sensory modalities of social stimuli

contribute to this response. Here, we show that CA2 place cell remapping is primarily driven by

the olfactory component of social stimuli. Exposure to social odors in the absence of a rat was

sufficient to induce remapping in CA2 place cells. CA2 place cells did not respond to social

interactions with another rat when social odors were reduced. The lack of remapping when

social odors were reduced was not explained by inadequate exploration of social stimuli. Lastly,

CA2 place cells that showed increased firing during exploration of social odors were

preferentially reactivated during sharp wave-ripples in subsequent rest. Our findings support the

notion that CA2 place cells may integrate social olfactory information with contextual information

to encode and consolidate social episodic memories (Oliva, 2022; Oliva et al., 2023). However,

future experiments that selectively silence CA2 are needed to support a direct link between CA2

place cell activity and behavioral performance on memory tasks requiring association of social

odors with specific contexts. Also, while social experiences did not induce significant global

remapping in CA2 place cells when social odors were lacking, the possibility remains that other

sensory modalities of a social experience are processed via different pathways and brain

structures and thereby contribute to social recognition and social memory processes.

The sensory modalities that drive social recognition vary across species. While humans depend

predominantly on auditory and visual cues for social recognition, rodents and other mammals

rely heavily on chemosensory cues in the form of olfactory or pheromonal signals (Popik and

van Ree, 1998). Indeed, olfactory bulb lesions or chemically induced anosmia impair individual

recognition in rats (Dantzer et al., 1990; Popik et al., 1991). The complex mix of chemosensory

signals embedded in rat urine convey information about a rat's social status and identity (Hurst,

2005). CA2 neurons differentially respond to urine from different conspecifics in mice (Hassan et

al., 2023). The ability of CA2 to respond to social odors in the absence of conspecifics, as seen

in our study, may allow rats to discern territory and identity information, which may drive social

behavior even when other rats are not present. The precise mechanisms underlying social odor

detection in CA2 neurons are unknown but may be related to the distinctive neurochemical and

structural features of CA2 neurons.

CA2 neurons contain an abundance of neuropeptide receptors, including oxytocin receptors

(OXTR) and vasopressin receptors (AVP1-b) (Mitre et al., 2016; Vaccari et al., 1998; Young and

Song, 2020). Genetic or pharmacological ablation of CA2 OXTRs impairs the persistence of

long-term social recognition memory and the ability to discriminate between social stimuli in

mice (Lee et al., 2008; Lin et al., 2018; Raam et al., 2017). AVP1-b activation in CA2 has been

associated with social aggression (Pagani et al., 2015) and the enhancement of social memory

(Smith et al., 2016). Long-range axonal projections from the paraventricular nucleus of the

hypothalamus (PVN) may be the source of neuropeptide release onto CA2 neurons during

social interactions. The PVN receives input from the olfactory system (Guevara-Aguilar et al.,

1988) and directly innervates CA2 (Cui et al., 2013; Zhang and Hernández, 2013).

Several other brain regions that are involved in social processing send direct inputs to CA2,

including the hypothalamic supramammillary nucleus (SuM) and the lateral entorhinal cortex

(LEC). SuM projections to CA2 are preferentially activated by novel social encounters (Chen et

al., 2020), making SuM inputs an unlikely candidate to induce CA2 place cell responses to a

familiar social odor. Conversely, neuronal activity in the LEC showed similar increases during

social exploration for both novel and familiar conspecifics (Lopez-Rojas et al., 2022). Recent

studies have suggested that the direct projection from LEC to CA2 is essential for social

memory (Dang et al., 2022; Lopez-Rojas et al., 2022). Whether LEC input to CA2 is selectively

enhanced during exploration of a social odor alone remains to be tested, but a more general

involvement of the LEC in odor processing is well documented (Igarashi et al., 2012; Kerr et al.,

2007).

CA2 also receives input from granule cells of the dentate gyrus (DG). Both mature and adult-

born granule cells directly synapse onto CA2 neurons via mossy fiber projections (Kohara et al.,

2014; Llorens-Martín et al., 2015; Laham et al., 2024). The DG is important for associating

odors with larger contexts in discrimination tests (Morris et al. 2012). In addition, recent work

has shown that adult-born granule cell projections to CA2 are necessary for the retrieval of

social memories formed during development (Laham et al., 2024). The exact relationship

between the DG and CA2 in social odor processing is currently unknown and warrants further

investigation.

Consistent with previous work, our results show that CA2 place cells that remap in response to

a familiar social odor do not return to their original activity, despite the removal of the social odor

in the last session (Alexander et al., 2016). It is possible that the persistent change in CA2

activity is a result of lingering social odors in the arena. Recording CA2 place cells for longer

than 20 minutes after the removal of the odor would address if CA2 place cell activity eventually

returns to baseline, although this would be complicated by the effect of time on CA2 place cells

(Mankin et al., 2015). However, this persistent change in CA2 place cell activity may also

suggest the involvement of synaptic plasticity. Interestingly, CA2 synapses are resistant to

standard long-term potentiation (LTP) compared to other hippocampal subfields (Chang et al.,

2007; Zhao et al., 2007). How, then, are long-lasting changes occurring in CA2 place cells? One

hypothesis is that the release of social neuropeptides, such as vasopressin and oxytocin, can

promote potentiation of CA2 neurons (Lin et al., 2018; Pagani et al., 2015; Tirko et al., 2018).

Selective activation of OXTRs or AVP1-b *in vitro* robustly depolarizes CA2 pyramidal neurons

and lowers the threshold for LTP at excitatory synapses (Dang et al., 2022; Tirko et al., 2018).

As a result, neuropeptide receptor activation may refine the responsiveness of CA2 pyramidal

neurons to synaptic input from upstream structures, thus creating appropriate conditions for

synaptic plasticity.

A question remains of how social information coded by CA2 is transmitted to downstream

regions that control behavior. The main output of dorsal CA2 is dorsal CA1. However, at the

single cell level, dorsal CA1 place cell activity was unaffected by social interactions (Alexander

et al., 2016). Dorsal CA2 neurons also project to ventral CA1, a region that is essential for social

memory (Okuyama et al., 2016). Recent work has shown that ventral CA1 place cells did not

remap to a social stimulus (Wu et al, 2023). However, ventral CA1 place cells became

significantly more spatially selective in an environment that contained a social stimulus

compared to an empty environment (Wu et al. 2023). Further, the activity of a subset of ventral

CA1 neurons was modulated by the presence of a conspecific (Rao et al. 2019; Wu et al. 2023).

Inhibition of dorsal CA2 projections to ventral CA1 impairs social memory (Meira et al., 2018;

Tsai et al., 2022), raising the possibility that social remapping in dorsal CA2 induces the

alterations in neuronal firing in ventral CA1 in response to social stimuli.

Interestingly, recent work has shown that sharp wave-ripples that originate in dorsal CA2 are

important for social memory consolidation (Oliva et al., 2016, 2020). While a portion of the sharp

wave-ripples initiated in dorsal CA2 propagate to other subregions of the dorsal hippocampus,

some do not propagate to dorsal CA1 or dorsal CA3 (Oliva et al., 2016). Given that individual

place cells in dorsal CA1 did not remap to a social stimulus (Alexander et al., 2016), it is

possible that the reactivation of CA2 place cells that respond to a social odor may be limited to

sharp wave-ripples that do not propagate to dorsal CA1. Instead, the function of these sharp

wave-ripples may be to propagate information to ventral CA1 in order to consolidate memories

of a social experience (Meira et al., 2018). Consistent with this idea, the rate of sharp wave-

ripples increased in ventral CA1 when conspecifics were present (Rao et al., 2019). A future

study examining the reactivation of dorsal CA2 place cells that respond to social stimuli while

simultaneously recording place cells in dorsal and ventral CA1 will be essential for our

understanding of how reactivation of CA2 place cells during sharp wave-ripples supports social

memory consolidation.

While hippocampal place cells are able to represent position and respond to social stimuli such

as odors at the single cell level, as evidenced by place cell remapping, population-level activity

in the hippocampus may incorporate the activity of both place cells and non-spatially modulated

cells to encode other features of an environment and ongoing behavior (Meshulam et al., 2017;

Nagelhaus et al., 2023; Stefanini et al., 2020). The relatively low CA2 cell yields per rat in our

data (see Table 1) made it difficult to examine population-level activity in CA2 across conditions.

While we did not see remapping at the single cell level for our social conditions that minimized

social odors and for our non-social odor condition, the possibility remains that these conditions

induced population-level changes in CA2 activity.

Another important consideration for this study is that rats were singly housed after surgery to

reduce trauma to the surgical site and protect drive implants. Social isolation can impair the

persistence of social recognition memory in both mice and rats (Kogan et al. 2000; Shahar-Gold

et al. 2013). This impairment may be due to a de-coupling of the olfactory bulb and dorsal

hippocampus in mice (Almeida-Santos et al. 2019). We attempted to minimize the effects of

social isolation by housing familiar conspecifics in adjacent cages, but we cannot rule out

unintended consequences of social isolation on CA2 responses to social stimuli.

Another important question left unanswered is the extent to which CA2 is specialized for social

memory. The present results show that a social odor, but not a non-social odor, was sufficient to

induce CA2 place cell remapping. However, due to the differences in ethological relevance

between social and non-social odors in this study, it remains unclear whether non-social odors

of comparable salience or ethological relevance would affect CA2 place cell activity. Future

studies examining the activity of CA2 place cells in response to ethologically relevant non-social

odors, such as predator urine, will help determine whether CA2 integrates other salient odors

into representations of space or whether the role of CA2 is specific to social information.

The current study shows how presentation of a social odor alone provides a well-controlled

social stimulus to induce changes in place cell activity in dorsal CA2. Use of a social odor can

eliminate confounds that could be introduced by the presence of a conspecific, such as

variations in locomotor behavior, social interactions, and ultrasonic vocalizations. Presentation

of a social odor thereby provides a valuable paradigm for assessing CA2 physiology in rodent

models of diseases and disorders, such as autism spectrum disorders, involving aberrant

processing of social stimuli.

Table 1. Number of CA2 place cells recorded in each rat and each condition during active

exploration sessions.

Animal ID	Control	Social Odor	Visual + Odor	Visual	Mirror	Non-social Odor
Rat 117	7	5	0	7	6	0
Rat 122	51	32	0	38	49	0
Rat 165	34	56	26	26	35	0
Rat 256	28	13	13	13	21	11
Rat 391	15	4	5	6	5	6
Rat 418	20	35	41	18	0	29
Total	**155**	**145**	**85**	**109**	**116**	**46**

660   Table 2. Number of CA2 place cells recorded in each rat and each condition during rest periods.

Animal ID	Control	Social Odor	Visual + Odor	Visual	Mirror	Non-social Odor
Rat 117	0	0	0	1	0	0
Rat 122	52	31	0	2	18	0
Rat 165	26	58	23	31	38	0
Rat 256	18	10	8	13	0	0
Rat 391	14	4	5	7	5	6
Rat 418	20	29	35	14	0	29
Total	130	132	71	68	61	35

662   Table 3. Pairwise comparisons of spatial correlations between conditions.

Pairwise contrast	t	df	Adjusted p-value
Control vs Social Odor	6.3	3250	4.265E-9***
Control vs Visual	1.8	3250	0.411
Control vs Visual + Odor	6.4	3250	2.421E-9***
Control vs Mirror	2.3	3250	0.166
Control vs Non-social Odor	1.4	3250	0.770
Social Odor vs Visual	-4.0	3250	0.001**
Social Odor vs Visual + Odor	1.2	3250	0.935
Social Odor vs Mirror	-3.6	3250	0.003**
Social Odor vs Non-social Odor	-2.8	3250	0.047*
Visual + Odor vs Visual	-4.5	3250	8.613E-5***
Visual + Odor vs Mirror	-4.1	3250	0.001**
Visual + Odor vs Non-social Odor	-3.5	3250	0.004**
Visual vs Mirror	0.4	3250	1.0
Visual vs Non-social Odor	0.1	3250	1.0
Mirror – Non-social Odor	-0.1	3250	1.0

CRediT authorship contribution statement

Emma Robson: Formal analysis, investigation, writing – original draft, visualization. **Margaret**

M. Donahue: Formal analysis, investigation, writing – original draft, visualization, funding

acquisition. **Alexandra J. Mably:** Conceptualization, methodology, formal analysis,

investigation. **Peyton G. Demetrovich:** Investigation. **Lauren T. Hewitt:** Investigation. **Laura**

Lee Colgin: Conceptualization, methodology, resources, writing – review and editing,

supervision, project administration, funding acquisition.

Acknowledgements

The authors thank Isabella Lee, Misty Hill, Ayomide Akinsooto, Sirisha Dhavala, and Alexa

Hassien for technical assistance with this project and Chenguang Zheng and Ernie Hwaun for

providing MATLAB code for some of the analyses. The authors acknowledge the Texas

Advanced Computing Center (TACC) at The University of Texas at Austin for providing data

storage resources that have contributed to the research described within this article. URL:

http://www.tacc.utexas.edu

Funding

This research was supported by the Department of Defense (grant number W81XWH1810314)

and the National Institutes of Health (grant numbers R56MH125655, R01MH131317, and

F31MH127933).

Data availability

Data will be made available on request.

References

Alexander, G. M., Farris, S., Pirone, J. R., Zheng, C., Colgin, L. L., Dudek, S. M., 2016. Social
and novel contexts modify hippocampal CA2 representations of space. *Nature Communications*,
7, 10300. https://doi.org/10.1038/ncomms10300

Almeida-Santos, A. F., Carvalho, V. R., Jaimes, L. F., de Castro, C. M., Pinto, H. P., Oliveira, T.
P. D., Vieira, L. B., Moraes, M. F. D., & Pereira, G. S., 2019. Social isolation impairs the
persistence of social recognition memory by disturbing the glutamatergic tonus and the olfactory

bulb-dorsal hippocampus coupling. *Scientific Reports*, *9*, 473. https://doi.org/10.1038/s41598-
018-36871-6

Breese, C. R., Hampson, R. E., Deadwyler, S. A., 1989. Hippocampal place cells: Stereotypy
and plasticity. *Journal of Neuroscience,* *9*(4), 1097–1111. https://doi.org/10.1523/jneurosci.09-
04-01097.1989

Chang, P. Y., Taylor, P. E., Jackson, M. B., 2007. Voltage imaging reveals the CA1 region at the
CA2 border as a focus for epileptiform discharges and long-term potentiation in hippocampal
slices. *Journal of Neurophysiology*, *98*(3), 1309–1322. https://doi.org/10.1152/jn.00532.2007

Chen, S., He, L., Huang, A. J. Y., Boehringer, R., Robert, V., Wintzer, M. E., Polygalov, D.,
Weitemier, A. Z., Tao, Y., Gu, M., Middleton, S. J., Namiki, K., Hama, H., Therreau, L.,
Chevaleyre, V., Hioki, H., Miyawaki, A., Piskorowski, R. A., McHugh, T. J., 2020. A hypothalamic
novelty signal modulates hippocampal memory. *Nature*, *586*(7828), 270–274.
https://doi.org/10.1038/s41586-020-2771-1

Colgin, L. L., Moser, E. I., Moser, M.-B., 2008. Understanding memory through hippocampal
remapping. *Trends in Neurosciences*, *31*(9), 469–477. https://doi.org/10.1016/j.tins.2008.06.008

Cui, Z., Gerfen, C. R., Young 3rd, W. S., 2013. Hypothalamic and other connections with dorsal
CA2 area of the mouse hippocampus. *Journal of Comparative Neurology*, *521*(8).
https://doi.org/10.1002/cne.23263

Dang, R., Zhou, Y., Zhang, Y., Liu, D., Wu, M., Liu, A., Jia, Z., Xie, W., 2022. Regulation of social
memory by lateral entorhinal cortical projection to dorsal hippocampal CA2. *Neuroscience*
Bulletin, *38*(3), 318–322. https://doi.org/10.1007/s12264-021-00813-6

Dantzer, R., Tazi, A., Bluthé, R.-M., 1990. Cerebral lateralization of olfactory-mediated affective
processes in rats. *Behavioural Brain Research*, *40*(1), 53–60. https://doi.org/10.1016/0166-
4328(90)90042-D

Ego-Stengel, V., Wilson, M. A., 2010. Disruption of ripple-associated hippocampal activity during
rest impairs spatial learning in the rat. *Hippocampus*, *20*(1), 1–10.
https://doi.org/10.1002/hipo.20707

Fenton, A. A., Csizmadia, G., Muller, R. U., 2000. Conjoint control of hippocampal place cell
firing by two visual stimuli: the effects of moving the stimuli on firing field positions. *Journal of*
General Physiology, *116*(2), 191–210. https://doi.org/10.1085/JGP.116.2.191

Gheusi, G., Goodall, G., Dantzer, R., 1997. Individually distinctive odours represent individual
conspecifics in rats. *Animal Behaviour*, *53*(5). https://doi.org/10.1006/anbe.1996.0314

Guevara-Aguilar, R., Jimenez-Montufar, L. L., Garcia-Diaz, D. E., Wayner, M. J., Armstrong, D.
L., 1988. Olfactory and visceral projections to the paraventricular nucleus. *Brain Research*
Bulletin, *20*(6), 799–801. https://doi.org/10.1016/0361-9230(88)90094-9

Hassan, S. I., Bigler, S., Siegelbaum, S. A., 2023. Social odor discrimination and its
enhancement by associative learning in the hippocampal CA2 region. *Neuron*, *111*(14), 2232-
2246.e5. https://doi.org/10.1016/j.neuron.2023.04.026

Hitti, F. L., Siegelbaum, S. A., 2014. The hippocampal CA2 region is essential for social memory.
Nature, *508*(7494). https://doi.org/10.1038/nature13028

Hsiao, Y.-T., Zheng, C., Colgin, L. L., 2016. Slow gamma rhythms in CA3 are entrained by slow
gamma activity in the dentate gyrus. *Journal of Neurophysiology*, *116*(6), 2594–2603.
https://doi.org/10.1152/jn.00499.2016

Hurst, J. L., 2005. Scent marking and social communication. In P. K. McGregor (Ed.), *Animal
Communication Networks* (pp. 219–244). Cambridge University Press.
https://doi.org/10.1017/CBO9780511610363.014

Husted, J. R., McKenna, F. S., 1966. The use of rats as discriminative stimuli. *Journal of the
Experimental Analysis of Behavior*, *9*(6). https://doi.org/10.1901/jeab.1966.9-677

Hwaun, E., Colgin, L. L., 2019. CA3 place cells that represent a novel waking experience are
preferentially reactivated during sharp wave-ripples in subsequent sleep. *Hippocampus*, *29*(10),
921–938. https://doi.org/10.1002/hipo.23090

Igarashi, K. M., Ieki, N., An, M., Yamaguchi, Y., Nagayama, S., Kobayakawa, K., Kobayakawa,
R., Tanifuji, M., Sakano, H., Chen, W. R., Mori, K., 2012. Parallel mitral and tufted cell pathways
route distinct odor information to different targets in the olfactory cortex. *Journal of
Neuroscience*, *32*(23), 7970–7985. https://doi.org/10.1523/jneurosci.0154-12.2012

Karlsson, M. P., Frank, L. M., 2009. Awake replay of remote experiences in the hippocampus.
Nature Neuroscience, *12*(7), 913–918. https://doi.org/10.1038/nn.2344

Kerr, K. M., Agster, K. L., Furtak, S. C., Burwell, R. D., 2007. Functional neuroanatomy of the
parahippocampal region: The lateral and medial entorhinal areas. *Hippocampus*, *17*(9), 697–
708. https://doi.org/10.1002/hipo.20315

Kogan, J. H., Franklandand, P. W., & Silva, A. J., 2000. Long-term memory underlying
hippocampus-dependent social recognition in mice. *Hippocampus*, *10*(1), 47–56.
https://doi.org/10.1002/(sici)1098-1063(2000)10:1<47::aid-hipo5>3.0.co;2-6

Kohara, K., Pignatelli, M., Rivest, A. J., Jung, H.-Y., Kitamura, T., Suh, J., Frank, D., Kajikawa,
K., Mise, N., Obata, Y., Wickersham, I. R., Tonegawa, S., 2014. Cell type-specific genetic and
optogenetic tools reveal hippocampal CA2 circuits. *Nature Neuroscience*, *17*(2).
https://doi.org/10.1038/nn.3614

Lee, H.-J., Caldwell, H. K., Macbeth, A. H., Tolu, S. G., Young, W. S., 2008. A conditional
knockout mouse line of the oxytocin receptor. *Endocrinology*, *149*(7), 3256–3263.
https://doi.org/10.1210/en.2007-1710

Lin, Y.-T., Hsieh, T.-Y., Tsai, T.-C., Chen, C.-C., Huang, C.-C., Hsu, K.-S., 2018. Conditional
deletion of hippocampal CA2/CA3a oxytocin receptors impairs the persistence of long-term

social recognition memory in mice. *Journal of Neuroscience*, *38*(5).

https://doi.org/10.1523/jneurosci.1896-17.2017

Lopez-Rojas, J., de Solis, C. A., Leroy, F., Kandel, E. R., Siegelbaum, S. A., 2022. A direct

lateral entorhinal cortex to hippocampal CA2 circuit conveys social information required for

social memory. *Neuron*, *110*(9), 1559-1572.e4. https://doi.org/10.1016/j.neuron.2022.01.028

Lu, L., Igarashi, K. M., Witter, M. P., Moser, E. I., & Moser, M. B., 2015. Topography of place

maps along the CA3-to-CA2 axis of the hippocampus. *Neuron*, *87*(5), 1078-1092.

https://doi.org/10.1016/j.neuron.2015.07.007

Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S., Leutgeb, J. K., 2015. Hippocampal CA2

activity patterns change over time to a larger extent than between spatial contexts. *Neuron*,

85(1). https://doi.org/10.1016/j.neuron.2014.12.001

McClelland, J. L., Goddard, N. H., 1996. Considerations arising from a complementary learning

systems perspective on hippocampus and neocortex. *Hippocampus*, *6*(6), 654–665.

https://doi.org/10.1002/(SICI)1098-1063(1996)6:6<654::aid-hipo8>3.0.co;2-g

Meira, T., Leroy, F., Buss, E. W., Oliva, A., Park, J., Siegelbaum, S. A., 2018. A hippocampal

circuit linking dorsal CA2 to ventral CA1 critical for social memory dynamics. *Nature*

Communications, *9*(1), 1–14. https://doi.org/10.1038/s41467-018-06501-w

Meshulam, L., Gauthier, J. L., Brody, C. D., Tank, D. W., & Bialek, W., 2017. Collective Behavior

of Place and Non-place Neurons in the Hippocampal Network. *Neuron*, *96*(5), 1178-1191.e4.

https://doi.org/10.1016/j.neuron.2017.10.027

Mitre, M., Marlin, B. J., Schiavo, J. K., Morina, E., Norden, S. E., Hackett, T. A., Aoki, C. J.,

Chao, M. V., Froemke, R. C., 2016. A distributed network for social cognition enriched for

oxytocin receptors. *Journal of Neuroscience*, *36*(8), 2517–2535.

https://doi.org/10.1523/jneurosci.2409-15.2016

Morris, A. M., Weeden, C. S., Churchwell, J. C., Kesner, R. P., 2013. The role of the dentate

gyrus in the formation of contextual representations. *Hippocampus, 23(2),* 162–168.

https://doi.org/10.1002/hipo.22078

Muller, R. U., Kubie, J. L., Ranck, J. B., 1987. Spatial firing patterns of hippocampal complex-

spike cells in a fixed environment. *Journal of Neuroscience, 7*(7), 1935–1950.

https://doi.org/10.1523/jneurosci.07-07-01935.1987

Nagelhus, A., Andersson, S. O., Cogno, S. G., Moser, E. I., & Moser, M.-B., 2023. Object-

centered population coding in CA1 of the hippocampus. *Neuron*, *111*(13), 2091-2104.e14.

https://doi.org/10.1016/j.neuron.2023.04.008Oettl, L. L., Kelsch, W., 2017. Oxytocin and

Olfaction. In: Hurlemann, R., Grinevich, V. (eds) *Behavioral Pharmacology of Neuropeptides:*

Oxytocin. Current Topics in Behavioral Neurosciences, vol 35. Springer, Cham.

https://doi.org/10.1007/7854_2017_8

O'Keefe, J., 1976. Place units in the hippocampus of the freely moving rat. *Experimental*
Neurology, *51*(1), 78–109. https://doi.org/10.1016/0014-4886(76)90055-8

O'Keefe, J., Conway, D. H., 1978. Hippocampal place units in the freely moving rat: Why they
fire where they fire. *Experimental Brain Research*, *31*(4), 573–590.
https://doi.org/10.1007/BF00239813

O'Keefe, J., Dostrovsky, J., 1971. The hippocampus as a spatial map: Preliminary evidence
from unit activity in the freely-moving rat. *Brain Research*, *34*, 171–175.
https://doi.org/10.1016/0006-8993(71)90358-1

Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S., Tonegawa, S., 2016. Ventral CA1 neurons
store social memory. *Science*. https://doi.org/10.1126/science.aaf7003

Oliva, A., 2022. CA2 physiology underlying social memory. *Current Opinion in Neurobiology*, *77*,
102642. https://doi.org/10.1016/j.conb.2022.102642

Oliva, A., Fernández-Ruiz, A., Buzsáki, G., Berényi, A., 2016. Role of hippocampal CA2 region
in triggering sharp-wave ripples. *Neuron*, *91*(6), 1342–1355.
https://doi.org/10.1016/j.neuron.2016.08.008

Oliva, A., Fernandez-Ruiz, A., Karaba, L. A., 2023. CA2 orchestrates hippocampal network
dynamics. *Hippocampus*, *33*(3), 241–251. https://doi.org/10.1002/hipo.23495

Oliva, A., Fernández-Ruiz, A., Leroy, F., Siegelbaum, S. A., 2020. Hippocampal CA2 sharp-wave
ripples reactivate and promote social memory. *Nature*, *587*(7833), 264–269.
https://doi.org/10.1038/s41586-020-2758-y

Pagani, J. H., Zhao, M., Cui, Z., Avram, S. K. W., Caruana, D. A., Dudek, S. M., Young, W. S.,
2015. Role of the vasopressin 1b receptor in rodent aggressive behavior and synaptic plasticity
in hippocampal area CA2. *Molecular Psychiatry*, *20*(4), 490–499.
https://doi.org/10.1038/mp.2014.47

Popik, P., van Ree, J. M., 1998. Neurohypophyseal peptides and social recognition in rats.
Progress in Brain Research, *119*, 415–436. https://doi.org/10.1016/s0079-6123(08)61585-x

Popik, P., Vetulani, J., Bisaga, A., van Ree, J. M., 1991. Recognition cue in the rat's social
memory paradigm. *Journal of Basic and Clinical Physiology and Pharmacology*, *2*(4), 315–327.
https://doi.org/10.1515/jbcpp.1991.2.4.315

Raam, T., McAvoy, K. M., Besnard, A., Veenema, A. H., Sahay, A., 2017. Hippocampal oxytocin
receptors are necessary for discrimination of social stimuli. *Nature Communications*, *8*(1), 2001.
https://doi.org/10.1038/s41467-017-02173-0

Ramadan, W., Eschenko, O., Sara, S. J., 2009. hippocampal sharp wave/ripples during sleep
for consolidation of associative memory. *PLoS ONE*, *4*(8), e6697.
https://doi.org/10.1371/journal.pone.0006697

Rao, R. P., Heimendahl, M. von, Bahr, V., Brecht, M., 2019. Neuronal responses to conspecifics
in the ventral CA1. *Cell Reports*, *27*(12). https://doi.org/10.1016/j.celrep.2019.05.081

Sawyer, T. F., Hengehold, A. K., Perez, W. A., 1984. Chemosensory and hormonal mediation of
social memory in male rats. *Behavioral Neuroscience*, *98*(5). https://doi.org/10.1037//0735-
7044.98.5.908

Shahar-Gold, H., Gur, R., & Wagner, S., 2013. Rapid and Reversible Impairments of Short- and
Long-Term Social Recognition Memory Are Caused by Acute Isolation of Adult Rats via Distinct
Mechanisms. *PLOS ONE*, *8*(5), e65085. https://doi.org/10.1371/journal.pone.0065085

Skaggs, W. E., McNaughton, B. L., Wilson, M. A., & Barnes, C. A., 1996. Theta phase
precession in hippocampal neuronal populations and the compression of temporal sequences.
Hippocampus, *6*(2), 149–172. https://doi.org/10.1002/(sici)1098-1063(1996)6:2<149::aid-
hipo6>3.0.co;2-KSmith, A. S., Williams Avram, S. K., Cymerblit-Sabba, A., Song, J., Young, W.
S., 2016. Targeted activation of the hippocampal CA2 area strongly enhances social memory.
Molecular Psychiatry, *21*(8), 1137–1144. https://doi.org/10.1038/mp.2015.189

Stefanini, F., Kushnir, L., Jimenez, J. C., Jennings, J. H., Woods, N. I., Stuber, G. D., Kheirbek,
M. A., Hen, R., & Fusi, S., 2020. A Distributed Neural Code in the Dentate Gyrus and in CA1.
Neuron, *107*(4), 703-716.e4. https://doi.org/10.1016/j.neuron.2020.05.022

Stevenson, E. L., Caldwell, H. K., 2014. Lesions to the CA2 region of the hippocampus impair
social memory in mice. *The European Journal of Neuroscience*, *40*(9), 3294–3301.
https://doi.org/10.1111/ejn.12689

Thompson, L. T., Best, P. J., 1990. Long-term stability of the place-field activity of single units
recorded from the dorsal hippocampus of freely behaving rats. *Brain Research*, *509*(2), 299–
308. https://doi.org/10.1016/0006-8993(90)90555-P

Thor, D. H., Holloway, W. R., 1982. Social memory of the male laboratory rat. *Journal of
Comparative and Physiological Psychology*, *96*(6). https://doi.org/10.1037/0735-7036.96.6.1000

Tirko, N. N., Eyring, K. W., Carcea, I., Mitre, M., Chao, M. V., Froemke, R. C., Tsien, R. W.,
2018. Oxytocin transforms firing mode of CA2 hippocampal neurons. *Neuron*, *100*(3), 593-
608.e3. https://doi.org/10.1016/j.neuron.2018.09.008

Tsai, T.-C., Fang, Y.-S., Hung, Y.-C., Hung, L.-C., Hsu, K.-S., 2022. A dorsal CA2 to ventral CA1
circuit contributes to oxytocinergic modulation of long-term social recognition memory. *Journal
of Biomedical Science*, *29*(1), 50. https://doi.org/10.1186/s12929-022-00834-x

Vaccari, C., Lolait, S. J., Ostrowski, N. L., 1998. Comparative distribution of vasopressin V1b
and oxytocin receptor messenger ribonucleic acids in brain. *Endocrinology*, *139*(12).
https://doi.org/10.1210/endo.139.12.6382

Wersinger, S. R., Ginns, E. I., O'Carroll, A.-M., Lolait, S. J., Young III, W. S., 2002. Vasopressin
V1b receptor knockout reduces aggressive behavior in male mice. *Molecular Psychiatry*, *7*(9),
975–984. https://doi.org/10.1038/sj.mp.4001195

Wersinger, S. R., Temple, J. L., Caldwell, H. K., Young, W. S., 3rd., 2008. Inactivation of the
oxytocin and the vasopressin (Avp) 1b receptor genes, but not the Avp 1a receptor gene,

differentially impairs the bruce effect in laboratory mice (mus musculus). *Endocrinology*, *149*(1).
https://doi.org/10.1210/en.2007-1056

Widloski, J., & Foster, D. J., 2022. Flexible rerouting of hippocampal replay sequences around
changing barriers in the absence of global place field remapping. *Neuron, 110*(9), 1547-1558.
https://doi.org/10.1016/j.neuron.2022.02.002

Wilson, M. A., McNaughton, B. L., 1994. Reactivation of hippocampal ensemble memories
during sleep. *Science, 265*(5172), 676–679. https://doi.org/10.1126/science.8036517

Wintzer, M. E., Boehringer, R., Polygalov, D., McHugh, T. J., 2014. The hippocampal CA2
ensemble is sensitive to contextual change. *Journal of Neuroscience*, *34*(8).
https://doi.org/10.1523/jneurosci.2563-13.2014

Wu, W.-Y., Yiu, E., Ophir, A. G., Smith, D. M., 2023. Effects of social context manipulation on
dorsal and ventral hippocampal neuronal responses. *Hippocampus, 33(7)*, 830–843.
https://doi.org/10.1002/hipo.23507

Young, W. S., Song, J., 2020. Characterization of oxytocin receptor expression within various
neuronal populations of the mouse dorsal hippocampus. *Frontiers in Molecular Neuroscience*,
13. https://doi.org/10.3389/fnmol.2020.00040

Zhang, L., Hernández, V. S., 2013. Synaptic innervation to rat hippocampus by vasopressin-
immuno-positive fibres from the hypothalamic supraoptic and paraventricular nuclei.
Neuroscience, *228*, 139–162. https://doi.org/10.1016/j.neuroscience.2012.10.010

Zhao, M., Choi, Y.-S., Obrietan, K., Dudek, S. M., 2007. Synaptic plasticity (and the lack thereof)
in hippocampal CA2 neurons. *Journal of Neuroscience*, *27*(44), 12025–12032.
https://doi.org/10.1523/jneurosci.4094-07.2007

Zheng, C., Bieri, K. W., Hsiao, Y.-T., Colgin, L. L., 2016. Spatial sequence coding differs during
slow and fast gamma rhythms in the hippocampus. *Neuron*, *89*(2), 398–408.
https://doi.org/10.1016/j.neuron.2015.12.005

Zhu, N., Zhang, Y., Xiao, X., Wang, Y., Yang, J., Colgin, L. L., Zheng, C., 2023. Hippocampal
oscillatory dynamics in freely behaving rats during exploration of social and non-social stimuli.
Cognitive Neurodynamics, *17*(2), 411–429. https://doi.org/10.1007/s11571-022-09829-8

Figure 1. Behavioral task. Rats freely explored a familiar open field arena for four 20-minute sessions per day. A stimulus cage, a standard rat housing cage, was placed in a corner of the arena. In the first (A) and final (A') sessions of experimental conditions (i.e., Social Odor, Visual + Odor, Visual, Mirror, and Non-social Odor), and in all four sessions of the Control condition, the cage contained only clean bedding. In the middle two sessions (B and B') of experimental conditions, a cage containing stimuli ("stimulus cage") was presented. In the Social Odor condition, a stimulus cage containing soiled bedding of a familiar rat or rats was presented. In the Visual + Odor condition, a stimulus cage containing a familiar rat and the rat's soiled bedding was presented. In the Visual condition, a familiar rat was placed in the stimulus cage with clean bedding and a filter-top lid to minimize social odors. In the Mirror condition, a familiar rat was placed in the stimulus cage with clean bedding, and the cage was lined with a one-way mirror attachment. This prevented the stimulus rat from seeing the implanted rat, thereby limiting reciprocal social interactions. In the Non-social Odor condition, a wooden block infused with hexyl acetate, a fruity-smelling odorant, was placed in the stimulus cage under clean bedding.

Figure 2. CA2 Histology. An example hippocampal section showing immunohistological identification of a tetrode track in CA2. To identify CA2, hippocampal sections were immunostained with a CA2 marker, Purkinje cell protein 4 (PCP4, red). DAPI nuclear staining is shown in blue. Scale bar, 200 μm.

Figure 3. Example CA2 place cell firing rate maps across the four recording sessions for all conditions. Color-coded firing rate maps are shown for all place cells recorded on a single example tetrode in each condition for one example rat. Rate maps are shown scaled to the maximal firing rate (shown inset) of each cell across all sessions. White pixels indicate places that were not visited by the rat during a session.

Figure 4. Remapping in CA2 in response to social stimuli. A-B. The estimated means of spatial correlation coefficients from our generalized linear mixed model are shown across each condition and session pair (A). The estimated means of spatial correlation coefficients from our generalized linear mixed model are shown across each condition for all session pairs combined (B). Error bars represent 95% confidence intervals. Statistics and associated p-values for specific pairwise comparisons of spatial correlations are shown in Table 3. C-G. Spatial correlation measures are shown for the entire sample of CA2 place cells for all pairs of sessions across all conditions. Each marker represents a spatial correlation value for an individual place cell. Different symbols are used for CA2 place cells recorded from different rats.

Figure 5. Exploration of stimuli across conditions. A. Heat maps show mean exploration times of different locations in the arena, with the stimulus cage shown in the top right corner of the arena. Time spent was calculated for the first 2 minutes of each session individually and then averaged within each rat. Heat maps shown are averaged across rats. B. Time spent exploring locations close to the cage (within 12 cm) was calculated for each session and then averaged within a rat. Individual dots represent the mean exploration time for each rat, and error bars represent 95% confidence intervals across all sessions within a rat. Rats increased their exploration time of the stimulus cage in session B for the Social Odor, Visual, and Mirror conditions compared to the Control condition (* indicates p-values < 0.05 and ** indicates p-values < 0.01).

Figure 6. CA2 place cells that responded to a social odor preferentially fired during sharp wave-ripples. A. The difference between firing rates of CA2 place cells during sharp wave-ripples in the rest periods following sessions A and B were positively correlated to the selectivity index for the Social Odor condition (B) but not the Control condition (A). The selectivity index indicates a

cell's preference for session A (value of -1 for maximum selectivity in session A) or session B (value of 1 for maximum selectivity in session B).

Supplementary Figure 1. Example of CA2 localization using the CA2 marker STEP. A. A Cresyl violet-stained brain section from an example rat (rat 117) showing CA3/CA2 (black arrows) and CA2/CA1 (red arrows) borders. Scale bar, 100 μm left and 50 μm right. B. Example tetrode track in CA2 from rat 117. Sections were immunostained with a CA2 marker, STEP, and developed with 3,3-diaminobenzidine (brown). Sections were counter-stained with cresyl-violet (purple/blue). Scale bar, 100 μm left and 50 μm right.

Supplementary Figure 2. No rate remapping was observed in CA2 in response to social stimuli. A-B. The estimated means of rate overlap values from a generalized linear mixed model are shown across each condition and session pair (A). The estimated means of rate overlap values from the generalized linear mixed model are shown across each condition for all session pairs combined (B). Error bars represent 95% confidence intervals. C-G. Rate overlap values are shown for the entire sample of CA2 place cells for all pairs of sessions across all conditions. Each marker represents a rate overlap value for an individual place cell. Different symbols are used for CA2 place cells recorded from different rats.

Supplementary Figure 3. Evaluation of CA2 place field shifts relative to the location of the stimulus cage. A. Shown are the distances measured between the positions of peak firing in Session B and the stimulus cage for CA2 place cells that "turn on", or gain a field, in Session B. B. Shown are the distances between the peak firing rate position in Session A and the location of the stimulus cage for CA2 place cells that "turn off", or lose a field, in Session B. C. Shown are the changes in distance from the stimulus cage of the peak firing rate positions in Sessions A and B for CA2 place cells that were active in both Sessions A and B but shifted their place

field locations. Individual markers represent measurements from individual place cells. Different symbols represent distance measurements for CA2 place cells recorded from different rats.

Supplementary Figure 4. CA2 place cell properties across sessions and conditions. (A) CA2 place cells conveyed similar spatial information across conditions. (B-C) CA2 place cell peak firing rates (B) and in-field firing rates (C) were similar across conditions. (D) The number of place fields per CA2 cell was similar across conditions.

A   B

Supplementary Figure 1

A

B

C   A-B

D   A-B'

E   B'-A'

F   A-A'

G   B-B'

Control
Social Odor
Visual + Odor
Visual
Mirror
Non-social Odor

rat 117
rat 122
rat 165
rat 256
rat 391
rat 418