**Title: Social odors drive hippocampal CA2 place cell responses to social stimuli**

**Running title: CA2 place cells respond to social odors**

Emma Robson [a b 1], Margaret M. Donahue [a c 1], Alexandra J. Mably [a b], Peyton G. Demetrovich [a c], Lauren T. Hewitt [a c], Laura Lee Colgin [a b c]

[a] Center for Learning and Memory, The University of Texas at Austin, Austin, TX 78712

[b] Department of Neuroscience, The University of Texas at Austin, Austin, TX 78712

[c] Institute for Neuroscience, The University of Texas at Austin, Austin, TX 78712

[1] These authors contributed equally.

Corresponding author: Laura Lee Colgin, The Center for Learning and Memory, The University of Texas at Austin, 1 University Station Stop C7000, Austin, TX 78712, USA. Email:

colgin@mail.clm.utexas.edu

**Abstract**

Hippocampal region CA2 is essential for social memory processing. Interaction with social stimuli induces changes in CA2 place cell firing during active exploration and sharp wave-ripples during rest following a social interaction. However, it is unknown whether these changes in firing patterns are caused by integration of multimodal social stimuli or by a specific sensory modality associated with a social interaction. Rodents rely heavily on chemosensory cues in the form of olfactory signals for social recognition processes. To determine the extent to which olfactory signals contribute to CA2 place cell responses to social stimuli, we recorded CA2 place cells in rats freely exploring environments containing social stimuli that included or lacked olfactory content. We found that CA2 place cell firing patterns significantly changed only when social odors were prominent. Also, place cells that increased their firing in the presence of social odors alone preferentially increased their firing during subsequent sharp wave-ripples. Our results suggest that olfactory cues are essential for changing CA2 place cell firing patterns during and after social interactions. These results support prior work suggesting CA2 performs social functions and shed light on processes underlying CA2 responses to social stimuli.

**Keywords**

hippocampus, place cells, CA2, social cognition, social memory, social odor

**1. Introduction**

Social recognition memory, or the ability to recognize and remember conspecifics, facilitates

social interactions that are essential for animals to survive. Rats can discriminate between

familiar conspecifics (Husted and McKenna, 1966; Thor and Holloway, 1982) based on their

distinct characteristics and olfactory signatures (Gheusi et al., 1997; Popik et al., 1991; Sawyer

et al., 1984). Many recent studies have implicated hippocampal area CA2 in social memory (for

a review, see Oliva, 2022). Lesioning or inactivating CA2 pyramidal neurons leads to impaired

social recognition memory in mice (Hitti and Siegelbaum, 2014; Stevenson and Caldwell, 2014)

and optogenetic silencing of CA2 activity shows that CA2 is crucial for encoding, consolidating,

and recalling social memories (Meira et al., 2018). Moreover, CA2 neurons are uniquely

enriched in receptors selective for a variety of social neuropeptides, including vasopressin and

oxytocin (Lee et al., 2008; Lin et al., 2018; Pagani et al., 2015; Raam et al., 2017; Wersinger et

al., 2002, 2008).

All subregions of the hippocampus contain place cells, neurons that are selectively activated in

a particular location in space (their "place fields") (O'Keefe, 1976; O'Keefe and Dostrovsky,

1971). Place cells alter their firing in response to significant environmental changes in a process

called "remapping" (for a review, see Colgin et al., 2008). During remapping, place fields may

appear, disappear, or change location. Neurophysiological recordings of place cells have shown

that place fields in neighboring hippocampal subregions CA1 and CA3 remain relatively stable

in unchanging environments (Muller et al., 1987; Thompson and Best, 1990). In contrast, CA2

place cells show gradual changes in firing patterns over time (Alexander et al., 2016; Mankin et

al., 2015). CA2 place cells also appear to be highly sensitive to small updates to familiar

environments (Wintzer et al., 2014). Previous work has shown that a significant proportion of

CA2 place cells remap when a familiar conspecific is presented (Alexander et al., 2016). This

remapping in response to social stimuli may support the ability of CA2 to encode social

memories (Hitti and Siegelbaum, 2014). However, it remains unclear whether CA2 place field

changes that occur in response to presentation of a conspecific rat are caused by integration of

multimodal social stimuli or primarily due to a specific sensory modality associated with the

social interaction.

Previous work has also shown that CA2 firing patterns during sleep are altered following a

social experience (Oliva et al., 2020). During awake rest or sleep, hippocampal place cells fire

during distinctive events in the hippocampal local field potential (LFP) known as sharp wave-

ripples. Place cells that were active during exploratory behaviors later reactivate during sharp

wave-ripples, and reactivated place cell ensembles are believed to support memory

consolidation (Ego-Stengel and Wilson, 2010; Ramadan et al., 2009; Wilson and McNaughton,

1994). CA2 place cells that represent a conspecific have been shown to reactivate during sharp

wave-ripples following a social experience (Oliva et al., 2020). Further, disrupting CA2 sharp

wave-ripples impairs social recognition memory (Oliva et al., 2020). Therefore, the reactivation

of CA2 place cells that encode a social experience may promote consolidation of social

memories (Oliva et al., 2023).

In this study, we recorded CA2 place cells in dorsal hippocampus of rats during free exploration

of a two-dimensional spatial environment and during subsequent rest periods. We compared

changes in CA2 place cell firing patterns across sessions in which different social stimuli

incorporating various sensory modalities were presented. Significant changes in CA2 place cell

firing patterns were observed when social odors were presented in the absence of a rat and

when a familiar rat was presented in a soiled home cage containing social odors (as in our prior

study; Alexander et al., 2016). However, no significant changes in CA2 place cell firing patterns

were observed when a familiar rat was presented in a clean and relatively odorless cage. The

results suggest that olfactory cues are the key sensory component of social experiences that

drives CA2 place cell remapping. Furthermore, CA2 place cells that increased their firing rates

in response to social odors preferentially increased their firing rates in sharp wave-ripples during

subsequent rest periods. These results improve our understanding of how CA2 place cells code

information related to social interactions.

**2.  Materials and Methods**

*2.1.   Subjects*

Six wild-type male Sprague-Dawley rats (Inotiv, USA) were used for this study. Rats were

between the ages of 4 and 10 months at the time of surgery. Before surgery, rats were double or

triple housed and were pre-trained to freely explore an open field enclosure. After surgery, rats

were singly housed in custom-built acrylic cages (40 cm x 40 cm x 40 cm) containing

enrichment material (wooden blocks, paper towel rolls, etc.) and maintained on a reverse light

cycle (light: 8 p.m. – 8 a.m.). Rats were housed next to their former cage mates after recovering

from surgery and throughout behavioral testing. Rats recovered from surgery for at least one

week before behavioral training resumed. All behavioral experiments were performed during the

dark cycle. When necessary to encourage spatial exploration, rats were placed on a food-

deprivation regimen that maintained them at ~90% of their free-feeding body weight. All

experiments were conducted according to the guidelines of the United States National Institutes

of Health Guide for the Care and Use of Laboratory Animals and under a protocol approved by

the University of Texas at Austin Institutional Animal Care and Use Committee.

*2.2.   Surgery and tetrode positioning*

Drives with 14 independently movable tetrodes were implanted in five of the rats. A drive with 21

independently movable tetrodes was implanted in one of the rats. Drives were implanted above

the right dorsal hippocampus (anterior-posterior -3.8 mm from bregma, medial-lateral -3.0 mm

from bregma). To stabilize the recording drives, eleven bone screws were affixed to the skull

and covered in dental acrylic. Two of the screws were connected to the recording drive ground.

Prior to surgery, tetrodes were built from 17 µm polyimide-coated platinum-iridium (90/10%) wire

(California Fine Wire, Grover Beach, California, USA). The tips of tetrodes designated for single-

unit recording were plated with platinum to reduce single-channel impedances to ~150 to

300 kOhms. All tetrodes were lowered ~1 mm on the day of surgery. Thereafter, tetrodes were

slowly lowered to the hippocampal pyramidal cell body layer over the course of several weeks.

One tetrode was designated for use as a reference for differential recording. All four wires of this

tetrode were connected to a single channel on the electrode interface board. The differential

recording reference tetrode was placed in an electrically quiet area approximately 1 mm above

the hippocampus and adjusted as needed to ensure quiescence. The reference signal was

duplicated using an MDR-50 breakout board (Neuralynx, Bozeman, MT, USA) and recorded

continuously to ensure that unit activity or volume conducted signals of interest were not

detected. Another tetrode was placed in the apical dendritic layer of CA1 to monitor LFPs and

guide placement of the other tetrodes using estimated depth and electrophysiological hallmarks

of the hippocampus (for example, sharp wave-ripples).

*2.3.   Data acquisition*

Data were acquired using a Digital Lynx system and Cheetah recording software (Neuralynx,

Bozeman, MT, USA). The recording setup has been described in detail previously (Hsiao et al.,

2016; Zheng et al., 2016). Briefly, LFP signals from one randomly chosen channel within each

tetrode were continuously recorded at a 2000 Hz sampling rate and filtered in the 0.1–500 Hz

band. Input amplitude ranges were adjusted before each recording session to maximize

resolution without signal saturation. Input ranges for LFPs generally fell within ±2,000 to

±3,000 µV. To detect unit activity, all four channels within each tetrode were bandpass filtered

from 600 to 6,000 Hz. Spikes were detected when the filtered continuous signal on one or more

of the channels exceeded a threshold set daily by the experimenter, which ranged from 55–

65 µV. Detected events were acquired with a 32,000 Hz sampling rate for 1 ms. For both LFPs

and unit activity, signals were recorded differentially against a dedicated reference channel (see

"Surgery and tetrode positioning" section above).

Videos of rats' behavior were recorded through the Neuralynx system with a resolution of 720 ×

480 pixels and a frame rate of 29.97 frames/s. Rat position and head direction were tracked via

an array of red and green or red and blue light-emitting diodes (LEDs) on a HS-54 or HS-27

headstage (Neuralynx, Bozeman, MT, USA), respectively.

*2.4.    Behavioral task*

Rats were familiarized to an open field arena (1 m x 1 m with 0.5 m wall height) for a minimum

of three days before recording started. Rats freely explored the open field arena for four 20-

minute sessions per day, with 10-min rest sessions preceding and following each active

exploration session. During active exploration, small pieces of sweetened cereal or cookies

were randomly scattered to encourage rats to explore the entirety of the arena. Rats had to

cover at least 60% of the arena across each of the four sessions for a day to be included for

further analysis. During each rest session, rats rested in a towel-lined, elevated flowerpot

outside of the arena. A plexiglass standard rat housing cage was placed in one corner of the

arena for all recording sessions. In the first and fourth sessions (A and A'), this cage contained

only clean bedding. In the middle two sessions (B and B') of the different experimental

conditions, the cage contained various types of social stimuli (Figure 1). In the Odor condition,

the soiled bedding from the former home cage of familiar rats (or a familiar rat) was used. In the

Visual condition, a familiar rat was placed in the stimulus cage with clean bedding and a filter-

top lid to minimize social odors. In the Visual + Odor condition, a familiar rat was presented in its

home cage containing soiled bedding. In the Mirror condition, a familiar rat was placed in the

stimulus cage with clean bedding and a filter-top lid, but the cage was lined with a one-way

mirror attachment. This one-way mirror attachment prevented the stimulus rat from seeing the

implanted rat, limiting visually driven reciprocal interactions between the two rats. In the Control

condition, a cage containing only clean bedding was presented in all four sessions.

*2.5.    Histology and tetrode localization*

Following recording, rats were perfused with 4% paraformaldehyde solution in phosphate-

buffered saline. Brains were cut coronally in 30 μm sections using a cryostat. For two rats,

brains were stained with cresyl violet. For four rats, brains were immunostained for the CA2

marker Purkinje Cell Protein 4 (PCP4) (Figure 2). For the immunostaining protocol, sections

were washed three times for 15 minutes in tris-buffered saline (TBS) followed by a 10-minute

water wash. Sections were then permeabilized and washed for 15 minutes in TBS containing

0.3% Triton-X followed by three 10-minute washes in TBS. Sections were blocked for 30

minutes in 10% normal goat serum in TBS. Sections were incubated overnight with rabbit anti-

PCP4 (1:200, Sigma-Aldrich Cat# HPA005792) diluted in TBS containing 0.05% Tween. The

next day, sections were washed twice for 10 minutes in TBS and incubated overnight with

secondary fluorescent antibody (in one rat: Alexa Flour™-555 anti-rabbit, Thermo Fisher

Scientific; in three rats: Alexa Flour™-555 anti-rabbit, Thermo Fisher Scientific). All washes and

incubations were performed at room temperature. Slides were mounted using DAPI

Fluoromount-G (Fisher Scientific). Tetrode recording sites were identified by comparing

locations across adjacent sections.

### 2.6. Spike sorting and unit selection

Spike sorting was performed manually using graphical cluster-cutting software (MClust, A.D.

Redish, University of Minnesota, Minneapolis, Minnesota) run in MATLAB (Mathworks). Spikes

were sorted using two-dimensional representations of waveform properties (i.e., energy, peak,

and peak-to-valley difference) from four channels. A single unit was accepted for further analysis

if the associated cluster was well isolated from, and did not share spikes with, other clusters on

the same tetrode. Units were also required to have a minimum 1 ms refractory period. Units with

mean firing rates above 5 Hz were considered putative interneurons and excluded from further

analysis. In order to be considered active in the arena, a unit had to reach a peak firing rate of at

least 1 Hz. In order to be included in the sharp wave-ripple firing analysis, a unit had to have

valid clusters in both the active exploration and rest sessions. CA2 cell yields for each condition

are reported in Tables 1 and 2.

### 2.7. Place cell remapping analyses

Methods used to create firing rate maps for each single unit were based on methods used in our

prior study (Alexander et al., 2016; example rate maps for the present study shown in Figure 3).

First, the arena was divided into 4 $cm^2$ bins. The number of spikes that occurred within each bin

was divided by the time spent in that bin to determine the firing rate. Only spikes that occurred

while the rat was traveling 5 cm/s or faster were included. The rate map was smoothed with a

two-dimensional Gaussian kernel (standard deviation = 6 cm). To determine if a place cell

remapped during presentation of a social stimulus, a Pearson correlation coefficient $R$ was

calculated for each unit between pairs of rate maps from control and social stimuli sessions (i.e.,

A-B, B-B', B'-A', A-B', and A-A' session pairs, see Figure 1). To determine if spatial correlation

coefficients differed across session pairs and conditions (i.e., Odor, Visual + Odor, Visual, Mirror,

Control), we used a generalized linear mixed model statistical analysis (IBM SPSS Statistics,

version 29.0.2.0). Condition and session pair were fixed factors, session pairs were repeated

measures within rats, and multiple place cells were nested within rats. A condition by session

pair interaction effect was also included in the model to determine whether differences between

session pairs varied across conditions. When a significant effect was observed, post-hoc

pairwise comparisons were performed with Bonferroni correction for multiple comparisons. The

estimated mean spatial correlations for each condition and session pair from the generalized

linear mixed model are shown in Figure 4A. The estimated mean spatial correlations for each

condition, collapsed across session pairs, are shown in Figure 4B. Individual spatial correlation

values for each cell are shown for each session pair and each condition in Figure 4C-G.

To determine if positions of place fields coherently shifted during remapping, we first identified

cells that remapped and sorted them into "turn on", "turn off", and "field shift" categories (Figure

5). Because place cells were classified as active if they reached a peak firing rate of at least 1

Hz, "turn on" cells were defined as cells that had a peak firing rate less than 1 Hz in Session A

and greater than 1 Hz in Session B. We then calculated the distance between the position of the

peak firing rate of the cell in Session B and the stimulus cage for "turn on" cells. "Turn off" cells

were defined as cells that had a peak firing rate above 1 Hz in Session A and below 1 Hz in

Session B. The distance between the position of the peak firing rate of the cell in Session A and

the stimulus cage was calculated for "turn off" cells. Cells were identified as "field shift" cells if

they reached a peak firing rate above 1 Hz in Sessions A and B but showed unstable place field

locations. We assessed place field stability as follows, using criteria defined in a prior study

(Widloski and Foster 2022). Cell IDs were randomly shuffled 1000 times within each condition

and rat. Spatial correlation coefficients were then calculated for a given unit from session A and

each shuffled rate map from session B. A place field was considered stable between sessions A

and B if the spatial correlation coefficient exceeded the 95$^{th}$ percentile of the shuffled

distribution. For "field shift" cells, we calculated the distances between the stimulus cage and

the positions of the peak firing rates of the cell in Sessions A and B. To determine if a cell's place

field moved closer to the stimulus cage when social stimuli were presented, we calculated the

difference between the distance estimates from Sessions A and B for "field shift" cells.

To quantify how selective a unit was for session A vs session B, we calculated a selectivity index

for each unit, as in Hwaun and Colgin (2019). The selectivity index was defined as $(\mu_B -$

$\mu_A)/(\mu_B + \mu_A)$, where $\mu_A$ was the mean firing rate in session A and $\mu_B$ was the mean firing rate in

session B. A value of -1 would indicate that a unit was exclusively active in session A, while a

value of 1 would indicate that a unit was exclusively active in session B.

*2.8.    Exploration time*

To investigate exploration of social stimuli, the arena was divided into 4 $cm^2$ bins, as in a prior

study (Zhu et al., 2023). The time spent in each bin during the first two minutes of every session

was determined for each day. These exploration maps were then smoothed with a two-

dimensional Gaussian kernel (standard deviation = 4 cm). Maps were averaged within and then

across rats and plotted as a heat map for each condition (Figure 6A). The time spent within 12

cm of the cage was then calculated. A generalized linear mixed model analysis (IBM SPSS

Statistics, V 29.0.2.0) was used to compare time spent investigating the stimulus cage across

different sessions and conditions, with conditions and sessions as fixed factors. Conditions were

repeated on different days within rats, with different days included as random factors within rats.

Sessions were repeated measures within conditions. Post-hoc tests were performed to compare

exploration times across conditions for the session of interest (i.e., Session B), using the

Bonferroni correction for multiple comparisons.

*2.9.    Sharp wave-ripple detection and analysis*

Sharp wave-ripples were detected from recordings obtained while rats rested in an elevated,

towel-lined flowerpot outside of the arena. Detection and analysis methods were similar to

previously published methods (Hwaun and Colgin, 2019). The LFP recorded on all tetrodes that

had CA2 place cells was band-pass filtered between 150 and 250 Hz. A Hilbert transform was

performed on the filtered LFP, and the absolute value of this signal was smoothed with a

Gaussian kernel (standard deviation = 25 ms). Potential sharp wave-ripple events were

detected when the signal exceeded at least 5 standard deviations above the mean and bounded

by crossings of the mean. Overlapping events were combined across tetrodes, so the time

interval of identified events could extend beyond sharp wave-ripples detected on a single

tetrode (as in Hwaun and Colgin, 2019; Karlsson and Frank, 2009). Potential sharp wave-ripple

events were kept for further analysis if they were between 50 and 500 ms in duration.

We then obtained the firing rate for each CA2 place cell around detected sharp wave-ripple

events by constructing a spike raster with a bin size of 1 ms and smoothing the raster with a

Gaussian kernel (standard deviation = 5 ms). To account for baseline firing rate differences

among individual cells, the firing rate was normalized by the average firing rate 400-100 ms

before ripple onset (Hwaun and Colgin, 2019). The peak normalized firing rate was obtained for

each unit by taking the maximum normalized firing rate after sharp wave-ripple onset. Because

a broad range of peak normalized firing rates was observed, we rescaled the distribution using a

$\log_{10}$ transformation of the peak normalized firing rates. To determine the extent to which sharp

wave-ripple-associated firing rates of individual cells were altered by presentation of a social

stimulus, we computed the difference between the peak normalized firing rates during the rest

periods after sessions A and B (i.e., peak normalized firing rate after session B - peak

normalized firing rate after session A) for the Control and Odor conditions. A positive value

indicates that a cell had higher sharp wave-ripple-associated firing in the rest session following

session B, and a negative value indicates that a cell had higher sharp wave-ripple-associated

firing in the rest session following session A. We used multiple linear regression (IBM SPSS

Statistics, V29.0.2.0) to assess the effects of session selectivity (as measured using the

selectivity index, see *Place cell remapping analyses* section) and experimental condition on the

difference in sharp wave-ripple-associated firing between rest sessions (as in Hwaun and

Colgin, 2019). The regression analysis was only performed for conditions that had more than

100 active cells during both run and rest sessions (Control condition: n = 130 cells from 5 rats,

Odor condition: n = 132 cells from 5 rats).

*2.10.   Data and Code Availability*

MATLAB (Mathworks) scripts were custom written for the analyses in this study, based on

algorithms that have been used in prior studies, as described above. Scripts and data are

available upon request.

**3.  Results**

In our prior study of CA2 place cells (Alexander et al., 2016), we compared CA2 place cell

responses to a rat in a soiled home cage and responses to the presentation of an object that

resembled a rat, namely a stuffed toy rat. However, the stuffed toy lacked some visual

components that occur during interactions with a live rat (e.g., motion content). Also, the toy rat

was presented in a clean cage and thus lacked the olfactory components of a social experience.

The major goal of the present study was to determine the extent to which different sensory

modalities associated with a social experience drive CA2 place cell remapping to social stimuli.

We employed an open field exploration behavior paradigm (see "*Behavioral task*" section of

Materials and Methods) in which different types of social stimuli were presented across sessions

for various experimental conditions (i.e., Odor, Visual, Visual + Odor, Mirror) and compared to a

Control condition in which no social stimuli were presented (Figure 1).

*3.1.    CA2 place cells remap to social odor stimuli*

The olfactory system is highly important for social recognition and social behaviors in rodents

(Oettl and Kelsch, 2018). Therefore, we hypothesized that olfactory cues would be a key

sensory component of CA2 place cell responses to social stimuli. In the Odor condition, rats

explored an environment in which social odors were presented, but another rat was not present.

This condition isolated the olfactory content of a social interaction and preserved the ethological

relevance of the presented stimuli. Example CA2 place cell firing rate maps during presentation

of social odors (Figure 3B) show that a subset of cells changed their firing patterns when social

odors were presented. To quantify CA2 place cell remapping, we compared firing rate changes

in cells recorded in the Odor condition (145 cells in 6 rats, Table 1) to firing rate changes in cells

recorded in the other conditions (see Figure 3 for example rate maps for all conditions and Table

1 for cell yields). We calculated the spatial correlation between rate maps from pairs of sessions

for each cell (Figure 4). We found that spatial correlation values for the Odor condition were

significantly lower than the Control, Visual, and Mirror conditions (generalized linear mixed

model, no significant interaction between session pair and condition, $F_{(16, 3025)} = 0.971$, $p =$

0.486 (Figure 4A); significant main effect of condition, $F_{(4, 3025)} = 15.4$, $p < 0.001$ (Figure 4B);

significant differences in post-hoc tests for Odor vs. Control ($t_{(3025)} = 6.3$, $p < 0.001$), Odor vs.

Visual ($t_{(3025)} = 4.0$, $p = 0.001$), and Odor vs. Mirror ($t_{(3025)} = 3.6$, $p = 0.001$)). In agreement

with our prior findings, CA2 place cells also showed significant remapping when a familiar

conspecific rat was presented in a soiled home cage containing social odors (Visual + Odor

condition, Figures 3C and 4, significant differences in post-hoc tests for Visual + Odor vs.

Control ($t_{(3025)} = 6.4$, $p < 0.001$), Visual + Odor vs. Visual ($t_{(3025)} = 4.5$, $p < 0.001$), and Visual

+ Odor vs. Mirror ($t_{(3025)} = 4.1$, $p = 0.001$). In contrast, significant changes in CA2 place cell

firing patterns were not observed when a stimulus rat was presented in a clean cage with a filter

top, a condition that maintained all visual components of social interactions and minimized

social odors (Visual condition, Figures 3D and 4, no significant differences in post-hoc test for

Visual vs. Control (t(3025) = 1.8, p = 0.2)). Similarly, CA2 place cell firing patterns did not

change significantly when a stimulus rat was presented in a clean cage with a filter top and a

one-way mirror to prevent visually mediated reciprocal interactions between rats (Mirror

condition, Figures 3E and 4, no significant differences in post-hoc test for Mirror vs. Control

(t(3025) = 2.3, p = 0.1)). Together, these results show that CA2 place cells remap when social

odors are presented but not during interactions with another rat when social odors are

minimized.

*3.2.    No coherent movement of place fields during remapping*

CA1 place fields have been reported to shift with salient stimuli (Fenton et al., 2000; O'Keefe

and Conway, 1978) or move toward locations of salient rewarding stimuli (Breese et al., 1989;

Fenton et al., 2000). However, our previous work showed that fields of CA2 place cells do not

move closer to a social stimulus during social remapping (Alexander et al., 2016). To test if CA2

place cells moved coherently towards or away from the social stimuli in our experiment, we first

identified place cells that remapped between sessions A and B. These cells were identified as

cells that either gained a place field ("turn on" cells), lost their place field ("turn off" cells), or

shifted their place field ("field shift" cells). For "turn on" cells, we calculated the distance between

the cell's peak firing rate position in the session in which social stimuli were presented (Session

B) and the position of the stimulus cage (Figure 5A). Similarly, we calculated the distance

between the peak firing rate position and the position of the stimulus cage in Session A for the

"turn off" cells (Figure 5B). If place cells preferentially gained or lost fields close to the stimulus

cage, we would have expected a skewed distribution towards low values. Instead, we observed

a broad distribution of distances from the stimulus cage. For "field shift" cells, we estimated the

distance between the stimulus cage and cells' peak firing rate positions in Sessions A and B and

then calculated the difference between these two distance estimates (Figure 5C). If place fields

shifted closer to the cage, we would have expected values to skew negatively. However,

differences between place field distances from the cage between Sessions A and B covered a

wide range of positive and negative values. Therefore, consistent with our previous findings

(Alexander et al., 2016), no coherent movement of place fields toward or away from the stimulus

cage was observed. This may be due to the nature of odor stimuli, as airborne odorants diffuse

across widespread locations within an environment. This may allow the rat to perceive odors

regardless of the exact location where odor stimuli are presented in the arena. It is also possible

that the introduction of social odors to an environment was sufficient to change the context as a

whole. This generalizability may help rats to associate conspecifics with larger areas of an

environment.

*3.3.    Rats explore all modalities of social stimuli*

It is possible that the lack of remapping to non-olfactory social stimuli may be due to inadequate

exploration of these stimuli. To test this possibility, we examined rats' stimuli exploration time

during the first 2 minutes of each exploration session for each experimental condition (Figure 6).

We found that stimulus cage exploration time during session B of most social conditions (i.e.,

Odor, Visual, and Mirror) significantly increased compared to session B of the Control condition

(generalized linear mixed model, interaction between session and condition, $F(12,260) = 2.589$,

$p = 0.003$; post-hoc tests, session B: significant difference between Control vs. Odor: $t(260) =$

$2.8$, $p = 0.03$; Control vs. Visual: $t(260) = 3.3$, $p = 0.009$; and Control vs. Mirror: $t(260) = 3.3$, $p =$

$0.01$; No significant difference between Control vs. Visual + Odor: $t(260) = 2.1$, $p = 0.2$). It is

important to note that stimulus cage exploration times significantly increased in the social

conditions in which CA2 place cells did not show significant remapping (i.e., Visual and Mirror

conditions), suggesting that the lack of place cell remapping in these conditions was not due to

reduced exploration of social stimuli. Note also that exploration times did not significantly differ

between Visual + Odor and Control conditions, although significant remapping was observed in

the Visual + Odor condition. However, the insignificant difference in exploration times may be

due to the lower number of rats in the Visual + Odor condition (i.e., 4 rats in Visual + Odor

condition compared to 5-6 rats in other conditions, see Table 1).

*3.4.    Place cells that respond to social odors are preferentially recruited into sharp wave-*

*ripples*

CA2 place cells that encode a social experience with a novel conspecific have been shown to

reactivate during sharp wave-ripples in mice (Oliva et al., 2020). Therefore, we aimed to

determine whether presentation of social odors alone was sufficient to drive preferential sharp

wave-ripple-associated reactivation of CA2 place cells that responded to social odors. We used

a selectivity index (see Materials and Methods) to classify cells according to their firing

preferences during the Control and Odor conditions. The selectivity index defined the extent to

which CA2 place cells fired during session A (i.e., the session in which an empty cage was

presented) compared to session B (i.e., the session in which a cage containing social odors was

presented in the Odor condition). We then estimated the normalized average peak firing rate of

each cell during sharp wave-ripples in rest sessions after session A and session B (see

Materials and Methods). Next, we used the difference between a cell's peak normalized firing

rate during the rest session after session B and the rest session after session A as a measure of

the extent to which a cell changed its sharp wave-ripple-associated firing after presentation of

social odors. CA2 place cells show unstable firing patterns in unchanged environments over

time (Mankin et al., 2015). Thus, we performed the same analysis for CA2 place cells recorded

in the Control condition to ensure that changes in sharp wave-ripple-associated firing were not

explained by the effect of time. We found that a cell's selectivity for social stimuli in active

exploratory sessions correlated with sharp wave-ripple-associated peak firing rate changes in

the Odor condition but not the Control condition (Figure 7; multiple linear regression, $F(3,243) =$

$2.661$, $p = 0.049$; interaction between selectivity index and condition, $t = 2.448$, $p = 0.015$,

Pearson correlation between selectivity index and firing rate changes for the Control condition R

= -0.102, p = 0.266, Pearson correlation between selectivity index and firing rate changes for

the Odor condition R = 0.227, p = 0.010). Specifically, cells that increased their firing rates

during exploration of a social odor preferentially increased their sharp wave-ripple-associated

firing rates during rest. These results suggest that the presentation of a social odor alone is

sufficient to preferentially reactivate CA2 place cells that code social stimuli.

**4. Discussion**

An increasing number of lesion, genetic, optogenetic, and pharmacogenetic manipulation

studies have implicated CA2 in social memory processing (Hitti and Siegelbaum, 2014; Meira et

al., 2018; Oliva et al., 2020; Smith et al., 2016; Stevenson and Caldwell, 2014). In this study, we

expanded on prior results showing that CA2 place cells remap during a social interaction

(Alexander et al., 2016) by investigating how various sensory modalities of social stimuli

contribute to this response. Here, we show that CA2 place cell remapping is primarily driven by

the olfactory component of social stimuli. Exposure to social odors in the absence of a rat was

sufficient to induce CA2 place cell remapping. CA2 place cells did not respond to social

interactions with another rat when social odors were reduced. The lack of remapping when

social odors were reduced was not explained by inadequate exploration of social stimuli. Lastly,

CA2 place cells that showed increased firing during exploration of social odors were

preferentially reactivated during sharp wave-ripples in subsequent rest. Our findings support the

notion that CA2 place cells integrate social olfactory information with contextual information to

encode and consolidate social episodic memories (Oliva, 2022; Oliva et al., 2023). While social

experiences did not induce significant CA2 place cell remapping when social odors were

lacking, the possibility remains that other sensory modalities of a social experience are

processed via different pathways and brain structures and are still able to support social

recognition and social memory processes.

The sensory modalities that drive social recognition vary across species. While humans depend

predominantly on auditory and visual cues for social recognition, rodents and other mammals

rely heavily on chemosensory cues in the form of olfactory or pheromonal signals (Popik and

van Ree, 1998). Indeed, olfactory bulb lesions or chemically induced anosmia impairs individual

recognition in rats (Dantzer et al., 1990; Popik et al., 1991). The complex mix of chemosensory

signals embedded in rat urine convey information about a rat's social status and identity (Hurst,

2005). CA2 neurons differentially respond to urine from different conspecifics in mice (Hassan et

al., 2023). The ability of CA2 to respond to social odors in the absence of conspecifics, as seen

in our study, may allow rats to discern territory and identity information, which may drive social

behavior even when other rats are not present. The precise mechanisms underlying social odor

detection in CA2 neurons are unknown but may be related to the distinctive neurochemical and

structural features of CA2 neurons.

CA2 neurons contain an abundance of neuropeptide receptors, including oxytocin receptors

(OXTR) and vasopressin receptors (AVP1-b) (Mitre et al., 2016; Vaccari et al., 1998; Young and

Song, 2020). Genetic or pharmacological ablation of CA2 OXTRs impairs the persistence of

long-term social recognition memory and the ability to discriminate between social stimuli in

mice (Lee et al., 2008; Lin et al., 2018; Raam et al., 2017). AVP1-b activation in CA2 has been

associated with social aggression (Pagani et al., 2015) and the enhancement of social memory

(Smith et al., 2016). Long-range axonal projections from the paraventricular nucleus of the

hypothalamus (PVN) may be the source of neuropeptide release onto CA2 neurons during

social interactions. The PVN receives input from the olfactory system (Guevara-Aguilar et al.,

1988) and directly innervates CA2 (Cui et al., 2013; Zhang and Hernández, 2013).

Several other brain regions that are involved in social processing send direct inputs to CA2,

including the hypothalamic supramammillary nucleus (SuM) and the lateral entorhinal cortex

(LEC). SuM projections to CA2 are preferentially activated by novel social encounters (Chen et

al., 2020), making SuM inputs an unlikely candidate to induce CA2 place cell responses to a

familiar social odor. Conversely, neuronal activity in the LEC showed similar increases during

social exploration for both novel and familiar conspecifics (Lopez-Rojas et al., 2022). Recent

studies have suggested that the direct projection from LEC to CA2 is essential for social

memory (Dang et al., 2022; Lopez-Rojas et al., 2022). Whether LEC input to CA2 is selectively

enhanced during exploration of a social odor alone remains to be tested, but a more general

involvement of the LEC in odor processing is well documented (Igarashi et al., 2012; Kerr et al.,

2007).

CA2 also receives inputs from the dentate gyrus (DG). The DG directly projects to CA2 via

mossy fibers that form excitatory synapses onto CA2 neurons (Kohara et al., 2014). The DG is

predicted to have a role in pattern separation, a process that reduces overlap between

ensembles of neurons to minimize interference (McClelland and Goddard, 1996). The DG is

important for associating odors with larger contexts in discrimination tests (Morris et al. 2012).

However, whether the DG responds selectively to social odors is unknown and may warrant

further investigation.

Consistent with previous work, our results show that CA2 place cells that remap in response to

a familiar social odor do not return to their original activity, despite the removal of the social odor

in the last session (Alexander et al., 2016). It is possible that the persistent change in CA2

activity is a result of lingering social odors in the arena. Recording CA2 place cells for longer

than 20 minutes after the removal of the odor would address if CA2 place cell activity eventually

returns to baseline, although this would be complicated by the effect of time on CA2 place cells

(Mankin et al., 2015). However, this persistent change in CA2 place cell activity may also

suggest the involvement of synaptic plasticity. Interestingly, CA2 synapses are resistant to

standard long-term potentiation (LTP) compared to other hippocampal subfields (Chang et al.,

2007; Zhao et al., 2007). How, then, are long-lasting changes occurring in CA2 place cells? One

hypothesis is that the release of social neuropeptides, such as vasopressin and oxytocin, can

promote potentiation of CA2 neurons (Lin et al., 2018; Pagani et al., 2015; Tirko et al., 2018).

Selective activation of OXTRs or AVP1-b *in vitro* robustly depolarizes CA2 pyramidal neurons

and lowers the threshold for LTP at excitatory synapses (Dang et al., 2022; Tirko et al., 2018).

As a result, neuropeptide receptor activation may refine the responsiveness of CA2 pyramidal

neurons to synaptic input from upstream structures, thus creating appropriate conditions for

synaptic plasticity.

A question remains of how social information coded by CA2 is transmitted to downstream

regions that control behavior. The main output of dorsal CA2 is dorsal CA1. However, at the

single cell level, dorsal CA1 place cell activity was unaffected by social interactions (Alexander

et al., 2016). Dorsal CA2 neurons also project to ventral CA1, a region that is essential for social

memory (Okuyama et al., 2016). Recent work has shown that ventral CA1 place cells did not

remap to a social stimulus (Wu et al, 2023). However, ventral CA1 place cells became

significantly more spatially selective in an environment that contained a social stimulus

compared to an empty environment (Wu et al. 2023). Further, the activity of a subset of ventral

CA1 neurons was modulated by the presence of a conspecific (Rao et al. 2019; Wu et al. 2023).

Inhibition of dorsal CA2 projections to ventral CA1 impairs social memory (Meira et al., 2018;

Tsai et al., 2022), raising the possibility that social remapping in dorsal CA2 induces the

alterations in neuronal firing in ventral CA1 in response to social stimuli.

Interestingly, recent work has shown that sharp wave-ripples that originate in dorsal CA2 are

important for social memory consolidation (Oliva et al., 2016, 2020). While a portion of the sharp

wave-ripples initiated in dorsal CA2 propagate to other subregions of the dorsal hippocampus,

some do not propagate to dorsal CA1 or dorsal CA3 (Oliva et al., 2016). Given that individual

place cells in dorsal CA1 did not remap to a social stimulus (Alexander et al., 2016), it is

possible that the reactivation of CA2 place cells that respond to a social odor may be limited to

sharp wave-ripples that do not propagate to dorsal CA1. Instead, the function of these sharp

wave-ripples may be to propagate information to ventral CA1 in order to consolidate memories

of a social experience (Meira et al., 2018). Consistent with this idea, the rate of sharp wave-

ripples increased in ventral CA1 when conspecifics were present (Rao et al., 2019). A future

study examining the reactivation of dorsal CA2 place cells that respond to social stimuli while

simultaneously recording place cells in dorsal and ventral CA1 will be essential for our

understanding of how reactivation of CA2 place cells during sharp wave-ripples supports social

memory consolidation.

An important question left unanswered is the extent to which CA2 is specialized for social

memory. While the present results show that a social odor is sufficient to induce CA2 place cell

remapping, it remains unknown whether other odors of similar salience or ethological relevance

have similar effects. Future studies examining the activity of CA2 place cells in response to

other relevant salient nonsocial odors, such as predator urine, will help determine whether CA2

integrates many salient odors into representations of space or whether the role of CA2 is

specific to social information.

The current study shows how presentation of a social odor alone provides a well-controlled

social stimulus to induce changes in place cell activity in dorsal CA2. Use of a social odor can

eliminate confounds that could be introduced by the presence of a conspecific, such as

variations in locomotor behavior, social interactions, and ultrasonic vocalizations. Presentation

of a social odor thereby provides a valuable paradigm for assessing CA2 physiology in rodent

models of diseases and disorders involving aberrant social behaviors, such as autism spectrum

disorders, to better understand how impairments in social behavior may arise.

Table 1. Number of CA2 place cells recorded in each rat and each condition during active

exploration sessions.

| Animal ID | Control | Odor | Visual | Visual + odor | Mirror |
|---|---|---|---|---|---|
| Rat 117 ("Lauren") | 7 | 5 | 7 | 0 | 6 |
| Rat 122 ("Chase") | 51 | 32 | 38 | 0 | 49 |
| Rat 165 ("Gus") | 34 | 56 | 26 | 26 | 35 |
| Rat 256 ("Viggo") | 28 | 13 | 13 | 13 | 21 |
| Rat 391 ("Daffodil") | 15 | 4 | 6 | 5 | 5 |
| Rat 418 ("Hugo") | 20 | 35 | 18 | 41 | 0 |
| **Total** | **155** | **145** | **109** | **85** | **116** |

526 Table 2. Number of CA2 place cells recorded in each rat and each condition during rest periods.

| Animal ID | Control | Odor | Visual | Visual + odor | Mirror |
|---|---|---|---|---|---|
| Rat 117 ("Lauren") | 0 | 0 | 1 | 0 | 0 |
| Rat 122 ("Chase") | 52 | 31 | 2 | 0 | 18 |
| Rat 165 ("Gus") | 26 | 58 | 31 | 23 | 38 |
| Rat 256 ("Viggo") | 18 | 10 | 13 | 8 | 0 |
| Rat 391 ("Daffodil") | 14 | 4 | 7 | 5 | 5 |

| | | | | | |
|---|---|---|---|---|---|
| Rat 418 ("Hugo") | 20 | 29 | 14 | 35 | 0 |
| **Total** | **130** | **132** | **68** | **71** | **61** |

**CRediT authorship contribution statement**

**Emma Robson**: Formal analysis, investigation, writing – original draft, visualization. **Margaret M. Donahue:** Formal analysis, investigation, writing – original draft, visualization, funding acquisition. **Alexandra J. Mably:** Conceptualization, methodology, formal analysis, investigation. **Peyton G. Demetrovich:** Investigation. **Lauren T. Hewitt:** Investigation. **Laura Lee Colgin:** Conceptualization, methodology, resources, writing – review and editing, supervision, project administration, funding acquisition.

**Acknowledgements**

The authors thank Isabella Lee, Misty Hill, Ayomide Akinsooto, Sirisha Dhavala, and Alexa Hassien for technical assistance with this project and Chenguang Zheng for providing MATLAB code for some of the analyses. The authors acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing data storage resources that have contributed to the research described within this article. URL: http://www.tacc.utexas.edu

**Funding**

This research was supported by the Department of Defense (grant number W81XWH1810314) and the National Institutes of Health (grant numbers R56MH125655, R01MH131317, and F31MH127933).

**Data availability**

Data will be made available on request.

## References

Alexander, G. M., Farris, S., Pirone, J. R., Zheng, C., Colgin, L. L., Dudek, S. M., 2016. Social and novel contexts modify hippocampal CA2 representations of space. *Nature Communications*, *7*, 10300. https://doi.org/10.1038/ncomms10300

Breese, C. R., Hampson, R. E., Deadwyler, S. A., 1989. Hippocampal place cells: Stereotypy and plasticity. *Journal of Neuroscience*, *9*(4), 1097–1111. https://doi.org/10.1523/jneurosci.09-04-01097.1989

Chang, P. Y., Taylor, P. E., Jackson, M. B., 2007. Voltage imaging reveals the CA1 region at the CA2 border as a focus for epileptiform discharges and long-term potentiation in hippocampal slices. *Journal of Neurophysiology*, *98*(3), 1309–1322. https://doi.org/10.1152/jn.00532.2007

Chen, S., He, L., Huang, A. J. Y., Boehringer, R., Robert, V., Wintzer, M. E., Polygalov, D., Weitemier, A. Z., Tao, Y., Gu, M., Middleton, S. J., Namiki, K., Hama, H., Therreau, L., Chevaleyre, V., Hioki, H., Miyawaki, A., Piskorowski, R. A., McHugh, T. J., 2020. A hypothalamic novelty signal modulates hippocampal memory. *Nature*, *586*(7828), 270–274. https://doi.org/10.1038/s41586-020-2771-1

Colgin, L. L., Moser, E. I., Moser, M.-B., 2008. Understanding memory through hippocampal remapping. *Trends in Neurosciences*, *31*(9), 469–477. https://doi.org/10.1016/j.tins.2008.06.008

Cui, Z., Gerfen, C. R., Young 3rd, W. S., 2013. Hypothalamic and other connections with dorsal CA2 area of the mouse hippocampus. *Journal of Comparative Neurology*, *521*(8). https://doi.org/10.1002/cne.23263

Dang, R., Zhou, Y., Zhang, Y., Liu, D., Wu, M., Liu, A., Jia, Z., Xie, W., 2022. Regulation of social memory by lateral entorhinal cortical projection to dorsal hippocampal CA2. *Neuroscience Bulletin*, *38*(3), 318–322. https://doi.org/10.1007/s12264-021-00813-6

Dantzer, R., Tazi, A., Bluthé, R.-M., 1990. Cerebral lateralization of olfactory-mediated affective processes in rats. *Behavioural Brain Research*, *40*(1), 53–60. https://doi.org/10.1016/0166-4328(90)90042-D

Ego-Stengel, V., Wilson, M. A., 2010. Disruption of ripple-associated hippocampal activity during rest impairs spatial learning in the rat. *Hippocampus*, *20*(1), 1–10. https://doi.org/10.1002/hipo.20707

Fenton, A. A., Csizmadia, G., Muller, R. U., 2000. Conjoint control of hippocampal place cell firing by two visual stimuli: the effects of moving the stimuli on firing field positions. *Journal of General Physiology*, *116*(2), 191–210. https://doi.org/10.1085/JGP.116.2.191

Gheusi, G., Goodall, G., Dantzer, R., 1997. Individually distinctive odours represent individual conspecifics in rats. *Animal Behaviour*, *53*(5). https://doi.org/10.1006/anbe.1996.0314

Guevara-Aguilar, R., Jimenez-Montufar, L. L., Garcia-Diaz, D. E., Wayner, M. J., Armstrong, D. L., 1988. Olfactory and visceral projections to the paraventricular nucleus. *Brain Research Bulletin*, *20*(6), 799–801. https://doi.org/10.1016/0361-9230(88)90094-9

Hassan, S. I., Bigler, S., Siegelbaum, S. A., 2023. Social odor discrimination and its
enhancement by associative learning in the hippocampal CA2 region. *Neuron*, *111*(14), 2232-
2246.e5. https://doi.org/10.1016/j.neuron.2023.04.026

Hitti, F. L., Siegelbaum, S. A., 2014. The hippocampal CA2 region is essential for social memory.
*Nature*, *508*(7494). https://doi.org/10.1038/nature13028

Hsiao, Y.-T., Zheng, C., Colgin, L. L., 2016. Slow gamma rhythms in CA3 are entrained by slow
gamma activity in the dentate gyrus. *Journal of Neurophysiology*, *116*(6), 2594–2603.
https://doi.org/10.1152/jn.00499.2016

Hurst, J. L., 2005. Scent marking and social communication. In P. K. McGregor (Ed.), *Animal*
*Communication Networks* (pp. 219–244). Cambridge University Press.
https://doi.org/10.1017/CBO9780511610363.014

Husted, J. R., McKenna, F. S., 1966. The use of rats as discriminative stimuli. *Journal of the*
*Experimental Analysis of Behavior*, *9*(6). https://doi.org/10.1901/jeab.1966.9-677

Hwaun, E., Colgin, L. L., 2019. CA3 place cells that represent a novel waking experience are
preferentially reactivated during sharp wave-ripples in subsequent sleep. *Hippocampus*, *29*(10),
921–938. https://doi.org/10.1002/hipo.23090

Igarashi, K. M., Ieki, N., An, M., Yamaguchi, Y., Nagayama, S., Kobayakawa, K., Kobayakawa,
R., Tanifuji, M., Sakano, H., Chen, W. R., Mori, K., 2012. Parallel mitral and tufted cell pathways
route distinct odor information to different targets in the olfactory cortex. *Journal of*
*Neuroscience*, *32*(23), 7970–7985. https://doi.org/10.1523/jneurosci.0154-12.2012

Karlsson, M. P., Frank, L. M., 2009. Awake replay of remote experiences in the hippocampus.
*Nature Neuroscience*, *12*(7), 913–918. https://doi.org/10.1038/nn.2344

Kerr, K. M., Agster, K. L., Furtak, S. C., Burwell, R. D., 2007. Functional neuroanatomy of the
parahippocampal region: The lateral and medial entorhinal areas. *Hippocampus*, *17*(9), 697–
708. https://doi.org/10.1002/hipo.20315

Kohara, K., Pignatelli, M., Rivest, A. J., Jung, H.-Y., Kitamura, T., Suh, J., Frank, D., Kajikawa,
K., Mise, N., Obata, Y., Wickersham, I. R., Tonegawa, S., 2014. Cell type-specific genetic and
optogenetic tools reveal hippocampal CA2 circuits. *Nature Neuroscience*, *17*(2).
https://doi.org/10.1038/nn.3614

Lee, H.-J., Caldwell, H. K., Macbeth, A. H., Tolu, S. G., Young, W. S., 2008. A conditional
knockout mouse line of the oxytocin receptor. *Endocrinology*, *149*(7), 3256–3263.
https://doi.org/10.1210/en.2007-1710

Lin, Y.-T., Hsieh, T.-Y., Tsai, T.-C., Chen, C.-C., Huang, C.-C., Hsu, K.-S., 2018. Conditional
deletion of hippocampal CA2/CA3a oxytocin receptors impairs the persistence of long-term
social recognition memory in mice. *Journal of Neuroscience*, *38*(5).
https://doi.org/10.1523/jneurosci.1896-17.2017

Lopez-Rojas, J., de Solis, C. A., Leroy, F., Kandel, E. R., Siegelbaum, S. A., 2022. A direct
lateral entorhinal cortex to hippocampal CA2 circuit conveys social information required for
social memory. *Neuron*, *110*(9), 1559-1572.e4. https://doi.org/10.1016/j.neuron.2022.01.028

Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S., Leutgeb, J. K., 2015. Hippocampal CA2
activity patterns change over time to a larger extent than between spatial contexts. *Neuron*,
*85*(1). https://doi.org/10.1016/j.neuron.2014.12.001

McClelland, J. L., Goddard, N. H., 1996. Considerations arising from a complementary learning
systems perspective on hippocampus and neocortex. *Hippocampus*, *6*(6), 654–665.
https://doi.org/10.1002/(SICI)1098-1063(1996)6:6<654::aid-hipo8>3.0.co;2-g

Meira, T., Leroy, F., Buss, E. W., Oliva, A., Park, J., Siegelbaum, S. A., 2018. A hippocampal
circuit linking dorsal CA2 to ventral CA1 critical for social memory dynamics. *Nature
Communications*, *9*(1), 1–14. https://doi.org/10.1038/s41467-018-06501-w

Mitre, M., Marlin, B. J., Schiavo, J. K., Morina, E., Norden, S. E., Hackett, T. A., Aoki, C. J.,
Chao, M. V., Froemke, R. C., 2016. A distributed network for social cognition enriched for
oxytocin receptors. *Journal of Neuroscience*, *36*(8), 2517–2535.
https://doi.org/10.1523/jneurosci.2409-15.2016

Morris, A. M., Weeden, C. S., Churchwell, J. C., Kesner, R. P., 2013. The role of the dentate
gyrus in the formation of contextual representations. *Hippocampus, 23(2),* 162–168.
https://doi.org/10.1002/hipo.22078

Muller, R. U., Kubie, J. L., Ranck, J. B., 1987. Spatial firing patterns of hippocampal complex-
spike cells in a fixed environment. *Journal of Neuroscience, 7*(7), 1935–1950.
https://doi.org/10.1523/jneurosci.07-07-01935.1987

Oettl, LL., Kelsch, W. (2017). Oxytocin and Olfaction. In: Hurlemann, R., Grinevich, V. (eds)
*Behavioral Pharmacology of Neuropeptides: Oxytocin*. Current Topics in Behavioral
Neurosciences, vol 35. Springer, Cham. https://doi.org/10.1007/7854_2017_8

O'Keefe, J., 1976. Place units in the hippocampus of the freely moving rat. *Experimental
Neurology*, *51*(1), 78–109. https://doi.org/10.1016/0014-4886(76)90055-8

O'Keefe, J., Conway, D. H., 1978. Hippocampal place units in the freely moving rat: Why they
fire where they fire. *Experimental Brain Research*, *31*(4), 573–590.
https://doi.org/10.1007/BF00239813

O'Keefe, J., Dostrovsky, J., 1971. The hippocampus as a spatial map: Preliminary evidence
from unit activity in the freely-moving rat. *Brain Research*, *34*, 171–175.
https://doi.org/10.1016/0006-8993(71)90358-1

Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S., Tonegawa, S., 2016. Ventral CA1 neurons
store social memory. *Science*. https://doi.org/10.1126/science.aaf7003

Oliva, A., 2022. CA2 physiology underlying social memory. *Current Opinion in Neurobiology*, *77*,
102642. https://doi.org/10.1016/j.conb.2022.102642

Oliva, A., Fernández-Ruiz, A., Buzsáki, G., Berényi, A., 2016. Role of hippocampal CA2 region
in triggering sharp-wave ripples. *Neuron*, *91*(6), 1342–1355.
https://doi.org/10.1016/j.neuron.2016.08.008

Oliva, A., Fernandez-Ruiz, A., Karaba, L. A., 2023. CA2 orchestrates hippocampal network
dynamics. *Hippocampus*, *33*(3), 241–251. https://doi.org/10.1002/hipo.23495

Oliva, A., Fernández-Ruiz, A., Leroy, F., Siegelbaum, S. A., 2020. Hippocampal CA2 sharp-wave
ripples reactivate and promote social memory. *Nature*, *587*(7833), 264–269.
https://doi.org/10.1038/s41586-020-2758-y

Pagani, J. H., Zhao, M., Cui, Z., Avram, S. K. W., Caruana, D. A., Dudek, S. M., Young, W. S.,
2015. Role of the vasopressin 1b receptor in rodent aggressive behavior and synaptic plasticity
in hippocampal area CA2. *Molecular Psychiatry*, *20*(4), 490–499.
https://doi.org/10.1038/mp.2014.47

Popik, P., van Ree, J. M., 1998. Neurohypophyseal peptides and social recognition in rats.
*Progress in Brain Research*, *119*, 415–436. https://doi.org/10.1016/s0079-6123(08)61585-x

Popik, P., Vetulani, J., Bisaga, A., van Ree, J. M., 1991. Recognition cue in the rat's social
memory paradigm. *Journal of Basic and Clinical Physiology and Pharmacology*, *2*(4), 315–327.
https://doi.org/10.1515/jbcpp.1991.2.4.315

Raam, T., McAvoy, K. M., Besnard, A., Veenema, A. H., Sahay, A., 2017. Hippocampal oxytocin
receptors are necessary for discrimination of social stimuli. *Nature Communications*, *8*(1), 2001.
https://doi.org/10.1038/s41467-017-02173-0

Ramadan, W., Eschenko, O., Sara, S. J., 2009. hippocampal sharp wave/ripples during sleep
for consolidation of associative memory. *PLoS ONE*, *4*(8), e6697.
https://doi.org/10.1371/journal.pone.0006697

Rao, R. P., Heimendahl, M. von, Bahr, V., Brecht, M., 2019. Neuronal responses to conspecifics
in the ventral CA1. *Cell Reports*, *27*(12). https://doi.org/10.1016/j.celrep.2019.05.081

Sawyer, T. F., Hengehold, A. K., Perez, W. A., 1984. Chemosensory and hormonal mediation of
social memory in male rats. *Behavioral Neuroscience*, *98*(5). https://doi.org/10.1037//0735-
7044.98.5.908

Smith, A. S., Williams Avram, S. K., Cymerblit-Sabba, A., Song, J., Young, W. S., 2016. Targeted
activation of the hippocampal CA2 area strongly enhances social memory. *Molecular*
*Psychiatry*, *21*(8), 1137–1144. https://doi.org/10.1038/mp.2015.189

Stevenson, E. L., Caldwell, H. K., 2014. Lesions to the CA2 region of the hippocampus impair
social memory in mice. *The European Journal of Neuroscience*, *40*(9), 3294–3301.
https://doi.org/10.1111/ejn.12689

Thompson, L. T., Best, P. J., 1990. Long-term stability of the place-field activity of single units
recorded from the dorsal hippocampus of freely behaving rats. *Brain Research*, *509*(2), 299–
308. https://doi.org/10.1016/0006-8993(90)90555-P

698 Thor, D. H., Holloway, W. R., 1982. Social memory of the male laboratory rat. *Journal of*
699 *Comparative and Physiological Psychology*, *96*(6). https://doi.org/10.1037/0735-7036.96.6.1000

700 Tirko, N. N., Eyring, K. W., Carcea, I., Mitre, M., Chao, M. V., Froemke, R. C., Tsien, R. W.,
701 2018. Oxytocin transforms firing mode of CA2 hippocampal neurons. *Neuron*, *100*(3), 593-
702 608.e3. https://doi.org/10.1016/j.neuron.2018.09.008

703 Tsai, T.-C., Fang, Y.-S., Hung, Y.-C., Hung, L.-C., Hsu, K.-S., 2022. A dorsal CA2 to ventral CA1
704 circuit contributes to oxytocinergic modulation of long-term social recognition memory. *Journal*
705 *of Biomedical Science*, *29*(1), 50. https://doi.org/10.1186/s12929-022-00834-x

706 Vaccari, C., Lolait, S. J., Ostrowski, N. L., 1998. Comparative distribution of vasopressin V1b
707 and oxytocin receptor messenger ribonucleic acids in brain. *Endocrinology*, *139*(12).
708 https://doi.org/10.1210/endo.139.12.6382

709 Wersinger, S. R., Ginns, E. I., O'Carroll, A.-M., Lolait, S. J., Young III, W. S., 2002. Vasopressin
710 V1b receptor knockout reduces aggressive behavior in male mice. *Molecular Psychiatry*, *7*(9),
711 975–984. https://doi.org/10.1038/sj.mp.4001195

712 Wersinger, S. R., Temple, J. L., Caldwell, H. K., Young, W. S., 3rd., 2008. Inactivation of the
713 oxytocin and the vasopressin (Avp) 1b receptor genes, but not the Avp 1a receptor gene,
714 differentially impairs the bruce effect in laboratory mice (mus musculus). *Endocrinology*, *149*(1).
715 https://doi.org/10.1210/en.2007-1056

716 Widloski, J., & Foster, D. J., 2022. Flexible rerouting of hippocampal replay sequences around
717 changing barriers in the absence of global place field remapping. *Neuron, 110*(9), 1547-1558.
718 https://doi.org/10.1016/j.neuron.2022.02.002

719 Wilson, M. A., McNaughton, B. L., 1994. Reactivation of hippocampal ensemble memories
720 during sleep. *Science, 265*(5172), 676–679. https://doi.org/10.1126/science.8036517

721 Wintzer, M. E., Boehringer, R., Polygalov, D., McHugh, T. J., 2014. The hippocampal CA2
722 ensemble is sensitive to contextual change. *Journal of Neuroscience*, *34*(8).
723 https://doi.org/10.1523/jneurosci.2563-13.2014

724 Wu, W.-Y., Yiu, E., Ophir, A. G., Smith, D. M., 2023. Effects of social context manipulation on
725 dorsal and ventral hippocampal neuronal responses. *Hippocampus, 33(7)*, 830–843.
726 https://doi.org/10.1002/hipo.23507

727 Young, W. S., Song, J., 2020. Characterization of oxytocin receptor expression within various
728 neuronal populations of the mouse dorsal hippocampus. *Frontiers in Molecular Neuroscience*,
729 *13*. https://doi.org/10.3389/fnmol.2020.00040

730 Zhang, L., Hernández, V. S., 2013. Synaptic innervation to rat hippocampus by vasopressin-
731 immuno-positive fibres from the hypothalamic supraoptic and paraventricular nuclei.
732 *Neuroscience*, *228*, 139–162. https://doi.org/10.1016/j.neuroscience.2012.10.010

Zhao, M., Choi, Y.-S., Obrietan, K., Dudek, S. M., 2007. Synaptic plasticity (and the lack thereof)
in hippocampal CA2 neurons. *Journal of Neuroscience*, *27*(44), 12025–12032.
https://doi.org/10.1523/jneurosci.4094-07.2007

Zheng, C., Bieri, K. W., Hsiao, Y.-T., Colgin, L. L., 2016. Spatial sequence coding differs during
slow and fast gamma rhythms in the hippocampus. *Neuron*, *89*(2), 398–408.
https://doi.org/10.1016/j.neuron.2015.12.005

Zhu, N., Zhang, Y., Xiao, X., Wang, Y., Yang, J., Colgin, L. L., Zheng, C., 2023. Hippocampal
oscillatory dynamics in freely behaving rats during exploration of social and non-social stimuli.
*Cognitive Neurodynamics*, *17*(2), 411–429. https://doi.org/10.1007/s11571-022-09829-8

Figure 1. Behavioral task. Rats freely explored a familiar open field arena for four 20-minute sessions per day. A stimulus cage, a standard rat housing cage, was placed in a corner of the arena. In the first and final sessions of experimental conditions (i.e., Odor, Visual + Odor, Visual, Mirror), and in all four sessions of the Control condition, the cage contained only clean bedding. In the middle two sessions of experimental conditions, a cage containing social stimuli was presented. In the Odor condition, a stimulus cage containing soiled bedding of a familiar rat or rats was presented. In the Visual + Odor condition, a cage containing a familiar rat and the rat's soiled bedding was presented. In the Visual condition, a familiar rat was placed in the stimulus cage with clean bedding and a filter-top lid to minimize social odors. In the Mirror condition, a familiar rat was placed in the stimulus cage with clean bedding, and the cage was lined with a one-way mirror attachment. This prevented the stimulus rat from seeing the implanted rat, thereby limiting reciprocal social interactions.

Figure 2. Histology. An example hippocampal section showing immunohistological identification of a tetrode track in CA2. To identify CA2, hippocampal sections were immunostained with a CA2 marker, Purkinje cell protein 4 (PCP4, red). DAPI nuclear staining is shown in blue.

Figure 3. Example CA2 place cell firing rate maps across the four recording sessions for all conditions. Color-coded firing rate maps are shown for all place cells recorded on a single example tetrode in each condition for one example rat (Rat 165). Rate maps are shown scaled to the maximal firing rate (shown inset) of each cell across all sessions. White pixels indicate places that were not visited by the rat during a session.

Figure 4. Changes in CA2 place cell firing patterns in response to social stimuli. A. The estimated means of spatial correlation coefficients from our generalized linear mixed model are shown across each condition and session pair. Error bars represent 95% confidence intervals.

B. The estimated means of spatial correlation coefficients from our generalized linear mixed model are shown across each condition for all session pairs combined. C-G. Spatial correlation measures are shown for the entire sample of CA2 place cells for all pairs of sessions across all conditions. Each marker represents a spatial correlation value for an individual place cell. Different symbols are used for CA2 place cells recorded from different rats.

Figure 5. CA2 place cells did not show a coherent place field shift related to the location of the stimulus cage. A. Shown are the distances measured between the positions of peak firing in Session B and the stimulus cage for CA2 place cells that "turn on", or gain a field, in Session B. B. Shown are the distances between the peak firing rate position in Session A and the location of the stimulus cage for CA2 place cells that "turn off", or lose a field, in Session B. C. Shown are the changes in distance from the stimulus cage of the peak firing rate positions in Sessions A and B for CA2 place cells that were active in both Sessions A and B but shifted their place field locations. Individual markers represent measurements from individual place cells. Different symbols represent distance measurements for CA2 place cells recorded from different rats.

Figure 6. Exploration of social stimuli. A. Heat maps show mean exploration times of different locations in the arena, with the stimulus cage shown in the top right corner of the arena. Time spent was calculated for the first 2 minutes of each session individually and then averaged within each rat. Heat maps shown are averaged across rats. B. Time spent exploring locations close to the cage (within 12 cm) was calculated for each session and then averaged within a rat. Individual dots represent the mean exploration time for each rat, and error bars represent 95% confidence intervals across all sessions within a rat. Rats increased their exploration time of the stimulus cage in session B for the Odor, Visual, and Mirror conditions compared to the Control condition.

Figure 7. CA2 place cells that responded to a social odor preferentially fired during sharp wave-ripples. A. The difference between peak normalized firing rates of CA2 place cells during sharp wave-ripples in the rest period following session B compared to peak normalized firing rates of CA2 place cells during sharp wave-ripples in the rest period following session A was positively correlated to the selectivity index for the Odor condition (B) but not the Control condition (A). The selectivity index indicates a cell's preference for session A (value of -1 for maximum selectivity in session A) or session B (value of 1 for maximum selectivity in session B).

A   Control

B   Odor

C   Visual + Odor

D   Visual

E   Mirror

Firing rate

Max