The asymmetric expression of HSPA2 in blastomeres governs the first embryonic cell-fate decision

Jiayin Gao [1,2,3,4,5,6,7,8†], Jiawei Wang[1,2,3,4,5,6,7,8†], Shiyu Liu[1,2,3,4,5,6,7,8], Jinzhu Song[1,2,3,4,5,6,7,8], Chuanxin Zhang[1,2,3,4,5,6,7,8], Boyang Liu[1,2,3,4,5,6,7,8*], Keliang Wu [1,2,3,4,5,6,7,8*]

Address: [1] Institute of Women, Children and Reproductive Health, Shandong University, 250012, China. [2] State Key Laboratory of Reproductive Medicine and Offspring Health, Shandong University, Jinan, Shandong, 250012, China. [3]National Research Center for Assisted Reproductive Technology and Reproductive Genetics, Shandong University, Jinan, Shandong, 250012, China. [4]Key Laboratory of Reproductive Endocrinology (Shandong University), Ministry of Education, Jinan, Shandong, 250012, China. [5]Shandong Technology Innovation Center for Reproductive Health, Jinan, Shandong, 250012, China. [6]Shandong Provincial Clinical Research Center for Reproductive Health, Jinan, Shandong, 250012, China. [7]Shandong Key Laboratory of Reproductive Medicine, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong, 250012, China. [8]Research Unit of Gametogenesis and Health of ART-Offspring, Chinese Academy of Medical Sciences (No.2021RU001), Jinan, Shandong, 250012, China.

[†] These authors contributed equally to this study.

*Corresponding author: Boyang Liu and Keliang Wu.

Correspondence address: byliu@sdu.edu.cn (B.L.), wukeliang@sdu.edu.cn (K.W.).

Abstract: The first cell-fate decision is the process by which cells of an embryo take on distinct lineage identities for the first time, thus representing the beginning of developmental patterning. Here, we demonstrate that the molecular chaperone heat shock protein A2 (HSPA2), a member of the 70 kDa heat shock protein (HSP70) family, is asymmetrically expressed in the late 2-cell stage of mouse embryos. The knockdown of *Hspa2* in one of the 2-cell blastomeres prevented its progeny predominantly toward the inner cell mass (ICM) fate. In contrast, the overexpression of *Hspa2* in one of the two-cell blastomeres did not induce blastomeres to differentiate towards the ICM fate. Furthermore, we demonstrated that HSPA2 interacts with CARM1 and its levels correlate with ICM-associated genes and that it interacts with the CARM1. Collectively, our results identify HSPA2 as a critical early regulator of the first cell-fate decision in mammalian 2-cell embryos.

Key words: first cell-fate decision, inner cell mass, HSPA2, CARM1, embryo

1. INTRODUTION

The preimplantation development of mammalian embryos is a complex and orderly process[1]. The first cell-fate decision is a key event in the preimplantation development in mammalian embryos. Specifically, the first cell-fate decision occurs at the morula stage following embryonic compaction and results in the division of embryonic cells into two distinct lineages: the trophectoderm (TE) and the inner cell mass (ICM). The TE is a single layer surrounding the fluid-filled cavity known as the blastocoel that provides extraembryonic structures such as the placenta, whereas the ICM, which is attached to the inside of the TE, contains pluripotent cells that give rise

to the fetus and extraembryonic tissues [2-4]. However, it remains an open question of

how these cells shape the first cell-fate decision of preimplantation development, and

when they begin to trigger the initial differentiation signals.

The asymmetry of the blastocyst may be traced back to the asymmetric nuclear

abundance of coactivator-associated arginine methyltransferase 1 (CARM1) at the 4-

cell stage [5-7]. The heterogeneous activity of CARM1 leads to the differential

methylation of H3 methylation of arginine 26 (H3R26me). More extensive

methylation of H3R26me can result in a higher increase in the expression levels of

ICM-specific genes (e.g., *Oct4*, *Nanog* and *Sox2*) and global chromatin accessibility,

which directs their progeny into the ICM fate [8]. Recent studies have further

suggested that asymmetry of the blastomere can even be traced back to the 2-cell

stage. For example, a long non-coding RNA, *LincGET*, is known to be

asymmetrically expressed in the nuclei of blastomeres in 2 and 4-cell embryos.

Specifically, *LincGET* and CARM1 are known to form a complex that promotes the

H3 methylation of arginine 26 (H3R26me) and activates the expression of genes in

the ICM [9]. In addition, HMGA1, a member of the high mobility group (HMG)

protein family, is also known to exhibit heterogeneity as early as the two-cell stage

and involves in cell fate differentiation [10]. Collectively, these studies suggest that

molecular heterogeneity takes place in the early stage of embryonic development and

is associated with cell fate differentiation.

Despite these findings, the specific factors that exhibit heterogeneity at the two-

cell stage and the mechanisms by which the cell fate and differentiation can be

decided still require further investigation. The molecular chaperone heat shock protein

A2 (HSPA2) is a member of the 70 kDa heat shock protein (HSP70) family that is

evolutionarily conserved in human and metazoan. It is originally identified in male

germ cells and described as a testis-specific protein that is essential for

spermatogenesis [11, 12]. The *Hspa2* gene is located on chromosome 14

(14q24.1)[13], and the aberrant expression of HSPA2 in testes was shown to induces

primary spermatocytes to arrest in meiosis I and undergo apoptosis, thus leading to

male infertility [14, 15]. HSPA2 is expressed in somatic tissues and during mouse

embryogenesis [16, 17], and high levels of HSPA2 expression play a critical role in

the genesis and progression of carcinoma [18, 19]. However, the role of HSPA2 in

lineage differentiation of embryonic cells is yet to be unveiled.

In this study, we separated blastomeres from the 4-cell stage and cultured to the

morula stage, analyzed the gene expression pattern by using Smart-seq2 to identify

differentially expressed genes. *Hspa2* was identified and was shown to be

asymmetrically distributed in two- to four-cell blastomeres. The knockdown of *Hspa2*

in one of the 2-cell blastomeres prevented its progeny predominantly towards the ICM

fate. Furthermore, HSPA2 and CARM1 were found to form a protein complex and

regulated ICM-specific gene expression. Collectively, these results suggest that

Hspa2 expression patterns can bias cell fate in the mouse embryo.

2. MATERIALS AND METHODS

2.1 Antibodies

In this study, we utilized CDX2 (ab76541), NANOG (ab17338) and H3R26me2

(ab127095) antibodies from Abcam; a CARM1 (12495s) antibody from Cell Signaling Technology; SOX2 (11064-1-AP), GAPDH (10494-1-AP), CARM1 (55246-1-AP), HSPA2 (12797-1-AP) and ACTIN (66009-1-Ig) antibodies from Proteintech; and an OCT4 (sc-5279) antibody from Santa Cruz Biotech.

2.2 Mouse embryo collection

Animal experiments were approved by the Ethics Committee of Shandong University. ICR mice (6–8 weeks-of-age) were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. and bred under standard SPF conditions with a 12 h dark and 12 h light cycle with food and water provided ad libitum). Superovulation was induced by an intraperitoneal injection of 5 IU of pregnant mare's serum gonadotrophin (PMSG) (110914564, Ningbo Sansheng Biological Technology Co., Ltd) followed 46–48 h later by 5 IU of human chorionic gonadotrophin (HCG) (110911282, Ningbo Sansheng Biological Technology Co., Ltd.). Oocytes were collected 15 h after hCG injection. First, the collected oocytes were fertilized in G-IVF (#10,136, Vitrolife) with epididymal sperm from adult males that had been capacitated in G-IVF for 1 h. All zygotes were cultured in G1 medium (Vitrolife, Sweden) at 37 °C in the presence of 6% CO_2 and 5% O_2. Early 2-, late 2-, 4-, and 8-cell embryos, and morula- and blastocyst stage embryos were collected after 22–26, 34-38, 48–50, 60-65, 70–75, and 96–100 h of culture, respectively. Single blastomeres were isolated from 2- and 4-cell embryos and cultured to the morula stage as described previously[20].

2.3 Cell culture and transfection

Mouse ESCs (mESCs) were cultured on plates coated with 0.1% gelatin (Oricell)

under feeder-free conditions with 5% CO2 at 37 ℃. For cell transfection, the mESCs

were transfected with si-*Hspa2* using Lipofectamine 3000 (Invitrogen) according to

the manufacturer's instructions. After transfection for 36 hours, cells were collected

for further analysis.

2.4 RT-qPCR analysis

Table 1. List of mouse primer pairs used for RT-qPCR analysis

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
Hspa2	AAGATTTCTTCAACGGCAAGGAG	GGGATCGTGGTGTTTCTCTTGAT
Oct4	GGGTGATGGGTCAGCAG	TCCGCAGAACTCGTATGC
Sox2	AACGCCTTCATGGTATGG	CTCGGTCTCGGACAAAAG
Nanog	CTGCTCCGCTCCATAACT	GGCTTTCCCTAGTGGCTT
Cdx2	GAAGGGGTGGTGGGTTC	AGGTTGGCTCTGGCATTT
Actin	GCTCTTTTCCAGCCTTCCTT	GTACTTGCGCTCAGGAGGAG
Hprt	CGAGATGTGATGAAGGAAATGG	AGCAGGTCAGCAAAGAATTTATAGC

For RT-PCR analysis, we used the Single Cell Sequence Specific Amplification Kit

(Vazyme, P621-01) and ChamQ Universal SYBR qPCR Master Mix (Vazyme, Q711-

03) in accordance with the manufacturer's instructions. Primers for the candidate

genes were designed by online Primer 3.0 software based on the sequence data

obtained from the NCBI database. Primer sequences are listed in Table 1. Gene

expression was quantified by the comparative CT method. Values are expressed as the

mean ± standard deviation (SD). Expression was normalized to Actin. The relative

amount of transcript present in each cDNA sample was calculated using the $2^{-\Delta\Delta CT}$

method. All experiments were repeated at least three times, and statistical analysis

was performed on individual experimental sets. All measurements were performed in

triplicate for each experiment.**2.5 Automated western immunoblotting**

Western immunoblots were performed on a PeggySue (ProteinSimple) system

using a Size Separation Master Kit with Split Buffer (12–230 kDa) in accordance with

the manufacturer's instructions. Capillary-based immunoassays were performed using

the Wes-Simple Western method (Proteintech: HSPA2 (12797-1-AP), 1:15 dilution;

GAPDH (10494-1-AP),1:35 dilution; SOX2 (11064-1-AP), 1:20 dilution; Abcam:

CDX2 (ab76541), 1:10 dilution; Santa Cruz Biotech: OCT4 (sc-5279), 1:15 dilution)

and an anti-rabbit or an-mouse detection module (ProteinSimple) [21]. Protein

expression was detected by chemiluminescence and quantified by area under the

curve (AUC) analysis using the Compass for Simple Western program

(ProteinSimple).

2.6 Microinjection and HSPA2 knockdown

Aliquots of 2–5 pl of siRNA at 20 μM concentration were microinjected into the

nucleus of zygote or one blastomere of the two-cell embryos using an Eppendorf

(Hamburg, Germany) micromanipulator under a Leica inverted microscope, and then

cultured in fresh G1 medium.

To knockdown Hspa2, two siRNAs were designed for each gene. The siRNAs of

Hspa2 are GGTGCAACAACTTAGTTTA, GCACTGCAGTGATATTAAA. For

microinjection in each KD group,20 μM of each siRNA was used. siRNAs were

ordered from RINOBIO.

2.7 Western blot analysis and co-immunoprecipitation (Co-IP)

Protein concentration was determined with a BCA protein assay (Bio-Rad,

Hemel Hempstead, UK). Proteins were then separated by 12% SDS-PAGE and

transferred to polyvinylidene difluoride membranes (Millipore, Bedford, MA, USA)

at 300mA for 90min. First, the membranes were incubated in TBST containing 5%

non-fat milk for 2h; then, the membranes were incubated with primary antibodies

(Abcam: CDX2 (ab76541), 1:1000 dilution; NANOG (ab17338) 1:800 dilution; Cell

Signaling Technology: CARM1 (12495s), 1:1000 dilution; Proteintech: SOX2

(11064-1-AP), 1:800 dilution; GAPDH (10494-1-AP), 1:1500 dilution; HSPA2

(12797-1-AP), 1:1000 dilution; ACTIN (66009-1-Ig), 1:700 dilution; Santa Cruz

Biotech: OCT4 (sc-5279), 1:800 dilution) overnight at 4°C. On the second day, the

membranes were washed three times with TBST and then incubated with horseradish

peroxidase-conjugated secondary antibody (1:2000 dilution; Proteintech) for 1 h.

149   The Pierce Co-IP kit (Thermo Fisher Scientific, USA) was used for co-IP assays.

150 In brief, mESCs were lysed using IP lysis buffer on ice. Next, the lysate was treated

151 with agarose resin at 4°C for 1 h followed by incubation with antibodies (Proteintech:

152 HSPA2(12797-1-AP), 1:100 dilution; CARM1(55246-1-AP), 1:60 dilution) at 4°C

153 overnight. The captured antigen was then eluted and SDS-PAGE was performed.

154 **2.8 RNA FISH and immunofluorescence**

155   Fluorescence *in situ* hybridization (FISH) was carried out using an RNA-FISH

Probe kit (GenePharma, Shanghai), according to the manufacturer's instruction. In

brief, embryos were collected and fixed in 4% paraformaldehyde in PBS-DEPC for

15 min at room temperature, permeabilized in 0.1%BufferA (permeabilizing solution)

for 15 min at room temperature, and incubated at 37°C in 2× Buffer C for 30 min. The

embryos were then dehydrated in 70%, 85%, and 100% EtOH and dried at room

temperature for 5 min each time. The embryos were denatured in buffer E

(hybridization buffer) containing 10 μM probe at 73°C for 5 min, incubated at 37°C

overnight, and washed in 0.1% Buffer F for 5 min to remove non-specific binding.

After washing in 2× Buffer C and 1× Buffer C (neutral solution) for 5 min, the nuclei

were stained with DAPI for 5 min

To perform immunofluorescence, we removed the zona pellucida with acid

Tyrode's solution (Sigma). After that, embryos were fixed in 4% paraformaldehyde in

PBS (KGB5001, KeyGEN BioTECH) at room temperature (RT). Embryos were then

permeabilized with 1% Triton-100 (T8787, SigmaAldrich) in PBS for 20 min.

Embryos were blocked for 30 min at RT on a shaker with a blocking buffer consisting

of 1% BSA in PBS containing 0.1% Tween-20 (P9416, Sigma-Aldrich) and 0.01%

Triton-100. Next, embryos were incubated with primary antibodies (Abcam: CDX2

(ab76541), 1:150 dilution; H3R26me2 (ab127095), 1:150 dilution; Santa Cruz

Biotech: OCT4 (sc-5279), 1:100 dilution) diluted in blocking buffer on a shaker at 4-°C

overnight. On the second day, embryos were washed three times with TBST and

incubated with secondary antibody (1:400 dilution; Sigma) and DAPI (D3571, 1:400

dilution; Life Technologies) for 30 min. First, embryos were rinsed three times for 10

min in washing buffer on a shaker at RT; then, the embryos were mounted on slides in

washing buffer and examined under a laser scanning confocal microscope (Dragonfly,

Andor Technology, UK). Fluorescence images were acquired at multiple $1-\mu m$

intervals in the z-axis with the use of a confocal microscope. Multiple images were

processed using ImageJ software for intensity measurement and cell counting

(https://imagej.nih.gov/ij/). The fluorescence intensity was quantified by

normalization to DAPI using the built-in ImageJ function. The regions of interest

(ROI) were selected using a custom Python script. The intensity was measured on

normalized sections using the ImageJ measure function. Data were normalized with

respect to background levels. To analyze the ICM/TE cell numbers, OCT4 and CDX2

signal at inner layers and outer layers were used to distinguish the ICM and TE cells.

2.9 Molecular docking analysis

For molecular docking analysis, we first generated a PPI network and identified

core targets. The proposed protein model was based on the crystal structure

downloaded from the AlphaFold Protein Structure Database

(https://alphafold.ebi.ac.uk/) and the two- and three-dimensional (2D/3D) structures

of potential active compounds were acquired from PubChem

(https://pubchem.ncbi.nlm.nih.gov/). Then, molecular docking analysis was

performed and visualized in Maestro11.9; the resultant docking score was saved and

then compared.

2.10 Proteomics and bioinformatic analysis

For proteomics and bioinformatic analysis, 4 embryos per sample were collected.

Four replicates were assessed for NC group and two replicates were assessed for KD

group. Protein samples were sonicated three times on ice using a high intensity

ultrasonic processor (Scientz) in lysis buffer (8 M urea, 1% protease inhibitor

cocktail). Inhibitors were also added to lysis buffer for PTM experiments. The

remaining debris was removed by centrifugation at 12,000 g at 4°C for 10 min. The

protein samples were then diluted by adding 200 mM of TEAB to urea concentrations

< 2 M. Finally, trypsin was added at a 1:50 trypsin-to-protein mass ratio for the first

digestion overnight and a 1:100 trypsin-to-protein mass ratio for a second 4 h-

digestion. Finally, the peptides were desalted with a Strata X SPE column. The eluted

peptides were then cleaned with C18 ZipTips (Millipore) in accordance with the

manufacturer's instructions, followed by analysis with LC–MS/MS. The resulting

MS/MS data were processed by MaxQuant with an integrated Andromeda search

engine (version 1.4.1.2). False discovery rate thresholds for proteins, peptides and

modification sites were specified at 1%.

COG, full name is Clusters of Orthologous Groups of proteins. The proteins that

make up each COG are assumed to be derived from an ancestral protein, Orthologs

are proteins that are derived from different species, evolved from vertical lineages

(species formation), and typically retain the same function as the original protein.

COG means "cluster of homologous proteins" in Chinese. COGs are divided into two

categories, one for prokaryotes and the other for eukaryotes. Those for prokaryotes

are generally referred to as COG databases; those for eukaryotes are generally

referred to as KOG databases. Compared to other databases, such as NCBI's COG

database, EggNOG provides a more comprehensive classification of species and more

homologous protein sequences, with phylogenetic tree construction and functional

annotation for each homologous gene cluster.

2.11 RNA-seq analysis

For RNA sequencing, 3 embryos per sample were collected. Three replicates were

assessed for NC group and three replicates were assessed for KD group. After being

denuded of the zona pellucida using EmbryoMax Acidic Tyrode's Solution (MR-004-

D, Sigma-Aldrich), RNA-seq libraries were generated following the Smart-seq2

protocol as described previously[22]. Sequencing libraries were generated using

TruePep™DNA Library Prep Kit V2 for Illumina® (TD502, Vazyme) according to

manufacturer's recommendations. The libraries were performed on an Illumina Hiseq

X-ten platform with 150 bp paired-end.

Filtering out low quality and joint sequences, the clean reads of each sample

obtained were mapped to the reference genome downloaded from the genome website

(https://ftp.ensembl.org/pub/release-

105/fasta/mus_musculus/dna/Mus_musculus.GRCm39.dna.primary_assembly.fa.gz)

using Hisat2. The expression values for each gene were calculated as Fragments Per

Kilobase of exon model per Million mapped fragments (TPM+1). DEG analysis was

conducted with edgeR package. In this project, the gene were identified as

differentially expressed genes (DEGs) if $|\log2FC| > 1$ and $P < 0.05$ (FC: fold change).

Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG)

enrichment analyses were s carried out for all the DEGs, utilizing the "cluster Profiler"

package.

2.12 Statistical analysis

Statistical analyzes were performed by GraphPad Prism software (version 9.3,

GraphPad Prism Software Inc., San Diego, CA) and the Statistics Package for Social

Science (SPSS 11.5; SPSS Inc., Chicago, IL, USA). Statistical significance (Figures

1E-J, 2G,I-M, 3B,3E,4B-D,4F,5B-E,5H, S1C-J, S2G) were calculated with two tailed

Student's t test. Levels of significance (Figures 1C) were calculated with t test. At

least three biological replicates were performed for each experiment and results are

represented as mean $\pm$ standard error of mean (SEM). $p < 0.05$ was considered

statistically significant.

2.13 Data access

The RNA-sequencing data used in this study were obtained from the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession number (GSE266025). Proteomic data were available via ProteomeX change with identifier PXD052193.

3. RESULTS

3.1 *Hspa2* is asymmetrically expressed between late two- and four-cell blastomeres

In our previous study, we established a culture system that allows blastomeres without their zona pellucida to develop in vitro until the blastocyst stage [20]. Previous studies have mainly focused on gene heterogeneity in 4-cell stage mouse embryos[5, 6]. Here, in order to expand the heterogeneity between blastomeres and identify factors that may involve in cell lineage differentiation, single blastomeres from the mouse 4-cell embryo were separated and cultured to the morula stage. By using Smart-seq2 technology, we analyzed the gene expression pattern of the morulae, which developed from the 4-cell blastomeres (Figure 1A). *Hspa2* was identified by differential gene expression analysis and showed distinct mRNA levels among the four morulae (Figure 1B). In addition, the level of *Nanog* and *Hspa2* mRNA showed stronger positive correlation (R=0.624) in gene expression than random gene pairs (R=0.029), indicating that *Nanog* expression levels increased with the increase of *Hspa2* (Figure 1C). When the late 2-cell embryo was divided into single blastomeres and cultured to the morula stage, we obtained similar results that *Hspa2* mRNA

distinctly distributed between the two morulae (Figure S1A, B). Next, we investigated the expression of *Hspa2* during preimplantation embryo development from meiosis II to the blastocyst stage. qRT-PCR analysis showed that *Hspa2* mRNA was stored maternally at the MII oocyte and decreased at the two-cell stage but remarkably increased from the 2-cell to the 4-cell stage, coinciding with zygotic genome activation (ZGA). There was a slight change in expression as embryos developed to the 8-cell and morula stages, and developed into blastocysts that had higher expression levels than before the 2-cell stage (Figure 1D).

To directly examine the heterogeneity among individual blastomeres, we separated 4-cell embryos into individual four blastomeres which were then analyzed by single-cell qRT-PCR and Jess TM Simple Western System based automated western immunoblotting with high sensitivity. The results showed that the mRNA and protein levels of HSPA2 differed among the four blastomeres (Figure 1E, F), and confirmed by fluorescence *in situ* hybridization (FISH) analysis (Figure 1G). We also identified heterogeneous HSPA2 expression during the late 2-cell stage (Figure 1H-J). For the early 2-cell stage, however, HSPA2 was symmetrically distributed between the two blastomeres (Figure S1C-E). These results indicating that the heterogeneity occurred as early as the late 2-cell stage. Thus, we hypothesized that the heterogeneous distribution of HSPA2 in late 2-cell plays a direct or indirect role in the lineage commitment of early embryonic stages.

3.2 The reduced expression of *Hspa2* leads to the reduced expression of ICM-marker

To test our hypothesis, we knocked down the *Hspa2* mRNA level by injecting

Hspa2-specific small interfering RNA (siRNA) into the cytoplasm of zygote, and NC-

FAM (negative control with FAM label) siRNA-injected embryos served as the

control (Figure 2A). The differentially expressed genes were detected by RNA-seq for

Hspa2 siRNA (Knockdown; KD) and control (NC) groups at the 8-cell stage (Figure

2B). In total, 96 up-regulated genes and 410 down-regulated genes were identified in

the *Hspa2*-KD group when compared to the NC group (Figure 2C, D). To investigate

the potential biological functions of the DEGs, we conducted GO and KEGG

enrichment analysis. The most significant GO enriched terms were closely related to

cell fate commitment, stem cell population maintenance, stem cell differentiation and

blastocyst development (Figure 2E). According to KEGG analysis, the down-

regulated genes were primarily associated with Hippo and Wnt signaling pathways

(Figure 2F), which are known to play an important role in cell proliferation,

differentiation, and fate decisions in embryonic development[23-26]. Furthermore, the

blastocyst formation rate in the *Hspa2-KD* group (52.3±2.5%) was significantly lower

than that in the NC group (81.6±2.0%) (Figure 2G, H), thus suggesting that HSPA2

plays an essential role in preimplantation embryonic development. In addition, we

found that the down-regulation of *Hspa2* resulted in a remarkable down-regulation of

specific ICM-marker genes (*Oct4*, *Sox2* and *Nanog*) at the 4-cell and blastocyst stages.

Meanwhile, the mRNA levels of a TE-marker gene, Cdx2, did not significantly

change between the groups (Figure 2I, J).   We then used automated Western

immunoblotting to detect the protein levels of ICM and TE markers in Hspa2-KD

embryos, and found that the expression levels of ICM-marker proteins (OCT4 and SOX2) were inhibited when HSPA2 was down-regulated at the 4-cell and blastocyst stages, while the TE-marker protein CDX2 was not significantly different between the groups (Figure 2K, L). Similarly, for mouse embryonic stem cells (mESCs), the knockdown of HSPA2 resulted in a significant reduction for the ICM-marker proteins, and no change for CDX2 (Figure 2M). These results suggest that HSPA2 plays a role in the regulation of ICM-specific gene expression.

3.3 The knockdown of *Hspa2* expression prevents blastomeres from an ICM fate

To investigate the functional role of HSPA2 in the first lineage segregation during early embryonic development, we examined the proportion of ICM and TE after Hspa2-KD. It was showed that the number of ICM and total cells in the blastocyst were significantly decreased, but the number of TE cells did not change significantly (Figure 3A, B). We then randomly co-injected green fluorescent protein (*Gfp*) mRNA with either *Hspa2*-siRNA or NC-FAM into one blastomere of the 2-cell embryos to trace the fate of its descendent cells into the blastocyst (Figure 3C). Knock-down *Hspa2* in one blastomere at the 2-cell stage also led to developmental defects (Figure S2A, B). Compared with the GFP+NC group (48.2%±7.4%), the GFP+*Hspa2*-KD group (33.8 %±4.3%) exhibited a significantly reduced ratio of progeny cells with an ICM fate. Moreover, the number of ICM-marker OCT4 positive cells was found to be significantly reduced in the GFP+*Hspa2*-KD group (27.1±3.8%) than in the GFP+NC group (37.0%±2.5%) (Figure 3D, E). This is consistent with our GO enriched terms of RNA-Seq in that the negative regulation of cell cycle, G1/S

transition of mitotic cell cycle, mitotic cell cycle phase transition and regulation of

mitotic cell cycle phase transition was found in *Hspa2*-KD group (Figure 2E).

Collectively, these results suggested that HSPA2 has a role in ICM lineage

establishment and its deficiency leads to partial defects in the ICM cells.

3.4 The overexpression of *Hspa2* does not induce blastomere cells to bias an ICM fate

To determine whether the redundant level of HSPA2 protein also affected first

lineage segregation, we overexpressed *Hspa2* by microinjecting *Hspa2* mRNA

(approximately 400ng/µL) into the zygote. Western blotting assays showed that the

HSPA2 was overexpressed in the blastocyst stage (Figure S3). We observed that the

blastocyst formation rate was not affected significantly (Figure 4A, B), thus showing

that the overexpression of *Hspa2* did not exert adverse effects on embryonic

development. In addition, we verified that the mRNA levels of the ICM-marker genes

(*Oct4, Sox2* and *Nanog*) and the TE-marker gene *Cdx2* at the 4-cell and blastocyst

stage did not differ significantly (Figure 4C, D). We also examined the number of

ICM cells and TE cells by immunostaining OCT4 and CDX2 in blastocysts from the

overexpression and control groups and found they were not altered significantly

(Figure 4E, F). To trace the fate of the descendent, we co-injected GFP mRNA along

with either *Hspa2* or NC mRNA randomly into one blastomere of 2-cell embryos and

counted the proportion of ICM and TE cells of blastocysts. We found that the

percentage of GFP cells that were present in the ICM at the blastocyst stage was

similar between the two groups (48.2±7.4 vs. 47.3±4.4), and the percentage of ICM

cells also did not differ significantly (37.0±2.5 vs. 36.2±1.9) (Figure 4G, H). These

observations indicated that the redundant level of HSPA2 were not sufficient to

induce blastomere cells to bias an ICM fate.

3.5 HSPA2 interacts with CARM1 and alters H3R26me2 levels

To further investigate the mechanism underlying the function of HSPA2 in cell

fate decision during early embryonic development, we next analyzed differentially

expressed proteins (DEPs) at the blastocyst stage in the *Hspa2*-KD and NC groups by

applying untargeted proteomics. Volcano map screening identified a total of 43

annotated DEPs, of which 15 proteins were significantly up-regulated and 28 proteins

were downregulated (Figure S4A). These DEPs were mainly involved in post-

translational modification by using Clusters of Orthologous Groups (COG)/Karyotic

Orthologous Groups (KOG) functional annotation statistics (Figure S4B). In order to

compare the functional similarities and differences between proteins with different

differential multiples, we divided the DEPs into four classifications: Q1 (<0.5), Q2

(0.5-0.667), Q3 (1.5-2.0), and Q4 (>2.0) according to differential expression multiples

(Figure S4C). Biological processes analysis showed that DEPs were associated with

positive regulation of the Wnt signaling pathway and cellular development (Figure

S4D). In terms of the cellular component, the DEPs were related to the methylome

and methyltransferase complex (Figure S4E). These results indicated that the DEPs

were mainly associated with the post-translational modification of proteins and cell

fate decision.

Coactivator-associated arginine methyltransferase 1 (CARM1), also known as

PRMT4, was the first protein arginine methyltransferase which was proved to be associated with transcriptional activation by methylating and/or regulating histone H3 and non-histone proteins, such as coactivators and transcription factors [27, 28]. Recent studies have demonstrated that CARM1 regulates early mouse development and plays a role in the determination of cell fate [29, 30]. CARM1 levels, along with its specific histone marks (H3R17me2a and H3R26me2a), were found to vary among the four-cell blastomeres and that high levels of CARM1 led to increased levels of H3R26me. This varied distribution led to biased subsequent fate of the blastomeres towards the ICM [6, 31]. Next, we considered whether HSPA2 regulates the decision of the first cell-fate through CARM1. Molecular docking revealed that HSPA2 docked efficiently with CARM1 via several H-bonds and pi-interactions, with strong binding affinity (Figure 5A). The Search Tool for the Retrieval of Interacting Genes (STRING) database was used to predict interactions between HSPA2 and CARM1. The results showed that HSPA2, CARM1 and ICM-specific proteins interacted with each other (Figure S4F). In addition, we found that the *Hspa2*-KD led to reduced CARM1 mRNA as well as protein levels at the 4-cell and blastocyst stages (Figure 5B-E). The knockdown of HSPA2 in mESCs also caused a significant reduction in the levels of CARM1 (Figure S4G).

To further investigate whether HSPA2 could interact with CARM1, we performed co-immunoprecipitation (co-IP) assays in mESCs using HSPA2 and CARM1 as bait. Co-IP analysis showed that HSPA2 and CARM1 proteins interacted effectively (Figure 5F, G). Moreover, we investigated whether HSPA2 could regulate

H3R26me2 modification, a known event that occurs downstream of CARM1 [6].

Immunofluorescence staining confirmed that CARM1-mediated H3R26me2 levels

were significantly reduced in the interference group at the 4-cell stage (Figure 5H).

Collectively, our data demonstrated that HSPA2 involved in the segregation of

embryonic cell lineage by forming a complex with CARM1 and through H3R26me2

modification.

DISCUSSION

During the development of the fertilized eggs to blastocyst stage, the

fundamental questions are to explain how the totipotent fertilized egg develops into

different cell types and when cellular heterogeneity occurs. Previous studies

demonstrated that molecular polarity and differential gene regulation by signaling

pathways drive the first cell-fate decision in mammals [26, 32-34]. The tracing of

embryonic lineages from the 2-cell stage onwards by live embryo labeling

demonstrated that the cellular heterogeneity is initiated early in the embryo, and thus

was proposed as cellular heterogeneity hypothesis [35]. H3R26me and its

methyltransferase CARM1exhibit asymmetric expression patterns in the 4-cell stage

of mouse embryos and plays an important role in regulating the first cell-fate decision.

[36, 37]. PRDM14 is also heterogeneously expressed in 4-cell stage mouse embryos

and interacts with CARM1 to drive progenies towards pluripotent ICM fate by

increasing the expression levels of H3R26me in 4-cell embryos [38]. Despite the

progress in our understanding of the mechanisms governing early cell fate decisions,

many important questions still remain unanswered. We do not yet fully comprehend

when the initial heterogeneities among cells arise and how are they translated into

divergent patterns of gene expression [39, 40].

Although two-cell blastomeres are generally considered to be totipotent, the

previous reports showing that when two-cell blastomeres are separated, in the

majority of cases, only one of the two-cell blastomeres develops into a mouse [41]. In

this study, we found that the expression levels of HSPA2 were symmetrically

distributed between the two blastomeres at the early 2-cell stage, but showed

heterogeneous distribution at the late 2-cell embryos. Interestingly, Hspa2 mRNA also

heterogeneous distribution at the late 2-cell and 4-cell embryos after microinjection of

siHspa2 in zygotes (Figure S1 F-G). This indicates that the heterogeneity of HSPA2 at

the late 2-cell embryo may be initiated concomitant with the zygotic genome

activation (ZGA) [7]. Alternatively, the asymmetry may be triggered by unequally

distributed unknown factors that already existed in the early 2-cell embryos or the

heterogeneous spatial distribution of maternal genes in the zygote.

The reduced expression of HSPA2 led to the reduced expression levels of ICM-

marker genes at both the mRNA and protein levels. Interfering with HSPA2

expression in one blastomere of the 2-cell embryos reduced the contribution of that

blastomere to ICM differentiation. ICM cells were still formed due to the incomplete

knockdown achieved or the possibility that redundant pathways exist. Notably, the

number of ICM and total cells in the blastocyst was significantly reduced when

compared to the control group (Figure 3B, E). We observed that the GO enriched

terms were also closely related to negative regulation of cell cycle, G1/S transition of

mitotic cell cycle, mitotic cell cycle phase transition and regulation of mitotic cell

cycle phase transition (Figure 2E). HSPA2 knockdown via siRNA reduced cell

proliferation and led to G1/S phase cell cycle arrest [42]. Thus, we guessed that

knockdown of HSPA2 results in lower cell number in ICM by interfering with the cell

cycle. Additionally, the lower cell number in ICM may also associated with failed

cytokinesis and the generation of binucleated or polyploid cells. Interestingly, half of

the ICM cells still form with HSPA2 deficiency (Figure 3B), possibly due to the

incomplete knockdown achieved or the possibility that redundant pathways exist.

However, the overexpression of HSPA2 did neither adversely affect embryo

development and nor induce blastomere bias towards an ICM fate. It may be because

that once the intracellular concentration of *Hspa2* reaches a threshold level, it was not

able to control the target gene expression. Furthermore, we also found that HSPA2

interacted with CARM1, and reduced HSPA2 level, led to the reduced expression of

CARM1, and thus reduced the levels of H3R26me2 modification at the 4-cell stage.

Collectively, these results suggest that HSPA2 physically binds to CARM1 to activate

ICM-specific gene expression through H3R26me2 modification (Figure 5I).

In fact, our RNA-seq results revealed that the differentially expressed genes

(DEGs) at blastocyst stage with HSPA2 knocking down were also enriched in Wnt

signaling pathway and Hippo signaling pathway(Figure 2E), which are also reported

to be involved in cell proliferation, cell polarity, and cell fate determination during

embryonic development. [26] [43]. This indicates that several alternative mechanisms

are compatible with the asymmetric hypothesis, including polarized cell division and

differential gene regulation by signaling.

In summary, our findings provide an exciting step forward for the first time that

the role of HSPA2 in governing the first cell-fate decision in the mouse embryo

lineage allocation. It may be beneficial for achieving a better understanding of

embryogenesis patterns and cell differentiation.

Compliance with Ethical Standards

The current study was carried out with the approval of the Ethic Committee of

Reproductive Medicine of Reproductive Hospital of Shandong University (2021-15),

and all experiments performed were abided by relevant regulations and guidelines of

the committees.

Acknowledgements

This work was supported by National Natural Science Foundation of China

(32170817); The National Key R&D Program of China (2021YFC27003); The

Fundamental Research Funds for the Central Universities (2023QNTD004,

2022JC006); Taishan Scholars Program of Shandong Province (tsqn201909194,

tsqn202211373); Innovative research team of high-level local universities in Shanghai

(SHSMU-ZLCX20210201); Shandong Provincial Natural Science Foundation

(2022HWYQ-034, ZR2021QC106)

Conflict of interest

The authors declare that they have no conflict of interests.

Author contribution

Jiayin Gao wrote the first draft of the manuscript. Jiawei Wang modified the language.

All authors participated in literature research, conception, design, and discussion of

the manuscript, and approved its final version.

References

[1] J. Mu, Z. Zhou, Q. Sang, L. Wang, The physiological and pathological mechanisms of early embryonic development, Fundam Res 2(6) (2022) 859-872.

[2] K. Cockburn, J. Rossant, Making the blastocyst: lessons from the mouse, J Clin Invest 120(4) (2010) 995-1003.

[3] J. Rossant, P.P. Tam, Blastocyst lineage formation, early embryonic asymmetries and axis patterning in the mouse, Development 136(5) (2009) 701-13.

[4] B. Liu, H. Zhao, K. Wu, J. Grosshans, Temporal Gradients Controlling Embryonic Cell Cycle, Biology (Basel) 10(6) (2021).

[5] M. Goolam, A. Scialdone, S.J.L. Graham, I.C. Macaulay, A. Jedrusik, A. Hupalowska, T. Voet, J.C. Marioni, M. Zernicka-Goetz, Heterogeneity in Oct4 and Sox2 Targets Biases Cell Fate in 4-Cell Mouse Embryos, Cell 165(1) (2016) 61-74.

[6] M.E. Torres-Padilla, D.E. Parfitt, T. Kouzarides, M. Zernicka-Goetz, Histone arginine methylation regulates pluripotency in the early mouse embryo, Nature 445(7124) (2007) 214-8.

[7] J. Shi, Q. Chen, X. Li, X. Zheng, Y. Zhang, J. Qiao, F. Tang, Y. Tao, Q. Zhou, E. Duan, Dynamic transcriptional symmetry-breaking in pre-implantation mammalian embryo development revealed by single-cell RNA-seq, Development 142(20) (2015) 3468-77.

[8] F. Sotomayor-Lugo, N. Iglesias-Barrameda, Y.M. Castillo-Aleman, I. Casado-Hernandez, C.A. Villegas-Valverde, A.A. Bencomo-Hernandez, Y. Ventura-Carmenate, R.A. Rivero-Jimenez, The Dynamics of Histone Modifications during Mammalian Zygotic Genome Activation, Int J Mol Sci 25(3) (2024).

[9] J. Wang, L. Wang, G. Feng, Y. Wang, Y. Li, X. Li, C. Liu, G. Jiao, C. Huang, J. Shi, T. Zhou, Q. Chen, Z. Liu, W. Li, Q. Zhou, Asymmetric Expression of LincGET Biases Cell Fate in Two-Cell Mouse Embryos, Cell 175(7) (2018) 1887-1901 e18.

[10] Y. Wang, X. Zheng, R. Cheng, J. Han, X. Ma, W. Xu, L. Gao, A. Lei, J. Liu, F. Quan, Y. Zhang, X. Liu, Asymmetric expression of maternal mRNA governs first cell-fate decision, FASEB J 35(11) (2021) e22006.

[11] D. Scieglinska, Z. Krawczyk, Expression, function, and regulation of the testis-enriched heat shock HSPA2 gene in rodents and humans, Cell Stress Chaperones 20(2) (2015) 221-35.

[12] M.J. Gomez-Torres, N. Huerta-Retamal, P. Saez-Espinosa, L. Robles-Gomez, M. Aviles, J. Aizpurua, Molecular Chaperone HSPA2 Distribution During Hyaluronic Acid Selection in Human Sperm, Reprod Sci 30(4) (2023) 1176-1185.

[13] H. Zhang, W. Chen, C.J. Duan, C.F. Zhang, Overexpression of HSPA2 is correlated with poor prognosis in esophageal squamous cell carcinoma, World J Surg Oncol 11 (2013) 141.

[14] L.L. Bonnycastle, C.E. Yu, C.R. Hunt, B.J. Trask, K.P. Clancy, J.L. Weber, D. Patterson, G.D. Schellenberg, Cloning, sequencing, and mapping of the human chromosome 14 heat shock protein gene (HSPA2), Genomics 23(1) (1994) 85-93.

[15] D.J. Dix, J.W. Allen, B.W. Collins, C. Mori, N. Nakamura, P. Poorman-Allen, E.H. Goulding, E.M. Eddy, Targeted gene disruption of Hsp70-2 results in failed meiosis, germ cell apoptosis, and male infertility, Proc Natl Acad Sci U S A 93(8) (1996) 3264-8.

[16] A.K. Murashov, D.J. Wolgemuth, Distinct transcripts are recognized by sense and antisense riboprobes for a member of the murine HSP70 gene family, HSP70.2, in various reproductive tissues, Mol Reprod Dev 43(1) (1996) 17-24.

[17] W. Rupik, A. Stawierej, I. Stolarczyk, W. Widlak, Promoter of the heat shock testis-specific
Hsp70.2/Hst70 gene is active in nervous system during embryonic development of mice, Anat
Embryol (Berl) 211(6) (2006) 631-8.
[18] M. Garg, D. Kanojia, S. Saini, S. Suri, A. Gupta, A. Surolia, A. Suri, Germ cell-specific heat
shock protein 70-2 is expressed in cervical carcinoma and is involved in the growth, migration,
and invasion of cervical cells, Cancer 116(16) (2010) 3785-96.
[19] M. Garg, D. Kanojia, A. Seth, R. Kumar, A. Gupta, A. Surolia, A. Suri, Heat-shock protein 70-2
(HSP70-2) expression in bladder urothelial carcinoma is associated with tumour progression and
promotes migration and invasion, Eur J Cancer 46(1) (2010) 207-15.
[20] J. Song, J. Zhang, X. Yuan, B. Liu, W. Tao, C. Zhang, K. Wu, Functional substitution of zona
pellucida with modified sodium hyaluronate gel in human embryos, J Assist Reprod Genet 39(11)
(2022) 2669-2676.
[21] G.G. Schiattarella, F. Altamirano, D. Tong, K.M. French, E. Villalobos, S.Y. Kim, X. Luo, N. Jiang,
H.I. May, Z.V. Wang, T.M. Hill, P.P.A. Mammen, J. Huang, D.I. Lee, V.S. Hahn, K. Sharma, D.A. Kass,
S. Lavandero, T.G. Gillette, J.A. Hill, Nitrosative stress drives heart failure with preserved ejection
fraction, Nature 568(7752) (2019) 351-356.
[22] S. Picelli, O.R. Faridani, A.K. Bjorklund, G. Winberg, S. Sagasser, R. Sandberg, Full-length
RNA-seq from single cells using Smart-seq2, Nat Protoc 9(1) (2014) 171-81.
[23] B. Swierczek-Lasek, D. Dudka, D. Bauer, T. Czajkowski, K. Ilach, W. Streminska, A. Kominek, K.
Piwocka, M.A. Ciemerych, K. Archacka, Comparison of Differentiation Pattern and WNT/SHH
Signaling in Pluripotent Stem Cells Cultured under Different Conditions, Cells 10(10) (2021).
[24] D. ten Berge, D. Kurek, T. Blauwkamp, W. Koole, A. Maas, E. Eroglu, R.K. Siu, R. Nusse,
Embryonic stem cells require Wnt proteins to prevent differentiation to epiblast stem cells, Nat
Cell Biol 13(9) (2011) 1070-5.
[25] A. de Jaime-Soguero, W.A. Abreu de Oliveira, F. Lluis, The Pleiotropic Effects of the
Canonical Wnt Pathway in Early Development and Pluripotency, Genes (Basel) 9(2) (2018).
[26] N. Nishioka, K. Inoue, K. Adachi, H. Kiyonari, M. Ota, A. Ralston, N. Yabuta, S. Hirahara, R.O.
Stephenson, N. Ogonuki, R. Makita, H. Kurihara, E.M. Morin-Kensicki, H. Nojima, J. Rossant, K.
Nakao, H. Niwa, H. Sasaki, The Hippo signaling pathway components Lats and Yap pattern
Tead4 activity to distinguish mouse trophectoderm from inner cell mass, Dev Cell 16(3) (2009)
398-410.
[27] M.R. Stallcup, J.H. Kim, C. Teyssier, Y.H. Lee, H. Ma, D. Chen, The roles of protein-protein
interactions and protein methylation in transcriptional activation by nuclear receptors and their
coactivators, J Steroid Biochem Mol Biol 85(2-5) (2003) 139-45.
[28] N. Troffer-Charlier, V. Cura, P. Hassenboehler, D. Moras, J. Cavarelli, Functional insights from
structures of coactivator-associated arginine methyltransferase 1 domains, EMBO J 26(20) (2007)
4391-401.
[29] A. Hupalowska, A. Jedrusik, M. Zhu, M.T. Bedford, D.M. Glover, M. Zernicka-Goetz, CARM1
and Paraspeckles Regulate Pre-implantation Mouse Embryo Development, Cell 175(7) (2018)
1902-1916 e13.
[30] M.R. Pawlak, C.A. Scherer, J. Chen, M.J. Roshon, H.E. Ruley, Arginine N-methyltransferase 1
is required for early postimplantation mouse development, but cells deficient in the enzyme are
viable, Mol Cell Biol 20(13) (2000) 4859-69.
[31] D.E. Parfitt, M. Zernicka-Goetz, Epigenetic modification affecting expression of cell polarity

and cell fate genes to regulate lineage specification in the early mouse embryo, Mol Biol Cell
21(15) (2010) 2649-60.

[32] M.H. Johnson, C.A. Ziomek, Induction of polarity in mouse 8-cell blastomeres: specificity,
geometry, and stability, J Cell Biol 91(1) (1981) 303-8.

[33] E. Korotkevich, R. Niwayama, A. Courtois, S. Friese, N. Berger, F. Buchholz, T. Hiiragi, The
Apical Domain Is Required and Sufficient for the First Lineage Segregation in the Mouse Embryo,
Dev Cell 40(3) (2017) 235-247 e7.

[34] Y. Hirate, H. Sasaki, The role of angiomotin phosphorylation in the Hippo pathway during
preimplantation mouse development, Tissue Barriers 2(1) (2014) e28127.

[35] K. Piotrowska, M. Zernicka-Goetz, Role for sperm in spatial patterning of the early mouse
embryo, Nature 409(6819) (2001) 517-21.

[36] R. Xu, C. Li, X. Liu, S. Gao, Insights into epigenetic patterns in mammalian early embryos,
Protein Cell 12(1) (2021) 7-28.

[37] W. Cui, A. Cheong, Y. Wang, Y. Tsuchida, Y. Liu, K.D. Tremblay, J. Mager, MCRS1 is essential
for epiblast development during early mouse embryogenesis, Reproduction 159(1) (2020) 1-13.

[38] A. Burton, J. Muller, S. Tu, P. Padilla-Longoria, E. Guccione, M.E. Torres-Padilla, Single-cell
profiling of epigenetic modifiers identifies PRDM14 as an inducer of cell fate in the mammalian
embryo, Cell Rep 5(3) (2013) 687-701.

[39] I. Chambers, J. Silva, D. Colby, J. Nichols, B. Nijmeijer, M. Robertson, J. Vrana, K. Jones, L.
Grotewold, A. Smith, Nanog safeguards pluripotency and mediates germline development,
Nature 450(7173) (2007) 1230-4.

[40] A.M. Singh, T. Hamazaki, K.E. Hankowski, N. Terada, A heterogeneous expression pattern for
Nanog in embryonic stem cells, Stem Cells 25(10) (2007) 2534-42.

[41] E. Casser, S. Israel, A. Witten, K. Schulte, S. Schlatt, V. Nordhoff, M. Boiani, Totipotency
segregates between the sister blastomeres of two-cell stage mouse embryos, Sci Rep 7(1) (2017)
8299.

[42] L. Cao, X. Yuan, F. Bao, W. Lv, Z. He, J. Tang, J. Han, J. Hu, Downregulation of HSPA2 inhibits
proliferation via ERK1/2 pathway and endoplasmic reticular stress in lung adenocarcinoma, Ann
Transl Med 7(20) (2019) 540.

[43] S.A. Morris, Y. Guo, M. Zernicka-Goetz, Developmental plasticity is bound by pluripotency
and the Fgf and Wnt signaling pathways, Cell Rep 2(4) (2012) 756-65.

Figure caption

Figure 1. *Hspa2* is asymmetrically expressed between mouse 2- and 4-cell blastomeres. (A)

Schematic overview. The 4-cell embryo divided into single blastomeres and cultured to the morula

stage. Created with biorender.com. **(B)** *Hspa2* expression is heterogeneous in morula developed

from 4-cell blastomeres. Three experimental replicates were performed. **(C)** *Nanog* and *Hspa2*

show stronger positive correlation (R=0.624) in gene expression than random gene pairs (n=100, n

means the number of random gene pairs). (D) qPCR analysis of the relative expression of Hspa2

mRNA form the meiosis II to blastocyst stages. The error bars represent SEM. About 20 embryos

of each stage were used and three experimental replicates were performed. **(E and H)** Single-cell

qRT-PCR results showing *Hspa2* distribution between 4- (E) (n=15), late 2- (H) (n=17) cell

blastomere. *Hprt* as the housekeeping gene. CT values are used for level analysis. Two-tailed

Student's t tests were used for the statistical analysis. Blastomeres are named B1 to B4 (E) or B1

and B2 (H) according to increasing *Hspa2* concentration. $*p < 0.05$. **(F and I)** Automated western

immunoblotting results showing HSPA2 distribution between 4- (F) and late 2- (I) cell

blastomeres. Blastomeres are named B1 to B4 (E) or B1 and B2 (H) according to decreasing

Hspa2 concentration. The band intensity was assessed. There experimental replicates were

performed. $*p < 0.05$. **(G and J)** FISH of *Hspa2* mRNA in early 4- (G) (n=17), late 2- (J) (n=22)

cell embryos. Blastomeres with low expression as white triangles. RNA-FISH fluorescence

quantified and normalized to the nucleus with the strongest staining per individual two- or four-

cell embryo. Blastomeres are named B1 to B4 (G) or B1 and B2 (J) according to increasing *Hspa2*

concentration. Three experimental replicates were performed. Bar, 25 μm. $*p < 0.05$.

Figure 2. Knockdown of *Hspa2* lead to decreased expression of ICM-mark at both the

mRNA and protein level. (A) Schematic diagram of microinjection of zygotic embryos. Created

with biorender.com. **(B)** Bar charts showing the expression of *Hspa2* in the negative control (NC)

group (injected with negative control siRNA) and siRNA targeting group based on RNA-seq. The

error bars denote the standard deviations of three experimental replicates of RNA-seq. $*p < 0.05$.

(C) Heat map of the hierarchical clustering analysis of the DEGs between NC and KD groups. **(D)**

Volcano plots of significantly differentially expressed genes (DEGs). Red, upregulated; Blue,

down-regulated**. (E)** GO enrichment analysis from down-regulated DEGs between NC and KD

groups. **(F)** Significantly down-regulated DEGs were analyzed by KEGG enrichment. **(G)**

Representative images of preimplantation embryos in the NC and KD groups from 2-cell to

blastocyst. Red arrowhead denotes degenerating blastocysts. Bar, 100 μm. **(H)** Bar plots showing

the developmental rates of NC group and KD group at the blastocyst stage.. Three experimental

replicates were performed. **(I and J)** qRT-PCR of 4-cell embryo and blastocyst showed that ICM-

marker gene (*Oct4*, *Sox2* and *Nanog*) mRNA was significantly lower in the interference compared

with the NC group. There were no significant differences in Cdx2 between the two groups.

*$p < 0.05$, **$p < 0.01$. **(K and L)** Automated western blotting analysis of ICM-marker protein

(OCT4, SOX2 and NANOG) expression in the interference and NC groups. Three experimental

replicates were performed. *$p < 0.05$. **(M)** Western blot analysis of ICM-marker protein (OCT4,

SOX2 and NANOG) and TE-marker protein (CDX2) expression in mESCs. The band intensity

was assessed with ImageJ. Three experimental replicates were performed. *$p < 0.05$.

Figure 3. Knockdown of *Hspa2* prevents blastomeres from undergoing an ICM fate. (A)

Immunofluorescence of OCT4 (green) and CDX2 (red) in NC (n=40) and si-*Hspa2* (n=36) groups.

OCT4 was used as an ICM marker. Nuclei were visualized with DAPI staining (blue). Bar, 50 μm.

(B) The number of ICM and TE. OCT4 was used as an ICM marker and CDX2 was used as a TE

marker. *$p < 0.05$. **(C)** Schematic diagram of microinjection of early 2-cell embryo single

blastomeres. Created with biorender.com. **(D)**OCT4 and GFP fluorescent staining of blastocysts.

OCT4 was used as an ICM marker. The results show that less OCT4-positive cells are GFP-

positive in embryos injected with GFP and *Hspa2 siRNA* (GFP+*si-Hspa2*). Three experimental

replicates were performed. Bar, 50 μm. **(E)** Analysis of the distribution of progeny of injected

blastomere at the blastocyst stage based on fluorescence staining. Total cells (n) are the total

number of cells in the blastocyst; inner cells (%) is the percentage of inner cells relative to the

total number of cells in the blastocyst; GFP cells that are inner (%) is the percentage of GFP-

positive inner cells relative to the total number of GFP-positive cells in the blastocyst. Data

derived from three independent experiments and presented as mean ± standard deviation. Three

experimental replicates were performed. *$p <0.05$.

Figure 4. Overexpression of *Hspa2* not induces blastocyst bias towards ICM fate. (A)

Representative images of preimplantation embryos in the NC and *Hspa2* mRNA groups from 2-

cell to blastocyst. Bar, 100 μm. **(B)** The blastocyst rate between NC (n=151) and overexpression

(n=150) groups. Three experimental replicates were performed. **(C and D)** qRT-PCR of 4-cell

embryo and blastocyst showed that ICM-marker gene (*Oct4, Sox2* and *Nanog*) and TE-marker

gene (*Cdx2*) were no significant differences in the overexpression compared with the control

group.   Three   experimental   replicates   were   performed.   *$p <0.05$,   ****$p <0.001$.   **(E)**

Immunofluorescence of OCT4 (green) and CDX2 (purple) in NC (n=38) and overexpression

(n=40) groups. OCT4 was used as an ICM marker. Nuclei were visualized with DAPI staining

(blue). Bar, 50 μm. **(F)** The number of ICM and TE. OCT4 was used as an ICM marker and

CDX2 was used as a TE marker. **(G)** OCT4 and GFP fluorescent staining of blastocysts. OCT4

was used as an ICM marker. The results show that the percentage of GFP cells that were present in

the ICM was similar between the two groups. Three experimental replicates were performed.

Three experimental replicates were performed. Bar, 50 μm. **(H)** Analysis of the distribution of

progeny of injected blastomere at the blastocyst stage based on fluorescence staining. Total cells

(n) are the total number of cells in the blastocyst; inner cells (%) is the percentage of inner cells

relative to the total number of cells in the blastocyst; GFP cells that are inner (%) is the percentage

of GFP-positive inner cells relative to the total number of GFP-positive cells in the blastocyst.

Data derived from three independent experiments and presented as mean ± standard deviation.

Three experimental replicates were performed.

Figure 5. HSPA2 and CARM1 form a complex. (A) Molecular docking analysis for HSPA2

binding to CARM1 proteins. **(B and C)** qRT-PCR of 4-cell embryo and blastocyst showed that

Carm1 was significantly lower in the interference compared with the control group. Three

experimental replicates were performed. *$p < 0.05$. **(D and E)** Automated western blotting

analysis of CARM1 expression in the interference and control groups. The band intensity was

assessed. Three experimental replicates were performed. *$p < 0.05$ **(F and G)** CO-IP analysis of

HSPA2 and CARM1 in mESCs. **(H)** H3R26me2 staining of 4-cell embryo shows that HSPA2

knockdown led to a dramatic decrease in H3R26me2 modification. Intensity relative to DAPI

signal was used. Three experimental replicates were performed. Bar, 50 μm.*$p < 0.05$. (I) The

model of HSPA2 physically binds to CARM1 to activate ICM-specific gene expression.

Supplemental Figure caption

Figure S1. *Hspa2* is asymmetrically expressed between mouse 2- and 4-cell blastomeres,

Related to Figure 1. (A) Schematic overview. Late 2-cell divided into single blastomeres and

cultured to the morula stage. Created with biorender.com. **(B)** *Hspa2* expression is heterogeneous

and the level of *Nanog* is positively correlated with the expression of *Hspa2* in morula developed

from late 2-cell blastomeres. Three experimental replicates were performed. **(C)** Single-cell

qRT-PCR results showing *Hspa2* distribution between early 2-cell (n=17) blastomere. *Hprt* as the

housekeeping gene. CT values are used for level analysis. Two-tailed Student's t tests were used

for the statistical analysis. Three experimental replicates were performed. **(D)** Automated western

immunoblotting results showing HSPA2 distribution between early 2-cell blastomeres.

Blastomeres are abbreviated named B. The protein level is normalized B1. Three experimental

replicates were performed. **(E)** FISH of *Hspa2* mRNA in early 2-cell embryos (n=40). RNA-

FISH fluorescence quantified and normalized to the nucleus with the strongest staining per

individual early 2-cell embryo. Three experimental replicates were performed. Bar, 25 μm. **(F and**

G) Single-cell qRT-PCR results showing *Hspa2* level in blastomeres at the late 2- (F) and 4-cell

(G) stage after microinjection of siHspa2 in zygotes. *Hprt* as the housekeeping gene. CT values

are used for level analysis. Two-tailed Student's t tests were used for the statistical analysis.

Blastomeres are named B1 to B4 (F) or B1 and B2 (G) according to increasing *Hspa2*

concentration. *p <0.05.

Figure S2. Knockdown of *Hspa2* prevents blastomeres from undergoing an ICM fate.

Related to Figure 3. (A) Representative images of preimplantation embryos in the NC and KD

groups from 2-cell to blastocyst. **(B)** Bar plots showing the developmental rates of NC group and

KD group at the blastocyst stage.

Figure S3. Overexpression of *Hspa2* not induces blastocyst bias towards ICM fate. Related

to Figure 4. (A) Western blot results confirming the overexpression of HSPA2 protein in the

blastocyst stages. Three experimental replicates were performed. OE, overexpression.

Figure S4. HSPA2 interacts with CARM1 and alters H3R26me2 levels. Related to Figure

5. (A) Volcano plot of the DEPs in *Hspa2* siRNA and control groups at blastocyst stage. **(B)**

COG/KOG enrichment analysis for DEPs. **(C)** Quantitative classification and numbers of DEPs

in each quantified part. The DEPs were divided into four quantified parts according to their

differential expression multiples: Q1 (<0.5), Q2 (0.5-0.667), Q3 (1.5-2.0), and Q4 (>2.0). **(D and**

E) Clustering analysis including biological process analysis (D) and cellular component (E) of

quantitative DEPs in each Q group. **(F)** The Protein-protein interactions (PPIs) network was

mapped   by   STRING   database   ([https://cn.string-](https://cn.string-db.org/cgi/network?taskId=bk9UNYyN0p0f&sessionId=bS81YN78aR)

[db.org/cgi/network?taskId=bk9UNYyN0p0f&sessionId=bS81YN78aR](https://cn.string-db.org/cgi/network?taskId=bk9UNYyN0p0f&sessionId=bS81YN78aR)). **(G)** Western blot analysis

of CARM1 expression in the knockdown and control groups. The band intensity was assessed

with ImageJ. Three experimental replicates were performed. ***$p < 0.001$.

Group	Blastocysts (n)	Total cells (n)	Inner cells (%)	GFP cells that are inner (%)
GFP+NC	49	48.6±7.9	37.0±2.5	48.2±7.4
GFP+*si-Hspa2*	37	46.6±5.1	27.1±3.8*	33.8±4.3*

A