Beyond Auditory Relay: Dissecting the Inferior Colliculus's Role in Sensory Prediction, Cognitive Decision-Making, and Reward Prediction

Xinyu Du[1, 2#], Haoxuan Xu[3, 4#], Peirun Song[1, 2], Yuying Zhai[1, 2], Hangting Ye[1, 2], Xuehui Bao[3, 4], Qianyue Huang[3, 4], Hisashi Tanigawa[3, 4], Zhiyi Tu[5], Pei Chen[5], Xuan Zhao[5], Josef P. Rauschecker [6] and Xiongjie Yu[1, 2, 3, 4, 5]

[1] Department of Anesthesia, Women's Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang Province, China;

[2] Zhejiang Provincial Key Laboratory of Precision Diagnosis and Therapy for Major Gynecological Diseases, Women's Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China.

[3] College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou, Zhejiang Province, China;

[4] Key Laboratory for Biomedical Engineering of Ministry of Education, Zhejiang University, China;

[5] Department of Anesthesiology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, China;

[6] Department of Neuroscience, Georgetown University, Washington, DC, 20007, U.S.A.

[#] Xinyu Du and Haoxuan Xu contributed equally to this work.

Corresponding Author: Xiongjie Yu (yuxiongj@zju.edu.cn)

Running Title: Multifaceted role in IC

Key Words: inferior colliculus; sound duration; decision making; prediction error; reward behavior.

Author contributions: Conceptualization, X.Y.; Methodology, X.Y., X.D., and H.X.; Investigation and Data Curation, X.D., H.X., P.S., Y.Z., H.Y., X.B., Q.H., H.T., Z.T., P.C. and X.Z.; Writing – Original Draft, X.Y. and X.D. and J.P.R.; Writing – Review & Editing, X.D., H.X. and J.P.R.; Funding Acquisition, X.Y. and Y.Z.; Resources, X.Y.; Software, X.D., H.X., P.S., Y.Z., H.Y., X.B., Q.H., H.T., Z.T., P.C. and X.Z.; Validation and Visualization, X.D. and H.X.; Supervision, X.Y.

Acknowledgments: We are grateful to Prof. Liping Wang for their invaluable comments on the early version of the manuscript, as well as to Xiaokai Kou and Fujin Gao for their assistance with the experiments. This work was supported by STI2030-Major Projects (2022ZD0204800 and 2022ZD0204600) to XY, the National Natural Science Foundation of China (32171044 to XY and 32100827 to YZ).

Declaration of interests: The authors declare no competing interests.

Abstract

The Inferior Colliculus (IC) has traditionally been regarded as an important relay in the auditory pathway, primarily involved in relaying auditory information from the brainstem to the thalamus. However, this study uncovers the multifaceted role of the IC in bridging auditory processing, sensory prediction, and reward prediction. Through extracellular recordings in monkeys engaged in a sound duration-based novelty detection task, we observed a "climbing effect" in neuronal firing rates, indicative of an enhanced response over sound sequences linked to sensory prediction rather than reward anticipation. Further exploration revealed a direct correlation between IC neuronal activity and behavioral choices, suggesting its involvement in decision-making processes. Additionally, our findings demonstrate reward prediction errors within the IC, highlighting its complex integration in auditory and reward processing. This research challenges conventional views of the IC, showcasing its integral role in cognitive and sensory processing and emphasizing its importance in integrated brain functions.

Introduction

The brain's architecture is a complex hierarchy, where functions become increasingly sophisticated from lower to higher levels. The cerebral cortex, for instance, is pivotal for higher-order cognitive functions such as decision-making, motivation, attention, learning, memory, problem-solving, and conceptual thinking (Kremkow and Alonso, 2018; Miterko et al., 2018). Conversely, subcortical sensory systems are traditionally viewed as mere conduits for sensory information (Sherman, 2016), despite suggestions of thalamic structures playing roles in advanced perceptual tasks (Jones, 2012; Sherman, 2005). Yet, the involvement of sensory neurons below the thalamo-cortical level in cognitive behaviors remains an underexplored domain. Within the auditory hierarchy, spanning seven stages from the cochlea to the auditory cortex (AC) (Pickles, 2015), the inferior colliculus (IC) emerges as a crucial intermediary, instrumental in processing spatial representations (Cohen and Knudsen, 1999; Ono and Ito, 2015) and novelty detection (Malmierca et al., 2009; Perez-Gonzalez et al., 2005). Utilizing the oddball paradigm, researchers have identified stimulus-specific adaptation (SSA) within the inferior colliculus (IC) (Ayala et al., 2016; Duque and Malmierca, 2015; Malmierca et al., 2009; Valdés-Baizabal et al., 2021; Valdés-Baizabal et al., 2017), where neurons decrease their response to frequently occurring sounds while maintaining robust reactions to deviant ones. This phenomenon, indicative of predictive coding, underscores the IC's relevance with sensory prediction. Exploration along the auditory pathway suggests SSA's origination from the IC (Carbajal and Malmierca, 2018; Khouri and Nelken, 2015;

Song et al., 2023). Intriguingly, the dynamic interplay between sensory prediction and

sensory encoding, particularly within lower sensory systems charged with the faithful

representation of external stimuli, remains an underexplored yet captivating area of

inquiry.

Traditionally viewed as a key player in auditory processing, the inferior colliculus

(IC) is situated in the auditory midbrain and serves as a vital conduit, channeling

inputs from the brainstem's auditory nuclei towards the thalamocortical auditory

pathway. Intriguingly, the IC is endowed with dopamine receptors and exhibits nerve

terminals positive for tyrosine hydroxylase, signifying a channel for dopaminergic

inputs originating from the subparafascicular thalamic nucleus (Harris et al., 2021;

Nevue et al., 2015; Nevue et al., 2016). This positions the IC as a crucial intersection

where auditory processing and reward mechanisms converge. Recent research

underscores the significant influence of dopamine on modulating auditory responses

within the IC (Gittelman et al., 2013; Hoyt et al., 2019). This modulation suggests a

complex interplay wherein dopamine has the potential to either suppress or enhance

neural activity in the IC, depending on various factors (Gittelman et al., 2013; Hoyt et

al., 2019). This amalgamation of sensory and reward coding within the IC paves new

paths for research, urging a deeper inquiry into how this midbrain entity integrates

auditory and reward information to mold behavioral processes, thereby challenging

the traditional auditory-centric view of the IC and opening avenues for understanding

its multifaceted role in cognitive and sensory modulation.

In this research, we embarked on a novelty detection task centered around sound

duration with trained monkeys, performing extracellular recordings in the IC. Our

observations unveiled a 'climbing effect'—a progressive increase in firing rate after

sound onset, not attributable to reward but seemingly linked to sensory experience

such as sensory prediction. Moreover, we identified signals of reward prediction error

and decision-making. These findings propose that the IC's role in auditory processing

extends into the realm of complex perceptual and cognitive tasks, challenging

previous assumptions about its functionality.

Results

Response dynamics of IC neurons in duration novelty detection

Neurons within IC (Supplementary Fig.1) are distinguished by their ability to

sustain firing, illustrating a consistent encoding of sound duration through tonic firing

throughout the entire auditory length in one example neuron (Fig. 1A). An analysis of

104 neurons (53 from Monkey B and 51 from Monkey J) demonstrated that these

neurons maintained sustained firing responses to sounds of varying durations—100,

500, and 1000 ms (Fig. 1B). This sustained firing behavior suggests the IC's role in

processing temporal aspects of auditory perception, highlighting its potential

involvement in functions related to duration perception.

Despite traditionally being viewed as a lower-level nucleus within the auditory

pathway in comparison to the auditory cortex, and rarely explored from a perceptual

standpoint, this study breaks new ground by examining the involvement of IC neurons

in duration perception. Monkeys were trained to perform a novelty detection task focusing on duration. Figures 1C and 1D outline the behavioral task, presenting repetitive sounds in blocks with oddball stimuli. Each block consisted of 3 to 6 repeated white-noise bursts of 300-ms duration at 600-ms interstimulus intervals, concluding with either a standard sound (control condition) or a deviant sound (Fig. 1D). Monkeys were required to press a button within 600 ms after the offset of the deviant stimulus to receive a water reward. Correct rejections occurred when the monkey refrained from pressing the button within 600 ms following a standard sound in control trials, where the deviant duration equaled the standard duration. The task difficulty was modulated by setting deviant durations at five levels: 300 (control), 357, 425, 506, and 600 ms (Fig. 1D), with the pressing ratio approaching zero in the control condition and increasing with the deviant duration, reaching 1 at the 600 ms deviant duration in one example session (Fig. 1E).

The initial analysis focused on the deviant response during correct trials, revealing that an example IC neuron exhibited continuous firing throughout the sound presentation (Fig. 2A). Following the peak response in the initial 0-60 ms window, the neuron's firing rate began to increase approximately 100 ms after the sound's onset, continuing until the sound's conclusion. This phenomenon, illustrated by aligning the responses to the sound's offset (Fig. 2B), indicated a proportional increase in firing rate with longer sound durations, suggesting a duration-dependent "climbing effect." To quantitatively assess this dynamic, the neuronal firing rate was examined across different sound durations within two distinct temporal windows: the "onset window"

(0-60 ms post-sound onset) and the "late window" (-100-0 ms relative to sound offset).

Notably, the firing rate of the example neuron in the late window escalated with increasing duration (p = 6.51e-23, ANOVA with post hoc test; red in Fig. 2C), while remaining constant in the onset window (p = 0.69, ANOVA with post-hoc test; black in Fig. 2C), contrasting sharply with the non-behavior condition (Supplementary Fig. 1) which showed no climbing effect.

To assess if the "climbing effect" was contingent on behavioral context, we compared responses across behavior and non-behavior conditions. Sound durations were analyzed at three levels (100, 500, and 1000 ms for the non-behavior condition as shown in Fig. 1A, B) and five levels (300, 357, 425, 506, and 600 ms for the behavior condition, illustrated in Fig. 1C, D). In the same neuronal population (n=99), the firing rate increased after the peak onset response for the behavior situation (red line of Supplementary Fig. 2A), while the firing rate stayed constant for the non-behavior condition from 100 ms to the end of the sound (blue line of Supplementary Fig. 2A). The "Response Dynamic Index" (RDI) was introduced to quantify the climbing effect, calculated as the normalized difference between responses in the late window (-100-0 ms relative to the sound's offset) and the after-peak window (100-200 ms relative to the sound's onset). This comparative analysis indicated a significantly higher RDI in the behavior condition (p = 4.40e-19, paired t-test, Supplementary Fig. 2B), indicating the climbing effect's reliance on behavioral context. Additionally, RDIs across various deviant sounds showed strong correlations (p < 0.001, Pearson Correlation Analysis, Supplementary Fig. 3) and

similar values across differing deviants (Supplementary Fig. 3). Detailed insights into the tonotopic distribution of the climbing effect within the IC are provided in Supplementary Fig. 4.

Exploration into whether the climbing effect manifested in responses to preceding standard stimuli showed that the same example neuron exhibited the climbing effect post-peak response to standard stimuli, which intensified as successive standard sounds progressed from the second to the seventh stimulation (Fig. 2D). The firing rate markedly decreased from the first to the second sound in both the late window (p = 3.33e-11, paired t-test, red in Fig. 2E) and onset window (p = 2.02e-40, paired t-test, black in Fig. 2E). The rate then increased in the late window (p = 3.05e-5, ANOVA with post hoc test: second versus fourth, p = 0.0398; fourth versus seventh, p = 1.00; red in Fig. 2E) but remained consistent in the onset window (p = 0.963, ANOVA with post hoc test, black in Fig. 2E) from the second to the seventh sound. Interestingly, no climbing effect was noted in the late window for the first stimulus, and the effect amplified with each subsequent standard presentation.

To monitor the dynamic of the climbing effect across repetitive sound presentations, we analyzed responses to the 300 ms sound across the entire oddball block in population. We normalized the firing rate relative to the response to the first stimulus in the block, considering both the onset and late windows (Fig. 3A), to account for variability across the neuronal population. In the onset window, responses remained relatively stable from the second to the seventh sound (black line, Fig. 3A). Conversely, in the late window, responses ascended from the second to the fourth

stimulus (ANOVA with post-hoc test: second versus fourth, p = 8.29e-4, red line, Fig.

3A) and maintained a stable level from the fourth to the seventh stimulus (p = 0.653,

ANOVA with post hoc test, red line, Fig. 3A). This pattern was mirrored in the

peri-stimulus time histograms (PSTHs), which consolidated responses from the

second to the seventh stimulus in one scale (left ordinate in Fig.3B) and the first

stimulus in another scale (right ordinate in Fig.3B) for comparative purposes. Onset

responses to the second through seventh stimuli completely overlapped, but responses

in the late window diverged. The response profile remained flat in the late window for

the first stimulus (black in Fig.3B), indicating a minimal climbing effect. Remarkably,

starting from the second stimulus, the climbing effect began to manifest and

progressively intensified, with a notable increase observed from the second to the

fourth stimulus. This effect appeared to stabilize and overlap in responses from the

fourth to the seventh sound (Fig. 3B), illustrating the nuanced modulation of neuronal

firing rates across successive sound presentations within the oddball sequence.

Reward effect on neuronal responses of IC neurons

The accumulation of the climbing effect alongside repetitive sound presentations

suggests a potential linkage to reward prediction, mirroring an increase in the

probability of receiving a reward as the sound sequence extends. A specialized

experimental setup was utilized to determine if the anticipation of a reward drives the

climbing effect. In this setup, a sequence of 1000-ms white noise bursts was played

across 150 trials (Fig. 4A), segmented into an initial 50-trial phase without reward

(early no-reward block), a subsequent 50-trial phase with a water reward dispensed

500 ms after the sound offset (reward block), and a concluding 50-trial phase without reward (late no-reward block). An exemplary IC neuron exhibited sustained firing throughout the 1000-ms sound across all trials (Fig. 4B), with a spike cluster coinciding with the reward delivery, indicating a reward-responsive behavior within the IC neuron (Fig. 4B). The PSTHs for these conditions revealed nearly identical responses during the sound period ($p = 0.098$, ANOVA with post hoc test, Fig. 4C), suggesting the absence of a climbing effect following the peak response.

Out of fifty-nine IC neurons tested using this protocol, twenty-four demonstrated significant reward responses ($p < 0.05$, paired t-test, Fig. 4D), while thirty-five showed no significant reward response (Fig. 4E). Across both neuron types, PSTHs following the onset peak response maintained a consistently flat profile (Fig. 4D, E). However, the RDI during the reward condition, calculated from the normalized difference between responses in the late window and the after-peak window, indicated a behavior-dependent climbing effect, with most values lying above the unitary line ($p = 9.75e-11$, paired t-test, Fig. 4F). Furthermore, an additional set of twenty-two neurons was analyzed using 500-ms sounds within the reward protocol to ensure consistency in sound duration. This analysis showed that the PSTH firing rate increased in the behavioral condition but remained flat in the reward condition following the peak response (100-500 ms) (Supplementary Fig. 5A). The majority of these neurons' RDIs exceeded the unitary line in pairwise comparisons ($p = 3.02e-5$, paired t-test, Supplementary Fig. 5B), reinforcing the notion that the climbing effect is contingent upon the behavioral context rather than the reward.

Reward prediction error in IC neuronal response

Notably, the reward response of the showcased neuron in Fig. 4B was more

pronounced at the start of the reward block. To explore this, we compared the first 15

trials to the last 15 trials within the reward block (Fig. 5A), revealing that the reward

response (0-200 ms relative to the reward onset) was significantly stronger during the

initial trials (p = 1.50e-2, ANOVA, Fig. 5A). Twenty-eight neurons exhibiting

significant reward responses were compiled; auditory responses were consistent

across initial and final trials (p = 0.534, ANOVA, Fig. 5B), yet the reward responses

in the initial trials were markedly higher (p = 7.00e-3, ANOVA, Fig. 5B). A

comparison on an individual neuronal basis showed most points below the unity line

(p = 3.20e-3, paired t-test, Fig. 5C), indicating a variance in reward responsiveness.

For another example neuron without a reward response (Fig. 5D), auditory and

reward responses between the first and last trials were indistinguishable for both

phases (p = 0.378 for auditory response, ANOVA, Fig. 5D; p = 0.637 for reward

response period, ANOVA, Fig. 5D). A collection of thirty-five neurons without

reward responses exhibited similar firing rates for both auditory and reward response

periods across trials (p = 0.506 for auditory response, ANOVA, Fig. 5E; p = 0.585 for

reward response period, ANOVA, Fig. 5E), with the majority of pairwise

comparisons aligning closely with the unity line (p = 0.62, paired t-test, Fig. 5F).

The heightened responses during the initial trials of the reward block suggest the

presence of a reward prediction error, marked by an unexpected reward at the start,

which gradually becomes predictable towards the block's end. This phenomenon was

further investigated by examining the responses in the duration novelty detection task.

In the control condition, where the reward was unpredictable, a distinct response was observed, contrasting with the predictable reward scenario in the deviant condition. In Fig. 6, two example IC neurons responded to the deviant sounds and subsequent rewards, each exhibiting a climbing effect post-peak response. The firing rate of these neurons returned to baseline post-sound for all durations, with a noticeable cluster of neuronal responses following the control condition sound presentation, absent in all four deviant conditions (top row of Fig. 6). When responses were aligned to reward delivery (the bottom row of Fig. 6), notable reward responses were only observed in the control condition ($p < 0.001$, ANOVA), underscoring the auditory IC neurons' capability to encode reward prediction errors.

Decision related signal of IC neurons in duration novelty detection

We have shown the neuronal dynamic of IC neurons in the duration novelty detection task for correct trails. Next, we will explore whether the IC, a lower-level nucleus of the auditory pathway, plays a role in decision making of a duration-related task. During a session where a monkey excelled in performing the task (Fig. 7A), trials were categorized based on the monkey's decisions—whether to press a button or not. Particularly in instances involving two specific deviant durations (357 ms and 425 ms), where the likelihood of pressing the button hovered around 50%, a sufficient number of trials were available to assess the correlation between neuronal activities and the monkey's decisions. This focus aims to delve deeper into how neuronal responses might inform or influence behavioral outcomes in these scenarios.

Analysis did not show the deviant response being influenced by the sequence of

preceding standard sounds (Supplementary Fig. 6), leading to an aggregation of

conditions for evaluation. In an illustrative neuron responding to the 357-ms deviant

sound, the activation was markedly stronger when the monkey identified the sound as

deviant and opted to press the button (indicated by the red color in Fig. 7B), in stark

contrast to when the monkey did not perceive the sound as deviant, thus choosing not

to press the button (illustrated in black color of Fig. 7B). Aligning this observation

with the moment of button press further accentuated the response disparity ($p =$

$1.32e-8$, ANOVA, right panel in Fig. 7B).

To quantitatively assess the decision related signal, an analysis incorporating

Receiver Operating Characteristic (ROC) was conducted (Fig. 7C), revealing a

pronounced difference in firing rates aligned with the act of pressing the button ($p =$

$2.58e-6$, ANOVA, Fig. 7C). The area under the ROC curve (AUC), signifying

detection probability, stood at 0.86 ($p < 0.001$, permutation test), indicating a high

likelihood that an ideal observer could discern between the two choices based solely

on the neuronal firing rate.

Across 69 recorded sessions showcasing reliable behavioral outcomes (Fig. 7D),

population-level PSTHs highlighted a tendency for heightened responses during trials

where the button was pressed ($p = 3.08e-4$, ANOVA, left panel of Fig. 7E), with a

notable difference preceding the button press ($p = 7.93e-7$, ANOVA, right panel of

Fig. 7E). The average detection probability across these sessions was 0.56, surpassing

the chance level of 0.5 ($p = 2.57e-4$, t-test, Fig. 7F), with fifteen out of sixty-nine

neurons showing significant detection probabilities. Specifically for the 425-ms sound

trials, eight out of sixty-eight neurons displayed a significant detection probability,

with an average value of 0.54, also above the chance level (p = 4.30e-2, t-test,

Supplementary Fig. 7).

Discussion

The study unveils novel insights into the inferior colliculus's (IC) role in auditory

processing, highlighting its involvement beyond traditional sensory encoding.

Neurons within the IC exhibit a sustained firing pattern, indicating a robust processing

of sound duration, which is crucial for temporal perception (Fig. 1). A

behavior-dependent "climbing effect" in neuronal firing rate suggests that IC's

response to auditory stimuli is modulated by the context of the behavior, with a

significant increase in response as the sound sequence progresses, implying a link to

sensory experience (Fig. 2, 3). This effect was further supported by experiments that

distinguished between behavior and reward conditions, showing that the climbing

effect is contingent upon the behavioral context rather than mere reward anticipation

(Fig. 4, 5). Additionally, IC neurons encoded reward prediction errors, demonstrating

a nuanced capability to adjust responses based on reward predictability (Fig. 6).

Lastly, evidence of decision signals in IC neurons correlated with the monkey's

decisions in a duration novelty detection task, indicating a direct involvement in

decision-making processes related to auditory perception (Fig. 7). Overall, our results

strongly suggest that the inferior colliculus is actively engaged in sensory experience,

decision making and reward prediction, shedding light on its intricate functions in

these processes.

Sensory experience and response dynamics in IC

The IC neurons showed a rising firing rate after the onset peak (Fig. 2A-D). The

climbing effect was strongly modulated by perception (Supplementary Fig.2),

echoing results from previous research (Metzger et al., 2006). This change was not

due to reward, as shown by an experiment controlling for reward effects (Fig. 4).

Reward only caused minor changes in the response profile (Fig. 4D). Therefore, the

climbing effect reflects perception rather than reward. Another intriguing aspect of the

profile change is its modulation along successive sound presentations in the oddball

paradigm. The rising profile becomes steeper from the first to the fourth stimulation

and then remains relatively stable from the fourth to the seventh stimulation (Fig. 3B).

This modulation by preceding sensory experiences challenges the traditional view of

the IC as merely a relay station, suggesting a more complex role in auditory

processing influenced by both ascending and descending neural pathways. The IC's

extensive descending network of connections, including those from the auditory

cortex (Adams, 1980; Andersen et al., 1980; Bajo and Moore, 2005; Coleman and

Clerici, 1987; Diamond et al., 1969; FitzPatrick and Imig, 1978; Winer et al., 2002),

thalamic nuclei (Adams, 1980; Kuwabara and Zook, 2000; Winer et al., 2002), and

limbic system areas (Adams, 1980; Coleman and Clerici, 1987; Larue et al., 2005;

Marsh et al., 2002), underscores its integration within broader sensory and cognitive

processes, potentially underpinning the observed climbing effect.

The accumulation of the climbing effect across successive sound presentations

appears to be intrinsically linked to the structure of oddball stimulation. Within the

oddball paradigm, both sensory and reward predictions intensify alongside the

recurrence of standard sounds, suggesting that the strength of these predictions could

significantly influence neuronal responses. Our experimentation with rewards has

effectively dismissed the role of reward prediction (Figs. 4 and 5), highlighting the

potential significance of sensory prediction in molding the climbing effect.

Historically, oddball stimuli have been a focal point for investigating novelty

detection within the auditory cortex (Carbajal and Malmierca, 2018; Malmierca et al.,

2015; Rui et al., 2018; Zhai et al., 2019), though many studies lacked incorporation of

a relevant behavioral task (Khouri and Nelken, 2015; Song et al., 2023). Sensory

prediction was indirectly assessed by contrasting predictive and non-predictive stimuli

in control experiments, hence not addressing sensory prediction in a direct, real-time

manner (Carbajal and Malmierca, 2018; Fishman and Steinschneider, 2012; Parras et

al., 2017). The progression of the climbing effect with successive sound presentations

could potentially offer a direct and real-time perspective on the impact of sensory

prediction. Notably, while the onset response remained consistent from the second

through the seventh sound, significant dynamic changes in the late response window

suggest a prolonged period is required for sensory prediction to exert its influence.

Nonetheless, a comprehensive understanding of this dynamic change necessitates

further detailed exploration.

Decision related signal in IC

Having established how behavioral modulation affects responses in IC, we now

delve into the relationship between IC responses and behavioral choices in a

trial-by-trial analysis within a duration novelty detection task. This approach provides

insight into the potential role of sensory neurons in perceptual tasks by comparing

neural responses and perceptual decisions concurrently in the same subject. While

numerous studies have explored this dynamic for cortical neurons (Tsunada et al.,

2016), the contribution of subcortical neurons to decision-making processes remains

less understood. To our knowledge, this study offers the direct correlation between IC

neuron activity and perceptual choice, observed in real-time during experiments with

animals (Fig. 1). These correlations are crucial for understanding the mechanisms by

which population encoding and decoding influence and direct behavior. Focusing on

the novelty detection task—a ubiquitous aspect of sensory systems and an ongoing

phenomenon in daily life—we employed an oddball paradigm with randomized

presentations of standard sound numbers (Fig. 1) to simulate natural occurrences of

novel stimuli. This approach allowed us to closely examine both the behavioral

responses of monkeys and the corresponding neuronal correlates in the IC.

When neural responses were grouped based on behavioral choices, the PSTHs

diverged (Fig. 7B), indicating a strong correlation between neural firing variations

and choice variations. Notably, within the overall population, the two PSTHs, sorted

by the monkeys' choices, diverged before the end of the sound, converged after its end,

and diverged once more prior to the movement decision (Fig. 7E). This pattern during

the sensory encoding phase underscores a potential critical role for IC in managing

duration-sensitive behaviors. Traditionally, decision signals are thought to be

predominantly driven by top-down inputs, as evidenced by previous studies

(Cumming and Nienborg, 2016; Nienborg et al., 2012; Nienborg and Cumming,

2009). While the decision-related signal in the IC may derive from these top-down

mechanisms, our findings also entertain the possibility that variations in IC neuronal

activity could directly modulate higher brain areas, influencing behavioral outcomes.

In either scenario, our findings clearly demonstrate the IC's significant involvement in

cognitive processing, extending well beyond its traditional role as a mere auditory

relay station.

Furthermore, neuronal responses to sound duration typically become more

transient throughout the auditory pathway with neurons in the thalamus and cortex

primarily showing onset responses (Anderson and Linden, 2011; deCharms and

Merzenich, 1996; DeWeese et al., 2003; He, 2001; Metzger et al., 2006; Nevue et al.,

2015; Rauschecker et al., 1995; Xia et al., 2000; Yasui et al., 1991; Zhai et al., 2019).

However, exceptions exist, as sustained neurons have been identified in the medial

geniculate body (MGB) (Allon et al., 1981; Bartlett and Wang, 2011) and the

auditory cortex (AC) (Lu et al., 2001; Malone et al., 2002; Rauschecker et al., 1995;

Wang et al., 2005). In stark contrast, the majority of Inferior Colliculus (IC) neurons

exhibit sustained responses to prolonged sounds across various species, including

frogs (Ratnam and Feng, 1998), bats (Luo et al., 2008), and rodents (Pérez-González

et al., 2006; Xia et al., 2000; Yin et al., 2008). Our examination of IC neurons in

macaque monkeys revealed a similar sustained firing pattern in response to

long-duration sounds (Fig. 1A-B), aligning with observations across diverse species.

This evolutionary conservation of sustained response characteristics in the IC suggests

its capacity to encode sound duration. However, the implications of this capability for

behavior have remained uncertain. The present study offers direct evidence

highlighting the significant role of the IC in decision-making processes related to

duration perception (Fig.7).

Reward effect and reward prediction error in IC

In addition to the role in sensory processing, another important finding in our

study is the existence of a reward effect and a reward prediction error in the IC. The

connection between the IC and the reward system is not very clear, but some studies

have suggested that dopamine may play a role in modulating auditory processing in

the inferior colliculus (Gittelman et al., 2013; Hoyt et al., 2019; Nevue et al., 2015).

Dopamine is a neurotransmitter that is involved in the reward system and other

functions (Hoyt et al., 2019). Dopamine may have heterogeneous effects on the

responses of many neurons in the IC, such as suppressing or enhancing neural activity,

depending on various factors (Hoyt et al., 2019; Nevue et al., 2015). However, the

exact mechanisms and functions of dopamine modulation in the inferior colliculus are

still not fully understood. In this study, we first noticed that some auditory neurons

responded to the reward (Fig. 4B, D), but the reward modulation effect on auditory

responses is slim in the awake monkey (Fig. 4B-E).

Like the response of the dopaminergic neurons in the reward system (Hollerman

and Schultz, 1998), the reward response of IC neurons displays prediction error

signals, which were uncovered in two experiments. Firstly, during the experiment in

which sound was followed by reward, the reward responses were significantly

stronger in the initial 15 trials than in the final 15 trials (Fig. 5A-C), as the reward is a

surprise at the beginning while becoming a routine at the end. Secondly, during the novelty detection task, as the number of standard sounds was randomly chosen and the monkeys were not informed about the end of the stimulation in the control trials, the reward in such trials was not expected, but then a strong response was detected (Fig. 6). By contrast, during the deviant trials, the monkey pressed the button, expected a reward, and no reward response was detected (Fig. 6). Collectively, these findings affirm the presence of reward prediction error signals in the IC, underscoring its sophisticated engagement in reward-related auditory processing dynamics.

Methods

Subjects and Apparatus.

Experiments were conducted in a sound-proof room. Acoustic stimuli were digitally generated using a computer-controlled Auditory Workstation (Tucker-Davis Technologies, TDT, Alachua, FL) and delivered via a loudspeaker (LS50, KEF, UK). The sound pressure was calibrated with a ¼´´condenser microphone (Brüel & Kjær 4954, Nærum, Denmark) and a PHOTON/RT analyzer (Brüel & Kjær). All stimuli were presented contralateral to the recording site.

Data were collected from two male rhesus monkeys (M. mulatta), which were chronically implanted with a plastic head-restraint ring and a recording grid, as described in detail in previous publications (Gong et al., 2024; Yu et al., 2012; Yu et

al., 2015). All procedures were approved by the Animal Care and Use Committee of Zhejiang University and were performed according to the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

We recorded extracellular neural activity using epoxy-coated tungsten microelectrodes (FHC, 2–4 MΩ). Electrodes were inserted into the inferior colliculus based on MRI structure (SF. 1a) through 26-gauge transdural guide tubes and were advanced by a remote-controlled microdrive (FHC). Raw neural activity was amplified and filtered (300 Hz to 3000 Hz). The spike train of the neuron was analyzed off-line. Single units were identified based on waveform shape, latency, and amplitude. Only well-isolated neurons were included in the analyses.

Auditory Stimulation

Tones (100 ms; 5-ms rise-fall time) with multiple combinations of frequencies and intensities were presented to determine the frequency response areas (FRAs). Tones were presented randomly with 5 repetitions at each frequency (0.5–48 kHz in 26 logarithmic steps) and intensity (0–70 dB SPL in 10 dB steps) and interstimulus intervals of 300 ms. We used the FRAs to determine the CF.

To assess the ability of sustained firing in the IC, we presented white noise bursts with three kinds of duration (100 ms, 500 ms, 1000 ms) at 2-s interstimulus intervals. The three sound durations were randomly presented at 60 dB SPL, each for 20 trials.

To explore the reward effect on the neuronal firing of IC neurons, 1000-ms white noise was presented 150 times at 60 dB SPL at 4-s interstimulus interval (Fig. 5A). The initial 50 trials consisted of no reward (early no-reward block), the middle 50

trials consisted of a reward of water 500 ms after the offset of each sound (reward

block); and the final 50 trials again consisted of no reward (late no-reward block). (At

the beginning of the experiment, we presented stimuli only 100 times, consisting of

early no-reward block and reward block and collected 4 IC neurons.)

Behavioral Protocol.

Animals were trained to perform a novelty detection task in two steps: 1) train the

monkeys to press the button after the sound; 2) train the monkeys to press the button

only after the deviant sound is presented. At both steps, the monkeys received a water

reward when they made the correct response.

During the recording session, the sound was presented in blocks (Fig. 1A-B). In

each block, 3 to 6 repeated white noise bursts (standard sound) were presented at

600-ms interstimulus interval, and the last stimulus was either the same sound (the

control condition) or a deviant sound (Fig. 1A-B). The monkey needed to press the

button immediately within 600 ms after the offset of the deviant stimulus to get the

reward of a drop of water. The task reaction time setting precluded the possibility that

the monkey made the decision just because it was the last stimulus in the block; in the

control condition, the monkey did not need to press the button within 600 ms after the

onset of the last stimulus to get the reward. The number of standard stimuli was

randomly selected from 3 to 6 to avoid that the monkey made the decision based on

counting the number of sounds and avoided that the monkey guessed when the

deviant would be presented, so that the monkey had to press the button based on the

detection of the current deviant stimulus. In each recording session, the standard

duration was kept the same (300 ms), but the deviant duration was systematically

changed to control the difficulty of the task according to the following formula:

Duration_{deviant} = Duration _{standard} * λ^n (n=0,1,2,3,4) (λ=1.19), where Duration_{deviant} and

Duration_{standard} were deviant and standard duration, respectively. Thus, in each session,

5 conditions were presented. Each condition was usually presented 30-40 times.

During the recording, all sounds were presented at 60 dB SPL with 5 ms rise-fall time

in the profile.

We assessed the behavior of the monkey according to two criteria: 1) the monkey

made the correct rejection. In the control condition, the ratio of pressing a button had

to be less than 0.1; 2) The monkey made the correct hit. In the most differential

condition, the ratio of pressing the button had to be greater than 0.85. Only the

behavior meeting these two criteria was kept for further analysis.

Data analysis

To characterize the climbing effect, we defined a factor termed the response dynamic

index (RDI). RDI is calculated according to the following formula:

$$RDI = \frac{Fir_{offset}^{[-100\ 0]} - Fir_{onset}^{[100\ 200]}}{Fir_{offset}^{[-100\ 0]} + Fir_{onset}^{[100\ 200]}}$$

$Fir_{offset}^{[-100\ 0]}$ and $Fir_{onset}^{[100\ 200]}$ were the neuronal firing rates within the late window

(-100-0 ms relative to the offset of the sound) and the after-peak window (100-200 ms

relative to the onset of the sound), respectively.

To define a reward-responsive neuron, we compared the response within 100 ms

after the reward and the response within 200 ms before the reward was offered in the

reward blocks (paired t-test). A neuron was considered reward-responsive only if the

difference reached significance ($p < 0.05$).

Behavioral performance was quantified by plotting the proportion of "pressing

button" choices as a function of duration difference between the deviant and standard

sounds in ratio. Psychometric data were fit with a Gaussian integral function:

$$p(r) = \frac{1}{b\sqrt{2\pi}} \int_0^r e^{-\frac{(x-a)^2}{2b^2}} \, dx$$

529   In this expression, p is the proportion of pressing button choices, r is the ratio

530   between deviant and standard duration.

533
534

 Figure Legends

 Figure 1 Sustained Firing in IC and Novelty Detection Paradigm Based on Sound

 Duration. (A) Raster plots of a representative neuron depicting the response profile to

 white noise stimuli with varying durations (100 ms, 500 ms, and 1000 ms). **(B)**

 PSTHs of neuronal population responses (n=104) to the stimuli described in **A** (black,

 100-ms white noise; red, 500-ms white noise; blue, 1000-ms white noise). **(C)**

 Schematic display of novelty detection behavior. A button is positioned in front of the

 monkey, and a loudspeaker is placed contralateral to the recording site at the height of

 the monkey's ear. Within each experimental block, a series of repeated 300-ms white

 noise bursts (WNB) serves as the standard stimulus, accompanied by a rare-duration

 WNB serving as the deviant stimulus. Following the presentation of the deviant

 stimulus, the primate is required to press the designated button within a 600-ms

 timeframe to obtain a water reward. **(D)** Oddball stimulation paradigm employed in

 all experimental blocks. While the duration of the standard sound remains constant at

 300 ms, the duration of the novel sound deviates from the standard sound across five

 distinct levels. This deviation is determined by the formula: $\text{Duration}_{\text{deviant}} =$

 $\text{Duration}_{\text{standard}} \times \lambda^{\text{n}}$ (where n=0,1,2,3,4; $\lambda = 1.19$). Furthermore, the number of

 standard stimuli within each block is randomly selected from a range of 3 to 6. Upon

 completion of the deviant sound, the primate is required to press the designated button

 within a 600-ms window to receive the reward. In the absence of a deviant sound

 (control condition), the reward is granted if the primate refrains from pressing the

 button within 600 ms after the onset of the final sound. **(E)** Cumulative Gaussian fits

of psychophysical data in one example session. The duration of the deviant sound is

plotted along the abscissa, while the ratio of button presses is plotted along the

ordinate.

Figure 2 Neuronal climbing effect in novelty detection behavior. (A) Peri-stimulus

time histograms (PSTHs) depicting example neuronal responses to different deviant

sounds aligned to deviant onset (light gray, 300-ms deviant sound; gray, 357-ms

deviant sound; black, 425-ms deviant sound; pink, 506-ms deviant sound; red, 600-ms

deviant sound). **(B)** The same neuronal responses as in **A**, but this time aligned to the

offset of the deviant stimulus. The PSTHs represent the neural activity before the

offset of each deviant sound. The color scheme remains consistent with **A** to indicate

the different deviant durations. **(C)** Firing rate to deviant stimuli in two windows: the

late window ([-100 0] ms relative to offset time) and the onset window ([0 60] ms

relative to onset time) (red, late window; black, onset window). **(D)** PSTH showing

neuronal responses to standard sounds (1-7) from the neuron described in **A**. **(E)**

Normalized firing rate of the neuron described in **A** to standard sounds in two

temporal windows (late window: [-100 0] ms relative to offset time; onset window: [0

60] ms relative to onset time) plotted as a function of the order of the sounds (red, late

window; black, onset window).

Figure 3 Relationship between neuronal climbing effect and order of sounds. (A)

Average normalized responses to standard sounds in the onset window and offset

window (red, late window; black, onset window. n=99). The left axis represents the

scale for responses in the late window, and the right axis represents the scale for

responses in the onset window. **(B)** PSTHs depicting the responses of the neuronal

population to the 300-ms sounds, ranging from the first to the sixth order in the block

(black, first; gray, second; blue, third; light blue, fourth; cyan, fifth; pink, sixth; red,

seventh). The left axis represents the response scale for sounds from the second to the

seventh, and the right axis represents the response scale for the first sound.

Figure 4 Influence of reward on inferior colliculus neurons. **(A)** Schematic

representation of the reward protocol. The protocol involves the presentation of a

1000-ms white noise stimulus at 60 dB SPL, repeated 150 times with a 4-second

interstimulus interval. The experimental design comprises three blocks: an initial

'early no-reward' block consisting of 50 trials without reward, a 'reward' block of 50

trials with water reward administered 500 ms after sound offset, and a final 'late

no-reward' block of 50 trials without reward. **(B)** Raster plots of a representative

neuron depicting the response patterns during the reward protocol. **(C)** PSTHs of the

neuron in **B** representing the neuronal responses during the reward block, early

no-reward block, and late no-reward block (red, reward block; black, early no-reward

block; blue, late no-reward block). **(D)** PSTHs of the neuronal population exhibiting

significant reward responses (n=24) representing the neuronal responses during the

reward block, early no-reward block, and late no-reward block (red, reward block;

black, early no-reward block; blue, late no-reward block). **(E)** Equivalent responses as

in **D** but from neurons with nonsignificant reward responses (n=35). **(F)** Scatterplots

of RDI in reward block versus RDI in behavior protocol (square, monkey B; circle,

monkey J; n=63).

Figure 5 Reward prediction error in inferior colliculus neurons. (A) PSTHs of the

representative neuron in Figure 6B depicting its response profiles during the initial 15

reward trials (red) and the final 15 reward trials (black). **(B)** PSTHs of the neuronal

population with significant reward responses (n=28) demonstrating the firing activity

of these neurons during the initial 15 reward trials (red) and the final 15 reward trials

(black). **(C)** Scatterplots of reward responses in initial 15 reward trials versus final 15

reward trials from neurons in **B**. The data points are color-coded, with solid circles

representing neurons that exhibit a significant difference in reward responses between

the initial and final reward trials. Conversely, open circles represent neurons wherein

the difference in reward responses between the two trial sets is nonsignificant. The

square and circle symbols correspond to monkey B and monkey J, respectively. **(D)**

PSTHs of a neuron with nonsignificant reward responses demonstrating the response

patterns of the neuron, similar to those presented in **A**. **(E-F)** Equivalent PSTHs and

scatterplots as in **B-C** but from neurons with nonsignificant reward responses (n=35).

Figure 6 Reward prediction error in novelty detection behavior. (A) PSTHs

illustrating neuronal responses aligned to deviant onset (top) and reward time

(bottom). **(B)** PSTHs demonstrating neuronal responses aligned to deviant onset (top)

and reward time (bottom) in another neuron.

Figure 7 Neuronal responses of decision making in novelty detection behavior.

(A) Cumulative Gaussian fits of psychophysical data in the example recording session.

The duration of the deviant sound is plotted along the abscissa, while the ratio of

button presses is plotted along the ordinate. **(B)** PSTHs showing example neuronal

responses to deviant sound with 357-ms duration (red, pressing button after deviant

sound; black, no pressing button after deviant sound). The left side of the subfigure

shows the PSTHs aligned to the onset of the deviant sound, while the right side shows

the PSTHs aligned to the time of button presses. **(C)** Top: Distribution of neuronal

responses to 357-ms deviant sound categorized based on the behavioral choice made

by the subject (red, pressing the button; black, no pressing the button). Bottom,

receiver operating characteristic (ROC) analysis of comparison between two

distributions. **(D)** Cumulative Gaussian fits to psychophysical data in population

(n=69). **(E)** PSTHs showing neuronal responses to deviant sound with 357 ms

duration in population (red, pressing button after deviant sound; black, no pressing

button after deviant sound; left, aligned to deviant onset; right, aligned to time of

pressing button). **(F)** Distribution of detection probability: This panel displays the

distribution of detection probability (DP), with significant DP indicated by black bars.

Figure S1: Neurons in the inferior colliculus with sustained response. (A) 7T MRI

with recording electrode: High-resolution 7T MRI scans displaying the recording

electrode in place within the brains of two monkeys (left, monkey J; right, monkey B).

(B) Localization of recording sites on fMRI image. The red dots overlaid on the fMRI

image indicate the locations of the recording sites within the inferior colliculus. The

fMRI image was acquired at a plane positioned 1 mm posterior to the ear bar zero

(EBZ). The red dots represent the projection of the recording sites onto this specific

plane.

Figure S2: Comparison of neuronal responses between behavior protocol and

non-behavior protocol. **(A)** PSTHs comparing neuronal responses in non-behavior

and behavior protocols (red: responses to 506-ms deviant sound in behavior protocol;

blue: responses to 500-ms sound in non-behavior protocol; n=99). The colored area

indicates the error band (standard error). **(B)** Scatterplots of RDI in non-behavior

protocol versus behavior protocol (square, monkey B; circle, monkey J). Each data

point represents the RDI value of an individual neuron, allowing for a comparison

between the two protocols.

Figure S3: Response dynamic index (RDI) of different deviant sounds. (A)

Scatterplots of RDI for 506-ms deviant sound versus RDI for 357-ms deviant sound

(n=99). **(B)** Scatterplots of RDI for 506-ms deviant sound versus RDI for 425-ms

deviant sound. **(C)** Scatterplots of RDI for 506-ms deviant sound versus RDI for

600-ms deviant sound.

Figure S4: Three-dimensional representation of recorded neurons based on

response dynamic index (RDI). (A) Three-dimensional representation of the

recorded neurons obtained from monkey B relative to EBZ. Each neuron is assigned a

distinct color based on its RDI. **(B)** Three-dimensional representation of the recorded

neurons obtained from monkey J relative to EBZ. Each neuron is color-coded

according to its RDI.

Figure S5: Influence of reward on neuronal responses. (A) PSTHs showing

neuronal population responses (n=22) in reward protocol with 500-ms sound (blue)

and behavior protocol (red). **(B)** Scatterplots of RDI in reward block with 500-ms

sound versus RDI in behavior protocol (square, monkey B; circle, monkey J).

Figure S6: Influence of sound numbers on neuronal responses to 357-ms deviant

sound. (A) PSTHs illustrating the neuronal responses to 357-ms deviant sounds with

different orders in the block when the monkey pressed the button. The PSTHs are

color-coded, with grey representing the fourth order, light gray representing the fifth

order, blue representing the sixth order, and red representing the seventh order. These

responses are obtained from the neuron depicted in Figure 8**B**. **(B)** PSTHs illustrating

the neuronal responses to 357-ms deviant sounds with different orders in the block

when the monkey didn't press the button. **(C)** Populational neuronal responses when

the monkey pressed the button (n=49). **(D)** Populational neuronal responses when the

monkey did not press the button (n=49).

Figure S7: Distribution of detection probability (DP) for 425-ms deviant sounds.

Within the distribution, significant DP values are indicated by the presence of black

bars.

References

Adams, J.C. (1980). Crossed and descending projections to the inferior colliculus. Neurosci Lett *19*, 1-5.

Allon, N., Yeshurun, Y., and Wollberg, Z. (1981). Responses of single cells in the medial geniculate body
of awake squirrel monkeys. Exp Brain Res *41*, 222-232.

Andersen, R.A., Snyder, R.L., and Merzenich, M.M. (1980). The topographic organization of
corticocollicular projections from physiologically identified loci in the AI, AII, and anterior auditory
cortical fields of the cat. J Comp Neurol *191*, 479-494.

Anderson, L.A., and Linden, J.F. (2011). Physiological differences between histologically defined
subdivisions in the mouse auditory thalamus. Hear Res *274*, 48-60.

Ayala, Y.A., Pérez-González, D., and Malmierca, M.S. (2016). Stimulus-specific adaptation in the inferior
colliculus: The role of excitatory, inhibitory and modulatory inputs. Biol Psychol *116*, 10-22.

Bajo, V.M., and Moore, D.R. (2005). Descending projections from the auditory cortex to the inferior
colliculus in the gerbil, Meriones unguiculatus. J Comp Neurol *486*, 101-116.

Bartlett, E.L., and Wang, X. (2011). Correlation of neural response properties with auditory thalamus
subdivisions in the awake marmoset. J Neurophysiol *105*, 2647-2667.

Carbajal, G.V., and Malmierca, M.S. (2018). The Neuronal Basis of Predictive Coding Along the Auditory
Pathway: From the Subcortical Roots to Cortical Deviance Detection. Trends Hear *22*,
2331216518784822.

Cohen, Y.E., and Knudsen, E.I. (1999). Maps versus clusters: different representations of auditory space
in the midbrain and forebrain. Trends in neurosciences *22*, 128-135.

Coleman, J.R., and Clerici, W.J. (1987). Sources of projections to subdivisions of the inferior colliculus
in the rat. J Comp Neurol *262*, 215-226.

Cumming, B.G., and Nienborg, H. (2016). Feedforward and feedback sources of choice probability in
neural population responses. Curr Opin Neurobiol *37*, 126-132.

deCharms, R.C., and Merzenich, M.M. (1996). Primary cortical representation of sounds by the
coordination of action-potential timing. Nature *381*, 610-613.

DeWeese, M.R., Wehr, M., and Zador, A.M. (2003). Binary spiking in auditory cortex. J Neurosci *23*,
7940-7949.

Diamond, I.T., Jones, E.G., and Powell, T.P. (1969). The projection of the auditory cortex upon the
diencephalon and brain stem in the cat. Brain Res *15*, 305-340.

Duque, D., and Malmierca, M.S. (2015). Stimulus-specific adaptation in the inferior colliculus of the
mouse: anesthesia and spontaneous activity effects. Brain Struct Funct *220*, 3385-3398.

Fishman, Y.I., and Steinschneider, M. (2012). Searching for the mismatch negativity in primary auditory
cortex of the awake monkey: deviance detection or stimulus specific adaptation? J Neurosci *32*,
15747-15758.

FitzPatrick, K.A., and Imig, T.J. (1978). Projections of auditory cortex upon the thalamus and midbrain
in the owl monkey. J Comp Neurol *177*, 573-555.

Gittelman, J.X., Perkel, D.J., and Portfors, C.V. (2013). Dopamine modulates auditory responses in the
inferior colliculus in a heterogeneous manner. J Assoc Res Otolaryngol *14*, 719-729.

Gong, Y., Song, P., Du, X., Zhai, Y., Xu, H., Ye, H., Bao, X., Huang, Q., Tu, Z., Chen, P.*, et al.* (2024). Neural
correlates of novelty detection in the primary auditory cortex of behaving monkeys. Cell reports *43*,
113864.

Harris, S., Afram, R., Shimano, T., Fyk-Kolodziej, B., Walker, P.D., Braun, R.D., and Holt, A.G. (2021).
Dopamine in Auditory Nuclei and Lemniscal Projections is Poised to Influence Acoustic Integration in
the Inferior Colliculus. Front Neural Circuits *15*, 624563.

He, J. (2001). On and off pathways segregated at the auditory thalamus of the guinea pig. J Neurosci
21, 8672-8679.

Hollerman, J.R., and Schultz, W. (1998). Dopamine neurons report an error in the temporal prediction
of reward during learning. Nat Neurosci *1*, 304-309.

Hoyt, J.M., Perkel, D.J., and Portfors, C.V. (2019). Dopamine Acts via D2-Like Receptors to Modulate
Auditory Responses in the Inferior Colliculus. eNeuro *6*.

Jones, E.G. (2012). The thalamus (Springer Science & Business Media).

Khouri, L., and Nelken, I. (2015). Detecting the unexpected. Curr Opin Neurobiol *35*, 142-147.

Kremkow, J., and Alonso, J.M. (2018). Thalamocortical Circuits and Functional Architecture. Annu Rev
Vis Sci *4*, 263-285.

Kuwabara, N., and Zook, J.M. (2000). Geniculo-collicular descending projections in the gerbil. Brain
Res *878*, 79-87.

Larue, D., Prieto, J., and Winer, J. (2005). The central gray and the inferior colliculus: an auditory-limbic
interface. Paper presented at: Assoc Res Otolaryngol Abstr.

Lu, T., Liang, L., and Wang, X. (2001). Temporal and rate representations of time-varying signals in the
auditory cortex of awake primates. Nat Neurosci *4*, 1131-1138.

Luo, F., Metzner, W., Wu, F., Zhang, S., and Chen, Q. (2008). Duration-sensitive neurons in the inferior
colliculus of horseshoe bats: adaptations for using CF-FM echolocation pulses. J Neurophysiol *99*,
284-296.

Malmierca, M.S., Anderson, L.A., and Antunes, F.M. (2015). The cortical modulation of
stimulus-specific adaptation in the auditory midbrain and thalamus: a potential neuronal correlate for
predictive coding. Front Syst Neurosci *9*, 19.

Malmierca, M.S., Cristaudo, S., Perez-Gonzalez, D., and Covey, E. (2009). Stimulus-specific adaptation
in the inferior colliculus of the anesthetized rat. J Neurosci *29*, 5483-5493.

Malone, B.J., Scott, B.H., and Semple, M.N. (2002). Context-dependent adaptive coding of interaural
phase disparity in the auditory cortex of awake macaques. J Neurosci *22*, 4625-4638.

Marsh, R.A., Fuzessery, Z.M., Grose, C.D., and Wenstrup, J.J. (2002). Projection to the inferior colliculus
from the basal nucleus of the amygdala. J Neurosci *22*, 10449-10460.

Metzger, R.R., Greene, N.T., Porter, K.K., and Groh, J.M. (2006). Effects of reward and behavioral
context on neural activity in the primate inferior colliculus. J Neurosci *26*, 7468-7476.

Miterko, L.N., Lackey, E.P., Heck, D.H., and Sillitoe, R.V. (2018). Shaping Diversity Into the Brain's Form
and Function. Front Neural Circuits *12*, 83.

Nevue, A.A., Elde, C.J., Perkel, D.J., and Portfors, C.V. (2015). Dopaminergic Input to the Inferior
Colliculus in Mice. Front Neuroanat *9*, 168.

Nevue, A.A., Felix, R.A., 2nd, and Portfors, C.V. (2016). Dopaminergic projections of the
subparafascicular thalamic nucleus to the auditory brainstem. Hear Res *341*, 202-209.

Nienborg, H., Cohen, M.R., and Cumming, B.G. (2012). Decision-related activity in sensory neurons:
correlations among neurons and with behavior. Annu Rev Neurosci *35*, 463-483.

Nienborg, H., and Cumming, B.G. (2009). Decision-related activity in sensory neurons reflects more
than a neuron's causal effect. Nature *459*, 89-92.

Ono, M., and Ito, T. (2015). Functional organization of the mammalian auditory midbrain. J Physiol Sci
65, 499-506.

Parras, G.G., Nieto-Diego, J., Carbajal, G.V., Valdes-Baizabal, C., Escera, C., and Malmierca, M.S. (2017).
Neurons along the auditory pathway exhibit a hierarchical organization of prediction error. Nat
Commun *8*, 2148.

Perez-Gonzalez, D., Malmierca, M.S., and Covey, E. (2005). Novelty detector neurons in the
mammalian auditory midbrain. Eur J Neurosci *22*, 2879-2885.

Pérez-González, D., Malmierca, M.S., Moore, J.M., Hernández, O., and Covey, E. (2006). Duration
selective neurons in the inferior colliculus of the rat: topographic distribution and relation of duration
sensitivity to other response properties. J Neurophysiol *95*, 823-836.

Pickles, J.O. (2015). Auditory pathways: anatomy and physiology. Handb Clin Neurol *129*, 3-25.

Ratnam, R., and Feng, A.S. (1998). Detection of auditory signals by frog inferior collicular neurons in
the presence of spatially separated noise. J Neurophysiol *80*, 2848-2859.

Rauschecker, J.P., Tian, B., and Hauser, M. (1995). Processing of complex sounds in the macaque
nonprimary auditory cortex. Science (New York, NY) *268*, 111-114.

Rui, Y.Y., He, J., Zhai, Y.Y., Sun, Z.H., and Yu, X. (2018). Frequency-dependent stimulus-specific
adaptation and regularity sensitivity in the rat auditory thalamus. Neuroscience *392*.

Sherman, S.M. (2005). Thalamic relays and cortical functioning. Prog Brain Res *149*, 107-126.

Sherman, S.M. (2016). Thalamus plays a central role in ongoing cortical functioning. Nat Neurosci *19*,
533-541.

Song, P., Zhai, Y., and Yu, X. (2023). Stimulus-specific adaptation (SSA) in the auditory system:
Functional relevance and underlying mechanisms. Neurosci Biobehav Rev *149*, 105190.

Tsunada, J., Liu, A.S., Gold, J.I., and Cohen, Y.E. (2016). Causal contribution of primate auditory cortex
to auditory perceptual decision-making. Nat Neurosci *19*, 135-142.

Valdés-Baizabal, C., Casado-Román, L., Bartlett, E.L., and Malmierca, M.S. (2021). In vivo whole-cell
recordings of stimulus-specific adaptation in the inferior colliculus. Hear Res *399*, 107978.

Valdés-Baizabal, C., Parras, G.G., Ayala, Y.A., and Malmierca, M.S. (2017). Endocannabinoid
Modulation of Stimulus-Specific Adaptation in Inferior Colliculus Neurons of the Rat. Sci Rep *7*, 6997.

Wang, X., Lu, T., Snider, R.K., and Liang, L. (2005). Sustained firing in auditory cortex evoked by
preferred stimuli. Nature *435*, 341-346.

Winer, J.A., Chernock, M.L., Larue, D.T., and Cheung, S.W. (2002). Descending projections to the
inferior colliculus from the posterior thalamus and the auditory cortex in rat, cat, and monkey. Hear
Res *168*, 181-195.

Xia, Y.F., Qi, Z.H., and Shen, J.X. (2000). Neural representation of sound duration in the inferior
colliculus of the mouse. Acta Otolaryngol *120*, 638-643.

Yasui, Y., Nakano, K., Kayahara, T., and Mizuno, N. (1991). Non-dopaminergic projections from the
substantia nigra pars lateralis to the inferior colliculus in the rat. Brain research *559*, 139-144.

Yin, S., Chen, Z., Yu, D., Feng, Y., and Wang, J. (2008). Local inhibition shapes duration tuning in the

inferior colliculus of guinea pigs. Hear Res *237*, 32-48.

Yu, X.J., Dickman, J.D., and Angelaki, D.E. (2012). Detection thresholds of macaque otolith afferents. J

Neurosci *32*, 8306-8316.

Yu, X.J., Dickman, J.D., DeAngelis, G.C., and Angelaki, D.E. (2015). Neuronal thresholds and

choice-related activity of otolith afferent fibers during heading perception. Proc Natl Acad Sci U S A

112, 6467-6472.

Zhai, Y.Y., Sun, Z.H., Gong, Y.M., Tang, Y., and Yu, X. (2019). Integrative stimulus-specific adaptation of

the natural sounds in the auditory cortex of the awake rat. Brain Struct Funct *224*, 1753-1766.

Fig. 1

A

B

C

Novelty?

stimulation

pressing button

water

90°

D

$Duration_{std} = 300\ ms$
$Duration_{dev} = Duration_{std} \times 1.19^{n}$
$(n = 0, 1, 2, 3, 4)$

3~6

$dev|_{n=0}$
$dev|_{n=1}$
$dev|_{n=2}$
$dev|_{n=3}$
$dev|_{n=4}$

E

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. S1

Fig. S2

Fig. S3

Fig. S4

Fig. S5

Fig. S6

Fig. S7