Untargeted metabolomics reveals anion and organ-specific biochemistry of salinity tolerance in willow

Eszter Sas[1], Adrien Frémont[1], Emmanuel Gonzalez[2,3], Mathieu Sarrazin[4], Simon Barnabé[5], Michel Labrecque[1,6], Nicholas James Beresford Brereton[7†] and Frédéric Emmanuel Pitre[1,6]

[1] Institut de recherche en biologie végétale (IRBV), Département de sciences biologiques, Université de Montréal - 4101 Sherbrooke East, Montreal, QC H1X 2B2, Canada
[2] Canadian Centre for Computational Genomics (C3G) - Department of Human Genetics, McGill University, 740 Dr. Penfield avenue, Montreal, QC H3A 0G1, Canada.
[3] Microbiome Research Platform - McGill Interdisciplinary Initiative in Infection and Immunity (MI4), Genome Centre, McGill University, Montreal, QC, Canada
[4] Centre d'Étude des Procédés Chimiques du Québec (CÉPROCQ), Collège de Maisonneuve, 6220 Rue Sherbrooke East, Montréal, QC H1N 1C1, Canada
[5] Département de chimie, biochimie et physique, Université du Québec à Trois- Rivières, 3351 boulevard des Forges, Trois-Rivières, QC G8Z 4M3, Canada
[6] Montreal Botanical Garden, 4101 Sherbrooke East, Montreal, QC H1X 2B2, Canada
[7] UCD School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland

† Corresponding author: nicholas.brereton@ucd.ie

Key words: Salt stress • Metabolomics • *Salix* • Sulfate / Sulfur • Sulphate / Sulphur • Chloride • Anion

1   Abstract

Willows can alleviate soil salinisation while generating sustainable feedstock for biorefinery, yet the metabolomic adaptations underlying their salt tolerance remain poorly understood. Testing two environmentally abundant salts, the response of *Salix miyabeana* was assessed after treatment with a moderate concentration of NaCl, and both moderate and high concentrations of Na_2SO_4 in a 12-week pot trial. Willows tolerated salts across all treatments (up to 9.1dS m^{-1} soil EC_e), maintaining photosynthesis and biomass while selectively partitioning ions, confining Na^+ to roots and accumulating Cl^- and SO_4^{2-} in the canopy, and adapting to osmotic stress via reduced stomatal conductance. Untargeted LC-MS/MS captured over 5,000 putative compounds, characterising the baseline willow metabolome, including 278 core compounds constitutively produced across organs. Comparative statistical analyses revealed widespread metabolic reprogramming in response to soil salinity, altering 28% of the overall metabolome, and highlighting organ-tailored regulation. Comparing both salt forms at equimolar sodium, generalised salinity responses were limited to 3% of the metabolome, predominantly in roots. Anion-specific metabolomic responses were more extensive, with NaCl reducing carbohydrates and TCA intermediates, thereby exerting pressure on carbon and energy resources, alongside the accumulation of root structuring compounds, antioxidants flavonoids, and fatty acids. In contrast, Na_2SO_4 salinity triggered accumulation of sulphur-containing larger peptides, suggesting that excess sulphate incorporation leverage ion toxicity to produce specialized salt-tolerance associated metabolites. This high-depth picture of the willow metabolome underscores the importance of capturing plant adaptations to salt stress at organ-scale and considering ion-specific contributions to soil salinity.

2   Introduction

Soil salinisation represents a major threat to agricultural productivity and environmental health on a global scale. It is estimated that over 20% of irrigated cropland is impacted by salt accumulation (FAO, 2015), reducing yields and critical ecosystem biodiversity (Zörb et al., 2019). Climate change and human activities are predicted to intensify both natural and anthropogenic soil salinisation processes, further threatening food security (Singh, 2021). While salt stress research typically investigates NaCl, Na_2SO_4, derived from anthropogenic sources like industrial sulphur emissions and agricultural practices, as well as natural volcanic and marine environments (Dick et al., 2008), represents another widespread salt. The use of salt-tolerant biomass crops able to sequester salt and improve soil health, could help reclaim and repurpose salt-impacted areas (Abreu et al., 2022; Quinn et al., 2015).

Soil salinisation results from soluble ions accumulation such as Na^+, K^+, Mg^{2+}, Ca^{2+}, Cl^-, SO_4^{2-} and CO_3^{2-}. When in contact with roots, these ions can induce both osmotic stress and ion toxicity (Munns and Tester, 2008). The osmotic component, primarily driven by any dissolved salt concentration, alters water potential gradients, disrupting cell turgor pressure and elongation. In response, stomatal closure limits water loss via transpiration but also impairs photosynthate supply, ultimately constraining growth through both water deficit and reduced carbon availability (Munns et al., 2020b). Ion toxicity, however, is dependent on the nature of the ion, with different elements inducing unique patterns of signalling, transport, oxidative damage, energy allocation, and hormone-regulated growth processes (Llanes et al., 2013; Reginato et al., 2021; Richter et al., 2019).

Chloride (Cl^-), an essential micronutrient for plants, contributes to vital physiological processes (Wege et al., 2017). As a major osmotically active solute, chloride fluxes are involved in vacuolar turgor required for cell expansion and are implicated in stomatal aperture regulation, contributing to water balance and gas exchange (Teakle and Tyerman, 2010). Additionally, chloride acts as a counter-ion, regulating membrane potential, intracellular pH gradients, electrical signalling in the cytoplasm (White and Broadley, 2001) and serves as a cofactor in the oxygen-evolving complex of photosystem II (Ifuku, 2015). However, in salt-impacted soils, excess chloride damages plants, leading to impaired photosynthesis, reduced NO_3^- uptake (Abdelgadir et al., 2005) and general enzymatic activity (Geilfus, 2018). In contrast, sulphate (SO_4^{2-}), the major bioavailable form of sulphur, is an essential macronutrient involved in physiological and biochemical processes (Prasad and Shivay, 2018; Watanabe and Hoefgen, 2019). Sulphur is required for the production of the amino acids cysteine and methionine, peptide derivatives such as glutathione, and disulphide bonds contributing to enzyme activity and stability. The production of chlorophyll, vitamins and certain plant defence specialised metabolites, such as glucosinolates, also depend upon sulphur (Chan et al., 2019; Rausch and Wachter, 2005). Plants regulate sulphate uptake and assimilation through sulphate specific transporters to avoid toxicity (Buchner et al., 2004), however in some crop species, such as *Brassica rapa*, excess sulphate ions can create nutrient deficiency by competing with the absorption of other ions, especially calcium and phosphorus (Reich et al., 2017).

Plants mediate salt stress through a coordinated series of physiological, structural, biochemical and molecular adjustments (Arif et al., 2020; Deinlein et al., 2014; Gupta and Huang, 2014; van Zelm et al., 2020; Yang and Guo, 2018). Although pathways vary between plant species (Sanchez et al., 2008) and depend on the type of ions involved (Reich et al., 2017; Richter et al., 2019), key tolerance mechanisms include osmotic regulation, ion homeostasis, antioxidant defence, and growth modulation. A primary adaptive response to osmotic stress is the accumulation of osmolytes, which regulate water status by reducing cell osmotic potential (D'Amelia et al., 2018). This common osmotic stress response is not exclusively induced by excess salt nor is dependent on salt type (Munns, 2002). Osmolytes include inorganic ions (Rodriguez et al., 1997; Wege et al., 2017) and organic compounds such as amino acids (e.g. proline), polyamines (e.g. spermidine), quaternary ammonium compounds (e.g. glycine betaine), sugars (e.g. sucrose, trehalose), and polyols (e.g. mannitol, sorbitol) (D'Amelia et al., 2018; Slama et al., 2015; Yang and Guo, 2018). While the biosynthesis of these osmoprotectants imposes a metabolic energy cost (Munns et al., 2020a), their functions extend beyond osmotic adjustment to detoxification processes such as antioxidant activity and chaperone-mediated protein stabilisation (Yancey, 2005). The upregulation of enzymatic (e.g. superoxide dismutase SOD, catalase CAT, peroxidase POX) and non-enzymatic (e.g. glutathione, ascorbate, flavonoids, anthocyanins) antioxidants is crucial for mitigating damage to cellular components such as lipids, proteins, and DNA caused by reactive oxygen species (ROS) (Gill and Tuteja, 2010a). More generally, plants avoid cytotoxicity and maintain ion homoeostasis through salt exclusion from roots, compartmentalisation into the vacuole or apoplast, selective transport and excretion through salt glands. Additionally, hormonal signalling (Ryu and Cho, 2015), modification of root architecture (Zou et al., 2022), xylem alteration and loading (Janz et al., 2012) and symbiotic associations with microorganisms such as arbuscular mycorrhizal fungi (Evelin et al., 2009) also play significant roles in enhancing plant tolerance to salinity.

Salicaceae, including poplars (*Populus sp.*) and willows (*Salix sp.*), are common phytoremediation and biorefinery crops on marginal lands (Brereton et al., 2017; Dimitriou and Aronsson, 2005; Gonzalez et al., 2018; Rodzkin and Volk, 2018; Sas et al., 2021; Volk et al., 2006), with high potential for growth on salinised soils. Some cultivars can maintain transpiration and growth under saline conditions up to 6-8 dS m^{-1} (Bilek et al., 2020; Chen and Polle, 2010; Huang et al., 2020; Mirck and Zalesny, 2015). This tolerance likely engages high metabolic diversity, including reactive oxygen species (ROS) protective phenolics and other antioxidants (Sui

and Wang, 2020; Zhou et al., 2020). However, understanding of the metabolomic toolkit employed beyond
these compounds, and their variation across different organs and salts, remains limited.

In this study, the shared and unique metabolomic adjustments to moderate NaCl, moderate Na_2SO_4 and high
Na_2SO_4 salinity treatment are assessed across multiple organs (Fig.1) to determine generalised and ion-specific
salt tolerance metabolome in willow.

3   Methods

3.1   Experimental design, plant growth, sample collection and physiological traits

The willow cultivar *Salix miyabeana* 'SX64' was grown from cuttings in a polyethylene grow tunnel within 15 L
pots filled with all-purpose growing soil (Promix^{TM}), sand and vermiculite (1:1:1 volume-based). After 56 days
of growth for root establishment, salt treatments were applied every 3 days during watering over 28 days to
progressively build up soil salinity and to avoid salt shock response (Shavrukov, 2013). The four treatments
were: tap water control (EC_{water}=0.3 mS cm^{-1}), 51mM NaCl ($EC_{solution}$=6.5 dS m^{-1}) and 26mM Na_2SO_4 ($EC_{solution}$=6.7
dS m^{-1}), both considered moderately saline, and 51mM Na_2SO_4 ($EC_{solution}$=13.4 dS m^{-1}), considered highly saline
(Mirck and Zalesny, 2015). The treatments were arranged in a randomised block design with seven replicate
plants (seven independent pots, n=7) per treatment (Fig.1). Leaves, stems, and roots were sampled after 28
days of treatment (84 days of growth). A representative subsample of each fraction was immediately flash-
frozen in liquid nitrogen and stored at -70°C for metabolomic analysis.

Shoot heights and numbers were measured weekly. Fresh weight (FW) at harvest, oven-dried biomass weight
(DW) and moisture content were measured for each organ. The LICOR-600 porometer/fluorometer was used
to measure photosynthetic parameters, including stomatal conductance and the quantum efficiency in light of
photosystem II ΦPSII (Genty et al., 1989). Soil salinity was inferred at harvest from electroconductivity of the
supernatant solution from a soil:water mixture (1:1) and converted to the electroconductivity of saturated soil
extract (EC_e) using the correlation EC_e=2.23 $EC_{(1:1)}$-0.58 (R^2=0.98) (Sonmez et al., 2008).

Elemental analysis of roots, stems, and leaves was performed on ground dry biomass (<500 μm) by Natural
Resources Analytical Laboratory at University of Alberta (AB, Canada). Briefly, 0.1±0.02 g of sample was
digested overnight, with 5 mL of Trace Metal Grade HNO_3 using the Microwave Accelerated Reaction System
(method EPA 3051a (modified)), diluted to 25 mL with milliQ water and analysed for mineral nutrients (Al, B,
Ca, Cu, Fe, K, Mg, Mn, Mo, Na, Ni, P, S, Zn) using a Thermo iCAP6300 Duo inductively coupled plasma-optical
emission spectrometer (ICP-OES). The anions Cl$^-$ and SO_4^{2-} were determined by extracting 0.2±0.01 g of dry
sample with 50 mL of 2% acetic acid using a reciprocal shaker for 30 min, followed by a 0.45 μm PTFE syringe
filtration. Chloride and sulphate were measured using the ferrithiocyante colorimetric method (EPA Method
325.2) and the barium chloride turbidimetric method (EPA Method 375.4), respectively.

3.2   Extraction of metabolites

Metabolites from willow roots, stems and leaves were extracted according to De Vos et al. (2007) with some
modifications. Briefly, plant tissues were freeze-dried for 72 hours and homogenised using a precooled bead
mill Tissuelyser II (Qiagen) at 30 Hz, for 2-4 cycles of 60 seconds each. From the dry tissue powder, 50±0.5 mg
was suspended in 1 mL of methanol:water (75:25%v/v), also containing the internal quantitative standard
resorcinol at 5 mmol L^{-1}. Samples were vortexed for 10 s, sonicated in a 135 W/42 Hz ultrasound bath for 10
min at room temperature and centrifuged at 11,000xg for 2 min at 20°C. The supernatant was filtered through
a 0.2 μm centrifuge filter, diluted by 4 with methanol 75% and placed in the LC-MS/MS autosampler
conditioned at 4°C.

3.3   Liquid chromatography-tandem mass spectrometry (LC-MS/MS) acquisition

Chromatographic separation was performed using an Agilent 1260 Infinity system, where 10 μL of extract was
injected onto a Zorbax Eclipse Plus C18 column (4.6 × 100 mm, 3.5 μm) at 30 °C. Mobile phase was solvent A
(water, 5% methanol, 0.1% formic acid) and solvent B (methanol, 0.1% formic acid) with a flow rate of 0.4 mL
min^{-1} and an 80-minute elution gradient: 100% A hold for 20 min, then a linear increase from 0% to 100% B
over 50 min and 100% B hold for 10 min. This gradient allowed detection of highly water-soluble (polar)
compounds within the first isocratic phase. Untargeted full-scan (100–1,300 Da) MS acquisition was performed
on extracts from all three organs (roots, stems and leaves) of each of the seven replicates (84 samples total),

blanks and quality control samples using an Agilent Q-TOF 6530B mass spectrometer equipped with an Agilent Jet Stream ion source operating in positive ion mode (ESI (+)). Gas temperature was 300 °C, drying gas flow was 5 L min^{-1}, and the nebuliser pressure was 45 psig. Sheath gas temperature was 250 °C with a gas flow of 11 L min^{-1}. Data-dependent MS/MS acquisition was performed on three biological organ replicates per treatment and acquired at collision energy of 20 and 35 eV. Precursor isolation used an intensity threshold set to 500, with a maximum of 4 precursors per cycle and 3 spectra per precursor released after 0.3 min.

3.4 Untargeted metabolite data processing

Raw LC-MS data were converted to mzML files by using msConvert (Chambers et al., 2012) and processed using MZmine 3.3.0 (Schmid et al., 2023) (Fig.1). Background noise cut-off was 1,000 for MS spectra and 0 for MS/MS spectra, with a minimum 5,000 peak intensity threshold for feature detection. Chromatogram deconvolution used the local minimum search algorithm, isotopes were grouped and features alignment across samples was performed with 20 ppm mass tolerance (80% weighting) and 1 min retention time tolerance (20% weighting). Feature quality control filtering applied isotope patterns (minimum two), minimal occurrence of four across all samples, and a maximal blank:sample average ratio (1:3). Features eluting within <0.2 min retention time and with peak shape correlated by >85% (MZmine metaCorrelate algorithm) were assigned to a feature correlation group and used for curation of adducts, complex formation, and in source fragments.

3.5 Metabolites annotation and molecular networking

The feature annotation pipeline used unsupervised and supervised searches across multiple databases integrated with cheminformatic tools (Fig.1). An initial unsupervised annotation step used *in silico* prediction with SIRIUS, combining raw formula and structural elucidation CSI:FingerID (Dührkop et al., 2019), as well as compound class annotation with CANOPUS (Dührkop et al., 2021). In parallel, all features were searched against a set of MS and MS/MS databases, including general databases available in the GNPS environment (including GNPS, MassBank and ReSpect) as well as experiment- and organism-specific databases (PlantCyc (Hawkins et al., 2021), CEU mass mediator (Gil-de-la-Fuente et al., 2019), an in-house library for Salicaceae based on literature as well as authentic standards). Annotation candidate information was standardised for comparison across tools, and the best hit for each feature selected after quality assessment and prioritisation based on integrated confidence/accuracy scores, MS/MS alignment scores, and MS match precision (minimal difference between experimental and predicted compound mass, adduct prediction) to generate a list of putative compounds from combined tools outputs.

In a second supervised step, annotation of significantly differentially abundant compounds was refined using data-driven and knowledge-based curation. Data-driven curation uses feature-based ion identity molecular networking (Z. Zhou et al., 2022) based on cosine and/or Spec2Vec metrics (Huber et al., 2021) of spectral similarity within subnetworks, and manual resolution of ambiguities. Knowledge-driven curation takes advantage of published compound lists and spectra from comparable experiments and plant species not deposited in public databases to improve annotation.

Each putative compound annotation is assigned a confidence level (1-4) (Schymanski et al., 2014; Sumner et al., 2007) and is reported in supporting information – Table S4, following best reporting practices (Alseekh et al., 2021). Chemical taxonomy was assigned to metabolites using the Natural Product Classification system (Kim et al., 2021). Molecular networks were generated in the GNPS analysis environment (Nothias et al., 2020) and visualisation used cosine metrics (at 0.6 threshold) within Cytoscape software (Shannon et al., 2003).

3.6 Statistical analysis

Physiological and metabolomic data were processed using R (R Core Team, 2022) and statistical significance was set at a Benjamin-Hochberg (BH) false discovery rate (FDR) adjusted p-value threshold of <0.05. Physiological comparisons between treatments used a two-way ANOVA, followed by a Tukey HSD *post hoc* test (n=7, except n = 3 for mineral element analysis). Metabolite peak areas were log-transformed and normalised using EigenMS (Karpievitch et al., 2014) prior to statistical analysis (Fig.1). To identify differentially abundant (DA) metabolites, pairwise comparisons of peak area for each treatment (n=7) were performed against controls using either a t-test or Wilcoxon rank sum test with FDR (BH) control, depending on whether assumptions of normality and equal variance were met, or through detection as structural zeros (He et al., 2014) (metabolites detected ≥4 replicates of one condition, and not detected in any replicates of the compared condition). For metabolite comparison across organs, a Kruskal-Wallis test was first performed before *post hoc* pairwise

202 comparisons between each organ pair (Wilcoxon rank sum tests with FDR control). The core metabolome was
203 tentatively defined as compounds that did not significantly vary across control organs and meet a metabolic
204 stability coefficient of variation of peak area below 30%. Principal component analysis (PCA) was performed
205 using MixOmics package (Rohart et al., 2017) and PERMANOVA on Bray-Curtis distances was used to test
206 significant differences between groups.

4   Results

4.1   Physiological effects of salt treatments and mineral elements accumulation

After 28 days of salinity treatments, soil electrical conductivity (EC_e) was 0.9 ± 0.2 dS m^{-1} in controls (Fig.2a) and
was significantly higher in both moderate NaCl and Na_2SO_4 treatments, at 6.5 ± 0.2 and 5.5 ± 0.5 dS m^{-1} (Tukey
HSD, $\alpha<0.05$). High Na_2SO_4 significantly increased soil conductivity further to 9.1 ± 0.5 dS m^{-1}.

Willows maintained a healthy phenotype throughout the experiment (Fig.2b) and salt treatments did not
induce significant differences in aboveground growth rates or organ biomass yields compared to controls
(Supporting information Fig.S1-S3), with total biomass averaging 131 ± 5 g dry matter (DM) (Fig.2c). Quantum
efficiency ΦPSII ranged from 0.44-0.61 and was not significantly different between treatments (Fig.2d). Salt-
treated willows showed significant reductions in stomatal conductance to water vapor (g_{sw}) ranging from 0.13-
0.19 mol m^{-2} s^{-1} (Fig.2e), compared to g_{sw} of 0.29 ± 0.04 mol m^{-2} s^{-1} in control willows. Leaf moisture content
(MC) was 1.97 ± 0.07 g_{H2O} g_{DM}^{-1} in controls and significantly increased to 2.28 ± 0.10 g_{H2O} g_{DM}^{-1} under moderate
NaCl treatment. Moderate Na_2SO_4 resulted in intermediate leaf MC, at 2.09 ± 0.04 g_{H2O} g_{DM}^{-1}, while high Na_2SO_4
treatment significantly decreased compared to both moderate salt treatments, measuring 1.75 ± 0.04 g_{H2O} g_{DM}^{-1}
$^{-1}$ (Fig.2f). None of the treatments induced significant differences in stem moisture content, which was
maintained at an average of 1.34 ± 0.03 g_{H2O} g_{DM}^{-1}.

Total sodium content was 2.78 ± 0.1 mmol(Na) in controls, 9.4 ± 0.3 mmol(Na) in moderate NaCl, 10.1 ± 0.1 in
moderate Na_2SO_4 and 13.0 ± 0.1 mmol(Na) in high Na_2SO_4-treated willows (Fig.3a). Comparison at the organ
scale (Fig.3b) revealed the highest sodium concentration in roots, 3-fold higher than controls in all salt
treatments. Total chloride content was significantly higher in NaCl treatment, with 26.3 ± 2.1 mmol(Cl) as
compared to an average of 6.8 ± 0.2 mmol(Cl) for controls and Na_2SO_4 treatments. In NaCl treatment, leaves
had the highest concentration of Cl at 575.4 ± 26.5 µmol(Cl) g_{DW}^{-1}, representing a 3.4-fold increase compared to
controls; while accumulation in stems and roots was below 190 µmol(Cl) g_{DW}^{-1}, but still 4.3 to 5.5-fold higher
than controls. Total sulphur content was significantly higher in trees of both Na_2SO_4 treatments, at similar
14.1 ± 0.5 and 15.7 ± 1.9 mmol(S), as compared to 7.4 ± 0.4 mmol(S) in controls and 8.5 ± 0.2 mmol(S) in NaCl.
Sulphur concentration significantly increased from 84.3 ± 6.1 µmol(S) g_{DW}^{-1} in control roots to 147.9 ± 6.1 µmol(S)
g_{DW}^{-1} in Na_2SO_4-treated roots and from 161.0 ± 11.1 µmol(S) g_{DW}^{-1} to 229.1 ± 7.5 µmol(S) g_{DW}^{-1} in leaves. At the
whole plant scale, the proportion of inorganic sulphur did not vary significantly across treatments, averaging
27% of the total sulphur present as inorganic SO_4^{2-} (Supporting information – Fig.S5). Total potassium content
was 21.3 ± 0.6 mmol(K) in controls and significantly higher in both moderate and high Na_2SO_4 treatments, with
similar content of 28.9 ± 0.2 and 28.4 ± 2.3 mmol(K) respectively. Potassium concentration tended to decrease
in salt-treated roots but increased in salt-treated leaves compared to controls. Across all four treatments, there
were no significant variations in total phosphorus, calcium and magnesium content (Fig.3a). However, calcium
concentration in leaves significantly decreased in both Na_2SO_4-treated plants compared to controls while leaf
magnesium concentration significantly increased in NaCl-treated plants only, compared to controls (Fig.3b).

4.2   Organ partitioning of the Salix metabolome

After 84 days of growth, untargeted metabolomic profiling of control willows, without salt treatment, revealed
a total of 5,043 metabolic compounds across all three organs, with 2,063 compounds in roots, 2,950
compounds in stems, and 3,601 compounds in leaves (Fig.4a). Shannon and inverse Simpson indices showed a
significant sequential increase in metabolic diversity from roots to stems to leaves (Fig.4b). Principal
Component Analysis (PCA) separated samples based on organ (Fig.4c), PC1 and PC2 explaining 52% and 26% of
variance, respectively, with significant differences between groups (*PERMANOVA*, p-value < 0.05). Leaves and
stems exclusively shared 481 compounds, stems and roots exclusively shared 371 compounds while leaves and
roots exclusively shared 83 compounds (Fig.4d).

A total of 1318 compounds were present in all three organs (Fig.4d), representing 73-96% of cumulative peak
area in each organ (Fig.4e). Differential abundance analysis between organs showed that, of these shared
compounds, 278 did not significantly differ. They were defined as "core metabolites" and included 29
flavonoids, 14 phenolic acids, 12 phenylpropanoids, 22 small peptides, 11 tryptophan alkaloids, 9
pseudoalkaloids, 10 nucleosides, 8 saccharides, 8 monoterpenoids and 4 fatty acids and conjugates (Fig.4f).
High-intensity compounds included adenosine, glutamate, 12-tetradecenoic acid, myrciacitrin III, apigenin 7-
O-neohesperidoside, kaempferol 3-O-glucoside (astragalin), salicortin, salicin, benzaldehyde, 4-o-
caffeoylshikimic acid, secoxyloganin, 7-O-p-coumaroyl loganin, nicotinamide, galactosylglycerol and fructose
(Supporting information - Tables S3,S4).

Conversely, 3,259 compounds were organ-specialised metabolites with significantly increased abundance in
one organ, including 1,719 compounds exclusively detected in leaves, 780 in stems and 291 in roots. Leaf-
specialised compounds were dominated by the terpenoid pathway (Fig.4g), with a high number of (seco)iridoid
monoterpenoids, diterpenoids, steroids, (apo)carotenoids, as well as the alkaloids pathway, particularly
tyrosine and tryptophan derived alkaloids. Stem-specialised compounds were dominated by
phenylpropanoids, lignans and fatty esters whereas root-specialised compounds were dominated by
saccharides and amino acids glycosides. Each organ also had a highly specific set of flavonoids and small
peptides. Interestingly, 19% of root-specialised compounds were highly water soluble (polar), compared to 5%
of stem-specialised and 10% of leaf-specialised compounds (Supporting information – Fig.S6).

4.3   Salix metabolomic response to moderate and high Na_2SO_4

Principle component analysis (PCA) separated samples by control, moderate and high Na_2SO_4 treatments in
roots (PERMANOVA, p-value <0.05), with PC1 explaining 17% and PC2 explaining 11% of the variance, while no
separation was observed between treatments in stems and leaves (Fig.5a). Moderate Na_2SO_4 led to 571
differentially abundant (DA) compounds across the whole plant, including 218 in roots, 105 in stems and 248
in leaves (Fig.5b,d). Comparatively, high Na_2SO_4 led to 976 DA compounds across the whole plant, including
574 in roots, 131 in stems and 271 in leaves (Fig.5c,d).

Taken together, the Na_2SO_4-salt response involved 989 DA compounds distributed across the main pathways
as 260 shikimates/phenylpropanoids, 133 alkaloids, 84 carbohydrates, 79 amino acids/peptides, 73 terpenoids,
51 fatty acids and 25 polyketides (Fig.5e). In response to moderate Na_2SO_4 treatment, DA compounds in roots
and stems were mostly depleted (58-60%), while DA compounds in leaves were mostly enriched (85%)
(Fig.5d,h). Higher Na_2SO_4 induced a larger metabolite enrichment across most pathways, except for further
depletion in carbohydrates, while stems maintained a reduced metabolite pattern and leaves maintained
increased metabolites across the major pathways.

Over the whole plant, 327 compounds were consistently altered in both moderate and high Na_2SO_4 treatments,
representing 21% of DA compounds in roots, 24% in stems and 53% in leaves (Fig.5f). This consistent response
was further reflected in a R^2=0.93 correlation between the fold changes (log_2) moderate and high Na_2SO_4
treatments compared to controls (Fig.5g).

In roots, moderate Na_2SO_4 reduced 45 compounds within shikimates/phenylpropanoids pathway, including 15
flavonoids, 9 phenolic acids and 5 lignans (Fig.5h). Comparatively, high Na_2SO_4 treatment increased 113
compounds from that pathway, including 52 flavonoids, 12 phenolic acids, 12 phenylpropanoids as well as 10
lignans. These included enrichment of important compounds such as 4-coumaroyl shikimate, 5-
hydroxyferulate, procyanidol, catechin 3-O-(1-hydroxy-6-oxo-2-cyclohexene-1-carboxylate), 8-
hydroxypinoresinol 4'-glucoside, guaicylglycerol-coniferyl ether, glucose-conjugated salicylic acid, and salicin.
Six sphingolipids, including phytosphingosine and 4-hydroxysphinganine, were consistently depleted by both
Na_2SO_4 concentrations, while 4 and 10 saccharides significantly reduced under moderate and high Na_2SO_4,
respectively. In contrast, 8 nucleosides and 20 small peptides increased only under high Na_2SO_4, as did 8
pseudoalkaloids, 8 tryptophan alkaloids, 6 tyrosine alkaloids, 12 monoterpenoids, 11 steroids and 9
triterpenoids. In stems, moderate and high Na_2SO_4 treatments enriched 3 and 7 small peptides, such as
aspartate and proline, while 7 and 8 small peptides were depleted, respectively. Additionally, 4 and 14
saccharides were depleted, including glucose and trehalose (Fig.5h), whereas 14 and 13 saccharides were
enriched in leaves. Common, dose-responsive compounds in leaves were dominated by 49
shikimates/phenylpropanoids, encompassing enrichment of 9 cinnamic acid derivatives, 5 salicinoids and 4
other simple phenolic acids, 4 flavonols and 4 chalcons. Commonly regulated metabolites also included 25

alkaloids and 20 amino acids/peptides, with the enrichment of 7 pseudoalkaloids (5 purine and 2 pteridine alkaloids), 3 tryptophan alkaloids and 15 small peptides (Fig.5h).

A total of 179 DA compounds altered by one or both concentrations of Na_2SO_4 (12% of all Na_2SO_4 metabolomic changes) contained sulphur, corresponding to 101 distinct compounds. Roots significantly accumulated 13 and 45 sulphur compounds (S-compounds) under moderate and high Na_2SO_4 respectively, but significantly reduced 4 and 20 S-compounds (Fig.5i). Additionally in leaves, 36 S-compounds were increased under one or both Na_2SO_4 concentrations, compared to 11 depleted S-compounds. Shared enriched S-compounds included dicarboxylic acid such as 2-(2-sulfoethyl)propanedioic acid or polyamines, such as N'-hydroxy-5-(2-methoxyethyl-sulfamoylamino) pentanimidamide, as well as stilbenes, such as resveratrol 4-sulfate. High Na_2SO_4 depleted S-containing compounds ranged from small peptides, such as threonine to larger alkaloids such as $C_{19}H_{16}N_8O_5S$ (CHEMBL164867). Notably, a subgroup of 16 DA S-compounds were in the same structural cluster (Fig.5j).

4.4   Salix metabolomic response to NaCl treatment

Comparing metabolomes from controls with NaCl-treated willows, PCA revealed a significant separation between control and NaCl treatment in roots (*PERMANOVA*, p-value <0.05), partial overlap in stems (*PERMANOVA*, p-value < 0.05) and no observed separation in leaves (Fig.6a). NaCl treatment led to DA in 1,063 DA compounds, with 475 in roots, 481 in stems and 107 in leaves (Fig.6b). Among these, 768 DA compounds were unique to one organ, while 135 were shared to two or all organs (Fig.6c). Notably, 98 DA compounds were exclusively shared between roots and stems and 24 were significantly depleted in all three organs due to NaCl treatment.

Across all organs, NaCl-responsive metabolites comprised 213 DA compounds within the shikimates/phenylpropanoids pathway, 118 alkaloids, 88 carbohydrates, 74 amino acids/peptides, 59 terpenoids, 48 fatty acids and 12 polyketides (Fig.6d). While shikimates/phenylpropanoid pathway compounds were the most diverse, carbohydrates represented the highest proportional change within a pathway, constituting >33% of total carbohydrates (Supporting information – Fig.S7). This was driven by significant depletion in 28 carbohydrates across multiple organs, including glucose, fructose, sucrose and valienol 1,7-diphosphate, as well as organ-specific depletion of 36 carbohydrates (Fig.6e,f). Tricarboxylic acid cycle intermediates, such as malate, citrate, 2-oxoglutarate, were also depleted across the entire plant as well as phosphorylated compounds, such as phosphocholine, phosphonoacetaldehyde, 1,7-diphospho-1-epi-valienol and 2-C-methyl-D-erythritol-2,4-cyclodiphosphate (Supporting information – Fig.S8).

Within organs, leaves had a limited response to NaCl treatment (Fig.6b,c), but depleted compounds included 2-oxoglutarate, malate, aspartate and choline, while accumulated compounds included populoside C, (indol-3-yl)acetyl-myo-inositol L-arabinoside (IAA derivative) and spermidine (Supporting information – Tables S3,S7). Stems and roots had a larger metabolic response, markedly within the shikimate/phenylpropanoid pathway, altering 115 and 108 compounds respectively (Fig.6e). This large group of differentially abundant phenolic compounds was primarily associated to flavonoids, phenylpropanoids, lignans and phenolic acids (including salicinoids). While enriched flavonoids number were similar in stems and roots, 20 and 23 DA compounds respectively, stems had more depleted flavonoids than roots, with 23 and 10 DA compounds respectively (Fig.6h). Salicinoids were net enriched in both stems and roots (Fig.6h). Stems had significant enrichment of myricetin 3-O-glucoside, kaempferide 3-O-glucoside, procyanidol, salicin and grandidentatin, alongside the depletion of eriodictyol 7-glucuronide, luteolin 7-(6''-benzoylglucoside, ramontoside and a hydroxycyclohexen-oyl ester catechin derivative. In roots, populin, tremuloidin and benzoyl-β-D-glucoside were enriched, while (iso-)salicin, salicylic acid glucoside, grandidentatin, catechin, procyandin C, pleurostimin 7-glucoside were depleted. Additionally, 21 lignans and phenylpropanoids significantly accumulated in roots, contrasting stems, where 14 lignans and phenylpropanoids were depleted (Fig.6h).

A balanced pattern of metabolomic change was observed in nitrogen metabolism in response to NaCl treatment, with enrichment of 147 DA compounds and depletion of 133 DA compounds. This was particularly evident as 31 small peptides increased across all organs, including proline, tryptophan and glutathione, while 34 small peptides decreased, including acetylproline, threonine and arogenate (Fig.6g). Other increasing N-containing compounds included purine nucleosides like adenosine, alkaloids such as dihydrosanguinarine and kinetin-7-N-glucoside, as well as sphingolipids, such as phytosphingosine and sphinganine. Concurrently,

terpenoids were enriched, especially monoterpenoids, with 15 enriched monoterpenoids in stems and 13 in roots (Fig.6i,j).

4.5 *Salix anion-specific metabolomics response to NaCl and Na₂SO₄ (at the same EC)*

Comparing the 1,063 and 571 DA compounds altered by moderate NaCl and moderate Na_2SO_4 revealed 103 DA compounds commonly regulated in roots, 53 in stems and 7 in leaves (Fig.7a,b). These generalised salinity responsive metabolites included 23 shikimates/phenylpropanoids, 10 alkaloids, 10 carbohydrates, 9 terpenoids, 9 amino acids and peptides and 5 fatty acids (11 other and 70 unknown) (Fig.7b). Commonly enriched compounds included (indol-3-yl)acetyl-myo-inositol L-arabinoside and 2-phenylacetamide in leaves, proline, grandidentatin and fraxiresinol 1-O-glucoside (a lignan) in stems, 2-thioethylidenebis(phosphonic acid), delphinidin 3-O-(6''-O-malonyl)-beta-D-glucoside-3',5'-di-O-beta-D-glucosideand and syringin in roots. Commonly depleted compounds included glucose, trehalose in stems, D-altro-heptulose, glycerophophoglycerol, benzoyl-beta-D-glucoside and N-(4-(1-cyanocyclopentyl)phenyl)-2-((pyridin-4-ylmethyl)amino)nicotinamide in roots, as well as d-glycero-l-galacto-octulose in both stems and roots.

Anion specific responses included 764 DA compounds unique to moderate NaCl treatment and 391 DA compounds unique to the moderate Na_2SO_4. Both salt treatments induced the accumulation of salt-specific monoterpenoids, as well as phenolic compounds, including flavonoids, phenylpropanoids, and phenolic acids. Moreover, NaCl treatment led to a net decrease of saccharides and nucleosides, whereas Na_2SO_4 treatment resulted in their net accumulation (Fig.7c). An additional contrasting effect was observed with sphingolipids, with increased in NaCl but decreased Na_2SO_4 treated trees.

An anion-specific effect could also be observed when comparing change in constitutively produced compounds that were core (present and relatively stable between organs) and organ-specialised (significantly different between organs), but also in salt-induced compounds (not detected in controls). In both moderate salt treatments, roots primarily reduced root-specialised compounds rather than core compounds, while significantly enriched compounds were predominantly newly synthesised (salt-induced), rather than root-specialised or core compounds (Fig.7d). Similarly, in stems, reduced compounds were largely stem-specialised rather than core compounds under both salt types, however enriched compounds were predominantly stem-specialised under NaCl, comprising 53 enriched DA compounds as compared to 20 salt-induced and 23 core enriched DA compounds. Conversely, Na_2SO_4 treatment resulted in 2 stem-specialised, 5 salt-induced and 3 core enriched DA compounds. In leaves, NaCl treatment reduced 13 leaf-specialised and 4 core DA compounds while enriching 26 leaf-specialised, 20 salt-induced and 4 core DA compounds. In contrast, leaves treated with Na_2SO_4 decreased 22 leaf-specialised and 4 core compounds while increasing 5 leaf-specialised and 63 core compounds.

5 Discussion

5.1 *Salix salinity tolerance response*

Soil salinity levels are commonly categorised based on EC_e as slightly saline (2-4 dS m^{-1}), moderately saline (4-8 dS m^{-1}), and highly saline (>8 dS m^{-1}) (Corwin and Scudiero, 2019; FAO, 1988). Here, the comparison of non-saline control, moderately saline NaCl and Na_2SO_4, and highly saline Na_2SO_4 (Fig.2a) enabled capture of the anion-specific effects (Cl$^-$ and SO$_4$$^{2-}$) and severity effects (moderate vs high Na_2SO_4) on willow (*Salix miyabeana* 'SX64').

Photosynthetic capacity and biomass productivity were maintained compared to water controls, even at high soil salinity levels of 9 dS m^{-1} (Fig.2c,d), supporting the suitability of willow for phytoremediation. This aligns with research by Hangs et al. (2011), reporting salinity tolerance in 37 willow varieties, where most cultivars could tolerate moderate soil salinity and the most salt-tolerant cultivars maintained yield under high soil salinity. Leaf moisture content increased under moderate salinity, alongside a >35% reduction of stomatal conductance (Fig.2e). This indicates reduced transpiration accompanied by osmotic adjustment to sustain turgor, potentially mitigating ion toxicity through absorbed salt dilution (Munns et al., 2020b; Nguyen et al., 2017). However, subsequent lower leaf moisture content suggests this water balance might not be maintained as salinity severity increase (Fig.2d).

Salt treatments did not reduce the acquisition of essential nutrients and even increased potassium uptake alongside salt ions, although these ions were selectively partitioned (Fig.3). Sodium accumulation was predominantly restricted to the roots and influx did not increase linearly with external concentration, suggesting a robust sodium exclusion mechanism (Fig.3a). Such exclusion can occur through root cell wall Na^+ binding (Byrt et al., 2018) or competing uptake of K^+ (Hauser and Horie, 2010), as proposed in the salt tolerant *Salix interior* (Major et al., 2017) and *Salix eriocephala* 'LAR' (Huang et al., 2023). In contrast, the high chloride uptake and transport throughout the plant (Fig.3) can be facilitated by passive anion channels (Skerrett and Tyerman, 1994), resulting in leaf chloride concentrations here reaching the upper optimal level for non-halophytes (~30-560 µmol(Cl) g_{DW}^{-1} (Geilfus, 2018)). This accumulation, coupled with increased leaf water content, suggests that Cl^- may serve as a readily available osmolyte, potentially conferring osmotic benefits similar to those observed in *Nicotiana tabacum* (Franco-Navarro et al., 2016). However, given that cytosolic chloride toxicity is thought to occur at concentrations of 5-20 mmol(Cl) $L(cytosol)^{-1}$ (Teakle and Tyerman, 2010), the accumulated Cl^- may also require compartmentalisation into vacuoles, apoplasts, or older leaves to mitigate cellular damage and optimise metabolic burden. In comparison to chloride, sulphate uptake was more constrained but still resulted in increased transport throughout the plant, where it was either stored as SO_4^{2-} or metabolised into organic sulphur compounds.

The maintenance of biomass yields despite reduced stomatal conductance, which can lead to decreased carbon assimilation (Munns et al., 2020a), suggests effective metabolic plasticity and ion management strategies, making this experimental system suitable for assessing willow metabolomic tolerance mechanisms without the confounding metabolic effects associated with severe salt-induced physiological damage.

5.2   High organ-specific diversity in the Salix metabolome

Untargeted metabolomic analysis of control trees revealed substantial diversity across the three main vegetative organs (roots, stems and leaves), with over 5,000 compounds detected, approximately 62% of which could be annotated at confidence levels 1-3 (Sumner et al., 2007), providing a detailed overview of the willow metabolome at whole plant scale (Fig.4). Previous metabolomic studies in single organs of *Salix* have captured up to 1,639 metabolites in bark (J. Zhou et al., 2022), 2,735 (Kaling et al., 2015), 600 (Jia et al., 2020) and 2,440 (Aliferis et al., 2015) metabolites in leaves, and 800 metabolites in roots (Xia et al., 2021). By integrating analyses of multiple plant organs, a group of common metabolites across organs were identified, representing 26% of metabolomic diversity and over 70% of the putative abundance (estimated from peak area) (Fig.4d). Of these, 287 metabolites had stable abundance across organs and represent its potential plant-wide basal, or "core", metabolism (Fig.4f); a recent metabolomic concept that can also denote evolutionarily conserved metabolites across species (Drapal et al., 2022; Dussarrat et al., 2022; McLaughlin et al., 2023). Many of these core metabolites were associated with central pathways like glycolysis, TCA cycle derivatives, and amino acid metabolism, reflecting primary metabolism across plant tissues. Others included bioactive phenolics found at high levels in Salicaceae, such as characteristic salicylates (e.g. salicin, salicortin) associated to antioxidant, antimicrobial and anti-herbivory functions in willow (Boeckler et al., 2011; Brereton et al., 2017; Meier et al., 1988; Pobłocka-Olech et al., 2007; Tyśkiewicz et al., 2019; Volf et al., 2015).

Beyond these common metabolites, willow organs had clear metabolic differences (Fig.4g). Leaves contained the most diverse metabolome, aligning with leaf function as a source of complex photosynthates. The leaf metabolome included diverse terpenoids involved in light harvesting, development, defence, and signalling (Pichersky and Raguso, 2018). Stems accumulated lignans and phenylpropanoids, reflecting woody tissue functions of structure and storage (Boeckler et al., 2011; Köhler et al., 2020). Additionally, the metabolome of stems had a large overlap with both leaves and roots, underscoring their function as conduits for biochemical transport and dialog between aerial and belowground processes (Fig.4d). Comparatively, despite critical functions of water and nutrient absorption, exudation, anchoring structure and storage, roots had the lowest metabolic diversity (Fig.4a,b). Roots were enriched in saccharides and flavonoids, thought to mediate soil interactions like microbial symbiosis (Endo et al., 2021), metal tolerance (Osawa et al., 2011), nutrient uptake (Cesco et al., 2012) and root architecture (Buer et al., 2010). The root metabolome was also distinctly richer in polar metabolites (Supporting information – Fig.S6) which can more readily diffuse through the soil environment within the water vector and could therefore play a role as exudates involved in shaping the rhizospheric biotic (Sasse et al., 2018) and abiotic (Badri and Vivanco, 2009) environment.

453 These distinct metabolomes reflect the different physiological roles of each organ, describing different
454 metabolic activity and degree of functional specialisation (Li et al., 2016) as well as serving as a baseline to
455 enable understanding of key responses to stimuli such as salt stress.

5.3 Sulphate salt induces dose-dependent metabolomic responses in Salix

Sulphate salinity induced extensive organ-specific metabolic reprogramming (Fig.5a-d,h), affecting 10% and
16% of all metabolites at moderate and high concentrations, respectively. Doubling the Na_2SO_4 concentration
in soil resulted in a twofold increase of metabolic changes in roots, but more limited responses in stems and
leaves. This localisation of metabolic response to the roots, which serve as the primary defence barrier against
excess ions and osmotic stress, is likely decisive in mitigating salinity stress impact aboveground. Notably,
consistent levels of salt ion uptake and accumulation despite increased soil salinity indicate effective ion
management (Fig.3), allowing leaves to maintain similar adaptive metabolomic strategies under both
sulphate salt treatments (Fig.5f,g). Previous studies reported sulphate salt tolerance in willows (Huang et al.,
2020) associated with ion imbalance restriction to roots, similarly preventing physiological impact on leaves at
moderate salinity levels.

In roots, Na_2SO_4 dose differentially effected shikimates/phenylpropanoids, with a reduction of flavonoids,
phenolic acids and lignans under moderate level, contrasting their net increase under higher sulphate salinity
(Fig.5h). This may indicate a developmental slowdown under mild stress, superseded by a required metabolic
burst under more severe salinity levels. The lower osmotic potential associated with higher salinity likely drove
the accumulation of lignans (Fig.5h) to enhance structural protection (as lignin precursors) and cavitation
protection (Reyt et al. 2021), while accumulation of antioxidative flavonoids can enhance secondary oxidative
stress tolerance (Nakabayashi et al., 2014). Under high Na_2SO_4, the sugar donor UDP-D-galactose significantly
increased alongside phenolics, indicating a disruption in glycosylation and flavonoid bioactivity regulation (Xiao
et al., 2014), while increases in iridoid monoterpenoids, such as lamiide, can prevent lipid peroxidation to
protect cell membrane integrity (Delaporte et al., 2002). The high diversity of salt-responsive metabolites
captured here implies additional functions beyond ROS scavenging, likely contributing to the observed Na_2SO_4
stress tolerance. Moreover, the reduction in sphingolipids, which regulate ions through early sensing and
signalling, could limit salt uptake below toxic thresholds (Jiang et al., 2019; Liu et al., 2022), coherent with
depletion of the essential precursor octanoyl-CoA (Neess et al., 2015; Sonar et al., 2020).

Primary metabolites declined in stems under both sulphur salt concentrations, including reductions in small
peptides, saccharides, and nucleosides (Fig.5h). A slowdown of developmental processes and primary
metabolism would be expected under salt stress, although there was no accompanying (short-term)
impairment to total biomass production (Fig.2c). The comparatively limited disruption of secondary
metabolites in stems indicates maintenance of homeostasis and primary resource allocation to support root
functions, effectively restricting salt stress propagation.

In leaves, the salt-responsive metabolites were broadly shared between both Na_2SO_4 levels, characterised by
dose-dependent accumulation, including saccharides, small peptides and precursors of phenolic compounds
(Fig.5f,g). Again, illustrating the buffering role of roots in protecting essential carbon fixation functions and
energy-related compound production, even under severe salt exposure.

An increase of sulphur-containing compound synthesis under SO_4^{2-} treatment and upregulation of sulphur
assimilation and transport pathways is a characterised plant adaptation mechanism to effectively utilise excess
SO_4^{2-} (Buchner et al., 2004; Davidian and Kopriva, 2010). Here, a substantial enrichment of small peptides
occurred in roots under sulphur salinity (Fig.5h), with around half incorporating sulphur. This accumulation of
sulphur-containing compounds was consistent throughout the plant at both concentrations (Fig.5i) and was
predominantly cysteine and glutathione-based small peptides and some fatty acyl thioesters and alkaloids. The
integration of sulphate ions into organic sulphur-containing compounds has the potential to reduce ion
imbalance *in planta* during salt stress and could represent a tolerance mechanism in willow. Moreover, the
increase of structurally related compounds here (Fig.5i), although their functional information is limited,
suggests activation of a bespoke, yet-to-be-described, metabolic response to Na_2SO_4. Watanabe et al. (2019)
highlighted a need for integrated sulphur-omics to overcome this shortfall in functional understanding of
sulphur-containing molecules in plants, of particular relevance to salinity tolerant non-model organisms such
as willow.

5.4   Salt anions shape Salix metabolomic tolerance responses

The salt response across all treatments involved an extensive proportion of the metabolome (28% of
metabolites), reflecting the highly plastic nature of plant metabolism under environmental challenge. While
activation of the calcium sensing signalling pathway by Na^+ (Munns and Tester, 2008) and disruption of the
Na^+/K^+ ratio (Zhang et al., 2018) are commonly considered to trigger salt responses in plants, anion driven
metabolomic responses have been less well explored. Comparison of metabolomic responses to moderate
NaCl and Na_2SO_4 treatments here, at equivalent soil electroconductivity and sodium concentrations, revealed
only 3% of metabolites were consistently responsive, highlighting extensive anion-specific changes (Fig.7).

The common, or generalised, salt metabolite changes were asymmetrically distributed across organs, with ~15-
fold more generalised salt responses in roots than in leaves (Fig.7b). These represent common metabolic
strategies to tolerate (anion agnostic) osmotic stress in roots at the root-soil interface and restrict sodium
translocation, while leaves might be more susceptible to the (anion dependent) ion toxicity of translocated
Cl^- and SO_4^{2-}. The generalised salt response included common accumulation of the phenylpropanoid precursor
2-phenylacetamide in leaves (a salt biomarker in the euhalophyte *Suaeda salsa* (Bao et al., 2023)), protective
molecules in stems such as the anthocyanin ternatin C5 (Kovinich et al., 2015) and a highly antioxidative
coumaric acid conjugate of salicin, grandidentatin (Si et al., 2009), as well as potential cell wall structural
components in roots, such as the monolignol precursors syringin.

NaCl treatment, unlike Na_2SO_4, induced a comparatively small metabolomic response in leaves (Fig.6a-c). Of
the limited responses, a significant accumulation of spermidine, a polyamine known for its acid- and ROS-
neutralising properties, could help protect cell membranes, reduce chlorophyll loss and leaf senescence (Gill
and Tuteja, 2010b). The auxin conjugate, (indol-3-ylacetyl)-myo-inositol L-arabinoside, also increased,
suggesting oxidative stress specific developmental regulation (Bajguz and Piotrowska, 2009; Ludwig-Müller,
2011). A larger change under NaCl salt exposure was the reconfiguration of carbon resources through the
shikimate/phenylpropanoids pathway (Fig.6d,h). Specifically, lignans and phenylpropanoids had organ-specific
responses: substantial accumulation in roots and significant depletion in stems (Fig.6h). As major lignin
precursors, these lignan and phenylpropanoid dynamics could represent the expected structural adaptations
within the roots and stems. In particular, lignin deposition can act as a barrier regulating ion uptake and
translocation to aerial parts (Byrt et al., 2018; Neves et al., 2010; Reyt et al., 2021).

Flavonoids, whose antioxidative capacity can facilitate localised neutralisation of ROS (Nakabayashi et al.,
2014), also had marked organ-specific responses, with substantial accumulation in roots and variable
regulation in the stems (Fig.6h). As flavonoids, phenolic acids, lignans and phenylpropanoids all derive from
the same substrates (phenylalanine and the pivotal intermediate cinnamic acid), the organ-specific shift
suggests a coordinated metabolic response promoting defence in the directly exposed roots while
simultaneously restricting the availability and energy expense of these compounds in the stems. By
comparison, moderate Na_2SO_4 did not induce similar extensive accumulation of phenolic compounds in roots
or stems but this response was triggered under higher Na_2SO_4 concentration.

Accumulation of osmoprotectants under salt stress is one of the most prevalent and well-described adaptive
strategies to reduce cell osmotic potential and stabilise cellular components (D'Amelia et al., 2018; Yang and
Guo, 2018). Surprisingly, except for stem proline enrichment under both NaCl and Na_2SO_4, the levels of
expected organic osmoprotectants, namely hydrophilic and low molecular weight sugars, polyamines or amino
acids, were largely reduced in roots and stems under NaCl while accumulating in leaves under Na_2SO_4, more
distal to the primary site of osmotic stress (Fig.5h, Fig.6f,g and Fig.7c). This response contrasts salt-tolerant
poplars which can accumulate soluble sugars as a tolerance mechanism (Jouve et al., 2004; Watanabe et al.,
2000) and might be a distinct strategy used by *S. miyabeana* to mitigate osmotic stress through the uptake of
inorganic ions, including root restricted Na^+ and anions (Cl^- and SO_4^{2-}) translocation to leaves. Similarly, Ottow
et al. (2005) reported preferential sodium uptake by *Populus euphratica*, where Na^+ served as a readily-
available osmoprotectant stored in the apoplast or bound in the cell wall, alongside reduced calcium and
soluble carbohydrate levels.

The anion-dependent regulation pattern of osmoprotectants also reflected broader differing metabolic
adjustments within central metabolism. Although reduction of stomatal conductance was similar between
both salts and photosystem II efficiency (Φ_{PSII}) preserved (Fig.2), the impact on foliar glucose and energy-
related compounds was distinct. NaCl induced the plant-wide reduction of glucose and multiple TCA cycle

intermediates (Supporting information – Fig.S8), representing impaired photosynthetic carbon fixation and
decreased flux through glycolysis and the TCA cycle, limiting mitochondrial respiration and overall energy
supply. A similar reconfiguration of resources and metabolic processes has been observed under salt stress in
barley roots (Wu et al., 2013), where glycolysis was inhibited, and in *Arabidopsis thalia* (Hartmann et al., 2015),
where a shift to non-oxidative fermentative energy metabolism can occur. This was coupled with reduced
levels of small peptides, nucleosides and other simple carbohydrates like fructose and sucrose in stems and
roots (Fig.6f,g and Fig.7c), possibly driven by increased carbon demand for secondary metabolism involved in
salt stress tolerance. By comparison, Na_2SO_4 induced an increase of glucose in leaves (Supporting information
– Fig.S8), alongside an accumulation of saccharides, small peptides and nucleosides, consistent even under
high Na_2SO_4. This suggests that willow can sustain chemical energy production in leaves in the presence of
sulphate, unlike with chloride, and plant-wide production of salt-tolerance supporting metabolites,
underscoring the importance of understanding anion toxicity in salt tolerance and implicating organic
compound sulphate incorporation in prevention of ionic imbalance. As an example, the accumulation of
sulphur-containing metabolites (Fig.5i,j), including 2,3-di-O-sulfopropylglucose, aligns with the known
mechanism of sugar sulphonation that promotes salt tolerance through ion regulation, osmotic tolerance, and
cell wall integrity in algae (Aquino et al., 2011).

Both chloride and sulphate salts activated the terpenoid pathway (Supporting information – Fig.S9), known to
mitigate salt stress in mangroves (Basyuni et al., 2009). NaCl induced significant accumulation of terpenoids
from the chloroplastic methylerythritol phosphate (MEP) pathway across the whole plant, including
diterpenoids as well as iridoid and secoiridoid monoterpenoids, reflecting transcriptomic changes observed in
salt-tolerant poplar (Chen et al., 2018). Na_2SO_4 treatment similarly enriched terpenoids through the MEP
pathway, but also activated the cytoplasmic mevalonate pathway in roots, also linked to salt tolerance in poplar
(Wei et al., 2020). This sulphate-salt-associated activation of the mevalonate pathway increased with higher
Na_2SO_4 levels, including enrichment of the hormone precursor abscisic aldehyde in roots and leaves. This
suggests an adaptive stress response where willows are positioned to quickly synthesise ABA, signalling
stomatal closure, root architecture changes, and osmoprotectant accumulation (Ryu and Cho, 2015), with
ABA-aldehyde oxidase mediated regulation. In contrast, ABA precursor enrichment was not observed with
NaCl, indicating chloride ion interference with signalling that may leave willows more vulnerable to salt stress.

Overall, willow responded to salt stress by modulating both core and organ-specialised metabolites (Fig.4 and
Fig.7d). Within each organ, there was a tendency for greater depletion of organ-specialised compounds, over
core metabolites, indicating higher resilience of core metabolism under salt stress (Fig.7d). Contrasting this,
Na_2SO_4 treatment resulted in the enrichment of core metabolites in leaves, while NaCl enriched organ-
specialised compounds. This anion-specific difference further highlights the nuanced metabolic mechanisms
deployed to face distinct salt stress conditions.

6 Conclusion

Untargeted metabolomics uncovered extremely high diversity of metabolites in willow, the majority of which
represent organ-specialised metabolites aligned with distinct physiological functions, but which also include
over 250 putative plant-wide stable, or core, metabolites. The leaves had the highest metabolite diversity, but
when subjected to different saline soil conditions, roots, as the frontline of exposure, had the most extensive
biochemical response, mitigating ion toxicity and osmotic pressure changes to enable plant tolerance.
Accompanying this was plant-wide reallocation of energy and metabolic resources, coupled with prioritisation
of antioxidant flavonoids, able to protect vulnerable leaf photosynthetic functions, constituting the signature
of a general willow salt response. However, the largely specialised nature of this response revealed that *in
planta*, salt anion type becomes an increasingly stronger discriminating force of metabolomic profile when
moving from roots to leaves. Indeed, the metabolic adjustment differed extensively in leaves, where Na_2SO_4
induces key components of the core metabolome to mitigate complex osmotic and toxic stress even as soil
salinity increases, whereas NaCl induces more organ-specialised metabolic tools. Salinity levels in agriculture
are currently categorised by soil EC, however, given the substantial anion-dependent differences observed in
plant metabolic responses, soil EC measurement might be insufficient to reflect the diversity of soil salinity
types, their actual osmotic potential, as well as the anion-specific plant metabolic toolkit necessary for plant
salt tolerance.

7   Declaration of Competing Interest

The authors declare that there is no conflict of interest.

8   Acknowledgements

We gratefully acknowledge the financial support provided from NSERC Discovery Grants (RGPIN-2017-05452 and RGPIN-2023-04863), Natural Resources Canada Forest Innovation Program Grant (CWFC1718-018 and CWFC1920-104), ECCC Environmental Damage Fund (EDF-PQ-2020b012), MITACS (IT23193), and the UCD Ad Astra Award Program. A special thank you is extended to Ariane Lafrenière, Marc-Olivier Brunette and Bruno Guerrier for their kind and effective support during harvest, and to Allan Harms from NRAL, University of Alberta, for his valuable expertise and sympathy.

9   Data availability statement

The data that supports the findings of this study are available in the supporting information of this article and on GNPS-MassIVE online dataset repository.

10   References

Abdelgadir, E.M., Oka, M., Fujiyama, H., 2005. Characteristics of nitrate uptake by plants under salinity. J. Plant Nutr. 28, 33–46. https://doi.org/10.1081/PLN-200042156

Abreu, M., Silva, L., Ribeiro, B., Ferreira, A., Alves, L., Paixão, S.M., Gouveia, L., Moura, P., Carvalheiro, F., Duarte, L.C., Fernando, A.L., Reis, A., Gírio, F., 2022. Low indirect land use change (ILUC) energy crops to bioenergy and biofuels: A review. Energies 15, 4348. https://doi.org/10.3390/en15124348

Aliferis, K.A., Chamoun, R., Jabaji, S., 2015. Metabolic responses of willow (*Salix purpurea* L.) leaves to mycorrhization as revealed by mass spectrometry and 1H NMR spectroscopy metabolite profiling. Front. Plant Sci. 6.

Alseekh, S., Aharoni, A., Brotman, Y., Contrepois, K., D'Auria, J., Ewald, J., C. Ewald, J., Fraser, P.D., Giavalisco, P., Hall, R.D., Heinemann, M., Link, H., Luo, J., Neumann, S., Nielsen, J., Perez de Souza, L., Saito, K., Sauer, U., Schroeder, F.C., Schuster, S., Siuzdak, G., Skirycz, A., Sumner, L.W., Snyder, M.P., Tang, H., Tohge, T., Wang, Y., Wen, W., Wu, S., Xu, G., Zamboni, N., Fernie, A.R., 2021. Mass spectrometry-based metabolomics: A guide for annotation, quantification and best reporting practices. Nat. Methods 18, 747–756. https://doi.org/10.1038/s41592-021-01197-1

Aquino, R.S., Grativol, C., Mourão, P.A.S., 2011. Rising from the sea: Correlations between sulfated polysaccharides and salinity in plants. PLOS ONE 6, e18862. https://doi.org/10.1371/journal.pone.0018862

Arif, Y., Singh, P., Siddiqui, H., Bajguz, A., Hayat, S., 2020. Salinity induced physiological and biochemical changes in plants: An omic approach towards salt stress tolerance. Plant Physiol. Biochem. 156, 64–77. https://doi.org/10.1016/j.plaphy.2020.08.042

Badri, D.V., Vivanco, J.M., 2009. Regulation and function of root exudates. Plant Cell Environ. 32, 666–681. https://doi.org/10.1111/j.1365-3040.2009.01926.x

Bajguz, A., Piotrowska, A., 2009. Conjugates of auxin and cytokinin. Phytochemistry 70, 957–969. https://doi.org/10.1016/j.phytochem.2009.05.006

Bao, J., Liu, Z., Ding, Z., Yisilam, G., Wang, Q., Tian, X., Bao, J., Liu, Z., Ding, Z., Yisilam, G., Wang, Q., Tian, X., 2023. Metabolomic analysis reveals key metabolites and metabolic pathways in *Suaeda salsa* under salt and drought stress. Funct. Plant Biol. https://doi.org/10.1071/FP23049

Basyuni, M., Baba, S., Inafuku, M., Iwasaki, H., Kinjo, K., Oku, H., 2009. Expression of terpenoid synthase mRNA and terpenoid content in salt stressed mangrove. J. Plant Physiol. 166, 1786–1800. https://doi.org/10.1016/j.jplph.2009.05.008

Bilek, M.A., Soolanayakanahally, R.Y., Guy, R.D., Mansfield, S.D., 2020. Physiological response of *Populus balsamifera* and *Salix eriocephala* to salinity and hydraulic fracturing wastewater: potential for phytoremediation applications. Int. J. Environ. Res. Public. Health 17, 7641. https://doi.org/10.3390/ijerph17207641

Boeckler, G.A., Gershenzon, J., Unsicker, S.B., 2011. Phenolic glycosides of the Salicaceae and their role as anti-herbivore defenses. Phytochemistry, Plant-Insect Interactions 72, 1497–1509. https://doi.org/10.1016/j.phytochem.2011.01.038

Brereton, N.J.B., Berthod, N., Lafleur, B., Pedneault, K., Pitre, F.E., Labrecque, M., 2017. Extractable phenolic yield variation in five cultivars of mature short rotation coppice willow from four plantations in Quebec. Ind. Crops Prod. 97, 525–535. https://doi.org/10.1016/j.indcrop.2016.12.049

Buchner, P., Takahashi, H., Hawkesford, M.J., 2004. Plant sulphate transporters: Co-ordination of uptake, intracellular and long-distance transport. J. Exp. Bot. 55, 1765–1773. https://doi.org/10.1093/jxb/erh206

Buer, C.S., Imin, N., Djordjevic, M.A., 2010. Flavonoids: New roles for old molecules. J. Integr. Plant Biol. 52, 98–111. https://doi.org/10.1111/j.1744-7909.2010.00905.x

Byrt, C.S., Munns, R., Burton, R.A., Gilliham, M., Wege, S., 2018. Root cell wall solutions for crop plants in saline soils. Plant Sci. 269, 47–55. https://doi.org/10.1016/j.plantsci.2017.12.012

Cesco, S., Mimmo, T., Tonon, G., Tomasi, N., Pinton, R., Terzano, R., Neumann, G., Weisskopf, L., Renella, G., Landi, L., Nannipieri, P., 2012. Plant-borne flavonoids released into the rhizosphere: impact on soil bio-activities related to plant nutrition. A review. Biol. Fertil. Soils 48, 123–149. https://doi.org/10.1007/s00374-011-0653-2

Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto, L., Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen, B., Baker, T.A., Brusniak, M.-Y., Paulse, C., Creasy, D., Flashner, L., Kani, K., Moulding, C., Seymour, S.L., Nuwaysir, L.M., Lefebvre, B., Kuhlmann, F., Roark, J., Rainer, P., Detlev, S., Hemenway, T., Huhmer, A., Langridge, J., Connolly, B., Chadick, T., Holly, K., Eckels, J., Deutsch, E.W., Moritz, R.L., Katz, J.E., Agus, D.B., MacCoss, M., Tabb, D.L., Mallick, P., 2012. A cross-platform toolkit for mass spectrometry and proteomics. Nat. Biotechnol. 30, 918–920. https://doi.org/10.1038/nbt.2377

Chan, K.X., Phua, S.Y., Van Breusegem, F., 2019. Secondary sulfur metabolism in cellular signalling and oxidative stress responses. J. Exp. Bot. 70, 4237–4250. https://doi.org/10.1093/jxb/erz119

Chen, P., Zuo, L., Yu, X., Dong, Y., Zhang, S., Yang, M., 2018. Response mechanism in *Populus × euramericana* cv. '74/76' revealed by RNA-seq under salt stress. Acta Physiol. Plant. 40, 96. https://doi.org/10.1007/s11738-018-2676-x

Chen, S., Polle, A., 2010. Salinity tolerance of *Populus*. Plant Biol. 12, 317–333. https://doi.org/10.1111/j.1438-8677.2009.00301.x

Corwin, D.L., Scudiero, E., 2019. Review of soil salinity assessment for agriculture across multiple scales using proximal and/or remote sensors, in: Sparks, D.L., Sparks, D.L. (Eds.), Advances in Agronomy. pp. 1–130.

D'Amelia, L., Dell'Aversana, E., Woodrow, P., Ciarmiello, L.F., Carillo, P., 2018. Metabolomics for crop improvement against salinity stress, in: Kumar, V., Wani, S.H., Suprasanna, P., Tran, L.-S.P., Kumar, V., Wani, S.H., Suprasanna, P., Tran, L.-S.P. (Eds.), Salinity Responses and Tolerance in Plants, Volume 2: Exploring RNAi, Genome Editing and Systems Biology. Cham, pp. 267–287.

Davidian, J.-C., Kopriva, S., 2010. Regulation of sulfate uptake and assimilation—the same or not the same? Mol. Plant 3, 314–325. https://doi.org/10.1093/mp/ssq001

De Vos, R.C., Moco, S., Lommen, A., Keurentjes, J.J., Bino, R.J., Hall, R.D., 2007. Untargeted large-scale plant metabolomics using liquid chromatography coupled to mass spectrometry. Nat. Protoc. 2, 778–791. https://doi.org/10.1038/nprot.2007.95

Deinlein, U., Stephan, A.B., Horie, T., Luo, W., Xu, G., Schroeder, J.I., 2014. Plant salt-tolerance mechanisms. Trends Plant Sci. 19, 371–379. https://doi.org/10.1016/j.tplants.2014.02.001

Delaporte, R.H., Sánchez, G.M., Cuellar, A.C., Giuliani, A., Palazzo de Mello, J.C., 2002. Anti-inflammatory activity and lipid peroxidation inhibition of iridoid lamiide isolated from *Bouchea fluminensis* (Vell.) Mold. (Verbenaceae). J. Ethnopharmacol. 82, 127–130. https://doi.org/10.1016/S0378-8741(02)00181-2

Dick, W.A., Kost, D., Chen, L., 2008. Availability of sulfur to crops from soil and other sources, in: Sulfur: A Missing Link between Soils, Crops, and Nutrition. John Wiley & Sons, Ltd, pp. 59–82. https://doi.org/10.2134/agronmonogr50.c5

Dimitriou, I., Aronsson, P., 2005. Willows for energy and phytoremediation in Sweden. Unasylva 56.

Drapal, M., Enfissi, E.M.A., Fraser, P.D., 2022. The chemotype core collection of genus *Nicotiana*. Plant J. 110, 1516–1528. https://doi.org/10.1111/tpj.15745

Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A.A., Melnik, A.V., Meusel, M., Dorrestein, P.C., Rousu, J., Böcker, S., 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. Nat. Methods 16, 299–302. https://doi.org/10.1038/s41592-019-0344-8

Dührkop, K., Nothias, L.-F., Fleischauer, M., Reher, R., Ludwig, M., Hoffmann, M.A., Petras, D., Gerwick, W.H., Rousu, J., Dorrestein, P.C., Böcker, S., 2021. Systematic classification of unknown metabolites using high-resolution fragmentation mass spectra. Nat. Biotechnol. 39, 462–471. https://doi.org/10.1038/s41587-020-0740-8

Dussarrat, T., Prigent, S., Latorre, C., Bernillon, S., Flandin, A., Díaz, F.P., Cassan, C., Van Delft, P., Jacob, D., Varala, K., Joubes, J., Gibon, Y., Rolin, D., Gutiérrez, R.A., Pétriacq, P., 2022. Predictive metabolomics of multiple Atacama plant species unveils a core set of generic metabolites for extreme climate resilience. New Phytol. 234, 1614–1628. https://doi.org/10.1111/nph.18095

Endo, I., Kobatake, M., Tanikawa, N., Nakaji, T., Ohashi, M., Makita, N., 2021. Anatomical patterns of condensed tannin in fine roots of tree species from a cool-temperate forest. Ann. Bot. 128, 59–71. https://doi.org/10.1093/aob/mcab022

Evelin, H., Kapoor, R., Giri, B., 2009. Arbuscular mycorrhizal fungi in alleviation of salt stress: a review. Ann. Bot. 104, 1263–1280. https://doi.org/10.1093/aob/mcp251

FAO, 2015. Status of the world's soil resources: Main report.

FAO, 1988. Salt-affected soils and their management (No. FAO Soils Bulletin 39).

Franco-Navarro, J.D., Brumós, J., Rosales, M.A., Cubero-Font, P., Talón, M., Colmenero-Flores, J.M., 2016. Chloride regulates leaf cell size and water relations in tobacco plants. J. Exp. Bot. 67, 873–891. https://doi.org/10.1093/jxb/erv502

Geilfus, C.-M., 2018. Chloride: from nutrient to toxicant. Plant Cell Physiol. 59, 877–886. https://doi.org/10.1093/pcp/pcy071

Genty, B., Briantais, J.-M., Baker, N.R., 1989. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. Biochim. Biophys. Acta BBA - Gen. Subj. 990, 87–92. https://doi.org/10.1016/S0304-4165(89)80016-9

Gil-de-la-Fuente, A., Godzien, J., Saugar, S., Garcia-Carmona, R., Badran, H., Wishart, D.S., Barbas, C., Otero, A., 2019. CEU Mass Mediator 3.0: A metabolite annotation tool. J. Proteome Res. 18, 797–802. https://doi.org/10.1021/acs.jproteome.8b00720

Gill, S.S., Tuteja, N., 2010a. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. Plant Physiol. Biochem. 48, 909–930. https://doi.org/10.1016/J.PLAPHY.2010.08.016

Gill, S.S., Tuteja, N., 2010b. Polyamines and abiotic stress tolerance in plants. Plant Signal. Behav. 5, 26–33. https://doi.org/10.4161/psb.5.1.10291

Gonzalez, E., Pitre, F.E., Pagé, A.P., Marleau, J., Guidi Nissim, W., St-Arnaud, M., Labrecque, M., Joly, S., Yergeau, E., Brereton, N.J.B., 2018. Trees, fungi and bacteria: tripartite metatranscriptomics of a root microbiome responding to soil contamination. Microbiome 6, 53. https://doi.org/10.1186/s40168-018-0432-5

Gupta, B., Huang, B., 2014. Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. Int. J. Genomics 2014, 701596–701596. https://doi.org/10.1155/2014/701596

Hangs, R.D., Schoenau, J.J., Van Rees, K.C.J., Steppuhn, H., 2011. Examining the salt tolerance of willow (*Salix* spp.) bioenergy species for use on salt-affected agricultural lands. Can. J. Plant Sci. 91, 509–517. https://doi.org/10.4141/cjps10135

Hartmann, L., Pedrotti, L., Weiste, C., Fekete, A., Schierstaedt, J., Göttler, J., Kempa, S., Krischke, M., Dietrich, K., Mueller, M.J., Vicente-Carbajosa, J., Hanson, J., Dröge-Laser, W., 2015. Crosstalk between two bZIP signaling pathways orchestrates salt-induced metabolic reprogramming in *Arabidopsis* roots. Plant Cell 27, 2244–2260. https://doi.org/10.1105/tpc.15.00163

Hauser, F., Horie, T., 2010. A conserved primary salt tolerance mechanism mediated by HKT transporters: a mechanism for sodium exclusion and maintenance of high K+/Na+ ratio in leaves during salinity stress. Plant Cell Environ. 33, 552–565. https://doi.org/10.1111/j.1365-3040.2009.02056.x

Hawkins, C., Ginzburg, D., Zhao, K., Dwyer, W., Xue, B., Xu, A., Rice, S., Cole, B., Paley, S., Karp, P., Rhee, S.Y., 2021. Plant Metabolic Network 15: A resource of genome-wide metabolism databases for 126 plants and algae. J. Integr. Plant Biol. 63, 1888–1905. https://doi.org/10.1111/jipb.13163

He, H., Tang, W., Wang, W., Crits-Christoph, P., 2014. Structural zeroes and zero-inflated models. Shanghai Arch. Psychiatry 26, 236–242. https://doi.org/10.3969/j.issn.1002-0829.2014.04.008

Huang, X., Hefer, C.A., Soolanayakanahally, R.Y., Guy, R.D., Mansfield, S.D., 2023. The overlooked ion: Unraveling the effects of magnesium-specific toxicity on willows under sulphate salinity. Environ. Exp. Bot. 105634. https://doi.org/10.1016/j.envexpbot.2023.105634

Huang, X., Soolanayakanahally, R.Y., Guy, R.D., Shunmugam, A.S.K., Mansfield, S.D., 2020. Differences in growth and physiological and metabolic responses among Canadian native and hybrid willows (*Salix* spp.) under salinity stress. Tree Physiol. 40, 652–666. https://doi.org/10.1093/treephys/tpaa017

Huber, F., Ridder, L., Verhoeven, S., Spaaks, J.H., Diblen, F., Rogers, S., Hooft, van der J.J.J., 2021. Spec2Vec: Improved mass spectral similarity scoring through learning of structural relationships. PLOS Comput. Biol. 17, e1008724. https://doi.org/10.1371/journal.pcbi.1008724

Ifuku, K., 2015. Localization and functional characterization of the extrinsic subunits of photosystem II: an update. Biosci. Biotechnol. Biochem. 79, 1223–1231. https://doi.org/10.1080/09168451.2015.1031078

Janz, D., Lautner, S., Wildhagen, H., Behnke, K., Schnitzler, J.-P., Rennenberg, H., Fromm, J., Polle, A., 2012. Salt stress induces the formation of a novel type of 'pressure wood' in two *Populus* species. New Phytol. 194, 129–141. https://doi.org/10.1111/j.1469-8137.2011.03975.x

Jia, H., Wang, L., Li, J., Sun, P., Lu, M., Hu, J., 2020. Comparative metabolomics analysis reveals different metabolic responses to drought in tolerant and susceptible poplar species. Physiol. Plant. 168, 531–546. https://doi.org/10.1111/ppl.13036

Jiang, Z., Zhou, X., Tao, M., Yuan, F., Liu, L., Wu, F., Wu, X., Xiang, Y., Niu, Y., Liu, F., Li, C., Ye, R., Byeon, B., Xue, Y., Zhao, H., Wang, H.-N., Crawford, B.M., Johnson, D.M., Hu, C., Pei, C., Zhou, W., Swift, G.B., Zhang, H., Vo-Dinh, T., Hu, Z., Siedow, J.N., Pei, Z.-M., 2019. Plant cell-surface GIPC sphingolipids sense salt to trigger Ca2+ influx. Nature 572, 341–346. https://doi.org/10.1038/s41586-019-1449-z

Jouve, L., Hoffmann, L., Hausman, J.-F., 2004. Polyamine, carbohydrate, and proline content changes during salt stress exposure of Aspen (*Populus tremula* L.): Involvement of oxidation and osmoregulation metabolism. Plant Biol. 6, 74–80. https://doi.org/10.1055/s-2003-44687

Kaling, M., Kanawati, B., Ghirardo, A., Albert, A., Winkler, J.B., Heller, W., Barta, C., Loreto, F., Schmitt-Kopplin, P., Schnitzler, J.-P., 2015. UV-B mediated metabolic rearrangements in poplar revealed by non-targeted metabolomics. Plant Cell Environ. 38, 892–904. https://doi.org/10.1111/pce.12348

Karpievitch, Y.V., Nikolic, S.B., Wilson, R., Sharman, J.E., Edwards, L.M., 2014. Metabolomics data normalization with EigenMS. PloS One 9, e116221. https://doi.org/10.1371/journal.pone.0116221

Kim, H.W., Wang, M., Leber, C.A., Nothias, L.-F., Reher, R., Kang, K.B., van der Hooft, J.J.J., Dorrestein, P.C., Gerwick, W.H., Cottrell, G.W., 2021. NPClassifier: A deep neural network-based structural classification tool for natural products. J. Nat. Prod. 84, 2795–2807. https://doi.org/10.1021/acs.jnatprod.1c00399

Köhler, A., Förster, N., Zander, M., Ulrichs, C., 2020. Compound-specific responses of phenolic metabolites in the bark of drought-stressed *Salix daphnoides* and *Salix purpurea*. Plant Physiol. Biochem. 155, 311–320. https://doi.org/10.1016/j.plaphy.2020.07.004

Kovinich, N., Kayanja, G., Chanoca, A., Otegui, M.S., Grotewold, E., 2015. Abiotic stresses induce different localizations of anthocyanins in *Arabidopsis*. Plant Signal. Behav. 10, e1027850. https://doi.org/10.1080/15592324.2015.1027850

Li, D., Heiling, S., Baldwin, I.T., Gaquerel, E., 2016. Illuminating a plant's tissue-specific metabolic diversity using computational metabolomics and information theory. Proc. Natl. Acad. Sci. 113, E7610–E7618. https://doi.org/10.1073/pnas.1610218113

Liu, Y., Wang, L., Li, X., Luo, M., 2022. Detailed sphingolipid profile responded to salt stress in cotton root and the GhIPCS1 is involved in the regulation of plant salt tolerance. Plant Sci. 316, 111174. https://doi.org/10.1016/j.plantsci.2021.111174

Llanes, A., Bertazza, G., Palacio, G., Luna, V., 2013. Different sodium salts cause different solute accumulation in the halophyte *Prosopis strombulifera*. Plant Biol. 15, 118–125. https://doi.org/10.1111/j.1438-8677.2012.00626.x

Ludwig-Müller, J., 2011. Auxin conjugates: Their role for plant development and in the evolution of land plants. J. Exp. Bot. 62, 1757–1773. https://doi.org/10.1093/jxb/erq412

Major, J.E., Mosseler, A., Malcolm, J.W., 2017. *Salix* species variation in leaf gas exchange, sodium, and nutrient parameters at three levels of salinity. Can. J. For. Res. 47, 1045–1055. https://doi.org/10.1139/cjfr-2017-0028

McLaughlin, S., Zhalnina, K., Kosina, S., Northen, T.R., Sasse, J., 2023. The core metabolome and root exudation
dynamics of three phylogenetically distinct plant species. Nat. Commun. 14, 1649.
https://doi.org/10.1038/s41467-023-37164-x
Meier, B., Julkunen-Tiitto, R., Tahvanainen, J., Sticher, O., 1988. Comparative high-performance liquid and gas-
liquid chromatographic determination of phenolic glucosides in *Salicaceae* species. J. Chromatogr. A
442, 175–186. https://doi.org/10.1016/S0021-9673(00)94467-4
Mirck, J., Zalesny, R.S., 2015. Mini-review of knowledge gaps in salt tolerance of plants applied to willows and
poplars. Int. J. Phytoremediation 17, 640–650. https://doi.org/10.1080/15226514.2014.950414
Munns, R., 2002. Comparative physiology of salt and water stress. Plant Cell Environ. 25, 239–250.
https://doi.org/10.1046/j.0016-8025.2001.00808.x
Munns, R., Day, D.A., Fricke, W., Watt, M., Arsova, B., Barkla, B.J., Bose, J., Byrt, C.S., Chen, Z.-H., Foster, K.J.,
Gilliham, M., Henderson, S.W., Jenkins, C.L.D., Kronzucker, H.J., Miklavcic, S.J., Plett, D., Roy, S.J.,
Shabala, S., Shelden, M.C., Soole, K.L., Taylor, N.L., Tester, M., Wege, S., Wegner, L.H., Tyerman, S.D.,
2020a. Energy costs of salt tolerance in crop plants. New Phytol. 225, 1072–1090.
https://doi.org/10.1111/nph.15864
Munns, R., Passioura, J.B., Colmer, T.D., Byrt, C.S., 2020b. Osmotic adjustment and energy limitations to plant
growth in saline soil. New Phytol. 225, 1091–1096. https://doi.org/10.1111/nph.15862
Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. Annu. Rev. Plant Biol. 59, 651–681.
https://doi.org/10.1146/annurev.arplant.59.032607.092911
Nakabayashi, R., Yonekura-Sakakibara, K., Urano, K., Suzuki, M., Yamada, Y., Nishizawa, T., Matsuda, F., Kojima,
M., Sakakibara, H., Shinozaki, K., Michael, A.J., Tohge, T., Yamazaki, M., Saito, K., 2014. Enhancement
of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. Plant
J. 77, 367–379. https://doi.org/10.1111/tpj.12388
Neess, D., Bek, S., Engelsby, H., Gallego, S.F., Færgeman, N.J., 2015. Long-chain acyl-CoA esters in metabolism
and signaling: Role of acyl-CoA binding proteins. Prog. Lipid Res. 59, 1–25.
https://doi.org/10.1016/j.plipres.2015.04.001
Neves, G.Y.S., Marchiosi, R., Ferrarese, M.L.L., Siqueira-Soares, R.C., Ferrarese-Filho, O., 2010. Root growth
inhibition and lignification induced by salt stress in soybean. J. Agron. Crop Sci. 196, 467–473.
https://doi.org/10.1111/j.1439-037X.2010.00432.x
Nguyen, H.T., Meir, P., Sack, L., Evans, J.R., Oliveira, R.S., Ball, M.C., 2017. Leaf water storage increases with
salinity and aridity in the mangrove *Avicennia marina*: Integration of leaf structure, osmotic adjustment
and access to multiple water sources. Plant Cell Environ. 40, 1576–1591.
https://doi.org/10.1111/pce.12962
Nothias, L.-F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., Protsyuk, I., Ernst, M., Tsugawa, H.,
Fleischauer, M., Aicheler, F., Aksenov, A.A., Alka, O., Allard, P.-M., Barsch, A., Cachet, X., Caraballo-
Rodriguez, A.M., Da Silva, R.R., Dang, T., Garg, N., Gauglitz, J.M., Gurevich, A., Isaac, G., Jarmusch, A.K.,
Kameník, Z., Kang, K.B., Kessler, N., Koester, I., Korf, A., Le Gouellec, A., Ludwig, M., Martin H., C.,
McCall, L.-I., McSayles, J., Meyer, S.W., Mohimani, H., Morsy, M., Moyne, O., Neumann, S., Neuweger,
H., Nguyen, N.H., Nothias-Esposito, M., Paolini, J., Phelan, V.V., Pluskal, T., Quinn, R.A., Rogers, S.,
Shrestha, B., Tripathi, A., van der Hooft, J.J.J., Vargas, F., Weldon, K.C., Witting, M., Yang, H., Zhang, Z.,
Zubeil, F., Kohlbacher, O., Böcker, S., Alexandrov, T., Bandeira, N., Wang, M., Dorrestein, P.C., 2020.
Feature-based molecular networking in the GNPS analysis environment. Nat. Methods 17, 905–908.
https://doi.org/10.1038/s41592-020-0933-6
Osawa, H., Endo, I., Hara, Y., Matsushima, Y., Tange, T., 2011. Transient proliferation of proanthocyanidin-
accumulating cells on the epidermal apex contributes to highly aluminum-resistant root elongation in
camphor tree. Plant Physiol. 155, 433–446. https://doi.org/10.1104/pp.110.166967
Ottow, E.A., Brinker, M., Teichmann, T., Fritz, E., Kaiser, W., Brosché, M., Kangasjärvi, J., Jiang, X., Polle, A.,
2005. *Populus euphratica* displays apoplastic sodium accumulation, osmotic adjustment by decreases
in calcium and soluble carbohydrates, and develops leaf succulence under salt stress. Plant Physiol.
139, 1762–1772. https://doi.org/10.1104/pp.105.069971
Pichersky, E., Raguso, R.A., 2018. Why do plants produce so many terpenoid compounds? New Phytol. 220,
692–702. https://doi.org/10.1111/nph.14178

Pobłocka-Olech, L., van Nederkassel, A.-M., Vander Heyden, Y., Krauze-Baranowska, M., Glód, D., Baczek, T.,
2007. Chromatographic analysis of salicylic compounds in different species of the genus *Salix*. J. Sep.
Sci. 30, 2958–2966. https://doi.org/10.1002/jssc.200700137

Prasad, R., Shivay, Y.S., 2018. Sulphur in soil, plant and human nutrition. Proc. Natl. Acad. Sci. India Sect. B Biol.
Sci. 88, 429–434. https://doi.org/10.1007/s40011-016-0769-0

Quinn, L.D., Straker, K.C., Guo, J., Kim, S., Thapa, S., Kling, G., Lee, D.K., Voigt, T.B., 2015. Stress-tolerant
feedstocks for sustainable bioenergy production on marginal land. BioEnergy Res. 8, 1081–1100.
https://doi.org/10.1007/s12155-014-9557-y

R Core Team, 2022. R: A Language and environment for statistical computing.

Rausch, T., Wachter, A., 2005. Sulfur metabolism: a versatile platform for launching defence operations. Trends
Plant Sci. 10, 503–509. https://doi.org/10.1016/j.tplants.2005.08.006

Reginato, M., Luna, V., Papenbrock, J., 2021. Current knowledge about Na2SO4 effects on plants: What is
different in comparison to NaCl? J. Plant Res. 134, 1159–1179. https://doi.org/10.1007/s10265-021-
01335-y

Reich, M., Aghajanzadeh, T., Helm, J., Parmar, S., Hawkesford, M.J., De Kok, L.J., 2017. Chloride and sulfate
salinity differently affect biomass, mineral nutrient composition and expression of sulfate transport
and assimilation genes in *Brassica rapa*. Plant Soil 411, 319–332. https://doi.org/10.1007/s11104-016-
3026-7

Reyt, G., Ramakrishna, P., Salas-González, I., Fujita, S., Love, A., Tiemessen, D., Lapierre, C., Morreel, K., Calvo-
Polanco, M., Flis, P., Geldner, N., Boursiac, Y., Boerjan, W., George, M.W., Castrillo, G., Salt, D.E., 2021.
Two chemically distinct root lignin barriers control solute and water balance. Nat. Commun. 12, 2320.
https://doi.org/10.1038/s41467-021-22550-0

Richter, J.A., Behr, J.H., Erban, A., Kopka, J., Zörb, C., 2019. Ion-dependent metabolic responses of *Vicia faba* L.
to salt stress. Plant Cell Environ. 42, 295–309. https://doi.org/10.1111/pce.13386

Rodriguez, H.G., Roberts, JKM., Jordan, W.R., Drew, M.C., 1997. Growth, water relations, and accumulation of
organic and inorganic solutes in roots of maize seedlings during salt stress. Plant Physiol. 113, 881–893.
https://doi.org/10.1104/pp.113.3.881

Rodzkin, A., Volk, T., 2018. Using schrub willow for biological reclamation of areas with raised soil salinity.
Presented at the 12th Eco-conference® on environmental protection of urban and suburban
settlements, Novi Sad, Serbia.

Rohart, F., Gautier, B., Singh, A., Cao, K.-A.L., 2017. mixOmics: An R package for 'omics feature selection and
multiple data integration. PLOS Comput. Biol. 13, e1005752.
https://doi.org/10.1371/journal.pcbi.1005752

Ryu, H., Cho, Y.-G., 2015. Plant hormones in salt stress tolerance. J. Plant Biol. 58, 147–155.
https://doi.org/10.1007/s12374-015-0103-z

Sanchez, D.H., Siahpoosh, M.R., Roessner, U., Udvardi, M., Kopka, J., 2008. Plant metabolomics reveals
conserved and divergent metabolic responses to salinity. Physiol. Plant. 132, 209–219.
https://doi.org/10.1111/j.1399-3054.2007.00993.x

Sas, E., Hennequin, L.M., Frémont, A., Jerbi, A., Legault, N., Lamontagne, J., Fagoaga, N., Sarrazin, M., Hallett,
J.P., Fennell, P.S., Barnabé, S., Labrecque, M., Brereton, N.J.B., Pitre, F.E., 2021. Biorefinery potential
of sustainable municipal wastewater treatment using fast-growing willow. Sci. Total Environ. 792,
148146. https://doi.org/10.1016/j.scitotenv.2021.148146

Sasse, J., Martinoia, E., Northen, T., 2018. Feed your friends: Do plant exudates shape the root microbiome?
Trends Plant Sci. 23, 25–41. https://doi.org/10.1016/j.tplants.2017.09.003

Schmid, R., Heuckeroth, S., Korf, A., Smirnov, A., Myers, O., Dyrlund, T.S., Bushuiev, R., Murray, K.J., Hoffmann,
N., Lu, M., Sarvepalli, A., Zhang, Z., Fleischauer, M., Dührkop, K., Wesner, M., Hoogstra, S.J., Rudt, E.,
Mokshyna, O., Brungs, C., Ponomarov, K., Mutabdžija, L., Damiani, T., Pudney, C.J., Earll, M., Helmer,
P.O., Fallon, T.R., Schulze, T., Rivas-Ubach, A., Bilbao, A., Richter, H., Nothias, L.-F., Wang, M., Orešič,
M., Weng, J.-K., Böcker, S., Jeibmann, A., Hayen, H., Karst, U., Dorrestein, P.C., Petras, D., Du, X., Pluskal,
T., 2023. Integrative analysis of multimodal mass spectrometry data in MZmine 3. Nat. Biotechnol. 41,
447–449. https://doi.org/10.1038/s41587-023-01690-2

Schymanski, E.L., Jeon, J., Gulde, R., Fenner, K., Ruff, M., Singer, H.P., Hollender, J., 2014. Identifying small
molecules via high resolution mass spectrometry: Communicating confidence. Environ. Sci. Technol.
48, 2097–2098. https://doi.org/10.1021/es5002105

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., Ideker, T.,
2003. Cytoscape: A software environment for integrated models of biomolecular interaction networks.
Genome Res. 13, 2498–2504. https://doi.org/10.1101/gr.1239303

Shavrukov, Y., 2013. Salt stress or salt shock: Which genes are we studying? J. Exp. Bot. 64, 119–127.
https://doi.org/10.1093/jxb/ers316

Si, C.-L., Kim, J.-K., Bae, Y.-S., Li, S.-M., 2009. Phenolic compounds in the leaves of *Populus ussuriensis* and their
antioxidant activities. Planta Med. 75, 1165–1167. https://doi.org/10.1055/s-0029-1185476

Singh, A., 2021. Soil salinization management for sustainable development: A review. J. Environ. Manage. 277,
111383. https://doi.org/10.1016/j.jenvman.2020.111383

Skerrett, M., Tyerman, S.D., 1994. A channel that allows inwardly directed fluxes of anions in protoplasts
derived from wheat roots. Planta 192, 295–305. https://doi.org/10.1007/BF00198563

Slama, I., Abdelly, C., Bouchereau, A., Flowers, T., Savouré, A., 2015. Diversity, distribution and roles of
osmoprotective compounds accumulated in halophytes under abiotic stress. Ann. Bot. 115, 433–447.
https://doi.org/10.1093/aob/mcu239

Sonar, P., Bellucci, L., Mossa, A., Heidarsson, P.O., Kragelund, B.B., Cecconi, C., 2020. Effects of ligand binding
on the energy landscape of acyl-CoA-binding protein. Biophys. J. 119, 1821–1832.
https://doi.org/10.1016/j.bpj.2020.09.016

Sonmez, S., Buyuktas, D., Okturen, F., Citak, S., 2008. Assessment of different soil to water ratios (1:1, 1:2.5,
1:5) in soil salinity studies. Geoderma, Antarctic Soils and Soil Forming Processes in a Changing
Environment 144, 361–369. https://doi.org/10.1016/j.geoderma.2007.12.005

Sui, D., Wang, B., 2020. Transcriptome analysis reveals complex defensive mechanisms in salt-tolerant and salt-
sensitive shrub willow genotypes under salinity stress. Int. J. Genomics 2020, e6870157.
https://doi.org/10.1155/2020/6870157

Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., Fan, T.W.-M., Fiehn, O., Goodacre,
R., Griffin, J.L., Hankemeier, T., Hardy, N., Harnly, J., Higashi, R., Kopka, J., Lane, A.N., Lindon, J.C.,
Marriott, P., Nicholls, A.W., Reily, M.D., Thaden, J.J., Viant, M.R., 2007. Proposed minimum reporting
standards for chemical analysis. Metabolomics 3, 211–221. https://doi.org/10.1007/s11306-007-0082-
2

Teakle, N.L., Tyerman, S.D., 2010. Mechanisms of Cl- transport contributing to salt tolerance. Plant Cell Environ.
33, 566–589. https://doi.org/10.1111/j.1365-3040.2009.02060.x

Tyśkiewicz, K., Konkol, M., Kowalski, R., Rój, E., Warmiński, K., Krzyżaniak, M., Gil, Ł., Stolarski, M.J., 2019.
Characterization of bioactive compounds in the biomass of black locust, poplar and willow. Trees 33,
1235–1263. https://doi.org/10.1007/s00468-019-01837-2

van Zelm, E., Zhang, Y., Testerink, C., 2020. Salt tolerance mechanisms of plants. Annu. Rev. Plant Biol. 71, 403–
433. https://doi.org/10.1146/annurev-arplant-050718-100005

Volf, M., Hrcek, J., Julkunen-Tiitto, R., Novotny, V., 2015. To each its own: differential response of specialist and
generalist herbivores to plant defence in willows. J. Anim. Ecol. 84, 1123–1132.
https://doi.org/10.1111/1365-2656.12349

Volk, T.A., Abrahamson, L.P., Nowak, C.A., Smart, L.B., Tharakan, P.J., White, E.H., 2006. The development of
short-rotation willow in the northeastern United States for bioenergy and bioproducts, agroforestry
and phytoremediation. Biomass Bioenergy, Proceedings of the 5th Biennial meeting of the Short
Rotation Woody Crops Operations Working Group in conjunction with International Energy Agency
Bioenergy, Task 30, Short Rotation Crops for Bioenergy Systems and International Union of Forest
Research Organization, Working Unit 1.09.01, Integrated Research in Temperate Short-Rotation Energy
Plantations 30, 715–727. https://doi.org/10.1016/j.biombioe.2006.03.001

Watanabe, M., Hoefgen, R., 2019. Sulphur systems biology: Making sense of omics data. J. Exp. Bot. 70, 4155–
4170. https://doi.org/10.1093/jxb/erz260

Watanabe, S., Kojima, K., Ide, Y., Sasaki, S., 2000. Effects of saline and osmotic stress on proline and sugar
accumulation in *Populus euphratica in vitro*. Plant Cell Tissue Organ Cult. 63, 199–206.
https://doi.org/10.1023/A:1010619503680

Wege, S., Gilliham, M., Henderson, S.W., 2017. Chloride: not simply a 'cheap osmoticum', but a beneficial plant
macronutrient. J. Exp. Bot. 68, 3057–3069. https://doi.org/10.1093/jxb/erx050

Wei, H., Movahedi, A., Xu, C., Sun, W., Li, L., Wang, P., Li, D., Zhuge, Q., 2020. Overexpression of PtHMGR
enhances drought and salt tolerance of poplar. Ann. Bot. 125, 785–803.
https://doi.org/10.1093/aob/mcz158

White, P.J., Broadley, M.R., 2001. Chloride in soils and its uptake and movement within the plant: A review.
Ann. Bot. 88, 967–988. https://doi.org/10.1006/anbo.2001.1540

Wu, D., Cai, S., Chen, M., Ye, L., Chen, Z., Zhang, H., Dai, F., Wu, F., Zhang, G., 2013. Tissue metabolic responses
to salt stress in wild and cultivated barley. PLOS ONE 8, e55431.
https://doi.org/10.1371/journal.pone.0055431

Xia, L., Xiaodong, M., Yunhe, C., Junxiang, L., Junzhu, Z., Feifei, Z., Zhenyuan, S., Lei, H., 2021. Transcriptomic
and metabolomic insights into the adaptive response of *Salix viminalis* to phenanthrene. Chemosphere
262, 127573. https://doi.org/10.1016/j.chemosphere.2020.127573

Xiao, J., Muzashvili, T.S., Georgiev, M.I., 2014. Advances in the biotechnological glycosylation of valuable
flavonoids. Biotechnol. Adv. 32, 1145–1156. https://doi.org/10.1016/j.biotechadv.2014.04.006

Yancey, P.H., 2005. Organic osmolytes as compatible, metabolic and counteracting cytoprotectants in high
osmolarity and other stresses. J. Exp. Biol. 208, 2819–2830. https://doi.org/10.1242/jeb.01730

Yang, Y., Guo, Y., 2018. Elucidating the molecular mechanisms mediating plant salt-stress responses. New
Phytol. 217, 523–539. https://doi.org/10.1111/nph.14920

Zhang, Y., Fang, J., Wu, X., Dong, L., 2018. Na+/K+ balance and transport regulatory mechanisms in weedy and
cultivated rice (Oryza sativa L.) under salt stress. BMC Plant Biol. 18, 375.
https://doi.org/10.1186/s12870-018-1586-9

Zhou, J., Guo, J., Chen, Q., Wang, B., He, X., Zhuge, Q., Wang, P., 2022. Different color regulation mechanism in
willow barks determined using integrated metabolomics and transcriptomics analyses. BMC Plant Biol.
22, 530. https://doi.org/10.1186/s12870-022-03909-x

Zhou, J., Huang, J., Tian, X., Zheng, J., He, X., 2020. Transcriptome analysis reveals dynamic changes in the salt
stress response in *Salix*. J. For. Res. 31, 1851–1862. https://doi.org/10.1007/s11676-019-00941-w

Zhou, Z., Luo, M., Zhang, H., Yin, Y., Cai, Y., Zhu, Z.-J., 2022. Metabolite annotation from knowns to unknowns
through knowledge-guided multi-layer metabolic networking. Nat. Commun. 13, 6656.
https://doi.org/10.1038/s41467-022-34537-6

Zörb, C., Geilfus, C.-M., Dietz, K.-J., 2019. Salinity and crop yield. Plant Biol. 21, 31–38.
https://doi.org/10.1111/plb.12884

Zou, Y., Zhang, Y., Testerink, C., 2022. Root dynamic growth strategies in response to salinity. Plant Cell Environ.
45, 695–704. https://doi.org/10.1111/pce.14205

1005

Figure 1: *Workflow for untargeted metabolomics approach.*

A randomized pot trial treated willows with salt solutions or water controls (7 replicates per treatment) to explore variation in anion-specific salt tolerance mechanisms across organs. Metabolites were extracted from root, stem and leaf samples and analysed by LC-MS/MS. Resulting chromatographic data were pre-processed using MZmine for features mass detection, cross-sample alignment and quality filters. Statistical analysis used a cross-sample aligned peak area table. After log transformation and Eigen MS normalization, univariate analysis was used to factorially compare feature peak areas across willow organs in controls, while pairwise comparisons were used to identify differentially abundant (DA) compounds between each salt treatment compared to controls. Parametric or non-parametric tests were used based on feature peak area normality and variance homogeneity before a false discovery rate correction was applied to account for multiple testing. Annotation strategy used precursors information and corresponding spectral fingerprints (when available) to query multiple databases and for *in silico* structure prediction to generate a list of annotation candidates (unsupervised annotation). After standardizing compounds identity information and assigning prioritization criteria, the best candidate was retained. Annotation was further refined for compounds identified as differentially abundant due to salt treatment and used data-driven (structural similarity networking) and knowledge-driven (literature) curation.

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

Figure 2: Physiological assessment of salt stress on *Salix miyabeana* 'SX64'.
(a) Following 56 days of root establishment, salinity in planted pots was progressively built-up over 28 days to reach the final saturated soil electroconductivity (EC), as inferred from a (1:1) soil-to-water extract. **(b)** Representative willow photographs, on harvest day, in response to salt treatments (white scale bars: 50 cm). The impact of moderate NaCl, moderate Na_2SO_4 and high Na_2SO_4 were quantified on **(c)** total biomass (dry weight), **(d)** photosystem II efficiency evaluated using the LI-COR 600 parameter phiPSII, **(e)** stomatal conductance gsw, **(f)** stems and leaves moisture contents. Boxes represent median and quartiles while whiskers extend to the largest value within 1.5 times the inter-quartile range (IQR) from the hinge. Dots represent individual plants (n=7). Significant differences (p<0.05) indicated by letters were determined by two-way ANOVA followed by Tukey's post hoc test.

1018

Figure 3: Elemental analysis of mineral nutrients.
Elemental analyses include sodium (Na), chloride (Cl), sulphur (S), potassium (K), phosphorus (P), calcium (Ca), and magnesium (Mg). **(a)** Total element content (mmol) combined from the leaves, stems and roots presented. The bars indicate mean values within each organ ± standard error (n=3). **(b)** Element concentration within a single organ (μmol g_{DW}^{-1}) is shown. The dots represent mean values ± standard error (n=3). Letters indicate significant differences (p<0.05) between control, moderate NaCl, moderate Na_2SO_4, and high Na_2SO_4 treatments and were determined using two-way ANOVA followed by Tukey's post hoc test. Mineral elements ratios (K^+/Na^+; Ca^{2+}/Na^+; K^+/Ca^{2+}) are presented in Supporting information – Fig.S4.

1019

Figure 4: Willow metabolic composition.
Organs (leaves, stems and roots) were separately collected from control willows and analysed for metabolomic profiling. **(a)** Total number of consistently detected metabolites within each major organ (present in at least 4

replicates out of 7). **(b)** Comparison of organ metabolic richness using Shannon and Inverse Simpson diversity indices (n=7), different letters indicate significant differences (ANOVA, Tukey HSD, p<0.05). **(c)** Principal component analysis depicting dissimilarity in organ composition based on Bray-Curtis distances, ellipses are for standard deviation. **(d)** Venn diagram illustrating the distribution of metabolites across organs as unique to one or shared between multiple. **(e)** Cumulative peak areas of metabolites present in each organ, categorised based on their exclusive presence in the respective organ or shared across multiple organs. **(f)** Cumulative peak area and number of core metabolites (compounds present across the entire plant, with no significant variation of abundance between organs) classified per metabolic pathways, with composition details of main superclasses. **(g)** Count of compounds significantly higher in one organ compared to the other two (ANOVA, Tukey HSD, p<0.05), revealing dominant organ-specialisation of some metabolic superclasses.

1020

Figure 5: Willow metabolic response to sodium sulphate (Na₂SO₄) salt and its variation with dose
(a) Principal component analysis (PCA) based on Bray-Curtis distance, illustrating metabolic profile dissimilarity between control, moderate Na₂SO₄ and high Na₂SO₄ at organ scale (n=7), including standard deviation ellipse

(95% confidence interval). **(b)** Organ distribution of differentially abundant (DA) compounds between control and moderate concentration Na_2SO_4-treated willows (p<0.05). **(c)** Organ distribution of differentially abundant compounds between control and high concentration Na_2SO_4-treated willows (p<0.05). **(d)** Overview of significantly enriched and depleted compounds across organs, comparing DA of control against moderate Na_2SO_4 and control against high Na_2SO_4. **(e)** Distribution of enriched and depleted DA compounds across metabolic pathways for moderate and high Na_2SO_4 treatments compared to controls. **(f)** Metabolic response overlap across both Na_2SO_4 concentrations, representing the number of DA compounds common and unique to a single dose DA, at organ scale. **(g)** Dose-dependent correlation of compounds consistently DA across Na_2SO_4 levels within the same organ (excluding salt-induced/salt-suppressed compounds with infinite fold change), representing the fold change (log2) of average peak areas (n=7) under moderate Na_2SO_4 relative to control against high Na_2SO_4 relative to control. **(h)** Sub-selection of main superclasses altered by moderate or high concentration of Na_2SO_4 compared to controls, illustrating enriched and depleted compounds distribution at organ scale. DA compounds shared between both treatment levels are highlighted by the darker section of the bars. **(i)** Distribution of enriched and depleted sulphur-containing DA compounds at organ scale, comparing both levels of Na_2SO_4. Shared compounds are highlighted by the darker section of the bars. **(j)** Molecular network of compounds MS/MS spectra similarities (cosine >0.6) computed in the GNPS environment and visualised in Cytoscape. Dots are color-coded based on the presence of sulphur (pink represents S-containing compounds) and dot size highlights significantly different compounds (larger dots are DA compounds). A major cluster containing 16 S-containing and structurally similar DA compounds is circled by an ellipse.

1021

Figure 6: Willow metabolic response to sodium chloride (NaCl) salt.
(a) Principal component analysis (PCA) based on Bray-Curtis distance, illustrating metabolic profile dissimilarity between control and moderate NaCl at organ scale (n=7), including standard deviation ellipse (95% confidence

interval) **(b)** Overview of significantly enriched and depleted compounds across organs. **(c)** Organ distribution of differentially abundant (DA) compounds between control and moderate concentration NaCl-treated willows (p<0.05). **(d)** Distribution of enriched and depleted DA compounds across metabolic pathways. **(e)** Metabolic profile at pathway level of DA compounds shared between multiple organs and unique to roots, stems or leaves. Sub-selection of main pathways or superclasses altered by moderate NaCl compared to controls, illustrating enriched and depleted compounds distribution at organ scale for **(f)** carbohydrates, including saccharides and nucleosides, **(g)** small peptides, **(h)** and 3 major superclass of shikimates and phenylpropanoids pathway: salicinoids, lignans, phenylpropanoids, flavonoids, as well as **(i)** terpenoids, including monoterpenoids and steroids. Dots are color-coded by regulation (blue: depleted, red: enriched by salt treatment compared to control). Base mean area reflects average peak area across all compared samples.

1022

Figure 7: General willow salt response and anion specific tailored responses.

(a) Venn diagrams showing DA metabolic responses overlap in roots, stems, and leaves under moderate NaCl and Na_2SO_4 stress. **(b**) Pathway profile of commonly enriched and depleted compounds in response to moderate salt stress. Fold change graph illustrated enriched and depleted compounds distribution at organ scale. **(c)** Superclass profiles of NaCl- and Na_2SO_4-specific responses. Superclasses are categorized as net enriched, equal, or net depleted based on the relative number of enriched versus depleted compounds. Differential patterns for nucleosides and saccharides are highlighted in bold. **(d)** Metabolic response to moderate NaCl and moderate Na_2SO_4 categorised based on compounds distribution across the whole plant: salt-induced compounds (not detected in controls), core compounds (present in all organs at stable state in controls), organ-specialised compounds: roots, stems or leaves (present in a higher abundance in a specific organ, based on controls). Bar graphs are split as depleted and enriched compounds to highlight patterning of the DA compounds.

1023

1024