1  ***In vitro* evolution of ciprofloxacin resistance in *Neisseria* commensals and derived**
2  **mutation population dynamics in natural *Neisseria* populations**
3

4  **Leah R. Robinson[1], Caroline J. Mcdevitt[1], Molly R. Regan[1], Sophie L. Quail[1], Crista B.**
5  **Wadsworth[1]**
6
7  **[1]Rochester Institute of Technology, Thomas H. Gosnell School of Life Sciences,**
8  **Rochester, New York, USA.**

9
**Abstract**

Commensal *Neisseria* are members of a healthy human oropharyngeal microbiome; however**,**
they also serve as a reservoir of antimicrobial resistance for their pathogenic relatives. Despite
their known importance as sources of novel genetic variation for pathogens, we still do not
understand the full suite of resistance mutations commensal species can harbor. Here, we use *in*
*vitro* selection to assess the mutations that emerge in response to ciprofloxacin selection in
commensal *Neisseria* by passaging 4 replicates of 4 different species in the presence of a
selective antibiotic gradient for 20 days; then categorized derived mutations with whole genome
sequencing. 10/16 selected cells lines across the 4 species evolved ciprofloxacin resistance (≥ 1
ug/ml); with resistance-contributing mutations primarily emerging in *DNA gyrase subunit A* and *B*
(*gyrA* and *gyrB*), *topoisomerase IV subunits C* and *E* (*parC* and *parE*), and the *multiple*
*transferable efflux pump repressor* (*mtrR*). Of note, these derived mutations appeared in the same
loci responsible for ciprofloxacin reduced susceptibility in the pathogenic *Neisseria*, suggesting
conserved mechanisms of resistance across the genus. Additionally, we tested for zoliflodacin
cross-resistance in evolved strain lines and found 6 lineages with elevated zoliflodacin minimum
inhibitory concentrations. Finally, to interrogate the likelihood of experimentally derived mutations
emerging and contributing to resistance in natural *Neisseria*, we used a population-based
approach and identified GyrA 91I as a substitution circulating within commensal *Neisseria*
populations and ParC 85C in a single gonococcal isolate. Small clusters of gonococcal isolates
had commensal-like alleles at *parC* and *parE*, indicating recent cross-species recombination
events.

31
32  **Keywords:** *Neisseria*, commensals, horizontal gene transfer, *in vitro* evolution
33
**Introduction**
Commensal species within the genus *Neisseria* have an intertwined relationship with their
pathogenic relatives *Neisseria gonorrhoeae* (Ngo, the pathogen responsible for the sexually
transmitted infection (STI) gonorrhea) and *N. meningitidis* (Nmen, the primary *Neisseria* species
responsible for meningococcemia and bacterial meningitis), through widespread horizontal gene
transfer (HGT) of both chromosomal[1–5] and plasmid DNA[6]. Genetic exchange across species'
boundaries has led to the acquisition of important phenotypic traits, such as antibiotic resistance[1–5], however**,** the commensal *Neisseria* are relatively underexplored for the phenotypic and
[5], however**,** the commensal *Neisseria* are relatively underexplored for the phenotypic and
genotypic diversity they harbor. Making up the most abundant genera within proteobacteria in oral
and pharyngeal samples (~10% of operational taxonomic units [OTUs]) in 100% of healthy human
adults and children[7–11], commensal species are a large portion of our oropharyngeal microbiomes;
and thus will be a persistent threat for rapid pathogen evolution through bystander selection and
DNA donation.
Though attention on the acquisition of antibiotic resistance through HGT between the
*Neisseria* has often focused on beta-lactam resistance acquisition through mosaic *penicillin-*
*binding protein 2* (*penA*)[4,12–15] and macrolide resistance through inheritance of mosaic *multiple*
*transferable efflux pump* (*mtr*) alleles[1,3,14,16,17]; quinolone resistance has also been documented to

be transferred through transformation and homologous recombination[5]. Ciprofloxacin is a
quinolone class antibiotic, which though once recommended as a first-line therapy to treat
gonorrhea from the mid- late- 1980's, was discontinued in the 1990's due to widespread
resistance emergence[18]. The first clinical failures were reported by 1990 to low dose treatment
(250 mg)[19], which prompted a higher dose recommendation (500 mg), however resistance to
high-dose treatment evolved rapidly in Asia[20]. The rate of ciprofloxacin resistance ≥ 1 ug/ml still
remains high, with over 31% of U.S. isolates surveyed in 2022 classified as resistant[21].
Ciprofloxacin and other quinolone antibiotics target bacterial type II topoisomerases, DNA gyrase
and topoisomerase IV, which disrupts DNA synthesis[22]. Work in *N. gonorrhoeae* has
demonstrated that DNA gyrase subunit A (encoded by *gyrA*) is the primary target and
topoisomerase IV subunit C (encoded by *parC)* is the secondary target for fluoroquinolones[23–25].
Unsurprisingly, reduced susceptibility to ciprofloxacin in gonococci tends to be conserved and
emerge in the quinolone resistance-determining regions (QRDRs; codons 67–106 in GyrA and
56–108 in ParC) of the drug targets which reduce fluoroquinolone binding affinity[26–28]. Particular
amino acid positions associated with reduced susceptibility include 91 and 95 in GyrA; and 85,
86, 87, 88, 91, and 116 in ParC; with genomic haplotypes harboring both *gyrA* and *parC* mutations
displaying higher MICs than those with a single mutation[29]. Alternative pathways to ciprofloxacin
reduced susceptibility exist however, with recent work in gonococci highlighting the emergence
of mutations in DNA gyrase subunit B (GyrB D429N and P739H) and topoisomerase IV subunit
E (ParE P456S) during *in vitro* selection[30,31].
Acquisition of ciprofloxacin reduced susceptibility in both pathogenic *Neisseria* species
has been associated with the inheritance of alleles from the commensals. For example, recent
spread of commensally-acquired *gyrA* and *parC* haplotypes in meningococcal populations has
played a major role in the establishment of quinolone resistant meningococci (QRM) globally[5,32,33].
The first records of commensally acquired quinolone resistance in meningococci are from the
United States in 2007 and 2008. Three cases of meningococci from serogroup B were identified
as ciprofloxacin-resistant (with MICs of 0.25 µg/mL) in two states (North Dakota and Minnesota)[34].
Currently the percentage of QRM with mosaic alleles is high, with over 53.4% and >70% of
resistant isolates in the U.S. and China harboring mosaic alleles at *gyrA* and *parC*[5]. Furthermore,
a global *in silico* analysis of >20,000 *Neisseria* isolates demonstrated that alleles at *gyrA*, *gyrB*,
and *parE* have been transferred from commensal *Neisseria* to *N. gonorrhoeae*[35]. These examples
underscore the importance of commensals in transferring quinolone resistance, and stress the
importance of characterizing the quinolone-resistance conferring mutations that commensals may
harbor.
Given the documented cross-species spread of quinolone resistance in natural *Neisseria*
populations[5,35], we set out to characterize the mutations which impart ciprofloxacin resistance in
*Neisseria* commensals using *in vitro* selection. Here, we passage four species of commensals
using a selective gradient of ciprofloxacin for 20 days, and characterize derived mutations post-
selection. As in our prior work[36,37], we query the identity and number of paths to reduced
susceptibility, and compare and contrast these paths across different commensal genomic
backgrounds. We also test for the potential for the emergence of zoliflodacin cross-resistance.
Zoliflodacin is a spiropyrimidinetrione antibiotic which also binds type II topoisomerases, and has
recently completed phase III trials for treatment of gonorrhea[38–40]. Though the mode of inhibition
and binding sites are distinct from those of the fluoroquinolones[41], we hypothesized that i)
mutations impacting topoisomerase conformation or ii) general influx or efflux related mutations
may emerge that could result in ciprofloxacin producing cross-resistance. Finally, we interrogate
the presence of selected mutations in natural populations of *Neisseria*, and search for evidence
of HGT from commensals to gonococci.
99
**Methods**

***Bacterial Isolates***
In this study, the following bacterial strains were examined; AR-0944 (*N. cinerea*), AR-
0945 (*N. elongata*), AR-0948 (*N. canis*), and AR-0953 (*N. subflava*), which were acquired from
the Centers for Disease Control and Prevention (CDC) and Food and Drug Association's (FDA)
Antibiotic Resistance (AR) Isolate Bank "*Neisseria* species MALDI-TOF Verification panel". These
strains were cultured on GC agar base (Becton Dickinson Co., Franklin Lakes, NJ, USA) media
plates along with 1% BBL IsoVitaleX Enrichment (Becton Dickinson Co.,Franklin Lakes, NJ, USA;
hereafter referred to as GCB-I media). These strains were grown for 18–24 hours at 37°C in a 5%
$CO_2$ incubator. Ancestral strain MIC values to ciprofloxacin were previously reported[42] (Table 1),
and confirmed prior to *in vitro* selection experiments. Stocks were stored at -80°C in a solution
containing trypticase soy broth (TSB) and 20% glycerol.
112
113 **Table 1. Ancestral ciprofloxacin MICs for strains used in this study**

| Strain | MIC (μg/mL) |
| --- | --- |
| AR-0944 (*N. cinerea*) | 0.032 |
| AR-0945 (*N. elongata*) | 0.25 |
| AR-0948 (*N. canis*) | 0.008 |
| AR-0953 (*N. subflava*) | 0.75 |

114
***In vitro selection of resistance to ciprofloxacin and antimicrobial susceptibility testing***
The selection for ciprofloxacin resistance was performed via passaging four replicates of
each of the aforementioned commensal strains in the presence of an ETEST strip to introduce a
selective antibiotic gradient. In short, cells on the interior and from a 1 cm perimeter on the zone
of inhibition (ZOI) were collected and suspended in TSB.  This solution was then spread onto a
fresh GCB-I plate, and incubated at 37°C with 5% $CO_2$. 48 hours later, minimum inhibitory
concentrations (MICs) were observed and recorded.  The repetition of this procedure occurred
for 11 passages over 21 days for each replicate.  Bacterial populations from each replicate were
passaged in the absence of antibiotic and struck out to collect single colonies, which were stocked
for future analysis.  Control strains were also passaged in the absence of ciprofloxacin with the
same procedure.
The MIC values for the aforementioned post-selection individual colony stocks were
measured with ETEST strips, which have generated comparable MIC results to the agar dilution
method[43]. Simply put, bacterial cells from plates incubated overnight were suspended in TSB to
a 0.5 McFarland standard, and then spread onto a GCB-I plate along with an Etest strip.  After an
18–24 hour incubation period in the aforementioned conditions, MIC values were recorded and
reduced susceptibility was determined using Clinical & Laboratory Standards Institute (CLSI)
guidelines for *N. gonorrhoeae* of ≥ 1 μg/mL[44]. The final MIC values reported are the mode of three
independent MIC tests and commonly agreed upon by two independent researchers.
134
***Genome sequencing and bioinformatics analysis for selected isolates***
Bacterial cells from overnight plates were lysed via suspension in TE buffer (10 mM Tris
[pH 8.0], 10 mM EDTA) with 0.5 mg/mL lysozyme and 3 mg/mL proteinase K (Sigma-Aldrich
Corp., St. Louis, MO).  DNA isolation was performed with PureLink Genomic DNA Mini Kit
(Thermo Fisher Corp., Waltham, MA) with RNase A treatment for the removal of unwanted RNA.
Isolated DNA was organized for subsequent sequencing utilizing the Nextera XT kit (Illumina
Corp., San Diego, CA), and uniquely dual-indexed and pooled. The finalized pool was then

sequenced using the Illumina MiSeq platform at the Rochester Institute of Technology Genomics
Core using V3 600 cycle cartridges (2x300bp).
The quality of each paired-end read library was analyzed using FastQC v0.11.9. Any
adaptor sequences or low-quality sequences were trimmed from read libraries using Trimmomatic
v0.39, using a phred quality score < 15 over a 4 bp sliding window as a cutoff. Reads shorter than
36 bp, and those missing a mate, were removed from subsequent analysis as well. The reference
genomes were assembled and annotated previously[36,37,42], and the read libraries were mapped
back to these reference assemblies using Bowtie2 v.2.2.4 using the "end-to-end" and "very-
sensitive" options[45]. Ultimately, Pilon v.1.16[46] identified any derived mutations; and custom scripts
were used to determine if these mutations were located in any previously annotated genomic
features. A common amino acid residue number scheme across species was determined by
alignment to the NZ_AP023069.1 gonococcal reference genome.
154
***Testing for zoliflodacin cross-resistance***
The presence of cross-resistance to zoliflodacin in the ciprofloxacin-selected replicates
was determined experimentally using agar dilution. In brief, GC agar base with 1% IsoVitaleX was
supplemented with zoliflodacin to produce plates containing 0.0, 0.06, 0.125, 0.25, 0.5, 1.0, 2.0,
4.0, and 8.0 µg/mL of drug. Bacterial cells from overnight plates were collected and suspended
in 1 mL TSB and spotted on plates. Plates were incubated at 37°C with 5% $CO_2$ for 16 hours, at
which time MICs were recorded. Three independent MIC readings were recorded for each strain,
with the mode reported as the final MIC. While the resistance breakpoint for species of *Neisseria*
in the presence of zoliflodacin has not yet been defined, we use a prior published breakpoint of
MIC of ≥ 4 µg/mL to define resistance[47].
165
***Comparative population analysis of the presence of derived mutations and HGT in natural***
***Neisseria populations***
Previous work by Manoharan-Basil *et al.* (2022) has demonstrated alleles of *gyrA*, *gyrB*,
and *parE* have been transferred from commensal *Neisseria* to *N. gonorrhoeae* in a population
dataset from the 1920's to 2020[48]. Here, we reevaluate cross-species sharing of these and *parC*
alleles using isolates deposited to the PubMLST database (https://www.pubmlst.org/Neisseria)[49],
but focusing on the more recent depositions from 2020 through 2024. We specifically interrogate
the presence of experimentally derived mutations in this dataset from our *in vitro* selection
approach, and also search for any evidence of HGT of these particular alleles if they were present.
2235 *gyrA* (NEIS1320)*,* 2302 *gyrB* (NEIS0204)*,* 2287 *parE* (NEIS1600)*,* and 2256 *parC*
(NEIS1525) alleles were downloaded from published genomes deposited to PubMLST, including
all human-associated commensal *Neisseria*: *N. benedictiae, N. bergeri, N. blantyrii, N. cinerea,*
*N. elongata, N. lactamica, N. maigaei, N. mucosa, N. oralis, N. polysaccharea, N. subflava, N.*
*uirgultaei* and *N. viridiae*. *N. gonorrhoeae* was included as the focal pathogen in this analysis,
however *N. meningitidis* was excluded*.* We limited the majority of the gonococcal dataset to the
last five years (2020-2024) to capture more recent commensal/pathogen transfer events, though
included commensals isolates from all years, and added some earlier *N. gonorrhoeae* isolates of
interest. Alleles for each gene were aligned using MAFFT v.7 with a UPGMA tree calculated[50].
iTOL was used to visualize and annotate the resultant trees[51]. Multispecies phylogenies were
assessed for admixture using the *genealogical sorting index* (*gsi*) with the *genealogicalSorting* R
package[52,53].
187
188 **Results**
189 ***Phenotypic results of In vitro selection***
190    The four commensal species selected to undergo experimental evolution (AR-0944 [*N.*
191 *cinerea*], AR-0945 [*N. elongata*], AR-0948 [*N. canis*], and AR-0953 [*N. subflava*]), represented

evolutionarily distinct starting points with known MIC values, as outlined in Table 1. Four replicate lineages of each of the four commensal species underwent the aforementioned passaging procedure. In response to ciprofloxacin selection, the average MICs of each species are as follows; *N. cinerea* 7.5 ± 3.4 µg/mL, *N. elongata* 1.38 ± 0.88 µg/mL, *N. canis* 0.5 ± 0.68 µg/mL, and *N. subflava* 12 ± 4.62 µg/mL (Figure 1; Supplementary Figure 1). The MIC values of 10 evolved lines surpassed the resistance breakpoint of ≥ 1 µg/mL as outlined by CLSI guidelines. The smallest changes in MIC were observed in *N. canis* and *N. elongata*, while all experimental replicates for *N. cinerea* and *N. subflava* surpassed the resistance breakpoint.

**Table 2. Characterization of MICs for ciprofloxacin-selected lineages**

| | Ciprofloxacin MICs (µg/mL) | | | |
|---|---|---|---|---|
| Species | Replicate 1 | Replicate 2 | Replicate 3 | Replicate 4 |
| AR-0944 | 4 | 12 | 8 | 6 |
| AR-0945 | DNG* | 2 | DNG* | 0.75 |
| AR-0948 | 1.5 | 0.19 | 0.032 | 0.19 |
| AR-0953 | 16 | 8 | 8 | 16 |

**\*DNG=Single colony picks did not grow.**

### *Comparative genomics and nominating derived mutations*

Selected lineages were whole genome sequenced (WGS) and aligned to the draft assemblies reported previously[36,37,42] to nominate derived polymorphisms. Mutations shared with control strains (with no drug selection) were omitted from subsequent analyses. 717 derived mutations were identified across all sequenced strains, averaging 44.81± 33.15 per evolved lineage. The most frequent mutation occurring in *N. cinerea* lineages was located within the *23s rDNA*, the genomic region encoding the 23s rRNA (present in 66% of replicates). Additional derived mutations of note included those in *parE* encoding one of the topoisomerase IV subunits (33%), *gyrA* encoding the DNA gyrase A subunit (33%), and *pilF,* an ATPase that promotes pilus elongation (33%; Figure 2; see Supplementary Table 1 for complete list). For, *N. elongata* the most frequent mutations were in *tdfG,* a TonB-dependent transporter (100%) and the region encoding the transfer RNA tRNA-Val(tac) (100%), followed by those in regions encoding the major outer membrane porin (*porB;* 75%), a component of the pilus involved in retraction (*pilC;* 75%), the repressor of the *mtr efflux pump* (*mtrR;* 25%), the N-acetylmuramyl-l-alanine amidase (*amiC;* 25%), and aconitate hydratase 2/2 methylisocitrate dehydratase (*acnB;* 25%) (Figure 2). For *N. canis* the most frequent mutations were in *gyrA* (75%), followed by those in *parC* encoding one of the topoisomerase IV subunits (25%), the a sensor-histidine kinase/response regulator (*ntrX;* 25%), and DNA gyrase subunit B (*gyrB*; 25%) (Figure 2). For *N. subflava* the most frequent mutations were in *porB* (100%) and *mtrR* (100%); with additional mutations of note in pilB, a cytoplasmic NTPase involved in modulating pilus extension and retraction (75%), *parC* (50%), and *parE* (25%; Figure 2; see Supplementary Table 1 for complete list).

For particular loci of note, mutations that emerged included: A R320S substitution in topoisomerase IV subunit E and a T91I substitution in DNA gyrase subunit A in *N. cinerea*; a T11N substitution in MtrR in *N. elongata*; substitutions in DNA gyrase subunit A (G89D, S91N, S91R), DNA gyrase subunit B (P739H), and topoisomerase IV subunit C (G85C) for N. canis; and MtrR (*504insA, 300delA*, T93P, G45S), topoisomerase IV subunit C (A123E, 477 insertion K)

233 **Table 3. Derived mutations emerging in selected commensal lineages, compared to known**
234 **ciprofloxacin resistance-contributing loci in *N. gonorrhoeae***

| Loci contributing to resistance *N. gonorrhoeae* | Specific mutations in *N. N. gonorrhoeae* | Derived mutations in commensals | | | |
|---|---|---|---|---|---|
| | | *N. canis* | *N. cinerea* | *N. elongata* | *N.subflava* |
| *gyrA* | S91F, D95A/N/G | G89D*, S91N*, S91R* | T91I | | |
| *gyrB* | D429N | P739H* | | | |
| *mtrR* | A39T, G45D, Y105H | | | T11N* | 504ins-a†, 300del-a†, T93P, G45S |
| *parC* | D86N, S87R/N, E91K | G85C* | | | A123E, 477insK |
| *parE* | G410V | | R320S | | 646insR |
| *porB* | E127G | | | | KO-indels† |

235 *Strains evolving these mutations did not cross the reduced susceptibility breakpoint of ≥ 1 µg/mL.

236 -Amino acid residue numbering scheme is based on alignment to the gonococcal reference
237 genome (NZ_AP023069.1).
238 †Putative DNA-level insertions or deletions of nucleotides.
239
### Testing for zoliflodacin cross-resistance in evolved cell lines
We tested evolved lineages for the emergence of cross-resistance to zoliflodacin after
ciprofloxacin selection. Though no strains displayed large changes in zoliflodacin MICs compared
to ancestral lineages (Table 4): 1 *N. elongata* replicate had an elevated MIC (from 1 to 4 µg/mL),
1 *N. canis* replicate had an elevated MIC (from 1 to 4 µg/mL), and all 4 *N. subflava* lineages had
elevated MIC (from 2 to 4 µg/mL). For the *N. elongata* isolate with an elevated zoliflodacin MIC,
it was the only lineage to inherit a MtrR T11N substitution post-selection. For the *N. canis* isolate
with an elevated zoliflodacin MIC, it was the only to inherit GyrA G89D and ParC G85C mutations.
Finally, for the *N. subflava* lineages, all inherited a MtrR mutation (300delA, T93P, or G45S);
coupled with *porB* indels which altered the reading frames; with 3 out of the 4 lineages also
inheriting either a K insertion at position 477 in ParC, ParC A123E, or a R insertion at position
646 in ParE.
252
253 **Table 4. Characterization of zoliflodacin MICs for ciprofloxacin-selected lineages**

| | Zoliflodacin MICs (µg/mL) | | | | |
|---|---|---|---|---|---|
| Species | Ancestral | Replicate 1 | Replicate 2 | Replicate 3 | Replicate 4 |
| AR-0944 | 2 | 2 | 1 | 1 | 0.25 |
| AR-0945 | 1 | DNG* | 1 | DNG* | 4 |
| AR-0948 | 1 | 4 | 1 | 1 | 1 |
| AR-0953 | 2 | 4 | 4 | 4 | 4 |

254 **\*DNG=Single colony picks did not grow.**
255
### Population analysis of derived mutations and HGT in natural Neisseria populations
We searched for evidence of *in vitro* evolved mutations and gene-level recombination
tracts in natural *Neisseria* populations using *Neisseria* isolates deposited to the PubMLST
database. For the *gyrA* locus, we observed an *in vitro* selected mutation at position 89 (G89D);
however, we found no evidence of amino acid substitutions at this position in natural populations
of *Neisseria,* with all isolates (n=2231) harboring a glycine at that position. At position 91 however,
we find a diversity of alleles encoding different amino acids in natural *Neisseria* populations with:
1 isolate harboring an alanine (A), 489 with a phenylalanine (F), 70 with an isoleucine (I), 1 with
a leucine (L), 531 with a serine (S) 1128 with a threonine (T), 10 with a valine (V), and 1 with a
tyrosine (T) (Figure 3). Of these mutations, 1 was uncovered in our experimental evolution
experiments 91I and was associated with elevated ciprofloxacin MICs. 91I in natural populations
of *Neisseria* was found in *N. lactamica*, *N. subflava*, and *N. polysaccharea* (Figure 3). We find no
evidence of 91F in commensals which is associated with ciprofloxacin resistance in gonococci
(Figure 3). Though we did not find any evolved mutations at position 95 in GyrA in our *in vitro*
selection, we characterized the amino acid diversity across natural populations as D95A/N/G
substitutions increase ciprofloxacin MICs in gonococci. Here we find, 369 isolates harboring an
alanine (A), 1708 with a aspartic acid (D), 118 with a glycine (G), 1 with a histidine (H), 25 with an
asparagine (N), and 10 with a tyrosine (Y) (Figure 3). Reduced susceptibility-associated amino

acids were found in *N. lactamica*, *N. subflava*, *N. polysaccharea* and *N. bergeri*. We find no
evidence of admixture between commensals and gonococci at the locus-level for *gyrA* however
in this analysis, with gonococci belonging to a monophyletic clade (*gsi* = 1).

In our selection experiment we uncovered a GyrB P739H substitution emerge post-
selection in *N. canis*. In natural *Neisseria* populations however, we did not find any evidence of
this mutation, with all isolations (n=2301) homozygous for a proline (P) at this position (Figure 4).
Additionally, we searched for the D429N substitution that is associated with zoliflodacin in
gonococci[31]. All isolated harbored an aspartic acid (D) at this position however, with no variation
observed (Figure 4). Finally, we find no evidence of commensal to pathogen allelic exchange at
the *gyrB* locus level (gonococcal *gsi* = 1).

At *parC*, a mutation encoding a G85C substitution emerged in *N. canis,* and A123E and
477 insertion K substitutions emerged in *N. subflava* after selection. In natural *Neisseria*
populations we find no variation at position 126 or 477. We next investigated the positions 85, 86,
87, and 91; as we found post-selection variation after experimental evolution at position 85, and
all three of the later residues are associated with ciprofloxacin resistance in gonococci. Here we
find: 2231 isolates with a glycine (G), 23 with a serine (S), 1 with a cysteine (C), and one with an
aspartic acid (D) at position 85; 2181 isolates harboring an aspartic acid (D), and 75 with a
asparagine (N) at residue 86; 1911 with a serine (S), 318 with an arginine (R), 21 with an
asparagine (N), and 6 with an isoleucine (I) at position 87; and 2208 with a glutamic acid (E), 33
with a glycine (G), 3 with a lysine (K), and 12 with a glutamine (Q) at position 91 (Figure 5). For
*parC*, we also identify a putative gene-level HGT event, with a clade of three gonococcal isolates
from the Netherlands collected in 2021 harboring more *N. polysaccharea* or *N. lactamica*-like
alleles (Figure 5). All three isolates had haplotypes of 86D 87S 91E. The 86N, 87R/N, and 91K
mutations associated with ciprofloxacin resistance in gonococci were limited to the gonococcal
clade and were not present in the commensal *Neisseria* (Figure 5). We also identify a single
gonococcal isolate collected in China with the ParC G85C substitution that emerged in *N. canis*
post-selection (Figure 5).

For ParE, substitutions emerging after ciprofloxacin selection included R320S in *N.*
*cinerea* and 646 R insertion in *N. subflava*. We found no evidence of variation at position 320 in
natural Neisseria populations, with all isolates (n=2286) harboring a R. We also looked for
evidence of variation at position 410 which has been associated with x resistance in gonococci,
however we also find no variation at this site in natural *Neisseria* populations. We do however find
HGT of commensal Neisseria *parE* alleles into a cluster of four gonococcal isolates from Scotland
isolated in 2023-24 (Figure 6).

**Discussion**
Cross-species transfer of alleles harboring ciprofloxacin resistance-associated mutations
has been observed in natural *Neisseria* populations[5,48], where commensals serve as the reservoir
for pathogen evolution. Thus, characterizing the mechanisms that impart ciprofloxacin resistance
in *Neisseria* commensals using *in vitro* selection may highlight important mutations to consider
for gonococcal surveillance programs and may also illuminate novel resistance-conferring
mutations that can arise in commensal genomic backgrounds. Previous work using *in vitro*
evolution has pointed to conserved mechanisms of resistance across the genus for azithromycin
and penicillin-based selection[36,37,42], here we expand our enquiry to ciprofloxacin for several
*Neissiera* commensals, and interrogate both the identity and diversity of paths to resistance
across different genomic backgrounds.

Ciprofloxacin, a quinolone-class antibiotic, targets the bacterial type II topoisomerases
DNA gyrase and topoisomerase IV, and disrupts DNA synthesis; therefore it is no surprise that in
gonococci, reduced susceptibility is often mediated by mutations in the primary and secondary
drug targets, GyrA and ParC respectively[12,20,24,25]. In particular, mutations falling within the
quinolone resistance-determining regions (QRDRs) of GyrA and ParC (codons 91-95 of GyrA and

86-91 of ParC) have been shown to impact ciprofloxacin MICs, with the substitutions S91F and
D95A/N/G in GyrA and D86N, S87R/N, and E91K ParC strongly associated with elevated MICs
in natural gonococcal populations[12,31]. Here in *Neisseria* commensals, we find amino acid-altering
mutations in these two loci: G89D, S91N, S91R, and T91I in GyrA; and G85C, A123E, and a K
insertion at position 477 in ParC (Table 3); the majority of which fall within the aforementioned
QRDRs. Of these mutations T91I has been found in quinolone resistant meningococci with gyrA
alleles that had been inherited from commensal *Neisseria* donors (*N. lactamica*, *N. cinerea*, and
*N. subflava*), and is associated with elevated ciprofloxacin MICs in conjunction with ParC
mutations[5]. Here, we find this 91I mutation in natural commensal populations of *Neisseria* (*N.
lactamica*, *N. subflava*, and *N. polysaccharea*; n=70 or ~ 3% of isolates*)* suggesting its availability
for HGT and resistance transfer, however we do not see this mutation in contemporary gonococci
(Figure 3). Finally, we also find the ParC G85C substitution which emerged post-selection in a
single gonococcal isolate. The isolate was collected in China (ciprofloxacin MIC of 2 µg/mL) and
had resistance-associated mutations at GyrA 91F and 95A, however had wildtype alleles at ParC
at other resistance associated sites (86D, 87S, 91E). Thus, we believe the ParC G85C
substitution is acting together with GyrA 91F and 95A to produced elevated ciprofloxacin MICs in
this strain.
We also find substitutions that emerged in the other subunits of DNA gyrase and
topoisomerase IV: GyrB P739H, ParE R320S, and ParE a R insertion at position 646. There was
no variation at these sites in the natural commensal or gonococcal populations included within
this analysis (Figure 4 and Figure 6). The GyrB P739H has been associated with intermediate
ciprofloxacin resistance (MICs: 0.032 to 0.125 mg/L) in gonococci in the laboratory, and has also
been observed in two commensal species that were not included in our population analysis (*N.
brasiliensis* (PATRIC ID-2666100.4) and an unidentified *Neisseria* isolate (PubMLST ID-
94179))[30]. Low rates of carriage likely suggest some fitness cost to maintenance of GyrB P739H,
though follow up experiments will be necessary to understand the impact on fitness for all
uncovered mutations.
Investigation of the haplotypes of each evolved lineage can help us to understand which
mutations may be contributing to observed ciprofloxacin resistance. For *N. cinerea*, though we
were unable to identify the mutations contributing to resistance in 3 strains, the fourth replicate
inherited both gyrA T91I and parE R320S (MIC = 6 µg/ml; Supplemental Table 2). We suspect
that both of these mutations contribute together to the high ciprofloxacin MICs in this strain, as in
meningococci both this mutation and ParC substitutions (D86N, S87R/I, E91G) must be inherited
together to produce elevated ciprofloxacin MICs[5]. In *N. canis*, a similar pattern exists where GyrA
or GyrB substitutions only marginal increase ciprofloxacin MICs compared to the ancestral strain,
however the strain inheriting both GyrA G89D and ParC G85C had a MIC above the breakpoint
(MIC = 1.5 µg/ml; Supplemental Table 2). For *N. subflava*, the largest contributors to elevated
MICs are less clear, with all strains inherited 2-3 likely contributors; however we speculate the
combination of reduced drug influx from porin knockouts and increased drug efflux from *mtrR*

mutations are likely causing the high MICs observed in these strains (≥ 8 µg/ml).

Emergence of ciprofloxacin-selected mutations in type II topoisomerases may be
concerning given the current development of zoliflodacin as a new treatment for gonorrhea.
Zoliflodacin has a novel target (GyrB) and mode of action compared to ciprofloxacin[41], however
we questioned if zoliflodacin cross-resistance could emerge after ciprofloxacin selection given
that mutations emerged in GyrB post-ciprofloxacin selection. Prior mutations associated with
elevated zoliflodacin MICs include GyrB D429V, GyrB S467N, and GyrB K450[31,54] which did not
emerge here (Table 3), however we wanted to test the impact of the derived GyrB P739H mutation
on zoliflodacin MIC. Interestingly, the *N. cinerea* strain that inherited GyrB P739H did not have a
higher zoliflodacin MIC than the ancestral strain; yet other strains from our selection experiment
did (Table 4). These strains included: A *N. elongata* strain (MIC increase from 1 to 4 µg/ml)

inheriting a MtrR T11N substitution, a *N. canis* strain (MIC increase from 1 to 4 µg/ml) inheriting
GyrA G89D and ParC G85C substitutions, and all four *N. subflava* strains (MIC increase from 2
to 4 µg/ml) which had a combination of influx, efflux, and ParC/ParE mutations (Table 4 and
Supplementary Table 2). Upon investigating the haplotypes of these strains ParC and ParE
mutations appear to have a limited impact on zoliflodacin MICs in *N. subflava* (Supplementary
Table 2); thus, we speculate the majority of cross-resistance observed here is emerging through
generalized mechanisms (i.e., drug influx or efflux), aside from the single *N. canis* isolate with the
GyrA G89D and ParC G85C substitutions. As GyrB variation in natural *Neisseria* populations is
rare ([54] and Figure 4), future studies may consider testing the impact of naturally occurring efflux
pump and porin mutations on zoliflodacin MIC when assessing the current and future potential for
resistance emergence rather than focusing on variation in the drug target alone.
In conclusion, this work again highlights the potential of commensal *Neisseria* in serving
as reservoirs of resistance for their pathogenic relatives. We find evidence of recent transfer of
*parC* and *parE* genes from commensals to pathogens through the observation of admixed
phylogenies (Figure 5 and 6). We also characterize mutations associated with ciprofloxacin
reduced susceptibility in commensals after selection, and find both previously described and novel
mutations emerging in the drug targets, the Mtr efflux pump system, and the major outer
membrane porin. Of these mutations we identify GyrA 91I as a substitution circulating within
commensal *Neisseria* populations, confirming prior results[5]; and a ParC G85C present in a single
ciprofloxacin-resistant gonococcal isolate. Finally, we observe cross-resistance to zoliflodacin
emerge, and speculate drug influx and/or efflux mutations are the likely underlying contributors.
396

**Acknowledgements**
Research by C.B.W. is supported by the College of Science and Thomas H. Gosnell School of
Life Sciences at RIT and the National Institute of General Medical Sciences of the National
Institutes of Health under Award Number R15AI174182. The content is solely the responsibility
of the authors and does not necessarily represent the official views of the National Institutes of
Health.

403
404 **Statement**
405 Whole genome sequencing libraries can be accessed through the NCBI's Sequence Read
406 Archive (SRA) with the accession numbers: SAMN41424178-SAMN41424193.
407
**Figure Captions**
**Figure 1.** *In vitro* selection of four commensal Neisseria species with ciprofloxacin. In brief, four
experimental replicates of each species were passaged for 20 days on selective gradients created
with Etest strips. Cells for each passage were selected by sweeping the zone of inhibition (ZOI)
and a 1 cm band in the bacterial lawn surrounding the ZOI. MIC trajectories are shown for (A) *N.*
*cinerea*, (B) *N. elongata*, (C) *N. canis*, and (D) *N. subflava*. Expanded scales are shown in panels
E-G for *N. cinerea*, *N. elongata*, and *N. canis* respectively.
415
416 **Figure 2.** CDs identity of *in vitro* derived mutations for drug-selected lineages. The percentage of
417 mutational hits within a given gene is displayed as a heatmap, with brighter blue coloration
418 indicating more frequent occurrence of a mutation within a CDS in replicate evolved lineages for
419 each species.
420
421 **Figure 3.** The ciprofloxacin reduced susceptibility-associated mutation GyrA 91I circulates within
422 commensal *Neisseria* populations, however has not been transferred to *Neisseria gonorrhoeae*.
423 The coloring in the inner ring represents the *Neisseria* species, followed by amino acid identity at
424 GyrA positions 89, 91, and 95 respectively. The derived 91I mutation was found in *N. lactamica*,
425 *N. subflava*, and *N. polysaccharea* (n=70 or ~ 3% of isolates*).*

426
427 **Figure 4.** GyrB does not show variation in the zoliflodacin reduced susceptibility mutation (D429V)
428 or derived P739H mutation in commensal populations. UPGMA phylogeny of the *gyrB* locus for
429 2302 PubMLST *Neisseria* isolates. The coloring in the inner ring represents the *Neisseria* species,
430 followed by amino acid identity at GyrB positions 429 and 739 respectively.
431
**Figure 5.** ParC 86N, 87R/N, and 91K mutations associated with ciprofloxacin resistance in
gonococci were limited to the gonococcal clade and were not present in the commensal *Neisseria*;
however, a single gonococcal isolate inherited the experimentally-derived ParC G85C substitution
that emerged in *N. canis* post-selection. UPGMA phylogeny of the *parC* locus for 2256 PubMLST
*Neisseria* isolates. The coloring in the inner ring represents the *Neisseria* species, followed by
amino acid identity at ParC positions 86, 87, and 91 and respectively. The subpanel highlights 3
gonococcal isolates with commensal-like *parC* alleles.
439
440 **Figure 6.** UPGMA phylogeny of the *parE* locus for 2287 PubMLST *Neisseria* isolates. The
441 coloring in the inner ring represents the *Neisseria* species, followed by amino acid identity at ParE
442 positions 320 and 410 respectively. The subpanel highlights 4 gonococcal isolates with
443 commensal-like *parE* alleles.
444

**References**
1. Wadsworth, C. B., Arnold, B. J., Sater, M. R. A. & Grad, Y. H. Azithromycin Resistance

through Interspecific Acquisition of an Epistasis-Dependent Efflux Pump Component and

Transcriptional Regulator in Neisseria gonorrhoeae. *mBio* **9**, e01419-18 (2018).

2. Goytia, M. & Wadsworth, C. B. Canary in the Coal Mine: How Resistance Surveillance in

Commensals Could Help Curb the Spread of AMR in Pathogenic *Neisseria*. *mBio* **13**,

e01991-22 (2022).

3. Rouquette-Loughlin, C. E. *et al.* Mechanistic Basis for Decreased Antimicrobial

Susceptibility in a Clinical Isolate of Neisseria gonorrhoeae Possessing a Mosaic-Like *mtr*

Efflux Pump Locus. *mBio* **9**, e02281-18 (2018).

4. Spratt, Brian. Hybrid penicillin-binding proteins in penicillin-resistant strains of Neisseria

gonorrhoeae. *Nature* **332**, 173–176 (1988).

5. Chen, M., Zhang, C., Zhang, X. & Chen, M. Meningococcal Quinolone Resistance

Originated from Several Commensal *Neisseria* Species. *Antimicrob. Agents Chemother.* **64**,

e01494-19 (2020).

6.  Yee, W.-X., Elsener, T., Cehovin, A., Maiden, M. C. J. & Tang, C. M. Evolution and

exchange of plasmids in pathogenic *Neisseria*. *mSphere* e00441-23 (2023)

doi:10.1128/msphere.00441-23.

7.  Dong, H. V. *et al.* Decreased Cephalosporin Susceptibility of Oropharyngeal Neisseria

Species in Antibiotic-using Men Who Have Sex With Men in Hanoi, Vietnam. *Clin. Infect.*

*Dis.* **70**, 1169–1175 (2020).

8.  Kenyon, C. R. & Schwartz, I. S. Effects of Sexual Network Connectivity and Antimicrobial

Drug Use on Antimicrobial Resistance in *Neisseria gonorrhoeae*. *Emerg. Infect. Dis.* **24**,

1195–1203 (2018).

9.  Vanbaelen, T. *et al.* Global epidemiology of antimicrobial resistance in commensal Neisseria

species: A systematic review. *Int. J. Med. Microbiol.* **312**, 151551 (2022).

10. Keijser, B. J. F. *et al.* Pyrosequencing analysis of the Oral Microflora of healthy adults. *J.*

*Dent. Res.* **87**, 1016–1020 (2008).

11. Burcham, Z. M. *et al.* Patterns of Oral Microbiota Diversity in Adults and Children: A

Crowdsourced Population Study. *Sci. Rep.* **10**, 2133 (2020).

12. Grad, Y. H. *et al.* Genomic Epidemiology of Gonococcal Resistance to Extended-Spectrum

Cephalosporins, Macrolides, and Fluoroquinolones in the United States, 2000–2013. *J.*

*Infect. Dis.* **214**, 1579–1587 (2016).

13. Ohnishi, M. *et al.* Is Neisseria gonorrhoeae Initiating a Future Era of Untreatable

Gonorrhea?: Detailed Characterization of the First Strain with High-Level Resistance to

Ceftriaxone. *Antimicrob. Agents Chemother.* **55**, 3538–3545 (2011).

14. Demczuk, W. *et al.* Genomic Epidemiology and Molecular Resistance Mechanisms of

Azithromycin-Resistant Neisseria gonorrhoeae in Canada from 1997 to 2014. *J. Clin.*

*Microbiol.* **54**, 1304–1313 (2016).

15. Ameyama, S. *et al.* Mosaic-Like Structure of Penicillin-Binding Protein 2 Gene ( *penA* ) in

Clinical Isolates of *Neisseria gonorrhoeae* with Reduced Susceptibility to Cefixime.

*Antimicrob. Agents Chemother.* **46**, 3744–3749 (2002).

16. Whiley, D. M. *et al.* Azithromycin-resistant Neisseria gonorrhoeae spreading amongst men

who have sex with men (MSM) and heterosexuals in New South Wales, Australia, 2017. *J.*

*Antimicrob. Chemother.* **73**, 1242–1246 (2018).

17. Banhart, S. *et al.* The Mosaic *mtr* Locus as Major Genetic Determinant of Azithromycin

Resistance of *Neisseria gonorrhoeae* —Germany, 2018. *J. Infect. Dis.* **224**, 1398–1404

(2021).

18. Unemo, M. & Shafer, W. M. Antibiotic resistance in *Neisseria gonorrhoeae* : origin,

evolution, and lessons learned for the future. *Ann. N. Y. Acad. Sci.* **1230**, (2011).

19. Gransden, WR, Warren, CA, Phillips, I, Hodges, M, & Barlow, D. Decreased susceptibility of

Neisseria gonorrhoeae to ciprofloxacin. *Lancet* **335**,.

20. Tanaka, M. *et al.* Antimicrobial Resistance of *Neisseria gonorrhoeae* and High Prevalence

of Ciprofloxacin-Resistant Isolates in Japan, 1993 to 1998. *J. Clin. Microbiol.* **38**, 521–525

(2000).

21. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance

2022: Enhanced Gonococcal Isolate Surveillance Project Profiles. (2024).

22. Fàbrega, A., Madurga, S., Giralt, E. & Vila, J. Mechanism of action of and resistance to

quinolones. *Microb. Biotechnol.* **2**, 40–61 (2009).

23. Lindbäck, E., Rahman, M., Jalal, S. & Wretlind, B. Mutations in *gyrA* , *gyrB* , *parC* , and *parE*

in quinolone-resistant strains of *Neisseria gonorrhoeae*. *APMIS* **110**, 651–657 (2002).

24. Shultz, T. R., Tapsall, J. W. & White, P. A. Correlation of In Vitro Susceptibilities to Newer

Quinolones of Naturally Occurring Quinolone-Resistant *Neisseria gonorrhoeae* Strains with

Changes in GyrA and ParC. *Antimicrob. Agents Chemother.* **45**, 734–738 (2001).

25. Trees, D. L. *et al.* Alterations within the quinolone resistance-determining regions of GyrA

and ParC of Neisseria gonorrhoeae isolated in the Far East and the United States. *Int. J.*

*Antimicrob. Agents* **12**, 325–332 (1999).

26. Alcal??, B. *et al.* Molecular Characterization of Ciprofloxacin Resistance of Gonococcal

Strains in Spain: *Sex. Transm. Dis.* **30**, 395–398 (2003).

27. Bodoev, I. N. & Il'ina, E. N. Molecular mechanisms of formation of drug resistance in

Neisseria gonorrhoeae: History and prospects. *Mol. Genet. Microbiol. Virol.* **30**, 132–140

(2015).

28. Yahara, K. *et al.* Emergence and evolution of antimicrobial resistance genes and mutations

in Neisseria gonorrhoeae. *Genome Med.* **13**, 51 (2021).

29. Collins, J. A., Oviatt, A. A., Chan, P. F. & Osheroff, N. Target-Mediated Fluoroquinolone

Resistance in *Neisseria gonorrhoeae* : Actions of Ciprofloxacin against Gyrase and

Topoisomerase IV. *ACS Infect. Dis.* **10**, 1351–1360 (2024).

30. González, N. *et al.* Alternative Pathways to Ciprofloxacin Resistance in Neisseria

gonorrhoeae: An In Vitro Study of the WHO-P and WHO-F Reference Strains. *Antibiotics*

**11**, 499 (2022).

31. Rubin, D. H., Mortimer, T. D. & Grad, Y. H. Neisseria gonorrhoeae diagnostic escape from a

gyrA-based test for ciprofloxacin susceptibility and the effect on zoliflodacin resistance: a

bacterial genetics and experimental evolution study. *Lancet Microbe* **4**, e247–e254 (2023).

32. Tsang, RS, Raymond, SW, Dennis, KS, Deng, S, & Hoang, L. Ciprofloxacin-resistant

Neisseria meningitidis in Canada: likely imported strains. *Can. J. Microbiol.* **63**, 265–268.

33. Gorla, M. C. *et al.* Emergence of resistance to ciprofloxacin in Neisseria meningitidis in

Brazil. *J. Med. Microbiol.* **67**, 286–288 (2018).

34. Wu, H. M. *et al.* Emergence of Ciprofloxacin-Resistant Neisseria meningitidis in North

America. *N Engl J Med* (2009).

35. Manoharan-Basil, S. S. *et al.* Evidence of Horizontal Gene Transfer of 50S Ribosomal

Genes rplB, rplD, and rplY in Neisseria gonorrhoeae. *Front. Microbiol.* **12**, 683901 (2021).

36. Raisman, J. C. *et al.* Evolutionary paths to macrolide resistance in a Neisseria commensal

converge on ribosomal genes through short sequence duplications. *PLOS ONE* **17**,

e0262370 (2022).

37. Frost, K. M. *et al.* Rolling the evolutionary dice: *Neisseria* commensals as proxies for

elucidating the underpinnings of antibiotic resistance mechanisms and evolution in human

pathogens. *Microbiol. Spectr.* e03507-23 (2024) doi:10.1128/spectrum.03507-23.

38. Bradford, P. A., Miller, A. A., O'Donnell, J. & Mueller, J. P. Zoliflodacin: An Oral

Spiropyrimidinetrione Antibiotic for the Treatment of *Neisseria gonorrheae* , Including Multi-

Drug-Resistant Isolates. *ACS Infect. Dis.* **6**, 1332–1345 (2020).

39. Taylor, S. N. *et al.* Single-Dose Zoliflodacin (ETX0914) for Treatment of Urogenital

Gonorrhea. *N. Engl. J. Med.* **379**, 1835–1845 (2018).

40. Global Antibiotic Research. Positive Results Announced in Largest Pivotal Phase 3 Trial of

a First-in-Class Oral Antibiotic to Treat Uncomplicated Gonorrhoea. (2023).

41. Basarab, G. S. *et al.* Responding to the challenge of untreatable gonorrhea: ETX0914, a

first-in-class agent with a distinct mechanism-of-action against bacterial Type II

topoisomerases. *Sci. Rep.* **5**, 11827 (2015).

42. Fiore, M. A., Raisman, J. C., Wong, N. H., Hudson, A. O. & Wadsworth, C. B. Exploration of

the Neisseria Resistome Reveals Resistance Mechanisms in Commensals That May Be

Acquired by N. gonorrhoeae through Horizontal Gene Transfer. *Antibiotics* **9**, 656 (2020).

43. Liu, H., Taylor, T. H., Pettus, K. & Trees, D. Assessment of Etest as an Alternative to Agar

Dilution for Antimicrobial Susceptibility Testing of Neisseria gonorrhoeae. *J. Clin. Microbiol.*

**52**, 1435–1440 (2014).

44. Clinical and Laboratory Standards Institute. *M100: Performance Standards for Antimicrobial*

*Susceptibility Testing*. (2020).

45. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,

357–359 (2012).

46. Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection

and Genome Assembly Improvement. *PLoS ONE* **9**, e112963 (2014).

47. Abdellati, S. *et al.* Gonococcal resistance to zoliflodacin could emerge via transformation

from commensal Neisseria species. An in-vitro transformation study. *Sci. Rep.* **14**, 1179

(2024).

48. Manoharan-Basil, S. S., González, N., Laumen, J. G. E. & Kenyon, C. Horizontal Gene

Transfer of Fluoroquinolone Resistance-Conferring Genes From Commensal Neisseria to

Neisseria gonorrhoeae: A Global Phylogenetic Analysis of 20,047 Isolates. *Front. Microbiol.*

**13**, 793612 (2022).

49. Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res.* **3**,

124 (2018).

50. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

51. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res.* **49**, W293–W296 (2021).

52. Cummings, M. P., Neel, M. C. & Shaw, K. L. A GENEALOGICAL APPROACH TO

QUANTIFYING LINEAGE DIVERGENCE. *Evolution* **62**, 2411–2422 (2008).

53. R Core Team. R: A language and environment for statistical computing. *R Found. Stat.*

*Comput. Vienna Austria 2015 Available HttpswwwR-Proj.*

54. Golparian, D. *et al.* GyrB *in silico* mining in 27 151 global gonococcal genomes from 1928–

2021 combined with zoliflodacin *in vitro* testing of 71 international gonococcal isolates with

different GyrB, ParC and ParE substitutions confirms high susceptibility. *J. Antimicrob.*

*Chemother.* **78**, 150–154 (2023).

**SPECIES**

- Neisseria gonorrhoeae
- Neisseria cinerea
- Neisseria lactamica
- Neisseria subflava
- Neisseria benedictiae
- Neisseria bergeri
- Neisseria blantyrii
- Neisseria mucosa
- Neisseria oralis
- Neisseria polysaccharea
- Neisseria viridiae
- Neisseria elongata

**gyrA89**

- G

**gyrA91**

- A
- F
- I
- L
- S
- T
- V
- Y

**gyrA95**

- A
- D
- G
- H
- N
- Y

Tree scale: 0.1

Tree scale: 0.1

SPECIES

- Neisseria gonorrhoeae
- Neisseria cinerea
- Neisseria lactamica
- Neisseria subflava
- Neisseria benedictiae
- Neisseria bergeri
- Neisseria blantyrii
- Neisseria mucosa
- Neisseria oralis
- Neisseria polysaccharea
- Neisseria viridiae
- Neisseria elongata
- Neisseriamaigaei

gyrB429

- D

gyrB739

- P

SPECIES

- Neisseria gonorrhoeae
- Neisseria cinerea
- Neisseria lactamica
- Neisseria subflava
- Neisseria benedictiae
- Neisseria bergeri
- Neisseria blantyrii
- Neisseria mucosa
- Neisseria oralis
- Neisseria polysaccharea
- Neisseria viridiae
- Neisseria elongata
- Neisseria maigaei

**parC86**

- D
- N

**parC87**

- S
- R
- N
- I

**parC91**

- E
- G
- K
- Q

Tree scale: 0.1

**SPECIES**

- Neisseria gonorrhoeae
- Neisseria cinerea
- Neisseria lactamica
- Neisseria subflava
- Neisseria benedictiae
- Neisseria bergeri
- Neisseria blantyrii
- Neisseria mucosa
- Neisseria oralis
- Neisseria polysaccharea
- Neisseria viridiae
- Neisseria elongata
- Neisseria maigaei

**parE320**
- R

**parE410**
- G