Title

A proteome-wide structural systems approach reveals insights into protein families of all human

herpesviruses

Authors

Timothy K. Soh[1, 2, 3, 4], Sofia Ognibene[1, 2, 3, 4], Saskia Sanders[1, 2, 3, 4], Robin Schäper[1, 2, 3, 4], Benedikt B.

Kaufer[5, 6], and Jens B. Bosse[1, 2, 3, 4, *]

[1] Centre for Structural Systems Biology, Hamburg, Germany

[2] Hannover Medical School, Institute of Virology, Hanover, Germany

[3] Cluster of Excellence RESIST (EXC 2155), Hanover Medical School, Hanover, Germany

[4] Leibniz Institute of Virology (LIV), Hamburg, Germany

[5] Institut für Virologie, Freie Universität Berlin, Robert von Ostertag-Straße 7-13, 14163 Berlin, Germany

[6] Veterinary Centre for Resistance Research (TZR), Freie Universität Berlin, 14163 Berlin, Germany

* To whom correspondence should be addressed. Tel: +49 040 8998 87645; Email: jens.bosse@cssb-

hamburg.de

Abstract

Structure predictions have become invaluable tools, but viral proteins are absent from the EMBL/DeepMind

AlphaFold database. Here, we provide proteome-wide structure predictions for all nine human

herpesviruses and analyze them in depth with explicit scoring thresholds. By clustering these predictions

into structural similarity groups, we identified new families, such as the HCMV UL112-113 cluster, which is

conserved in alpha- and betaherpesviruses. A domain-level search found protein families consisting of

subgroups with varying numbers of duplicated folds. Using large-scale structural similarity searches, we

identified viral proteins with cellular folds, such as the HSV-1 US2 cluster possessing dihydrofolate

reductase folds and the EBV BMRF2 cluster that might have emerged from cellular equilibrative nucleoside

transporters. Our HerpesFolds database is available at https://www.herpesfolds.org/herpesfolds and

displays all models and clusters through an interactive web interface. Here, we show that system-wide

structure predictions can reveal homology between viral species and identify potential protein functions.

Introduction

The *Orthoherpesviridae* family comprises alpha-, beta-, and gammaherpesviruses containing nine human herpesviruses (see Table 1 for virus abbreviations and their respective subfamilies). These are medically important human pathogens, including viruses such as human cytomegalovirus (HCMV), the leading infectious cause of congenital disorders in the USA. Due to the large number of encoded genes (between 70 and 181, depending on the species), experimental structural information for most viral proteins is not available, severely hampering their functional analysis. Moreover, herpesvirus gene and protein nomenclature are very inconsistent between species. Even worse, some proteins of the same species have multiple names based on their genome position, molecular weight, expression in the infected cell, or packaging into the virion. This situation makes it difficult even for experts to draw parallels between viruses. While some corresponding orthologues between species can be deduced from the UniProtKB annotation, this database is incomplete and sometimes inconsistent. Finally, herpesviruses also have a penchant for duplicating genes, further complicating annotation.

With the development of machine learning-based prediction software[1,2], the accuracy of the structural information has increased so that it can often be directly used for building wet-lab testable hypotheses, making it an invaluable tool for molecular biologists[3]. DeepMind, in cooperation with EMBL-EBI, has published over 200 million protein structures to facilitate this process, covering most of UniProtKB in an open-access database (https://alphafold.ebi.ac.uk). While this is an extraordinary resource for many fields, it currently excludes viral proteins such as the human herpesvirus proteomes.

Herpesvirus phylogeny has classically been sequence-based, such as a recent study that classified the human herpesvirus proteomes into homology families[4]. However, the current revolution in structure prediction and rapid structural similarity search algorithms[5] has vastly expanded the pool of sequences that can be interrogated by structure-based phylogeny[6-8]. When applied to large proteins or large phylogenetic distances, *e.g.* across the *Duplodnaviria* realm, this approach clustered species more consistently than the sequence-based ones[9]. Moreover, a recent study using this approach showed that orthopoxviruses acquired cellular proteins, with some losing their original function[10]. Furthermore, structure prediction of the glycoproteins of different flaviviruses identified novel and common mechanisms in the fusion machinery[8].

Here, we predicted the structures of all nine human herpesvirus proteomes, totaling 844 individual proteins. We critically evaluated their accuracy with stringent quality scores and thresholds. We clustered their predicted structures into structurally similar groups and analyzed them using a structural systems virology approach. Many similarities suggest a common ancestor and thus a homologous relationship, whether orthologous, *i.e.* due to speciation, or paralogous, *i.e.* due to gene duplication. We found new members to previous groups identified by sequence, such as the US22 family of HCMV. We could also identify unexpected groups with distinct folds, such as the UL112-113 family harboring a unique beta-barrel domain that is conserved in seven of the nine human herpesviruses. A detailed domain-level similarity search

identified several subclusters of proteins harboring different numbers of duplicated domains, deletions, and

acquisitions. Using structural similarity searches against cellular proteins, we propose new functions for

uncharacterized proteins. Examples include the HSV-1 and HSV-2 US2 proteins, which potentially have

dihydrofolate reductase activity. We also found cases of exaptation, such as the BMRF2 family of

transmembrane proteins that share structural similarity to nucleoside transporters but are likely inactive and

now have functions in cell-cell spread.

In this work, we create a structural database that is accessible through an open-access and searchable

web interface: https://www.herpesfolds.org/herpesfolds. It offers interactive displays of the predicted

structures and structural clusters. HerpesFolds is a curated database that groups the protein predictions

based on structural similarity while linking their established names and groups. It also serves as a reference

that can translate the complicated herpesvirus nomenclature for the expert and non-expert alike and sheds

light on the relationships of all nine human herpesvirus proteomes. This work will aid in developing wet-lab

testable hypotheses and gain new insights into herpesvirus biology and pathology.

Results

Most herpesvirus proteins can be confidently predicted

To generate proteome-wide predictions of prototypic strains of all 9 human herpesviruses (Table 1), we

initially used LocalColabFold[11], an implementation of AlphaFold2 that mainly uses a 40-60x faster multiple

sequence alignments (MSA) generator (**Error! Reference source not found.**A). For each protein, 5

models were generated. For most proteins, the associated predicted local distance difference test (pLDDT)

plots of all five models were very similar. This indicated that the models converged in all five runs to a

similar solution, indicating that the best solution was found (**Error! Reference source not found.**A). To

categorize the models objectively, we established three thresholds based on the pLDDT consistency

between models and the predicted template modeling (pTM) scores. See the methods section for more

details. Of the 844 proteins, 249 (30%) failed these quality scores (**Error! Reference source not found.**B

and **Error! Reference source not found.** Sheet 1). We used standard AlphaFold[1] (**Error! Reference**

source not found. Sheet 2) to improve these predictions using the slower but more thorough jackhmmer

MSA generator algorithm as well as LocalColabFold[11] with additional recycles (**Error! Reference source**

not found. Sheet 3). This resulted in good predictions for 95 (38%) of the previously rejected models.

Therefore, the final dataset consisted of 690 proteins (82% of 844 proteins) with consistent models as

indicated by the pLDDT plot (**Error! Reference source not found.** Sheet 4). We compared the proteins

that failed and passed the quality scores for each threshold criterion (**Error! Reference source not**

found.C-E). Most proteins that did not pass had low pTM and a small percentage of sequence with a

pLDDT >0.70. This correlation suggested that proteins low in folded regions, as indicated by a pLDDT >0.70,

had lower prediction confidence. The pLDDT score can also predict whether a protein region is structured

or intrinsically disordered. As shown before, a pLDDT >0.70 indicates a well-modeled region with a Cα root mean square displacement likely less than 1.5 Å[12]. Conversely, pLDDT scores <0.50 likely indicate intrinsically disordered regions (IDR)[1,13,14]. Consequently, the AlphaFold database has been used to predict protein disorder, such as in MobiDB[15]. IDRs have recently been implicated in essential processes such as replication compartment formation of HCMV by liquid-liquid phase separation[16]. Therefore, we determined which well-predicted proteins likely contain disordered regions by quantifying the percentage of pLDDT values each model has <0.50 or >0.70. As shown in **Error! Reference source not found.**A, 197 (23%) of the predicted proteins likely contain a disordered region (cut-off >44% of total sequence), and 640 (76%) of the proteins are likely structured (cut-off >33% of total sequence) (**Error! Reference source not found.** Sheet 5). This can be compared to the 1-8% disordered proteins in bacterial proteomes, 2-11% in archaea, 23-28% in eukaryotes, and 45% in humans[15,17,18]. Examples of N-terminal or C-terminal structured domains in a disordered protein are illustrated (**Error! Reference source not found.**B). We also found a subset of 100 proteins that did not pass the pTM threshold but showed converging model pLDDT plots, indicating that the predictions came to a final solution. These proteins contained extensive sequence stretches with a pLDDT <0.50, which indicated that these proteins are well-predicted, disordered proteins. Finally, to indicate transmembrane domains as well as signal peptides, all predictions were further annotated with DeepTMHMM (**Error! Reference source not found.** Sheet 6).

To illustrate the accuracy of the general prediction workflow, we compared our predictions to experimental data that were released after the AlphaFold 2.3 training cut-off of 2021-09-30. Our predictions aligned very well with experimental structures, as depicted for HCMV UL77[19] (**Error! Reference source not found.**). The Cα RMSD (310 atoms) between the predicted and experimental HCMV UL77 structure (7NXP[19]) was 0.509 Å, indicating a highly consistent model. These observations are in line with a previous community assessment that concluded that AlphaFold models are, on average, as good as experimental structures[3].

Table 1. Viral proteins used for structure prediction.

Subfamily	Virus	Virus abbreviation	Strain	Proteome ID or Reference Sequence	Number of proteins
alpha	herpes simplex virus 1	HSV-1	17	UP000009294	77
alpha	herpes simplex virus 2	HSV-2	HG52	UP000001874	74
alpha	varicella-zoster virus	VZV	Dumas	UP000002602	70
beta	human cytomegalovirus	HCMV	Merlin	UP000000938	181
beta	human herpesvirus 6A	HHV-6A	U1102	NC_001664	90
beta	human herpesvirus 6B	HHV-6B	Z29	NC_000898	97
beta	human herpesvirus 7	HHV-7	RK	NC_001716	84
gamma	Epstein-Barr virus	EBV	B95-8	UP000153037	87
gamma	Kaposi's sarcoma-associated herpesvirus	KSHV	GK18	UP000000942	86

HerpesFolds is a searchable online structure database

To make our proteome-wide predictions accessible and easily searchable, we created a database at https://www.herpesfolds.org/herpesfolds (**Error! Reference source not found.**F). The main table contains the individual herpesvirus protein names and their UniProtKB accession numbers. Based on the previous knowledge[20,21], we generated a chart where each row contains proteins with known homology (**Error! Reference source not found.** Sheet 1). This way, the database also serves as a reference guide. This feature is important as the nomenclature of human herpesviruses is highly inconsistent and difficult to interpret even for experts. We also integrated the common names based on the UniProtKB annotation. Moreover, each protein entry links to the UniProtKB webpage and an individual webpage containing an animated model and the pLDDT and PAE graphs (**Error! Reference source not found.**G). If predictions were rerun, the best model is displayed, and if the models of all runs failed the thresholds, a warning is displayed at the top of the page. Furthermore, transmembrane domains as detected by DeepTMHMM are indicated and a second model lacking the signal peptide is provided. As an important feature, we also integrated a Foldseek[5] search link kindly provided by the Steinegger lab that directly submits the predicted model to the Foldseek server and displays structurally similar proteins. Finally, we interactively display the structural similarity network (**Error! Reference source not found.**H).

Clustering of known and unexpected structural homologs

The identification of herpesvirus orthologs has classically been through sequence alignment, but this approach is restricted by its need for some extent of conserved sequence similarity. To find more distant relationships that might represent homologs between different viral species and determine the potential function of orphan genes, we employed structural similarity searches. We used both DALI[22] as an established algorithm and the newer Foldseek[5] algorithm and ran all models against each other. We used a threshold of DALI Z-score of >8 and E-value <0.001 for Foldseek (see materials and methods) (**Error! Reference source not found.** Sheet 4 and 5, respectively). Overall, the resulting structural clusters were similar, with 615 proteins in protein similarity clusters for DALI and 632 for Foldseek. One example of a difference is the tegument protein 3 (TEG3) group. Foldseek grouped only the alphaherpesvirus TEG3 proteins with the gammaherpesvirus BGLF3.5/ORF35 proteins. In contrast, DALI also found a similarity of the roseolovirus (HHV-6A, HHV-6B, and HHV-7) U68 proteins with this group. Since Foldseek performed slightly better overall and is computationally much more efficient, we used it for the remainder of this work. The resulting structural similarity connections were used to cluster the herpesvirus proteins into groups of structural similarity (**Error! Reference source not found.**A). All clusters are available for interactive exploration at https://www.herpesfolds.org/herpesclusters.

Based on the UniProtKB[20] and previous work[4,21], 39 protein groups were expected to be found in all 9

species. A more recent sequence analysis based on domain architecture identified 23 proteins with a

consistent domain structure in all 9 species and 169 homology groups[4]. Using structural similarity searches,

we identified 25 groups with exactly 1 protein from each human herpesvirus and 89 homology groups. Many

of these clusters matched these previous attempts to categorize herpesvirus proteins in groups based on

experimental data and sequence analysis, confirming our approach. An example are the homologs of the

alphaherpesvirus UL25 proteins, which are essential capsid-associated proteins and part of the portal cap[19].

We also found clusters of proteins with known and verified enzymatic functions, such as the herpesvirus

serine/threonine kinases, named after the prototypic HSV-1 US3 and UL13 proteins (**Error! Reference**

source not found.B). As only the protein models that passed the quality scores were used, not all groups

contain exactly one protein from each species. This was seen for the proliferating cell nuclear antigen

(PCNA)-like protein, large tegument protein, and tegument protein 7 clusters. For example, HCMV UL44 is

missing from the PCNA group because it was deemed a low-confidence model.

Surprisingly, the DNA polymerase catalytic subunit and DNA helicase/primase complex-associated protein

were found to be structurally similar (**Error! Reference source not found.**C). While these proteins have

different biological functions, this clustering can be explained by partial sequence similarity in the C-

terminus[23,24]. Furthermore, they are both similar to nucleases, with HSV-1 UL8 being homologous to the

exonuclease domain of B-family DNA polymerases[24] and HSV-1 UL30 having RNase H activity[25]. Since all

viruses have both of these genes, this suggests that while these genes likely share a common ancestor,

they have diverged in sequence and function.

The small capsomere-interacting protein (SCP) is conserved in all herpesviruses. However, they clustered

as distinct groups based on the subfamily (**Error! Reference source not found.**D). Genetic drift may have

led to structurally distinct features, as the last common ancestor likely predates the divergence of the

subfamilies. This was seen with both the sequence and structure based clustering.

A particularly interesting cluster of so far-unannotated proteins consisted of the likely homologs of HCMV

UL112-113. HCMV UL112-113 is essential for viral replication by mediating replication compartment

formation through phase separation at viral genomes, thereby recruiting the viral polymerase[16]. The poorly

characterized roseolovirus U79 proteins and the alphaherpesvirus UL4/ORF56 proteins cluster with UL112-

113. They shared an undescribed and conserved predicted beta-barrel (**Error! Reference source not**

found.A). These herpesviruses also form replication compartments and have a similar subcellular

localization[16,26], and their potential UL112-113 homologs would be prime candidates for further investigation.

Another cluster contained the gammaherpesvirus KSHV ORF73, also known as LANA, and EBV EBNA1

proteins that harbor a known DNA binding domain[27,28] (**Error! Reference source not found.**B). Our

structure predictions indicate that this DNA binding fold is conserved amongst all human herpesviruses,

and we further analyzed these proteins by generating AlphaFold 3 predictions of dimers[29] with DNA (**Error!**

Reference source not found.). The alphaherpesvirus UL3/ORF58 proteins showed similar surface charge
distributions and DNA binding sites as KSHV and EBV. In contrast, the betaherpesvirus UL122 and U86
proteins, which have known IE2 transcriptional regulation activity[30,31], bound to DNA with a different
orientation in the predictions. Surprisingly, the betaherpesvirus proteins UL117 and U84 were not predicted
to bind DNA, which is consistent with the less prominent positive surface charge. Perhaps these proteins
have lost the DNA binding activity and acquired a different function.

The HCMV phosphoprotein UL25 formed a group with its roseolovirus U14 homologs and clustered with
the HCMV UL35 proteins, which likely arose via gene duplication, constituting the pp85 superfamily[32].
Strikingly, we also found HCMV UL32 and its roseolovirus U11 homologs to be structurally similar. UL32,
also known as pp150, is a protein conserved in the betaherpesviruses. It binds directly to capsids as an
inner tegument protein while bridging the full virion tegument via its disordered C-terminus[33]. It will be
essential to dissect if the structurally similar UL25 and UL35 can also bind capsids and potentially regulate
pp150 abundance at pentons, potentially confusing the reconstruction of virus capsid cryo-EM maps. Finally,
the poorly studied KSHV ORF48 and likely the EBV BBRF2 protein, which was excluded from the clustering
due to low prediction scores, also clustered with the pp150 homologs. ORF48 is part of virions[34], and it
should be assessed if it can bind to the viral capsid as pp150 does.

Since some protein predictions have unstructured regions, it is possible that these flexible regions lead to
false negative results by Foldseek. To evaluate this possibility, all predicted structures were cut into smaller
pieces based on where they are likely structured, *i.e.* pLDDT >0.7. Foldseek clustering was then repeated
and compared to clustering based on full-length predictions (**Error! Reference source not found.** Sheet
9). While this method resulted in an interesting novel cluster containing EBV BALF1, EBV BHRF1, and
KSHV ORF16, which are all known apoptosis regulators, most novel clusters identified using this method
were likely false-positives such as HSV-1 UL49 and UL49A. UL49 is a tegument protein involved in immune
evasion, and UL49A is a glycoprotein, respectively. Another example is a cluster containing HCMV UL96
and HSV-1 UL14, which show little unique or similar tertiary structure. For this reason, we focused on the
analysis of full-length proteins.

We also compared our results to sequence-based clustering by MMseqs2 and HHblits (**Error! Reference
source not found.** Sheet 2 and 3, respectively). Importantly, several structure-based clusters contained
additional proteins when compared to sequence-based clustering methods, such as the UL112-113 cluster
and the KSHV ORF73 (LANA) cluster, where all the identified proteins of this cluster have functions related
to DNA binding. However, we also identified cases where structure predictions failed the quality scores and
HHblits-derived clusters were more complete. For example, all pAP proteins, which are splice variants of
the scaffold protein, failed the structure prediction quality scores and are absent from the structural cluster.
Since sequence-based methods can complement these instances where the structure prediction is of low
quality, we provide an option to view both the structure and sequence-based clusters at
https://www.herpesfolds.org/herpesclusters. **Error! Reference source not found.** Sheet 10 lists the

respective HHblits and Foldseek scores for each protein pair as well as all interactions above their
respective thresholds, which were found only by HHblits, only Foldseek, or both.

Immunoglobulin-like domains are common structural elements

One of the largest clusters we identified using structural clustering via Foldseek consists of viral proteins
that contain immunoglobulin (Ig)-like domains. This cluster merged several previously defined protein
families. One of the included groups is the HCMV RL11 family[35]. They share an RL11 domain (RL11D),
formed by a conserved tryptophan and two cysteines, and are structurally similar to immunoglobulin
domains. The RL11 proteins are non-essential in cell culture[36] but have high sequence variability between
isolates with some playing a role in modulating immune responses, such as UL11[37]. While there are 14
accepted members of this family, they did not form an isolated structural group in our clustering (**Error!**
Reference source not found.C). Unexpectedly, EBV BILF2 and KSHV K14 but also the well-studied HSV-
1 glycoproteins UL44 (gC), US6 (gD), US7 (gI), and US8 (gE) were structurally similar to multiple members
of this family. EBV BILF2 is encoded in a genome region unique to *lymphocryptoviruses*, and KSHV K14 is
also unique to KSHV. Furthermore, HSV-1 and HSV-2 US8 are located in a different part of the genome
than HCMV. In addition to the low sequence identity, these data suggest that these Ig-like domain-
containing proteins were acquired independently rather than from a common ancestor. Furthermore, gene
duplication continued after speciation. While most RL11 members have the same 6-strand antiparallel beta-
fold, RL13 has an additional beta-strand (**Error! Reference source not found.**D), potentially representing
a distinct evolutionary branch and function.

Identification of domain-level duplications and deletions

We performed the initial structural similarity search with full-length proteins. While this will identify
commonalities between proteins, it will, by definition, not identify differences. To address how the domain
architecture of a protein can differ between species, we performed a structural similarity search with
overlapping structural snippets using a sliding window approach (see materials and methods) (**Error!**
Reference source not found. Sheet 6). We identified 2806 structurally similar connections between
proteins, which includes 183 new connections for proteins relative to the full-length analysis (**Error!**
Reference source not found. Sheet 1 and 2). This analysis identified 24 internal duplications, where
domains were likely duplicated in the same protein; 35 repetitive acquisitions, where domains of a different
protein matched the query protein multiple times at different positions; and 13 domain additions, where
multiple domains of a protein matched different proteins (**Error! Reference source not found.**A and **Error!**
Reference source not found. Sheet 3).

We found that HSV-1 UL15 was structurally similar to the previously identified HSV-2 UL15 but also HSV-
1 UL9 (**Error! Reference source not found.**B). UL15 is part of the tripartite terminase, essential for
packaging viral genomes into newly formed capsids, and UL9 is the replication origin-binding protein. The
fragment that aligns is oriented in the opposite direction relative to the core in UL15 versus UL9, which
suggests that this region is not involved in an external interaction and has a conserved structural integrity
function.

An important example of domain duplications is in the large betaherpesvirus US22 family cluster. No
experimental structure of this large family of proteins is available, and sequence-based methods indicate
that the family consists of at least 12 members in HCMV[38,39]. Using full-length structural similarity searches,
we identified HCMV proteins that share the same conserved fold (**Error! Reference source not found.**C).
Importantly, this group consists of four different domain architectures depending on whether the protein
contains 0, 1, 2, or 4 copies of the US22 domain (**Error! Reference source not found.**D). The majority of
proteins are similar to US22 with two domains, while the following members have only one domain: HCMV
UL26, HCMV UL26_UL26.21, HCMV UL27, HCMV UL38, HHV-6A U19, HHV-6B U19, and HHV 7 U19.
We also found a subgroup that lack a US22 homology domain consisting of HHV-6A U4, HHV-6B U4, and
HHV-7 U4. They are connected to the main group through the HHV-6A U7 and HHV-7 U7 proteins, which
appear to have an additional domain relative to US22, which the U4 proteins consist of exclusively. The
proteins with four copies of the US22 domain are HCMV UL29, HHV-6A U7, and HHV-7 U7. Oddly, HHV-
6B U7, unlike the other roseoloviruses, only has two domains, suggesting a divergence in this species.
Furthermore, many group members were identified to contain an internal duplication and/or repetitive
acquisition. The low RMSDs between the two domains of US22 and between the domains from different
proteins are consistent with a domain duplication (**Error! Reference source not found.**D). These data
suggest that the US22 family is bigger than previously thought and that domain additions and deletions are
particularly common in this family.

Viral exaptation of cellular proteins

Many of the identified clusters contained understudied proteins. We expanded the structural similarity
search to cellular proteins using Foldseek to determine their potential biological activity from the PDB
database (**Error! Reference source not found.** Sheet 1), and the predicted AlphaFold structures of the
Swiss-Prot database (**Error! Reference source not found.** Sheet 3). The PDB contains experimentally
verified structures, but it is limited in diversity. In contrast, the AlphaFold predictions of the Swiss-Prot
database enable comparisons to a much more extensive predicted database that is less biased by
experimental limitations. We combined the significant hits from the two databases and identified structurally
similar cellular proteins. To find putative functions for the viral protein groups, the functional description of
the homologous proteins was mined for keywords (**Error! Reference source not found.** Sheets 2 and 4).

Unfortunately, the depositors' descriptions of structures in the PDB database are not systematic. To handle
these variations, we calculated the frequencies of all words in the descriptions, and the words with the
highest frequencies, which likely represent the protein function, were manually chosen as the annotation.
Each viral homology group identified above was then given a HerpesFolds annotation. Since the PDB, but
not the AlphaFold database, contains structures of viral proteins, it is important to note that some hits in the
PDB database are because they were aligning with themselves.

Some predicted viral protein structures did not map to any other protein fold in the searched databases,
indicating that these are unique folds (**Error! Reference source not found.** Sheet 5). We found that even
though the viral alkaline nucleases are not similar to cellular nucleases, they are consistent within the
herpesviruses (**Error! Reference source not found.**A). Experimental structures have been solved for
EBV[40] and KSHV[41,42]. While the central core was found to show some similarity to λ-exonuclease, the viral
proteins have additional terminal extensions. Another striking example was the cytoplasmic envelopment
protein 2, which is conserved in all herpesviruses and for which no experimental structure exists (**Error!
Reference source not found.**A). Still, this protein group was very well predicted as indicated by its pLDDT
and PAE scores. Moreover, the most distant phylogenetic pair, VZV ORF44 and HHV-6B U65, had an
RMSD of 4.3 Å against the 112 Cα atoms despite a sequence identity of only 14%. These results indicate
that AlphaFold can predict herpesvirus proteins with high confidence *de novo* even if no structurally similar
dataset exists. Since the CEP2 proteins are crucially involved in virion envelopment[43], it will be important
to analyze their role in modulating the cellular secretory system through this unique fold.

Using this workflow, we could annotate several poorly studied viral proteins and identify potential functions.
One example is HSV-1 US2, which is structurally similar to dihydrofolate reductases (DHFRs) (**Error!
Reference source not found.**B-D). The most significant cellular target was the DHFR from *Mesocricetus
auratus* (commonly known as the golden or Syrian hamster) (PDB 3EIG[44]). Furthermore, HSV-1 US2 was
structurally similar to KSHV ORF2, a known DHFR homolog with enzymatic activity[45,46]. While KSHV ORF2
contains the catalytic residues identified in *E. coli*[47,48], HSV-1 and HSV-2 US2 do not contain them (**Error!
Reference source not found.**). Since no DHFR activity has been reported for the US2 proteins, this fold
was likely exaptated. Functional assays will be needed to verify if the US2 proteins still have DHFR activity.

We also identified a cluster containing EBV BMRF2, KSHV ORF58, VZV ORF15, and HSV-1 UL43 (with
HSV-2 UL43 being absent due to low-quality scores). Importantly, EBV BMRF2, KSHV ORF58, and VZV
ORF15 predictions were significantly similar to the human equilibrative nucleoside transporter (ENT) 4
(**Error! Reference source not found.**E-G). This cellular protein is involved in transporting metabolites and
regulating adenosine concentrations[49]. However, no experimental data currently supports a role for any of
these viral proteins in transmembrane transport. Instead, EBV BMRF2 can change cellular morphology by
modulating the actin cytoskeleton[50] and is essential for efficient cell-cell spread[51]. In a closely related
alphaherpesvirus, UL43 colocalizes with proteins involved in membrane fusion and potentially interacts with
the gM/gN complex[52], which is engaged in syncytia formation[53]. These commonalities suggest a function

for this viral protein cluster in cell-cell spread, which does not align with the transmembrane transporter activity of the structurally similar cellular proteins. Moreover, comparative cross-sections of the viral proteins and ENT4 illustrate that they do not have a channel necessary for metabolite transport[54] (**Error! Reference source not found.**A). Several transporters depend on accessory proteins for correct trafficking to the plasma membrane, such as the lysosomal transporter MFSD1, which depends on GLMP[55]. This is also true for EBV BMRF2, which needs BDLF2[50] for correct trafficking to the plasma membrane, and KSHV ORF58, which needs ORF27[34]. Interestingly, AlphaFold predictions of the viral heterodimers suggest that even if channels were present in BMRF2 and ORF58, they would be blocked by the partner protein (**Error! Reference source not found.**B), arguing against a conserved transporter activity as recently proposed[56]. Instead, the structure predictions likely indicate that the similarity to ENT4 may stem from "radical" exaptation, where the transporter activity was lost and replaced by a role in viral cell-to-cell spread.

All human herpesviruses encode for deoxyuridine triphosphatases (dUTPases)[57] called HSV-1 UL50, HSV-2 UL50, VZV ORF8, HCMV UL72, U45 of roseoloviruses, KSHV ORF54 and EBV BLLF3. Cellular dUTPases are homotrimers that form three active centers at their interfaces. In contrast, the herpesvirus dUTPases are monomers consisting of a fusion of two dUTPase domains, as shown for EBV BLLF3[58]. While the respective alpha- and gammaherpesvirus proteins still have enzymatic activity, the betaherpesvirus homologs have lost them and are involved in immune modulation[59]. Human beta- and gammaherpesviruses code for 17 more dUTPase-related proteins (DURPs), likely due to gene duplications[57]. Using domain-level search, we found that all DURPs were predicted to harbor a three-domain architecture in contrast to the two domains of the viral dUTPases. However, the separate domains diverged quite significantly from the canonical dUTPase fold as indicated by their RMSD (**Error! Reference source not found.**H), which is in line with a recent report on the crystal structure of the HCMV DURP UL82 (pp71), which describes that it has lost its dUTPase function[60].

Herpesviruses have been suggested to induce autoimmunity through molecular mimicry[61,62]. Several studies have attempted to identify viral proteins that might induce autoimmunity due to a similarity to a cellular protein[63,64]. Protein sequence-based analyses could not identify viral proteins that would induce autoimmunity to known cellular proteins[63], such as myelin for multiple sclerosis[64]. Our structural similarity search identified no direct structural match of myelin to a viral protein. Interestingly, multiple members of the immunoglobulin domain-containing group, such as HSV-1 US8, mapped to myelin protein zero-like protein 1, and others, such as EBV BARF1, mapped to myelin-associated glycoprotein. These myelin-related proteins could be involved and warrant further investigation.

Discussion

Understanding protein structures is critical to comprehending pathogen-host interactions. Unfortunately, experimental structural data is lacking for most human herpesvirus proteins. Machine learning-based

structure prediction tools have made huge strides in accuracy and provide quality scores for their

predictions[1,2,11]. Accordingly, recent community evaluations have found "that AF2 [AlphaFold2]-predicted

structures, on average, tend to give results that are as good as those derived from experimental structures"[3]

while also cautioning that "AlphaFold predictions are valuable hypotheses and accelerate but do not replace

experimental structure determination"[12].

The human herpesviruses encode for at least 844 proteins, and for most, no experimental structure has

been solved. Here, we applied proteome-wide predictions to provide a structural systems view of all

annotated human herpesvirus proteins and developed stringent scoring workflows to validate our

predictions. We generated clusters of proteins with similar folds using all-versus-all full-length structural

similarity searches. Importantly, we identified previously unknown protein clusters sharing conserved folds,

such as the UL112-113 family that carries a beta-barrel-like fold and likely plays an essential role in genome

replication, while confirming previously known clusters, such as viral G-protein coupled receptors and

kinases. In comparison, previous work has used sequence-based approaches to cluster herpesvirus

proteins into related families. Importantly, neither a sequence-based alignment algorithm (MMseqs2) nor

an HMM-HMM-based algorithm (HHblits) could uncover all of the novel and biologically important

relationships that we found using a structural approach, such as the finding that the alphaherpesvirus UL3

proteins are new members of the KSHV ORF73 (LANA) cluster, which are DNA-binding proteins and have

too little sequence conservation to be identified by MMseqs2 or HHblits.

Still, structural clustering might not always indicate direct evolutionary relatedness. The large Ig-like domain

cluster appears to be a group of proteins with a common fold, likely the result of independent acquisitions

rather than originating from a single common ancestor or gene duplication. Using domain-level searches,

we identified differences between structurally related proteins such as viral dUTPases and the US22 family.

Both consist of subgroups with different numbers of duplicated domains that correlate with their described

activity. Finally, we searched for cellular proteins with matching folds. We identified cases of potential

exaptation, such as the membrane transporter ENT4 by human alpha-and gammaherpesviruses and

unique folds that do not match any cellular protein, such as the CEP2 proteins.

This body of data provides many valuable hypotheses for experimental validation. To accelerate the use of

this data and research on human herpesvirus, we have generated an easy-to-use, searchable web interface

for the community, accessible at https://www.herpesfolds.org/herpesfolds. The database groups proteins

by structural relationship and allows interactive evaluation of all structures. PDB files of each prediction can

be downloaded or directly submitted to the Foldseek structural similarity search server to find similar cellular

proteins in up-to-date databases.

This work is a first step based on high-confidence monomeric predictions of viral proteins. It neglects that

some proteins may fold differently in complex with others, adopt multiple conformations, or need to form

oligomers to function, such as gB[65]. Future work will need to take into account protein complex composition

and stoichiometry. However, accomplishing this task is challenging as this information is unavailable for
most viral proteins and generating all likely and possible combinations is computationally highly demanding.
The next logical step will be predicting viral protein-protein interaction networks. Since post-translational
modifications and protein-ligand complexes[66,67] or nucleic acids[68] will likely play a role in many such
complexes, the recently released RoseTTAFold-All-Atom[69] and AlphaFold 3[70] algorithms will be of great
use.

Herpesviruses have large bicistronic genomes, and for example, ribosomal profiling has identified 100s of
additional transcripts and ORFs for HSV-1[71], HCMV[72], EBV[73] and KSHV[74,75]. Structure predictions of these
could provide insights into their functions and roles during infection and pathogenesis and should be the
next step.

The methodology presented here applies to any virus or virus family. It can easily be extended to analyze
all reading frames of a given organism, not only the annotated ORFs. As the databases of predicted and
experimental viral protein structures grow, more connections between different viral pathogens and their
hosts will be made. The speed at which these *in silico* models can be generated will drive rapid hypothesis
design and subsequent experimental confirmation.

Methods

Viruses

The strains and associated reference numbers of the nine human herpesvirus proteomes used for
prediction are noted in Table 1. In addition to the full-length proteins, splice variants were also included. A
complete list of the predicted proteins can be found in **Error! Reference source not found.** Sheet 4.

Software

Sequence alignments were performed with MMseqs2 release 15-6f452[76]
(https://github.com/soedinglab/MMseqs2) and HHblits v3.3.0[77] (https://github.com/soedinglab/hh-suite).
HHblits HMM generation used the UniRef30 database from 2023_02. The significance threshold used for
MMseqs2 was E-value <0.1[76] and for HHblits was E-value <0.001
(https://toolkit.tuebingen.mpg.de/tools/hhblits). Protein structure predictions were made with
LocalColabFold v1.5.1[11] (https://github.com/YoshitakaMo/localcolabfold) and AlphaFold[1] using AlphaFold
version 2.3.0 (https://github.com/google-deepmind/alphafold). Fasta files were downloaded from
https://www.uniprot.org, and structure predictions were performed in accordance with the respective
documentation detailed in the respective GitHub releases. For each protein, 5 models were generated using
3 recycles and a stop-at-score of 100. Proteins that failed the quality scores were rerun with LocalColabFold
with 20 recycles. AlphaFold 3[70] predictions were performed through the AlphaFold Server at
https://alphafoldserver.com. Protein topology was identified with DeepTMHMM[78] using their provided Demo
Colab (https://dtu.biolib.com/DeepTMHMM). The custom python script "DeepTMHMM_export_domains.py"
consolidated the DeepTMHMM outputs into a single file and "DeepTMHMM_make_truncated_fasta.py"
generated fasta files of the proteins without the identified signal peptide. Structural similarity searches were
performed with local install versions of DALI[22] (DaliLite.v5 from http://ekhidna2.biocenter.helsinki.fi/dali/)
and Foldseek[5] version 2ad017897d3dab66dd33ea675e92215bdfb4a64d
(https://github.com/steineggerlab/foldseek). Networks were visualized with Gephi v0.10.1 (http://gephi.org/).
UCSF ChimeraX version 1.8 (https://www.cgl.ucsf.edu/chimerax) was used for visualization of structures,
and PyMOL 2.5.8 (https://www.pymol.org) was used for root mean square deviation (RMSD) calculations.
Data analysis was carried out with custom-written Python scripts, which were partly generated with the
assistance of large language models and thoroughly tested manually. The complete set of scripts can be
accessed through the project's GitHub repository: https://github.com/QuantitativeVirology/Herpesfolds.

Evaluation of model quality

All proteins were initially predicted using LocalColabFold[11] with 3 recycles as it is more time-efficient. To

assess the quality of the resulting models, 3 thresholds were used. Failing any single threshold led to the

model being flagged as "fail". We interrogated whether the model optimization had converged by testing

the consistency of the predicted local distance difference test (pLDDT) scores associated with every model

(**Error! Reference source not found.**A and B). To do so, we first calculated the mean and standard

deviation of the pLDDT values per model and then the standard deviation of these 5 values, resulting in

"StDev of mean" and "StDev of StDev". To set thresholds, we fit a Gaussian curve to their respective

histograms and set the cutoff at the mean of the Gaussian plus 2 times its standard deviation (**Error!**

Reference source not found.C and D). Therefore, a "StDev of mean" >3.2 or a "StDev of StDev" >1.9 was

deemed a low-quality model. We also evaluated the predicted template modeling score (pTM) to validate

model confidence further. To set a global threshold at which a prediction likely constitutes a folded protein,

we chose 106 herpesvirus proteins representing the protein length distribution of the herpesvirus proteomes

and scrambled their amino acid sequences. These scrambled sequences were used for model prediction

and treated as a negative dataset, assuming that a random residue order should not fold. Receiver

Operating Characteristic (ROC) analysis yielded an area under the curve of 91.3%, and the Youden index

was used to set a threshold (**Error! Reference source not found.**E). Following this analysis, a pTM

<0.3150 was deemed a low-quality model. We reanalyzed all models that failed the initial threshold in

AlphaFold[1] and LocalColabFold[11] with 20 recycles and rescored them. For example, **Error! Reference**

source not found.F shows an improved model run in AlphaFold with higher pLDDT scores compared to

Error! Reference source not found.B. The corresponding scripts can be found in our HerpesFolds GitHub

repository in the file "score_colabfold.py" at https://github.com/QuantitativeVirology/Herpesfolds.

Defining disordered and structured content of models

The pLDDT value at a given residue indicates whether a region is predicted to be disordered (pLDDT <0.50)

or structured (pLDDT >0.70). For the models that passed all quality scores, we generated histograms of

the percentage of the protein below or above these values and fit a Gaussian to both curves. Since the

majority of proteins are expected to be structured, a threshold, >44%, for proteins containing a disordered

region was set at the mean of the distribution + 2x StDev for the percentage of protein sequence below

0.50, and a threshold, >33%, for proteins with a structured region was set at the mean - 2x StDev for the

percentage of protein sequence above a pLDDT of 0.70. The script for extracting pLDDT values from

the .json files associated with every prediction can be found in our HerpesFolds GitHub repository in the

file "score_colabfold.py" at https://github.com/QuantitativeVirology/Herpesfolds.

Extracting pLDDT domains from predicted structures

The structured domains from a predicted structure were extracted based on the pLDDT scores. The regions
with a pLDDT >0.7 were identified. If adjacent regions were connected by a disordered region, *i.e.* pLDDT
<0.7, that was shorter than 100 residues, the adjacent regions along with the disordered region were
counted as a single domain. The corresponding script can be found as "extract_pLDDT_domains.py" in our
HerpesFolds GitHub repository at https://github.com/QuantitativeVirology/Herpesfolds.

Structural similarity search with DALI and Foldseek

Foldseek aligns protein structures with decreased computation time compared to alternative algorithms[5]. It
provides different alignment algorithms, 3Di+AA Gotoh-Smith-Waterman and Tmalign, and measures of
significance, such as the Expect-value (E-value) and probability of homology (prob). We tested both
alignment algorithms with different thresholds and compared the protein clusters, **Error! Reference source**
not found. Sheet 7. The Tmalign results did not agree with the sequence based clustering and known
literature. The E-value and prob thresholds with 3Di+AA Gotoh-Smith-Waterman gave similar results, and
thus we chose to continue with an E-value threshold of 0.001 to be consistent with the literature[10], **Error!**
Reference source not found. Sheet 8.

To cluster viral proteins into groups of structural similarity, we employed DALI as an established algorithm
and Foldseek as a recent algorithm designed to work with vast predicted datasets with decreased
computation times. With DALI, we used the recommended threshold of Z-score >8 for probable homologs[79],
and with Foldseek, we used the previously published threshold of <0.001[10]. Since the results from both
algorithms were similar, we only used Foldseek for virus-host and domain-level searches as DALI is orders
of magnitude slower, which would have made these analyses computationally too expensive otherwise.
Similarity searches against cellular proteins were done using the PDB database (**Error! Reference source**
not found. Sheet 1), and the predicted AlphaFold structures of the Swiss-Prot database (**Error! Reference**
source not found. Sheet 3).

Clustering of structure predictions by similarity

The Foldseek and DALI outputs are a list of query and target proteins with the associated significance value.
We used a threshold of E-value <0.001 for Foldseek and a Z-score >8 for DALI to determine the significant
results. Both reciprocal comparisons for each protein pair needed to be significant for inclusion in the
clustering, *i.e.* A matches B and B matches A. If protein A was structurally similar to B and B similar to C,
then A, B, and C were clustered. Once all clusters were identified, we converted the protein names to their
row position in **Error! Reference source not found.** Sheet 1. This system allowed us to quickly determine

whether a structural cluster contained proteins expected to group, *i.e.* all row numbers are the same, or not,

i.e. the row numbers are different, based on UniProtKB annotations[20] and previously known homologies[21].

The associated script can be found in our HerpesFolds GitHub repository in the file

"analyze_foldseek_for_structural_similarity_cluster.py"   at

https://github.com/QuantitativeVirology/Herpesfolds.

Analysis of sliding window structural similarity

To search for alterations at the domain level between protein predictions, we generated "structural snippets"

from all predictions using a sliding window approach. We then ran these structural snippets in an all-versus-

all Foldseek search to identify conserved folds. We empirically tested which window size would result in the

highest number of significant hits using Foldseek. Trying a window size of 25, 50, 100, 150, and 200 amino

acids, we found the most significant hits with a window size of 150. The sliding window step size was tested

at 5, 10, 15, 20, and 25 amino acids. We used a step size of 20 amino acids for the final analysis as it

resulted in the maximal number of protein species detected. This approach resulted in over 12000 structural

snippets of 150 residues in length, each overlapping by 130 residues with the preceding fragment and

covering the whole folded human herpesvirus proteome. All structural fragments were compared pairwise

with Foldseek, accounting for about 1.6 million tested combinations. We next assembled "structural contigs"

of overlapping frames that passed the Foldseek threshold of an E-value <0.001. If a structural snippet

matched another structural snippet and they were not overlapping in the same original protein, then the pair

was deemed a "hit". If several structural snippets overlapped in sequential order, they were used to build

larger domains. This analysis resulted in a list of structured areas, which we used to determine if a query

protein has a different domain architecture than a structurally similar protein in this data set. If a snippet of

a query protein matched a different part of the query protein, we deemed it to have an "internal duplication".

If a target snippet matched the query protein at multiple locations, we considered the query protein to have

a "repetitive acquisition". If the query protein was structurally similar to a target protein but the query protein

had at least 1 domain that was not found in the target protein, we deemed the protein to have a "domain

addition". The associated script can be found in our HerpesFolds GitHub repository in the files

"make_sliding_window_pdb.py", "Sliding_Window_Analysis.ipynb","analyze_foldseek_sliding_window.py",

and "analyze_sliding_window_domains.py" at https://github.com/QuantitativeVirology/Herpesfolds.

Keyword analysis of Foldseek results

Most Foldseek searches against cellular databases resulted in multiple significant matches, each

containing protein descriptions and matching protein identifiers. We filtered for the most enriched keywords

to extract likely biological functions from these descriptions. The following terms, in addition to single
characters, single and double digits, and punctuation, were filtered out as they are too generic:

crystal, a, after, allosteric, allosteric, alpha, an, and, angstrom, angstroms, antibody, apo, as, at, atomic,
based, basic, beta, between, bound, by, by, c1, c-c, cell, chain, chi, class, class, coli, complex, complexed,
complexes., compound, conserved, cryoem, cryo-em, crystal, c-terminal, c-x-c, de, dehydration, deletion,
delta, design, different, discovery, domain, e., e.coli, edition, element, engineered, epsilon, escherichia, eta,
fab, for, form, fragment, from, full, full-length, functional, gamma, herpes, herpesvirus, high, holoenzyme, ii,
iii, implications, in, inhibitor, inhibitors, ion, iota, its, kappa, lambda, ligand, ligand-binding, loader, loop, low-
ph, mode, molecule, monoclonal, motif, mu, mutant, natural, neutralizing, novel, novo, nu, of, omega,
omicron, on, one, open, open, opposite, peptide, phi, pi, probable, protein, psi, region, resolution, reveal,
rho, sigma, significance, site, specificity, structural, structure, structures, substrate, symmetry, tau,
template-primer, the, the, therapeutic, theta, to, type, uncharacterized, unit, upsilon, using, variant, virus,
with, xfel, xi, x-ray, zeta

The used script can be found in our HerpesFolds GitHub repository at
https://github.com/QuantitativeVirology/Herpesfolds in the files "Keyword_Frequency_SP_PDB.ipynb".
This script was written with the assistance of a large language model.

Data Availability

569 Predictions and interactive clustering is available through a web interface at
570 https://www.herpesfolds.org/herpesfolds and https://www.herpesfolds.org/herpesclusters. Moreover, all
571 predictions and ChimeraX sessions are downloadable at https://zenodo.org/records/13284140.

Code Availability

574 Code is available at the project's GitHub repository: https://github.com/QuantitativeVirology/Herpesfolds.

References

1   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A.,
Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E.,
Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals,
O., Senior, A. W., Kavukcuoglu, K., Kohli, P. & Hassabis, D. Highly accurate protein structure
prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

2   Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R., Wang, J., Cong, Q.,
Kinch, L. N., Schaeffer, R. D., Millan, C., Park, H., Adams, C., Glassman, C. R., DeGiovanni, A.,
Pereira, J. H., Rodrigues, A. V., van Dijk, A. A., Ebrecht, A. C., Opperman, D. J., Sagmeister, T.,
Buhlheller, C., Pavkov-Keller, T., Rathinaswamy, M. K., Dalwadi, U., Yip, C. K., Burke, J. E., Garcia,
K. C., Grishin, N. V., Adams, P. D., Read, R. J. & Baker, D. Accurate prediction of protein
structures and interactions using a three-track neural network. *Science* **373**, 871-876,
doi:10.1126/science.abj8754 (2021).
3   Akdel, M., Pires, D. E. V., Pardo, E. P., Janes, J., Zalevsky, A. O., Meszaros, B., Bryant, P., Good, L.
L., Laskowski, R. A., Pozzati, G., Shenoy, A., Zhu, W., Kundrotas, P., Serra, V. R., Rodrigues, C. H.
M., Dunham, A. S., Burke, D., Borkakoti, N., Velankar, S., Frost, A., Basquin, J., Lindorff-Larsen, K.,
Bateman, A., Kajava, A. V., Valencia, A., Ovchinnikov, S., Durairaj, J., Ascher, D. B., Thornton, J.
M., Davey, N. E., Stein, A., Elofsson, A., Croll, T. I. & Beltrao, P. A structural biology community
assessment of AlphaFold2 applications. *Nat Struct Mol Biol* **29**, 1056-1067, doi:10.1038/s41594-
022-00849-w (2022).
4   Zmasek, C. M., Knipe, D. M., Pellett, P. E. & Scheuermann, R. H. Classification of human
Herpesviridae proteins using Domain-architecture Aware Inference of Orthologs (DAIO).
Virology **529**, 29-42, doi:10.1016/j.virol.2019.01.005 (2019).
5   van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L. M., Soding, J. &
Steinegger, M. Fast and accurate protein structure search with Foldseek. *Nat Biotechnol* **42**, 243-
246, doi:10.1038/s41587-023-01773-0 (2024).
6   Abrescia, N. G., Bamford, D. H., Grimes, J. M. & Stuart, D. I. Structure unifies the viral universe.
Annu Rev Biochem **81**, 795-822, doi:10.1146/annurev-biochem-060910-095130 (2012).
7   Ng, W. M., Stelfox, A. J. & Bowden, T. A. Unraveling virus relationships by structure-based
phylogenetic classification. *Virus Evol* **6**, veaa003, doi:10.1093/ve/veaa003 (2020).
8   Oliver, M. R., Toon, K., Lewis, C. B., Devlin, S., Gifford, R. J. & Grove, J. Structures of the Hepaci-,
Pegi-, and Pestiviruses envelope proteins suggest a novel membrane fusion mechanism. *PLoS
Biol* **21**, e3002174, doi:10.1371/journal.pbio.3002174 (2023).
9   Evseev, P., Gutnik, D., Shneider, M. & Miroshnikov, K. Use of an Integrated Approach Involving
AlphaFold Predictions for the Evolutionary Taxonomy of Duplodnaviria Viruses. *Biomolecules* **13**,
doi:10.3390/biom13010110 (2023).
10   Mutz, P., Resch, W., Faure, G., Senkevich, T. G., Koonin, E. V. & Moss, B. Exaptation of
Inactivated Host Enzymes for Structural Roles in Orthopoxviruses and Novel Folds of Virus
Proteins Revealed by Protein Structure Modeling. *mBio* **14**, e0040823, doi:10.1128/mbio.00408-
23 (2023).
11   Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. & Steinegger, M. ColabFold:
making protein folding accessible to all. *Nat Methods* **19**, 679-682, doi:10.1038/s41592-022-
01488-1 (2022).
12   Terwilliger, T. C., Liebschner, D., Croll, T. I., Williams, C. J., McCoy, A. J., Poon, B. K., Afonine, P.
V., Oeffner, R. D., Richardson, J. S., Read, R. J. & Adams, P. D. AlphaFold predictions are valuable
hypotheses and accelerate but do not replace experimental structure determination. *Nat
Methods* **21**, 110-116, doi:10.1038/s41592-023-02087-4 (2024).
13   Necci, M., Piovesan, D., Predictors, C., DisProt, C. & Tosatto, S. C. E. Critical assessment of
protein intrinsic disorder prediction. *Nat Methods* **18**, 472-481, doi:10.1038/s41592-021-01117-
3 (2021).
14   Piovesan, D., Monzon, A. M. & Tosatto, S. C. E. Intrinsic protein disorder and conditional folding
in AlphaFoldDB. *Protein Sci* **31**, e4466, doi:10.1002/pro.4466 (2022).

15　Piovesan, D., Del Conte, A., Clementel, D., Monzon, A. M., Bevilacqua, M., Aspromonte, M. C.,
Iserte, J. A., Orti, F. E., Marino-Buslje, C. & Tosatto, S. C. E. MobiDB: 10 years of intrinsically
disordered proteins. *Nucleic Acids Res* **51**, D438-D444, doi:10.1093/nar/gkac1065 (2023).
16　Caragliano, E., Bonazza, S., Frascaroli, G., Tang, J., Soh, T. K., Grunewald, K., Bosse, J. B. & Brune,
W. Human cytomegalovirus forms phase-separated compartments at viral genomes to facilitate
viral replication. *Cell Rep* **38**, 110469, doi:10.1016/j.celrep.2022.110469 (2022).
17　Dunker, A. K., Obradovic, Z., Romero, P., Garner, E. C. & Brown, C. J. Intrinsic protein disorder in
complete genomes. *Genome Inform Ser Workshop Genome Inform* **11**, 161-171 (2000).
18　Oldfield, C. J., Cheng, Y., Cortese, M. S., Brown, C. J., Uversky, V. N. & Dunker, A. K. Comparing
and combining predictors of mostly disordered proteins. *Biochemistry* **44**, 1989-2000,
doi:10.1021/bi047993o (2005).
19　Naniima, P., Naimo, E., Koch, S., Curth, U., Alkharsah, K. R., Stroh, L. J., Binz, A., Beneke, J. M.,
Vollmer, B., Boning, H., Borst, E. M., Desai, P., Bohne, J., Messerle, M., Bauerfeind, R., Legrand,
P., Sodeik, B., Schulz, T. F. & Krey, T. Assembly of infectious Kaposi's sarcoma-associated
herpesvirus progeny requires formation of a pORF19 pentamer. *PLoS Biol* **19**, e3001423,
doi:10.1371/journal.pbio.3001423 (2021).
20　The UniProt, C. UniProt: the universal protein knowledgebase. *Nucleic Acids Res* **45**, D158-D169,
doi:10.1093/nar/gkw1099 (2017).
21　in *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*　(eds A. Arvin *et al.*)　(2007).
22　Holm, L. & Sander, C. Dali: a network tool for protein structure comparison. *Trends Biochem Sci*
20, 478-480, doi:10.1016/s0968-0004(00)89105-7 (1995).
23　Bermek, O. & Williams, R. S. The three-component helicase/primase complex of herpes simplex
virus-1. *Open Biol* **11**, 210011, doi:10.1098/rsob.210011 (2021).
24　Kazlauskas, D. & Venclovas, C. Herpesviral helicase-primase subunit UL8 is inactivated B-family
polymerase. *Bioinformatics* **30**, 2093-2097, doi:10.1093/bioinformatics/btu204 (2014).
25　Weisshart, K., Kuo, A. A., Hwang, C. B., Kumura, K. & Coen, D. M. Structural and functional
organization of herpes simplex virus DNA polymerase investigated by limited proteolysis. *J Biol*
Chem **269**, 22788-22796 (1994).
26　Jahedi, S., Markovitz, N. S., Filatov, F. & Roizman, B. Colocalization of the herpes simplex virus 1
UL4 protein with infected cell protein 22 in small, dense nuclear structures formed prior to
onset of DNA synthesis. *J Virol* **73**, 5132-5138, doi:10.1128/JVI.73.6.5132-5136.1999 (1999).
27　Bochkarev, A., Barwell, J. A., Pfuetzner, R. A., Bochkareva, E., Frappier, L. & Edwards, A. M.
Crystal structure of the DNA-binding domain of the Epstein-Barr virus origin-binding protein,
EBNA1, bound to DNA. *Cell* **84**, 791-800, doi:10.1016/s0092-8674(00)81056-9 (1996).
28　Stevenson, J. P. Correspondence: Fish diseases research. *Vet Rec* **95**, 325-326,
doi:10.1136/vr.95.14.325 (1974).
29　Hellert, J., Weidner-Glunde, M., Krausze, J., Lunsdorf, H., Ritter, C., Schulz, T. F. & Luhrs, T. The
3D structure of Kaposi sarcoma herpesvirus LANA C-terminal domain bound to DNA. *Proc Natl*
Acad Sci U S A **112**, 6694-6699, doi:10.1073/pnas.1421804112 (2015).
30　Ball, C. B., Li, M., Parida, M., Hu, Q., Ince, D., Collins, G. S., Meier, J. L. & Price, D. H. Human
Cytomegalovirus IE2 Both Activates and Represses Initiation and Modulates Elongation in a
Context-Dependent Manner. *mBio* **13**, e0033722, doi:10.1128/mbio.00337-22 (2022).
31　Gravel, A., Tomoiu, A., Cloutier, N., Gosselin, J. & Flamand, L. Characterization of the immediate-
early 2 protein of human herpesvirus 6, a promiscuous transcriptional activator. *Virology* **308**,
340-353, doi:10.1016/s0042-6822(03)00007-2 (2003).
32　Szymanska-de Wijs, K., Dezeljin, M., Bogdanow, B. & Messerle, M. Viral determinants influencing
intra- and intercellular communication in cytomegalovirus infection. *Curr Opin Virol* **60**, 101328,
doi:10.1016/j.coviro.2023.101328 (2023).

33   Bogdanow, B., Gruska, I., Muhlberg, L., Protze, J., Hohensee, S., Vetter, B., Bosse, J. B., Lehmann,
M., Sadeghi, M., Wiebusch, L. & Liu, F. Spatially resolved protein map of intact human
cytomegalovirus virions. *Nat Microbiol* **8**, 1732-1747, doi:10.1038/s41564-023-01433-8 (2023).
34   Nabiee, R., Syed, B., Ramirez Castano, J., Lalani, R. & Totonchy, J. E. An Update of the Virion
Proteome of Kaposi Sarcoma-Associated Herpesvirus. *Viruses* **12**, doi:10.3390/v12121382
(2020).
35   Davison, A. J., Akter, P., Cunningham, C., Dolan, A., Addison, C., Dargan, D. J., Hassan-Walker, A.
F., Emery, V. C., Griffiths, P. D. & Wilkinson, G. W. G. Homology between the human
cytomegalovirus RL11 gene family and human adenovirus E3 genes. *J Gen Virol* **84**, 657-663,
doi:10.1099/vir.0.18856-0 (2003).
36   Dunn, W., Chou, C., Li, H., Hai, R., Patterson, D., Stolc, V., Zhu, H. & Liu, F. Functional profiling of
a human cytomegalovirus genome. *Proc Natl Acad Sci U S A* **100**, 14223-14228,
doi:10.1073/pnas.2334032100 (2003).
37   Osanyinlusi, S. A., Zischke, J., Jacobs, R., Weissinger, E. M., Schulz, T. F. & Kay-Fedorov, P. C.
Human Cytomegalovirus pUL11, a CD45 Ligand, Disrupts CD4 T Cell Control of Viral Spread in
Epithelial Cells. *mBio* **13**, e0294622, doi:10.1128/mbio.02946-22 (2022).
38   Chee, M. S., Bankier, A. T., Beck, S., Bohni, R., Brown, C. M., Cerny, R., Horsnell, T., Hutchison, C.
A., 3rd, Kouzarides, T., Martignetti, J. A. & et al. Analysis of the protein-coding content of the
sequence of human cytomegalovirus strain AD169. *Curr Top Microbiol Immunol* **154**, 125-169,
doi:10.1007/978-3-642-74980-3_6 (1990).
39   Menard, C., Wagner, M., Ruzsics, Z., Holak, K., Brune, W., Campbell, A. E. & Koszinowski, U. H.
Role of murine cytomegalovirus US22 gene family members in replication in macrophages. *J
Virol* **77**, 5557-5570, doi:10.1128/jvi.77.10.5557-5570.2003 (2003).
40   Buisson, M., Geoui, T., Flot, D., Tarbouriech, N., Ressing, M. E., Wiertz, E. J. & Burmeister, W. P.
A bridge crosses the active-site canyon of the Epstein-Barr virus nuclease with DNase and RNase
activities. *J Mol Biol* **391**, 717-728, doi:10.1016/j.jmb.2009.06.034 (2009).
41   Dahlroth, S. L., Gurmu, D., Schmitzberger, F., Engman, H., Haas, J., Erlandsen, H. & Nordlund, P.
Crystal structure of the shutoff and exonuclease protein from the oncogenic Kaposi's sarcoma-
associated herpesvirus. *FEBS J* **276**, 6636-6645, doi:10.1111/j.1742-4658.2009.07374.x (2009).
42   Lee, H., Patschull, A. O. M., Bagneris, C., Ryan, H., Sanderson, C. M., Ebrahimi, B., Nobeli, I. &
Barrett, T. E. KSHV SOX mediated host shutoff: the molecular mechanism underlying mRNA
transcript processing. *Nucleic Acids Res* **45**, 4756-4767, doi:10.1093/nar/gkw1340 (2017).
43   Maninger, S., Bosse, J. B., Lemnitzer, F., Pogoda, M., Mohr, C. A., von Einem, J., Walther, P.,
Koszinowski, U. H. & Ruzsics, Z. M94 is essential for the secondary envelopment of murine
cytomegalovirus. *J Virol* **85**, 9254-9267, doi:10.1128/JVI.00443-11 (2011).
44   Volpato, J. P., Yachnin, B. J., Blanchet, J., Guerrero, V., Poulin, L., Fossati, E., Berghuis, A. M. &
Pelletier, J. N. Multiple conformers in active site of human dihydrofolate reductase F31R/Q35E
double mutant suggest structural basis for methotrexate resistance. *J Biol Chem* **284**, 20079-
20089, doi:10.1074/jbc.M109.018010 (2009).
45   Sarid, R., Flore, O., Bohenzky, R. A., Chang, Y. & Moore, P. S. Transcription mapping of the
Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) genome in a body cavity-based
lymphoma cell line (BC-1). *J Virol* **72**, 1005-1012, doi:10.1128/JVI.72.2.1005-1012.1998 (1998).
46   Nicholas, J., Zong, J. C., Alcendor, D. J., Ciufo, D. M., Poole, L. J., Sarisky, R. T., Chiou, C. J., Zhang,
X., Wan, X., Guo, H. G., Reitz, M. S. & Hayward, G. S. Novel organizational features, captured
cellular genes, and strain variability within the genome of KSHV/HHV8. *J Natl Cancer Inst
Monogr*, 79-88, doi:10.1093/oxfordjournals.jncimonographs.a024179 (1998).
47   Howell, E. E. Searching sequence space: two different approaches to dihydrofolate reductase
catalysis. *Chembiochem* **6**, 590-600, doi:10.1002/cbic.200400237 (2005).

48   Sawaya, M. R. & Kraut, J. Loop and subdomain movements in the mechanism of Escherichia coli
dihydrofolate reductase: crystallographic evidence. *Biochemistry* **36**, 586-603,
doi:10.1021/bi962337c (1997).

49   Barnes, K., Dobrzynski, H., Foppolo, S., Beal, P. R., Ismat, F., Scullion, E. R., Sun, L., Tellez, J.,
Ritzel, M. W., Claycomb, W. C., Cass, C. E., Young, J. D., Billeter-Clark, R., Boyett, M. R. &
Baldwin, S. A. Distribution and functional characterization of equilibrative nucleoside
transporter-4, a novel cardiac adenosine transporter activated at acidic pH. *Circ Res* **99**, 510-519,
doi:10.1161/01.RES.0000238359.18495.42 (2006).

50   Loesing, J. B., Di Fiore, S., Ritter, K., Fischer, R. & Kleines, M. Epstein-Barr virus BDLF2-BMRF2
complex affects cellular morphology. *J Gen Virol* **90**, 1440-1449, doi:10.1099/vir.0.009571-0
(2009).

51   Walston, J. J., Hayman, I. R., Gore, M., Ferguson, M., Temple, R. M., Liao, J., Alam, S., Meyers, C.,
Tugizov, S. M., Hutt-Fletcher, L. & Sample, C. E. The Epstein-Barr Virus Glycoprotein BDLF2 Is
Essential for Efficient Viral Spread in Stratified Epithelium. *J Virol* **97**, e0152822,
doi:10.1128/jvi.01528-22 (2023).

52   Klupp, B. G., Altenschmidt, J., Granzow, H., Fuchs, W. & Mettenleiter, T. C. Identification and
characterization of the pseudorabies virus UL43 protein. *Virology* **334**, 224-233,
doi:10.1016/j.virol.2005.01.032 (2005).

53   El Kasmi, I. & Lippe, R. Herpes simplex virus 1 gN partners with gM to modulate the viral fusion
machinery. *J Virol* **89**, 2313-2323, doi:10.1128/JVI.03041-14 (2015).

54   Wright, N. J. & Lee, S. Y. Structures of human ENT1 in complex with adenosine reuptake
inhibitors. *Nat Struct Mol Biol* **26**, 599-606, doi:10.1038/s41594-019-0245-7 (2019).

55   Massa Lopez, D., Thelen, M., Stahl, F., Thiel, C., Linhorst, A., Sylvester, M., Hermanns-Borgmeyer,
I., Lullmann-Rauch, R., Eskild, W., Saftig, P. & Damme, M. The lysosomal transporter MFSD1 is
essential for liver homeostasis and critically depends on its accessory subunit GLMP. *Elife* **8**,
doi:10.7554/eLife.50025 (2019).

56   Nomburg, J., Doherty, E. E., Price, N., Bellieny-Rabelo, D., Zhu, Y. K. & Doudna, J. A. Birth of
protein folds and functions in the virome. *Nature* **633**, 710-717, doi:10.1038/s41586-024-07809-
y (2024).

57   Davison, A. J. & Stow, N. D. New genes from old: redeployment of dUTPase by herpesviruses. *J
Virol* **79**, 12880-12892, doi:10.1128/JVI.79.20.12880-12892.2005 (2005).

58   Tarbouriech, N., Buisson, M., Seigneurin, J. M., Cusack, S. & Burmeister, W. P. The monomeric
dUTPase from Epstein-Barr virus mimics trimeric dUTPases. *Structure* **13**, 1299-1310,
doi:10.1016/j.str.2005.06.009 (2005).

59   Williams, M. V., Cox, B. & Ariza, M. E. Herpesviruses dUTPases: A New Family of Pathogen-
Associated Molecular Pattern (PAMP) Proteins with Implications for Human Disease. *Pathogens*
6, doi:10.3390/pathogens6010002 (2016).

60   Eberhage, J., Bresch, I. P., Ramani, R., Viohl, N., Buchta, T., Rehfeld, C. L., Hinse, P., Reubold, T.
F., Brinkmann, M. M. & Eschenburg, S. Crystal structure of the tegument protein UL82 (pp71)
from human cytomegalovirus. *Protein Sci* **33**, e4915, doi:10.1002/pro.4915 (2024).

61   Zhao, Z. S., Granucci, F., Yeh, L., Schaffer, P. A. & Cantor, H. Molecular mimicry by herpes simplex
virus-type 1: autoimmune disease after viral infection. *Science* **279**, 1344-1347,
doi:10.1126/science.279.5355.1344 (1998).

62   Soldan, S. S. & Lieberman, P. M. Epstein-Barr virus and multiple sclerosis. *Nat Rev Microbiol* **21**,
51-64, doi:10.1038/s41579-022-00770-5 (2023).

63   Begum, S., Aiman, S., Ahmad, S., Samad, A., Almehmadi, M., Allahyani, M., Aljuaid, A., Afridi, S.
G. & Khan, A. Molecular Mimicry Analyses Unveiled the Human Herpes Simplex and Poxvirus

772   Epitopes as Possible Candidates to Incite Autoimmunity. *Pathogens* **11**,
773   doi:10.3390/pathogens11111362 (2022).

774   64   Lunemann, J. D., Jelcic, I., Roberts, S., Lutterotti, A., Tackenberg, B., Martin, R. & Munz, C.
775   EBNA1-specific T cells from patients with multiple sclerosis cross react with myelin antigens and
776   co-produce IFN-gamma and IL-2. *J Exp Med* **205**, 1763-1773, doi:10.1084/jem.20072397 (2008).

777   65   Heldwein, E. E., Lou, H., Bender, F. C., Cohen, G. H., Eisenberg, R. J. & Harrison, S. C. Crystal
778   structure of glycoprotein B from herpes simplex virus 1. *Science* **313**, 217-220,
779   doi:10.1126/science.1126548 (2006).

780   66   Hekkelman, M. L., de Vries, I., Joosten, R. P. & Perrakis, A. AlphaFill: enriching AlphaFold models
781   with ligands and cofactors. *Nat Methods* **20**, 205-213, doi:10.1038/s41592-022-01685-y (2023).

782   67   Pang, M., He, W., Lu, X., She, Y., Xie, L., Kong, R. & Chang, S. CoDock-Ligand: combined template-
783   based docking and CNN-based scoring in ligand binding prediction. *BMC Bioinformatics* **24**, 444,
784   doi:10.1186/s12859-023-05571-y (2023).

785   68   Baek, M., McHugh, R., Anishchenko, I., Jiang, H., Baker, D. & DiMaio, F. Accurate prediction of
786   protein-nucleic acid complexes using RoseTTAFoldNA. *Nat Methods* **21**, 117-121,
787   doi:10.1038/s41592-023-02086-5 (2024).

788   69   Krishna, R., Wang, J., Ahern, W., Sturmfels, P., Venkatesh, P., Kalvet, I., Lee, G. R., Morey-
789   Burrows, F. S., Anishchenko, I., Humphreys, I. R., McHugh, R., Vafeados, D., Li, X., Sutherland, G.
790   A., Hitchcock, A., Hunter, C. N., Kang, A., Brackenbrough, E., Bera, A. K., Baek, M., DiMaio, F. &
791   Baker, D. Generalized biomolecular modeling and design with RoseTTAFold All-Atom. *Science*
792   **384**, eadl2528, doi:10.1126/science.adl2528 (2024).

793   70   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L.,
794   Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C. C., O'Neill, M., Reiman, D.,
795   Tunyasuvunakool, K., Wu, Z., Zemgulyte, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A.,
796   Cherepanov, A., Congreve, M., Cowen-Rivers, A. I., Cowie, A., Figurnov, M., Fuchs, F. B.,
797   Gladman, H., Jain, R., Khan, Y. A., Low, C. M. R., Perlin, K., Potapenko, A., Savy, P., Singh, S.,
798   Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E. D., Zielinski, M., Zidek, A., Bapst,
799   V., Kohli, P., Jaderberg, M., Hassabis, D. & Jumper, J. M. Accurate structure prediction of
800   biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500, doi:10.1038/s41586-024-
801   07487-w (2024).

802   71   Whisnant, A. W., Jurges, C. S., Hennig, T., Wyler, E., Prusty, B., Rutkowski, A. J., L'Hernault, A.,
803   Djakovic, L., Gobel, M., Doring, K., Menegatti, J., Antrobus, R., Matheson, N. J., Kunzig, F. W. H.,
804   Mastrobuoni, G., Bielow, C., Kempa, S., Liang, C., Dandekar, T., Zimmer, R., Landthaler, M.,
805   Grasser, F., Lehner, P. J., Friedel, C. C., Erhard, F. & Dolken, L. Integrative functional genomics
806   decodes herpes simplex virus 1. *Nat Commun* **11**, 2038, doi:10.1038/s41467-020-15992-5
807   (2020).

808   72   Stern-Ginossar, N., Weisburd, B., Michalski, A., Le, V. T., Hein, M. Y., Huang, S. X., Ma, M., Shen,
809   B., Qian, S. B., Hengel, H., Mann, M., Ingolia, N. T. & Weissman, J. S. Decoding human
810   cytomegalovirus. *Science* **338**, 1088-1093, doi:10.1126/science.1227919 (2012).

811   73   Bencun, M., Klinke, O., Hotz-Wagenblatt, A., Klaus, S., Tsai, M. H., Poirey, R. & Delecluse, H. J.
812   Translational profiling of B cells infected with the Epstein-Barr virus reveals 5' leader ribosome
813   recruitment through upstream open reading frames. *Nucleic Acids Res* **46**, 2802-2819,
814   doi:10.1093/nar/gky129 (2018).

815   74   Arias, C., Weisburd, B., Stern-Ginossar, N., Mercier, A., Madrid, A. S., Bellare, P., Holdorf, M.,
816   Weissman, J. S. & Ganem, D. KSHV 2.0: a comprehensive annotation of the Kaposi's sarcoma-
817   associated herpesvirus genome using next-generation sequencing reveals novel genomic and
818   functional features. *PLoS Pathog* **10**, e1003847, doi:10.1371/journal.ppat.1003847 (2014).

75 Shekhar, R., O'Grady, T., Keil, N., Feswick, A., Amador, D. A. M., Tibbetts, S. A., Flemington, E. K.
& Renne, R. High-density resolution of the Kaposi's sarcoma associated herpesvirus
transcriptome identifies novel transcript isoforms generated by long-range transcription and
alternative splicing. *Nucleic Acids Res* **52**, 7720-7739, doi:10.1093/nar/gkae540 (2024).
76 Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028, doi:10.1038/nbt.3988 (2017).
77 Remmert, M., Biegert, A., Hauser, A. & Soding, J. HHblits: lightning-fast iterative protein
sequence searching by HMM-HMM alignment. *Nat Methods* **9**, 173-175,
doi:10.1038/nmeth.1818 (2011).
78 Hallgren, J., Tsirigos, K. D., Pedersen, M. D., Armenteros, J. J. A., Marcatili, P., Nielsen, H., Krogh,
A. & Winther, O. DeepTMHMM predicts alpha and beta transmembrane proteins using deep
neural networks. *bioRxiv*, doi:10.1101/2022.04.08.487609 (2022).
79 Holm, L. Using Dali for Protein Structure Comparison. *Methods Mol Biol* **2112**, 29-42,
doi:10.1007/978-1-0716-0270-6_3 (2020).
80 Hu, J., Garber, A. C. & Renne, R. The latency-associated nuclear antigen of Kaposi's sarcoma-
associated herpesvirus supports latent DNA replication in dividing cells. *J Virol* **76**, 11677-11687,
doi:10.1128/jvi.76.22.11677-11687.2002 (2002).

Acknowledgements

We would like to thank the Topf lab at CSSB for help in the initial phases of this project, Milot Midarta and
Martin Steinegger for help with Foldseek and for providing links to Foldseek in HerpesFolds and Lisa Holm
for advice in using DALI as a local installation. Finally, we thank Christian Löw for his help in evaluating if
the BMRF2 protein cluster codes for functional transporters. This research was partly supported by the
Maxwell computational resources operated at Deutsches Elektron-Synchrotron DESY, Hamburg,
Germany.

Work in the Bosse lab was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) under Germany's Excellence Strategy EXC 2155—project no. 390874280, the DFG-funded
RTG 2771 Humans and Microbes, project no. 453548970 (Bosse), the DFG-funded RTG 2887, project
number 49735088, by the Wellcome Trust through a Collaborative Award (209250/Z/17/Z) and the Leibniz
ScienceCampus InterACt, funded by the BWFGB Hamburg and the Leibniz Association (W75/2022)
InterACt and "Hamburg- X Infektionsforschung". Moreover, the Bosse and Kaufer labs are both funded
through the DFG Research Unit FOR5200 DEEP-DV (443644894) project BO 4158/5-1 and KA 3492/12-1
awarded to JBB and BBK respectively. BKK is further by the ERC consolidator grant (ERC-CoG ENDo-
HERPES, 101087480).

Author Contributions

TKS: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Software,
Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing

SO: Data curation, Software

SS: Data curation, Software

RS: Data curation, Software

BK: Data curation, Writing – original draft, Writing – review & editing, Funding acquisition, Project
administration

JBB: Conceptualization, Data curation, Funding acquisition, Methodology Project administration,
Supervision, Writing – original draft, Writing – review & editing

Competing Interests

The authors declare no competing interests.

Figure 1. **Structure predictions of the proteome of all human herpesviruses.** A. Pipeline of model generation and quality scoring. The structures of 844 proteins covering the proteomes of all nine human herpesviruses were initially predicted with ColabFold with 3 recycles. After scoring, models that failed any quality score were rerun with AlphaFold and ColabFold with 20 recycles. B. Percentage of each viral proteome that passed the quality score. C. Histogram of the pTM scores of the initial ColabFold models that passed or failed the quality scores. D-E. Histogram of the percentage of each protein with a pLDDT score below 0.50 (D) or above 0.70 (E). In both cases, a Gaussian distribution was fit to identify the distribution for the proteins that passed the quality scores. F. HerpesFolds online database available at https://www.herpesfolds.org/herpesfolds. Structure predictions are organized and searchable by homology and structural similarity. G. Example webpage of the displayed information for each predicted model. H. Searchable clustering of all human herpesvirus proteins based on structure (Foldseek) or sequence (HHblits) similarity is available at https://www.herpesfolds.org/herpesclusters.

Figure 2. **Structural similarity search identifies protein clusters.** A. Graphical display of all structurally similar protein clusters. The 690 herpes proteins that passed the quality score were analyzed pairwise by Foldseek for structural similarity. The color scheme for the viral species is used throughout this work. The name of the protein is written in the center of each node and is legible upon zooming in. Visualization and interaction is recommended at https://www.herpesfolds.org/herpesclusters. B. Cluster of viral serine/threonine-protein kinases US3 and UL13. C. Cluster of viral DNA helicase/primase complex-associated protein (HEPA) and DNA polymerase catalytic subunit (DPOL). D. Clusters of small capsomere-interacting protein (SCP). The alpha-, beta-, and gammaherpesviruses clustered separately. The corresponding UCSF ChimeraX session can be found at https://zenodo.org/records/13284140.

Figure 3. **Structural clustering identifies differently grouped protein groups.** A. Structural alignment of the HCMV UL112-113 cluster. Only the conserved beta-barrel-like domain is shown for clarity of HCMV UL112-113, HHV-6A U79_U80, HHV-6B U79, HHV-7 U79, HSV-1 UL4, HSV-2 UL4, and VZV ORF56. B. Structural alignment of the LANA DNA binding domain-containing cluster with only the matching domain shown for clarity of KSHV ORF73, EBV EBNA1, HHV-6A U84, HCMV UL117, HCMV UL122, HCMV UL123, HHV-6A U86, HHV-6A U90_U87U86, HHV-6B U84, HHV-6B U90_U86, HHV-7 U84, HHV-7 U86, HSV-1 UL3, HSV-2 UL3, and VZV ORF58. C. Ig-like domain-containing cluster harboring the HCMV RL11 protein family. The canonical members are marked with a black outline. D. Structural alignment of RL11 with RL13. The blue arrow points to the additional beta-strand. The corresponding UCSF ChimeraX sessions can be found at https://zenodo.org/records/13284140.

Figure 4. **Domain-level similarity search identifies conserved domain duplications.** A. Domain-level similarities identified using a sliding window approach. The "SW Unique Hits" are the number of proteins that had a structurally similar query-target pair in the sliding window analysis that was absent from the full-length analysis. "Internal Duplications" are proteins where parts of the protein are similar to another part of itself. "Repetitive Acquisition" are proteins where a piece of a different protein matches the query protein multiple times at different positions. "Domain Addition" refers to proteins with multiple domains matching different proteins. Illustrative cartoons are shown above the corresponding bar. B. HSV-1 UL15 is aligned to the known homolog HSV-2 UL15 and the structurally similar target HSV-1 UL9, which was only identified in the domain-level search. The part of HSV-1 UL15 that aligns with HSV-1 UL9 is colored blue. C. The cluster of HCMV proteins that contain HCMV US22. The proteins that contain 2 domains, like the canonical US22, are shown in the HCMV color. The proteins with 1 domain are shown in light red, and the protein that has 4 domains is shown in dark red. D. The tertiary structure of the core domain is similar between the proteins in this group. The top row is the full-length protein, the bottom row is a structural alignment of the core domain, and the RMSD against the 91 Cα of the core domain alignment is written below. HCMV US22 has 2 domains, shown in yellow and blue, which are aligned. HCMV UL29 has 4 domains and was aligned to US22. HCMV US22 was aligned to HCMV US23, which has 2 domains, and HCMV US23 was aligned to HCMV UL26, which has 1 domain. In the case of multiple domains, the best-fitting domain combination is shown. The corresponding UCSF ChimeraX sessions can be found at https://zenodo.org/records/13284140.

Figure 5. **Annotation of viral protein functions from structural similarity search.** A. Structural alignment of the alkaline nuclease (AN) and cytoplasmic envelopment protein 2 (CEP2) from all human herpesviruses. CEP2 was found in all human herpesviruses but had no significant cellular hit. B. Cluster of KSHV ORF2, HSV-1 US2, and HSV-2 US2 containing the cellular hits. Each cellular node (blue) represents a specific protein. C. Summary of B for clarity. D. Structural alignment of the proteins in C. P04753 dihydrofolate reductase (DHFR) was the most significant cellular hit. E. Cluster of EBV BMRF2, KSHV ORF58, HSV-1 UL43, and VZV ORF15 containing the cellular hits. Each cellular node (blue) represents a specific protein. F. Summary of E for clarity. G. Structural alignment of the proteins in F. The most significant cellular hit was Q8R139 equilibrative nucleoside transporter 4 (ENT4). H. Structural architecture of the classical DUT HSV-1 UL50 and the DURP HHV-6A U54. The rainbow-colored viral protein illustrates how nuclepolypeptide folds back on itself. The most significant cellular hit, AF-P70583 DUT from *Rattus norvegicus*, is shown multiple times so that it can be aligned to each domain in the viral protein. The RMSD against the 205 Cα for each alignment is shown on the right of the overlay. The disordered termini were hidden for clarity. The corresponding UCSF ChimeraX sessions can be found at https://zenodo.org/records/13284140.

A proteome-wide structural systems approach reveals insights into protein families of all human herpesviruses

Supplementary Fig. 1. **Developing thresholds for evaluating model quality.** A. pLDDT plot of HCMV UL114 as an example of a protein where the pLDDT curves of the 5 models are consistent. B. pLDDT plot of HHV-7 U29 as an example of a protein where the pLDDT curves of the 5 models are inconsistent. C. Histogram of the standard deviation (StDev) of the mean of the pLDDT for each model for all LocalColabFold models. The threshold (black vertical bar) is set at the mean + 2x StDev, 3.2. D. Histogram of the StDev of the StDev of the pLDDT for each model for all LocalColabFold models. The threshold (black vertical bar) is set at the mean + 2x StDev, 1.9. E. ROC analysis of the pTM of all LocalColabFold models that passed the quality scores versus a negative control harboring scrambled amino acid sequences. The area under the curve is 91.3%. F. pLDDT curve of HHV-7 U29 as an example of a protein that passed the quality score thresholds after AlphaFold was used.

Supplementary Fig. 2. **Herpesviruses express a large fraction of proteins that contain a disordered domain.** A. The percentage of proteins that contain a structured or disordered domain are shown for each species as well as the consolidated proteome. B. HHV-7 U14 is an example of a protein with a structured N-terminus while HHV-6B U90_U86 has a structured C-terminus. The pLDDT and PAE plots are also shown to illustrate how the structured and disordered regions appear. The predicted structure is shown as a rainbow colored ribbon diagram where blue is the N-terminus and red is the C-terminus.

Supplementary Fig. 3. **Alignment of predicted HCMV UL77 structure to the experimental structure that was deposited post-AlphaFold training.** The deposited structure for HCMV UL77 (PDB 7NXP[19]) (blue) was aligned with the AlphaFold prediction (red), and the comparison has an RMSD of 0.509 Å against the 310 Cα. Only the residues present in the deposited structure are shown in the predicted model. The corresponding UCSF ChimeraX session can be found at https://zenodo.org/records/13284140.

Supplementary Fig. 4. **Predicted positive surface charge and DNA binding activity of the KSHV ORF73 and EBV EBNA1 cluster.** The proteins in this cluster were structurally predicted as dimers of the conserved core with the DNA binding sequence of KSHV ORF73[80] with AlphaFold 3. The crystal structure of KSHV ORF73 with DNA (PDB: 4UZB[29]) was included for comparison. The average pairwise ipTM between each protein and DNA strand is shown above the structure. The dashed line separates the different binding phenotypes: KSHV ORF73, HCMV UL122 (different orientation), and HCMV UL117 (no binding). Positive surface charge is displayed as blue, and negative charge is displayed as red. The corresponding UCSF ChimeraX session can be found at https://zenodo.org/records/13284140.

Supplementary Fig. 5. **Alignment of E. coli DHFR to KSHV ORF2 and HSV-1 US2.** A. The *E. coli* DHFR (PDB 1RX2[48]) (blue) was aligned to KSHV ORF2 (purple) and HSV-1 US2 (green), and the cofactor folic acid is shown in black. The atomic structure of the catalytic residues of the *E. coli* DHFR (M20, P21, W22, D27, F31, R44, R57, G95, G96, T113) and the potential catalytic residues that are in similar positions in ORF2 and US2 are shown. B. An enlarged view of the catalytic site is shown. C. The protein sequences were aligned based on the structure-based alignment or with Clustal Omega. The *E. coli* catalytic residues are marked with red asterisks. The potential catalytic residues in the herpesvirus proteins are from the structured based alignment. The corresponding UCSF ChimeraX session can be found at https://zenodo.org/records/13284140.

Supplementary Fig. 6. **The herpesvirus ENTs are unlikely to have transporter activity.** A. Cross-sections of surface plots of the monomer predictions of ENT4 (AF-Q8R139-F1-v4) and the structurally similar viral proteins. The white arrow points to the metabolite channel in ENT4. B. Structure predictions of the known EBV (BMRF2 pink and BDLF2 white) and KSHV (ORF58 purple and ORF27 light purple) heterodimers. The corresponding UCSF ChimeraX sessions can be found at https://zenodo.org/records/13284140.