MLL4 regulates postnatal palate growth and midpalatal suture development

Jung-Mi Lee[1,*], Hunmin Jung[1], Bruno de Paula Machado Pasqua[1], Yungki Park[2], Qinghuang Tang[1], Shin Jeon[3,4], Soo-Kyung Lee[4], Jae W. Lee[4], and Hyuk-Jae Edward Kwon[1,*]

[1]Department of Oral Biology, School of Dental Medicine, University at Buffalo, The State University of New York, Buffalo, NY 14214, U.S.A.;
[2]Department of Biochemistry, Jacobs School of Medicine and Biomedical Sciences, Institute for Myelin and Glia Exploration, University at Buffalo, The State University of New York, Buffalo, NY 14203, U.S.A.;
[3]Department of Biological Sciences, College of Arts and Sciences, FOXG1 Research Center, University at Buffalo, The State University of New York, Buffalo, NY 14260, U.S.A.;
[4]Department of Systems Pharmacology & Translational Therapeutics, Institute for Immunology, University of Pennsylvania, PA 19104, U.S.A.

*Corresponding authors:

Hyuk-Jae Edward Kwon, D.D.S., Ph.D.
Department of Oral Biology
School of Dental Medicine
University at Buffalo, The State University of New York
547 Biomedical Research Building
Buffalo, NY 14214-8024, U.S.A.
hekwon@buffalo.edu

Jung-Mi Lee, Ph.D.
Department of Oral Biology
School of Dental Medicine
University at Buffalo, The State University of New York
540 Biomedical Research Building
Buffalo, NY 14214-8024, U.S.A.
jungmile@buffalo.edu

1 **Abstract**
2

MLL4, also known as KMT2D, is a histone methyltransferase that acts as an important epigenetic regulator in

various organogenesis programs. Mutations in the *MLL4* gene are the major cause of Kabuki syndrome, a

human developmental disorder that involves craniofacial birth defects, including anomalies in the palate. This

study aimed to investigate the role of MLL4 and the underlying mechanisms in the development and growth of

the palate. We generated a novel conditional knockout (cKO) mouse model with tissue-specific deletion of *Mll4*

in the palatal mesenchyme. Using micro-computed tomography (CT), histological analysis, cell mechanism

assays, and gene expression profiling, we examined palate development and growth in the *Mll4*-cKO mice.

Gross craniofacial examination at adult stages revealed mild midfacial hypoplasia and midline defects of the

palate in *Mll4*-cKO mice, including a widened midpalatal suture and disrupted midline rugae pattern. Micro-CT-

based time-course skeletal analysis during postnatal palatogenesis through adulthood demonstrated a

transverse growth deficit in overall palate width in *Mll4*-cKO mice. Whole-mount and histological staining at

perinatal stages identified that the midline defects in the *Mll4*-cKO mice emerged as early as one day prior to

birth, presenting as a widened midpalatal suture, accompanied by increased cell apoptosis in the suture

mesenchyme. Genome-wide mRNA expression analysis of the midpalatal suture tissue revealed that MLL4 is

essential for the timely expression of major cartilage development genes, such as *Col2a1* and *Acan*, at birth.

Immunofluorescence staining for osteochondral differentiation markers demonstrated a marked decrease in

the chondrogenic marker COL2A1, while the expression of the osteogenic marker RUNX2 remained

unchanged, in the *Mll4*-cKO midpalatal suture. Additionally, SOX9, a master regulator of chondrogenesis,

exhibited a significant decrease in protein expression. Indeed, time-course histological analysis during

postnatal palate growth revealed retardation in the development of the suture cartilage in *Mll4*-cKO mice.

Taken together, our results demonstrate that MLL4 is essential for orchestrating key cellular and molecular

events that ensure proper midpalatal suture development and palate growth.

Introduction

The palate is a central structure in the craniofacial complex, playing a crucial role in breathing, feeding, speech, and maintaining overall craniofacial health. Thus, palatogenesis—the development and growth of the palate—is a vital process in establishing the craniofacial complex. Palatogenesis can be broadly divided into two phases: embryonic palatogenesis and postnatal palatogenesis, which occurs very similarly in humans and mice (Dixon et al., 2011). The first phase, embryonic palatogenesis, occurs during the early stages of fetal development and involves the initial formation of a pair of palatal shelves, which are bilateral outgrowths from the left and right maxillary processes, at embryonic day (E) 11 in mice. These bilateral palatal shelves then undergo remodeling, grow toward the midline, and fuse with each other between E12 and E15, establishing the foundational structure of the palate (Bush and Jiang, 2012). The second phase, postnatal palatogenesis, involves skeletal growth, which occurs along the anteroposterior and transverse directions, with the palate reaching its full adult dimension by 5 weeks of age in mice. A key feature in postnatal palatogenesis is the role of the palatal sutures, which remain as active growth sites postnatally, allowing for the transverse growth of the palate and the accommodation of the growing craniofacial complex during the growth spurts of childhood and adolescence (Griffiths et al., 1967, Latham, 1971). For example, infant patients that undergo surgical repair of cleft palate frequently exhibit signs of midfacial growth arrest, whereas those with unrepaired cleft palate often display normal jaw dimensions and occlusion (Fudalej et al., 2008, Mølsted et al., 2005, Brattström et al., 2005, Wolford et al., 2008, Ye et al., 2012). Similarly, in developing mice, surgical disruption of the midpalatal suture can result in a narrow palate and midfacial hypoplasia (Li et al., 2015). Both phases are crucial for the functional integrity of the palate, and their regulation is orchestrated by a complex interplay of genetic and epigenetic factors, ensuring the precise expression of developmental genes at the right time and place (Li et al., 2017). Any disruption in these factors may lead to severe craniofacial birth defects such as orofacial clefts and midfacial hypoplasia (Garland et al., 2020, Reynolds et al., 2020, Cox, 2004, Suzuki et al., 2016). This direct clinical relevance has driven substantial research efforts to understand embryonic palatogenesis. In contrast, postnatal palatogenesis remains significantly understudied, especially regarding the underlying mechanisms, which has contributed to disparity in research focus. Thus, there is a critical need to expand

research into postnatal palatogenesis to address issues related to functional impairments and orthodontic concerns that can arise later in life and to better understand the complete picture of craniofacial development.

Epigenetic regulation, including histone modification, plays a pivotal role in orchestrating craniofacial development and growth (Shull and Artinger, 2024). By regulating chromatin structure and accessibility, epigenetic regulation ensures precise temporal and spatial expression of genes essential for establishing the craniofacial structures in health and disease, influencing cellular mechanisms, such as proliferation, migration, and differentiation. In this regard, mixed lineage leukemia 4 (MLL4), also known as histone-lysine *N*-methyltransferase 2D (KMT2D), is a plausible candidate that may serve as a key epigenetic regulator in palatogenesis. MLL4 is a histone H3 lysine 4 (H3K4) methyltransferase that plays a crucial role in modulating gene expression patterns that are fundamental to the establishment and maintenance of cell identity and lineage specificity during organogenesis (Froimchuk et al., 2017). Mutations in *MLL4* cause Kabuki syndrome, which is a human developmental disorder characterized by distinctive facial features including eversion of lower lateral eyelids and elongated palpebral fissures, postnatal growth retardation, intellectual disability, and skeletal, craniofacial, and dermatoglyphic abnormalities (Boniel et al., 2021). Within the craniofacial complex, Kabuki syndrome is associated with orofacial anomalies, including various palate anomalies and midfacial dysplasia (Matsune et al., 2001). Previous studies using various animal models have shown that MLL4 plays an important role in craniofacial development by regulating neural crest formation, migration, and differentiation (Bögershausen et al., 2015, Schwenty-Lara et al., 2020, Van Laarhoven et al., 2015). Notably, mice with tissue-specific deletion of *Mll4* in the neural crest cell-lineage, driven by either *Wnt1-Cre* or *Sox10-Cre*, showed cleft palate (Shpargel et al., 2020). This suggests that palatal shelf fusion during embryonic palatogenesis requires MLL4 function in the cranial neural crest-derived cells, which give rise to the palate mesenchyme. However, deletion of *Mll4* in these mice occurs considerably earlier than the initiation of palatogenesis (at E11). Therefore, it remains uncertain whether the cleft palate phenotype is a result of *Mll4* deficiency during or prior to the initiation of palatogenesis. Moreover, due to the neonatal lethality, these neural crest-specific *Mll4* deletion mice were limited to investigating only the embryonic palatogenesis. Thus, there is a critical need to understand the comprehensive role and underlying mechanism of MLL4 in palatogenesis in a time- and tissue-specific manner.

In this study, to elucidate the comprehensive role and the underlying mechanism by which MLL4 regulates palatogenesis, we generated a novel mouse model with palate mesenchyme- specific and palatogenesis-specific *Mll4* deletion (*Mll4^{fl/fl};Osr2-Cre*). Using micro-computed tomography (CT), histology, cell mechanism assays, and gene expression analysis approaches, we examined the development and growth of the palate in our mouse model. Our results revealed that *Mll4* deficiency results in retardation in the developmental of the midpalatal suture and transverse growth deficits during postnatal palatogenesis. Our study provides critical insights into the intrinsic role of MLL4 in palatogenesis, which informs the physiological developmental mechanism and suggests therapeutic strategies for congenital orofacial anomalies.

Materials and Methods

Mouse Strains

Osr2-Cre (originally described as '*Osr2KI*-Cre') (Chen et al., 2009, Shpargel et al., 2020), and *Mll4*$^{fl/fl}$ (Lee et

al., 2013) lines were used for generating an *Mll4* conditional knockout (cKO) model. *ROSA*$^{mT/mG}$ (hereafter

referred to as *mTmG*) reporter line (Muzumdar et al., 2007) was used to label palate mesenchyme cells

expressing eGFP by crossing with *Osr2-Cre* line. Mice were maintained by intercrossing or by crossing with

C57BL/6J inbred mice. All animal procedures were approved by the Institutional Animal Care and Use

Committee at the University at Buffalo.

Micro-CT Scanning and 3-Dimensional Reconstruction

The heads of mice aged 1, 3, 6, and 10 weeks were fixed in 10% formalin overnight and subsequently washed

with PBS. Micro-CT scans were performed using a SCANCO μCT 100 (SCANCO Medical, Switzerland) (voxel

size, 10 μm). The scanned images were calibrated to milligrams of hydroxyapatite per cubic centimeter

(mgHA/cm^3) for accurate density measurement. Each image was manually adjusted for rotation and alignment

along each axis using the FIJI software (Schindelin et al., 2012). To eliminate background noise, areas with

measurements below 200.03 mgHA/cm^3 were removed. Each image was then cropped to focus on the specific

region of interest within the maxillary jaw. The Fijiyama plugin was used to manually register the images,

aligning them for further analysis (Fernandez and Moisy, 2021). Finally, the 3Dscript plugin was used for 3-

dimensional (3D) rendering (Schmid et al., 2019).

Histology and Immunofluorescence Staining

Mice were euthanized at predetermined stages. Head tissue specimens were fixed in 4% paraformaldehyde

(PFA), dehydrated through an ethanol series, embedded in paraffin, and sectioned at 5–7 μm thickness, as

described previously (Lee et al., 2022). Specimens harvested at E18.5, postnatal day (P) 0.5, and 1, 3, 6, and

10 weeks were decalcified in 14% ethylenediaminetetraacetic acid (EDTA) for 1, 3, 6, and 10 weeks,

respectively. For histological analysis, sections were stained with Azan's trichrome staining, as previously

described (Lee et al., 2022), and were imaged on an Axioscope compound upright light microscope (for 20X and 40X magnifications) (Zeiss, Germany). An SZX16 stereomicroscope (Olympus, Japan) was used for 5X magnification images, as well as gross images. For immunofluorescence staining, antigen retrieval was performed in an antigen retrieval buffer, using a pressure cooker for 15 min at low pressure, followed by washing in PBS, blocking at room temperature in a blocking buffer (2% goat serum, 5% BSA, 1% Triton-X 100, 0.1% Tween-20 in PBS) for 1 hr, and incubation with primary antibodies, diluted in blocking buffer, overnight at 4°C. On the second day, the sections were washed in PBS, incubated with secondary antibodies, diluted in blocking buffer, at room temperature for 1 hr, washed in PBS, and mounted with Vectashield mounting media with DAPI (Vector Laboratories, USA; H-1200). The primary antibodies used were: homemade guinea pig anti-MLL4 (1:300) (Huisman et al., 2021), rabbit monoclonal anti-SOX9 (1:300; Abcam, UK; ab185230), mouse monoclonal anti-COL2A1 (1:50, Santa Cruz, USA; sc-52658), and rabbit monoclonal anti-RUNX2 (1:300; Cell Signaling, USA; #12556). Immunofluorescence images were acquired using a THUNDER Imager fluorescence microscope (Leica Microsystems, Germany). To eliminate non-specific signals from auto-fluorescence, we acquired additional images usng a non-stained CFP (cyan) filter and subtracted these signals following an established scientific method (Van de Lest et al., 1995).

TUNEL Assay

The DeadEnd™ Fluorometric TUNEL System (Promega, USA; G3250) was used to assess apoptosis in paraffin sections, according to the manufacturer's protocol.

Skeletal Preparations

Mouse heads, harvested at predetermined stages, were fixed in 100% ethanol for 3 days, stained in Alcian Blue solution (30 mg Alcian Blue dissolved in 20 ml glacial acetic acid plus 80 ml 95% ethanol) for 2 days, re-fixed in 100% ethanol for 24 hr, cleared in 2% KOH for 3 hr, and counterstained in Alizarin Red solution (50 mg Alizarin Red dissolved in 1 L of 1% KOH) for 24 hr, followed by tissue clearing in 1% KOH/20% glycerol for 3 days. Images were acquired using a standard stereomicroscope (Amscope, USA).

RNA Sequencing and Bioinformatics Analysis

Total RNA was extracted from micro-dissected midpalatal suture tissue samples using the RNeasy Micro Plus Kit (Qiagen, Germany; 74034). Four biological replicates per genotype (control and *Mll4*-cKO), were used for sequencing, with one sample in each genotype pooled from three individuals. Library construction and high-throughput sequencing were performed by Novogene company (China) using the Illumina NovaSeq 6000 platform. Sequencing reads were analyzed using Salmon (Patro et al., 2017) to quantify gene expression at the transcript level. The resulting outputs from Salmon were analyzed using DESeq2 (Love et al., 2014) to obtain gene-level expression data and identify differentially expressed genes (DEGs) between *Mll4*-cKO and littermate control samples. DEGs were defined by a log2 fold change (log2FC) $\geq$ 0.4 and an adjusted *p*-value $\leq$ 0.001. Volcano plots were generated using the EnhancedVolcano package (Blighe et al., 2024), and gene ontology (GO) analysis was performed with clusterProfiler (Wu et al., 2021). Raw expression data results are included in the Supplementary Material. The original RNA sequencing (RNA-seq) data files have been deposited in the National Center for Biotechnology Information Gene Expression Omnibus (NCBI GEO) database under the accession number PRJNA1148613.

ChIP Sequencing Re-Analysis

The ChIP-sequencing (ChIP-seq) datasets for UTX and H3K4me3 in cranial neural crest cells (GSE103849) were re-analyzed using a previously described data processing workflow (Shpargel et al., 2017). Briefly, sequencing reads were aligned to the mouse mm10 reference genome using Bowtie2 (Langmead and Salzberg, 2012) and converted to BAM files using SAMtools (Li et al., 2009). Reads mapping to the X and Y chromosomes were excluded. Filtered BAM files from biological replicates were merged and subjected to peak calling with MACS (Zhang et al., 2008). Peaks were annotated using the ChIPseeker R package (Yu et al., 2015). For visualization in Integrative Genomics Viewer (IGV), the filtered BAM files were further processed using BEDTools (Quinlan and Hall, 2010) and converted to BigWig format using the UCSC command-line tool bedGraphToBigWig (Kent et al., 2002). The intersection of *Mll4*-cKO DEGs with UTX and H3K4m3 target genes, along with functional enrichment analysis, was performed in R using clusterProfiler (Wu et al., 2021).

Statistics

All statistical analyses were performed using Prism 10 software (GraphPad Software, USA). Pairwise differential expression was analyzed using Student's t-test. Statistical significance was defined as p-values $\leq$ 0.05 (*), $\leq$ 0.01 (**), and $\leq$ 0.001 (***).

Results

Generating a novel *Mll4* deficiency model for studying the palate tissue-autonomous role of MLL4

during palate development and growth

For the purpose of investigating the role of MLL4 specifically during palatogenesis, we used the Cre/loxP

system driven by *Osr2-Cre*. *Osr2-Cre* is first activated at E11, when palatogenesis is initiated, and remains

active in the palate mesenchyme cells (Lan et al., 2001, Lan et al., 2004). The *Osr2-Cre;mTmG* line allows us

to confirm the activation of *Osr2-Cre* by detecting GFP signals using fluorescence microscopy. When the

bilateral palatal shelves are actively undergoing morphogenesis, for example at E13.5, *Osr2-Cre* was activated

in the developing palatal shelves, which occupies the center of the developing face (Fig. 1A, B). Thus, we were

able to generate a novel mouse model with palatogenesis-specific deletion of *Mll4* in the palate mesenchyme

cells (*Mll4$^{fl/fl}$;Osr2-Cre*, or *Mll4*-cKO). Surprisingly, compared to the previously reported neural crest-specific

Mll4 deletion (*Mll4$^{fl/fl}$;Wnt1-Cre* and *Mll4$^{fl/fl}$;Sox10-Cre*) mice, which exhibited cleft palate and neonatal lethality

(Shpargel et al., 2020), our palatogenesis-specific *Mll4*-cKO mice did not show cleft palate or lethality at birth

and could grow up into adult mice that did not show much noticeable changes when grossly examined for

exterior appearance at 10 weeks (Fig. 1C, G). However, close examination of the orofacial skeleton

dimensions by using micro-CT analysis revealed that the facial angle, length, and width in the maxilla showed

mild but significant decreases in the *Mll4*-cKO mice (Fig. 1D–F, H–J, N), which were signs of mild midfacial

hypoplasia. Additional examination of the intra-oral aspect of the palate revealed several minor differences

between the control and *Mll4*-cKO mice (Fig. 1K, L). Compared to controls, the midpalatal suture appeared

more widened in the *Mll4*-cKO palate (Fig. 1K′, L′, arrowheads), with minor changes observed in the rugae

patterning (Fig. 1K′, L′, arrows). To test the possibility that these mild midfacial hypoplasia and palate

phenotypes that we observed in the *Mll4*-cKO mice may have resulted from delay in their overall development

and growth, we measured the body weight both at 10 weeks and at birth but did not find any significant

difference (Fig. 1M). We also compared the body weight between male and female mice at these two stages

but found no significant difference (data not shown). Our initial observations suggested that, separate from its

early role in neural crest cell differentiation, MLL4 may have other unknown functions in postnatal palate

development and growth.

Thus, to gain better understanding of the midfacial and palate phenotypes observed from gross

examination of the *Mll4*-cKO mice (Fig. 1), we further examined the palate skeleton at 10 weeks using our

micro-CT results. By examining the virtual 3D models that were created by reconstructing the planar X-ray

images, we found that the overall palate width showed narrowing in the *Mll4*-cKO mice, which was consistent

with the decrease in the facial width that we observed earlier. Anatomical landmarks for assessing the palate

width, such as the incisive foramens, palatal processes of maxilla, and palatal processes of palatine, all

showed constricted appearances in the virtual 3D model of the *Mll4*-cKO mice (Fig. 2A, B). Using virtual slicing

of the 3D models, we were able to further generate digital coronal/frontal cross sections of the palate. These

virtual cross sections enabled us to perform comprehensive quantitative analysis of the overall horizontal

dimension of the palate along its anteroposterior axis by precisely measuring the palate width at four

representative section levels: the incisive foramen (in slice 1), the palatal processes of maxilla (in slice 2), the

transpalatal suture (in slice 3), and the palatal processes of palatine (in slice 4) (Fig. 2C, D). At all four section

levels along the anteroposterior axis, the palate width showed significant decrease in the *Mll4*-cKO mice,

compared to the control mice (Fig. 2E1–E4). These results suggested that MLL4 plays an important role in

transverse growth of the palate during postnatal palatogenesis.

Thus, to address the question of whether MLL4 is actually required for proper growth of the palate

during postnatal palatogenesis, we conducted a time-course quantitative analysis of the horizontal dimension

of the palate skeleton using micro-CT-generated virtual coronal/frontal cross sections at 1, 3, and 6 weeks. For

this approach, we used the same four representative coronal/frontal cross section levels along the

anteroposterior axis of the palate, which we used at 10 weeks. At 1 week, we found that palate growth was not

significantly changed in the *Mll4*-cKO mice at the incisive foramen level (Fig. 2E1). In contrast, at the rest of the

levels, the first (M1), second (M2), and third molar (M3) levels, palate growth showed minor, but significant

decrease (Fig. 2E2–E4). Later, at 3, 6, and 10 weeks, palate growth showed significant decrease at all four

levels along the anteroposterior axis in the *Mll4*-cKO mice, compared to the control mice (Fig. 2E1–E4). These

results suggested that MLL4 is indeed required for proper transverse growth of the palate during postnatal

palatogenesis.

MLL4 is essential for the initial formation of the midpalatal suture around birth

We next asked the question of how MLL4 promotes palate growth. Based on our finding that the *Mll4*-cKO

mice showed significant narrowing of the palate during postnatal palatogenesis (Fig. 2), we reasoned that the

key biological events for palate growth that depend on MLL4 function may occur at younger ages, possibly as

early as, or even prior to, birth. Thus, to identify the primary developmental mechanism underlying the role of

MLL4 in palate growth, we examined the palate anatomy in the control and *Mll4*-cKO mice around birth,

searching for the earliest signs of disruption in the *Mll4*-cKO palate. By first using skeletal staining of the

developing cranium and palate tissues, we found that the overall skeletal pattern was similar between the

control and *Mll4*-cKO mice (Fig. 3A, B). However, where we had initially observed midline defects in the rugae

pattern and a widened midpalatal suture in the *Mll4*-cKO mice (Fig. 1), we found that the width of the

midpalatal suture was significantly wider in the *Mll4*-cKO mice, compared to the controls, prior to birth (at

E18.5) (Fig. 3A, B, G). In parallel, histological examination of the palate, by using trichrome staining of the

coronal/frontal paraffin sections, confirmed that the width of the midpalatal suture, situated between the

bilateral bony palatal processes, was significantly increased in the *Mll4*-cKO mice, compared to the control

mice (Fig. 3C, D, C′, D′, G). To determine the cellular mechanisms underlying the wider midpalatal suture in

the *Mll4*-cKO mice, we performed both cell proliferation, using EdU incorporation assays, and cell apoptosis,

using TUNEL staining. While rarely any cells in the midpalatal suture were EdU-positive for cell proliferation in

both the control and *Mll4*-cKO mice (data not shown), TUNEL-positive cell apoptosis showed minor but

significant increase in the *Mll4*-cKO midpalatal suture, compared to controls, at E18.5 (Fig. 3E, F, E′, F′, H).

Our findings that the earliest signs of disrupted biological events were specifically discovered in the midpalatal

suture of the perinatal *Mll4*-cKO palate suggested that MLL4 is essential for the establishment of the midpalatal

suture around birth.

Mll4 **deficiency results in dysregulation of timely expression of chondrogenic genes at birth**

To gain insight into the molecular mechanism underlying the histological and cellular alterations observed in the perinatal *Mll4*-cKO palate tissue, we micro-dissected the midpalatal suture tissues and performed RNA-seq analysis on control and *Mll4*-cKO mice at birth (P0.5). Our results identified a total of 117 DEGs between the control and *Mll4*-cKO midpalatal suture tissues, with 40 genes upregulated and 77 genes downregulated in the *Mll4*-cKO group (adjusted *p*-value ≤ 0.001, log2FC ≥ 0.4) (Fig. 4A). Notably, among the DEGs were genes

associated with cartilage development, or chondrogenesis, including extracellular matrix genes *Col2a1*, *Col9a1*, *Col9a2*, *Acan* (Aggrecan), and *Comp* (Cartilage oligomeric matrix protein), as well as *Mia/Cdrap1* (Melanoma inhibitory activity) and *Snorc* (Small novel rich in cartilage). GO analysis identified the top 10 most enriched biological processes, along with a list of downregulated genes associated with each process (Fig. 4B–D). This analysis suggested that cartilage development, endochondral bone morphogenesis, and chondrocyte differentiation are among the highly enriched biological processes potentially regulated by MLL4.

In parallel, we selected several representative markers of osteochondral progenitor cell differentiation during chondrogenesis and endochondral ossification for immunofluorescence staining on paraffin sections of the perinatal palate (at P0.5) in control and *Mll4*-cKO mice. Among these markers, COL2A1 was selected from the DEG list (Fig. 4) to validate our RNA-seq results. Additionally, RUNX2 and SOX9 were chosen as critical transcription factors regulating osteochondral progenitor cell differentiation: RUNX2 marks osteoblast differentiation, while SOX9 marks chondrocyte differentiation, providing insights into the developmental processes occurring in the midpalatal suture. We also verified the expression of MLL4 in the perinatal *Mll4*-cKO palate using immunofluorescence staining. Our results revealed that the expression of COL2A1 and SOX9, both early markers of chondroblast differentiation, was markedly reduced in the *Mll4*-cKO midpalatal suture (Fig. 5B1, D2, E), compared to controls (Fig. 5A1, C2, E). In contrast, the expression of RUNX2, an osteogenic differentiation marker, was similar between the control and *Mll4*-cKO midpalatal suture (Fig. 5A2, B2, E). These findings suggest that the primary molecular events dependent on MLL4 function in the perinatal midpalatal suture are associated with chondrogenic cell differentiation, cartilage development, and potentially endochondral ossification. Additionally, the markedly reduced intensity of MLL4 in the perinatal *Mll4*-cKO palate confirms the effective deletion of the *Mll4* gene in the *Mll4*-cKO palate (Fig. 5C1, D1).

***Mll4* deficiency leads to retardation of the development of the midpalatal suture**

Finally, to verify whether the biological processes that were identified from our RNA-seq analysis were actually

impacted as a result of *Mll4* deficiency during postnatal palatogenesis, we performed histological examination

of the midpalatal suture at birth and beyond (up to 1 week), using trichrome staining of the paraffin sections

(Fig. 6A–D). Interestingly, after birth (at P0.75), a newly formed layer of cartilage tissue capped the medial

surface of each of the bilateral palatal processes, flanking the midpalatal suture in the control mice (Fig. 6A,

arrowheads). In contrast, the newborn *Mll4*-cKO mice lacked these midpalatal suture-flanking cartilage tissue

layers on the medial surfaces of the bilateral palatal processes (Fig. 6B). By histologically monitoring the

postnatal development of the midpalatal suture, we found that, at 1 week in the control mice, the midpalatal

suture cartilage layers showed uniform enlargement of the consisting chondrocytes (Fig. 6C, arrowheads). In

contrast, the 1-week old *Mll4*-cKO mice just started to develop the midpalatal suture cartilage layers, which

were only partially formed (Fig. 6D, arrowheads). Our findings suggested that this defective phenotype

observed in the *Mll4*-cKO midpalatal suture cartilage was due to significant retardation of chondrogenic

differentiation as a result of *Mll4* deficiency in the developing palate. Thus, we further performed additional

time-course examination of the midpalatal suture development at 3, 6, and 10 weeks. Interestingly, at all three

stages (3, 6, and 10 weeks), the bilateral cartilage layers in both the control and *Mll4*-cKO midpalatal sutures

were shown to be adhered to each other in the midline (Fig. 6E–J). However, the interface between the

bilateral cartilage layers was less straight in the *Mll4*-cKO suture, compared to the control suture (Fig. 6E–J).

Also at all three stages, the overall width of the midpalatal suture cartilage was markedly increased in the *Mll4*-

cKO mice (brackets in Fig. 6F, H, J) than in the control mice (brackets in Fig. 6E, G, I). Importantly, compared

to the horizontally organized chondrocyte 'columns' in the control group, the *Mll4*-cKO suture chondrocytes

appeared disorganized and were under-differentiated, especially at 10 weeks (Fig. 6I, J, I′, J′). Taken together,

these results suggest that *Mll4* deficiency results in retardation in the developmental of the midpalatal suture

during postnatal palatogenesis. And by integrating our findings in the developing midpalatal suture (Fig. 6) and

in the growing palate skeleton (Fig. 2), our data suggest that *Mll4* is associated with midpalatal suture

development and palate growth during postnatal palatogenesis.

Discussion

A novel role of MLL4 in determining palate width during postnatal palatogenesis using a newly

developed *Mll4* deficiency/Kabuki syndrome mouse model

In this study, we generated for the first time a unique mouse model demonstrating that MLL4 plays an essential

role in determining palate width (Fig. 1). By using *Osr2-Cre* to delete *Mll4* tissue-specifically in the palate

mesenchyme from the onset of palatogenesis (at E11) (Lan et al., 2001, Lan et al., 2004), we examined the

intrinsic role of MLL4 during the active process of palatogenesis (Fig. 2). Unlike previously reported neural

crest-specific *Mll4* deletion models driven by *Wnt1-Cre* or *Sox10-Cre*, which exhibit cleft palate and neonatal

lethality (Shpargel et al., 2020), our *Mll4*-cKO mice did not show cleft palate or neonatal lethality. This allowed

us to extend our examinations beyond prenatal stages. Importantly, the narrow palate phenotype observed in

our *Mll4*-cKO mice closely resembles a common craniofacial phenotype of Kabuki syndrome: the high-arched,

narrow palate, reported in 72% (64/89) of individuals with Kabuki syndrome (Matsumoto and Niikawa, 2003,

Porntaveetus et al., 2018). This prevalence is twice as high as the 35% (68/196) frequency of cleft lip/palate

individuals in Kabuki syndrome (Matsumoto and Niikawa, 2003), which was studied using the aforementioned

neural crest-specific *Mll4* deletion models (Shpargel et al., 2020). Our findings further emphasize the clinical

relevance of the narrow palate phenotype, especially when accompanied by midfacial hypoplasia, which poses

potential risks for various dental and orofacial complications, such as crowded or impacted teeth, breathing

difficulties, and further midfacial underdevelopment. Taken together, our unique *Mll4*-cKO model not only

highlights a significant craniofacial problem in Kabuki syndrome but also sheds light on an understudied yet

critical developmental process that warrants urgent attention. These results underscore the multifaceted role of

MLL4 in craniofacial development and disease.

MLL4 is essential for proper development of the midpalatal suture

Here, we establish MLL4 as a crucial factor in midpalatal suture development, as evidenced by the significant

developmental delays observed in the *Mll4*-cKO mice beginning in the perinatal stages (Fig. 3) and persisting

throughout postnatal palatogenesis into adulthood (Fig. 6). Whole gene expression profiling of midpalatal

suture tissues revealed that *Mll4* is required for the timely expression of a large number of genes related to

chondrogenesis, including *Col2a1* (Bell et al., 1997, Lefebvre et al., 1997, Ng et al., 1997), *Col9a1* (Genzer

and Bridgewater, 2007, Zhang et al., 2003, Zhou et al., 1998), *Col9a2* (Bernard et al., 2003), *Acan* (Sekiya et

al., 2000, Han and Lefebvre, 2008), *Comp* (Liu et al., 2007), *Mia*/*Cdrap1* (Xie et al., 1999), and *Snorc* (Jaiswal

et al., 2020) (Fig. 4). Interestingly, *Sox9* did not appear in our DEG list, possibly due to (1) variability introduced

during tissue micro-dissection, given its small size of the midpalatal suture and the potential for contamination

from adjacent tissues, or (2) the relatively low steady-state expression level of *Sox9* compared to other

chondrogenic genes (e.g., RPKM values: *Sox9* = 37.6 vs. *Col2a1* = 787.8 and *Acan* = 308.2), which may

obscure detectable differences. Nevertheless, immunofluorescence analysis confirmed that SOX9 expression

was markedly downregulated in the mesenchyme cells in the *Mll4*-cKO midpalatal suture (Fig. 5), suggesting

that MLL4 may indeed regulate *Sox9* expression.

Notably, the seven downregulated chondrogenic genes in the Mll4-cKO midpalatal suture mesenchyme

are all known SOX9 targets, highlighting its role as a master regulator of chondrogenesis (Lefebvre et al.,

2019). Recent studies suggest that SOX9 functions as a pioneer factor, recruiting epigenetic co-factors such

as MLL4 to epigenetically prime and remodel chromatin to a transcriptionally accessible state during cell fate

determination (Yang et al., 2023). Therefore, it is reasonable to speculate that *Mll4* deficiency may affect

midpalatal suture development through two mechanisms: (1) lack of MLL4 as a transcriptional co-activator of

SOX9 and (2) downregulation of SOX9 itself. Both mechanisms would impair SOX9's ability to activate

downstream chondrogenic genes, as observed in the *Mll4*-cKO midpalatal suture. Furthermore, the known

auto-regulatory function of SOX9 in chondrocytes (Mead et al., 2013) could exacerbate the chondrogenesis

defects in our model.

To further explore the potential mechanism by which MLL4 regulates chondrogenic gene expression,

we re-analyzed published UTX and H3K4me3 ChIP-seq datasets in O9-1 cranial neural crest cells (Shpargel et

al., 2017). Given that MLL4 coordinates with UTX to mediate H3K4 trimethylation (H3K4me3), a histone

modification essential for transcription (Wang et al., 2017, Shpargel et al., 2017), we integrated these datasets

with our *Mll4*-cKO DEGs (Fig. 4). Approximately 90% (105/117) of the DEGs overlapped with H3K4me3 target

genes (Fig. 7A) and were enriched for skeletal system-related biological processes (Fig. 7B). Among these, 66

genes were co-bound by UTX, implicating them as direct transcriptional targets of MLL4 (Fig. 7A). Intriguingly, these potential MLL4 direct target genes included *Comp* and *Snorc*, as well as other chondrogenesis-related (*Trpv4*, *Scx*, *Matn3*, and *Prkg2*) and osteogenesis-related genes (*Wnt10b* and *Hhip*) (Suppl. Figs. 1 and 2). Although *Sox9* did not appear in our DEG list, immunofluorescence results clearly indicated reduced SOX9 expression in the *Mll4*-cKO samples (Fig. 5). Incorporating *Sox9* as a "potential" DEG in our analysis revealed that it is targeted by both UTX and H3K4me3 (Fig. 7A; Suppl. Fig. 1), strongly suggesting that *Sox9* is a key MLL4 target. Together, these results indicate that MLL4 activates a subset of chondrogenic genes, potentially through the regulation of *Sox9*, via H3K4me3 deposition in collaboration with UTX. This proposed molecular mechanism is a schematic diagram (Fig. 7C).

Another factor contributing to gene expression changes in *Mll4*-deficient midpalaltal sutures is increased cell death. While *Mll4*-cKO mice exhibited a statistically significant increase in TUNEL-positive cells, the percentages of TUNEL-positive nuclei in both control and *Mll4*-cKO mice were relatively low (Fig. 3), indicating that apoptosis is rare in this context. Thus, the widened midpalatal suture phenotype in *Mll4*-cKO mice is likely attributed to deficient development due to impaired chondrogenic gene expression (Fig. 4). Analysis of DEGs for overlap with the Hallmark Apoptosis gene set (Liberzon et al., 2015) identified two downregulated genes, *Igfbp6* and *Lgals3*, but no upregulated genes. SOX9, which is known to regulate apoptosis in neural crest cells (Cheung et al., 2005), may contribute to the observed increase in TUNEL-positive cells (Fig. 3E–G). However, given the low percentage of apoptotic cells, increased apoptotic cells likely reflects an indirect consequence of developmental retardation in the midpalatal suture, rather than being the primary cause of the phenotype.

The chondrogenic retardation observed in our model aligns with previously reported defects in the cartilaginous primordia of the cranial base in neural crest-specific *Mll4* deletion mice and in the growth plates of long bones in heterozygous *Mll4* loss-of-function mice (Fahrner et al., 2019, Shpargel et al., 2020). Also, common downregulated genes were identified when comparing our DEG list with an RNA-seq dataset from earlier palatal shelves (at E14.25) in the neural crest-specific *Mll4* deletion mice (Shpargel et al., 2020), which include *Acan*, *Col2a1*, *Col9a1*, *Col9a2*, *Col9a3*, and *Mia*. These findings suggest a potential consensus regulatory mechanism downstream of MLL4 in different states of cell differentiation at different stages of

palatogenesis. Among these genes, *Acan* and *Col2a1* have been identified as causal genes for cleft palate (Xu

et al., 2021), highlighting a shared genetic basis for two distinct phenotypes—narrow palate and cleft palate—

manifesting under different molecular contexts at their onset. This is consistent with previous studies showing

that high-arched, narrow palate and cleft palate may share common genetic causes (Conley et al., 2016,

Tabler et al., 2013). However, it remains unclear why bilateral palatal shelves fail to fuse only in the neural

crest-specific *Mll4* deletion mice (resulting in cleft palate) but not in our palate mesenchyme-specific *Mll4*-cKO

mice. This may reflect an earlier requirement for MLL4 during the migration and differentiation of neural crest

cells prior to their incorporation into the palate mesenchyme, potentially explaining the cleft palate phenotype in

neural crest-specific *Mll4* mutants.

For future studies, it will be important to investigate whether the H3K4 methyltransferase activity of

MLL4 is specifically required for its role in postnatal palatogenesis. Recent findings (Xie et al., 2023) suggest

that MLL4 has both H3K4 methyltransferase-dependent and -independent roles in development and

differentiation. Notably, while H3K4 methylation appears largely dispensable for enhancer activation during

embryonic stem cell differentiation, it is crucial for lineage-selective processes in early embryonic development.

Exploring these distinct mechanisms in the context of postnatal palatogenesis may provide valuable insights

into the regulatory functions of MLL4 and its role in tissue-specific development.

MLL4 and the midpalatal suture: a central growth site and mechanosensitive regulator of craniofacial

development

The midpalatal suture has long been regarded as a critical skeletal structure that is anatomically situated at the

very center of the craniofacial complex, playing a significant role in the development and growth of the palate

and the maxillary jaw (Latham, 1971, Li et al., 2015, Rice, 2008). Therefore, it is intriguing to discover that the

two major phenotypes observed in the *Mll4*-cKO mice were a growth deficit in the palate and developmental

retardation of the midpalatal suture. While the causal relationship between midpalatal suture developmental

retardation and lateral growth deficit remains to be tested in future studies, our findings invite the possibility

that MLL4 regulates growth of the palate and the midface by modulating midpalatal suture. This highlights a

functional link between midpalatal suture development and palate growth and supports the potential central role of the midpalatal suture as a growth site driving craniofacial development.

In addition to serving as a growth site, the midpalatal suture functions as a mechanosensitive structure that responds to environmental forces, such as suckling, mastication, and skeletal growth (Roth et al., 2022, Li et al., 2016). For example, in orthodontic dentistry, rapid palatal expansion procedures apply mechanical force to the maxillary jaw, widening the midpalatal suture and inducing tensile forces that promote new bone formation, thereby increasing maxillary width. These responses have been partially attributed to *Piezo2*-positive chondrogenic mesenchyme cells in the midpalatal suture (Gao et al., 2022). Therefore, it would be interesting to test whether *Mll4* deficiency impacts the mechanosensory properties of the midpalatal suture, possibly through *Piezo2*-positive chondrogenic mesenchyme cells. This can be explored by simulating mechanical stress, such as performing a palatal expansion procedure in *Mll4*-cKO mice (Hou et al., 2007, Katebi et al., 2012). Such studies would provide valuable insights into the role of MLL4 in craniofacial development and the biological responses of the midpalatal suture to mechanical cues.

Data Availability Statement

Raw expression data results from the RNA-seq are included in the Supplementary Material, and the original RNA-seq data files were deposited in the NCBI GEO database under the accession number PRJNA1148613.

Ethics Statement

The animal study was reviewed and approved by Institutional Animal Care and Use Committee (IACUC) at the University at Buffalo.

Author Contributions

J-ML acquired funding, conceptualized, designed, administered, and supervised the research, performed experiments, acquired data, conducted data analysis and visualization, wrote and edited the manuscript. HJ performed experiments, acquired data, conducted data analysis and visualization, and edited the manuscript. BdPMP performed experiments, acquired data, conducted data analysis and visualization, and edited the manuscript. YP conducted data analysis and visualization and edited the manuscript. QT conducted re-analysis and visualization and edited the manuscript. SJ provided resources. S-KL acquired funding for the project, provided resources, and edited the manuscript. JWL acquired funding for the project, provided resources, and edited the manuscript. H-JEK acquired funding, conceptualized, designed, administered, and supervised the research, performed experiments, acquired data, conducted data analysis and visualization, wrote and edited the manuscript. All authors contributed to and approved the publication of the manuscript.

Funding

This study was supported by the National Institutes of Health's (NIH's) National Center for Advancing Translational Sciences (KL2TR001413 and UL1TR001412 to H-JEK), National Institute of Dental and Craniofacial Research (T32DE023526 to J-ML; R03DE030985 to H-JEK), and National Institute of Neurological Disorders and Strokes (R21NS123775 to YP; R01NS118748 to JWL and S-KL; R01NS100471 and R01NS111760 to S-KL).

Conflict of Interest

All authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

We thank Andrew D. McCall at the Optical Imaging and Analysis Facility (OIAF), School of Dental Medicine, The State University of New York at Buffalo, for assistance in acquiring micro-CT data and gross, histology, and stereo immunofluorescence images, which were all conducted at the OIAF. We thank Yang Chai at the University of Southern California for providing the *Osr2-Cre* mouse strain. We thank Eun-Myung Mary Song and Monique M. Kapur-Mauleon for assistance with experiments. We would also like to thank the members of the Kwon lab for their helpful comments and feedback.

References

BELL, D. M., LEUNG, K. K., WHEATLEY, S. C., NG, L. J., ZHOU, S., LING, K. W., SHAM, M. H., KOOPMAN, P., TAM, P. P. & CHEAH, K. S. 1997. SOX9 directly regulates the type-II collagen gene. *Nat Genet,* 16**,** 174-8.

BERNARD, P., TANG, P., LIU, S., DEWING, P., HARLEY, V. R. & VILAIN, E. 2003. Dimerization of SOX9 is required for chondrogenesis, but not for sex determination. *Human Molecular Genetics,* 12**,** 1755-1765.

BLIGHE, K., S, R. & M, L. 2024. *EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling* [Online]. Available: https://github.com/kevinblighe/EnhancedVolcano [Accessed].

BÖGERSHAUSEN, N., TSAI, I. C., POHL, E., KIPER, P., BELEGGIA, F., PERCIN, E. F., KEUPP, K., MATCHAN, A., MILZ, E., ALANAY, Y., KAYSERILI, H., LIU, Y., BANKA, S., KRANZ, A., ZENKER, M., WIECZOREK, D., ELCIOGLU, N., PRONTERA, P., LYONNET, S., MEITINGER, T., STEWART, A. F., DONNAI, D., STROM, T. M., BODUROGLU, K., YIGIT, G., LI, Y., KATSANIS, N. & WOLLNIK, B. 2015. RAP1-mediated MEK/ERK pathway defects in Kabuki syndrome. *J Clin Invest,* 125**,** 3585-99.

BONIEL, S., SZYMAŃSKA, K., ŚMIGIEL, R. & SZCZAŁUBA, K. 2021. Kabuki Syndrome-Clinical Review with Molecular Aspects. *Genes (Basel),* 12.

BRATTSTRÖM, V., MØLSTED, K., PRAHL-ANDERSEN, B., SEMB, G. & SHAW, W. C. 2005. The Eurocleft study: intercenter study of treatment outcome in patients with complete cleft lip and palate. Part 2: craniofacial form and nasolabial appearance. *Cleft Palate Craniofac J,* 42**,** 69-77.

BUSH, J. O. & JIANG, R. 2012. Palatogenesis: morphogenetic and molecular mechanisms of secondary palate development. *Development,* 139**,** 231-43.

CHEN, J. Q., LAN, Y., BAEK, J. A., GAO, Y. & JIANG, R. L. 2009. Wnt/beta-catenin signaling plays an essential role in activation of odontogenic mesenchyme during early tooth development. *Developmental Biology,* 334**,** 174-185.

CHEUNG, M., CHABOISSIER, M. C., MYNETT, A., HIRST, E., SCHEDL, A. & BRISCOE, J. 2005. The transcriptional control of trunk neural crest induction, survival, and delamination. *Dev Cell,* 8**,** 179-92.

CONLEY, Z. R., HAGUE, M., KUROSAKA, H., DIXON, J., DIXON, M. J. & TRAINOR, P. A. 2016. A quantitative method for defining high-arched palate using the mutant mouse as a model. *Developmental Biology,* 415**,** 296-305.

COX, T. C. 2004. Taking it to the max: the genetic and developmental mechanisms coordinating midfacial morphogenesis and dysmorphology. *Clin Genet,* 65**,** 163-76.

DIXON, M. J., MARAZITA, M. L., BEATY, T. H. & MURRAY, J. C. 2011. Cleft lip and palate: understanding genetic and environmental influences. *Nat Rev Genet,* 12**,** 167-78.

FAHRNER, J. A., LIN, W. Y., RIDDLE, R. C., BOUKAS, L., DELEON, V. B., CHOPRA, S., LAD, S. E., LUPERCHIO, T. R., HANSEN, K. D. & BJORNSSON, H. T. 2019. Precocious chondrocyte differentiation disrupts skeletal growth in Kabuki syndrome mice. *JCI Insight,* 4.

FERNANDEZ, R. & MOISY, C. 2021. Fijiyama: a registration tool for 3D multimodal time-lapse imaging. *Bioinformatics,* 37**,** 1482-1484.

FROIMCHUK, E., JANG, Y. & GE, K. 2017. Histone H3 lysine 4 methyltransferase KMT2D. *Gene,* 627**,** 337-342.

FUDALEJ, P., OBLOJ, B., MILLER-DRABIKOWSKA, D., SAMARCEW-KRAWCZAK, A. & DUDKIEWICZ, Z. 2008. Midfacial growth in a consecutive series of preadolescent children with complete unilateral cleft lip and palate following a one-stage simultaneous repair. *Cleft Palate Craniofac J,* 45**,** 667-73.

GAO, L., XU, T., ZHANG, L., LI, Y., YAN, T., YU, G. & CHEN, F. 2022. Midpalatal Suture: Single-Cell RNA-Seq Reveals Intramembrane Ossification and Piezo2 Chondrogenic Mesenchymal Cell Involvement. *Cells,* 11.

GARLAND, M. A., REYNOLDS, K. & ZHOU, C. J. 2020. Environmental mechanisms of orofacial clefts. *Birth Defects Res,* 112**,** 1660-1698.

GENZER, M. A. & BRIDGEWATER, L. C. 2007. A Col9a1 enhancer element activated by two interdependent SOX9 dimers. *Nucleic Acids Res,* 35**,** 1178-86.

GRIFFITHS, D. L., FURSTMAN, L. & BERNICK, S. 1967. Postnatal development of the mouse palate. *Am J Orthod,* 53**,** 757-68.

HAN, Y. & LEFEBVRE, V. 2008. L-Sox5 and Sox6 drive expression of the aggrecan gene in cartilage by securing binding of Sox9 to a far-upstream enhancer. *Mol Cell Biol,* 28**,** 4999-5013.

HOU, B., FUKAI, N. & OLSEN, B. R. 2007. Mechanical force-induced midpalatal suture remodeling in mice. *Bone,* 40**,** 1483-1493.

HUISMAN, C., KIM, Y. A., JEON, S., SHIN, B., CHOI, J., LIM, S. J., YOUN, S. M., PARK, Y., K, C. M., KIM, S., LEE, S. K., LEE, S. & LEE, J. W. 2021. The histone H3-lysine 4-methyltransferase Mll4 regulates the development of growth hormone-releasing hormone-producing neurons in the mouse hypothalamus. *Nat Commun,* 12**,** 256.

JAISWAL, P. K., ALJEBALI, L., GAUMOND, M. H., OH, C. D., YASUDA, H. & MOFFATT, P. 2020. Biochemical characteristics of the chondrocyte-enriched SNORC protein and its transcriptional regulation by SOX9. *Sci Rep,* 10**,** 7790.

KATEBI, N., KOLPAKOVA-HART, E., LIN, C. Y. & OLSEN, B. R. 2012. The mouse palate and its cellular responses to midpalatal suture expansion forces. *Orthod Craniofac Res,* 15**,** 148-58.

KENT, W. J., SUGNET, C. W., FUREY, T. S., ROSKIN, K. M., PRINGLE, T. H., ZAHLER, A. M. & HAUSSLER, D. 2002. The human genome browser at UCSC. *Genome Res,* 12**,** 996-1006.

LAN, Y., KINGSLEY, P. D., CHO, E. S. & JIANG, R. L. 2001. Osr2, a new mouse gene related to Drosophila odd-skipped, exhibits dynamic expression patterns during craniofacial, limb, and kidney development. *Mechanisms of Development,* 107**,** 175-179.

LAN, Y., OVITT, C. E., CHO, E. S., MALTBY, K. M., WANG, Q. R. & JIANG, R. L. 2004. Odd-skipped related 2 (Osr2) encodes a key intrinsic regulator of secondary palate growth and morphogenesis. *Development,* 131**,** 3207-3216.

LANGMEAD, B. & SALZBERG, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods,* 9**,** 357-9.

LATHAM, R. A. 1971. The development, structure and growth pattern of the human mid-palatal suture. *J Anat,* 108**,** 31-41.

LEE, J. E., WANG, C., XU, S., CHO, Y. W., WANG, L., FENG, X., BALDRIDGE, A., SARTORELLI, V., ZHUANG, L., PENG, W. & GE, K. 2013. H3K4 mono- and di-methyltransferase MLL4 is required for enhancer activation during cell differentiation. *Elife,* 2**,** e01503.

LEE, J. M., QIN, C., CHAI, O. H., LAN, Y., JIANG, R. & KWON, H. E. 2022. MSX1 Drives Tooth Morphogenesis Through Controlling Wnt Signaling Activity. *J Dent Res,* 101**,** 832-839.

LEFEBVRE, V., ANGELOZZI, M. & HASEEB, A. 2019. SOX9 in cartilage development and disease. *Current Opinion in Cell Biology,* 61**,** 39-47.

LEFEBVRE, V., HUANG, W., HARLEY, V. R., GOODFELLOW, P. N. & DE CROMBRUGGHE, B. 1997. SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(II) collagen gene. *Mol Cell Biol,* 17**,** 2336-46.

LI, C., LAN, Y. & JIANG, R. 2017. Molecular and Cellular Mechanisms of Palate Development. *J Dent Res,* 96**,** 1184-1191.

LI, H., HANDSAKER, B., WYSOKER, A., FENNELL, T., RUAN, J., HOMER, N., MARTH, G., ABECASIS, G. & DURBIN, R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics,* 25**,** 2078-9.

LI, J., JOHNSON, C. A., SMITH, A. A., HUNTER, D. J., SINGH, G., BRUNSKI, J. B. & HELMS, J. A. 2016. Linking suckling biomechanics to the development of the palate. *Sci Rep,* 6**,** 20419.

LI, J. T., JOHNSON, C. A., SMITH, A. A., SALMON, B., SHI, B., BRUNSKI, J. & HELMS, J. A. 2015. Disrupting the intrinsic growth potential of a suture contributes to midfacial hypoplasia. *Bone,* 81**,** 186-195.

LIBERZON, A., BIRGER, C., THORVALDSDÓTTIR, H., GHANDI, M., MESIROV, J. P. & TAMAYO, P. 2015. The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst,* 1**,** 417-425.

LIU, C. J., ZHANG, Y., XU, K., PARSONS, D., ALFONSO, D. & DI CESARE, P. E. 2007. Transcriptional activation of cartilage oligomeric matrix protein by Sox9, Sox5, and Sox6 transcription factors and CBP/p300 coactivators. *Front Biosci,* 12**,** 3899-910.

LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.

MATSUMOTO, N. & NIIKAWA, N. 2003. Kabuki make-up syndrome: A review. *American Journal of Medical Genetics Part C-Seminars in Medical Genetics,* 117c**,** 57-65.

MATSUNE, K., SHIMIZU, T., TOHMA, T., ASADA, Y., OHASHI, H. & MAEDA, T. 2001. Craniofacial and dental characteristics of Kabuki syndrome. *Am J Med Genet,* 98**,** 185-90.

MEAD, T. J., WANG, Q., BHATTARAM, P., DY, P., AFELIK, S., JENSEN, J. & LEFEBVRE, V. 2013. A far-upstream (-70 kb) enhancer mediates Sox9 auto-regulation in somatic tissues during development and adult regeneration. *Nucleic Acids Res,* 41**,** 4459-69.

MØLSTED, K., BRATTSTRÖM, V., PRAHL-ANDERSEN, B., SHAW, W. C. & SEMB, G. 2005. The Eurocleft study: intercenter study of treatment outcome in patients with complete cleft lip and palate. Part 3: dental arch relationships. *Cleft Palate Craniofac J,* 42**,** 78-82.

MUZUMDAR, M. D., TASIC, B., MIYAMICHI, K., LI, L. & LUO, L. Q. 2007. A global double-fluorescent cre reporter mouse. *Genesis,* 45**,** 593-605.

NG, L. J., WHEATLEY, S., MUSCAT, G. E. O., CONWAYCAMPBELL, J., BOWLES, J., WRIGHT, E., BELL, D. M., TAM, P. P. L., CHEAH, K. S. E. & KOOPMAN, P. 1997. SOX9 binds DNA, activates transcription, and coexpresses with type II collagen during chondrogenesis in the mouse. *Developmental Biology,* 183**,** 108-121.

PATRO, R., DUGGAL, G., LOVE, M. I., IRIZARRY, R. A. & KINGSFORD, C. 2017. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods,* 14**,** 417-419.

PORNTAVEETUS, T., ABID, M. F., THEERAPANON, T., SRICHOMTHONG, C., OHAZAMA, A., KAWASAKI, K., KAWASAKI, M., SUPHAPEETIPORN, K., SHARPE, P. T. & SHOTELERSUK, V. 2018. Expanding the Oro-Dental and Mutational Spectra of Kabuki Syndrome and Expression of KMT2D and KDM6A in Human Tooth Germs. *Int J Biol Sci,* 14**,** 381-389.

QUINLAN, A. R. & HALL, I. M. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics,* 26**,** 841-2.

REYNOLDS, K., ZHANG, S., SUN, B., GARLAND, M. A., JI, Y. & ZHOU, C. J. 2020. Genetics and signaling mechanisms of orofacial clefts. *Birth Defects Res,* 112**,** 1588-1634.

RICE, D. P. 2008. Craniofacial sutures. Development, disease and treatment. Preface. *Front Oral Biol,* 12**,** xi.

ROTH, D. M., SOUTER, K. & GRAF, D. 2022. Craniofacial sutures: Signaling centres integrating mechanosensation, cell signaling, and cell differentiation. *Eur J Cell Biol,* 101**,** 151258.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.

SCHMID, B., TRIPAL, P., FRAAß, T., KERSTEN, C., RUDER, B., GRÜNEBOOM, A., HUISKEN, J. & PALMISANO, R. 2019. 3Dscript: animating 3D/4D microscopy data using a natural-language-based syntax. *Nat Methods,* 16**,** 278-280.

SCHWENTY-LARA, J., NEHL, D. & BORCHERS, A. 2020. The histone methyltransferase KMT2D, mutated in Kabuki syndrome patients, is required for neural crest cell formation and migration. *Hum Mol Genet,* 29**,** 305-319.

SEKIYA, I., TSUJI, K., KOOPMAN, P., WATANABE, H., YAMADA, Y., SHINOMIYA, K., NIFUJI, A. & NODA, M. 2000. SOX9 enhances aggrecan gene promoter/enhancer activity and is up-regulated by retinoic acid in a cartilage-derived cell line, TC6. *Journal of Biological Chemistry,* 275**,** 10738-10744.

SHPARGEL, K. B., MANGINI, C. L., XIE, G., GE, K. & MAGNUSON, T. 2020. The KMT2D Kabuki syndrome histone methylase controls neural crest cell differentiation and facial morphology. *Development,* 147.

SHPARGEL, K. B., STARMER, J., WANG, C., GE, K. & MAGNUSON, T. 2017. UTX-guided neural crest function underlies craniofacial features of Kabuki syndrome. *Proc Natl Acad Sci U S A,* 114**,** E9046-e9055.

SHULL, L. C. & ARTINGER, K. B. 2024. Epigenetic regulation of craniofacial development and disease. *Birth Defects Res,* 116**,** e2271.

SUZUKI, A., SANGANI, D. R., ANSARI, A. & IWATA, J. 2016. Molecular mechanisms of midfacial developmental defects. *Dev Dyn,* 245**,** 276-93.

TABLER, J. M., BARRELL, W. B., SZABO-ROGERS, H. L., HEALY, C., YEUNG, Y., PERDIGUERO, E. G., SCHULZ, C., YANNAKOUDAKIS, B. Z., MESBAHI, A., WLODARCZYK, B., GEISSMANN, F., FINNELL, R. H., WALLINGFORD, J. B. & LIU, K. J. 2013. Mutant Mice Reveal Shared Mechanisms between Ciliopathies and FGF-Related Syndromes. *Developmental Cell,* 25**,** 623-635.

VAN DE LEST, C. H., VERSTEEG, E. M., VEERKAMP, J. H. & VAN KUPPEVELT, T. H. 1995. Elimination of autofluorescence in immunofluorescence microscopy with digital image processing. *J Histochem Cytochem,* 43**,** 727-30.

VAN LAARHOVEN, P. M., NEITZEL, L. R., QUINTANA, A. M., GEIGER, E. A., ZACKAI, E. H., CLOUTHIER, D. E., ARTINGER, K. B., MING, J. E. & SHAIKH, T. H. 2015. Kabuki syndrome genes KMT2D and

KDM6A: functional analyses demonstrate critical roles in craniofacial, heart and brain development. *Hum Mol Genet,* 24**,** 4443-53.

WANG, S. P., TANG, Z., CHEN, C. W., SHIMADA, M., KOCHE, R. P., WANG, L. H., NAKADAI, T., CHRAMIEC, A., KRIVTSOV, A. V., ARMSTRONG, S. A. & ROEDER, R. G. 2017. A UTX-MLL4-p300 Transcriptional Regulatory Network Coordinately Shapes Active Enhancer Landscapes for Eliciting Transcription. *Mol Cell,* 67**,** 308-321.e6.

WOLFORD, L. M., CASSANO, D. S., COTTRELL, D. A., EL DEEB, M., KARRAS, S. C. & GONCALVES, J. R. 2008. Orthognathic surgery in the young cleft patient: preliminary study on subsequent facial growth. *J Oral Maxillofac Surg,* 66**,** 2524-36.

WU, T., HU, E., XU, S., CHEN, M., GUO, P., DAI, Z., FENG, T., ZHOU, L., TANG, W., ZHAN, L., FU, X., LIU, S., BO, X. & YU, G. 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb),* 2**,** 100141.

XIE, G., LEE, J. E., SENFT, A. D., PARK, Y. K., JANG, Y., CHAKRABORTY, S., THOMPSON, J. J., MCKERNAN, K., LIU, C., MACFARLAN, T. S., ROCHA, P. P., PENG, W. & GE, K. 2023. MLL3/MLL4 methyltransferase activities control early embryonic development and embryonic stem cell differentiation in a lineage-selective manner. *Nat Genet,* 55**,** 693-705.

XIE, W. F., ZHANG, X., SAKANO, S., LEFEBVRE, V. & SANDELL, L. J. 1999. Trans-activation of the mouse cartilage-derived retinoic acid-sensitive protein gene by Sox9. *Journal of Bone and Mineral Research,* 14**,** 757-763.

XU, H. D., YAN, F. F., HU, R. F., SUZUKI, A., IWAYA, C., JIA, P. L., IWATA, J. & ZHAO, Z. M. 2021. CleftGeneDB: a resource for annotating genes associated with cleft lip and cleft palate. *Science Bulletin,* 66**,** 2340-2342.

YANG, Y. H., GOMEZ, N., INFARINATO, N., ADAM, R. C., SRIBOUR, M., BAEK, I., LAURIN, M. & FUCHS, E. 2023. The pioneer factor SOX9 competes for epigenetic factors to switch stem cell fates. *Nature Cell Biology,* 25**,** 1185-+.

YE, B., RUAN, C., HU, J., YANG, Y., THOMAS, J. & ZHANG, G. 2012. A comparative study on the measurements of palatal shelf area and gradient for adult patients with unoperated cleft palate. *Cleft Palate Craniofac J,* 49**,** 561-5.

YU, G., WANG, L. G. & HE, Q. Y. 2015. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics,* 31**,** 2382-3.

ZHANG, P., JIMENEZ, S. A. & STOKES, D. G. 2003. Regulation of human COL9A1 gene expression. Activation of the proximal promoter region by SOX9. *J Biol Chem,* 278**,** 117-23.

ZHANG, Y., LIU, T., MEYER, C. A., EECKHOUTE, J., JOHNSON, D. S., BERNSTEIN, B. E., NUSBAUM, C., MYERS, R. M., BROWN, M., LI, W. & LIU, X. S. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol,* 9**,** R137.

ZHOU, G., LEFEBVRE, V., ZHANG, Z. P., EBERSPAECHER, H. & DE CROMBRUGGHE, B. 1998. Three high mobility group-like sequences within a 48-base pair enhancer of the gene are required for cartilage-specific expression. *Journal of Biological Chemistry,* 273**,** 14989-14997.

Figure Legends

Figure 1. Generation of a novel *Mll4*-deficient Kabuki syndrome mouse model for palate studies. A

conditional *Mll4*-deficiency mouse model (*Mll4^{fl/fl}^;Osr2-Cre*, or *Mll4*-cKO) was generated, using the Cre/loxP

system driven by *Osr2-Cre* for studying postnatal palate development. (A, B) Osr2-Cre activation in the

developing midface (A) and palate (B), demonstrated in the *Osr2-Cre;mTmG* mouse at E13.5. ps, palatal shelf

(in B). (C–J) Craniofacial phenotype analysis, including gross examination (C, G) and micro-CT analysis (D–F;

H–J): facial angle (D, H), facial length (E, I), and facial width (F, J) all show mild but significant decrease in the

Mll4-cKO (H–J) compared to control (D–F) in adult mice at 10 weeks. (K, L) Orofacial phenotypes in the control

(K, K′) and *Mll4*-cKO (L, L′) adult mice at 10 weeks: *Mll4*-cKO mice exhibit midline defects, such as widened

midpalatal suture (arrowheads) and discontinuous ruga (3, arrows), along with disrupted rugae patterning (3–

6). (M) Body weight comparison between the control and *Mll4*-cKO mice shows no significant difference in

either adult (*n*=9 and 5 for control and cKO, respectively) or newborn mice (*n*=10 and 6 for control and cKO,

respectively). (N) Measurements of facial angle, length, and width from panels D–F and H–J (*n*=6 for both

control and cKO). Scale bars: 1 mm (in panels A–L).

Figure 2. *Mll4* is essential for proper growth of the palate during postnatal palatogenesis. (A–D) Skeletal

phenotypes were examined, using micro-CT analysis, in control (A, C) and *Mll4*-cKO (B, D) adult mice at 10

weeks. Virtual 3D (A, B) and coronal/frontal sections (C, D) along the anteroposterior axis at four

representative levels—for the incisive foramen (if) (slice 1), the palatine process of maxilla (ppm) (slice 2),

transpalatal suture (tps) (slice 3), and the palatal process of the palatine (ppp) (slice 4)—were examined. (E)

Palate width was measured in each for the four sections that represent the incisive foramen (slice 1), the

palatine process of maxilla (slice 2), transpalatal suture (slice 3), and the palatal process of the palatine (slice

4) at 1, 3, 6, and 10 weeks. *n*=6 or more for both controls and cKOs at all stages. Scale bars: 1 mm (in panels

A–D).

Figure 3. *Mll4* is required for initial formation of the midpalatal suture prior to birth. (A–D, G) Skeletal

staining (A, B) and trichrome staining results (C, D), with quantitative analysis of the midpalatal suture width

(G). Bone is stained magenta (in A, B) or blue (in C, D). The overall skeletal patterning was similar between the

control (A, C) and *Mll4*-cKO palate (B, D) prior to birth (at E18.5). However, specifically in the newly

established midpalatal suture (arrowheads in A, B; brackets in C′, D′), *Mll4*-cKO mice (D, D′) showed wider

suture width, compared to the control mice (C, C′, G). (E, F, E′, F′, H) TUNEL staining at E18.5 revealed that

Mll4 deficiency resulted in subtle but significant increase in cell apoptosis in the midpalatal suture (brackets in

E, F, E′, F′; H). *n*=6 for both control and cKO in panels G and H.

Figure 4. MLL4 regulates expression of genes associated with skeletal development in the newborn

midpalatal suture. (A–D) Genome-wide mRNA expression analysis was performed using the control and *Mll4*-

cKO midpalatal suture tissues at birth (P0.5). (A) Volcano plot showing that total 117 genes were differentially

expressed (adjusted *p*-value ≤ 0.001, log2FC ≥ 0.4)—40 upregulated genes and 77 downregulated genes. (B–

D) Top 10-enriched biological processes (B) and heatmap (C), identified through Gene Ontology (GO) analysis

from a list of significantly downregulated genes in the *Mll4*-cKO group, compared to the control group (D).

Figure 5. *Mll4* is essential for timely differentiation of chondrogenic cells in the midpalatal suture at

birth. (A–D) Immunofluorescence staining for expression of osteochondroprogenitor markers COL2A1, SOX9,

and RUNX2, as well as MLL4, in the control (A, C) and *Mll4*-cKO midpalatal suture (B, D) at birth (P0.5):

Chondrogenic markers, COL2A1 and SOX9, are markedly reduced in the *Mll4*-cKO midpalatal suture (B1, D2),

compared to control (A1, C2). In contrast, osteogenic marker, RUNX2, showed similar expression in the control

and *Mll4*-cKO suture tissue (A2, B2). MLL4 was markedly decreased in the *Mll4*-cKO palate (D1), compared to

control (C1). Brackets (in A–D) indicate the midpalatal suture. (E) Quantification of RUNX2-positive and SOX9-

positive cell ratios, as well as DAPI-positive nuclei counts in the midpalatal suture (*n*=6 for both control and

cKO). Scale bars: 50 μm (in panels A–D).

Figure 6. *Mll4* is required for proper development of the midpalatal suture during postnatal

palatogenesis. (A–J, I′, J′) Trichrome staining of the midpalatal suture in the control and *Mll4*-cKO mice,

showing that *Mll4* deficiency results in significant retardation in the postnatal development of the midpalatal

suture cartilage. (A, B) After birth (P0.75), the bilateral cartilage layers become present in the control

midpalatal suture (A, arrowheads) but remain completely absent in the *Mll4*-cKO midpalatal suture (B). (C, D)

At 1 week, the bilateral midpalatal suture cartilage layers become closer to each other and show uniform

enlargement of the chondrocyte cells in the control mice (C, arrowheads), whereas the suture cartilage layers

just start to partially appear in the *Mll4*-cKO mice (D, arrowheads). (E–J) At 3 weeks, the bilateral suture

cartilage layers adhere to each other in the midline in both the control and *Mll4*-cKO midpalatal sutures, but the

Mll4-cKO suture (F) is wider than the control suture (E), which becomes more obvious at 6 and 10 weeks (G–

J). (I, J, I′, J′) At 10 weeks, the control suture cartilage remains present and narrow (I), and its chondrocytes

are well differentiated and organized (I′), whereas the *Mll4*-cKO midpalatal suture cartilage remains wide (J),

and its chondrocytes are under-differentiated and disorganized (J′). Brackets (in A–J) indicate the midpalatal

suture. Scale bars: 50 μm (in panels A–D).

Figure 7. Proposed mechanism by which MLL4 regulates chondrogenic gene expression during

midpalatal suture development in postnatal palatogenesis. (A) Venn diagram illustrating the integration of

Mll4-cKO DEGs (including *Sox9*) and genes identified from re-analysis of UTX and H3K4me3 ChIP-seq

datasets (Shpargel et al., 2017). (B) GO enrichment analysis of *Mll4*-cKO DEGs associated with H3K4me3,

highlighting their involvement in various skeletal system-related biological processes. (C) Schematic diagram

illustrating the deduced molecular model involving MLL4, UTX, H3K4me3 and potential direct and indirect

target genes. MLL4 activates chondrogenic gene expression, potentially through targeting *Sox9*, in association

with UTX and by inducing H3K4me3. Created in https://BioRender.com.

Figure 1. Generation of a novel *Mll4*-deficient Kabuki syndrome mouse model for palate studies. A conditional *Mll4*-deficiency mouse model (*Mll4$^{fl/fl}$;Osr2-Cre*, or *Mll4*-cKO) was generated, using the Cre/loxP system driven by *Osr2-Cre* for studying postnatal palate development. (A, B) Osr2-Cre activation in the developing midface (A) and palate (B), demonstrated in the *Osr2-Cre;mTmG* mouse at E13.5. ps, palatal shelf (in B). (C–J) Craniofacial phenotype analysis, including gross examination (C, G) and micro-CT analysis (D–F; H–J): facial angle (D, H), facial length (E, I), and facial width (F, J) all show mild but significant decrease in the *Mll4*-cKO (H–J) compared to control (D–F) in adult mice at 10 weeks. (K, L) Orofacial phenotypes in the control (K, K′) and *Mll4*-cKO (L, L′) adult mice at 10 weeks: *Mll4*-cKO mice exhibit midline defects, such as widened midpalatal suture (arrowheads) and discontinuous ruga (3, arrows), along with disrupted rugae patterning (3–6). (M) Body weight comparison between the control and *Mll4*-cKO mice shows no significant difference in either adult (*n*=9 and 5 for control and cKO, respectively) or newborn mice (*n*=10 and 6 for control and cKO, respectively). (N) Measurements of facial angle, length, and width from panels D–F and H–J (*n*=6 for both control and cKO). Scale bars: 1 mm (in panels A–L).

1
2
Figure 2. *Mll4* is essential for proper growth of the palate during postnatal palatogenesis. (A–D) Skeletal
phenotypes were examined, using micro-CT analysis, in control (A, C) and *Mll4*-cKO (B, D) adult mice at 10
weeks. Virtual 3D (A, B) and coronal/frontal sections (C, D) along the anteroposterior axis at four
representative levels—for the incisive foramen (if) (slice 1), the palatine process of maxilla (ppm) (slice 2),
transpalatal suture (tps) (slice 3), and the palatal process of the palatine (ppp) (slice 4)—were examined. (E)
Palate width was measured in each for the four sections: at the incisive foramen (E1), first molar (M1) (slice 2),
second molar (M2) (slice 3), and third molar (M3) (slice 4) at 1, 3, 6, and 10 weeks. $n \geq 6$ for both controls and
cKOs at all stages. Scale bars: 1 mm (in panels A–D).

Figure 3. *Mll4* is required for initial formation of the midpalatal suture prior to birth. (A–D, G) Skeletal
staining (A, B) and trichrome staining results (C, D), with quantitative analysis of the midpalatal suture width
(G). Bone is stained magenta (in A, B) or blue (in C, D). The overall skeletal patterning was similar between the
control (A, C) and *Mll4*-cKO palate (B, D) prior to birth (at E18.5). However, specifically in the newly
established midpalatal suture (arrowheads in A, B; brackets in C′, D′), *Mll4*-cKO mice (D, D′) showed wider
suture width, compared to the control mice (C, C′, G). (E, F, E′, F′, H) TUNEL staining at E18.5 revealed that
Mll4 deficiency resulted in subtle but significant increase in cell apoptosis in the midpalatal suture (brackets in
E, F, E′, F′; H). *n*=6 for both control and cKO in panels G and H.

A Differential Gene Expression
Ctrl vs. *Mll4*-cKO

B Over-Representation Analysis, Biological Processes

C

D

Description	Gene ID	Count
Connective tissue development	Ihh, Trpv4, Hsd17b1, Matn3, Cnmd, Col2a1, Col9a1, Snorc, Prkg2, Acan, Comp, Scx, Mustn1	13
Cartilage development	Ihh, Trpv4, Matn3, Cnmd, Col2a1, Col9a1, Snorc, Prkg2, Acan, Comp, Scx, Mustn1	12
Extracellular matrix organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
External encapsulating structure organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
Extracellular structure organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
Skeletal system morphogenesis	Ihh, Trpv4, Col2a1, Wnt10b, Col9a1, Acan, Comp, Scx, Hhip	9
Chondrocyte differentiation	Ihh, Col2a1, Prkg2, Acan, Comp, Scx, Mustn1	7
Endochondral bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp, Scx	6
Bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp, Scx	6
Cartilage development involved in endochondral bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp	5

1
2
Figure 4. MLL4 regulates expression of genes associated with skeletal development in the newborn
midpalatal suture. (A–D) Genome-wide mRNA expression analysis was performed using the control and *Mll4*-
cKO midpalatal suture tissues at birth (P0.5). (A) Volcano plot, showing that total 117 genes were differentially
expressed (adjusted *p*-value ≤ 0.001, log2FC ≥ 0.4)—40 upregulated genes and 77 downregulated genes. (B–
D) Top 10-enriched biological processes (B) and heatmap (C), identified through Gene Ontology (GO) analysis
from a list of significantly downregulated genes in the *Mll4*-cKO group, compared to the control group (D).

Figure 5. *Mll4* is essential for timely differentiation of chondrogenic cells in the midpalatal suture at birth. (A–D) Immunofluorescence staining for expression of osteochondroprogenitor markers COL2A1, SOX9, and RUNX2, as well as MLL4, in the control (A, C) and *Mll4*-cKO midpalatal suture (B, D) at birth (P0.5): Chondrogenic markers, COL2A1 and SOX9, are markedly reduced in the *Mll4*-cKO midpalatal suture (B1, D2), compared to control (A1, C2). In contrast, osteogenic marker, RUNX2, showed similar expression in the control and *Mll4*-cKO suture tissue (A2, B2). MLL4 was markedly decreased in the *Mll4*-cKO palate (D1), compared to control (C1). Brackets (in A–D) indicate the midpalatal suture. (E) Quantification of RUNX2-positive and SOX9-positive cell ratios, as well as DAPI-positive nuclei counts in the midpalatal suture (*n*=6 for both control and cKO). Scale bars: 50 μm (in panels A–D).

Figure 6. *Mll4* is required for proper development of the midpalatal suture during postnatal palatogenesis. (A–J, I′, J′) Trichrome staining of the midpalatal suture in the control and *Mll4*-cKO mice, showing that *Mll4* deficiency results in significant retardation in the postnatal development of the midpalatal suture cartilage. (A, B) After birth (P0.75), the bilateral cartilage layers become present in the control midpalatal suture (A, arrowheads) but remain completely absent in the *Mll4*-cKO midpalatal suture (B). (C, D) At 1 week, the bilateral midpalatal suture cartilage layers become closer to each other and show uniform enlargement of the chondrocyte cells in the control mice (C, arrowheads), whereas the suture cartilage layers just start to partially appear in the *Mll4*-cKO mice (D, arrowheads). (E–J) At 3 weeks, the bilateral suture cartilage layers adhere to each other in the midline in both the control and *Mll4*-cKO midpalatal sutures, but the *Mll4*-cKO suture (F) is wider than the control suture (E), which becomes more obvious at 6 and 10 weeks (G–J). (I, J, I′, J′) At 10 weeks, the control suture cartilage remains present and narrow (I), and its chondrocytes are well differentiated and organized (I′), whereas the *Mll4*-cKO midpalatal suture cartilage remains wide (J), and its chondrocytes are under-differentiated and disorganized (J′). Brackets (in A–J) indicate the midpalatal suture. Scale bars: 50 μm (in panels A–D).

1
2
Figure 7. Proposed mechanism by which MLL4 regulates chondrogenic gene expression during
midpalatal suture development in postnatal palatogenesis. (A) Venn diagram illustrating the integration of
Mll4-cKO DEGs (including *Sox9*) and genes identified from re-analysis of UTX and H3K4me3 ChIP-seq
datasets (Shpargel et al., 2017). (B) GO enrichment analysis of *Mll4*-cKO DEGs associated with H3K4me3,
highlighting their involvement in various skeletal system-related biological processes. (C) Schematic diagram
illustrating the deduced molecular model involving MLL4, UTX, H3K4me3 and potential direct and indirect
target genes. MLL4 activates chondrogenic gene expression, potentially through targeting *Sox9*, in association
with UTX and by inducing H3K4me3. Created in https://BioRender.com.

UTX targets   H3K4me3 targets

687   10,854   4,835

66+1

0   39

Mll4-cKO DEG, plus Sox9

Sox9, Comp, Snorc, Trpv4, Scx, Matn3, Prkg2, Wnt10b,* and *Hhip
9 (out of the 67) genes: Enriched for skeletal system-related GO terms.
The 67 genes are direct targets of UTX & H3K4me3 (in O9-1 NCC cell line)
and also differentially expressed in the *Mll4*-cKO midpalatal suture.

Sox9

Comp

Snorc

Trpv4

Scx

Matn3

Wnt10b

Prkg2

Hhip

1
2
3 **Supplemental Figure 1. Integrative Genomics Viewer (IGV) browser tracks of UTX and H3K4me3 ChIP-**
4 **seq results.** Representative chondrogenesis-related (*Sox9*, *Comp*, *Snorc*, *Trpv4*, *Scx*, *Matn3*, and *Prkg2*) and
5 osteogenesis-related (*Wnt10b* and *Hhip*) genes identified by integrating *Mll4*-cKO midpalatal suture DEGs with
6 UTX and O9-1 cranial neural crest UTX and H3K4me3 ChIP-seq re-analyses (Shpargel et al., 2020), showing
7 MLL4-mediated H3K4 trimethylation in coordination with UTX.

Ihh, Cnmd, Col2a1, Col9a1, and *Acan*

5 (out of the 39) genes: Enriched for skeletal system-related GO terms. The 39 genes are direct targets of H3K4me3 only (in O-9 NCC cell line) and also differentially expressed in the *Mll4*-cKO midpalatal suture.

Supplemental Figure 2. Integrative Genomics Viewer (IGV) browser tracks of UTX and H3K4me3 ChIP-seq results. Representative chondrogenesis-related (*Ihh*, *Cnmd*, *Col2a1*, *Col9a1*, and *Acan*) genes identified by integrating *Mll4*-cKO midpalatal suture DEGs with UTX and O9-1 cranial neural crest UTX and H3K4me3 ChIP-seq re-analyses (Shpargel et al., 2020), showing MLL4-mediated H3K4 trimethylation independent of UTX.

Control   *Mll4*-cKO

A   B
if   if
ppm — slice 1   ppm
tps — slice 2   tps
— slice 3
— slice 4
ppp   ppp   1 mm

C   D
slice 1
slice 2
slice 3
slice 4   1 mm

E   ● Ctrl   ▲ cKO

E1 Incisive foramen level   E2 M1 level
    ns ** ** ****   ** **** **** ****

E3 M2 level   E4 M3 level
    *   **** *** ***   **** *** **** ****

weeks   weeks

	Control	*Mll4*-cKO

G **Midpalatal suture**

Whole mount

Width (μm)

**

Ctrl   cKO

Section

Width (μm)

*

Ctrl   cKO

H **Apoptosis**

% of TUNEL⁺ / total nuclei

**

Ctrl   cKO

A

Differential Gene Expression
Ctrl vs. *Mll4*-cKO

- Downregulated
- No change
- Upregulated

Log₁₀ *P* (y-axis)
Log2 fold change (x-axis)

Labeled genes: Col2a1, Cnmd, Matn3, Comp, Col9a2, Hapln1, Snorc, Mia, Mustn1, Acan, Ihh

Total = 54,347 genes

B Over-Representation Analysis, Biological Processes

Connective tissue development
Cartilage development
Extracellular matrix organization
External encapsulating structure organization
Extracellular structure organization
Skeletal system morphogenesis
Chondrocyte differentiation
Endochondral bone morphogenesis
Bone morphogenesis
Cartilage development involved in endochondral bone morphogenesis

p.adjust
0.00005
0.00010
0.00015
0.00020

Count

GeneRatio (x-axis)

C

Heatmap genes (top to bottom): Scube3, Hhip, Cnmd, Col2a1, Col9a1, Col9a3, Col9a2, S100b, Matn3, Comp, Snorc, Des, Id3, Id1, Tns2, Lgals3, Prkg2, Cryab, Jph2, Scx, P3h2, Acan, Trpv4, Wnt10b, Hsd17b1, Agt, Mustn1, Mia, Notum, Ihh

Columns: cKO 1, cKO 2, cKO 3, cKO 4, Ctrl 1, Ctrl 2, Ctrl 3, Ctrl 4

Scale: 1.5, 1, 0.5, 0, -0.5, -1, -1.5

D

Description	Gene ID	Count
Connective tissue development	Ihh, Trpv4, Hsd17b1, Matn3, Cnmd, Col2a1, Col9a1, Snorc, Prkg2, Acan, Comp, Scx, Mustn1	13
Cartilage development	Ihh, Trpv4, Matn3, Cnmd, Col2a1, Col9a1, Snorc, Prkg2, Acan, Comp, Scx, Mustn1	12
Extracellular matrix organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
External encapsulating structure organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
Extracellular structure organization	Ihh, Matn3, Col2a1, Col9a1, Col9a3, Col9a2, Acan, Comp, Agt, Scx, Lgals3, Mia	12
Skeletal system morphogenesis	Ihh, Trpv4, Col2a1, Wnt10b, Col9a1, Acan, Comp, Scx, Hhip	9
Chondrocyte differentiation	Ihh, Col2a1, Prkg2, Acan, Comp, Scx, Mustn1	7
Endochondral bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp, Scx	6
Bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp, Scx	6
Cartilage development involved in endochondral bone morphogenesis	Ihh, Trpv4, Col2a1, Col9a1, Comp	5

Control   *Mll4*-cKO

P0.75

A   B

50 μm

1 wk

C   D

50 μm

3 wks

E   F

50 μm

6 wks

G   H

50 μm

10 wks

I   J

50 μm

I'   J'

50 μm