Within-species variation eclipses between-species differences in *Pan* consolation

Jake S. Brooker[a], Christine E. Webb[b], Stephanie Kordon[a], Frans B. M. de Waal[c], Zanna Clay[a]

[a]Department of Psychology, Durham University, South Road, Durham, DH1 3LE, UK;
[b]Department of Human Evolutionary Biology, Harvard University, 11 Divinity Avenue,
Cambridge, MA 02138, USA; [c]Department of Psychology, Emory University, 36 Eagle Row,
Atlanta, GA 30322, USA

Corresponding authors: Zanna Clay, zanna.e.clay@durham.ac.uk; Jake Brooker,
jake.s.brooker@durham.ac.uk

Abstract

Empathy and its subcomponents are well documented throughout the animal kingdom, indicating
the deep evolutionary origins of this socioemotional capacity. A key behavioural marker of
empathy is consolation, or unsolicited bystander affiliation directed towards distressed others.
Consolation has been observed in our closest living relatives, bonobos (*Pan paniscus*) and
chimpanzees (*P. troglodytes*). However, systematic comparisons are absent, despite potential for
interspecific differences. Bonobos are often considered less aggressive, more emotionally
sensitive, and more socially tolerant than chimpanzees—key characteristics purported to drive
consolation. Furthermore, social and individual factors also appear to drive intraspecific
variation in empathy. To address within- and between-species variability in *Pan* consolation, we
systematically tested the consolatory tendencies of large sanctuary-living groups of both species.
Bonobos and chimpanzees exhibited similar consolation tendencies; however, within-species
analyses revealed further similarities and variation. Bonobo consolation was most often directed
towards and received by younger individuals, while chimpanzee consolation was most often
directed towards close social partners. In addition, males and females of both species showed
decreased consolation with age, with some evidence for chimpanzee males consoling more than
young females. Our findings support the notion that within-species variation in *Pan* socio-

27 emotional abilities is greater than between-species differences, highlighting the presence of
28 striking behavioural diversity across our two closest cousins.

29   Keywords: *bonobo; chimpanzee; consolation; empathy; Pan*

30   Word count: 9702

Introduction

Empathy, broadly defined as sharing and/or understanding others' emotional states, is a
cornerstone of the human experience [1,2]. From politics and marketing to day-to-day social
interactions, empathy allows us to effectively cooperate and communicate as well as strengthen
our social relationships [3]. Cross-cultural and personality studies have revealed remarkable
individual and cultural variation in empathic concern and perspective-taking [4–7]. Some
longitudinal studies indicate that individual differences in human empathic responding are
relatively stable [5–7]. As variation has been reported on individual and cultural levels, it has
been suggested that both personality and cultural norms may influence empathy. Likewise, it has
often been suggested that humans exhibit a uniquely advanced capacity for empathy, related to
our advanced socio-cognitive capacities including perspective taking, self-other differentiation,
emotion regulation and cognitive appraisal [8]. However, comparative research has revealed that
empathy likely manifests through bottom-up processes from foundational components with a
potential ancient ancestry in our evolutionary lineage [9]. These processes are thus proposed to
be regulated by sophisticated top-down information processing mechanisms.

In the animal kingdom, many behaviours possibly related to empathy have been reported,
mainly in mammals [10,11] though there is also some evidence in birds (e.g., consolation in
Corvus corax; [12]). Empathy-driven behaviours emerge in the form of sub-components that
represent differing expressions of emotional responsiveness and understanding [3,13]. Some of
these sub-components, such as emotional contagion, have been observed across many social
mammals, indicating a deep evolutionary history [10,13]. Furthermore, some behavioural
prosocial manifestations of empathy, such as consoling others in distress, have been observed in
rodents [14], elephants [15], and some primate and great ape species (for a review, see [13]).

Consolation, defined as unsolicited affiliation offered by an uninvolved bystander
towards a distressed conspecific [16], is often considered a more complex empathic behaviour.
Consolation, sometimes referred to as 'sympathetic concern' [17], is thought to require both a
cognitive appreciation (and even understanding) of another's state combined with a prosocial
orientation to improve it, such as by providing reassuring contact [18]. Among non-human
animals, our two closest relatives, the bonobos and chimpanzees have demonstrated the capacity
for consolation in a range of settings, including captivity, sanctuaries, and even the wild [19–23].
Among wild chimpanzees, both the western and eastern subspecies have been observed to
console others, including at Taï National Park, Ivory Coast [22] and the Mahale Mountains
National Park, Tanzania [24] respectively. Another study of the eastern subspecies in Budongo
Forest, Uganda, showed that consolation only occurred after 3.3% of observed conflicts, which
was not significantly more than matched controls [25]. In contrast, Kutsukake and Castles [24]
observed consolation after around 22% of conflicts in the Mahale M-group of eastern
chimpanzees. This difference between the Mahale and Budongo groups indicate possible within-
species behavioural flexibility and cross-community variation in consolatory tendencies. Whilst
there are no studies of consolation in wild bonobos, case studies of targeted prosocial behaviour,
such as conspecifics attempting to remove snares from group members and searching for lost
group members [26] suggest consolation is likely. Moreover, consolation has been observed in
sanctuary-settings [19], including by wild-born immature individuals, suggesting that
consolation is likely part of the wild bonobo repertoire.

Evidence that immature apes engage in consolation challenges assumptions that
consolation is dependent on sophisticated cognitive mechanisms. In both chimpanzees and
bonobos, consolation is offered most by younger individuals [19,27], suggesting that empathic
orientation towards other's states may be intrinsic and adaptive. In humans, concern for others
including consolation has been reported even within the first year of life, as early as nine months
[28]. Empathic behaviours, like consolation, may strengthen interpersonal bonds, especially for
dyads that demonstrate enduring cooperative relationships. For this reason, social closeness,
familiarity, or similarity has been shown to be a key predictor of empathic tendencies in humans
[29,30] and among nonhuman apes (bonobos: [19,31]; chimpanzees: [21,27,32]).

83   Bonobos and chimpanzees develop long term enduring bonds with other group members
84   and form strong support networks [33]; yet there are reported differences in their social
85   tendencies and socio-emotional orientations. For example, bonobos appear to have enhanced
86   attentiveness to conspecific social and emotional expressions than chimpanzees [34,35].
87   Apparent differences in sociability, socio-sexuality, and aggressive tendencies between the two
88   species may have biological and neurological foundations. Firstly, chimpanzees, but not
89   bonobos, have been shown to have deletion of the DupB region in the AVPR1A gene, which
90   includes a microsatellite called RS3 [36]. Variation in RS3 is linked with variation in social
91   bonding, and increased levels in bonobos may support their reported xenophilia. Neurological
92   research shows that bonobos have twice the density of serotonergic axons in the amygdala than
93   chimpanzees [37]. This region is associated with social cognition and emotional regulation
94   amongst others, and this variation between species may mediate the differences in their social
95   structures and behaviour. The ability to regulate one's own emotions—i.e., suppressing personal
96   arousal upon exposure to an arousing situation—can vary between individuals [38]. However,
97   emotional regulation appears to be crucial for empathy to emerge [39]. In addition, bonobos have
98   more grey matter in the right dorsal amygdala and right anterior insula than chimpanzees, as well
99   as a larger pathway linking the amygdala with the ventral anterior cingulate cortex [40]. These
100   regions are associated with perceiving distress in oneself and others, as well as control of
101   reactive and proactive aggressive impulses. These differences, as well as lower purported levels
102   of aggression in bonobos, and therefore decreased risk of injury [41,42], may facilitate enhanced
103   production of prosocial empathic behaviours, such as consolation and targeted helping,
104   compared to chimpanzees.

105   Social tolerance is also thought to influence empathic tendencies, according to the *Social*
Constraints Hypothesis [43]. For example, evidence of consolation in the Tonkean macaques
(*Macaca tonkeana*; [44]) and not in other more despotic macaque species (for a review, see
[13]), indicates that social tolerance may facilitate empathic responses that might otherwise be
inhibited due to intimidation, fear, or aggression. Chimpanzees are often considered to be more
intolerant and despotic than bonobos [33], following evidence that chimpanzees have formal
signals of subordinacy, through submissive gestures [45], whereas bonobos do not [46]. These
expressions of submission are often used to assess chimpanzee hierarchies, which commonly
follow a linear pattern [47–49]. Bonobos appear to have a more flexible, egalitarian dominance

style [50–52], which may be facilitated by their reduced aggressivity and increased sociability.

Furthermore, chimpanzees tend to be xenophobic to other communities [53]. In contrast, whilst

wild bonobo communities can be vocally hostile to other groups, numerous populations regularly

have peaceful intergroup encounters, even cases of food sharing, and exchanges of sex,

grooming, and play [54,55]. Some experimental paradigms have indicated that bonobos

outperform chimpanzees on cooperation [56] and theory-of-mind-related tasks [57]. Where

empathy expressions are related to understanding others, emotions, and social tolerance, one

might expect bonobos to be more consolatory than chimpanzees.

However, there is evidence of within-species variation. Several semi-wild chimpanzee

communities have been shown to vary in interindividual tolerance measured by cofeeding

proximity [58]. Similarly, a recent comparison of multiple captive and semi-wild populations

revealed that levels of social tolerance overlapped between bonobos and chimpanzees [59].

Furthermore, as previously mentioned, consolation appeared relatively absent in one population

of eastern chimpanzees at Budongo [25] and yet present in their Mahale counterparts [24]. These

communities have been demonstrated to vary considerably in social tolerance and hierarchical

steepness, whereby Budongo chimpanzees have steeper, and more despotic hierarchies compared

to Mahale [60]. Whilst these wild groups have not been systematically compared, these findings

indicate that consolation may show inter-group variation and emerge flexibly if conditions

permit. As population-level variation in group social tolerance can differ within species,

consolatory tendencies may vary across groups living in similar conditions.

Whilst some socio-cognitive abilities of bonobos and chimpanzees have been compared,

direct systematic comparisons of their consolation tendencies have yet to occur. As coalitionary

support and intermediate despotism are key factors of bonobo and chimpanzee societies,

comparing them may reveal whether reduced physical aggression and increased emotional

orientation in bonobos predict an increased likelihood to console. Consolation does appear to be

a feature of natural *Pan* social living; however, wild studies are scarce due to methodological

constraints. Therefore, comparing great apes in semi-wild sanctuary settings can provide a

balance of natural ecological surroundings with improved observational conditions. Based on

evidence that bonobos show enhanced social awareness and emotional sensitivity, we tested the

hypothesis that bonobos are more empathic than chimpanzees, predicting that consolation would

be more prevalent in bonobos than chimpanzees. Furthermore, following the hypothesis that

empathy is socially-biased [2], we predicted higher rates of consolation between dyads that are

socially bonded, through kinship and close affiliative tendencies.

Consolatory tendencies tend to decrease in each species with age [19,27], whilst previous

studies indicate a lack of general sex differences in consolation in bonobos and chimpanzees.

However, as males remain in their natal groups in each species [46,47], they might be expected

to invest in building long-term social bonds already at a young age. In bonobos, males tend not

to have strong social bonds beyond their mothers. They have been shown to form bonds with

other females that extend their alliance relationships and improve reproductive success, however

these are typically females with elevated rank positions, therefore reducing the likelihood that

they would be victims of conflict [61,62]. In chimpanzees, adult males are less likely to be

victims of conflicts than females and immature males, implying that their closest social partners,

other adult males, will have fewer opportunities to offer consolation. As such, and in line with

previous findings [19,27], we predicted that, regardless of sex, younger individuals of both

species will show the highest tendencies to console. In addition, we predicted an interaction

between bystander age and sex in both species, whereby younger individuals of both sexes

would show the highest consolatory tendencies whereas in adulthood, older males of each

species would show lower rates than females. By comparing between- and within-species

influences on an empathic behaviour like consolation in our closest living relatives, we aim to

improve our understanding of the origins of human empathy and emotional responsiveness.

Methods

Subjects and housing

We conducted observations of bonobos at Lola ya Bonobo Sanctuary (hereafter, "Lola ya

Bonobo") in the Democratic Republic of the Congo during July-September 2019. We conducted

observations of chimpanzees at Chimfunshi Wildlife Orphanage Trust (hereafter, "Chimfunshi")

in the Copperbelt Province of Zambia during March-August 2019. Lola ya Bonobo houses three

groups of bonobos in enclosures ranging from 15–20 hectares with rainforest, lake, swamp,

stream, and open grass area, whilst Chimfunshi is home to four groups of chimpanzees that are

accessible for observational research, living in outdoor miombo woodland enclosures ranging

from 20–80 hectares. At Lola ya Bonobo, bonobos sleep together in dormitories. At Chimfunshi,

chimpanzees nest independently unless kept inside for monitoring or medical intervention.

Both sanctuaries house wild-born individuals, orphaned and rescued from the pet and

bushmeat trades, as well as sanctuary-born individuals, all of whom are supported by caregiving

and veterinary staff. In both sanctuary environments, the apes could roam, forage, and nest

independently, whilst supported by an onsite caregiving team. The groups at Lola ya Bonobo

sleep at night in a managed indoor dormitory and otherwise roam in their forested enclosures

during the daytime. At both sites, groups are provisioned at least twice per day with a variety of

fruits and vegetables.

At Lola ya Bonobo, we observed all inhabitants of Groups 1 and 2 (hereafter: B1 and

B2), which housed $N = 22$ and $N = 18$ bonobos, respectively. At Chimfunshi we observed all

inhabitants of Group 2 (hereafter: C2), the largest group which comprised $N = 50$ chimpanzees at

the start of data collection. The age and sex composition of our sample is provided in Table S1

(ESM). We logged 800 hours of observations at Chimfunshi and 600 hours of observations at

Lola ya Bonobo.

Data collection

Victim focal follows

Consolation has typically been recorded using the post-conflict/matched-control method [63].

This involves conducting focal follows of a victim for a standardised period—usually 5–10-

minutes—after a conflict or spontaneous distress period, while recording the initiator and

behaviour of all affiliative interactions that occur involving the focal victim. Post-conflict (PC)

and post-distress (PD) periods are then compared with matched control (MC) recordings

observed at a similar time and circumstance a day later. The PC/MC method has already reliably

demonstrated consolation in multiple bonobo and chimpanzee communities compared with

matched controls [19–21,43], including in some of these sample populations [19]. Thus, we

decided to only collect post-conflict and post-distress events to ensure a large enough sample to

compare individual and social influences on consolation tendencies.

We used focal all-occurrence sampling [64] to collect $N = 150$ PCs and $N = 10$ PDs in

C2. In B1, we observed $N = 28$ PCs and $N = 36$ PDs and in B2, we observed $N = 36$ PCs and $N = 16$ PDs.

16 PDs. Our total sample of events was therefore $N = 276$ across both species and all three

groups. Previous studies of consolation show that most consolatory interactions occur during the

first minute after distress [19]. We therefore conducted victim focals for the first 5-minutes

following distress to optimise our limited observation time. We defined agonistic encounters by

the presence of at least one of the following behaviours: high-contact aggression [hit, slap, kick,

trample, bite], low-contact aggression [poke, push, push away, brush aside], chase, or threat

[threat bark, swagger, display, flail arm, stamp] [45]. PCs were only analysed if the victim

elicited a clear victim response, identified as the occurrence of bared-teeth scream, whimper,

tantrum, or flee from aggression [45] following an agonistic encounter. If these victim response

behaviours occurred spontaneously, they were coded as PD events. During the 5-min focal-

follows, we noted the presence of all bystanders within 1-metre and 5-metres of the focal subject

at the onset of the event. We recorded all PC/PD victim focals using Panasonic HC-V770

Camcorders with detachable Sennheiser MKE 400 directional shotgun microphones.

Dyadic social affinity and gregariousness

We also collected focal scan follows [64] of social tendencies to measure dyadic affiliation levels

and individual gregariousness. We collected social focals outside of feeding times for both

species. As we measured affiliation on a dyadic level, we accumulated information on

individuals who were not currently the focal subject if they associated with the focal. The data

set consisted of $N = 706$ focal follows for B1 (*range* = 28–41; *M* = 32.1; *SD* = 11.0), $N = 587$

focal follows for B2 (*range* = 16–44; *M* = 34.2; *SD* = 11.3), and $N = 684$ focal follows for the C2

(*range* = 9–18; *M* = 14.3; *SD* = 2.6). Each focal follow lasted 10-minutes and behaviours were

recorded at 1-minute scan points, upon which we recorded all social interactions the focal was

involved in including grooming, play, and sex, as well as all individuals within 1-metre. As the

social behaviours required proximity of 1-metre to the focal, we only coded either behaviour or

proximity once per dyad per scan during observations. However, when we compared dyadic

scores for behaviour and proximity when independent of one another, they were strongly

correlated (r(375) = .43, $P < .001$). We thus incorporated all social behaviour observations into

one score for proximity and used that to compute unique dyadic affiliation scores for every

possible dyad. We divided the number of scan points each dyad interacted for by the total

number of scan points both individuals of the dyad was observed for. We also computed

individual gregariousness scores by dividing the total number of scans each focal subject was

observed within proximity to at least one other individual by the total numbers of scans they

were observed for. At Chimfunshi Wildlife Orphanage, we recorded social focal follows using

the tablet app Zoo Monitor [65], which contributed towards long-term data collection of social

relationships at this site. At Lola ya Bonobo, we recorded social focals by hand due to software

incompatibility. In addition, we were interested in the effect of kinship, and defined each dyad as

'kin' or 'non-kin' depending on whether they shared a maternal genetic relationship or not,

respectively. This therefore included all mother-infant, sibling-sibling, and grandparent-

grandchild pairs as 'kin'. Some dyads at Chimfunshi represented a genetic uncle-nephew

dynamic and were also included as 'kin'. Inter-observer reliability revealed almost perfect

agreement between observers (partner identity: bonobo $\kappa > 0.80$; chimpanzee $\kappa > 0.96$; social

behaviour: bonobo $\kappa > 0.95$; chimpanzee $\kappa > 0.93$).

Hierarchies

We assigned categorical rank values (high, medium, low) for each ape. For both species, we used

the R package 'EloRating' [66] to create dominance scores based on dyadic agonistic

interactions involving high-contact aggression, low-contact aggression, and chases [66]. Each

agonism type was assigned a different optimised weighting, known as a K-value, based on

intensity and likelihood of winning probabilities. We only included individuals four-years and

older with at least six observed dyadic agonistic interactions in this ELO analysis. For the

chimpanzees, the ordered rank output was consistent with categorical rankings acquired from the

keepers and care staff. Hence, we evenly divided this output into 'high', 'medium', and 'low'

rank, and used the keeper allocations to assign categorical ranks to individuals for whom we

lacked sufficient ELO data for. For the bonobos, the ELO analysis revealed ordered results that

were broadly but not wholly consistent with keeper and researcher observations. Whilst human

judgements of social rank may be influenced by observer bias, these slight inconsistencies were

likely contributed to by a relatively lower quantity of agonistic observations over our relatively

short study period, therefore affording our ELO analysis lower power. However, whilst the rank

order was inconsistent, categorisation into 'high', 'medium', and 'low' was fairly consistent with

keeper and researcher observations. As we did not have substantial or prior quantitative data to support the ELO analysis, we decided to assign rank categories based on agreed deliberation between two long-term experienced observers of the bonobos at Lola ya Bonobo, supported by views of the sanctuary staff. For both species, we assigned mother ranks to infants (apes between 0 and 2 years of age). In both cases, observer rankings were based on observations of dynamics surrounding food and resource competition.

Ethics

This study was approved by the Senior Veterinary Advisory Team of Lola ya Bonobo Sanctuary, the Chimfunshi Research Advisory Board, and the Animal Welfare Ethics and Research Board of Durham University. Data collection comprised purely naturalistic observations and adhered to the legal requirements of the Democratic Republic of the Congo and Zambia, as well as the International Primatological Society's Principles for the Ethical Treatment of Nonhuman Primates.

Analysis

We conducted all-occurrence coding of affiliative and agonistic encounters during PC/PD periods in ELAN [67,68]. Consolation was defined as the onset of an affiliative interaction involving contact during the 5-minute follow that was spontaneously initiated by a bystander. Initiation was defined as the solicitation of an affiliative interaction either by gesturing or initiating physical contact with a partner. Thus, interactions initiated by the victim—e.g., the victim approached the bystander—were excluded as they do not represent spontaneous approaches by the bystander, thereby these interactions are not considered consolation [69]. Affiliative contact behaviours included body kiss, contact sit, embrace, finger/hand in mouth, embrace, genital inspection, genito-genital contact, grasp hand, groom, hunch-over, mount, mount walk, mouth kiss, pat, play, rump-rump touch, and touch [45,70]; see Table S2). Inter-coder reliability of consolation occurrence and consoler identity indicated almost perfect agreement (consolation occurrence: bonobo $\kappa = 0.86$, chimpanzee $\kappa = 0.85$; consoler identity: bonobo $\kappa = 1.00$, chimpanzee $\kappa = 0.97$).

Summary of statistical approaches

We used a Bayesian mixed models approach to assess whether consolation was more likely to occur in the bonobo or chimpanzee groups sampled, along with within-species variation in consolation tendencies according to bystander and victim characteristics: sex, age, rank, gregariousness, and group, as well as the bystander-victim social relationship. We collapsed our observations into two formats: event-level, where each observation row represents one PC or PD event; and dyad-level, where each observation row represents a unique bystander-victim combination within an event. This allowed us to comparably investigate general species tendencies at the event-level and then to assess the influence of individual (e.g., bystander age, bystander sex) and social factors (e.g., kinship, affiliation score) on consolation at the dyad-level. We z-transformed all covariate predictors and dummy coded and centred all factor predictors prior to their inclusion as random slopes.

We fitted all models in RStudio (version 1.3.1093; [71]) using the function brm of the package brms (version 2.21.0; [72]). Each model included four Markov chain Monte Carlo (MCMC) chains, with 10,000 iterations per chain, of which we specified 2,000 iterations as warm-up to ensure sampling calibration. This resulted in 32,000 posterior samples in total for each model. In all models, we used default priors (weakly informative priors with a student's t-distribution of 3 degrees of freedom and a scale parameter of 2.5) to ensure that our inferences were driven primarily by the data. Flat priors cover a wide range of parameter values without favouring any specific region. This allowed our already complex models incorporating many fixed and random effects to explore a broad parameter space, ensuring that no potential values are excluded a priori. Furthermore, flat priors emphasize the likelihood function, allowing the posterior distribution to closely follow the shape of the likelihood. Lastly, this approach also facilitates comparison with frequentist methods, enhancing the interpretability and generalisability of our findings.

For all models, we used multiple measures to summarise the posterior distributions for each variable and report evidence of an effect: 1) we characterised uncertainty by two-sided credible intervals (89% CrI) [73], denoting the range of probable values in which the true value could fall [74]; and 2) we computed the probability of direction (hereafter: *pd*) [75]. The 89% CrI reveals the range within which an effect falls with 89% probability [73], while the *pd*

indicates the probability that a parameter is strictly positive or negative [75]. For all models,

diagnostics revealed an accurate reflection of the original response values by the posterior

distributions, as R-hat statistics were < 1.01, the numbers of effective samples >1000, and

MCMC chains had no divergent transitions (see electronic supplementary material, hereafter:

ESM, Figure S1) [72]. Furthermore, diagnostics also revealed an accurate reflection of the

original response values by the posterior distributions (see ESM, Figure S2).

Event-level analyses: Testing species differences

We fitted two Bayesian Generalised Linear Mixed Models (GLMMs) to investigate whether

consolation was more likely to occur in bonobos or chimpanzees. The sample for this model

consisted of $N = 276$ events (PC $N = 214$; PD $N = 62$) of $N = 90$ apes (bonobo $N = 40$;

chimpanzee $N = 50$). The first model (hereafter: *Model 1.1*) fitted a Bernoulli distribution

(consolation occurrence: 1 = yes; 0 = no) and the second model (hereafter: *Model 1.2*) fitted a

Poisson distribution (count of consolations per event). The fixed effects structure for these

models included species and the number of bystanders. As population sizes varied across the

three groups, and consolation may be more likely if there are more individuals present, we

included the number of bystanders as a control effect to improve the accuracy of the estimated

fixed effect of species. The random effects structure consisted of random intercepts for aggressor

identity and victim identity, as well as random slopes for number of bystanders within both

aggressor identity and victim identity.

Dyad-level analyses: Testing individual and social factors

To test how individual and social factors influenced consolatory tendencies, we fitted two

GLMMs to our dyad-level data for each species separately [76]. Each observation row

represented a particular bystander-victim combination present during a given victim follow

event. Two bonobos and three chimpanzees were excluded due to long periods of absence from

their main groups (bonobos: 62–95% of observation time; chimpanzees: 54–65% of observation

time). The sample for the bonobo model (hereafter: *Model 2.1*) consisted of $N = 1656$

observation rows from $N = 38$ individuals observed across $N = 116$ events within $N = 2$ groups.

The sample for the chimpanzee model (hereafter: *Model 2.2*) consisted of $N = 5668$ observation

rows from $N = 47$ individuals observed across $N = 160$ within $N = 1$ group.

Our response variable constituted a binary outcome fitted with a Bernoulli distribution
according to whether the bystander solicited affiliative contact to the victim focal during the 5-
minute event (consolation occurrence: 1 = yes; 0 = no). Full models included individual factors
of the bystander (age, rank, sex), the victim (age, rank, and sex), and the social relationship of
the dyad (kinship) as fixed effects. Unfortunately, due to inconsistencies in the social affiliation
data collection protocol between Lola ya Bonobo and Chimfunshi, we were unable to include
quantitative measures for gregariousness and dyadic affiliation score in the bonobo model. The
chimpanzee model did however included fixed effects for bystander gregariousness and dyadic
affiliation score. For both species, we also included the interaction of bystander age and
bystander sex as a fixed effect and the bonobo model additionally included group as a fixed
effect. The random effects structure for each model consisted of random intercepts for the
individual identities of the aggressor, bystander, and victim as random effects as well as the
event ID. We also included theoretically identifiable random slopes of victim age and sex within
bystander identity and bystander age and sex within victim identity.

Results

Event-level results: Do bonobos console more than chimpanzees?

Across observations, consolation occurred at least once in 127 of the 276 events (46.0%). On
average, $M = 0.64$ ($SD = 0.81$) consolatory approaches occurred per event in bonobos, and $M =$
0.75 ($SD = 1.00$) consolatory approaches per event in chimpanzees. *Model 1.1* revealed no effect
of species on the probability of consolation occurring at least once in an event ($b = 0.08$, $SD =$
0.33, 89% credible interval (CrI) [-0.94, 1.11], *pd*: 0.55; see Figure 1). Additionally, *Model 1.2*
showed no effect of species on the count of consolers that responded per event ($b = 0.08$, $SD =$
0.33, 89% CrI [-0.44, 0.60], *pd* = 0.60). Importantly, in each model, the inclusion of the number
of bystanders also did not influence the outcome variables (*Model 1.1*: $b = 0.13$, $SD = 0.15$, 89%
CrI [-0.36, 0.62], *pd* = 0.66; *Model 1.2*: $b = 0.13$, $SD = 0.15$, 89% CrI [-0.12, 0.37], *pd* = 0.80),
indicating consolation likelihood does not increase if more possible consolers are present. Output
from the event-level models can be seen in Table S3 (ESM).

Figure 1. Barplot (left) and conditional probability plot (right) showing no credible difference between bonobos and chimpanzees in their relative tendency for a PC or PD event to feature 1+ consolation approach towards a distressed victim. Results obtained from *Model 1.1*. Whiskers on barplot show one standard error above and below mean. Probability plot shows conditional effect of species with 89% credible intervals.

Dyad-level results: Which within-species factors drive consolation tendencies?

A full table of the output for bonobos (*Model 2.1*) and chimpanzees (*Model 2.2*) can be seen in Table S4 (ESM).

Social effects

We found a very strong effect of kin dyads consoling more than non-kin dyads in both species (bonobos: b = -2.35, SD = 0.62, 89% CI [-3.37, -1.37], pd = 1.00; chimpanzees: b = -0.86, SD =

0.41, 89% CrI [-1.50, -0.19], *pd* = 0.98; see Figure 2). We also found a very strong positive

effect of affiliation level on consolation in chimpanzees (*b* = 0.20, *SD* = 0.09, 89% CrI [0.05,

0.35], *pd* = 0.98; see Figure 2). Furthermore, in chimpanzees, we found no clear effect of

bystander gregariousness (chimpanzees: *b* = -0.16, *SD* = 0.21, 89% CrI [-0.49, 0.17], *pd* = 0.79).

Finally, for bonobos we found no clear evidence for an effect of group (*b* = -0.49, *SD* = 0.52,

89% CrI [-1.34, 0.32], *pd* = 0.83).

Figure 2. Predicted probability plots showing credible effect of kinship in bonobos (1a) and

chimpanzees (1b) and a credible effect of dyadic affiliation level in chimpanzees (2). For all

plots, Y-axis = predicted probability of consolation. (1a-1b) X-axis = kinship. Square points and

error bars correspond to posterior means and the upper and lower 89% credible intervals,

respectively. (2) X-axis = z-transformed dyadic affiliation level. Shaded area corresponds to 89%

credibility intervals. Dyadic affiliation level not tested in bonobos.

Victim characteristics

In bonobos, we found a very strong negative effect of victim age where younger bonobos were

consoled more than older bonobos (b = -1.59, SD = 0.74, 89% CrI [-2.88, -0.55], pd = 1.00; see

Figure 2), but no credible effect in chimpanzees (b = 0.28, SD = 0.25, 89% CrI [-0.13, 0.68], pd

= 0.87; see Figure 2). We found no credible effects for victim sex in either species (bonobos: b =

1.08, SD = 0.97, 89% CrI [-0.36, 2.68], pd = 0.88; chimpanzees: b = -0.56, SD = 0.65, 89% CrI [-

1.62, 0.41], pd = 0.81). Victim rank had no credible effect on consolation tendency in either

bonobos or chimpanzees (see ESM, Table S4).

Figure 3. Predicted probability plot showing credible main effect of victim age in bonobos (left)

and no credible effect in chimpanzees (right). X-axis = z-transformed age. Y-axis = predicted

probability of consolation. Shaded area corresponds to 89% credibility intervals.

Bystander characteristics

We found no clear effect of bystander age and bystander sex interacting in bonobos (b = -0.81,

SD = 0.89, 89% CrI [-2.32, 0.50], pd = 0.67), however, there was evidence of a strong positive

effect of bystander age (b = -0.74, SD = , 89% CrI [-1.64, 0.12], pd = 1.00; see Figure 4)

indicating that younger bystanders may console more than older bystanders. However, for

chimpanzees we found moderate evidence for an interaction between bystander age and

bystander sex (b = -0.76, SD = 0.61, 89% CrI [-1.77, 0.19], pd = 0.90; see Figure 4). This

interaction indicates that indicated that younger individuals of both species consoled more than

older individuals, with young males consoling more than females. However, males appear to

decrease in their consolation tendency at a steeper rate than females. Conditional probability

plots of these interactions are shown in Figure S3 (ESM). Bystander rank had no credible effect

on consolation tendency in chimpanzees (see ESM, Table S4). However, high-ranking bonobos

consoled victims more than low-ranking bonobos (low vs. high: $b = -1.01$, $SD = 0.50$, 89% CrI [-

1.81, -0.24], $pd = 0.98$) but there was no credible difference between medium- and high-ranking

bystanders (medium vs. high: $b = -0.43$, $SD = 0.51$, 89% CrI [-1.44, 0.57], $pd = 0.81$).

Figure 4. Scatterplot showing credible main effect of bystander age in bonobos (top) and
moderate evidence for an interaction between bystander age and sex in chimpanzees (bottom).
X-axis = age in years. Y-axis = consolation/total opportunities to console. Females = blue
circles; males = red triangles. Point size reflects number of events the individual was present as a
bystander. Note, this plot provides a simple visualisation of the aggregated raw data using
frequentist 95% confidence limits for ease of comparison. See Figure S3 (ESM) for predicted
probability plots.

Discussion

Systematic cross-species comparisons are vital for elucidating the evolutionary origins for

complex socio-emotional behaviours, including empathy. Human research has revealed

substantial individual and cultural variation in emotional responding and this study indicates that

this may be the case for our closest living relatives. Due to purported increased emotional

sensitivity, reduced aggressivity, and higher levels of social tolerance [33,41], it has been

suggested that bonobos may be more empathic than chimpanzees. However, our results indicate

that the two species do not significantly differ in consolation tendency—considered a

behavioural expression of empathy—both in terms of the behaviour's overall occurrence and the

number of consolers that respond per event. Our results reflect recent findings that contrast with

these previously held assumed species dichotomisations, such as in aggression [77] and social

tolerance [59]. We did, however, find several within-species trends in potential drivers of

consolatory tendencies. Expressions of empathy and other emotion-related behaviour are known

to vary in wild chimpanzee communities of the same subspecies [24,25], which parallels findings

across diverse human cultures [4]. The presence of intraspecies group variation in consolation in

the sanctuary bonobos complements this literature and suggests that empathic tendencies may be

shaped by broader individual and socio-cultural circumstances.

Our findings complement previous consolation research in *Pan* by showing that

bystander age is an important predictor of consolation in bonobo and chimpanzee communities

[19,27]. Although underlying mechanisms cannot be detected in an observational study, it is

possible that young individuals may console more than older individuals as they are typically

afforded more social tolerance before they become fully embedded into their respective group's

dominance dynamics [78]. Furthermore, younger individuals may be less inhibitive with their

responses in these contexts. We did not measure individual's capacity for inhibition in this study.

However, an interesting follow-up would be to compare these two species on inhibition, such as

through delayed gratification tasks [79], and investigate if this measure predicts an individual's

relative tendency to console.

Consistent with the theory that social closeness predicts empathic responses [2,80],

kinship predicted consolation in both species. However, we were only able to test dyadic

affiliation levels in chimpanzees, finding that consolation was biased towards closer social partners generally. Previous investigations on these bonobo groups revealed strong influences of social relationship on consolation occurrences [19]. Temporal variation in consolatory efforts may reflect empathy as a phenomenon that emerges flexibly when social conditions permit [13].

It is possible that consolation may serve different functions in bonobo and chimpanzee societies. Chimpanzees may be more selective in their responses as wild and captive communities have been well-documented to exhibit political social lives, centred around strong adult male-male alliances and displays of dominance and subordinacy [33]. This may be further evidenced by a possible age-sex interaction in chimpanzees, whereby young males were the most consolatory and older individuals were the least consolatory. Enhanced responsiveness in young male chimpanzees may represent a drive to form and reinforce strong social bonds with their close peers and older individuals they will spend their lives with [81]. Furthermore, adult male chimpanzees were much less likely to be followed as distressed focals ($N = 17/160$) compared to adult females ($N = 103/160$). Therefore, opportunities for adult male chimpanzees to console their closest social partners were respectively low and may additionally explain the direction of the age-sex interaction whereby older males as unlikely to console as adult females. In the Kanyawara community at Kibale Chimpanzee Project in Uganda, young male chimpanzees are typically more likely to be victims of aggression, which appears to be due to their tendencies to elicit the most aggression themselves [82]. Regardless, social bonds formed during male chimpanzee development predict dominance trajectories [83]. Thus, this effect at a young age in chimpanzees may be caused by younger males watching their peers being aggressed often during a period when they are building foundational relationships for later life [84].

In contrast, in bonobos, males typically only form strong relationships with their mothers [85] and occasionally other females for mating and alliance formation [61]. Despite males being the philopatric sex for both species, only chimpanzee communities tend to feature strong male-male relationships [33]. Furthermore, the political dynamics in bonobo societies appear to be relatively more subtle, and thus difficult to study [54]. Instead, bonobos may be less selective and respond out of care and emotional sensitivity towards others [35,86]. Bonobos tended to console victims of both sexes and all ranks evenly. Furthermore, in line with recent findings that paedomorphic signalling increases likelihood of receiving consolation [87], younger bonobo

victims were generally more likely to be consoled. However, higher-ranked bonobo bystanders
did offer consolation more than those of low rank, indicating that lower social standing may
inhibit individuals to approach distressed peers. A deeper, longitudinal comparison with more
groups of each species is needed to reveal the extent to which consolation has a political
association across age, rank, and sex combinations in each species.

In this study, we imposed strict criteria of inclusion for consolation, where any behaviour
following a solicitation from the victim was not counted. For example, if a victim immediately
approached a bystander and solicited an interaction with them, any subsequent interactions
initiated by the bystander during the PC/PD period were not coded as consolatory approaches.
As consolation is linked with empathy, if bystanders are not the party to solicit contact then it is
not possible to infer that they may have been motivated by the victim's distress to respond
[43,88]. Therefore, there is also a possibility that victim-initiated interactions provided
reassurance and comfort to the victim, yet these encounters were not counted in our analyses. A
wider investigation into bystander-victim interactions following distress could reveal whether the
intraspecies trends found in this study are associated with empathic tendencies, or wider conflict
management responses in general [89].

Intraspecies variation is vital for comparative research across taxa [90]. Studies of
primates have revealed substantial intergroup behavioural variation, such as bartering in long-
tailed macaques (*Macaca fascicularis*; [91]), activity and ranging in guerezas (*Colobus guereza*;
[92]), and even the connectedness of social networks in vervets (*Chlorocebus pygerythrus*; [93]).
The *Pan* apes similarly can vary between groups regarding expressions of various social and
ecological behaviours. Examples include, but are by no means limited to, grooming traditions
[94] and communication in chimpanzees [95], and hunting behaviour and tool use in bonobos
[96]. In addition, bonobos and chimpanzees also show marked intraspecies variation in levels of
social tolerance [58,59,97], a factor considered to facilitate empathy [43,44].

Whilst there were no overall species differences between the bonobos and chimpanzees
in this sample, individual variation and social relationships appear to drive consolation. The
Social Constraints Hypothesis [43] posits that empathy is facilitated by social tolerance, and has
been suggested to explain why some primates express empathic behaviour and others do not
[43,44]. Bonobos have previously been purported to be more socially tolerant than chimpanzees

[41,56], however recent analyses indicate that greater variation lies at the group-level in the *Pan*

apes [59,77]. Consolation appears to be primarily driven by individual and social characteristics,

indicating that group compositional changes may lead to variation in empathic tendencies over

time, similar to recent studies on social tolerance [98,99]. For example, groups may score

differently on consolation when numbers of juveniles and kin-related pairs fluctuate. In our

sample, when controlling for all other factors, there was no credible difference between bonobo

groups in consolation tendency. To further elucidate any group-level variation in consolation

across the *Pan* apes [43,44], future research should assess tolerance levels and risk of aggression

of multiple groups and compare these group-specific factors with relative tendencies to console.

In sum, our findings support the notion that within-species community variation among

Pan behavioural tendencies may be more significant than between-species differences

[33,90,100]. Both bonobos and chimpanzees are highly flexible and adaptable species, and under

particular conditions and pressures may exhibit greater or reduced tendencies to express

empathy-related behaviours like consolation. In humans, we see individual- and group-level

variation in behaviours such as communication, prosociality, conformity, and empathy and other

socio-emotional responses that may be facilitated by certain social dynamics and cultural norms

[4,101–103]. The same may be true in our closest living relatives. Further research into species

comparisons should always integrate group-level differences (e.g., composition, collective

temperaments) that may promote or hinder empathic expressions such as consolation. Comparing

more groups of the same species and investigating within-group drivers of various empathy-

related behaviours may reveal a deep ancestral history of such cultural flexibility in emotional

responsiveness, and what features promote or hinder the expression of empathy.

Acknowledgements

We thank the keepers, veterinarians, and all other staff of Chimfunshi Wildlife Orphanage Trust

and Lola ya Bonobo, including Innocent Mulenga, Felix Chinyama, Thomson Mbilishi, Lameck

Musaka, Joseph Kasongo, Thalita Calvi, Fanny Minesi, Raphaël Belais, Stany Mokando, and

Jean-Claude Nzumbi. We thank Zoë Goldsborough and Heritier Izansone for their assistance

with data collection and Emma Doherty and Georgia Sandars for help with reliability coding. We

also thank Edwin van Leeuwen for his support and assistance with this project, particularly with

social data collection and analysis. Funding was provided by the Templeton World Charity

Foundation, and CW would like to thank the Carper Foundation for additional grant support.

Conflict of interest

565 We declare we have no competing interests.

Data accessibility

567 All data and R code supporting this article are included as electronic supplementary material.

References

1. Eisenberg N, Strayer J. 1990 *Empathy and Its Development*. CUP Archive.

2. Preston SD, de Waal FBM. 2002 Empathy: Its ultimate and proximate bases. *Behavioral and*
Brain Sciences **25**, 1–20. (doi:10.1017/S0140525X02000018)

3. de Waal FBM. 2007 The 'Russian doll' model of empathy and imitation. In *On being moved:*
From mirror neurons to empathy (ed S Bråten), pp. 49–69. Amsterdam, Netherlands: John
Benjamins Publishing Company. (doi:10.1075/aicr.68.06waa)

4. Chopik WJ, O'Brien E, Konrath SH. 2017 Differences in Empathic Concern and Perspective
Taking Across 63 Countries. *Journal of Cross-Cultural Psychology* **48**, 23–38.
(doi:10.1177/0022022116673910)

5. Eisenberg N, Carlo G, Murphy B, van Court P. 1995 Prosocial Development in Late
Adolescence: A Longitudinal Study. *Child Development* **66**, 1179–1197.
(doi:10.1111/j.1467-8624.1995.tb00930.x)

6. Eisenberg N, Guthrie IK, Murphy BC, Shepard SA, Cumberland A, Carlo G. 1999
Consistency and Development of Prosocial Dispositions: A Longitudinal Study. *Child*
Development **70**, 1360–1372. (doi:10.1111/1467-8624.00100)

7. Knafo A, Zahn-Waxler C, Van Hulle C, Robinson JL, Rhee SH. 2008 The developmental
origins of a disposition toward empathy: Genetic and environmental contributions. *Emotion*
8, 737–752. (doi:10.1037/a0014179)

8. Decety J, Norman GJ, Berntson GG, Cacioppo JT. 2012 A neurobehavioral evolutionary
perspective on the mechanisms underlying empathy. *Progress in Neurobiology* **98**, 38–48.
(doi:10.1016/j.pneurobio.2012.05.001)

9. Decety J, Lamm C. 2006 Human Empathy Through the Lens of Social Neuroscience.
TheScientificWorldJOURNAL **6**, 1146–1163. (doi:10.1100/tsw.2006.221)

10. Adriaense JEC, Koski SE, Huber L, Lamm C. 2020 Challenges in the comparative study of
empathy and related phenomena in animals. *Neuroscience & Biobehavioral Reviews* **112**,
62–82. (doi:10.1016/j.neubiorev.2020.01.021)

11. Pérez☐Manrique A, Gomila A. 2018 The comparative study of empathy: sympathetic
concern and empathic perspective-taking in non-human animals. *Biological Reviews* **93**,
248–269. (doi:10.1111/brv.12342)

12. Fraser ON, Bugnyar T. 2010 Do Ravens Show Consolation? Responses to Distressed Others.
PLoS One **5**, e10605. (doi:10.1371/journal.pone.0010605)

13. Brooker JS, Webb CE, Clay Z. 2022 Primate empathy: A flexible and multi-componential
phenomenon. *Primate Cognitive Studies*. , 505–532. (doi:10.31234/osf.io/bpw5x)

14. Burkett JP, Andari E, Johnson ZV, Curry DC, de Waal FBM, Young LJ. 2016 Oxytocin-
dependent consolation behavior in rodents. *Science* **351**, 375–378.
(doi:10.1126/science.aac4785)

15. Plotnik JM, de Waal FBM. 2014 Asian elephants (Elephas maximus) reassure others in
distress. *PeerJ* **2**, e278. (doi:10.7717/peerj.278)

16. de Waal FBM, van Roosmalen A. 1979 Reconciliation and consolation among chimpanzees.
Behavioral Ecology and Sociobiology **5**, 55–66. (doi:10.1007/BF00302695)

17. Batson CD, Fultz J, Schoenrade PA. 1987 Distress and Empathy: Two Qualitatively Distinct
Vicarious Emotions with Different Motivational Consequences. *Journal of Personality* **55**,
19–39. (doi:10.1111/j.1467-6494.1987.tb00426.x)

18. de Waal FBM. 2008 Putting the altruism back into altruism: the evolution of empathy.
Annual Review of Psychology **59**, 279–300. (doi:10.1146/annurev.psych.59.103006.093625)

19. Clay Z, de Waal FBM. 2013 Bonobos respond to distress in others: Consolation across the
age spectrum. *PLoS One* **8**, e55206. (doi:http://dx.doi.org/10.1371/journal.pone.0055206)

20. Palagi E, Paoli T, Tarli SB. 2004 Reconciliation and consolation in captive bonobos (Pan
paniscus). *American Journal of Primatology* **62**, 15–30. (doi:10.1002/ajp.20000)

21. Romero T, Castellanos MA, de Waal FBM. 2010 Consolation as possible expression of
sympathetic concern among chimpanzees. *Proceedings of the National Academy of Sciences*
107, 12110–12115. (doi:10.1073/pnas.1006991107)

22. Wittig R, Boesch C. 2003 The Choice of Post-conflict Interactions in Wild Chimpanzees
(Pan troglodytes). *Behaviour* **140**, 1527–1559. (doi:10.1163/156853903771980701)

23. Brooker JS, Webb CE, de Waal FBM, Clay Z. 2024 The expression of empathy in human's
closest relatives, bonobos and chimpanzees: current and future directions. *Biological*
Reviews **n/a**. (doi:10.1111/brv.13080)

24. Kutsukake N, Castles DL. 2004 Reconciliation and post-conflict third-party affiliation
among wild chimpanzees in the Mahale Mountains, Tanzania. *Primates* **45**, 157–165.
(doi:10.1007/s10329-004-0082-z)

25. Arnold K, Whiten A. 2001 Post-conflict behaviour of wild chimpanzees (Pan troglodytes
schweinfurthii) in the Budongo Forest, Uganda. *Behaviour* **138**, 649–690.

26. Tokuyama N, Emikey B, Bafike B, Isolumbo B, Iyokango B, Mulavwa MN, Furuichi T.
2012 Bonobos apparently search for a lost member injured by a snare. *Primates* **53**, 215–
219. (doi:10.1007/s10329-012-0298-2)

27. Webb CE, Romero T, Franks B, de Waal FBM. 2017 Long-term consistency in chimpanzee
consolation behaviour reflects empathetic personalities. *Nat Commun* **8**, 1–8.
(doi:http://dx.doi.org/10.1038/s41467-017-00360-7)

28. Davidov M, Paz Y, Roth-Hanania R, Uzefovsky F, Orlitsky T, Mankuta D, Zahn-Waxler C.
2021 Caring babies: Concern for others in distress during infancy. *Developmental Science*
24, e13016. (doi:10.1111/desc.13016)

29. Lindegaard MR, Liebst LS, Bernasco W, Heinskou MB, Philpot R, Levine M, Verbeek P.
2017 Consolation in the aftermath of robberies resembles post-aggression consolation in
chimpanzees. *PLoS One* **12**. (doi:10.1371/journal.pone.0177725)

30. Norscia I, Palagi E. 2011 Yawn Contagion and Empathy in Homo sapiens. *PLoS One* **6**,
e28472. (doi:10.1371/journal.pone.0028472)

31. Palagi E, Norscia I. 2013 Bonobos Protect and Console Friends and Kin. *PLoS One* **8**,
e79290. (doi:10.1371/journal.pone.0079290)

32. Fraser ON, Stahl D, Aureli F. 2008 Stress reduction through consolation in chimpanzees.
PNAS **105**, 8557–8562. (doi:10.1073/pnas.0804141105)

33. Gruber T, Clay Z. 2016 A Comparison Between Bonobos and Chimpanzees: A Review and
Update. *Evolutionary Anthropology: Issues, News, and Reviews* **25**, 239–252.
(doi:10.1002/evan.21501)

34. Kano F, Hirata S, Call J. 2015 Social Attention in the Two Species of Pan: Bonobos Make
More Eye Contact than Chimpanzees. *PLoS One* **10**, e0129684.
(doi:10.1371/journal.pone.0129684)

35. Kret ME, Jaasma L, Bionda T, Wijnen JG. 2016 Bonobos (Pan paniscus) show an attentional
bias toward conspecifics' emotions. *Proceedings of the National Academy of Sciences of the*
United States of America **113**, 3761–3766. (doi:10.1073/pnas.1522060113)

36. Staes N, Stevens JMG, Helsen P, Hillyer M, Korody M, Eens M. 2014 Oxytocin and
Vasopressin Receptor Gene Variation as a Proximate Base for Inter- and Intraspecific

Behavioral Differences in Bonobos and Chimpanzees. *PLOS ONE* **9**, e113364.

(doi:10.1371/journal.pone.0113364)

37. Stimpson CD, Barger N, Taglialatela JP, Gendron-Fitzpatrick A, Hof PR, Hopkins WD,

Sherwood CC. 2016 Differential serotonergic innervation of the amygdala in bonobos and

chimpanzees. *Social Cognitive and Affective Neuroscience* **11**, 413–422.

(doi:10.1093/scan/nsv128)

38. Eisenberg N, Fabes RA, Murphy B, Karbon M, Maszk P, Smith M, O'Boyle C, Suh K. 1994

The relations of emotionality and regulation to dispositional and situational empathy-related

responding. *Journal of Personality and Social Psychology* **66**, 776–797. (doi:10.1037/0022-

3514.66.4.776)

39. Lockwood PL, Seara-Cardoso A, Viding E. 2014 Emotion regulation moderates the

association between empathy and prosocial behavior. *PLoS One* **9**, e96555.

(doi:10.1371/journal.pone.0096555)

40. Rilling JK, Scholz J, Preuss TM, Glasser MF, Errangi BK, Behrens TE. 2012 Differences

between chimpanzees and bonobos in neural systems supporting social cognition. *Social*

Cognitive and Affective Neuroscience **7**, 369–379. (doi:10.1093/scan/nsr017)

41. Hare B, Wobber V, Wrangham R. 2012 The self-domestication hypothesis: evolution of

bonobo psychology is due to selection against aggression. *Animal Behaviour* **83**, 573–585.

(doi:10.1016/j.anbehav.2011.12.007)

42. Wilson ML *et al.* 2014 Lethal aggression in Pan is better explained by adaptive strategies

than human impacts. *Nature* **513**, 414–417. (doi:10.1038/nature13727)

43. de Waal FBM, Aureli F. 1996 Consolation, reconciliation, and a possible cognitive

difference between macaques and chimpanzees. In *Reaching Into Thought: The minds of the*

great apes (eds AE Russon, KA Bard, ST Parker), pp. 80–110. Cambridge, UK: Cambridge

University Press.

44. Palagi E, Dall'Olio S, Demuru E, Stanyon R. 2014 Exploring the evolutionary foundations of
empathy: consolation in monkeys. *Evolution and Human Behavior* **35**, 341–349.
(doi:10.1016/j.evolhumbehav.2014.04.002)

45. Nishida T, Zamma K, Matsusaka T, Inaba A, McGrew WC. 2010 *Chimpanzee Behavior in*
the Wild: An Audio-Visual Encyclopedia. Tokyo, Japan: Springer Science & Business Media.

46. Kano T. 1992 *The Last Ape: Pygmy Chimpanzee Behavior and Ecology*. Stanford, CA:
Stanford University Press.

47. Goodall J. 1986 *The Chimpanzees of Gombe: Patterns of Behavior*. Cambridge, MA:
Belknap Press of Harvard University Press. See
https://repository.library.georgetown.edu/handle/10822/811357.

48. Newton-Fisher NE. 2004 Hierarchy and social status in Budongo chimpanzees. *Primates* **45**,
81–87. (doi:10.1007/s10329-003-0064-6)

49. Nishida T, Kano T, Goodall J, McGrew WC, Nakamura M. 1999 Ethogram and ethnography
of Mahale chimpanzees. *Anthropol Sci* **107**, 141–188. (doi:10.1537/ase.107.141)

50. Fruth B, Hohmann G. 2003 Intra- and Inter-Sexual Aggression By Bonobos in the Context of
Mating. *Behaviour* **140**, 1389–1413. (doi:10.1163/156853903771980648)

51. Furuichi T. 1989 Social interactions and the life history of female Pan paniscus in Wamba,
Zaire. *Int J Primatol* **10**, 173–197. (doi:10.1007/BF02735199)

52. White FJ. 1996 Comparative socio–ecology of Pan paniscus. In *Great Ape Societies* (eds LF
Marchant, T Nishida, WC McGrew), pp. 29–42. Cambridge, UK: Cambridge University
Press. (doi:10.1017/CBO9780511752414.005)

53. Wilson ML, Wrangham RW. 2003 Intergroup Relations in Chimpanzees. *Annual Review of*
Anthropology **32**, 363–392. (doi:10.1146/annurev.anthro.32.061002.120046)

54. Furuichi T. 2011 Female contributions to the peaceful nature of bonobo society.
Evolutionary Anthropology: Issues, News, and Reviews **20**, 131–142.
(doi:10.1002/evan.20308)

55. Lucchesi S, Cheng L, Deschner T, Mundry R, Wessling EG, Surbeck M. 2020 Better
together? How intergroup associations affect energy balance and feeding behavior in wild
bonobos. *Behav Ecol Sociobiol* **75**, 2. (doi:10.1007/s00265-020-02943-9)

56. Hare B, Melis AP, Woods V, Hastings S, Wrangham R. 2007 Tolerance Allows Bonobos to
Outperform Chimpanzees on a Cooperative Task. *Current Biology* **17**, 619–623.
(doi:10.1016/j.cub.2007.02.040)

57. Herrmann E, Hare B, Call J, Tomasello M. 2010 Differences in the Cognitive Skills of
Bonobos and Chimpanzees. *PLoS One* **5**, e12438. (doi:10.1371/journal.pone.0012438)

58. Cronin KA, van Leeuwen EJC, Vreeman V, Haun DBM. 2014 Population-level variability in
the social climates of four chimpanzee societies. *Evolution and Human Behavior* **35**, 389–
396. (doi:10.1016/j.evolhumbehav.2014.05.004)

59. van Leeuwen EJC, Staes N, Brooker JS, Kordon S, Nolte S, Clay Z, Eens M, Stevens JMG.
2023 Group-specific expressions of co-feeding tolerance in bonobos and chimpanzees
preclude dichotomous species generalizations. *iScience* **26**, 108528.
(doi:10.1016/j.isci.2023.108528)

60. Kaburu SSK, Newton-Fisher NE. 2015 Egalitarian despots: hierarchy steepness, reciprocity
and the grooming-trade model in wild chimpanzees, Pan troglodytes. *Animal Behaviour* **99**,
61–71. (doi:10.1016/j.anbehav.2014.10.018)

61. Surbeck M, Deschner T, Schubert G, Weltring A, Hohmann G. 2012 Mate competition,
testosterone and intersexual relationships in bonobos, Pan paniscus. *Animal Behaviour* **83**,
659–669. (doi:10.1016/j.anbehav.2011.12.010)

62. Surbeck M *et al.* 2017 Sex-specific association patterns in bonobos and chimpanzees reflect
species differences in cooperation. *Royal Society Open Science* **4**, 161081.
(doi:10.1098/rsos.161081)

63. de Waal FBM, Yoshihara D. 1983 Reconciliation and Redirected Affection in Rhesus
Monkeys. *Behaviour* **85**, 224–241. (doi:10.1163/156853983X00237)

64. Altmann J. 1974 Observational Study of Behavior: Sampling Methods. *Behaviour* **49**, 227–
266. (doi:10.1163/156853974X00534)

65. Wark JD, Cronin KA, Niemann T, Shender MA, Horrigan A, Kao A, Ross MR. 2019
Monitoring the Behavior and Habitat Use of Animals to Enhance Welfare using the
ZooMonitor App. *AB&C* **6**, 158–167. (doi:10.26451/abc.06.03.01.2019)

66. Neumann C, Kulik L. 2020 EloRating: Animal Dominance Hierarchies by Elo Rating.

67. 2019 ELAN.

68. Wittenburg P, Brugman H, Russel A, Klassmann A, Sloetjes H. 2006 ELAN: a professional
framework for multimodality research. pp. 1556–1559.

69. Das M, Penke Z, van Hooff JARAM. 1998 Postconflict Affiliation and Stress-Related
Behavior of Long-Tailed Macaque Aggressors. *International Journal of Primatology* **19**, 53–
71. (doi:10.1023/A:1020354826422)

70. de Waal FBM. 1988 The communicative repertoire of captive bonobos (Pan paniscus),
compared to that of chimpanzees. *Behaviour* **106**, 183–251.
(doi:10.1163/156853988X00269)

71. RStudio Team. 2020 RStudio: Integrated Development for R.

72. Bürkner P-C. 2017 brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal*
of Statistical Software **80**, 1–28. (doi:10.18637/jss.v080.i01)

73. McElreath R. 2020 *Statistical Rethinking: A Bayesian Course with Examples in R and STAN.*
2nd edn. New York: Chapman and Hall/CRC. (doi:10.1201/9780429029608)

74. Kruschke J. 2014 *Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan.*
Academic Press.

75. Makowski D, Ben-Shachar MS, Chen SHA, Lüdecke D. 2019 Indices of Effect Existence
and Significance in the Bayesian Framework. *Front. Psychol.* **10**.
(doi:10.3389/fpsyg.2019.02767)

76. Baayen RH, Davidson RJ, Bates DM. 2008 Mixed-effects modeling with crossed random
effects for subjects and items. *Journal of Memory and Language* **59**, 390–412.
(doi:10.1016/j.jml.2007.12.005)

77. Mouginot M, Wilson ML, Desai N, Surbeck M. 2024 Differences in expression of male
aggression between wild bonobos and chimpanzees. *Current Biology* **34**, 1780-1785.e4.
(doi:10.1016/j.cub.2024.02.071)

78. de Waal FBM. 1986 The Integration of Dominance and Social Bonding in Primates. *The*
Quarterly Review of Biology **61**, 459–479. (doi:10.1086/415144)

79. Beran MJ, Savage-Rumbaugh ES, Pate JL, Rumbaugh DM. 1999 Delay of gratification in
chimpanzees (Pan troglodytes). *Developmental Psychobiology* **34**, 119–127.
(doi:10.1002/(SICI)1098-2302(199903)34:2<119::AID-DEV5>3.0.CO;2-P)

80. Anderson C, Keltner D. 2002 The role of empathy in the formation and maintenance of
social bonds. *Behavioral and Brain Sciences* **25**, 21–22. (doi:10.1017/S0140525X02230010)

81. Mitani JC. 2009 Male chimpanzees form enduring and equitable social bonds. *Animal*
Behaviour **77**, 633–640. (doi:10.1016/j.anbehav.2008.11.021)

82. Sabbi KH, Emery Thompson M, Machanda ZP, Otali E, Wrangham RW, Muller MN. 2021
Sex differences in early experience and the development of aggression in wild chimpanzees.
Proceedings of the National Academy of Sciences **118**, e2017144118.
(doi:10.1073/pnas.2017144118)

83. Bray J, Gilby IC. 2020 Social relationships among adult male chimpanzees (Pan troglodytes
schweinfurthii): variation in the strength and quality of social bonds. *Behav Ecol Sociobiol*
74, 112. (doi:10.1007/s00265-020-02892-3)

84. Sandel AA, Langergraber KE, Mitani JC. 2020 Adolescent male chimpanzees (Pan
troglodytes) form social bonds with their brothers and others during the transition to
adulthood. *American Journal of Primatology* **82**, e23091. (doi:10.1002/ajp.23091)

85. Surbeck M, Mundry R, Hohmann G. 2011 Mothers matter! Maternal support, dominance
status and mating success in male bonobos (Pan paniscus). *Proceedings of the Royal Society*
B: Biological Sciences **278**, 590–598. (doi:10.1098/rspb.2010.1572)

86. Clay Z, de Waal FBM. 2013 Development of socio-emotional competence in bonobos.
Proceedings of the National Academy of Sciences **110**, 18121.
(doi:10.1073/pnas.1316449110)

87. Heesen R, Austry DA, Upton Z, Clay Z. 2022 Flexible signalling strategies by victims
mediate post-conflict interactions in bonobos. *Philosophical Transactions of the Royal*
Society B: Biological Sciences **377**, 20210310. (doi:10.1098/rstb.2021.0310)

88. de Waal FBM, Preston SD. 2017 Mammalian empathy: behavioural manifestations and
neural basis. *Nature Reviews Neuroscience* **18**, 498–509. (doi:10.1038/nrn.2017.72)

89. Clay Z, Furuichi T, de Waal FBM. 2016 Obstacles and catalysts to peaceful coexistence in
chimpanzees and bonobos. *Behaviour* **153**, 1293–1330. (doi:10.1163/1568539X-00003335)

90. Kaufhold SP, van Leeuwen EJC. 2019 Why intergroup variation matters for understanding
behaviour. *Biology Letters* **15**, 20190695. (doi:10.1098/rsbl.2019.0695)

91. Brotcorne F, Giraud G, Gunst N, Fuentes A, Wandia IN, Beudels-Jamar RC, Poncin P,
Huynen M-C, Leca J-B. 2017 Intergroup variation in robbing and bartering by long-tailed
macaques at Uluwatu Temple (Bali, Indonesia). *Primates* **58**, 505–516. (doi:10.1007/s10329-
017-0611-1)

92. Fashing PJ. 2001 Activity and Ranging Patterns of Guerezas in the Kakamega Forest:
Intergroup Variation and Implications for Intragroup Feeding Competition. *International*
Journal of Primatology **22**, 549–577. (doi:10.1023/A:1010785517852)

93. Borgeaud C, Sosa S, Bshary R, Sueur C, van de Waal E. 2016 Intergroup Variation of Social
Relationships in Wild Vervet Monkeys: A Dynamic Network Approach. *Frontiers in*
Psychology **7**.

94. van Leeuwen EJC, Cronin KA, Haun DBM, Mundry R, Bodamer MD. 2012 Neighbouring
chimpanzee communities show different preferences in social grooming behaviour.
Proceedings of the Royal Society B: Biological Sciences **279**, 4362–4367.
(doi:10.1098/rspb.2012.1543)

95. Crockford C, Herbinger I, Vigilant L, Boesch C. 2004 Wild Chimpanzees Produce Group-
Specific Calls: a Case for Vocal Learning? *Ethology* **110**, 221–243. (doi:10.1111/j.1439-
0310.2004.00968.x)

96. Hohmann G, Fruth B. 2003 Culture in Bonobos? Between☐Species and Within☐Species
Variation in Behavior. *Current Anthropology* **44**, 563–571. (doi:10.1086/377649)

97. Cronin KA, De Groot E, Stevens JMG. 2015 Bonobos show limited social tolerance in a
group setting: a comparison with chimpanzees and a test of the relational model. *Folia*
Primatol. **86**, 164–177. (doi:10.1159/000373886)

98. DeTroy SE, Ross CT, Cronin KA, van Leeuwen EJC, Haun DBM. 2021 Cofeeding tolerance
in chimpanzees depends on group composition: A longitudinal study across four
communities. *iScience* **24**, 102175. (doi:10.1016/j.isci.2021.102175)

99. Staes N, Vermeulen K, van Leeuwen EJC, Verspeek J, Torfs JRR, Eens M, Stevens JMG.
2022 Drivers of Dyadic Cofeeding Tolerance in Pan: A Composite Measure Approach.
Biology (Basel) **11**, 713. (doi:10.3390/biology11050713)

100.   McGrew WC, Marchant LF, Scott SE, Tutin CEG. 2001 Intergroup Differences in a
Social Custom of Wild Chimpanzees: The Grooming Hand☐Clasp of the Mahale Mountains.
Current Anthropology **42**, 148–153. (doi:10.1086/318441)

101.   House BR, Silk JB, Henrich J, Barrett HC, Scelza BA, Boyette AH, Hewlett BS,
McElreath R, Laurence S. 2013 Ontogeny of prosocial behavior across diverse societies.

Proceedings of the National Academy of Sciences **110**, 14586–14591.

(doi:10.1073/pnas.1221217110)

102.   Hua X, Greenhill SJ, Cardillo M, Schneemann H, Bromham L. 2019 The ecological

drivers of variation in global language diversity. *Nat Commun* **10**, 2047.

(doi:10.1038/s41467-019-09842-2)

103.   van Leeuwen EJC, Cohen E, Collier-Baker E, Rapold CJ, Schäfer M, Schütte S, Haun

DBM. 2018 The development of human social learning across seven societies. *Nat Commun*

9, 2076. (doi:10.1038/s41467-018-04468-2)

843