Loss of PINK1 causes age-dependent mitochondrial trafficking deficits in nigrostriatal dopaminergic neurons through aberrant activation of p38 MAPK

Jingyu Zhao,[1,2] Yuanxin Chen,[2,3] Lianteng Zhi,[2] Qing Xu,[4] Hui Zhang,[2,3] and Chenjian Li[1]

Abstract

Mutations in *PTEN-induced putative kinase 1* (*PINK1*) gene cause early-onset, autosomal-recessive familial inherited Parkinson's disease (PD). Previous studies have shown age-related declines in dopamine release and ATP levels in the striatum of *Pink1*[-/-] mice, but the underlying mechanisms remain unclear. While the role of PINK1 in mitochondrial transport has been studied in cell culture and *Drosophila*, its impacts on axonal transport in dopaminergic neurons and the consequent synaptic and axonal degeneration *in vivo* remain critical questions that warrant further investigation. In this study, we developed a TH-Mito-Dendra2 transgenic mouse model to label mitochondria in dopaminergic neurons with Dendra2 protein. Using this model, we found that loss of PINK1 causes age-dependent deficits in mitochondrial axonal trafficking, with reduced anterograde movement and increased static mitochondria in acute brain slices. Pharmacological induction of reactive oxygen species (ROS) and calcium release hindered mitochondrial mobility. Additionally, *Pink1* knockout mice showed increased levels of mitochondrial calcium and ROS. Furthermore, *Pink1*[-/-] mice showed p38 MAPK hyperactivation. Treatment with the p38 inhibitor SB202190 restored mitochondrial motility and increased anterograde transport. Taken together, our findings suggest that loss of PINK1 disrupts mitochondrial trafficking by disturbing calcium homeostasis and ROS production via the p38 pathway for PD pathogenesis.

Author affiliations:

1 The MOE Key Laboratory of Cell Proliferation and Differentiation, School of Life Sciences, Peking University, Beijing, China, 100871

2 Department of Neuroscience, Thomas Jefferson University, Philadelphia, United States, 19107

3 Department of Physiology & Pharmacology, The University of Georgia, Athens, United States, 30602

4 Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology, Saudi Arabia, Thuwal, Kingdom of Saudi Arabia, 23955-6900

Correspondence to:

Chenjian Li, PhD

Professor

The MOE Key Laboratory of Cell Proliferation and Differentiation, School of Life Sciences, Peking University, Beijing, China, 100871

E-mail: Li_chenjian@pku.edu.cn

Hui Zhang, PhD

Associate Professor

Center for Neurological Disease Research, Department of Physiology & Pharmacology, College of Veterinary Medicine, The University of Georgia, 501 D.W. Brooks Drive, Athens, GA 30602

Email: Hui.Zhang1@UGA.edu

Running title: PINK1 Loss Leads to Age-Dependent Mitochondrial Trafficking Deficits

Keywords: Parkinson's disease; mitochondrial axonal transport; acute brain slices; calcium homeostasis; oxidative stress

Introduction

Parkinson's disease (PD) is a major neurodegenerative disorder affecting people above 65 years old, characterized by dopamine neuron loss in the substantia nigra pars compacta (SNc) leading to striatal dopamine deficiency.[1] While the pathogenesis of PD is complex, mounting evidence implicates mitochondrial dysfunction as a key initiating factor.[2] Mitochondrial abnormalities in PD include altered morphology, impaired energy production, increased oxidative stress, dysregulated dynamics and calcium homeostasis, and disrupted mitophagy. However, it remains unclear whether these mitochondrial defects are primary causes or downstream consequences in PD. The identification of *PINK1* as a PD gene provided direct evidence linking mitochondrial dysfunction to PD pathogenesis.[3] PINK1 regulates key mitochondrial processes like mitophagy, calcium homeostasis, and fission in neurons, underscoring its relevance in studying mitochondrial dysfunction's role in PD. PINK1 loss-of-function models demonstrate mitochondrial defects like increased fission,[4] depolarization,[5] impaired electron transport chain complexes I and III,[6,7] and calcium overload.[8] Additionally, PINK1 interacts with Miro/Milton, implicating its role in mitochondrial trafficking.[9]

Neurons, especially synapses, are highly energy demanding. A human cortical neuron utilizes ~4.7 billion ATPs/second at rest,[10] underscoring the importance of mitochondrial trafficking. Following evidence that PINK1 can interact with trafficking adaptor proteins,[9] several studies have explored the relationship between PINK1 and mitochondrial stand mitochondrial trafficking in different models. One study demonstrated that overexpression of PINK1 or PARKIN in rat hippocampal axons significantly reduced mitochondrial motility. For impaired mitochondria, PINK1 phosphorylates Miro, leading to PARKIN-dependent Miro degradation and subsequent dissociation of kinesin from mitochondria, thus sequestering depolarized mitochondria.[11] Another study demonstrated that D-AKAP1/PKA (D-AKAP1, dual-specificity A Kinase Anchoring protein 1; PKA, Protein Kinase A) can restore the mitochondrial trafficking deficiency in cortical neurons from PINK1 knockout (KO) mice.[12] While these studies support the critical role of PINK1/Parkin pathway in regulating mitochondrial trafficking[13], most research utilizes *in vitro* systems or *Drosophila* models, which differ from *in vivo* conditions. Additionally, most studies are not specific to dopaminergic neurons. From this perspective, further investigations closer to *in vivo* settings and focused on dopaminergic neurons are needed.

Here, we generated a new Mito-Dendra2 tool mouse model to specifically label mitochondria in dopaminergic neurons and established a new approach to directly observe mitochondrial trafficking in nigrostriatal dopaminergic neurons in acute brain slices from a *Pink1*[-/-] mouse model. We found an age-dependent impairment of mitochondrial motility and anterograde transport in dopaminergic neurons lacking PINK1 mediated through the aberrant p38 pathway. These results provide a possible explanation for our previous findings of decreased dopamine release and ATP production in the striatum of *Pink1*[-/-] mice at early middle-age.[14]

Materials and methods

Animals

Mice were housed in a temperature and humidity controlled specific pathogen-free (SPF) facility under a 12-hour light/dark cycle schedule with food and water available ad libitum. Mice were on FVB/N mouse background (Taconic). All experimental protocols were approved by the Animal Care and Use Committees of Thomas Jefferson University, Peking University, and The University of Georgia.

Construction, modification and confirmation of TH-MitoDendra2-BAC

BAC clone covering the entire tyrosine hydroxylase (*Th*) gene was obtained from GENSAT RPCI23-350E13 (http://genome.ucsc.edu/cgi-bin/hgTracks?clade=mammal&org=Mouse&db=mm9&position=RP23-350E13).

Dendra2-SV40 polyA from Clontech pCMV vector was ligated with a mitochondrial targeting sequence.[15] The mitochondrial targeting sequence (MTS) was from human subunit 8A of cytochrome c oxidase (COX8A): 5'ATGTCCGTCCTGACGCCGCTGCTGCTGCGGGGCTTGACAGGCTCGGCCCGGCG GCTCCCAGTGCCGCGCGCCAAGATCCATTCGTTGGGGGATCCACCGGTCGCCACC 3'. Mito-Dentra2 was inserted to the exon1 of *Th* gene with homologous recombination. Homologous arms, A arm (642 bp) and B arm (538 bp), from the first exon of *Th* were ligated to 5' and 3' ends of the Mito-Dendra2 cassette by PCR amplification respectively and inserted into the shuttle vector following the standard protocol.[16] The shuttle vectors were introduced into the RP23-350E13 competent cells to construct the co-integrated BACs using

electroporation. Recombinants were identified by chloramphenicol and ampicilin resistant selection followed by PCR analysis.[16] Purified TH-MitoDendra2 BACs were used to generate TH-MitoDendra2 transgenic mice.

TH-MitoDendra2-BAC transgenic mice

TH-MitoDendra2-BACs were used for pronuclear injection in the Taconic FVB/NTac background. Founders and progeny carrying the TH-MitoDendra2 transgene were identified by PCR analysis of tail DNA and direct sequencing. The expression pattern of Mito-Dendra2 was confirmed using TH immunofluorescence in TH-MitoDendra2-BAC transgenic mice. The transgenic mice were maintained by breeding with wild-type (WT) Taconic FVB/NTac mice.

Immunofluorescence

Mice were anesthetized with ketamine/xylazine (100 mg/10 mg per kg) and then perfused with ice-cold 4% paraformaldehyde in phosphate-buffered saline (PBS) and postfixed for 24 hours. Coronal sections, 30 μm thick, were obtained using a Thermo Scientific Microm HM550 Cryostat. Serial sections were collected into 10 wells, with each well containing every 10th section. Floating sections were stored in antifreeze buffer (30% glycerol and 30% ethylene glycol in PBS) at $-20°C$. The sections were stained with a polyclonal anti-tyrosine hydroxylase (TH) antibody (Millipore, MAB5280) at a 1:1000 dilution for 36 hours at 4°C, followed by incubation with a Dylight 594-conjugated secondary antibody (1:500, Life Technologies). Fluorescent images were captured using a Leica SP3800 laser scanning confocal microscope.

Acute sagittal slice preparation

Animals were anesthetized with either isoflurane (1-2%) or ketamine/xylazine (100 mg/10 mg per kg) and then underwent transcardiac perfusion with ice-cold, pre-oxygenated cutting solution containing (in mM): 125 NaCl, 2.5 KCl, 26 NaHCO3, 3.7 MgSO4, 0.3 KH2PO4, and 10 glucose, pH 7.4). The brains were immediately dissected out in cold, pre-oxygenated cutting solution. Sagittal nigrostriatal projection slices were sectioned at a thickness of 300 μm using a vibrating microtome (VT1200, Leica, Solms, Germany) with an angle of approximately 21°. The slices were then recovered in oxygenated artificial cerebrospinal fluid

(aCSF) containing (in mM): 125 NaCl, 2.5 KCl, 26 NaHCO$_3$, 2.4 CaCl$_2$, 1.3 MgSO$_4$, 0.3 KH$_2$PO$_4$, 10 glucose, 2 HEPES, pH7.4, at room temperature for > 30 minutes.[14]

Two-photon laser microscopy imaging of mitochondrial trafficking in brain slices

Parasagittal slices containing nigrostriatal projections were obtained from TH-MitoDendra2 transgenic mice. The brain slices were positioned within a recording chamber and superfused with oxygenated aCSF at a flow rate of 1.5 mL/min. Optical imaging of Dendra signals were acquired using an Ultima multiphoton laser scanner (Bruker) for BX51/61 Olympus microscope with a Titanium-sapphire laser (Chameleon-Ultra2, Coherent Laser Group), equipped with a 20× 1.0 NA water immersion objective (XLUMPLFL20XW, Olympus). The Dendra2 protein was excited at a wavelength of 900 nm, and its fluorescence was collected at 500-560 nm. Images were captured in 8-bit resolution and 4 μs dwell time, with regions of interest set at 512 × 512 pixels. The sampling rate ranged from 0.2 to 0.5 seconds per frame. All experiments were conducted at 33-35 °C.

Images were processed using the open-source software Fiji (NIH) and MATLAB (Version 8.5.0 R2015a, MathWorks). Intensity-based alignment was performed to register images at different time points. To measure mitochondrial motility velocity, the plugins KymographClear and KymographDirect were used to generate kymographs representing moving distance versus moving duration. The number and percentage of mitochondria in different states—static, retrograde, and anterograde—were quantified and measured.

Stereotaxic injection of AAV

3-4-month-old male mice of *Pink1*$^{+/+}$ and *Pink1*$^{-/-}$ were anesthetized with isoflurane (1-2%) or ketamine/xylazine (100 mg/10 mg/kg). A small craniotomy, approximately 0.8 mm in diameter, was made over the right midbrain (−3.4 mm caudal, +1.0 mm lateral). To achieve expression of mito-GCaMP6 in dopaminergic neurons, 0.1-0.2 μl of AAV-GCaMP6-mito virus (10^13 vg/ml, a gift from Dr. James Surmeier's lab at Northwestern University) was pressure-injected through a syringe into the midbrain at a depth of -4.2 mm ventral from the dura surface. Following the injections, the incisional wound on the skull was sealed with tissue adhesive (3M Vetbond). The mice were placed on a heating pad until fully recovered

and then returned to the isolation room. Carprofen (5 mg/kg, Sigma Aldrich) was administered intraperitoneally once per day for at least 3 days to reduce pain.

2PLSM mito-GCaMP6 imaging in acute brain slices

Mitochondrial Ca^{2+} levels were measured with the mitochondrially targeted Ca^{2+}-sensitive probe mito-GCaMP6.[17] Two-three weeks after AAV virus injection, mice were prepared for coronal brain slice sectioning following transcardiac perfusion with ice-cold, oxygenated cutting solution. After a recovery period of 30-45 minutes, the coronal brain slices were placed in a chamber perfused with oxygenated aCSF. To measure free Ca^{2+} levels in mitochondria, the slices were sequentially treated with the following solutions: (1) regular aCSF, (2) 500 µM EGTA + aCSF without Ca^{2+} for 30-40 minutes, (3) 500 µM EGTA + 1 µM ionomycin + aCSF without Ca^{2+} for 30 minutes, and (4) 1 µM Ionomycin + aCSF with high Ca^{2+} (3 mM) for 20 minutes. Time series images were acquired every 5 minutes to determine the minimal and maximal fluorescence intensity. The fluorescence intensity at the end of treatment (2) was defined as the minimum fluorescence (Fmin), and the fluorescence intensity at the end of treatment (4) was defined as the maximum fluorescence (Fmax). The relative free calcium level was calculated using the equation: (F - Fmin) / (Fmax - Fmin).

Immunoblot

Tissues were homogenized and lysed with RIPA buffer (Millipore Sigma, R0278) supplemented with fresh protease inhibitors (Roche, 11873580001) and phosphatase inhibitors (Roche, 04906845001). The homogenate was further lysed for 1 hour at 4°C and cleared by centrifugation at 12,000 × g for 15 minutes at 4°C. The supernatants were collected, and protein concentrations were determined using a BCA assay kit (Thermo Scientific Pierce, 23227). Protein samples (30-50 µg) were separated using Mini-PROTEAN® TGX™ Precast Gels (Bio-Rad). After blocking, the blots were incubated with primary antibodies overnight at 4°C. Following washing with 1× TBST (0.1% Tween-20), the blots were incubated with secondary antibodies (1:10000, IRDye® 800RD, Li-COR 926-32211; IRDye® 680RD, Li-COR, 926-68070) for 1 hour and visualized using the Odyssey Infrared Imager (Li-COR, 9120). The relative protein abundance was quantified using Image Studio software (Li-COR).

TH-Immunogold labeling and electron microscopy

Mice were perfused intracardially with freshly prepared fixatives. The fixative consisted of 3.75% acrolein and 2% paraformaldehyde (PFA) in 0.1 M phosphate buffer, pH 7.4 (PB). After fixation, brains were sectioned at 50 μm in PB using a vibratome subsequently fixed at 4°C overnight in 4% PFA/PB. Sections were treated with 50 mM glycine in $1\times$ PBS for 30 minutes to reduce free aldehydes and rinsed extensively with PB three times. The sections were then permeabilized with Tris-buffered saline (TBS) containing 3% goat serum, 1% bovine serum albumin (BSA), and 0.04% Triton X-100 for 30 minutes. The primary mouse anti-TH antibody (#MAB5280, Millipore Sigma) was diluted 1:100 in PBS with 3% goat serum and 1% BSA. The sections were incubated with the primary antibody at 4°C for 48 hours. After washing three times (10 minutes each) with PBS, the sections were labeled with 0.8 nm gold-conjugated goat anti-mouse secondary antibody (#800.177, Aurion) diluted in a buffer containing 3% goat serum, 0.8% BSA, and 0.1% cold fish gelatin in PBS. The sections were incubated with the secondary antibody at 4°C for 48 hours. Following one wash with dilution buffer and six washes with PBS, sections were post-fixed with 2.5% glutaraldehyde in PBS. After several washes with PBS, the sections were exposed to Enhancement Conditioning Solution (ECS; #500.055, Aurion) diluted 1:10 with ddH2O, for four times at 10 minutes each. Ultra-small gold particles were enlarged using R-Gent SE-EM Silver Enhancement Reagents (#500.044, Aurion) for 150 minutes. The sections were then rinsed in ECS before being transferred to PBS. Tissue sections were transferred to PBS in Coors dishes and fixed in 1% osmium tetroxide in PB for 60 minutes on ice. After rinsing in ddH2O, the sections were sequentially dehydrated in 30%, 50%, 70%, 85%, and 95% ethanol solutions, followed by two exposures to 100% ethanol and then two exposures to acetone. Tissue sections were incubated overnight in a 1:1 mixture of epoxy resin (#14120, EMbed 812, Electron Microscopy Sciences) and acetone before being transferred to pure EMbed 812 for 3 hours (three times, 1 hour each). The sections were then embedded in EMbed 812 and polymerized for 24 hours at 65°C. Plastic blocks were trimmed and sectioned with a Leica ultramicrotome (EM UC7, Leica) equipped with a 35° diamond knife (Diatome). Serial ultrathin sections (75 nm) were collected on single-slot formvar-coated copper grids. After post-staining with uranyl acetate and lead citrate, the grids were examined under a transmission electron microscope (Tecnai G^2 Spirit BioTWIN, FEI) equipped with a digital camera (Orius 832, Gatan) at 120kV.

Antibodies and Reagents

The antibodies and reagents used in the study include: Anti-Tyrosine Hydroxylase (rabbit, 1:5000 for IF, Pel-Freez, P40101-0), Anti-Tyrosine Hydroxylase (mouse, 1:500 for IF, Millipore, MAB5280), Anti-Kinesin heavy chain (mouse, 1:500 for WB, Millipore, MAB1614), Anti-Dynein medium chain (mouse, 1:1000 for WB, Millipore, MAB1618), Anti-Miro1 (rabbit, 1:1000 for WB, Antibodies_online, ABIN635090), MAPK Family Antibody (rabbit, 1:1000 for WB, Cell signaling, 9926), Phospho-MAPK Family Antibody (rabbit, 1:1000 for WB, Cell signaling, 9910), Anti-MnSOD (rabbit, 1:1000 for WB, Abcam, ab68155), Anti-MnSOD (rabbit, 1:100 for IF, Abcam, ab13534), and Anti-β-actin (rabbit, 1:1000 for WB, Cell signaling, 4970T). Secondary antibodies used were DyLight 594 Antibody (Goat anti-rabbit, 1:500 for IF, Invitrogen, 35560), Alexa Fluor 488nm (Goat anti-mouse, 1:1000 for IF, Invitrogen, A-11029), Alexa Fluor 594nm (Goat anti-rabbit, 1:1000 for IF, Invitrogen, A-11012), IRDye® 800CW (Goat anti-rabbit, 1:5000 for WB, Li-COR, 926-32211), and IRDye® 680RD (Goat anti-mouse, 1:5000 for WB, Li-COR, 926-68070). Additional reagents included A23187 (Sigma-Aldrich, C7522), CGP37157 (Sigma-Aldrich, C8874), menadione (Sigma-Aldrich, M5625), and ionomycin (Sigma-Aldrich, I3909).

Data and Statistical Analysis

Data were presented as mean ± SEM unless otherwise specified. Statistical analyses were conducted using Prism 9.0 (GraphPad), including the unpaired t test, Mann–Whitney test and one-way ANOVA. Differences were considered statistically significant when $P < 0.05$. Statistical analysis was based on the number of slices or experiments unless otherwise noted. "N" denotes the number of mice, and "n" denotes the number of experiments.

Results

Generation and characterization of TH-MitoDendra2 BAC transgenic mice

To specifically label mitochondria in dopaminergic neurons, we generated a bacterial artificial chromosome (BAC) clone encompassing the entire tyrosine hydroxylase (*Th*) gene (Figure 1A). A mitochondria-targeting photo-convertible fluorescent protein Dendra2 gene[15,16] was inserted in the *Th* exon 1 (Figure 1A). The co-localization of Mito-Dendra2 and the immunofluorescence for TH confirmed the specificity of Mito-Dendra2 in dopaminergic neurons within the SNc and ventral tegmental (VTA) area (Figure 1B-1E). We also verified the specificity of Mito-Dendra2 signals at the dopaminergic terminals in the striatum and olfactory bulb (Figure S1) and the results showed Mito-Dendra2 only expressed in TH positive neurons and neurites. Moreover, the co-localization of Mito-Dendra2 signals with MitoTracker staining verified the subcellular localization of Mito-Dentra2 in mitochondria (Figure 1F). Using this model, we can visualize the continuous and intact projection bundles from dopaminergic neurons in the SNc to the striatum in the sagittal view (Figure 1E, S2). These results demonstrate that Mito-Dendra2 is specifically localized in the mitochondria of dopaminergic neurons and their projections, providing a novel and powerful tool for studying mitochondrial trafficking in these neurons.

Investigation of mitochondrial trafficking in acute brain slices with two-photon laser microscopy (2PLSM)

To evaluate mitochondrial trafficking in *Pink1*[-/-] mice, we crossed them with Mito-Dendra2 transgenic mice to generate *Pink1*[-/-] /MD+ transgenic mice (Figure 2A). We chose to use acute sagittal brain slices (Figure 2A), which closely mimic *in vivo* conditions, to monitor mitochondrial trafficking. Two-photon laser scanning microscopy (2PLSM) was employed to track mitochondrial movement. Using time-lapse imaging, we visualized and recorded videos showing mitochondrial trafficking along the axon bundles (Figure 1E, Figure 2A, and B). Mitochondrial movement was primarily classified into three states: stationary, anterograde (moving towards the terminals), and retrograde (moving towards the cell body). As depicted in Figure 2B and Video S1, we clearly captured mitochondrial movement at sequential time

points. After collecting mitochondrial trafficking data, we used the software Fiji with two plugins, KymographDirect and KymographClear,[18] to analyze the differences in the percentage and velocity of moving mitochondria (Figure 2C).

Age-dependent mitochondrial axonal transport deficits in *Pink1*[-/-] mice

After establishing the procedure, we evaluated the alterations in mitochondrial trafficking between *Pink1*[-/-] mice and age-matched wild-type (WT) littermates at different ages (Figure 3, Tables S1-S3). The percentages of stationary, anterograde, and retrograde mitochondria were similar in 2-4-month-old *Pink1*[-/-] mice and WT littermates (Figures 3A1-A2, 3B1-B2, 3C, Table S1). The average velocities of anterograde and retrograde movement were also comparable (Figure 3D, Table S1). Approximately 84-85% of mitochondria were stationary in both genotypes at 2-4 months, which is higher than typically reported in cultured cells.[13,19] In these mice, about 11% of mitochondria moved anterogradely and 4-5% moved retrogradely, differing from the more balanced 15% anterograde and 15% retrograde movement seen in cultured cells.

In 5-7-month-old *Pink1*[-/-] mice, the proportion of stationary mitochondria increased significantly ($90.79 \pm 0.69\%$ (*Pink1*[-/-]) *vs* $85.18 \pm 0.65\%$ (WT), $P < 0.0001$). Anterograde movement decreased from $9.95 \pm 0.94\%$ to $5.63 \pm 0.77\%$ ($P = 0.0019$), while retrograde movement showed a slight decrease ($3.58 \pm 0.52\%$ (*Pink1*[-/-]) *vs* $4.88 \pm 0.75\%$ (WT), $P = 0.1624$; Figures 3A3-A4, 3B3-B4, 3E, Table S2). Additionally, the anterograde velocity decreased significantly (0.52 ± 0.02 µm/s (*Pink1*[-/-]) *vs* 0.60 ± 0.03 µm/s (WT), $P = 0.0335$) (Figure 3F, Table S2). This reduction in anterograde movement may result in fewer mitochondria being transported to axonal terminals.

In *Pink1*[-/-] mice older than 10 months, the percentage of static mitochondria increased significantly compared to WT littermates ($91.72 \pm 1.07\%$ (*Pink1*[-/-]) *vs* $86.55 \pm 1.88\%$ (WT), $P = 0.0251$), while anterograde movement decreased to $2.54 \pm 0.60\%$ ($P = 0.008$) (Figures 3A5-A6, 3B5-B6, 3G, Table S3). In these older *Pink1*[-/-] mice, retrograde mitochondrial transport exceeded anterograde transport, reaching $6.24 \pm 1.00\%$ (Figure 3G, Table S3). Since retrograde-moving mitochondria are typically associated with damage[19], this suggests an increased proportion of dysfunctional mitochondria in *Pink1*[-/-] mice. Furthermore, the

anterograde velocity was significantly decreased to 0.36 ± 0.03 µm/s ($P = 0.004$) compared to age-matched WT controls (0.48 ± 0.02 µm/s), while retrograde velocity remained unchanged (Figure 3H, Table S3).

In summary, $Pink1^{-/-}$ mice exhibited age-dependent deficits in mitochondrial axonal trafficking. These deficits, which were not present in 2-4-month-old mice, emerged around 5 months of age and persisted beyond 10 months of age. The trafficking abnormalities included an increased proportion of static mitochondria, a decreased proportion of antegrade motile mitochondria, and slower anterograde velocities. These findings differ from previous reports in cultured neurons[11] and $Drosophila$[20], highlighting potential differences between $in\ vivo$ and $in\ vitro$ models, as well as between invertebrate and vertebrate systems.

Mitochondrial calcium elevation and transport deficits in $Pink1^{-/-}$ mice

Mitochondrial axonal transport is a tightly regulated process involving many factors. Cytosolic Ca^{2+} can modulate mitochondrial motility via the Miro protein.[21] Furthermore, mitochondria themselves play a pivotal role in calcium homeostasis.[22] we first evaluated mitochondrial calcium levels in 4-month-old $Pink1^{-/-}$ mice and WT littermates using AAV9-TH-mito-GCaMP6, a modified GCaMP6 virus that is specifically expressed in mitochondria in dopaminergic neurons. Figure 4A and 4B show that the relative $[Ca^{2+}]_{mt}$ (relative fluorescence intensity) was significantly increased in $Pink1^{-/-}$ mice ($78.6 \pm 4.3\%$) compared to WT littermates ($35.7 \pm 4.6\%$, $P = 0.0001$). Next, we investigated the effect of elevated mitochondrial calcium on mitochondrial trafficking. We incubated WT acute brain slices with 5 µM CGP37157, an inhibitor of the mitochondrial Na☐/Ca²☐ exchanger (NCLX) that causes mitochondrial calcium overload.[23] Mitochondrial motility was significantly decreased after 2 hours of CGP37157 incubation (Figures 4C and 4D, Video S2 and S3). The proportion of static mitochondria increased from $85.26 \pm 0.69\%$ to $90.29 \pm 1.17\%$ ($P = 0.0048$), while anterograde movement declined from $8.69 \pm 1.07\%$ to $5.62 \pm 0.85\%$ ($P = 0.0432$) (Figure 4D).

This phenomenon of altered mitochondrial motility following $[Ca^{2+}]_{mt}$ elevation was reminiscent of the findings observed in 5- to 7-month-old and above 10-month-old $Pink1^{-/-}$

mice, suggesting a similar underlying mechanism contributing to the mitochondrial trafficking defects in *Pink1*[-/-] mice. The velocities of anterograde and retrograde mitochondrial transport remained unaffected by the calcium perturbations in this experimental context (Figure 4E).

Reduced mitochondrial anterograde movement by ROS elevation

Mitochondrial calcium overload impairs normal mitochondrial function, leading to increased reactive oxygen species (ROS) generation and decreased ATP production.[24,25] To elucidate the relationship between mitochondrial calcium homeostasis and reactive oxygen species (ROS) generation in acute brain slices, we utilized CGP37157 to induce mitochondrial calcium overload in brain slices derived from roEGFP transgenic mice and evaluate the ROS alteration[26] after treatment (Figure 5). Our results showed a significant increase in $[Ca^{2+}]_{mt}$, which led to an elevation in ROS levels ($P = 0.0005$, Figure 5C).

Subsequently, to investigate the impact of elevated ROS levels on mitochondrial trafficking dynamics, we exposed acute brain slice preparations from WT mice (5-7-month-old) to 50 µM menadione (Figure 6A-6D). Menadione is a widely employed compound known to exacerbate endogenous ROS production in cultured cell systems.[27] Following a 20-minute incubation with 50 µM menadione, the proportion of stationary mitochondria significantly increased from 88.71 ± 1.49% to 96.75 ± 0.86% ($P = 0.0036$, Figure 6C). Anterograde moving mitochondria were reduced by nearly 70%, from 6.70 ± 1.14% to 2.20 ± 0.73% ($P = 0.0202$), while retrograde movement also declined (4.59 ± 0.72% (aCSF control), 1.05 ± 0.18% (MED), $P = 0.0177$, Figure 6A, 6B, and 6C, Video S4 and S6). The average velocity of both anterograde and retrograde movement remained unchanged after MED treatment (Figure 6D). To rule out the possibility that decreased motility was due to DMSO toxicity, we also evaluated the effect of DMSO treatment on mitochondrial trafficking. The results showed no significant change in the percentage of moving mitochondria and their velocity between aCSF control and DMSO treatment (Figure 6C and 6D, Video S4 and S5). Additionally, we examined the effect of H_2O_2 treatment, another reagent commonly used to induce ROS generation and observed similar results on mitochondrial trafficking (data not shown). These findings suggest that artificially increasing endogenous ROS levels in WT

brain slices can impair mitochondrial motility, particularly affecting anterograde movement, similar to the deficits observed in *Pink1*[-/-] mice.

We then evaluated ROS levels in *Pink1*[-/-] mice. Manganese superoxide dismutase (MnSOD), a crucial mitochondrial antioxidant enzyme, detoxifies oxygen free radicals generated during mitochondrial respiration. Evidence indicates that oxidative stress significantly upregulates MnSOD expression levels.[28] We quantified MnSOD expression levels in the striatum and SNc regions of *Pink1*[-/-] mice and WT littermates. Western blot analysis revealed a significant upregulation of MnSOD levels in 5-7-month-old *Pink1*[-/-] mice compared to WT littermates in both the striatum and SNc regions (Figure 6E and 6F), with an approximate 2-fold increase in striatal MnSOD expression ($P = 0.0019$) and a 1.75-fold elevation in the region of substantia nigra ($P = 0.0144$). Complementary immunofluorescence staining corroborated these findings, showing significantly higher MnSOD immunoreactivity in dopaminergic neurons from *Pink1*[-/-] mice compared to WT controls (Figure 6H and 6I). Quantitative analysis revealed an approximate 4-fold increase in relative MnSOD fluorescence intensity in *Pink1*[-/-] mice ($P < 0.0001$). Collectively, these data provide compelling evidence for elevated oxidative levels in PINK1 knockout mice. The excessive ROS generation in acute brain slices from WT mice induced by menadione treatment was associated with impaired mitochondrial motility. These findings underscore a potential pathogenic link between PINK1-mediated mitochondrial dysfunction, dysregulated redox homeostasis, and disrupted mitochondrial trafficking, which may contribute to the degeneration of dopaminergic neurons in PD.

p38 MAPK pathway hyperactivation by loss of PINK1

We have already demonstrated that the of PINK1 causes mitochondrial calcium overload in dopaminergic neurons, resulting in a significant increase in intracellular ROS levels. Excessive ROS could lead to defects in mitochondrial axonal transport. However, the specific mechanisms by which excessive ROS affect mitochondrial movement in neurons remain to be fully elucidated. Previous studies have shown that intracellular ROS can inhibit the function of the motor-adaptor complex that regulates mitochondrial motility through p38α MAPK pathway, thus reducing mitochondrial motility *in vitro*.[27]

We then analyzed p38 phosphorylation level in *Pink1$^{-/-}$* mice (6-7-month-old) (Figure 7A and 7B). Immunoblot results showed that p38 phosphorylation levels were approximately threefold higher in *Pink1$^{-/-}$* mice than in WT mice in the striatum ($P = 0.0286$). The substantia nigra region exhibited a similar increase in p38 phosphorylation levels ($P = 0.0079$). Elevated p38 phosphorylation indicates heightened activation of the p38 pathway. Collectively, these results suggest hyperactivation of the p38 MAPK pathway in *Pink1$^{-/-}$* mice.

We extended our analysis to examine the phosphorylation levels of two other MAPK family pathways: JNK (Figure S3) and ERK (Figure S4). JNK phosphorylation levels were also elevated in *Pink1$^{-/-}$* mice, while ERK phosphorylation levels remained unchanged. Both p38 and JNK pathways play crucial roles in cellular stress responses. The observed hyperactivation of these pathways in *Pink1$^{-/-}$* mice aligns with our previous results and provides additional support for the oxidative stress hypothesis in PINK1 deficiency (Figure 6H).

Improved mitochondrial motility in *Pink1$^{-/-}$* mice by p38 inhibition

To validate whether inhibiting p38 activity could alleviate mitochondrial trafficking deficits in *Pink1$^{-/-}$* mice, we treated acute brain slices from these mice with the p38 inhibitor SB202190 (Figure 7C-7F, Video S7-S8). Incubating *Pink1$^{-/-}$* mice brain slices in aCSF containing 50 μM SB202190 for one hour significantly increased mitochondrial motility in dopaminergic neurons (Figure 7E). The proportion of immobile mitochondria decreased from $90.93 \pm 0.51\%$ to $84.59 \pm 0.61\%$ ($P < 0.0001$). The percentage of anterograde-moving mitochondria significantly increased from $4.96 \pm 0.86\%$ to $11.12 \pm 1.36\%$ ($P = 0.0037$). Retrograde mitochondrial movement was not significantly affected (aCSF: $4.11 \pm 0.78\%$ *vs.* SB202190: $4.29 \pm 1.15\%$). Additionally, SB202190 treatment did not alter the velocity of either anterograde or retrograde mitochondrial movement (Figure 7F).

We also verified the effectiveness of SB202190 treatment in *Pink1$^{-/-}$* mouse brain slices through immunoblot analysis, which showed a significant ~62% reduction in p38 phosphorylation (Figure S6). This substantial decrease in p38 phosphorylation led to a significant rescue of mitochondrial motility defects in dopaminergic neurons of *Pink1$^{-/-}$* mice,

particularly by increasing the percentage of anterograde movement. However, SB202190 treatment did not improve the decreased anterograde velocity in *Pink1*[-/-] mice.

We also analyzed the protein expression involved in axonal mitochondrial transport (Figure S5). The adaptor protein Miro1 showed a significant reduction in *Pink1*[-/-] mice compared to WT littermates: approximately a 15% decrease in the striatum (P = 0.0286) and approximately a 24% decrease in the substantia nigra (P = 0.0317). This reduction in Miro1 may lead to decreased mitochondrial attachment to microtubules and reduced motility. However, we observed no significant changes in the expression of the motor proteins kinesin and dynein in *Pink1*[-/-] mice.

Defective mitochondria at dopaminergic neuron terminals in a *Pink1*[-/-] mice

Pink1[-/-] mice exhibited age-dependent deficits in striatal dopamine release and ATP levels [14]. This observation, along with the impaired mitochondrial axonal transport in *Pink1*[-/-] dopaminergic neurons, led us to hypothesize that reduced anterograde movement of mitochondria in *Pink1*[-/-] dopaminergic neurons might result in a depletion of healthy mitochondria at their axon terminals. To test this hypothesis, we employed tyrosine hydroxylase (TH) immunogold electron microscopy to directly assess mitochondrial morphology and quantify mitochondrial numbers. In WT mice (8-9-month-old), dopaminergic neurons exhibited relatively intact mitochondrial cristae, indicating overall healthy mitochondria (Figure 8A). In contrast, *Pink1*[-/-] mice displayed notable mitochondrial abnormalities, including rounded mitochondrial shapes, indistinct cristae, and vacuolated internal structures, suggesting mitochondrial fragmentation (Figure 8B). In *Pink1*[-/-] mice, there was a significant decrease in the proportion of healthy mitochondria within dopaminergic terminals (P = 0.0079). However, the overall percentage of healthy mitochondria in the striatum remained unchanged (Figure 8C, P = 0.4524). Compared with WT mice, the circularity (P < 0.0001) and aspect ratio (P < 0.0001) of mitochondria in the dopaminergic neurons of *Pink1*[-/-] mice were significantly reduced (Figure 8D), indicating that the mitochondria at the terminals of dopaminergic neurons in *Pink1*[-/-] mice tended to be more rounded.

Taken together, our findings indicate that the loss of PINK1 leads to a significant increase in mitochondrial calcium levels in the SNc dopaminergic neurons by disrupting specific regulatory mechanisms. This disturbance in mitochondrial calcium homeostasis results in increased oxidative stress and ROS production, activate the p38 signaling cascade. Previous studies have shown that p38 can directly phosphorylate kinesin heavy chains (KHCs), thereby inhibiting axonal transport.[27,29–32] Based on these observations, we propose a working model (Figure 8G): the increase in ROS caused by loss of PINK1 activates the p38 MAPK pathway, leading to p38-mediated phosphorylation of the anterograde motor protein KIF5. This phosphorylation disrupts the mitochondrial anterograde transport complex, reducing the percentage of mitochondria undergoing anterograde movement. Consequently, the decreased anterograde trafficking of mitochondria results in a lower quantity of healthy mitochondria at the terminals of dopaminergic neurons. This deficit may lead to a reduction in ATP content at the neuron terminals, causing an energetic shortfall and decreased in dopamine release.

Discussion

Here, we developed and validated a novel approach that synergistically combines Mito-Dendra2 mice, acute brain slice preparations, and 2PLSM to investigate mitochondrial axonal transport within dopaminergic neurons. This strategy allowed us to robustly interrogate mitochondrial transport kinetics in an *ex vivo* setting that more accurately recapitulates the pathophysiological scenario of PD in dopaminergic neurons, owing to the following key advantages: **(i)** Cell-type specificity: We focused specifically on the nigrostriatal dopaminergic neurons, which are the right and most vulnerable neuronal population in PD pathogenesis. **(ii)** Preserved neuronal network: Utilization of acute brain slices from mice provides a more integrated and physiologically relevant neuronal network that closely mimics *in vivo* conditions. Notably, the relative thickness of these slices helps maintain the coherence of the neuronal circuitry, including interactions among neurons, astrocytes, and glial cells, for a considerable duration, despite potential mechanical damage from sectioning. **(iii)** Sustained neuronal functionality and pharmacological interventions: This approach not only preserves the intrinsic functional activity of dopaminergic neurons for an extended duration but also enables us to perform pharmacological manipulations and assess their effects on neuronal function and mitochondrial dynamics within an integrated *ex vivo* system.

Using this method, we investigated the mitochondrial trafficking alteration in our previous reported *Pink1*[-/-] mice.[14] Our findings reveal that loss of PINK1 causes age-dependent mitochondrial trafficking impairments in dopaminergic neurons, reducing healthy mitochondrial presence at terminals and provides a plausible mechanistic explanation for our previous finding.[14] Corroborating our findings, previous studies have also demonstrated that genetic ablation of *Pink1* in mouse primary cortical and midbrain neurons led to a significant reduction in the proportion of mitochondria undergoing anterograde movement along dendrites.[12] Interestingly, some studies investigating loss of PINK1 in different models or neuronal subtypes have reported contrasting findings. Researches in rat primary hippocampal and *Drosophila* demonstrated that PINK1 deficiency led to an increase in the proportion of motile mitochondria.[11,33] These divergent findings suggest that the effects of PINK1 deficiency on mitochondrial dynamics are context-dependent, varying across species, cell types, and developmental stages. This underscores the critical need to elucidate the molecular mechanisms underlying the differential susceptibility of specific neuronal populations to ubiquitous genetic perturbations.

Regarding the underlying mechanism of these deficits, we observed elevated ROS levels and p38 hyperactivation in *Pink1*[-/-] mice. Pharmacological inhibition of p38 rescued mitochondrial motility. Previous studies have elucidated a potential mechanism for how p38 activation specifically impacts mitochondrial axonal transport dynamics and they have proposed that heightened p38 activity can lead to an increase in the phosphorylation status of the KIF5 protein, a key component of the kinesin motor complex.[27,29–32] Notably, this enhanced phosphorylation of KIF5 results in a reduced binding affinity for microtubules, without affecting the overall expression levels of the kinesin motor protein itself.[27,29–32] Further validation is still needed in *Pink1*[-/-] mice.

Collectively, our studies introduce a novel approach to specifically investigate mitochondrial trafficking dynamics within dopaminergic neurons in Parkinson's disease mouse models. Applying this approach to *Pink1*[-/-] mice, we observed a unique phenotype of mitochondrial trafficking in dopaminergic neurons that differed from previous findings. This method can be further refined and adapted for other disease contexts and animal models in future studies. By

integrating this approach with various tool mouse models, it offers the potential to investigate the transport dynamics of diverse organelles or proteins beyond mitochondria in different cellular environments.

Data availability

Enquiries about data availability should be directed to the authors.

Acknowledgements

This work was supported by Thomas Jefferson University, Peking University School of Life Sciences and The University of Georgia (UGA). The authors wish to thank Dr. Ying-Chun Hu, Zhen-Yang Kong, and Hong-Zhang Zhou for their expert technical assistance with EM sample preparation and image analysis at the Core Facilities of the School of Life Sciences, Peking University. Additionally, the authors thank Dr. D. James Surmeier at Northwestern University for providing the AAV-TH-GCamp6 virus and TH-Mito-roGFP mice, Dr. Jerry Cheng (New York Institute of Technology) for his valuable suggestions on the statistical analysis, and Dr. Shiquan Cui at UGA for his feedback on the manuscript.

Funding

This study was supported by fundings from Hereditary Disease Foundation (CJL), Peking University School of Life Sciences (CJL), Thomas Jefferson University startup funds (HZ), UGA startup funds (HZ), and Parkinson's Disease and Alzheimer's Disease Innovative Research Fund (H.Z.).

Competing interests

The authors report no competing interests.

Supplementary material

Supplementary material is available at *Brain* online.

References

1. Poewe W, Seppi K, Tanner CM, et al. Parkinson disease. *Nat Rev Dis Prim*. 2017;3(1):1-21.

2. Malpartida AB, Williamson M, Narendra DP, Wade-Martins R, Ryan BJ. Mitochondrial Dysfunction and Mitophagy in Parkinson's Disease: From Mechanism to Therapy. *Trends Biochem Sci*. 2020;0(0):1-15.

3. Winklhofer KF, Haass C. Mitochondrial dysfunction in Parkinson's disease. *Biochim Biophys Acta - Mol Basis Dis*. 2010;1802(1):29-44.

4. Pryde KR, Smith HL, Chau KY, Schapira AHV. PINK1 disables the anti-fission machinery to segregate damaged mitochondria for mitophagy. *J Cell Biol*. 2016;213(2):163-171.

5. Exner N, Lutz AK, Haass C, Winklhofer KF. Mitochondrial dysfunction in Parkinson's disease: Molecular mechanisms and pathophysiological consequences. *EMBO J*. 2012;31(14):3038-3062.

6. Vilain S, Esposito G, Haddad D, et al. The Yeast Complex I Equivalent NADH Dehydrogenase Rescues pink1 Mutants. Larsson N-G, ed. *PLoS Genet*. 2012;8(1):e1002456.

7. Park JS, Davis RL, Sue CM. Mitochondrial Dysfunction in Parkinson's Disease: New Mechanistic Insights and Therapeutic Perspectives. *Curr Neurol Neurosci Rep*. 2018;18(5).

8. Kostic M, Ludtmann MHR, Bading H, et al. PKA Phosphorylation of NCLX Reverses Mitochondrial Calcium Overload and Depolarization, Promoting Survival of PINK1-Deficient Dopaminergic Neurons. *Cell Rep*. 2015;13(2):376-386.

9. Weihofen A, Thomas KJ, Ostaszewski BL, Cookson MR, Selkoe DJ. Pink1 forms a multiprotein complex with miro and milton, linking Pink1 function to mitochondrial trafficking. *Biochemistry*. 2009;48(9):2045-2052.

10. Zhu XH, Qiao H, Du F, et al. Quantitative imaging of energy expenditure in human brain. *Neuroimage*. 2012;60(4):2107-2117.

11. Wang X, Winter D, Ashrafi G, et al. PINK1 and Parkin target miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell*.

2011;147(4):893-906.

12. Das Banerjee T, Dagda RKRY, Dagda M, et al. PINK1 regulates mitochondrial trafficking in dendrites of cortical neurons through mitochondrial PKA. *J Neurochem*. 2017;142(4):545-559.

13. Sheng ZH. Mitochondrial trafficking and anchoring in neurons: New insight and implications. *J Cell Biol*. 2014;204(7):1087-1098.

14. Zhi L, Qin Q, Muqeem T, et al. Loss of PINK1 causes age-dependent decrease of dopamine release and mitochondrial dysfunction HHS Public Access. *Neurobiol Aging*. 2019;75:1-10.

15. Wang X, Su B, Siedlak SL, et al. Amyloid-beta overproduction causes abnormal mitochondrial dynamics via differential modulation of mitochondrial fission/fusion proteins. *Proc Natl Acad Sci U S A*. 2008;105(49):19318-19323.

16. Gong S, Yang XW, Li C, Heintz N. Highly Efficient Modification of Bacterial Artificial Chromosomes (BACs) Using Novel Shuttle Vectors Containing the R6Kγ Origin of Replication. *Genome Res*. 2002;12(12):1992-1998.

17. Zampese E, Wokosin DL, Gonzalez-Rodriguez P, et al. Ca2+ channels couple spiking to mitochondrial metabolism in substantia nigra dopaminergic neurons. *Sci Adv*. 2022;8(39):8701.

18. Mangeol P, Prevo B, Peterman EJG. KymographClear and KymographDirect: Two tools for the automated quantitative analysis of molecular and cellular dynamics using kymographs. *Mol Biol Cell*. 2016;27(12):1948-1957.

19. Schwarz TL. Mitochondrial trafficking in neurons. *Cold Spring Harb Perspect Biol*. 2013;5(6).

20. Liu W, Acín-Peréz R, Geghman KD, Manfredi G, Lu B, Li C. Pink1 regulates the oxidative phosphorylation machinery via mitochondrial fission. *Proc Natl Acad Sci U S A*. 2011;108(31):12920-12924.

21. Wang X, Schwarz TL. The Mechanism of Ca2+-Dependent Regulation of Kinesin-Mediated Mitochondrial Motility. *Cell*. 2009;136(1):163-174.

22. Giorgi C, Marchi S, Pinton P. The machineries, regulation and cellular functions of mitochondrial calcium. *Nat Rev Mol Cell Biol 2018 1911*. 2018;19(11):713-730.

doi:10.1038/s41580-018-0052-8

23. Ruiz A, Alberdi E, Matute C. CGP37157, an inhibitor of the mitochondrial Na+/Ca2+ exchanger, protects neurons from excitotoxicity by blocking voltage-gated Ca2+ channels. 2014;5(4):e1156-e1156.

24. Santulli G, Xie W, Reiken SR, Marks AR. Mitochondrial calcium overload is a key determinant in heart failure. *Proc Natl Acad Sci U S A*. 2015;112(36):11389-11394.

25. Peng TI, Jou MJ. Oxidative stress caused by mitochondrial calcium overload. *Ann N Y Acad Sci*. 2010;1201:183-188.

26. Msackyi M, Chen Y, Tsering W, Wang N, Zhang H. Dopamine Release Neuroenergetics in Mouse Striatal Slices. *Int J Mol Sci*. 2024;25(9).

27. Debattisti V, Gerencser AA, Saotome M, Das S, Hajnóczky G. ROS Control Mitochondrial Motility through p38 and the Motor Adaptor Miro/Trak. *Cell Rep*. 2017;21(6):1667-1680.

28. Candas D, Li JJ. MnSOD in Oxidative Stress Response-Potential Regulation via Mitochondrial Protein Influx. *Antioxid Redox Signal*. 2014;20(10):1599.

29. Padzik A, Deshpande P, Hollos P, et al. KIF5C S176 Phosphorylation Regulates Microtubule Binding and Transport Efficiency in Mammalian Neurons. *Front Cell Neurosci*. 2016;10(MAR2016).

30. Brady ST, Morfini GA. Regulation of motor proteins, axonal transport deficits and adult-onset neurodegenerative diseases. *Neurobiol Dis*. 2017;105:273-282.

31. Morfini GA, Bosco DA, Brown H, et al. Inhibition of Fast Axonal Transport by Pathogenic SOD1 Involves Activation of p38 MAP Kinase. *PLoS One*. 2013;8(6):e65235.

32. Berth SH, Lloyd TE. Disruption of axonal transport in neurodegeneration. *J Clin Invest*. 2023;133(11).

33. Liu S, Sawada T, Lee S, et al. Parkinson's Disease–Associated Kinase PINK1 Regulates Miro Protein Level and Axonal Transport of Mitochondria. Copenhaver GP, ed. *PLoS Genet*. 2012;8(3):e1002537.

Figure legends

Figure 1 Generation and characterization of TH-Mito-Dendra2 (MD) BAC transgenic mouse. **(A)** Depiction of the TH-MitoDendra2 BAC construct. To modify RPCI23-350E13 BAC, a targeting cassette was generated with A arm, B arm and mitochondrial-targeting dendra2 insert. The mito-dentra2 and Th gene exon1 share the same initiation ATG codon. A and B represent the homologous recombination regions. MTS: mitochondrial targeting sequence. **(B)** A coronal section of TH-Mito-Dendra2 (MD) BAC transgenic mouse brain showing the histological overview of Mito-Dentra2 expression pattern. Tyrosine hydroxylase (TH) signal (red) co-localizes with Mito-Dentra2 fluorescence (green) in midbrain areas. **(C)** Mito-Dendra2 (green) expressing cells are all TH (red) immunoreactive in the substantia nigra (SN). **(D)** Mito-Dendra2 (green) expressing cells are all TH (red) immunoreactive in the ventral tegmental area (VTA). **(E)** A sagittal section of TH-MitoDendra2 BAC mouse brain showing continuous and intact nigrostriatal axonal projection of dopaminergic neurons. **(F)** MitoDendra2 is localized in mitochondria, as labeled with MitoTracker, in the striatum of TH-mitodendra2-BAC transgenic mice.

Figure 2 Investigation of mitochondrial movement in acute brain slices using two-photon laser microscopy (2PLSM). **(A)** Schematic illustration of mitochondrial trafficking imaging in acute brain slices. **(B)** Image sequences of mitochondrial trafficking. B1-B6 represent images of mitochondria moving at the different time points: 0, 5, 10, 15, 20 and 25 seconds, respectively. White circles 1 and 2 denote the anterograde movement of mitochondria, blue circle 3 denotes the retrograde movement of mitochondria. **(C)** Kymograph generation and mitochondria trafficking measurement. C1: Representative image of mitochondria expressing Dendra2 protein. C2: Maximum intensity projection of a series of mitochondrial trafficking images. C3: Pseudo color map of panel C2 to facilitate outlining the trace of mitochondrial movement. C4: Kymograph map generated from the images, showing anterograde, retrograde, and static traces. Red lines indicate the anterograde trace of mitochondrial movement, green lines indicate the retrograde movement, and blue lines represent static mitochondria.

Figure 3 Age-dependent mitochondrial axonal transport deficits in *Pink1*^{-/-} mice. (A) Representative in vivo two-photon microscopy images of mitochondrial trafficking in dopaminergic neuron axons at the initial time point. Images depict *Pink1*$^{+/+}$ and *Pink1*$^{-/-}$ mice at 2-4 months (A1-A2), 5-7 months (A3-A4), and above 10 months (A5-A6) of age, respectively. **(B)** Representative kymographs illustrating mitochondrial trafficking patterns in *Pink1*$^{+/+}$ and *Pink1*$^{-/-}$ mice across different age groups. **(C)** Quantification of mitochondrial motility states in 2-4-month-old *Pink1*$^{+/+}$ (N = 3, n = 8) and *Pink1*$^{-/-}$ (N = 3, n = 12) mice. Data points represent mean percentages per brain slice, with colors denoting individual animals. *P*-values were determined using the unpaired t-test. **(D)** SuperPlots illustrating anterograde (left) and retrograde (right) mitochondrial velocities in 2-4-month-old mice. In scatter plots, each data point represents an individual mitochondrial velocity measurement. In summary plot, each data point represents the mean velocity per animal, with colors distinguishing individual mice. *P*-values were determined using the Mann-Whitney U test. **(E)** Quantification of mitochondrial motility states in 5-7-month-old *Pink1*$^{+/+}$ (N = 3, n = 10) and *Pink1*$^{-/-}$ (N = 5, n = 12) mice. *P*-values were determined using the unpaired t-test. ****P < 0.01, *****P < 0.0001. **(F)** SuperPlots illustrating anterograde (left) and retrograde (right) mitochondrial velocities in 5-7-month-old mice. *P*-values were determined using the Mann-Whitney U test. *P < 0.05. **(G)** Quantification of mitochondrial motility states in older than 10-month-old *Pink1*$^{+/+}$ (N = 3, n = 5) and *Pink1*$^{-/-}$ (N = 4, n = 9) mice. *P*-values were determined using the unpaired t-test. *P < 0.05, ** P < 0.01. **(H)** SuperPlots illustrating anterograde (left) and retrograde (right) mitochondrial velocities in mice older than 10 months. *P*-values were determined using the Mann-Whitney U test. ***P < 0.001.

Figure 4 Mitochondrial calcium elevation and transport deficits in acute brain slices. (A) Representative fluorescence images of TH-mito-GCaMP6 in acute brain slices from approximately 3-month-old *Pink1*$^{+/+}$ (upper panel) and *Pink1*$^{-/-}$ mice (lower panel). Images depict mitochondrial $[Ca^{2+}]$ under baseline conditions (left column), after 20 minutes of Ca^{2+}-free aCSF perfusion (500 µM EGTA; middle column) and following 30 minutes of 3 mM Ca^{2+} aCSF perfusion (right column). **(B)** Quantification of mitochondrial $[Ca^{2+}]$ in *Pink1*$^{+/+}$ (N = 3, n = 8) and *Pink1*$^{-/-}$ (N = 3, n = 12) mice. Data points represent mean fluorescence intensity per brain slice, with colors denoting individual animals. Statistical significance was determined using the Mann-Whitney U test. ***P = 0.0001. **(C)** Representative images and

corresponding kymographs illustrating mitochondrial dynamics before (C1 and C3) and after (C2 and C4) 2-hour incubation with 5 µM CGP37157. (D) Quantification of mitochondrial motility states following 5 µM CGP37157 treatment. Ctrl: N = 3, n = 6; CGP37157: N = 3, n = 7. P-values were determined using the unpaired t-test. *P < 0.05, **P < 0.01. (E) SuperPlots of anterograde (left) and retrograde (right) mitochondrial velocities in control and CGP37157-treated groups. P-values were determined using the Mann-Whitney U test.

Figure 5 NCLX inhibition-induced mitochondrial Ca^{2+} overload leading to enhanced oxidative levels. (A-B) Representative images of SN dopaminergic neurons expressing roEGFP. Neurons were imaged at excitation wavelengths of 900 nm (reduced state, A1 and B1) and 800 nm (oxidized state, A2 and B2) under two conditions: (A) regular aCSF, (B) aCSF supplemented with 10 µM CGP37157. **(C)** Quantification of oxidative stress levels following 60-minute treatment with 10 µM CGP37157. Each data point represents the mean ROS level of an individual neuron, with colors denoting neurons from distinct brain slices. P-value was determined using the unpaired t-test. ***P < 0.001. Ctrl: n = 20; CGP37157: n = 19, where n represents the number of individual neurons analyzed.

Figure 6 Reduced mitochondrial anterograde movement by ROS elevation. (A) Representative images depicting mitochondrial distribution under various treatment conditions at the initial imaging point: (A1) aCSF, (A2) 0.2% DMSO in aCSF, and (A3) 50 µM menadione (MED) in aCSF. **(B)** Kymographs illustrating mitochondrial trafficking patterns following different treatments: (B1) aCSF, (B2) 0.2% DMSO in aCSF, and (B3) 50 µM menadione (MED) in aCSF. **(C)** Quantitative analysis of mitochondrial motility under different treatment conditions. P-values were determined using the unpaired t-test. *P < 0.05, ** P < 0.01. Regular aCSF control: N = 3, n = 8; DMSO control: N = 3, n = 6; menadione: N = 3, n = 4. **(D)** Effect of menadione treatment on mitochondrial velocity in anterograde (left) and retrograde (right) directions. **(E)** Representative immunoblots showing MnSOD expression in the striatum (left) and substantia nigra (right) of 5-7-month-old $Pink1^{+/+}$ and $Pink1^{-/-}$ mice. **(F)** Quantification of MnSOD expression levels in striatum (left) and substantia nigra (right) of 5-7-month-old $Pink1^{+/+}$ and $Pink1^{-/-}$ mice. Each color represents an individual animal. P-values were determined using the unpaired t-test. *P < 0.05, **P < 0.01. **(G)** Immunofluorescence images of TH-positive dopaminergic neurons in the substantia nigra of

5-7-month-old *Pink1$^{+/+}$* (G1) and *Pink1$^{-/-}$* (G2) mice. **(H)** Immunofluorescence images of MnSOD in the substantia nigra of 5-7-month-old *Pink1$^{+/+}$* (H1) and *Pink1$^{-/-}$* (H2) mice. **(I)** Quantification of relative MnSOD fluorescence intensity in dopaminergic neurons of 5-7-month-old *Pink1$^{+/+}$* and *Pink1$^{-/-}$* mice. Statistical significance was determined using the Mann-Whitney U test. In scatter plots, each data point represents an individual brain slice. In summary plot, each data point represents the mean fluorescence intensity per animal, with colors distinguishing individual mice. ****$P < 0.0001$.

Figure 7 p38 MAPK pathway hyperactivation by loss of *Pink1* and improved mitochondrial motility in *Pink1$^{-/-}$* mice by p38 inhibition. (A) Representative immunoblots showing phospho-p38 and total p38 expression levels in the striatum (left) and substantia nigra (right) of 6-7-month-old *Pink1$^{+/+}$* and *Pink1$^{-/-}$* mice. **(B)** Quantification of phospho-p38 and total p38 expression levels in striatum (left) and substantia nigra (right). Each data point represents an individual animal. *P*-values were calculated using Mann-Whitney U-test. **$P < 0.01$, ****$P < 0.0001$. **(C)** Representative images depicting mitochondrial distribution under different treatment conditions at the initial imaging point: (C1) aCSF, (C2) 50 μM p38 inhibitor (SB202190) in aCSF. **(D)** Kymographs illustrating mitochondrial trafficking patterns: (D1) aCSF, (D2) 50 μM SB202190 treatment. **(E)** Percentage quantification of mitochondrial movement status after SB202190 incubation. *P*-values were determined using the unpaired t-test. **$P < 0.01$, ****$P < 0.0001$. Regular aCSF control: N = 3, n = 6; SB202190: N = 3, n = 7. **(F)** Impact of SB202190 treatment on mitochondrial velocity in anterograde (left) and retrograde (right) directions. *P*-values were calculated using the Mann-Whitney U-test.

Figure 8 Defective mitochondria at dopamine neuron terminals in *Pink1$^{-/-}$* mice. (A-B) TH immunogold electron micrographs illustrating mitochondrial morphology in dopaminergic terminals of (A) *Pink1$^{+/+}$* and (B) *Pink1$^{-/-}$* mice. Blue arrows indicate TH-positive signals in dopamine terminals. Red arrows denote mitochondria within or proximal to dopamine terminals. **(C)** Quantitative analysis of mitochondrial status in dopaminergic terminals and surrounding striatal regions of *Pink1$^{+/+}$* and *Pink1$^{-/-}$* mice. Each data point represents an individual mouse. Statistical significance was determined using the Mann-Whitney U test. **$P < 0.01$, N = 5. **(D)** Quantification of mitochondrial circularity and aspect

ratio in dopaminergic neuron terminals of *Pink1*$^{+/+}$ and *Pink1*$^{-/-}$ mice. In scatter plots, each data point represents individual mitochondria. In summary plot, each data point represents the mean value per animal, with colors distinguishing individual mice. *P*-values were calculated using Mann-Whitney U test. ****$P < 0.001$, N = 5. Morphometric analysis of mitochondrial circularity and aspect ratio in dopaminergic neuron terminals of *Pink1*$^{+/+}$ and *Pink1*$^{-/-}$ mice. In scatter plots, each data point represents an individual mitochondrion. In summary plot, each data point represents the mean value per animal, with colors distinguishing individual mice. Statistical analysis was performed using the Mann-Whitney U test. ****$P < 0.001$, N = 5. **(E)** Proposed mechanistic model: *Pink1* gene deletion leads to aberrant activation of the p38 pathway, resulting in impaired mitochondrial axonal transport in dopaminergic neurons.

Zhao J *et al.* Figure 2.

Zhao J *et al.* Figure 5.

Zhao J *et al.* Figure 6.

Zhao J *et al.* Figure 7.