MONOPTEROS isoform *MP11ir* role during somatic embryogenesis in *Arabidopsis thaliana*

Barbara Wójcikowska[1,2 # *,] Samia Belaidi[1,3#], Victoria Mironova[4], Helene Robert Boisivon[1*]

[1] Hormonal Crosstalk in Plant Development, Mendel Centre for Plant Genomics and Proteomics,

CEITEC MU - Central European Institute of Technology, Masaryk University, Brno, Czech

Republic

[2] Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences,

University of Silesia in Katowice, Katowice, Poland

[3] Laboratory of Functional Genomics and Proteomics, National Centre for Biomolecular Research,

Faculty of Science, Masaryk University, Brno, Czechia

[4] Department of Plant and Animal Biology, Radboud Institute for Biological and Environmental

Sciences (RIBES), Radboud University, The Netherlands

# These authors contributed equally to this work.

* Authors to whom correspondence should be addressed (barbara.wojcikowska@us.edu.pl and

helene.robert.boisivon@ceitec.muni.cz).

ORCID

BW 0000-0003-0287-9859

SB 0009-0006-4574-4005

VM 0000-0003-3438-0147

HSR 0000-0002-6610-836X

Running title

MP/ARF5 and MP11ir required for somatic embryogenesis

8 Main Figures

1 Supplementary Table and 8 Supplementary Figures

7,502 words

Abstract

Auxin is crucial for plant morphogenesis, including *in vivo* and *in vitro* embryo development. Exogenous auxin application is necessary for inducing embryogenic responses in *in vitro* cultured explants of Arabidopsis and other plants. Thus, components of auxin transport, signaling, and metabolism are key to somatic embryo formation. AUXIN RESPONSE FACTOR (ARF) transcription factors, which bind to auxin response elements to control the auxin-responsive gene expression, are vital in somatic embryo regeneration. ARFs are often repressed by AUXIN/INDOLE-3-ACETIC ACIDs (Aux/IAAs). MONOPTEROS (MP)/ARF5 is especially important in the embryogenic transition, being highly expressed during somatic embryogenesis; its mutant cannot develop somatic embryos. The *MP11ir* transcript, an alternatively spliced variant of *MP/ARF5*, produces a truncated protein missing the Phox and Bem1p (PB1) domain, crucial for ARF-Aux/IAA dimerization. This makes the MP11ir isoform insensitive to Aux/IAA repression, suggesting auxin-independent regulation. High levels of *MP11ir* transcript are observed during auxin- and trichostatin A-dependent induction of somatic embryogenesis. Both MP/ARF5 and MP11ir are essential for embryo regeneration in the *mpS319* mutant. However, overexpressing truncated MP/ARF5 protein (ΔARF5) lacking the PB1 domain inhibits somatic embryogenesis, leading to callus formation instead of somatic embryos. Overexpressing *ΔARF5*, lacking MP/ARF5 protein (*mp/arf5* mutant), and blocking of MP/ARF5 action with auxin-resistant BODENLOS (BDL) protein affect the expression of genes involved in auxin biosynthesis, like *TRYPTOPHAN AMINOTRANSFERASE OF ARABIDOPSIS 1* (*TAA1*), *TAA1-RELATED 1* (*TAR1*), *YUCCA3* (*YUC3*), *YUC5* and *YUC8*, which may be potential targets of MP11ir and/or MP/ARF5. Consequently, *ΔARF5* overexpression alters auxin homeostasis and endogenous auxin levels, hindering embryogenic transition.

Keywords: Alternative splicing; *Arabidopsis thaliana*; Auxin; AUXIN RESPONSE FACTOR; Auxin signalling; *In vitro* culture; Isoform; MONOPTEROS; MP11ir; Somatic embryogenesis; Transcription factor

Introduction

Auxin plays a vital role in various aspects of plant development, including plant growth and morphogenesis in response to environmental signals through its transport, metabolism, and

signaling (Caumon and Vernoux, 2023; Cohen and Strader, 2024; Liu et al., 2024). Recognized as

a key trigger of developmental changes, auxin is a critical inducer of somatic embryogenesis (SE)

in many plant species (Elhiti and Stasolla, 2022). SE is a unique plant-specific developmental

process that demonstrates the pluripotency of somatic cells and is widely exploited for *in vitro* plant

regeneration. This technique is used in biotechnology for mass micropropagation of plants

(Mehbub et al., 2022), plant biodiversity conservation (Ballesteros et al., 2024), and the production

of transgenic plants (Lee and Wang, 2023; McFarland and Kaeppler, 2024; Yang et al., 2024).

With the rising global population, decreased crop yields due to climate change, and increasing

environmental pollution, the importance of *in vitro* plant propagation techniques is growing

(Hasnain et al., 2022; Krasteva et al., 2021). Understanding the molecular mechanisms of SE can

enhance its efficiency and its optimization for recalcitrant species (Chen et al., 2022). Exogenous

auxin treatment of explants modifies chromatin accessibility and the transcriptome of somatic cells

(Wang et al., 2020; Wu et al., 2022). This modulation affects many genes, including those encoding

transcription factors (TFs), proteins associated with hormone transport, metabolism, signaling, and

stress response. The induction of SE is also associated with the expression of the genes encoding

the core components of the nuclear auxin signaling pathway, such as AUXIN RESPONSE

FACTORs (ARFs) and AUXIN/INDOLE-3-ACETIC ACIDs (Aux/IAAs) (Quintana-Escobar et

al., 2019). Thus, the crosstalk between auxin and these transcriptional regulators is fundamental to

the SE regulatory network.

MONOPTEROS (MP), known as AUXIN RESPONSE FACTOR 5 (ARF5), is pivotal in nuclear

auxin signaling and has been identified in many plant species (Chandler, 2016). The function of

MP/ARF5 is well-characterized, thanks to a wide range of tools like allelic mutants, overexpressing

and reporter lines in *Arabidopsis thaliana,* allowing for comprehensive functional, genetic, and

microscopic analyses (Hardtke et al., 2004; Odat et al., 2014; Rademacher et al., 2011; Schlereth

et al., 2010; Wójcikowska et al., 2023; Wu et al., 2015). Knowledge of the MP/ARF5 action is

being transferred to commercially important species (Liu et al., 2018a; Xu et al., 2019; Yue et al.,

2020). MP/ARF5 is crucial for many aspects of *in vivo* plant development, including flower, ovule,

and pollen development (Cucinotta et al., 2021; Liu et al., 2018b; Wu et al., 2015), zygotic

embryogenesis (Möller et al., 2017), root and shoot meristem establishment (Cole et al., 2009;

Krogan et al., 2016; Schlereth et al., 2010), organ polarity maintenance (Bhatia et al., 2016), and

vascularization patterns (Mattsson et al., 2003; Przemeck et al., 1996). MP/ARF5 is also critical

for *in vitro* plant morphogenesis. *MP/ARF5* is essential for an effective SE process because its expression is stimulated during SE; it is active in regions (cotyledons and SAM) where somatic embryos emerge; and, more importantly, explants of *mp/arf5* mutant cannot produce somatic embryos (Wójcikowska and Gaj, 2017).

The MP/ARF5 protein has four distinct domains: a B3-type DNA binding domain (DBD), dimerization domains (DD1, DD2), a middle region (MR) region, and a Phox and Bem1 (PB1) domain (Guilfoyle, 2015). MP/ARF5 binds to the AuxRE *cis*-elements in the promoter of target genes. As a transcription factor, MP/ARF5 primarily acts as a positive regulator of gene expression (Boer et al., 2014; Freire-Rios et al., 2020). In Arabidopsis, MP/ARF5 binds as a dimer to AuxRE motifs organized as inverted repeats (IR), direct repeats (DR), and everted repeats (ER). IR consists of two AuxREs on opposite DNA strands facing each other, DR has two consecutive AuxREs on the same DNA strand, and ER features two AuxREs on opposite directions and DNA strands. The DD1 and DD2 domains flank the DBD, acting as a caliper for binding to IR motifs. For docking on ER and DR motifs, the MP/ARF5-DBD dimer relies on the MR or PB1 domains (Cancé et al., 2022). Additionally, AuxRE elements in the MP/ARF5 promotor indicate that its expression is regulated by cellular auxin level (Lau et al., 2011).

Three scenarios outline MP/ARF5 action. First, at low auxin levels, MP/ARF5 interacts with BODENLOS/INDOLE-3-ACETIC ACID 12 (BDL/IAA12) via PB1 domains, recruiting TOPLESS/TOPLESS RELATED (TPL/TPR), transcriptional co-repressor proteins, and HDA19 histone deacetylase enzyme. HDA19 removes the acetyl group from the histone H3 and H4 tails, condensing the chromatin and blocking the expression of auxin-responsive genes (Chandler, 2016). In the presence of auxin, BDL is recruited by the SCF[TIR/AFBs] complex, ubiquitinated, and degraded by the proteasome, releasing MP/ARF5 to form a complex with SWI/SNF chromatin remodeling ATPases BRAHMA (BRM) and SPLAYED (SYD). It unlocks the chromatin to provide histone acetyltransferases (HATs) and transcription factors access to the *cis*-elements of MP/ARF5-regulated promoters, thus regulating gene transcription (Wu et al., 2015). Second, above-threshold auxin levels may potentially cause MP/ARF5 protein oligomerization with MP/ARF5 or other ARFs, inhibiting the expression of auxin-responsive genes, as observed *in vitro* (Korasick et al., 2014; Nanao et al., 2014) and *in vivo* for ARF19 (Powers et al., 2019). This scenario is plausible as plants with *MP/ARF5* constitutive overexpression (Hardtke et al., 2004) phenocopy *mp/arf5* mutants (Przemeck et al., 1996). Third, the MP11ir isoform, produced by alternative splicing (AS)

of the *MP/ARF5* transcript during ovule development, lacks the PB1 domain due to the retention of intron 11, resulting in the translation of a truncated protein. This isoform is insensitive to Aux/IAA repression and could function independently of auxin. Ectopic expression of *MP11ir* partially complements *mp/arf5* mutant phenotypes during reproductive development, indicating that some MP/ARF5 functions do not require interaction with Aux/IAA repressors. MP11ir activates target genes in ovular regions with low auxin concentrations (Cucinotta et al., 2021).

To understand the role of MP11ir in the embryogenic transition, we analyzed its transcript level in Arabidopsis explants cultured *in vitro* and treated with SE inducers, such as auxins 2,4-dichlorophenoxyacetic acid (2,4-D), 1-naphthaleneacetic acid (NAA), and indole-3-acetic acid (IAA), and trichostatin A (TSA). High *MP11ir* transcript levels were associated with auxin- and TSA-dependent SE inductions. MP11ir can partially rescue the *mp/arf5* phenotype, improving its capacity for embryogenic transition. However, both MP/ARF5 and MP11ir are required to fully complement the *mp/arf5* phenotype. Overexpression of the truncated MP/ARF5 protein (ΔARF5, mimicking MP11ir) inhibits SE and alters the expression of auxin biosynthetic genes of the indole-3-pyruvic acid (IPyA) pathway: *TRYPTOPHAN AMINOTRANSFERASE OF ARABIDOPSIS 1* (*TAA1*), *TAA1-RELATED 1* (*TAR1*), *YUCCA 3* (*YUC3*), *YUC5* and *YUC8*. This modulation changes endogenous auxin levels, negatively impacting embryogenic transition. We hypothesized that *TAA1*, *YUC3,* and *YUC8* may be directly controlled by MP11ir. These findings suggest that MP11ir is involved in SE induction alongside the canonical MP/ARF5, contributing to the regulation of auxin biosynthesis and homeostasis during this process.

Results

The presence of *MP11ir* transcript is specific for auxin-dependent embryogenic transition

MP11ir was first detected in ovules, where it likely regulates target genes in cells with low auxin concentrations (Cucinotta et al., 2021). Given the critical role of auxin signaling proteins in the embryogenic transitions, we investigated whether *MP11ir* would be active during SE induced by exogenously applied synthetic auxin 2,4-D. We hypothesized that if *MP11ir* regulates its target genes in cells with low auxin concentrations, its presence would be minimal in cells undergoing embryogenic reprogramming, as those cells accumulate exogenous 2,4-D and endogenous IAA (Schulz and Segobye, 2016; Wójcikowska and Gaj, 2015). To test this, we analyzed *MP11ir* transcript levels in various samples: 10-day-old seedlings, flowers, leaves, immature zygotic

embryos (IZEs), and IZE explants during auxin-dependent SE induction. *MP11ir* was not detected in leaves but was present in other tissues, notably flowers, as previously observed (Fig. S1) (Cucinotta et al., 2021). Quantitative analysis revealed that *MP11ir* was most abundant during the auxin-dependent embryogenic transition, with 11-18 times higher expression than in seedlings (Fig. 1A). It suggests that *MP11ir* may regulate target gene expression in both low (Cucinotta et al., 2021) and high auxin concentrations (our results), indicating a role in auxin-independent control transcription during SE.

The presence of the *MP11ir* transcript depends on the type and concentration of exogenously applied auxin

To investigate the correlation between *MP11ir* transcript levels and auxin presence, we treated IZEs with various auxins, i.e., 2,4-D, NAA, and IAA, at increasing concentrations (0, 2.5, 5.0, and 7.5 µM) for ten days. Different auxins at the same concentrations had varied effects on *MP11ir* levels. We observed a positive correlation between *MP11ir* levels and increasing concentrations of 2,4-D (Spearman correlation coefficient 0.87; $p < 0.01$), NAA (Spearman correlation coefficient 0.92; $p < 0.01$), but not IAA (Fig. 1 B-D). The application of 2,4-D had the most substantial promoting effect on *MP11ir* levels. A 2.5 µM concentration of 2,4-D increased *MP11ir* transcript levels tenfold compared to untreated explants, while a 7.5 µM concentration of 2,4-D (three times higher) resulted in an 18-fold increase (Fig. 1B, E-H). The highest concentration of NAA (7.5 µM) only increased *MP11ir* levels sevenfold (Fig. 1C, I-L). IAA did not significantly affect MP11ir transcript levels (Fig. 1D, M-P). The strength of SE induction also varied with the type of applied auxins (Fig. S2). In Arabidopsis IZE culture, 2,4-D is the most effective for SE induction (Fig. S2A, D-G), followed by NAA (Fig. S2B, H-K), with IAA being the least effective (Fig. S2C, L-O). We found a positive correlation between the increase of *MP11ir* transcript levels and the strength of SE induction for 2,4-D (Spearman correlation coefficient 0.80; $p < 0,01$) and NAA (Spearman correlation coefficient 0.75; $p < 0,01$) but not for IAA. This data suggests that the type and concentration of auxin influence *MP11ir* levels and subsequently affect the efficiency of the SE process.

The presence of *MP11ir* transcript is specific for the embryogenic transition independently of the presence of auxin

Synthetic auxin 2,4-D is the most common and effective inducer of SE in Arabidopsis and other plants (Wójcik et al., 2020). To determine whether MP11ir is specific for embryogenic transition rather than merely responding to the application of exogenous auxins, which may also act as stressors, we assessed the impact of trichostatin A (TSA), an inhibitor of histone deacetylases (HDAC) known to induce SE, on the levels of *MP11ir* transcript. IZE explants treated with 1 µM TSA rapidly and massively regenerate somatic embryos (Wójcikowska et al., 2018). We measured *MP11ir* levels on the 5th and 10th days of TSA-dependent SE induction and found significantly elevated levels of *MP11ir*, 55 and 84 times higher than in seedlings, respectively (Fig. 2Q). These findings suggest that MP11ir is specifically associated with the developmental switch from somatic to embryonic cell fate, highlighting its role in the embryonic transition (Fig. 2R).

The *MP/ARF5* transcript is detected during the SE process

The transcript coding for the canonical MP/ARF5 protein was also detectable during the SE process, consistent with current knowledge (Wójcikowska and Gaj, 2017). Its expression pattern was highly similar to *MP11ir*. The *MP/ARF5* transcript significantly accumulated in 2,4-D and NAA-treated IZE explants (all tested concentrations) and in IAA-treated explants under 5.0 µM concertation (Fig. S3A-C). We observed a significant up-regulation of *MP/ARF5* expression during auxin-dependent (Fig. S3D) and TSA-dependent (Fig. S3E) induction of the SE process. During the SE process induced by auxin- and TSA, the *MP/ARF5* transcript is predominantly synthesized over *MP11ir* (Fig. S3F, G). In the auxin-induced embryogenic transition, the *MP/ARF5* transcript was more than 100 times higher than *MP11ir*, while in TSA-induced SE, it was nearly 18 times higher (Fig. S3F). These results indicate that both *MP/ARF5* and *MP11ir* transcripts are synthesized during somatic embryo formation, denoting their involvement in embryogenic transition.

Both MP/ARF5 and MP11ir are necessary to fully rescue the SE-related phenotypes of the *mpS319* mutant

Mutations in the *MP/ARF5* gene dramatically reduce the capacity of IZE explants for SE. The *mpS319* mutant showed impaired SE efficiency (20%) and productivity (2 somatic embryos per explant) compared to Col-0 (85% efficiency and 4 embryos per explant, respectively) (Fig. 2A-D, H-J). Mutations in *MP/ARF5* affect zygotic embryo development, potentially impacting the

capacity of the IZE to undergo SE. To address this, we inhibited the MP/ARF5-dependent signaling

pathway by expressing an auxin-insensitive BDL/IAA12 (bdl) protein in IZE. Overexpression of

DEX-induced bdl in IZE completely shut down the embryogenic transition. The explants of the

35S::bdl-GR line treated with DEX did not respond to the 2,4-D SE-inducing agent and developed

into seedlings. In contrast, Col-0 and the *35S::BDL-GR* transgenic line expressing a native BDL

protein, which is ubiquitinated and degraded by the proteasome in the presence of auxin,

regenerated somatic embryo massively and independently of the presence of DEX (Fig. 2N-X).

These results confirm the findings obtained with the *mpS319* line.

It was previously shown that MP11ir could partially rescue the phenotype of the *mpS319* mutant

during ovule development (Cucinotta et al., 2021). To investigate the function of MP11ir and its

involvement in the SE process, we performed a complementation test by expressing *MP11ir* or/and

MP/ARF5 under the control of the *MP/ARF5* promoter (*MP::MP11ir* and *MP::MP* transgenes) in

the *mpS319* mutant. The presence of MP11ir improved the SE capability of the *mpS319* mutant by

2.3 times (Fig. 2A-B, E, K). Interestingly, the canonical MP/ARF5 also did not fully rescue the

mpS319 SE defects, with 48% of explants responding to the SE induction in *MP::MP mpS319*

explants (Fig. 2A-B, F, L). It appears that both *MP/ARF5* and *MP11ir* transcripts must be present

to fully rescue the *mpS319* phenotype (Fig. 2A-B, G, M). The complementation test showed that

both MP11ir and MP/ARF5 are required to fully rescue the *mpS319* SE defects, and both the

canonical and truncated MP/ARF5 proteins are necessary for effective somatic embryo formation.

MP11ir localization during the SE process

To detect MP11ir protein during the auxin-dependent SE process, we used the transgenic line

pMP::MP11ir-GFP (MR). The MP11ir was detected in various stages and regions during SE

induction. MP11ir was present in freshly isolated IZEs in the cotyledonary stage of development,

primarily localized in the vasculature (Fig. 3A). After 24 hours of SE induction, MP11ir was

observed on the adaxial side of the cotyledon as well as near the shoot apical meristem (SAM)

region (Fig. 3B). On the 5th day of SE induction, the localization of MP11ir remained in the adaxial

side of cotyledon and SAM (Fig. 3C). In contrast, MP/ARF5 showed a different pattern of

expression. It was not detected in freshly isolated IZEs (Fig. S4A). After 24 hours of auxin-

dependent SE induction, MP/ARF5 was expressed in single cells of hypocotyl, SAM, and the

abaxial side of the cotyledon (Fig. S4). Both proteins MP11ir and MP/ARF5 were present during

the SE process but localized in different regions within the explants, suggesting that MP11ir and

MP/ARF5 might have distinct roles during the SE process.

Overexpression of a truncated MP/ARF5 inhibits the SE process.

To better understand the complex roles of full-length and truncated MP/ARF5 proteins in SE, we

investigated the effects of ectopic expressing a truncated MP/ARF5 protein lacking the PB1

domain (ΔARF5). ΔARF5 mimics the MP11ir protein, thereby helping to dissect the functional

relevance of ARF-ARF or ARF-Aux/IAA interactions for SE. We utilized a β-estradiol inducible

XVE::ΔARF5 DR5::GFP transgenic line to overexpress ΔARF5. We observed that overexpression

of *ΔARF5* inhibits the SE process when explants are cultured on a 2,4-D-supplemented medium

(Fig. 4). Specifically, explants overexpressing *ΔARF5* treated with 1 µM β-estradiol and various

concentrations of 2,4-D (1.0, 2.5, 5.0 µM 2,4-D) developed callus tissue rather than somatic

embryos. This outcome underscores that the constitutive overexpression of *ΔARF5* affects

morphogenesis and disrupts the transition to embryonic development. These results indicate that

the cell-type-specific expression of MP11ir is crucial, and its ubiquitous overexpression impedes

SE induction.

Truncated MP/ARF5 and canonical MP/ARF5 proteins influence genes involved in auxin

biosynthesis

The inhibitory effect of overexpressing *ΔARF5* on the SE process prompted us to investigate its

potential influence on endogenous auxin levels. SE inhibition often correlates with suboptimal

auxin concentrations during SE induction with IZE. Indeed, the overexpression *LEAFY*

COTELYDON 2 (*LEC2*) gene, a master regulator of the embryogenic process during auxin-

dependent SE induction, resulted in SE inhibition due to increased endogenous auxin levels

(Wójcikowska and Gaj, 2015; Wójcikowska et al., 2013). Similarly, the combined treatment of

exogenous auxin and TSA similarly inhibits SE in IZEs (Wójcikowska et al., 2018). To explore

the link between ΔARF5-induced phenotypes and fluctuations in endogenous auxin levels, we

examined the expression of the auxin biosynthetic genes from the IPyA pathway: *TRYPTOPHAN*

AMINOTRANSFERASE of ARABIDOPSIS 1 (*TAA1*)*, TAA1-related 1* (*TAR1*) and *TAR2*, and flavin

monooxygenases *YUCCA 1-11* (*YUC1-11*). These genes are expressed during the SE process and

pivotal for auxin production during SE formation (Bai et al., 2013; Karami et al., 2023; Li et al.,

2022). Using the *XVE::ΔARF5 DR5::GFP* transgenic line, we induced *ΔARF5* overexpression in explants cultured on 2,4-D supplemented medium for 1, 5, and 10 days with 1 µM β-estradiol. Our analysis revealed significant alterations in the expression of *TAA1*, *TAR1*, *YUC3*, *YUC5*, and *YUC8* genes upon *ΔARF5* overexpression (Fig. 5A-C). Specifically, *TAA1*, *TAR1*, *YUC5*, and *YUC8* showed increased expression levels after 24 hours of *ΔARF5* overexpression (Fig. 5A), while *YUC3* and *YUC8* were upregulated on the 5th day (Fig. 5B). Overexpression of *ΔARF5* for ten days sustained the up-regulation of *TAA1* and *YUC5* genes but let to down-regulation of *YUC3* and no change in *TAR1* and *YUC8* expression (Fig. 5C). Further analysis in the *35S::bdl-GR* line, where the DEX-dependent expression of the auxin-insensitive bdl inhibits MP/ARF5 activity, and in the *mp/arf5* mutant confirmed that *TAA1*, *TAR1*, *YUC3, YUC5* and *YUC8* are sharply down-regulated during conditions of reduced MP/ARF5 function (Fig. 5D-F). The expression profiles of the remaining genes in *XVE::ΔARF5 DR5::GFP* (Fig. S5), *35S::bdl-GR,* and *mp/arf5* lines (data not presented) were mostly characterized by constant or reduced expression.

Notably, the promoters of *TAA1*, *TAR1*, and *YUC5* were found to harbor binding peaks for MP/ARF5 and *YUC8* promoter for ARF2, as detected by DAP-seq analysis (O'Malley et al., 2016) (Fig. S6). All the genes deregulated in response to ΔARF5 have at least one ARF-binding peak. These results collectively suggest that ΔARF5, and by extension MP11ir, positively regulate the expression of genes involved in auxin biosynthesis, specifically *TAA1*, *TAR1*, *YUC3*, *YUC5,* and *YUC8* genes.

The localization of TAA1, TAR1, YUC3, YUC5, and YUC8 enzymes during induction of the SE process is regulated by MP/ARF5 and MP11ir

To elucidate the spatial distribution of the enzymes involved in the IPyA-dependent auxin biosynthesis during the embryogenic transition, we utilized translational reporter lines for TAA1, TAR1, TAR2, and YUCs proteins (Fig. S7). The TAA1, TAR1, YUC3, YUC5, and YUC8 enzymes were selected based on the RT-qPCR results (Fig. 5). To validate the regulation of their expression by ΔARF5, and thereby MP11ir, and MP/ARF5, we crossed the translational reporter lines for TAA1, TAR1, YUC3, YUC5, and YUC8 into *XVE::ΔARF5 DR5::GFP* and *35::bdl-GR* transgenic lines, and *mp/arf5* mutants to investigate the localization of GUS signals. Explants were induced into SE with 2,4-D and 1 µM β-estradiol or 10 µM DEX to induce *ΔARF5* or bdl

overexpression, respectively, and sampled on the 5th and 10th days of the culture. Additionally,

mp/arf5 IZE (0 day) and explants on the 5th and 10th day of the SE process were analyzed.

Among the enzymes analyzed, TAA1, YUC3, YUC5, and YUC8 exhibited distinct responses to

ΔARF5 overexpression (Fig. 6). Notably, YUC8 showed robust accumulation in the proximal

regions of cotyledons and hypocotyl upon *ΔARF5* overexpression, as evidenced by intense GUS

signals (Fig. 6A). Conversely, bdl overexpression led to a significant decrease in GUS signal

intensity for YUC8 (Fig. 6B). In *mp/arf5* mutant, the GUS signal for YUC8 was absent (Fig. 6C),

confirming the MP- and MP11ir-dependent expression of the *YUC8* gene. TAA1 was broadly

active throughout the IZE explants during embryogenic transition (5th and 10th day), with a slight

increase of the GUS signal upon *ΔARF5* overexpression (Fig. 6A). However, changes in GUS

intensity were modest upon bdl overexpression or in *mp/arf5* mutant (Fig. 6B, C). YUC3 exhibits

ubiquitous localization in the hypocotyl region of explants, which is enhanced in the cotyledons

upon *ΔARF5* overexpression (Fig. 6A). In contrast, bdl overexpression deeply reduced the intensity

of the GUS signal (Fig. 6B), which was completely absent in *mp/arf5* mutant IZEs or reduced at

5th and 10th days of SE induction (Fig. 6C). YUC5 showed a specific accumulation at the ends of

cotyledon only on the 10th day-old SE under *ΔARF5* overexpression (Fig. 6A), with no changes

observed during bdl overexpression (Fig. 6B). In *mp/arf5* mutant, the GUS signal for YUC5 was

restricted to single cells (Fig. 6C). In contrast, TAR1 localization was unchanged upon *ΔARF5,*

and bdl overexpression, and in *mp/arf5* mutant.

These findings demonstrated the cell-type specific and temporal regulation of the TAA1, YUC3,

YUC5, and YUC8 auxin biosynthetic enzymes by ΔARF5 (MP11ir) and MP/ARF5 proteins during

the SE process.

Influence of MP11ir on endogenous auxin levels

We hypothesized that MP11ir might regulate the endogenous auxin levels through TAA1, YUC3,

YUC5, and YUC8 during SE. We explored this hypothesis using three approaches. First, we

assessed local auxin responses in the *XVE::ΔARF5 DR5::GFP* explants during the auxin-

independent embryogenic transition (1st and 5th days of TSA treatment) to avoid activation of the

DR5 signal by the exogenously applied auxin. Explants were cultured on ET medium with or

without β-estradiol. GFP signals indicate auxin-responsive gene transcription in the root apical

meristem and the tip of cotyledons in non-induced explants (Fig. 7A, E). After 24 hours of induced

ΔARF5 overexpression, an enhanced GFP signal was observed in the hypocotyl and throughout the cotyledons (Fig. 7B, E). In 5-day-old explants, *ΔARF5* overexpression intensified the GFP signal in the hypocotyl and cotyledons (Fig. 7C, D, F). Second, we quantified indolic compounds in a 3-week-old *in vitro* cultures of *XVE::ΔARF5 DR5::GFP* explants treated and not with β-estradiol (Fig. 7G). The explants overexpressing *ΔARF5*, exhibiting callus formation, showed a 1.37-fold increase in indolic compound levels compared to non-induced explants. Finally, to determine whether the formation of calli might result from high auxin responses and indolic compound levels, we analyzed the SE process in Col-0 and the *XVE::ΔARF5 DR5::GFP* line treated with different concentrations of yucasin (100 and 150 µM), an inhibitor of YUC activity (Tsugafune et al., 2017). (Non-)induced Col-0 (-E, +E) and non-induced *XVE::ΔARF5 DR5::GFP* (-E) explants treated with yucasin showed highly reduced SE efficiency (1.3 to 1.5 times lower) and productivity (no more than two somatic embryos per reacting explant were formed) (Fig. S8). The results align with published literature (Karami et al., 2023; Li et al., 2022). We then investigated whether treating *XVE::ΔARF5 DR5::GFP* IZE explants with yucasin and inducing *ΔARF5* overexpression with β-estradiol could reverse the phenotype, enabling the explants to regenerate embryos instead of callus tissue. *XVE::ΔARF5 DR5::GFP* IZE explants treated with yucasin (100 and 150 µM) formed somatic embryos, but very rarely (15.4% and 3.4%, respectively) and with low productivity (1 somatic embryo per reacting explant) (Fig. S8).

This finding supports the notion that *ΔARF5* overexpression amplifies local auxin responses by modulating auxin production and signaling during SE, ultimately forming calli instead of embryonic structures. However, other genetic components deregulated by *ΔARF5* overexpression are likely inhibiting SE capacity.

Discussion

MP/*ARF5* is upregulated in plant embryogenic cells and serves as a potential molecular marker for this developmental stage. Various transcriptomic studies support the involvement of *MP*/*ARF5* in the SE process in various plants (Awada et al., 2023; Capote et al., 2019; Chen et al., 2020; Gliwicka et al., 2013; Indoliya et al., 2016; Juárez-González et al., 2019; Qi et al., 2021; Quintana-Escobar et al., 2019; Shi et al., 2016; Thibaud-Nissen et al., 2003; Wickramasuriya and Dunwell, 2015; Xu et al., 2024; Zeng et al., 2024; Zhu et al., 2023). Additionally, the presence of *MP*/*ARF5* transcripts in regions critical for SE induction, such as cotyledons and SAM, suggests that

MP/ARF5 may be involved in the embryogenic transition (Wójcikowska and Gaj, 2017). Two

mp/*arf5* mutant alleles, *arf5* (SALK_001058C) (Wójcikowska and Gaj, 2017) and *mpS319*

(SALK_021319) (present work), exhibit impaired SE processes. This inhibition is linked to the

differential expression of SE master regulators (Wójcikowska and Gaj, 2017). Similarly, the *Osarf5*

mutant shows partial defects in forming scutellum-derived embryogenic callus and a significantly

reduced expression of *LEAFY COTELYDON 1* (*OsLEC1*), indicating that the OsARF5-mediated

auxin signaling pathway might regulate *OsLEC1* (Guo et al., 2023a).

Expressing auxin-insensitive bdl protein during SE induction revealed that SE inhibition in the

mp/arf5 mutant is not due to embryonic defects but is triggered by MP/ARF5 inactivity in IZE.

Many *EMBRYO-DEFECTIVE* (*EMB*) genes involved in zygotic embryo development and

morphogenesis, such as *LEC1*, *LEC2*, *FUSCA 3*, *DICER-LIKE 1*, *WUSCHEL RELATED*

HOMEOBOX 5 (*WOX5*), and *PIN1*, are also critical for SE processes (Ledwoń and Gaj, 2009;

Ledwoń and Gaj, 2011; Meinke, 2020; Su et al., 2015; Su et al., 2009; Szyrajew et al., 2017).

However, the overexpression of the auxin-insensitive bdl protein had a more substantial impact on

inhibiting embryogenic transition than the *mp/arf5* mutation, suggesting that mutated bdl protein

might repress other ARFs besides MP/ARF5 (Piya et al., 2014).

MP/ARF5's transcriptional targets likely include genes encoding proteins essential to the SE

process. AGAMOUS-LIKE 15 (AGL15) and AGL18, crucial for SE (Harding et al., 2003; Paul et

al., 2022; Thakare et al., 2008; Zheng and Perry, 2014), might regulate *MP*/*ARF5* expression as the

agl15 agl18 double mutant, highly impaired in SE, shows a significant reduction of *MP*/*ARF5*

mRNA levels (Zheng et al. 2009). Also, inhibition of PHYTOCHROME-INTERACTING

FACTOR 4 (PIF4) induces several auxin responses (*ARF5, ARF8,* and *ARF16*) and biosynthetic

(*YUC1, YUC2*, *YUC6*, *CYP79B2* and *AMIDASE 1* (*AMI1*)) genes, resulting in the formation of the

somatic embryos (Mira et al., 2023).

Despite these insights, the detailed mechanism of action of MP/ARF5 and its MP11ir isoform

during SE induction and somatic embryo formation remains unknown. Lines expressing truncated

ARF5 proteins without the PB1 domain could elucidate the complex action of MP11ir by

mimicking its function. Truncated ΔARF5 and MP^{abn} proteins partially rescue *mp/arf5* phenotypic

defects but also cause unique leaf defects (Garrett et al., 2012; Krogan et al., 2012). Ectopic ΔARF5

protein stimulates the formation of new shoots *in vivo* (Ma et al., 2019), whereas *mpS319* mutants

show reduced shoot regeneration frequency (Zhang et al., 2021). During *in vitro* culture, ΔARF5

increases the shoot organogenesis frequencies in Arabidopsis root-derived explants and activated auxin signaling. Visualization and cytometric quantification of the DR5::GFP nuclear auxin signaling reporter indicated increased GFP signal in roots and protoplasts overexpressing *ΔARF5* (Gonzalez et al., 2021). A similar trend was observed after inducing *ΔARF5* expression during the embryogenic transition. MP/ARF5 is involved in shoot apical meristem (SAM) activity and maintenance by directly controlling the expression of ARABIDOPSIS HISTIDINE PHOSPHOTRANSFER PROTEIN 6 (AHP6) (Besnard et al., 2014), TARGET OF MONOPTEROS (TMOs) (Schlereth et al., 2010), DORNRÖSCHEN/ENHANCER OF SHOOT REGENERATION 1 (DRN/ESR1) (Cole et al., 2009), *WUSCHEL RELATED HOMEOBOXs* (*WOXs*) (Lee et al., 2022; Tvorogova et al., 2019), essential for shoot regeneration or SE (Fambrini et al., 2022). The shoot regeneration-promoting effect of ΔARF5 may overcome organogenic resistance in recalcitrant explants and species (Ckurshumova and Berleth, 2015; Gonzalez et al., 2021). However, *ΔARF5* overexpression did not enhance embryogenic capacity (our results) but instead led to callus formation, similar to *NcARF5* overexpression in *Neolamarckia cadamba*, which triggers rapid callus proliferation (Ma et al., 2024).

Studies on embryo-derived SE systems showed that SEs initiate from the callus edge, where auxin accumulates, as visualized using PIN-FORMED 1 (PIN1) and DR5 reporters (Kadokura et al., 2018; Su et al., 2015; Su et al., 2009). MP/ARF5 directly and positively regulates PIN1, responsible for auxin efflux in polar auxin transport (PAT) (Krogan et al., 2016). Auxin is present in the *pin1* mutant meristem, ensured by vascular tissue transport or induction of auxin biosynthesis (Banasiak et al., 2019). Therefore, *MP/ARF5* is proposed as a primary regulator of auxin transport and biosynthesis. Our study identifies auxin biosynthetic genes, *TAA1*, *TAR1*, *YUC3*, *YUC5,* and *YUC8*, potentially positively regulated by ΔARF5, and thus MP11ir, during *in vitro* embryogenic transition. We propose that MP/ARF5 and MP11ir proteins directly or indirectly control the TAA1/TARs-YUCs-mediated auxin biosynthetic pathway (our work) and polar auxin transport during embryogenesis (Robert et al., 2015; Schlereth et al., 2010). This hypothesis aligns with findings that MP/ARF5 affects *YUC1* and *YUC8* expression to specify the ground tissue in early globular zygotic embryos (Möller et al., 2017). IPyA-dependent auxin biosynthesis is crucial for effective SE induction and somatic embryo identity and growth (Li et al., 2022). Yucasin treatment, inhibiting YUC activity (Tsugafune et al., 2017), reduces SE efficiency and productivity in Arabidopsis (Karami et al., 2023; Li et al., 2022) and coffee culture (Uc-Chuc et al., 2020). The

importance of auxin biosynthesis in the SE process is evidenced by the decreased embryogenic response in *yuc* mutants. Multiple *yuc* mutants, *yuc3 yuc8* (Li et al., 2022), *yuc1 yuc4 yuc10 yuc11* (Bai et al., 2013), *yuc3 yuc5 yuc7 yuc8 yuc9* (Sakamoto et al., 2022), display disrupted SE or callus formation compared to single mutants (Sakamoto et al., 2022; Wójcikowska et al., 2013). *YUC* genes are direct targets of TFs, such as LEC1, LEC2, BABY BOOM (BBM), and AT-HOOK MOTIF CONTAINING NUCLEAR LOCALIZED 15 (AHL15), which play pivotal roles in embryogenic transition, demonstrated by decreased SE in their mutant and spontaneous induction of somatic embryo formation when overexpressed (Boutilier et al., 2002; Gaj et al., 2005; Karami et al., 2021; Ledwoń and Gaj, 2009; Lotan et al., 1998; Stone et al., 2001; Zuo et al., 2002). During SE induction, LEC1 activates *YUC4* and *YUC10* (Hu et al., 2018; Junker et al., 2012), while LEC2 positively regulates *YUC1*, *YUC4*, and *YUC10* expression (Stone et al., 2008; Wójcikowska et al., 2013). BBM induces local and ectopic expression of *YUC3* and *YUC8* coupled with IAA biosynthesis and activation of the *WOX2* embryo marker (Li et al., 2022). During SE induction, AHL15 activates *YUC6*, *YUC7*, *YUC8*, and *YUC9* expression (Karami et al., 2023), confirming YUCs as crucial molecular elements of a network governing the SE process.

A recent study identified target genes of MP/ARF5 and truncated ΔARF5 proteins in 3-day-old seedlings (un)treated with auxin (Xie et al., 2022). This ChIP-seq analysis suggests that deleting the PB1 domain renders the protein auxin-insensitive and alters its binding profile. Among genes involved in the IPyA auxin biosynthesis pathway, *YUC3* and *YUC5* promoters are recognized and bound by ΔARF5, while the *YUC6* promoter is targeted by ΔARF5 and MP/ARF5. Differences in the preferential AuxRE motifs bound by ΔARF5 and MP/ARF5 suggest that the PB1 domain may affect homodimer formation and DNA binding, indicating that ΔARF5 and MP/ARF5 may differ in their molecular actions and thus impact plant development differently (Xie et al., 2022).

Transcriptomic data have shown that many splice isoforms formed due to alternative splicing (AS) show cell-specific expression (Klepikova et al., 2016; Martín et al., 2021). Among AS events, intron retention events are functionally underestimated in plants. Given the importance of AS in responding to various stresses, it is likely required for proper environmental response in plants (Guo et al., 2023b). Limited functional evidence supports their specific expression patterns or distinct ability to rescue mutant loss-of-function phenotypes. For example, *ARF8* undergoes tissue-specific AS in flower, where the *ARF8.4* variant, which retains the eighth intron and has a premature stop codon, is over-represented. Overexpression of *ARF8.4*, but not other *ARF8*

isoforms, reverts stamen elongation defects associated with the *arf8* knockout mutation (Ghelli et al., 2023; Ghelli et al., 2018). Another example is the *BIGPETAL* (*BPE*) gene, which encodes a petal-specific *BPEp* transcript that retains its last intron, in addition to the canonical *BPEub* transcript uniformly expressed in all organs (Szécsi et al., 2006). The C-terminal sequence exclusively encoded by *BPEp* is required to interact with ARF8. The *arf8* loss-of-function mutant phenocopies the petal defects of the *bpe* knockout mutants, suggesting that BPEp, unlike the canonical BPEub protein, is required for petal development (Kashkan et al., 2022; Varaud et al., 2011).

In conclusion, both MP/ARF5 proteins, canonical MP/ARF5 and MP11ir isoform, seem to be involved in the SE process (Fig. 8A). The different expression patterns of MP/ARF5 and MP11ir suggest that they act in different cells, probably with different endogenous auxin levels. The ectopic expression of *ΔARF5,* mimicking MP11ir, results in a global increase in endogenous auxin content in explants through direct or indirect activation of *TAA1*, *TAR1*, *YUC3*, *YUC5,* and *YUC8* genes. Disruption of auxin homeostasis in explant cells leads to SE inhibition and callus formation (Fig. 8B). Blocking MP/ARF5 action through the bdl repressor results in decreased levels of *YUC3* and *YUC8* during SE induction, inhibiting the SE process and promoting seedling development (Fig. 8C). The absence of both MP/ARF5 and MP11ir in *mp/arf5* mutant leads to the absence of *YUC3*, *YUC5,* and *YUC8* during SE induction, resulting in SE inhibition, callus formation or impaired seedling development (Fig. 8D). In summary, this study provides an elegant and straightforward experimental setup that elucidates the role of MP/ARF5 and MP11ir in SE induction, acting through the control auxin biosynthetic genes. These findings suggest that MP/ARF5 and MP11ir are central regulators of auxin biosynthesis during embryogenic processes.

Materials and methods

Plant material

The seeds of *Arabidopsis thaliana* (L.) Heynh Columbia (Col-0) wild type, *mpS319* mutant (N21319), reporter lines *pTAA1::TAA1-GUS* (N72246), *pTAR1::TAR1-GUS* (N72247), *pTAR2::TAR2-GUS* (N72248), *pYUC1::YUC1-GUS* (N72250), *pYUC2::YUC2-GUS* (N72252), *pYUC3::YUC3-GUS* (N72254), *pYUC4::YUC4-GUS* (N72256), *pYUC5::YUC5-GUS* (N72259), *pYUC6::YUC6-GUS* (N72261*), pYUC7::YUC7-GUS* (N72262), *pYUC8::YUC8-GUS* (N72264), *pYUC9::YUC9-GUS* (N72266), *pYUC01::YUC10-GUS* (N72268), and *pYUC11::YUC11-GUS*

(N72271) were supplied by NASC (The Nottingham Arabidopsis Stock Centre) (Alonso et al.,

2003; Brumos et al., 2019). Seeds of the *MP::MP mpS319, MP::MP11ir mpS319*, and

pMP::MP11ir-GFP (MR) transgenic lines were provided by Lucia Colombo (Dipartimento di

BioScienze, Università degli Studi di Milano, Milano, Italy). Dolf Weijers (Laboratory of

Biochemistry, Wageningen University, and Research, Wageningen, The Netherlands) generously

donated seeds of the *pMP::MP-GFP*, *35S::BDL-GR*, *35S::bdl-GR* lines. The seeds of the

XVE::ΔARF5 DR5::GFP line were provided by Bastiaan Bargmann (School of Plant and

Environmental Sciences, Virginia Polytechnic Institute and State University, Blacksburg, USA).

The generation of *MP::MP mpS319* x *MP::MP11ir mpS319*, and *XVE::ΔARF5 DR5::GFP*;

35S::bdl-GR; arf5 x *pTAA1::TAA1-GUS*, *pTAR1::TAR1-GUS*, *pYUC3::YUC3-GUS*,

pYUC5::YUC5-GUS, *pYUC8::YUC8-GUS* was performed by crossing.

Plant growth and *in vitro* culture conditions

Seeds were sterilized with sodium chloride (20% commercial bleach) and plated onto ½ MS

medium (Murashige and Skoog, 1962), (Duchefa Biochemie; #M0222), 10 g L^{-1} sucrose (Penta #

57-50-1) and 8 g L^{-1} agar (Duchefa Biochemie; #P1001). Seed-derived plants grown *in vitro* were

used as a source of seedlings, and *in vivo* were used as a source of leaves, flowers, and siliques for

IZE isolation. Plants were grown in soil with vermiculite (4:1) at 22 °C under a 16 h photoperiod

of 100 µM m^{-2} s^{-1} white, fluorescent light.

Somatic embryogenesis induction

IZEs at the cotyledonary stage of development were used as explants for the *in vitro* culture. The

IZEs were cultured on an E5 medium containing 3.2 g L^{-1} of B5 micro and macro-elements

(Duchefa Biochemie; #G0210) (Gamborg et al., 1968), 20 g L^{-1} sucrose, 5.0 µM of 2,4-D (2,4-

dichloro phenoxy acetic acid, Sigma-Aldrich #D7299) and 8 g L^{-1} agar (Duchefa Biochemie;

#P1001), pH 5.8 (Gaj, 2001). To induce the SE process without auxin, the IZEs were cultured on

an ET medium containing 3.2 g L^{-1} of B5 micro and macro-elements (Duchefa Biochemie;

#G0210) (Gamborg et al., 1968), 20 g L^{-1} sucrose, 1 µM of TSA (Trichostatin A, Sigma Aldrich;

#T1952) and 8 g L^{-1} agar (Duchefa Biochemie; #P1001), pH 5.8 (Wójcikowska et al., 2018). Plant

materials that grow in sterile conditions were kept at 23 °C under a 16 h photoperiod of 40 µM m^{-2}

s^{-1} white, fluorescent light. The explant capacity for SE was evaluated in an IZE-derived culture

that was induced for 21 days on an induction media, and two parameters were calculated – SE efficiency (percentage of explants that developed somatic embryos), and productivity (average number of somatic embryos produced by embryogenic explant). Thirty explants were analyzed in at least three biological replicates of the experimental combination. Around 10,000 IZEs were isolated for all the experiments. Assuming the average embryo isolation rate (100 IZEs per hour), the time to obtain the research material for each experiment was 100 hours.

IZE treatments

IZEs at the cotyledonary stage were treated with different 2,4-D (Duchefa Biochemie; #D0911), IAA (Duchefa Biochemie; #I0901) and NAA (Duchefa Biochemie; #N0903) (0.0, 2.5, 5.0, 7.5 µM) concentrations for ten days to evaluate the level of *MP/ARF5* and *MP11ir* transcripts. IZEs were cultured on a medium with 2,4-D and treated with different yucasin (Merck; #23711-26-4) concentrations (0, 100, 150 µM) for 3 weeks to evaluate the effect of YUC activity inhibition on the embryogenic capacity. Explants were first precultured for 1 day on a medium without yucasin and then transferred to a medium with different yucasin concentrations to allow for IZE survival. Treating IZE explants with yucasin from the beginning of the culture (without preculture) resulted in the death of the explants (Karami et al., 2023).

Induction of BDL, bdl, and *ΔARF5* expression in transgenic plants

The action of BDL and bdl was controlled by adding dexamethasone (water-soluble DEX, Sigma; #D2915) to the media at a concentration of 10 or 30 µM. The expression of *ΔARF5* was controlled by adding 1 µM β-estradiol (ethanol-soluble E, Sigma; # E8875) to the media.

RNA isolation, reverse transcription, qualitative and quantitative PCR

RNA was isolated from different samples, depending on culture age, tissue from 300 (0, 1st day), 150 (5th day) to 50 (10th day) explants were used for RNA extraction. A NucleoSpin RNA Plant Kit (Macherey-Nagel) was used to isolate the RNA from the samples. The RNAs were treated with RQ1 RNase-free DNase I (Promega) to control DNA contamination according to the manufacturer's instructions. The first-strand cDNA was produced from 1,000 ng of RNA using a RevertAid First Strand cDNA Synthesis Kit (Fermentas). The product of the reverse transcription was used for RT-PCR and RT-qPCR reactions to detect and quantify gene transcripts. Undiluted

cDNA (1 µL) was used for RT-PCR. cDNA diluted with water at a 1:4 ratio (2.5 µL) was used for

the RT-qPCR reactions. RT-qPCR was carried out in a 10 µL reaction volume using a LightCycler®

480 SYBR^{TM} Green I Master (Roche) kit, a LightCycler® 96 Multiwell Plate 96, and

MultiwellSealing Foil (Roche). The relative RNA levels were calculated and normalized to the

internal control of the AT4G27090 (*TIN*) gene encoding the 60S ribosomal protein (Thellin et al.,

1999). Fold change values were calculated using the comparative $2^{-\Delta\Delta Ct}$ method, where $\Delta\Delta Ct$

represents ΔCt $^{\text{reference condition}}$ − ΔCt $^{\text{compared condition}}$. 40-ΔCt method was used to compare the

expression levels of *MP* and *MP11ir* (Czechowski et al., 2004). The plant tissues for gene

expression analysis were produced in three biological replicates, and two technical replicates were

analyzed. Primer sequences for *MP11ir*, *MP*, *TIN*, *TAA1*, *TARs*, and *YUCCAs* are in Suppl. Tab.

S1.

Fixation

The explants were fixed using a fixative solution (4% paraformaldehyde in 1X PBS with 0.05%

Triton-X100 (PBS-T)) and incubated in a 1.5 mL fixative in a microcentrifuge tube. The tubes

were covered with aluminum foil throughout the procedure to be light-tight. The samples were

vacuum infiltrated for ~ 30 min on ice to remove air bubbles. Then, the samples were incubated

overnight at 4 °C with slow rotation. The following day, samples were washed three times with 1X

PBS-T at room temperature for one hour each.

GUS staining

577   The GUS enzyme activity was performed as described by Bassuner et al. (2007). The explants were

578   stained in a standard X-Gluc solution at 37 °C for 12 h. The experiments were repeated two times

579   for each studied combination, and at least 30 somatic embryos were subjected to GUS signal

580   detection in each repetition. The GUS staining pattern was analyzed using a microscope Zeiss

581   Axioscope.A1 ZEN.

ClearSee alpha clearing

584   The ClearSee alpha solution was prepared according to Kurihara et al. (2021). The ClearSee

585   solution was supplemented with 50 mM sodium sulfite as an antioxidant. The clearing was

586   extended to 7 days with gentle shaking at room temperature for fixed explants.

Renaissance SR2200 staining

A 0.1% (v/v) Renaissance SR2200 (Renaissance Chemicals, UK) solution was used to stain the samples. After clearing, the staining step was performed by mixing the dye with the mounting medium ClearSee alpha (Attuluri et al., 2022). After staining, the samples were washed for one hour in the same solution used for the staining procedure.

Microscopy

Expression pattern analysis was performed on a ZEISS LSM 700 confocal microscope with a correction collar $10 \times$ magnification objective at 0.6 zoom at 1024×1024 pixel resolution. Two laser lines were used: 405 nm for imaging Renaissance combined with 488 nm for GFP imaging. The images were captured in 12 bits. The laser intensity and detector gain were kept constant for all samples.

Content of indolic compounds

A colorimetric technique that enabled the detection of indolic compounds, including IAA, was applied (Bric et al., 1991; Wójcikowska et al., 2013). Each analysis was carried out in five biological replicates.

ARF binding regions

We employed the DAP-Seq profiles for the MP/ARF5 and ARF2 transcription factors (O'Malley et al., 2016) available at the PlantCistrome database (http://neomorph.salk.edu/PlantCistromeDB). We analyzed (-1500; +100) *TAA1/TAR* and *YUC* gene promoters and detected TGTCnn sequences and ARF-binding DAP-Seq peaks. We utilized a suite of R packages including biomaRt, dplyr, and ggplot2 for the computational analysis and data visualization.

Statistical analysis

Phenotype scoring and quantification data were collected in EXCEL (Microsoft Corp.) and, from there, imported into STATISTICA (TIBCO Software Inc., Palo Alto, CA, USA) for statistical analysis and to generate graphs. For one-on-one comparisons, The Student's T-test was used ($p < 0.05$). For intercomparison of more than two data points, the two-way ANOVA analysis ($p < 0.05$)

618 followed by Tukey's HSD test ($p < 0.05$, and $p < 0.01$) was used to determine any significant
619 differences between the compared combinations. The graphs show the means with the standard
620 deviations (SD).
621

Supplementary Data
Supplementary Figure S1. MP11ir is expressed during somatic embryogenesis.
Supplementary Figure S2. The induction of somatic embryogenesis depends on 2,4-D and NAA,
not IAA.
Supplementary Figure S3. Expression of MP/ARF5 during induction of somatic embryogenesis.
Supplementary Figure S4. Expression pattern of MP/ARF5 during somatic embryogenesis
induction.
Supplementary Figure S5. Expression level of auxin biosynthetic genes in *XVE::ΔARF5*
DR5::GFP somatic embryos.
Supplementary Figure S6. Analysis of the *TAA1*, *TAR1*, and *YUC* promoters of MP/ARF5 biding
sites.
Supplementary Figure S7. Expression pattern of TAA1, TAR1, TAR2 and YUC1-11 during
somatic embryogenesis induction.
Supplementary Figure S8. Effects of repressing auxin biosynthesis in *XVE::ΔARF5 DR5::GFP*
during somatic embryogenesis induction.
Supplementary Table S1. The primers used in experiments.
638

639 **Funding**
640 B.W. was supported by the Polish National Agency for Academic Exchange
641 (BPN/BEK/2021/1/00278/U/00001). This work was supported by the Ministry of Education,
642 Youth and Sports of the Czech Republic with the European Regional Development Fund-Project
643 "Centre for Experimental Plant Biology" (No. CZ.02.1.01/0.0/0.0/16_019/0000738).
644

645 **Acknowledgments**
646 We acknowledge the CEITEC core facilities Plant Sciences and CELLIM, supported by the MEYS
647 CR (LM2023050 Czech-BioImaging). We are grateful to Lucia Colombo (Dipartimento di
648 BioScienze, Università degli Studi di Milano, Milano, Italy), Dolf Weijers (Laboratory of

Author contributions

B.W. and H.R.B. designed and supervised the project. B.W., S.B., and V.M. conducted the experiments and analyzed the data. B.W. and H.R.B. wrote the manuscript. All authors discussed the results and approved the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., Gadrinab, C., Heller, C., Jeske, A., Koesema, E., Meyers, C.C., Parker, H., Prednis, L., Ansari, Y., Choy, N., Deen, H., Geralt, M., Hazari, N., Hom, E., Karnes, M., Mulholland, C., Ndubaku, R., Schmidt, I., Guzman, P., Aguilar-Henonin, L., Schmid, M., Weigel, D., Carter, D.E., Marchand, T., Risseeuw, E., Brogden, D., Zeko, A., Crosby, W.L., Berry, C.C., Ecker, J.R., 2003. Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science 301, 653-657.

Attuluri, V.P.S., Sánchez López, J.F., Maier, L., Paruch, K., Robert, H.S., 2022. Comparing the efficiency of six clearing methods in developing seeds of Arabidopsis thaliana. Plant Reprod 35, 279-293.

Awada, R., Lepelley, M., Breton, D., Charpagne, A., Campa, C., Berry, V., Georget, F., Breitler, J.C., Léran, S., Djerrab, D., Martinez-Seidel, F., Descombes, P., Crouzillat, D., Bertrand, B.,

Etienne, H., 2023. Global transcriptome profiling reveals differential regulatory, metabolic and hormonal networks during somatic embryogenesis in Coffea arabica. BMC Genomics 24, 41.

Bai, B., Su, Y.H., Yuan, J., Zhang, X.S., 2013. Induction of somatic embryos in Arabidopsis requires local YUCCA expression mediated by the down-regulation of ethylene biosynthesis. Mol Plant 6, 1247-1260.

Ballesteros, D., Martínez, M.T., Sánchez-Romero, C., Montalbán, I.A., Sales, E., Moncaleán, P., Arrillaga, I., Corredoira, E., 2024. Current status of the cryopreservation of embryogenic material of woody species. Frontiers in Plant Science 14, 1337152.

Banasiak, A., Biedroń, M., Dolzblasz, A., Berezowski, M.A., 2019. Ontogenetic Changes in Auxin Biosynthesis and Distribution Determine the Organogenic Activity of the Shoot Apical Meristem in pin1 Mutants. International Journal of Molecular Sciences 20, 180.

Bassuner, B.M., Lam, R., Lukowitz, W., Yeung, E.C., 2007. Auxin and root initiation in somatic embryos of Arabidopsis. Plant Cell Rep 26, 1-11.

Besnard, F., Refahi, Y., Morin, V., Marteaux, B., Brunoud, G., Chambrier, P., Rozier, F., Mirabet, V., Legrand, J., Lainé, S., 2014. Cytokinin signalling inhibitory fields provide robustness to phyllotaxis. Nature 505, 417-421.

Bhatia, N., Bozorg, B., Larsson, A., Ohno, C., Jönsson, H., Heisler, M.G., 2016. Auxin Acts through MONOPTEROS to Regulate Plant Cell Polarity and Pattern Phyllotaxis. Curr Biol 26, 3202-3208.

Boer, D.R., Freire-Rios, A., van den Berg, W.A., Saaki, T., Manfield, I.W., Kepinski, S., López-Vidrieo, I., Franco-Zorrilla, J.M., de Vries, S.C., Solano, R., Weijers, D., Coll, M., 2014. Structural basis for DNA binding specificity by the auxin-dependent ARF transcription factors. Cell 156, 577-589.

Boutilier, K., Offringa, R., Sharma, V.K., Kieft, H., Ouellet, T., Zhang, L., Hattori, J., Liu, C.M., van Lammeren, A.A., Miki, B.L., Custers, J.B., van Lookeren Campagne, M.M., 2002. Ectopic expression of BABY BOOM triggers a conversion from vegetative to embryonic growth. Plant Cell 14, 1737-1749.

Bric, J.M., Bostock, R.M., Silverstone, S.E., 1991. Rapid in situ assay for indoleacetic acid
production by bacteria immobilized on a nitrocellulose membrane. Applied and environmental
Microbiology 57, 535-538.

Brumos, J., Zhao, C., Gong, Y., Soriano, D., Patel, A.P., Perez-Amador, M.A., Stepanova, A.N.,
Alonso, J.M., 2019. An Improved Recombineering Toolset for Plants. The Plant Cell 32, 100-122.

Cancé, C., Martin-Arevalillo, R., Boubekeur, K., Dumas, R., 2022. Auxin response factors are keys
to the many auxin doors. New Phytol 235, 402-419.

Capote, T., Usié, A., Barbosa, P., Ramos, M., Morais-Cecílio, L., Gonçalves, S., 2019.
Transcriptome dynamics of cork oak (Quercus suber) somatic embryogenesis reveals active gene
players in transcription regulation and phytohormone homeostasis of embryo development. Tree
Genetics & Genomes 15, 52.

Caumon, H., Vernoux, T., 2023. A matter of time: auxin signaling dynamics and the regulation of
auxin responses during plant development. Journal of Experimental Botany 74, 3887-3902.

Chandler, J.W., 2016. Auxin response factors. Plant Cell Environ 39, 1014-1028.

Chen, Y., Xu, X., Liu, Z., Zhang, Z., XuHan, X., Lin, Y., Lai, Z., 2020. Global scale transcriptome
analysis reveals differentially expressed genes involve in early somatic embryogenesis in
Dimocarpus longan Lour. BMC Genomics 21, 4.

Chen, Z., Debernardi, J.M., Dubcovsky, J., Gallavotti, A., 2022. The combination of morphogenic
regulators BABY BOOM and GRF-GIF improves maize transformation efficiency. bioRxiv,
2022.2009.2002.506370.

Ckurshumova, W., Berleth, T., 2015. Overcoming recalcitrance-Auxin response factor functions
in plant regeneration. Plant signaling & behavior 10, e993293.

Cohen, J.D., Strader, L.C., 2024. An auxin research odyssey: 1989–2023. The Plant Cell 36, 1410-
1428.

Cole, M., Chandler, J., Weijers, D., Jacobs, B., Comelli, P., Werr, W., 2009. DORNROSCHEN is

a direct target of the auxin response factor MONOPTEROS in the Arabidopsis embryo.

Development 136, 1643-1651.

Cucinotta, M., Cavalleri, A., Guazzotti, A., Astori, C., Manrique, S., Bombarely, A., Oliveto, S.,

Biffo, S., Weijers, D., Kater, M.M., Colombo, L., 2021. Alternative Splicing Generates a

MONOPTEROS Isoform Required for Ovule Development. Curr Biol 31, 892-899.e893.

Czechowski, T., Bari, R.P., Stitt, M., Scheible, W.-R., Udvardi, M.K., 2004. Real-time RT-PCR

profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel

root- and shoot-specific genes. The Plant Journal 38, 366-379.

Elhiti, M., Stasolla, C., 2022. Transduction of Signals during Somatic Embryogenesis. Plants

(Basel) 11.

Fambrini, M., Usai, G., Pugliesi, C., 2022. Induction of Somatic Embryogenesis in Plants:

Different Players and Focus on WUSCHEL and WUS-RELATED HOMEOBOX (WOX)

Transcription Factors. Int J Mol Sci 23.

Freire-Rios, A., Tanaka, K., Crespo, I., van der Wijk, E., Sizentsova, Y., Levitsky, V., Lindhoud,

S., Fontana, M., Hohlbein, J., Boer, D.R., Mironova, V., Weijers, D., 2020. Architecture of DNA

elements mediating ARF transcription factor binding and auxin-responsive gene expression in

Arabidopsis. Proc Natl Acad Sci U S A 117, 24557-24566.

Gaj, M.D., 2001. Direct somatic embryogenesis as a rapid and efficient system for in vitro

regeneration of Arabidopsis thaliana. Plant Cell, Tissue and Organ Culture 64, 39-46.

Gaj, M.D., Zhang, S., Harada, J.J., Lemaux, P.G., 2005. Leafy cotyledon genes are essential for

induction of somatic embryogenesis of Arabidopsis. Planta 222, 977-988.

Gamborg, O.L.c., Miller, R.A., Ojima, K., 1968. Nutrient requirements of suspension cultures of

soybean root cells. Experimental cell research 50, 151-158.

Garrett, J.J., Meents, M.J., Blackshaw, M.T., Blackshaw, L.C., Hou, H., Styranko, D.M., Kohalmi,
S.E., Schultz, E.A., 2012. A novel, semi-dominant allele of MONOPTEROS provides insight into
leaf initiation and vein pattern formation. Planta 236, 297-312.

Ghelli, R., Brunetti, P., Marzi, D., Cecchetti, V., Costantini, M., Lanzoni-Rossi, M., Scaglia
Linhares, F., Costantino, P., Cardarelli, M., 2023. The full-length Auxin Response Factor 8 isoform
ARF8.1 controls pollen cell wall formation and directly regulates TDF1, AMS and MS188
expression. Plant J 113, 851-865.

Ghelli, R., Brunetti, P., Napoli, N., De Paolis, A., Cecchetti, V., Tsuge, T., Serino, G., Matsui, M.,
Mele, G., Rinaldi, G., Palumbo, G.A., Barozzi, F., Costantino, P., Cardarelli, M., 2018. A Newly
Identified Flower-Specific Splice Variant of AUXIN RESPONSE FACTOR8 Regulates Stamen
Elongation and Endothecium Lignification in Arabidopsis. Plant Cell 30, 620-637.

Gliwicka, M., Nowak, K., Balazadeh, S., Mueller-Roeber, B., Gaj, M.D., 2013. Extensive
modulation of the transcription factor transcriptome during somatic embryogenesis in Arabidopsis
thaliana. PLoS One 8, e69261.

Gonzalez, J.H., Taylor, J.S., Reed, K.M., Wright, R.C., Bargmann, B.O., 2021. Temporal control
of morphogenic factor expression determines efficacy in enhancing regeneration. Plants 10, 2271.

Guilfoyle, T.J., 2015. The PB1 domain in auxin response factor and Aux/IAA proteins: a versatile
protein interaction module in the auxin response. Plant Cell 27, 33-43.

Guo, F., Wang, H., Lian, G., Cai, G., Liu, W., Zhang, H., Li, D., Zhou, C., Han, N., Zhu, M., Su,
Y., Seo, P.J., Xu, L., Bian, H., 2023a. Initiation of scutellum-derived callus is regulated by an
embryo-like developmental pathway in rice. Communications Biology 6, 457.

Guo, X., Wang, T., Jiang, L., Qi, H., Zhang, Z., 2023b. PlaASDB: a comprehensive database of
plant alternative splicing events in response to stress. BMC plant biology 23, 225.

Harding, E.W., Tang, W., Nichols, K.W., Fernandez, D.E., Perry, S.E., 2003. Expression and
maintenance of embryogenic potential is enhanced through constitutive expression of
AGAMOUS-Like 15. Plant Physiol 133, 653-663.

Hardtke, C.S., Ckurshumova, W., Vidaurre, D.P., Singh, S.A., Stamatiou, G., Tiwari, S.B., Hagen,

G., Guilfoyle, T.J., Berleth, T., 2004. Overlapping and non-redundant functions of the Arabidopsis

auxin response factors MONOPTEROS and NONPHOTOTROPIC HYPOCOTYL 4.

Development 131, 1089-1100.

Hasnain, A., Naqvi, S.A.H., Ayesha, S.I., Khalid, F., Ellahi, M., Iqbal, S., Hassan, M.Z., Abbas,

A., Adamski, R., Markowska, D., 2022. Plants in vitro propagation with its applications in food,

pharmaceuticals and cosmetic industries; current scenario and future approaches. Frontiers in plant

science 13, 1009395.

Hu, Y., Zhou, L., Huang, M., He, X., Yang, Y., Liu, X., Li, Y., Hou, X., 2018. Gibberellins play

an essential role in late embryogenesis of Arabidopsis. Nature plants 4, 289-298.

Indoliya, Y., Tiwari, P., Chauhan, A.S., Goel, R., Shri, M., Bag, S.K., Chakrabarty, D., 2016.

Decoding regulatory landscape of somatic embryogenesis reveals differential regulatory networks

between japonica and indica rice subspecies. Sci Rep 6, 23050.

Juárez-González, V.T., López-Ruiz, B.A., Baldrich, P., Luján-Soto, E., Meyers, B.C., Dinkova,

T.D., 2019. The explant developmental stage profoundly impacts small RNA-mediated regulation

at the dedifferentiation step of maize somatic embryogenesis. Sci Rep 9, 14511.

Junker, A., Mönke, G., Rutten, T., Keilwagen, J., Seifert, M., Thi, T.M., Renou, J.P., Balzergue,

S., Viehöver, P., Hähnel, U., Ludwig-Müller, J., Altschmied, L., Conrad, U., Weisshaar, B.,

Bäumlein, H., 2012. Elongation-related functions of LEAFY COTYLEDON1 during the

development of Arabidopsis thaliana. Plant J 71, 427-442.

Kadokura, S., Sugimoto, K., Tarr, P., Suzuki, T., Matsunaga, S., 2018. Characterization of somatic

embryogenesis initiated from the Arabidopsis shoot apex. Dev Biol 442, 13-27.

Karami, O., Philipsen, C., Rahimi, A., Nurillah, A.R., Boutilier, K., Offringa, R., 2023.

Endogenous auxin maintains embryonic cell identity and promotes somatic embryo development

in Arabidopsis. Plant J 113, 7-22.

Karami, O., Rahimi, A., Mak, P., Horstman, A., Boutilier, K., Compier, M., van der Zaal, B., Offringa, R., 2021. An Arabidopsis AT-hook motif nuclear protein mediates somatic embryogenesis and coinciding genome duplication. Nat Commun 12, 2508.

Kashkan, I., Timofeyenko, K., Růžička, K., 2022. How alternative splicing changes the properties of plant proteins. Quant Plant Biol 3, e14.

Klepikova, A.V., Kasianov, A.S., Gerasimov, E.S., Logacheva, M.D., Penin, A.A., 2016. A high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. The Plant Journal 88, 1058-1070.

Korasick, D.A., Westfall, C.S., Lee, S.G., Nanao, M.H., Dumas, R., Hagen, G., Guilfoyle, T.J., Jez, J.M., Strader, L.C., 2014. Molecular basis for AUXIN RESPONSE FACTOR protein interaction and the control of auxin response repression. Proc Natl Acad Sci U S A 111, 5427-5432.

Krasteva, G., Georgiev, V., Pavlov, A., 2021. Recent applications of plant cell culture technology in cosmetics and foods. Engineering in Life Sciences 21, 68-76.

Kriechbaumer, V., Botchway, S.W., Hawes, C., 2016. Localization and interactions between Arabidopsis auxin biosynthetic enzymes in the TAA/YUC-dependent pathway. Journal of Experimental Botany 68, 4195-4207.

Krogan, N.T., Ckurshumova, W., Marcos, D., Caragea, A.E., Berleth, T., 2012. Deletion of MP/ARF5 domains III and IV reveals a requirement for Aux/IAA regulation in Arabidopsis leaf vascular patterning. New Phytol 194, 391-401.

Krogan, N.T., Marcos, D., Weiner, A.I., Berleth, T., 2016. The auxin response factor MONOPTEROS controls meristem function and organogenesis in both the shoot and root through the direct regulation of PIN genes. New Phytol 212, 42-50.

Kurihara, D., Mizuta, Y., Nagahara, S., Higashiyama, T., 2021. ClearSeeAlpha: advanced optical clearing for whole-plant imaging. Plant and Cell Physiology 62, 1302-1310.

Lau, S., De Smet, I., Kolb, M., Meinhardt, H., Jürgens, G., 2011. Auxin triggers a genetic switch.
Nat Cell Biol 13, 611-615.

Ledwoń, A., Gaj, M.D., 2009. LEAFY COTYLEDON2 gene expression and auxin treatment in
relation to embryogenic capacity of Arabidopsis somatic cells. Plant Cell Rep 28, 1677-1688.

Ledwoń, A., Gaj, M.D., 2011. LEAFY COTYLEDON1, FUSCA3 expression and auxin treatment
in relation to somatic embryogenesis induction in Arabidopsis. Plant Growth Regulation 65, 157-
167.

Lee, K., Kim, J.H., Park, O.S., Jung, Y.J., Seo, P.J., 2022. Ectopic expression of WOX5 promotes
cytokinin signaling and de novo shoot regeneration. Plant Cell Rep 41, 2415-2422.

Lee, K., Wang, K., 2023. Strategies for genotype-flexible plant transformation. Current Opinion in
Biotechnology 79, 102848.

Li, M., Wrobel-Marek, J., Heidmann, I., Horstman, A., Chen, B., Reis, R., Angenent, G.C.,
Boutilier, K., 2022. Auxin biosynthesis maintains embryo identity and growth during BABY
BOOM-induced somatic embryogenesis. Plant Physiol 188, 1095-1110.

Liu, L., Yahaya, B.S., Li, J., Wu, F., 2024. Enigmatic role of auxin response factors in plant growth
and stress tolerance. Frontiers in Plant Science 15, 1398818.

Liu, S., Zhang, Y., Feng, Q., Qin, L., Pan, C., Lamin-Samu, A.T., Lu, G., 2018a. Tomato AUXIN
RESPONSE FACTOR 5 regulates fruit set and development via the mediation of auxin and
gibberellin signaling. Sci Rep 8, 2971.

Liu, Z., Miao, L., Huo, R., Song, X., Johnson, C., Kong, L., Sundaresan, V., Yu, X., 2018b. ARF2–
ARF4 and ARF5 are essential for female and male gametophyte development in Arabidopsis. Plant
and Cell Physiology 59, 179-189.

Lotan, T., Ohto, M.-a., Yee, K.M., West, M.A., Lo, R., Kwong, R.W., Yamagishi, K., Fischer,
R.L., Goldberg, R.B., Harada, J.J., 1998. Arabidopsis LEAFY COTYLEDON1 is sufficient to
induce embryo development in vegetative cells. Cell 93, 1195-1205.

Ma, Y., Miotk, A., Šutiković, Z., Ermakova, O., Wenzl, C., Medzihradszky, A., Gaillochet, C., Forner, J., Utan, G., Brackmann, K., Galván-Ampudia, C.S., Vernoux, T., Greb, T., Lohmann, J.U., 2019. WUSCHEL acts as an auxin response rheostat to maintain apical stem cells in Arabidopsis. Nat Commun 10, 5093.

Ma, Y., Zhao, B., Zhang, D., Ouyang, K., Li, J., 2024. Transcriptomic and functional analyses identify genes associated with embryogenic callus formation in Neolamarckia cadamba. Industrial Crops and Products 216, 118686.

Martín, G., Márquez, Y., Mantica, F., Duque, P., Irimia, M., 2021. Alternative splicing landscapes in Arabidopsis thaliana across tissues and stress conditions highlight major functional differences with animals. Genome biology 22, 1-26.

Mattsson, J., Ckurshumova, W., Berleth, T., 2003. Auxin signaling in Arabidopsis leaf vascular development. Plant Physiol 131, 1327-1339.

McFarland, F.L., Kaeppler, H.F., 2024. History and current status of embryogenic culture-based tissue culture, transformation and gene editing of maize (Zea mays L.). The Plant Genome, e20451.

Mehbub, H., Akter, A., Akter, M.A., Mandal, M.S.H., Hoque, M.A., Tuleja, M., Mehraj, H., 2022. Tissue Culture in Ornamentals: Cultivation Factors, Propagation Techniques, and Its Application. Plants (Basel) 11.

Meinke, D.W., 2020. Genome-wide identification of EMBRYO-DEFECTIVE (EMB) genes required for growth and development in Arabidopsis. New Phytologist 226, 306-325.

Mira, M.M., Day, S., Ibrahim, S., Hill, R.D., Stasolla, C., 2023. The Arabidopsis Phytoglobin 2 mediates phytochrome B (phyB) light signaling responses during somatic embryogenesis. Planta 257, 88.

Möller, B.K., Ten Hove, C.A., Xiang, D., Williams, N., López, L.G., Yoshida, S., Smit, M., Datla, R., Weijers, D., 2017. Auxin response cell-autonomously controls ground tissue initiation in the early Arabidopsis embryo. Proc Natl Acad Sci U S A 114, E2533-e2539.

Murashige, T., Skoog, F., 1962. A Revised Medium for Rapid Growth and Bio Assays with
Tobacco Tissue Cultures. Physiologia Plantarum 15, 473-497.

Nanao, M.H., Vinos-Poyo, T., Brunoud, G., Thévenon, E., Mazzoleni, M., Mast, D., Lainé, S.,
Wang, S., Hagen, G., Li, H., Guilfoyle, T.J., Parcy, F., Vernoux, T., Dumas, R., 2014. Structural
basis for oligomerization of auxin transcriptional regulators. Nat Commun 5, 3617.

O'Malley, R.C., Huang, S.C., Song, L., Lewsey, M.G., Bartlett, A., Nery, J.R., Galli, M., Gallavotti,
A., Ecker, J.R., 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape.
Cell 165, 1280-1292.

Odat, O., Gardiner, J., Sawchuk, M.G., Verna, C., Donner, T.J., Scarpella, E., 2014.
Characterization of an allelic series in the MONOPTEROS gene of Arabidopsis. Genesis 52, 127-
133.

Paul, P., Joshi, S., Tian, R., Diogo Junior, R., Chakrabarti, M., Perry, S.E., 2022. The MADS-
domain factor AGAMOUS-Like18 promotes somatic embryogenesis. Plant Physiol 188, 1617-
1631.

Piya, S., Shrestha, S.K., Binder, B., Stewart, C.N., Jr., Hewezi, T., 2014. Protein-protein interaction
and gene co-expression maps of ARFs and Aux/IAAs in Arabidopsis. Front Plant Sci 5, 744.

Powers, S.K., Holehouse, A.S., Korasick, D.A., Schreiber, K.H., Clark, N.M., Jing, H., Emenecker,
R., Han, S., Tycksen, E., Hwang, I., Sozzani, R., Jez, J.M., Pappu, R.V., Strader, L.C., 2019.
Nucleo-cytoplasmic Partitioning of ARF Proteins Controls Auxin Responses in Arabidopsis
thaliana. Mol Cell 76, 177-190.e175.

Przemeck, G.K., Mattsson, J., Hardtke, C.S., Sung, Z.R., Berleth, T., 1996. Studies on the role of
the Arabidopsis gene MONOPTEROS in vascular development and plant cell axialization. Planta
200, 229-237.

Qi, S., Zhao, R., Yan, J., Fan, Y., Huang, C., Li, H., Chen, S., Zhang, T., Kong, L., Zhao, J., Zhang,
J., 2021. Global Transcriptome and Coexpression Network Analyses Reveal New Insights Into
Somatic Embryogenesis in Hybrid Sweetgum (Liquidambar styraciflua × Liquidambar
formosana). Front Plant Sci 12, 751866.

Quintana-Escobar, A.O., Nic-Can, G.I., Galaz Avalos, R.M., Loyola-Vargas, V.M., Gongora-
Castillo, E., 2019. Transcriptome analysis of the induction of somatic embryogenesis in Coffea
canephora and the participation of ARF and Aux/IAA genes. PeerJ 7, e7752.

Rademacher, E.H., Möller, B., Lokerse, A.S., Llavata-Peris, C.I., van den Berg, W., Weijers, D.,
2011. A cellular expression map of the Arabidopsis AUXIN RESPONSE FACTOR gene family.
Plant J 68, 597-606.

Robert, H.S., Grunewald, W., Sauer, M., Cannoot, B., Soriano, M., Swarup, R., Weijers, D.,
Bennett, M., Boutilier, K., Friml, J., 2015. Plant embryogenesis requires AUX/LAX-mediated
auxin influx. Development 142, 702-711.

Sakamoto, Y., Kawamura, A., Suzuki, T., Segami, S., Maeshima, M., Polyn, S., De Veylder, L.,
Sugimoto, K., 2022. Transcriptional activation of auxin biosynthesis drives developmental
reprogramming of differentiated cells. Plant Cell 34, 4348-4365.

Schlereth, A., Möller, B., Liu, W., Kientz, M., Flipse, J., Rademacher, E.H., Schmid, M., Jürgens,
G., Weijers, D., 2010. MONOPTEROS controls embryonic root initiation by regulating a mobile
transcription factor. Nature 464, 913-916.

Schulz, B., Segobye, K., 2016. 2, 4-D transport and herbicide resistance in weeds. Journal of
experimental botany 67, 3177-3179.

Shi, X., Zhang, C., Liu, Q., Zhang, Z., Zheng, B., Bao, M., 2016. De novo comparative
transcriptome analysis provides new insights into sucrose induced somatic embryogenesis in
camphor tree (Cinnamomum camphora L.). BMC Genomics 17, 26.

Stone, S.L., Braybrook, S.A., Paula, S.L., Kwong, L.W., Meuser, J., Pelletier, J., Hsieh, T.F.,
Fischer, R.L., Goldberg, R.B., Harada, J.J., 2008. Arabidopsis LEAFY COTYLEDON2 induces
maturation traits and auxin activity: Implications for somatic embryogenesis. Proc Natl Acad Sci
U S A 105, 3151-3156.

Stone, S.L., Kwong, L.W., Yee, K.M., Pelletier, J., Lepiniec, L., Fischer, R.L., Goldberg, R.B.,
Harada, J.J., 2001. LEAFY COTYLEDON2 encodes a B3 domain transcription factor that induces
embryo development. Proc Natl Acad Sci U S A 98, 11806-11811.

Su, Y.H., Liu, Y.B., Bai, B., Zhang, X.S., 2015. Establishment of embryonic shoot–root axis is
involved in auxin and cytokinin response during Arabidopsis somatic embryogenesis. Frontiers in
plant science 5, 123838.

Su, Y.H., Zhao, X.Y., Liu, Y.B., Zhang, C.L., O'Neill, S.D., Zhang, X.S., 2009. Auxin-induced
WUS expression is essential for embryonic stem cell renewal during somatic embryogenesis in
Arabidopsis. Plant J 59, 448-460.

Szécsi, J., Joly, C., Bordji, K., Varaud, E., Cock, J.M., Dumas, C., Bendahmane, M., 2006.
BIGPETALp, a bHLH transcription factor is involved in the control of Arabidopsis petal size. The
EMBO journal 25, 3912-3920.

Szyrajew, K., Bielewicz, D., Dolata, J., Wójcik, A.M., Nowak, K., Szczygieł-Sommer, A.,
Szweykowska-Kulinska, Z., Jarmolowski, A., Gaj, M.D., 2017. MicroRNAs Are Intensively
Regulated during Induction of Somatic Embryogenesis in Arabidopsis. Front Plant Sci 8, 18.

Thakare, D., Tang, W., Hill, K., Perry, S.E., 2008. The MADS-domain transcriptional regulator
AGAMOUS-LIKE15 promotes somatic embryo development in Arabidopsis and soybean. Plant
physiology 146, 1663-1672.

Thellin, O., Zorzi, W., Lakaye, B., De Borman, B., Coumans, B., Hennen, G., Grisar, T., Igout, A.,
Heinen, E., 1999. Housekeeping genes as internal standards: use and limits. Journal of
biotechnology 75, 291-295.

Thibaud-Nissen, F., Shealy, R.T., Khanna, A., Vodkin, L.O., 2003. Clustering of microarray data
reveals transcript patterns associated with somatic embryogenesis in soybean. Plant Physiol 132,
118-136.

Tsugafune, S., Mashiguchi, K., Fukui, K., Takebayashi, Y., Nishimura, T., Sakai, T., Shimada, Y.,
Kasahara, H., Koshiba, T., Hayashi, K.-i., 2017. Yucasin DF, a potent and persistent inhibitor of
auxin biosynthesis in plants. Scientific Reports 7, 13992.

Tvorogova, V.E., Fedorova, Y.A., Potsenkovskaya, E.A., Kudriashov, A.A., Efremova, E.P.,
Kvitkovskaya, V.A., Wolabu, T.W., Zhang, F., Tadege, M., Lutova, L.A., 2019. The WUSCHEL-

related homeobox transcription factor MtWOX9-1 stimulates somatic embryogenesis in Medicago
truncatula. Plant Cell, Tissue and Organ Culture (PCTOC) 138, 517-527.

Uc-Chuc, M.A., Pérez-Hernández, C., Galaz-Ávalos, R.M., Brito-Argaez, L., Aguilar-Hernández,
V., Loyola-Vargas, V.M., 2020. YUCCA-Mediated Biosynthesis of the Auxin IAA Is Required
during the Somatic Embryogenic Induction Process in Coffea canephora. Int J Mol Sci 21.

Varaud, E., Brioudes, F., Szécsi, J., Leroux, J., Brown, S., Perrot-Rechenmann, C., Bendahmane,
M., 2011. AUXIN RESPONSE FACTOR8 regulates Arabidopsis petal growth by interacting with
the bHLH transcription factor BIGPETALp. Plant Cell 23, 973-983.

Wang, F.X., Shang, G.D., Wu, L.Y., Xu, Z.G., Zhao, X.Y., Wang, J.W., 2020. Chromatin
Accessibility Dynamics and a Hierarchical Transcriptional Regulatory Network Structure for Plant
Somatic Embryogenesis. Dev Cell 54, 742-757.e748.

Wang, K., Gao, E., Liu, D., Wu, X., Wang, P., 2022. The ER network, peroxisomes and actin
cytoskeleton exhibit dramatic alterations during somatic embryogenesis of cultured citrus cells.
Plant Cell, Tissue and Organ Culture (PCTOC) 148, 259-270.

Wickramasuriya, A.M., Dunwell, J.M., 2015. Global scale transcriptome analysis of Arabidopsis
embryogenesis in vitro. BMC Genomics 16, 301.

Wójcik, A.M., Wójcikowska, B., Gaj, M.D., 2020. Current Perspectives on the Auxin-Mediated
Genetic Network that Controls the Induction of Somatic Embryogenesis in Plants. Int J Mol Sci
21.

Wójcikowska, B., Belaidi, S., Robert, H.S., 2023. Game of thrones among AUXIN RESPONSE
FACTORs—over 30 years of MONOPTEROS research. Journal of Experimental Botany 74, 6904-
6921.

Wójcikowska, B., Botor, M., Morończyk, J., Wójcik, A.M., Nodzyński, T., Karcz, J., Gaj, M.D.,
2018. Trichostatin A Triggers an Embryogenic Transition in Arabidopsis Explants via an Auxin-
Related Pathway. Front Plant Sci 9, 1353.

Wójcikowska, B., Gaj, M.D., 2015. LEAFY COTYLEDON2-mediated control of the endogenous

hormone content: implications for the induction of somatic embryogenesis in Arabidopsis. Plant

Cell, Tissue and Organ Culture (PCTOC) 121, 255-258.

Wójcikowska, B., Gaj, M.D., 2017. Expression profiling of AUXIN RESPONSE FACTOR genes

during somatic embryogenesis induction in Arabidopsis. Plant Cell Rep 36, 843-858.

Wójcikowska, B., Jaskóła, K., Gąsiorek, P., Meus, M., Nowak, K., Gaj, M.D., 2013. LEAFY

COTYLEDON2 (LEC2) promotes embryogenic induction in somatic tissues of Arabidopsis, via

YUCCA-mediated auxin biosynthesis. Planta 238, 425-440.

Wu, L.Y., Shang, G.D., Wang, F.X., Gao, J., Wan, M.C., Xu, Z.G., Wang, J.W., 2022. Dynamic

chromatin state profiling reveals regulatory roles of auxin and cytokinin in shoot regeneration. Dev

Cell 57, 526-542.e527.

Wu, M.F., Yamaguchi, N., Xiao, J., Bargmann, B., Estelle, M., Sang, Y., Wagner, D., 2015. Auxin-

regulated chromatin switch directs acquisition of flower primordium founder fate. Elife 4, e09269.

Xie, M., Huang, L., Song, L., O'Neil, R., Lewsey, M., Chen, H., Chen, H., Zhuo, R., Shokhirev,

M., Alonso, J., Ecker, J., 2022. Defining in vivo transcriptional responses to auxin. Research

Square.

Xu, C., Shen, Y., He, F., Fu, X., Yu, H., Lu, W., Li, Y., Li, C., Fan, D., Wang, H.C., 2019. Auxin-

mediated Aux/IAA-ARF-HB signaling cascade regulates secondary xylem development in

Populus. New Phytologist 222, 752-767.

Xu, L., Liu, Y., Zhang, J., Wu, W., Hao, Z., He, S., Li, Y., Shi, J., Chen, J., 2024. Genomic survey

and expression analysis of LcARFs reveal multiple functions to somatic embryogenesis in

Liriodendron. BMC Plant Biology 24, 94.

Yang, X., Lin, Q., Udayabhanu, J., Hua, Y., Dai, X., Xin, S., Wang, X., Huang, H., Huang, T.,

2024. An optimized somatic embryo transformation system assisted homozygous edited rubber

tree generation method mediated by CRISPR/Cas9. bioRxiv, 2024.2003. 2014.585007.

Yue, P., Lu, Q., Liu, Z., Lv, T., Li, X., Bu, H., Liu, W., Xu, Y., Yuan, H., Wang, A., 2020. Auxin-

activated MdARF5 induces the expression of ethylene biosynthetic genes to initiate apple fruit

ripening. New phytologist 226, 1781-1795.

Zeng, J., Deng, Y., Iqbal, S., Zhang, J., Wu, K., Ma, G., Li, L., Dai, G., Deng, R., Fang, L., Zeng,

S., 2024. Direct somatic embryogenesis and related gene expression networks in leaf explants of

Hippeastrum 'Bangkok Rose'. Horticultural Plant Journal 10, 556-572.

Zhang, M.M., Zhang, H.K., Zhai, J.F., Zhang, X.S., Sang, Y.L., Cheng, Z.J., 2021. ARF4 regulates

shoot regeneration through coordination with ARF5 and IAA12. Plant Cell Rep 40, 315-325.

Zheng, Q., Perry, S.E., 2014. Alterations in the transcriptome of soybean in response to enhanced

somatic embryogenesis promoted by orthologs of Agamous-like15 and Agamous-like18. Plant

Physiol 164, 1365-1377.

Zhu, X., Xu, Z., Wang, G., Cong, Y., Yu, L., Jia, R., Qin, Y., Zhang, G., Li, B., Yuan, D., Tu, L.,

Yang, X., Lindsey, K., Zhang, X., Jin, S., 2023. Single-cell resolution analysis reveals the

preparation for reprogramming the fate of stem cell niche in cotton lateral meristem. Genome Biol

24, 194.

Zuo, J., Niu, Q.W., Frugis, G., Chua, N.H., 2002. The WUSCHEL gene promotes vegetative-to-

embryonic transition in Arabidopsis. Plant J 30, 349-359.

1026

1027   **Figure legend**

1028

Figure 1. *MP11ir* is expressed during somatic embryogenesis induction.

(**A**) The level of *MP11ir* transcript in IZE explants and during auxin-dependent (5.0 µM 2,4-D) SE

induction (5th and 10th day of *in vitro* culture). Values significantly different from the control

culture (10-day-old seedlings) are marked with an asterisk. The relative expression level was

normalized to internal control (At4g27090) and calibrated to the 10-day-old seedling. A two-way

ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were

significantly different from 10-day-old seedlings (*$p < 0.05$; **$p < 0.01$); (n = 3; means ± SD are

given). (**B-P**) The level of *MP11ir* transcript in the IZE explants that had been cultured for ten days

on medium without exogenous auxin, control condition (**E, I, M**), and treated with different 2,4-D

(**B**), NAA (**C**) and IAA (**D**) concentrations 2.5 (**F, J, N**), 5.0 (**G, K, O**), and 7.5 µM (**H, L, P**).

Values significantly different from the control culture (explants untreated with 2,4-D, NAA, or

IAA) are marked with an asterisk. The relative expression level was normalized to internal control

(At4g27090) and calibrated to the 10-day-old IZE control culture untreated with exogenous auxin.

A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any

values that were significantly different from untreated IZE explants (*$p < 0.05$; **$p < 0.01$);(n = 3;

means ± SD are given). (**Q**) The *MP11ir* transcript level in IZE explants and during TSA-dependent

(1.0 µM TSA) SE induction (5th and 10th day of *in vitro* culture). Values significantly different

from the control culture (10-day-old seedlings) are marked with an asterisk. A two-way ANOVA

analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were

significantly different from 10-day-old seedlings (*$p < 0.05$; **$p < 0.01$);(n = 3; means ± SD are

given). (**R**) Together with data presented in Supplementary Figure S2, we hypothesize a positive

correlation between *MP11ir* levels and SE capacity. Size bars indicate 1 mm (**E-P**).

1051

Figure 2. Both MP/ARF5 and MP11ir are required to rescue the reduced SE capacity of

mpS319.

(**A-M**) The embryogenic capacity (**A, B**) of the Col-0 (**C, H**), *mpS319* mutant (**D, I, J**),

MP::MP11ir (**E, K**), *MP::MP* (**F, L**) and *MP::MP11ir MP::MP* (**G, M**) transgenic lines in *mpS319*

background on the E5 medium evaluated in a 21-day-old culture. Values that were significantly

different from the control culture (Col-0) are marked with an asterisk, and from *mpS319* mutant

are marked with a hashtag (n ≥ 3; means ± SD are given). A two-way ANOVA analysis ($p < 0.05$)

followed by Tukey's HSD was used to determine any values that were significantly different (*,#p

< 0.05; **,##$p < 0.01$); (n ≥ 3; means ± SD are given). (**N-X**) The embryogenic capacity (**N, O**) of

the Col-0 (**P, S, V**), *35S::bdl-GR* (**Q, T, W**), *35S::BDL-GR* (**R, U, X**) transgenic lines on the E5

medium (**P, Q, R**) and E5 medium supplemented with 10 µM (**S, T, U**) and 30 µM (**V, W, X**) of

DEX. SE capacity was evaluated in 21-day-old cultures. Values significantly different from the

control culture (Col-0) untreated with DEX are marked with an asterisk (n ≥ 3; means ± SD are

given). A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine

any values that were significantly different (*$p < 0.05$; **$p < 0.01$); (n = 3; means ± SD are given).

Size bars indicate 1 mm for explant images and 1 cm for plate images.

1068

Figure 3. Expression pattern of MP11ir during SE induction

1069

1070 Spatio-temporal localization of MP11ir in IZE explants (**A**), explants cultured on the SE induction

1071 medium with 5 μM 2,4-D by 1 (**B**), and 5 (**C**) days of SE culture. d, day of SE culture. IZE explants

1072 were cultured from *pMP::MP11ir-GFP (MR)* transgenic lines. The GFP fluorescence is shown in

1073 green. Size bars indicate 200 μm (**A-C**).

1074

Figure 4. Overexpressing *ΔARF5* impairs SE induction

1075

(**A, B**) The embryogenic capacity of *XVE::ΔARF5 DR5::GFP* explants treated with different 2,4-

D concentrations evaluated in a 21-day-old culture. The significantly impaired SE efficiency (**A**)

and productivity (**B**) of the *ΔARF5*-expressing explants are presented. (**C**) Explants overexpressing

ΔARF5 and cultured on auxin-media could not regenerate somatic embryos and produced callus.

ΔARF5 overexpression was induced with 1 μM estradiol (+E). Values that were significantly

different from the control culture (Col-0) are marked with an asterisk (n ≥ 3; means ± SD are given).

A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any

values that were significantly different (*$p < 0.05$); (n = 3; means ± SD are given). Size bars

indicate 1 mm for explant images and 1 cm for plate images.

1085

Figure 5. The expression levels of *TAA1*, *TAR1*, *YUC3*, *YUC5*, and *YUC8* are altered when

1086

MP-dependent transcriptional activities are impaired.

(**A-F**) Relative expression level of *TAA1*, *TAR1*, *YUC3*, *YUC5*, and *YUC8* auxin biosynthetic genes

in the 1- (**A**), 5- (**B**), and 10-day-old (**C**) embryogenic cultures of the *XVE::ΔARF5 DR5::GFP*

transgenic line treated with 1 μM β-estradiol (+E), 5- and 10-day-old embryogenic cultures of the

35S::bdl-GR transgenic line treated with 10 μM dexamethasone (+DEX) (**D**), and 0- (**E**) and 10-

day-old (**F**) embryogenic cultures of *mp/arf5* mutant. The relative expression level was normalized

to internal control (At4g27090) and calibrated to the *XVE::ΔARF5 DR5::GFP* culture of the same

age and untreated with 1 μM β-estradiol (-E) (**A-C**) or *35S::bdl-GR* culture of the same age and

untreated with 10 μM dexamethasone (-DEX) (**D**), or Col-0 embryogenic culture of the same age

(**E, F**). Value significantly different from the *XVE::ΔARF5 DR5::GFP* (-E) (**A-C**), *35S::bdl-GR* (-

DEX) (**D**) or Col-0 (**E, F**) culture of the same age are marked with an asterisk (*$p < 0.05$; n = 3 ±

SD).

1099

Figure 6. The expression pattern of TAA1, TAR1, YUC3, YUC5, and YUC8 is altered when MP-dependent transcriptional activities are impaired.

Spatio-temporal localization of TAA1, TAR1, YUC3, YUC5, and YUC8 enzymes during SE process under *in vitro* culture in *XVE::ΔARF5 DR5::GFP* (**A**) *35S::bdl-GR* (**B**) or *mp/arf5* background (**C**). IZE explants of reporter lines were cultured on an E5 medium as a control (**A, B, C**), E5 medium with 1 µM β-estradiol for *ΔARF5* overexpression (+E) (**A**), E5 medium with 10 µM dexamethasone (+DEX) (**B**). The tissue was sampled on the 5- and 10-day-old *in vitro* cultures (**A, B**), and in IZEs, explants on the 5- and 10-day-old *in vitro* cultures (**C**). The expression is visible in blue from *pTAA1::TAA1-GUS*, *pTAR1::TAR1-GUS*, *pYUC3::YUC3-GUS*, *pYUC5::YUC5-GUS*, and *pYUC8::YUC8-GUS* reporter lines crossed into the respective genetic backgrounds. Size bars indicate 200 µm.

1111

Figure 7. Overexpressing *ΔARF5* increases endogenous auxin levels.

(**A-D**) GFP-monitored auxin maxima in *XVE::ΔARF5 DR5::GFP* explants cultured for one (**A, B**) and five days (**C, D**) on an ET medium (1.0 µM TSA) without (-E) (**A, C**) or with β-estradiol (+E) (**B, D**). The GFP fluorescence signal is displayed in green. (**E, F**) The fluorescence signal was quantified in one- (**E**) and five-day-old (**F**) explants. (**G**) Level of indolic compounds (µg/g of fresh tissue) in 3-week-old IZE-culture of *XVE::ΔARF5 DR5::GFP* transgenic line with 1 µM β-estradiol-induced *ΔARF5* overexpression. A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different (*$p < 0.05$; **$p < 0.01$). Size bars indicate 200 µm (**A-D**) and 1 mm (**G**).

1121

Figure 8. Proposed model of MP/ARF5 and MP11ir action during embryogenic transition through control of auxin-biosynthesis genes expression.

(**A**) The canonical MP/ARF5 protein and MP11ir isoform are required for the efficient embryogenic transition via controlling auxin biosynthesis genes. (**B**) Overexpression of truncated *ΔARF5*, mimicking MP11ir, activates expression of *TAA1 TAR1*, *YUC3*, *YUC5,* and *YUC8,* which disrupts auxin homeostasis, leads to suboptimal endogenous auxin levels, inhibits the SE process, and induces the formation of callus. (**C**) Blocking MP/ARF5 action through bdl repressor results in decreased *YUC3* and *YUC8* levels during SE induction, inhibits of SE process, and induces

seedling development. (**D**) The lack of MP/ARF5 and MP11ir in *mp/arf5* mutant leads to the

absence of *YUC3*, *YUC5*, and *YUC8* expression during SE induction and inhibits of SE process,

callus formation, or seedling development. AuxRE – Auxin Response Elements; BRM/SYD –

BRAHMA/SPLAYED; MP/ARF5 – MONOPTEROS; MP11ir – MP/ARF5 isoform; TFs -

TRANSCRIPTION FACTOR; HAT – HISTONE ACETYLASES; TAA1 - TRYPTOPHAN

AMINOTRANSFERASE   OF   ARABIDOPSIS   1,   TAR1   -   TRYPTOPHAN

AMINOTRANSFERASE   RELATED   1;   TPL/TPR   –   TOPLESS/TPL-RELATED;   YUC   –

YUCCA; Ac – acetyl group; SE – somatic embryogenesis. Created with BioRender.com.

1138

Figure 1. *MP11ir* is expressed during somatic embryogenesis induction.
(**A**) The level of *MP11ir* transcript in IZE explants and during auxin-dependent (5.0 µM 2,4-D) SE induction (5th and 10th day of *in vitro* culture). Values significantly different from the control culture (10-day-old seedlings) are marked with an asterisk. The relative expression level was normalized to internal control (At4g27090) and calibrated to the 10-day-old seedling. A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different from 10-day-old seedlings (*$p < 0.05$; **$p < 0.01$); (n = 3; means ± SD are given). (**B-P**) The level of *MP11ir* transcript in the IZE explants that had been cultured for ten days on medium without exogenous auxin, control condition (**E, I, M**), and treated with different 2,4-D (**B**), NAA (**C**) and IAA (**D**) concentrations 2.5 (**F, J, N**), 5.0 (**G, K, O**), and 7.5 µM (**H, L, P**). Values significantly different from the control culture (explants untreated with 2,4-D, NAA, or IAA) are marked with an asterisk. The relative expression level was normalized to internal control (At4g27090) and calibrated to the 10-day-old IZE control culture untreated with exogenous auxin. A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different from untreated IZE explants (*$p < 0.05$; **$p < 0.01$);(n = 3; means ± SD are given).

(**Q**) The *MP1Iir* transcript level in IZE explants and during TSA-dependent (1.0 µM TSA) SE induction (5th and 10th day of *in vitro* culture). Values significantly different from the control culture (10-day-old seedlings) are marked with an asterisk. A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different from 10-day-old seedlings (*$p < 0.05$; **$p < 0.01$);(n = 3; means ± SD are given). (**R**) Together with data presented in Supplementary Figure S2, we hypothesize a positive correlation between *MP1Iir* levels and SE capacity. Size bars indicate 1 mm (**E-P**).

Figure 2. Both MP/ARF5 and MP11ir are required to rescue the reduced SE capacity of *mpS319*.

(**A-M**) The embryogenic capacity (**A, B**) of the Col-0 (**C, H**), *mpS319* mutant (**D, I, J**), *MP::MP11ir* (**E, K**), *MP::MP* (**F, L**) and *MP::MP11ir MP::MP* (**G, M**) transgenic lines in *mpS319* background on the E5 medium evaluated in a 21-day-old culture. Values that were significantly different from the control culture (Col-0) are marked with an asterisk, and from *mpS319* mutant are marked with a hashtag (n ≥ 3; means ± SD are given). A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different (*, #$p < 0.05$; **, ##$p < 0.01$); (n ≥ 3; means ± SD are given). (**N-X**) The embryogenic capacity (**N, O**) of the Col-0 (**P, S, V**), *35S::bdl-GR* (**Q, T, W**), *35S::BDL-GR* (**R, U, X**) transgenic lines on the E5 medium (**P, Q, R**) and E5 medium supplemented with 10 µM (**S, T, U**) and 30 µM (**V, W, X**) of DEX. SE capacity was evaluated in 21-day-old cultures. Values significantly different from the control culture (Col-0) untreated with DEX are marked with an asterisk (n ≥ 3; means ± SD are given). A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different (*$p < 0.05$; **$p < 0.01$); (n = 3; means ± SD are given). Size bars indicate 1 mm for explant images and 1 cm for plate images.

Figure 3. Expression pattern of MP11ir during SE induction

Spatio-temporal localization of MP11ir in IZE explants (**A**), explants cultured on the SE induction medium with 5 μM 2,4-D by 1 (**B**), and 5 (**C**) days of SE culture. d, day of SE culture. IZE explants were cultured from *pMP::MP11ir-GFP (MR)* transgenic lines. The GFP fluorescence is shown in green. Size bars indicate 200 μm (**A-C**).

Figure 4. Overexpressing *ΔARF5* impairs SE induction.
(**A, B**) The embryogenic capacity of *XVE::ΔARF5 DR5::GFP* explants treated with different 2,4-D concentrations evaluated in a 21-day-old culture. The significantly impaired SE efficiency (**A**) and productivity (**B**) of the *ΔARF5*-expressing explants are presented. (**C**) Explants overexpressing *ΔARF5* and cultured on auxin-media could not regenerate somatic embryos and produced callus. *ΔARF5* overexpression was induced with 1 µM estradiol (+E). Values that were significantly different from the control culture (Col-0) are marked with an asterisk (n ≥ 3; means ± SD are given). A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different (*$p < 0.05$); (n = 3; means ± SD are given). Size bars indicate 1 mm for explant images and 1 cm for plate images.

Figure 5. The expression levels of *TAA1*, *TAR1*, *YUC3*, *YUC5*, and *YUC8* are altered when MP-dependent transcriptional activities are impaired.

(**A-F**) Relative expression level of *TAA1*, *TAR1*, *YUC3*, *YUC5,* and *YUC8* auxin biosynthetic genes in the 1- (**A**), 5- (**B**), and 10-day-old (**C**) embryogenic cultures of the *XVE::ΔARF5 DR5::GFP* transgenic line treated with 1 µM β-estradiol (+E), 5- and 10-day-old embryogenic cultures of the *35S::bdl-GR* transgenic line treated with 10 µM dexamethasone (+DEX) (**D**), and 0- (**E**) and 10-day-old (**F**) embryogenic cultures of *mp/arf5* mutant. The relative expression level was normalized to internal control (At4g27090) and calibrated to the *XVE::ΔARF5 DR5::GFP* culture of the same age and untreated with 1 µM β-estradiol (-E) (**A-C**) or *35S::bdl-GR* culture of the same age and untreated with 10 µM dexamethasone (-DEX) (**D**), or Col-0 embryogenic culture of the same age (**E, F**). Value significantly different from the *XVE::ΔARF5 DR5::GFP* (-E) (**A-C**), *35S::bdl-GR* (-DEX) (**D**) or Col-0 (**E, F**) culture of the same age are marked with an asterisk (*$p < 0.05$; n = 3 ± SD).

Figure 6. The expression pattern of TAA1, TAR1, YUC3, YUC5, and YUC8 is altered when MP-dependent transcriptional activities are impaired.

Spatio-temporal localization of TAA1, TAR1, YUC3, YUC5, and YUC8 enzymes during SE process under *in vitro* culture in *XVE::ΔARF5 DR5::GFP* (**A**) *35S::bdl-GR* (**B**) or *mp/arf5* background (**C**). IZE explants of reporter lines were cultured on an E5 medium as a control (**A, B, C**), E5 medium with 1 μM β-estradiol for *ΔARF5* overexpression (+E) (**A**), E5 medium with 10 μM dexamethasone (+DEX) (**B**). The tissue was sampled on the 5- and 10-day-old *in vitro* cultures (**A, B**), and in IZEs, explants on the 5- and 10-day-old *in vitro* cultures (**C**). The expression is visible in blue from *pTAA1::TAA1-GUS*, *pTAR1::TAR1-GUS*, *pYUC3::YUC3-GUS*, *pYUC5::YUC5-GUS*, and *pYUC8::YUC8-GUS* reporter lines crossed into the respective genetic backgrounds. Size bars indicate 200 μm.

Figure 7. Overexpressing *ΔARF5* increases endogenous auxin levels.
(**A-D**) GFP-monitored auxin maxima in *XVE::ΔARF5 DR5::GFP* explants cultured for one (**A, B**) and five days (**C, D**) on an ET medium (1.0 μM TSA) without (-E) (**A, C**) or with β-estradiol (+E) (**B, D**). The GFP fluorescence signal is displayed in green. (**E, F**) The fluorescence signal was quantified in one- (**E**) and five-day-old (**F**) explants. (**G**) Level of indolic compounds (μg/g of fresh tissue) in 3-week-old IZE-culture of *XVE::ΔARF5 DR5::GFP* transgenic line with 1 μM β-estradiol-induced *ΔARF5* overexpression. A two-way ANOVA analysis ($p < 0.05$) followed by Tukey's HSD was used to determine any values that were significantly different (*$p < 0.05$; **$p < 0.01$). Size bars indicate 200 μm (**A-D**) and 1 mm (**G**).

Figure 8. Proposed model of MP/ARF5 and MP11ir action during embryogenic transition through control of auxin-biosynthesis genes expression.
(**A**) The canonical MP/ARF5 protein and MP11ir isoform are required for the efficient embryogenic transition via controlling auxin biosynthesis genes. (**B**) Overexpression of truncated *ΔARF5*, mimicking MP11ir, activates expression of *TAA1 TAR1*, *YUC3*, *YUC5*, and *YUC8*, which disrupts auxin homeostasis, leads to suboptimal endogenous auxin levels, inhibits the SE process, and induces the formation of callus. (**C**) Blocking MP/ARF5 action

through bdl repressor results in decreased *YUC3* and *YUC8* levels during SE induction, inhibits of SE process, and induces seedling development. (**D**) The lack of MP/ARF5 and MP11ir in *mp/arf5* mutant leads to the absence of *YUC3*, *YUC5*, and *YUC8* expression during SE induction and inhibits of SE process, callus formation, or seedling development. AuxRE – Auxin Response Elements; BRM/SYD – BRAHMA/SPLAYED; MP/ARF5 – MONOPTEROS; MP11ir – MP/ARF5 isoform; TFs - TRANSCRIPTION FACTOR; HAT – HISTONE ACETYLASES; TAA1 - TRYPTOPHAN AMINOTRANSFERASE OF ARABIDOPSIS 1, TAR1 - TRYPTOPHAN AMINOTRANSFERASE RELATED 1; TPL/TPR – TOPLESS/TPL-RELATED; YUC – YUCCA; Ac – acetyl group; SE – somatic embryogenesis. Created with BioRender.com.