Cryptic Cryptophytes – revision of the genus *Goniomonas*

Maria Sachs[1], Frank Nitsche[1], Hartmut Arndt[1]*

[1]University of Cologne, Institute of Zoology, Department of General Ecology and

Limnology, Zuelpicher Straße 47b, 50674 Cologne, Germany

***corresponding author:**

H. Arndt, Institute of Zoology, University of Cologne, Zuelpicher Straße 47b, 50674

Cologne, Germany. Telephone number: +49 221 470 3100; email: Hartmut.Arndt@uni-

koeln.de

Abstract

Cryptomonad protists are ubiquitously distributed over marine and freshwater habitats. As an
exception to the coloured cryptomonads, the heterotrophic cryptomonads of the genus
Goniomonas have an ancestral phylogenetic position. They lack any kind of chloroplast and
most likely represent a basal group to those cryptomonad groups having obtained their
chloroplast by secondary endosymbiosis. Earlier studies have shown a deep divergence
between freshwater and marine clades of goniomonads that comprise large genetic distances
between members within the group and also between the two groups of marine and freshwater
taxa. Still, marine and freshwater species carry the same genus name, and until up to date only
few species have been described. We therefore restructured goniomonad systematics based not
only on a separation of marine and freshwater taxa, but, taking the large genetic distances into
account, also several new genera are described. Based on morphological as well as
phylogenetic data (18S rDNA sequences), this leads to the formation of the freshwater genera
Limnogoniomonas n. g., *Goniomonas* and *Aquagoniomonas* n. g. and the marine genera
Neptunogoniomonas n. g., *Marigoniomonas* n. g., *Thalassogoniomonas* n. g.,
Poseidogoniomonas n. g. and *Cosmogoniomonas* n. g. To give the restructuring process a stable
basis, we additionally propose a neotype for *Goniomonas truncata*.

Keywords: Cryptophyta; Cryptista; heterotrophic flagellates; n. g.; n. sp.; salinity;

INTRODUCTION

Cryptomonad flagellates are small, mostly bean shaped, non-colonial protists that encompass at least 200 species (Novarino, 2003). The group is globally distributed in marine, brackish and freshwater habitats (Archibald, 2020). Cryptomonads played a crucial role in understanding evolutionary processes as phototrophic representatives are the outcome of secondary endosymbiosis (Cenci et al., 2018; McFadden, 2018). Phototrophic cryptomonads had engulfed red algae which themselves were originally created through the engulfment of a cyanobacterium by a eukaryotic phagotroph during primary endosymbiosis (Keeling, 2004). There is one prominent exception - the cryptomonad genus *Goniomonas*. When analysing sediment associated protists, this genus is among the 20 most commonly observed heterotrophic flagellate morphotypes (Patterson and Lee, 2000) and have been recorded in many different habitats (Arndt et al., 2000). Goniomonads do neither possess plastids nor a nucleomorph. For a long time, it was not clear whether the genus has never been involved in the endosymbiosis and would therefore represent a very basal, ancestral heterotrophic cryptomonad or whether it had secondarily lost plastids. Cenci et al. (2018) sequenced the entire genome of the most recently described *Goniomonas* species *G. avonlea*. On gene level, no evidence could be found that *G. avonlea* ever possessed a plastid indicating that *Goniomonas* is indeed a very basal phagotroph, a "pre-secondary-endosymbiosis cryptomonad" (McFadden, 2018). Apart from the lack of plastids, all goniomonads share a number of morphological traits, such as a laterally flattened, anteriorly more or less truncated cell body and a horizontal row of ejectisomes in the anterior part of the cell (Novarino, 2003). Unlike other cryptomonads which were described to have a "recoiling" swimming behaviour (Novarino, 2003), goniomonads display a rather "jerky" movement (Kim and Archibald, 2013), and move on the substrate along irregular tracks while rotating. Until the taxon was transferred to the currently used name *Goniomonas* by Stein (1878), it was described under the protonym *Monas truncata* (Fresenius, 1858). Larsen and Patterson (1990) discussed the history of using the genus name in the past and described two new marine species. Up to now, five different species have been described, with *G. avonlea* (Kim and Archibald, 2013) being the most recent. Apart from the type species *G. truncata* and *G. brasiliensis* (Jarreta de Castro and Bicudo, 2023) which were isolated from freshwater, the three other species are marine species (*G. amphinema*, *G. pacifica,* see Larson and Patterson, 1990*)* and *G. avonlea* (Kim and Archibald, 2013). Molecular phylogenetic analyses revealed a deep divergence between the goniomonad species from marine and freshwaters (von der Heyden et al., 2004). Studies on other protist

groups have shown similar differences in ecological clusters together with molecular

differentiations, for example choanoflagellate taxa such as *Codosiga* and *Hartaetosiga* (Carr

et al., 2017). The high divergence among the genus *Goniomonas* could imply that there are still

multiple undescribed species and that the genus is generally under-split from a taxonomical

point of view (von der Heyden et al., 2004). The taxonomic situation becomes even more

complex since type material other than drafts of microscopic images are missing. Studies by

Shalchian-Tabrizi et al. (2008) and Shiratori and Ishida (2016) have already indicated a large

hidden diversity within cryptomonads, including heterotrophic lineages.

In this study, we describe ten new species from different aquatic habitats including marine,

brackish and freshwater environments and propose the revision of the genus *Goniomonas* into

Goniomonas sensu strictu, which includes several freshwater species and a newly created

neotype for *Goniomonas truncata*, as well as five new genera within both the freshwater and

marine clusters based mainly on their large genetic distances. Morphological descriptions are

supported by light and electron micrographs and genetic divergences based on 18S rDNA

phylogeny.

MATERIALS AND METHODS

Sediment and plankton samples were taken at diverse sampling sites (Table 1). Sediment

samples were taken with a Multi-Corer system (MUC), while plankton samples were either

taken with a CTD-rosette system from different depth or were taken with flasks from surface

waters. Subsamples of a few millilitres were suspended in ambient autoclaved water (approx.

20 ml) and transferred to 50-ml tissue culture flasks (Sarstedt, Nümbrecht, Germany). Isolation

of cells from raw cultures was carried out by micromanipulation under an inverted microscope

or by by the liquid aliquot methods pipetting a few microlitres of raw cultures in either 0.8- or

0.4-ml multi-well plates containing ambient autoclaved water. Cultures flasks and wells were

supplied with autoclaved Quinoa grains to support the growth of autochthonous bacteria.

Mono-cultures of strains HFCC 22, 157, 272, 841 and 1666 with marine/brackish origin were

further cultivated in Schmalz-Pratt medium (Suppl. Table 1). Salinity was adjusted to original

salinity of the sampling site. Freshwater strains HFCC 235, 251, 23, 5007, 162 were cultivated

in WC medium (Guillard and Lorenzen, 1972). To stimulate growth of associated bacteria,

either autoclaved wheat or quinoa grains were added to the cultures as a carbon source.

Morphological analyses

Light microscopical investigations were carried out using an inverted microscope (Zeiss, Axio Observer; water immersion condenser; 100x/1.4NA oil immersion objective and differential interference contrast). High resolution videos were recorded from cultures in Petri dishes with coverslip bases using an Allen Video-enhanced system (Hamamatsu C6489, Argus-20) to supress noise and amplify contrast. Frame by frame evaluation of videos was carried out using VirtualDub (www.virtualdub.org) and editing by ImageJ (Abramoff et al., 2004). Morphological measurements were carried out using Axio Vision Rel. 4.8 (Zeiss, Germany).

For electron microscopical investigations, cultures were fixated with 0.1 M cacodylate buffered glutaralaldehyde solution (2.5%) for one hour at 4°C. Fixated cells were washed twice with cacodylate buffer and once with distilled water and then stained with 0.5% osmium tetroxide solution for ten minutes. After two additional washing steps with distilled water, the cells were step-wise dehydrated with an increasing ethanol series (30, 50, 70, 80, 90, 96%; each step 15 min) followed by ethanol/HMDS (1:1) and dried in pure HDMS (Hexamethyldisilazan, Carl Roth, Germany). Before examination by SEM (Fei Quanta 250 FEG at the Biocenter, University of Cologne), cells were sputtered with a gold layer (12 nm).

Molecular studies

Molecular analyses followed procedures described by Schoenle et al. (2020). Strains were centrifuged (4000 x g for 20 min at 4°C, Megafuge 2.0 R, Heraeus Instruments) and genomic DNA was extracted using the Quick-gDNA^{TM} MiniPrep (Zymo Research, USA). For sequencing partial SSU rDNA, three different sets of primers were used: 18S-For (5'-AACCTGGTTGATCCTGCCAGT 3') and 18S-Rev (5'TGATCCTTCTGCAGGTTCACCTAC 3') (Medlin et al., 1988), 590For (5'-CGGTAATTCCAGCTCCAATAGC-3') in combination with 1300 Rev (5'-CACCAACTAAGAACGGCCATGC-'3) (Wylezich et al., 2002) and the set 82F (5'-GAAACTGCGAATGGCTC-3') (López-García et al., 2002) in combination with Rev-1 (5'-ACCTACGGAAACCTTGTTACG-3') (newly assigned) were used for amplification with a primer concentration of 1 µM and the help of a PCR Mastermix (VWR Life Science, Red Taq DNA Polymerase, Hassrode, Belgium). For the first two primer sets the following steps were applied: initial denaturation step at 98°C for 2 min, followed by 35 cycles of 98°C for 30 s, 55 °C for 45 s and 72°C for 2.30 min and a final elongation step for 10 min at 72°C. For the third

primer set, the following steps were used: initial denaturation at 98°C for 2 min, 35 cycles at 98°C for 30 s, at 52°C for 45 s, at 48°C for 30 s and at 72°C for 2.30 min, final elongation step at 72°C for 10 min. Length of PCR products were checked on a 1% agarose gel and subsequently purified with the PCR Purification Kit (Jena Bioscience, Jena, Germany). Purified PCR products were sequenced at GATC Biotech, Germany.

Phylogenetic analyses

The 18S rDNA of the 26 new goniomonad sequences were aligned with 25 closely related *Goniomonas* sequences from GenBank, as well as six outgroup sequences of the CRY1 group (following Shiratori and Ishida, 2016) that includes both freshwater and marine strains, as well as *Hemiarma marina*. Sequences were aligned using the MAFFT algorithm (Katoh and Standley, 2013) and were slightly manually corrected with the help of BioEdit (Hall, 1999). The corrected alignment was analysed with the Maximum likelihood (ML) method using RAxML v.8.2.12 (Stamatakis, 2014) applying the (GTR) model + Γ + I model of rate heterogeneity with 1000 replicates. Pairwise distances were calculated with BioEdit (Hall, 1999). Bayesian analysis was carried out with MrBayes on XSEDE (3.2.7a) tool (Huelsenbeck and Ronquist, 2001; Ronquist et al., 2012) using the (GTR) model + Γ + I with a sampling frequency for the Markov chain set to 10. The stop value of topological convergence diagnostic was set to 0.01. The number of generations/cycles for the MCMC algorithm was set initially to 1,000,000 and was reached after 256,000 generations. The final version of the tree was edited using MEGA-X (Kumar et al., 2018), FigTree (Version 1.4.4, https://github.com/rambaut/figtree/releases), and Inkscape (Version 1.0.1) (https://www.inkscape.org/).

RESULTS

Morphology

All investigated "*Goniomonas*-like" strains investigated in this study shared the following morphological characteristics: an anteriorly truncated cell body, rounded at the posterior end, cell covered with periplast plates, two flagella arising laterally at the cell anterior and a band of ejectisomes appearing transversely at the anterior end of the cell. Those basic characteristics varied especially between the freshwater and marine clade, with freshwater strains generally

being larger in cell size and more elongated in shape, whereas marine strains were overall smaller and more roundish.

Two out of ten analysed strains for this study (HFCC 235 and 251) could be assigned to the genus *Goniomonas* (Figure 1, A-G resp. H-K) isolated from freshwater. The cells were colourless and had a length of 5.9–10.8 µm and width of 4.4–7.9 µm (Fig 2). All isolated strains were free swimming in a "waggling, jerky" mode across the substrate. Two flagella of a subequal length (3.9–6.7 µm length for the anterior flagellum and 3.2-5.6 µm for the posterior flagellum) arise from a groove at the very anterior margin of the cell and are orientated in the same anterior-laterally direction. The cell shape was always laterally compressed, slightly elongated ovoid and sometimes pear shaped (HFCC 251). The elongated shape makes the left and right margin of the cells appearing almost parallel. The posterior end is rounded, while the anterior end is generally angularly truncated. Cells carried four to six irregularly shaped periplast plates on the ventral and dorsal side. Along the junctions of the plates, small ejectisomes arise. Cells carry a band of ejectisomes that follows the angle of the anterior end of the cell and is between 1.5–2.8 µm long.

Two other goniomonad strains (HFCC 23 and 5007) were assigned to the newly designed freshwater clade *Aquagoniomonas* n.g. (Figure 1, L-P resp. Q-T). The cells were colourless, 6.0–9.7 µm long and 3.9–5.9 µm wide. The length of the anterior flagellum was 3.8–7.6 µm and the posterior flagellum 2.0–5.6 µm long. Cells were laterally compressed, elongated, rounded towards the posterior end and angularly truncated at the anterior end. Cells carried four irregularly shaped periplast plates on the dorsal side and five on the ventral side. A band of ejectisomes (1.2–2.5 µm, n=8; Figure 2) is visible following the angle of the anterior margin. Both species are free swimming and swim in a "waggling, jerky" mode across the substrate.

Morphologically similar, though genetically distinct were the following clades: Strain HFCC 162 (Figure 3) was assigned to the new genus *Limnogoniomonas* n.g. Morphological observations for strain HFCC 22 assigned to the genus *Neptunogoniomonas* n. g. (Figure 4, A-D) show a similar cell size as *Limnogoniomonas* n. g. Both strains are free living and display a similar swimming behaviour like *Goniomonas* and *Aquagoniomonas* n. g. Cells are laterally compressed and slightly elongated. They carry two flagella of unequal length of which one possesses mastigonemes.

Strains HFCC 841 and 272 both fall into the newly created genus *Marigoniomonas* n. g. (Figure 4, E-I and J-M). Cells of this genus were in the range of 3.3–5.0 µm in length and 2.9–5.0 µm

in width, both almost only half the length of the freshwater genera. Both strains carry a slightly shorter anterior flagellum with 4.1–6.3 µm length and a posterior flagellum between 3.2–8.5 µm in length. In relation to the cell size, the flagella are much longer compared to the freshwater strains. A band of ejectisomes with 1.2–1.9 µm length can be seen at the anterior end of the cell. Bot strains show a roundish rather compact cell shape, not elongated in contrast to the freshwater genera. Cells were rounded and slightly pointed towards the posterior end and truncated and obtuse at the anterior end. Strain HFCC 272 has slightly shorter flagella than strain HFCC 841. Flagella of strain HFCC 272 additionally appeared to be rather thick (Figure 4, L, M), in contrast to all other strains investigated in this study. Cells of strain HFCC 841 carry five rectangular periplast plates on the dorsal side and four on the ventral side. Periplast plates could not be analysed for strain HFCC 272 due to difficulties with the preparation.

Another genetically distinct strain, HFCC 1666, was placed into the newly described genus *Thalassogoniomonas* n. g. (Figure 4, N-P). Since there is only one species described for this genus so far, no range of morphological data can be given. The same applies for strain HFCC 157 for the newly created genus *Poseidogoniomonas* n. g. (Figure 4, Q-U).

Phylogenetic analysis

The analysis of the partial 18S rDNA data shows a clear separation between the freshwater and marine clades. The freshwater cluster is formed by the genera *Goniomonas* and the newly described genera *Limnogoniomonas* n. g. and *Aquagoniomonas* n. g. The marine cluster is formed by *Marigoniomonas* n. g., *Thalassogoniomonas* n. g. *Poseidogoniomonas* n. g. and *Cosmogoniomonas* n. g., and there is one genus, up to now only recorded from brackish water, *Neptunogoniomonas* n. g. (Figure 5).

Among the newly isolated and analysed strains, two cluster within the freshwater clade (HFCC 251, 235) of *Goniomonas*, one in the freshwater clade of *Limnogoniomonas* n. g. (HFCC 162), and two in the freshwater clade of *Aquagoniomonas* n. g. (HFCC 23 and 5007). Among the marine isolates one clusters in the clade *Neptunogoniomonas* n. g. (HFCC 22), two in the clade *Marigoniomonas* gen. nov (HFCC 841 and 272), one in the clade *Thalassogoniomonas* n. g. (HFCC 1666) and one in the clade *Poseidogoniomonas* n. g. (HFCC 157). All clades are well supported by Bayesian analysis, yet the marine clades partly show weaker support.

Among the freshwater clades, the neotype of *Goniomonas truncata* clusters together with two other sequences isolated from the River Rhine. A pairwise comparison revealed 0.0% difference to strain HFCC 232 and 3.8% difference to strain HFCC 251 indicating the existence of two distinct species in this cluster, strongly supported by bootstrap values (Suppl. Table 2).

The large clade of the newly described genus *Aquagoniomonas* n. g. houses eleven sequences from strains originating from different freshwater environments. The newly described species, *Aquagoniomonas mylnikovii* gen. et n. sp. isolated from a pond in Borok, Russia, is closely related to a sequence from GenBank (LC000677), to which it has a pairwise distance of 0.5%, and it clusters closely together with a another GenBank sequence from an UK freshwater habitat (AY705740) and a strain from our own collection (HFCC 163) isolated from the river Rhine with a relatively small genetic distance (p-distance = 0.6% on the SSU rDNA level, Suppl. Table 2). The clade consists of several undescribed and undetermined strains from various geographic origins (originally assigned to *Goniomonas* despite their large genetic distances). Within this clade two strains from our own collection appear, both coming from the Rio Loa in the Atacama Desert in Chile. Strain HFCC 5007 is closely related to strain 5008 (0.6% pairwise distance on the SSU rDNA) and has a pairwise distance of 0.8% to GenBank sequence LC000677. Within this clade itself, genetic differences can be high, up to 9.7 % (Suppl. Table 2).

In the clade named *Limnogoniomonas* n. g., strain HFCC 162 isolated from a pond on the island Fiji, clusters closely together with two sequences formerly designated as *Goniomonas truncata* (pairwise distance to DQ980479 only 0.17 %; Suppl. Table 3).

The marine species clustered more speparate from each other, with a shorter branch and are probably more basal than isolates from freshwater habitats. Strain HFCC 22 from brackish environment (Baltic Sea) clusters together with a GenBank sequence (MK177625) originating from a freshwater pond on an island in Korea with only a relatively small genetic difference (0.3%, Suppl. Table 4).

The clade that consists of the most recently described "*Goniomonas*" *avonlea* and two other sequences, one originating from our own collection (HFCC 1525, isolated by Sabine Schiwitza from Denmark brackish coastal waters) and one sequence that has been (probably wrongly) assigned to "*G. amphinema*", LC647565, also branched quite basal. On the SSU rDNA, HFCC 1525 has 0.4% pairwise distance to "*G.*" *avonlea* (JQ434475), but is identical on the LSU (not

included in this study). In close vicinity to this cluster, the branch with strain HFCC 22 forms the newly created genus *Neptunogoniomonas* n. g. Strains HFCC 1525 and HFCC 22 show a pairwise distance of 3.1 % to each other (Suppl. Table 4). We also added several sequences (BEAP) originating from organisms sampled in the Mediterranean Ocean kindly provided by Daniel Richter and his group (Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona) to our phylogenetic analysis. Of those sequences, five formed an individual cluster supported with high bootstrap support. These potentially new species and a potentially new genus have to be described in future studies when morphological data are available.

The large variety of marine species is restructured into clades with four new genera. As the potentially most basal genus, a clade consisting of sequences either belonging to the species "*Goniomonas* (aff.) *amphinema*" and "*Goniomonas*" *pacifica* appeared. The *amphinema*/*pacifica* clade is enlarged by a strain originating from our own collection, HFCC 662, isolated from the Maldives, with only minor pairwise distances (0.1%) on the SSU rDNA level to strain LC674566 that was originally assigned to "*G.*" *pacifica*. This sequence is also very close to the sequence defined as "*G*. aff. *amphinema*" (EU047707) with only two nucleotides being different (0.1% pairwise distance, Suppl. Table 5).

Another clade with strong bootstrap support belongs to the newly created genus *Marigoniomonas* n. g. It comprises the closely related strains HFCC 841 and 5003 (pairwise distance of 0.3%; Suppl. Table 6) both originated from the coast of the Atacama Desert in Chile and are highly similar on the basis of the SSU rDNA. The strain HFCC 272 from the North Atlantic clusters besides the two sequences from the Atacama Desert with a pairwise distance of 3.4% to strain HFCC 841. Six other sequences from marine sites form a well-supported further clade named *Thalassogoniomonas* n. g. with distances among this clade of 1.4% (Suppl. Table 6).

Poseidogoniomonas n. g. is the last branch among the marine clades comprising five sequences of which one belongs to our own collection and the other four originate from the Mediterranean Ocean (Richter and Zavadska, pers. communic., see above). The strain HFCC 157 serves as a type species for this clade. The genetic diversity in the clade is relatively large with a pairwise distance of ≥3.4% (Suppl. Table 6).

DISCUSSION

Divergence of freshwater and marine strains and new genera

During this study we investigated ten different cryptophyte strains with a *Goniomonas*-like morphology. Apart from designating a valid neotype for *G. truncata*, we split the taxonomically not well resolved genus into three fresh and five marine/brackish water genera and described ten new species based on morphological and/or phylogenetic findings.

A possible splitting of the genus *Goniomonas* into further clades has already been proposed in earlier studies (Kim and Archibald, 2013; Von der Heyden et al., 2004). Von der Heyden et al. (2004) compared the branching depth between freshwater and marine species to be similar to what separates most of the prymnesiophyte genera. We found the largest genetic difference (38% on basis of the 18S rDNA) between the newly defined neotype of *Goniomonas truncata* (HFCC 235) and *Poseidogoniomonas azorensis* n. g. et n. sp. (HFCC 157), a strain from one newly described marine genus. But even between an established species like the formerly known *G.* (aff.) *amphinema* (EU047707, now *Cosmogoniomonas amphinema* comb. nov.) and *G. truncata* we found a pairwise distance of 21%. The formerly known *G. amphinema*, *G. pacifica* and *G. avonlea* were therefore sorted to different genera, *Cosmogoniomonas* n. g. resp. *Neptunogoniomonas* n. g. Earlier studies focussing on the 18S rDNA phylogeny of goniomonads (Deane et al., 2002; Von der Heyden et al., 2004) showed that the major difference between the freshwater and marine clades is based on several inserts which some freshwater species possess and which are absent in marine goniomonads. This pattern is confirmed by our newly described species. Deane et al. (2002) and Von der Heyden et al. (2004) found also that one sequence (UO3072) from freshwater is lacking the insert typical for all other freshwater goniomonads. The newly described genus *Limnogoniomonas* n. g. includes this specific sequence. Additionally, all other sequences that appear in the *Limnogoniomonas*-clade are lacking this insert, resulting in an individual branch, apart from the other freshwater genera. This branching as well as the large genetic pairwise distance of the type sequence HFCC 162 to the new neotype of *G. truncata* (HFCC 235) of 19.7% may validate the erection of a new genus *Limnogoniomonas* n. g. The. clade does also house a sequence (ON067811) originating from a study focussing on cryptic and aplastic cryptophytes (Šimek et al., 2022). The study designates, but not scientifically describes the strain as *Pseudogoniomonas* and sorts it as a sister clade to the aplastidic CRY1 lineage (Shiratori and Ishida, 2016; Shalchian-Tabrizi et al., 2008) and *Goniomonas*. However, the sequence was deposited in GenBank not under the name *Pseudogoniomonas* but *Goniomonas truncata*. Although this is initially misleading, it

ultimately corresponds to our argument above, since the study separates *Pseudogoniomonas*
from *Goniomonas* as we do here for *Limnogoniomonas* n.g.

Apart from the divergence between freshwater and marine/brackish strains, there was also a
remarkable genetic diversity even within the different genera. In our study, the *Goniomonas*
freshwater clade houses four sequences, including one of a newly described species. Between
the neotype and *Goniomonas rhenensis* n. sp. we calculated a pairwise distance on basis of the
18S rDNA of approx. 3.8%. For species in the genus *Aquagoniomonas* n. g. a pairwise distance
to 6% (between sequences of HFCC659 and AY360458) and up to 9.67 % (between sequences
AY705739 and AY360458) were recorded. Within the marine genera, we found the following
maximum pairwise distance with regard to the 18S rDNA gene: *Neptunogoniomonas* n. g. -
highest between HFCC 22 and JQ434475 with 3.5 %, *Marigoniomonas* n. g. highest between
HFCC 841 and HFCC 272 with 3.4 %, *Thalassogoniomonas* n. g. highest between AY705738
and HFCC 1666 with 1.76 % and for *Poseidogoniomonas* n. g. sequence BEAP0122b and
HFCC 157 with 3.4 %.

Isolates originating from rather remote islands (*Marigoniomonas madeiriensis* n. g. et n. sp.
from Madeira; *Thalassogoniomonas helgolandica* n. g. et n. sp. from Helgoland and
Poseidogoniomonas azorensis n. g. et n. sp. from Azores) each form an individual branch. This
may be signs for an allopatric speciation in remote areas/islands (Hernandez-Hernandez et al.,
2021) as it was postulated for percolomonads and other protist species (Arndt et al, 2020;
Hohlfeld et al., 2022). Jet, genetic differences between different protist genera can widely vary
depending on the genus and family. It is therefore difficult to set up boundaries or thresholds
up to where a sequence does still belong to a genus or not. Without being interested in intense
splitting or lumping, we consider our separation reasonable, but are aware that further studies
may change the picture again.

The observed deep sequence divergence stands in contrast to the rather conserved
morphological features of the established genera that does hardly allow such a clear
differentiation on the first glance based on their morphology. Our study shows that features
like cell size and even position and length of flagella that have been used to distinguish e.g. the
species *Cosmogoniomonas pacifica* comb. nov. and *C. amphinema* comb. nov. from each other,
have lost their discriminative power as these characters are also found in other, genetically
distinct species. The position, direction and length of flagella as described for "*G.* aff."
amphinema (Martin-Cereceda et al., 2010) can also be found in species like *Neptunogoniomons*

juergensii n. g. et n. sp., *Marigoniomonas chilensis* n. g. et n. sp. and *Poseidogoniomonas*
azorensis n. g. et n. sp., that have at least 6% pairwise distance in the 18S rDNA to the former
species. Earlier studies have already noted that even without a close link to 18S rDNA data,
different genera of Cryptophyta can show surprisingly similar morphologic features (Deane et
al., 2002). Hence, it is necessary to lay a stronger focus on molecular data to define species in
this phylogenetic group of organisms and to use morphological features as supporting
information. Still, we can observe morphological similarities within the newly described
genera.

Designation of a neotype for *Goniomonas truncata*

In public databases around 25-30 sequences are available for phylogenetic analyses of the
genera investigated in this study (on basis of the 18S rDNA). After our phylogenetic analyses,
four of these sequences fall into the *Goniomonas* clade. Since its first description (Fresenius,
1858) the species *Goniomonas truncata* has been analyzed in multiple studies (Bernard et al.,
2000; Ekelund and Patterson, 1997; Hill, 1991; Kugrens and Lee, 1991; Patterson and Lee,
2000; Vørs, 1993), but every time without any genetic material to underpin its taxonomic
position. Together with other studies (von der Heyden et al., 2004), our studies show how
genetically diverse the genus actually is. It is therefore hard to judge, whether the publications
listed above really analyzed *G. truncata* itself, or lumped multiple species into one ever-
increasing description. Kim and Archibald (2013) have completed a helpful table, showing
how many studies actually described each species of the genus.

The designation of strain HFCC 235 as the neotype for *Goniomonas truncata* was done, as the
morphology of this strain resembles the original description and the sampling origin is close to
the area where the original studies have been carried out by Fresenius. Stein (1878), placed the
species from *Monas truncata* into the new genus *Goniomonas truncata*. The original
description by Fresenius (1858) lists the following morphological features: cells are found in
freshwater, are colourless, oval to roundish and 6-10 µm in length, cells are laterally
compressed. In addition, the anterior end is truncated and carries two flagella that originate on
the side and have a length of about the total cell length. Close to the anterior margin lies a band
of a narrow structure (today we know that these are ejectisomes) and close to this band a
contractile vacuole. The swimming behaviour is described as trembling with a frequent change
in direction. The strain HFCC 235 fits well to this original description. With a cell length of

5.9–9.3 µm, an ovoid cell shape and two flagella with a similar length of 4.5–6.3 µm

respectively 4.0–5.6 µm. Depending on the individual cell, the characters fall within the

description given by Fresenius (1858). It also has a band of ejectisomes on the anterior margin

of the cell. Additionally, HFCC 235 comes from a freshwater environment and was sampled

in Germany, which is assumingly the sampling origin of both Fresenius´ and Steins´ strains.

One could argue that Fresenius gave a rather unspecific description of a goniomonad and that,

apart from the length of flagella in ratio to the cell size, his descriptions are quite universal for

this genus. Nonetheless, we find this strain appropriate due to the reasons mentioned above and

also on the basis of the illustrations that both Fresenius (1858) and Stein (1878) provided.

Combining phylogenetic and morphological data

With *Limnogoniomonas* n. g. we named a clade currently consisting of 5 strains with

Limnogoniomonas fijiensis n. g. et n. sp. as the type species. *Limnogoniomonas fijiensis* n. g.

et n. sp. differs from *Goniomonas truncata* by having a larger cell length of 7.6 - 11.4 µm but

being similar in width, making the cells much more elongated than the ones of *G. truncata*. In

addition, *Limnogoniomonas fijiensis* n. g. et n. sp. has longer flagella than all *Goniomonas*

species described within this study.

Within the freshwater clades, three isolates from the river Rhine cluster closely together (HFCC

251, 232 and 235). Another one (HFCC 163) appears quite far away and forms the

Aquagoniomonas n. g. clade. HFCC 235 was designated as the neotype for the *G. truncata* and

HFCC 251 was described as the new species *G. rhenensis* n. sp. With a pairwise distance of

3.8 %, the type species is its closest relative. *G. rhenensis* n. sp. displays larger (both in length

and width) cells like *G. truncata*, while the anterior and posterior flagella are of similar length.

G. rhenensis n. sp. carries six irregularly shaped periplast plates on the dorsal side and four on

the ventral side, while *G. truncata* carries five and respectively six periplast plates. HFCC 232

has an identical sequence to HFCC 235 and can be considered as a re-isolation of *G. truncata*

from the neotype locality.

The strain HFCC 23, isolated near Borok, Russia, was described as the new species

Aquagoniomonas mylnikovii n. g. et n. sp.. It serves as the type species for this new genus. On

basis of the 18S rDNA it shows a pairwise distance of 0.47 % to its closest relatives, the two

undescribed sequences LC000677 and AY705460 isolated in Japan and UK, respectively. In

distinction to *Goniomonas*, *Aquagoniomonas mylnikovii* n. g. et n. sp. has a similar cell length

but is wider and has a longer posterior flagellum. The cell size is smaller than of *Goniomonas rhenensis* n. sp. and it has a shorter anterior and posterior flagellum than the latter. It carries four periplast plates on the dorsal and five on the ventral side. HFCC 5007 was designated as *Aquagoniomonas atacamiensis* n. g. et n. sp. and has its type locality in the Rio Loa in Chile. This strain has a pairwise distance of 10.4 % to *G. truncata* and 0.61 % to its closest relative, a strain also originating from the Rio Loa, but was found in the mouth area of the river. *Aquagoniomonas atacamiensis* n. g. et n. sp. is of similar cell size as the type species *Aquagoniomonas mylnikovii* n. g. et n. sp. but carries longer flagella. Both species have four periplast plates on the dorsal and five on the ventral side.

The genus *Neptunogoniomonas* n. g. comprises three sequences from public databases and two sequences from our own collection. Phylogenetically, the genus is subdivided into two branches. One branch contains two species from GenBank, including the species formerly known as *Goniomonas avonlea*, now *Neptunogoniomonas avonlea* n. comb. One sequence originating from Denmark just at the transition zone between North Sea and Baltic (HFCC 1525) and one sequence with marine origin (LC647565). The other branch contains strain HFCC 22, originating from the Baltic Sea, described here as *Neptunogoniomonas juergensii* n. g. et n. sp. It branches together with a sequence from a freshwater pond in Korea (MK177625). Thus, this genus contains strains from brackish, marine and freshwater environment. This is quite unusual as all other sequences in the lower half of the phylogenetic tree (Figure 5) have a strictly marine background, maybe that further studies on this clade might reveal a further splitting. Aside from this fact, the morphology of *Neptunogoniomonas juergensii* n. g. et n. sp. slightly differs from the one of *B.* a*vonlea* n. comb. With a cell length of 5.9-9.3 µm and cell width of 4.8-7.1 µm it is smaller than the latter, but the ratio of flagella length compared to the cell length seems to be similar. Both species carry mastigonemes on one flagellum whereas the other one appears to be smooth. Both species carry four rectangular periplast plates on the dorsal side and three on the ventral side. The sequences show a pairwise distance of 3.4 % to each other (on basis of 18S rDNA).

The strains HFCC 841 and HFCC 272 form an individual branch together with another sequence from our own collection, HFCC 5003, and a more distant sequence from a public database (KX431494). Together they form the new genus *Marigoniomonas* n. g., with *Marigoniomonas chilensis* n. g. et n. sp. being designated as the type species. Interestingly, HFCC 841 branches closely together with HFCC 5003, both strains originated from coastal waters in Chile. To its closest relative, HFCC 5003, *Marigoniomonas chilensis* n. g. et n. sp.

has only minor differences (0.26 %) on basis of the 18S rDNA, and to the next sequence in the

branch, HFCC 272, a difference of 3.5 %. *Marigoniomonas chilensis* n. g. et n. sp. and

Marigoniomonas madeiriensis n. g. et n. sp. both have a rather small and very similar cell size,

both have an anterior flagellum of almost the same length, but *Marigoniomonas madeiriensis*

n. g. et n. sp. has a shorter posterior flagellum. While *Marigoniomonas chilensis* gen. et n. sp.

carries mastigonemes on one flagellum, we could not observe those in *M. madeiriensis* n. g. et

n. sp. The flagella of *M. madeiriensis* n. g. et n. sp. on the other hand, appeared to be

substantially thicker than the ones of *M. chilensis* n. g. et n. sp. Regardless of multiple trials to

fix the cells for EM studies, it was not possible to determine the exact number of periplast

plates for *M. madeiriensis* n. g. et n. sp.

The sister clade to *Marigoniomonas* n. g. houses a variety of sequences from all over the world.

Some were designated as "*G. amphinema*", assumingly on basis of morphological

observations, even though they appear genetically quite distant from the clade where *G.* aff-

amphinema (EU047707) appears. Outside of this clade the newly designated type species

HFCC 1666, *Thalassogoniomonas helgolandica* n. g. et n. sp. branches in some distance. It is

closest related to strain AY360454 to which it has a pairwise distance of 1.44%. While there

are no other described species in the clade, we can make no comparison to any other species

within this genus.

The strain HFCC 157 is designated as a type species for the new genus *Poseidogoniomonas* n.

g. *Poseidogoniomonas azorensis* n. sp. originated from the Azores islands. It clusters together

with four sequences from the Mediterranean Sea that are not described within this study, but

are added for a better resolution of the tree. It is closest related to sequence BEAP122b with

3.4 % pairwise distance and large genetic differences on basis of 18S rDNA when compared

to other genera.

The last clade consists of a mixture of sequences either traditionally assigned as *G.* (aff.)

amphinema or *G. pacifica*. Both species have been described in several publications (Al-

Qassab et al., 2002; Ekelund and Patterson, 1997; Larsen and Patterson, 1990; Martin-Cereceda

et al., 2010; Patterson and Lee, 2000; Vørs, 1993). Although our genetic analyses revealed that

this clade exhibits a large genetic diversity, its members shared a high degree of morphological

similarity. By defining "*G.* aff. *amphinema*" and the associated sequence EU047707 as

Cosmogoniomonas amphinema n. comb., we established a solid foundation for our taxonomic

work. The comprehensive species description by Martin-Cereceda et al. (2010) serves as an

ideal reference for comparing this species with others. "*G. pacifica*", but unfortunately, it lacks ultrastructural descriptions. Within our phylogenetic analysis two sequences that were assigned to "*G. pacifica*". With a pairwise distance of 0.12%, one of these sequences (LC647566) shows only minor pairwise differences to *Cosmogoniomonas amphinema* n. comb. (EU047707) and it is likely that it has been falsely assigned and belongs to the latter. Consequently, we chose to redefine the remaining sequence (AF508277) as a reference for *Cosmogoniomonas pacifica* n. comb. as it originates from Australia, and was sampled from the Pacific Ocean.

Our study, as other studies before, demonstrates the large diversity within the genus *Goniomonas* and offers a solution by splitting the genus into several individual genera. Still, we are aware that this might not be the last taxonomic revision that will be made for goniomonads. The large intraspecific differences hint at a much larger diversity and, by adding more sequences, may allow an even finer resolution (and splitting) either on the genus or species level. As already mentioned, it is hard to draw a line on basis of genetic differences, especially when the morphological bauplan is similar. Whereas differences within the marine genera ranged around 4 %, they were higher within freshwater genera such as *Aquagoniomonas* n. g.

As with many protist taxa, the diversity of goniomonads is still largely unknown, but, particularly with the help of modern sequencing techniques we are slowly taking steps to get an idea about their diversity. Nevertheless, modern taxonomy, which complements molecular databases, is the backbone for all metabarcoding studies. It is crucial to keep the taxonomy up to date, otherwise much of the diversity will be lost or remain cryptic. As more and more sequences are generated, it is important to critically examine previous taxonomic decisions and revise them if necessary.

TAXONOMIC SUMMARY

Assignment: Eukaryota, Cryptista**,** Cryptophyta, Goniomonadea, Goniomonadida

TAXONOMY OF NOVEL GENERA AND SPECIES

Goniomonas **Stein 1878, n. emend. Sachs et Arndt**

Diagnosis. A free living, colourless, bi-flagellated heterotrophic cryptophyte only found in freshwater habitats, bacterivorous. Cells are laterally compressed, slightly longer than wide, ovoid to pear shaped. Anterior end slightly angled, posterior end rounded. Cells carry two flagella of subequal length that can carry fine hairs or appendages. The cell body is covered by irregularly shaped, elongated periplast plates that have a dorso-ventral orientation.

Type species. *G. truncata*

Goniomonas truncata **Stein 1878, emend. Sachs et Arndt**

Diagnosis. Free living, colourless, bi-flagellated cryptophyte from freshwater environment. Laterally compressed cells, slightly elongated, ovoid to pear-shaped cell body, rounded towards posterior end, tapered, angularly truncated anterior end, with 5.9 – 9.3 µm length (average 7.3 µm, SD=1.0, n=22) and 4.4–7.1 µm width (average 5.5 µm, SD = 0.6, n = 22) of cell body. Two flagella of subequal length arise from an anterior groove and are directed anterior-laterally with the anterior flagellum of 4.5–6.3 µm (average 5.4, SD=0.6, n=17) length and the posterior flagellum being 4.0–5.6 µm (average 4.9, SD=0.5, n=16) long. Both flagella are equipped with fine hairs, the shorter flagellum carries additional spikey appendages. Cells carry five irregularly shaped periplast plates on the dorsal side and six on the ventral side. A band of ejectisomes is visible on the anterior margin of the cell with 1.7–2.8 µm (average 2.3, SD=0.4, n=7) in length.

Neotype location. Strain HFCC 235 was obtained from the River Rhine, Cologne, 50°49'34"N 6°35'34"E

Neotype material. The type culture is currently deposited in the Heterotrophic Flagellate Collection Cologne, as strain HFCC 235, specimen shown in Figure 1, A

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Goniomonas rhenensis **n. sp. Sachs, Nitsche et Arndt**

Diagnosis. Free living, colourless, bi-flagellated cryptophyte from freshwater environment. Laterally compressed cells, slightly elongated, ovoid to pear-shaped cell body, rounded towards posterior end, tapered, angularly truncated anterior end, with 7.8–10.8 µm (average

9.3 µm, SD = 0.4, n =20) length and 5.3–7.9 µm (average 6.4 µm, SD = 0.8, n = 20) width of cell body. Two flagella of subequal length arise from an anterior groove and are directed anterior-laterally with the anterior flagellum of 3.9–6.7 µm (average 5.8 µm, SD=0.7, n = 20) length and the posterior flagellum 3.2–5.5 µm (average 4.6 µm, SD=0.5, n =20) long. Cells carry six irregularly shaped periplast plates on the dorsal side and four on the ventral side. A slim band of ejectisomes is visible on the anterior margin of the cell with 1.6–1.9 µm (average 1.7 µm, SD = 0.1, n = 6) in length.

Type location. Strain HFCC 251 was obtained from the River Rhine, Cologne, 50°49'34"N 6°35'34"E

Type material. The type culture is currently deposited in the Heterotrophic Flagellate Collection Cologne, as strain HFCC 251, specimen shown in Figure 1, K

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology: The species name (lat. "from the Rhine") is based on the type locality of the strain, the River Rhine

Aquagoniomonas **n. g. Sachs et Arndt**

Diagnosis. A free living, colourless, bi-flagellated heterotrophic cryptophyte from freshwater habitats, bacterivorous. Cell body laterally compressed, elongated, lean to slightly ovoid shape. Posterior end slightly tapered and rounded, proximal end truncated and angled. Cells carry two smooth flagella of subequal length, anterior flagellum slightly longer than posterior. The cell body is covered with irregularly shaped, elongated periplast plates, four on the dorsal side and five on the ventral side, that have a dorso-ventral orientation.

Type species. *Aquagoniomonas mylnikovii* n. g. et n. sp.

Etymology. The genus name refers to the origin of the strains in this genus, the freshwater environment (lat. "aqua" - water)

Aquagoniomonas mylnikovii **Sachs et Arndt n. sp.**

Diagnosis. Free living, colourless, bi-flagellated cryptophyte from freshwater environment. Laterally compressed cells, elongated, lean cell body, rounded, slightly tapered towards posterior end, angularly truncated anterior end, with 6.1–9.7 µm (average 7.7 µm, SD = 1.1, n = 19) length and 4.0–5.6 µm (average 4.7, SD = 0.5, n = 20) width of cell body. Two flagella of subequal length arise from an anterior groove and are directed anterior-laterally with the anterior flagellum being slightly longer with 3.8–6.7 µm (average 5.5 µm, SD = 0.8, n = 8) length and the posterior flagellum 2.0–5.5 µm (average 4.0 µm, SD=1.2, n =8) long. Cells carry four irregularly shaped periplast plates on the dorsal side and five on the ventral side. A prominent band of ejectisomes is visible on the anterior margin of the cell with 1.8–2.5 µm (average 2.0 µm, SD = 0.2, n = 8) in length.

Type location. Pond, Borok near Jaroslawl, Russia, 58°03'27"N 38°15'37"E

Type material. The type culture is currently deposited in the Heterotrophic Flagellate Collection Cologne, as strain HFCC 23, specimen shown in Figure 1, N-O

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. This species is dedicated to and in remembrance of Alexander P. Mylnikov for his unparalleled contributions to protozoological research. He isolated this strain 30 years ago.

Aquagoniomonas atacamiensis **n. sp. Sachs, Nitsche et Arndt**

Diagnosis: Free living, colourless, bi-flagellated cryptophyte from freshwater environment. Laterally compressed cells, slightly elongated ovoid cell body, rounded towards posterior end, angularly truncated anterior end, with 6.0–9.0 µm (average 7.7 ± 0.8 µm; n = 15) length and 3.9–5.9 µm (average 5.0 ± 0.6 µm, n = 15) width of cell body. Two flagella of subequal length arise from an anterior groove and are directed anterior-laterally with the anterior flagellum being slightly longer with 7.5–7.6 µm length than the posterior flagellum being 5.6 µm long. Cells carry four irregularly shaped periplast plates on the dorsal side and five on the ventral side. A band of ejectisomes following the angle of the anterior margin is visible and 1.3–2.4 µm in length.

Type location. Atacama, Rio Loa mouth, Chile, 21°25'41"S, 70°03'15"W

Type material. The type culture is currently deposited in the Heterotrophic Flagellate Collection Cologne, as strain HFCC 5007, specimen shown in Figure 1, T

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology: The species name refers to the geographic origin of the strain. The Rio Loa runs over 400 km through the Atacama Desert until it disembogues into the Pacific Ocean and is dedicated to the friendly people of the Atacama region.

Limnogoniomonas **n. g. Sachs et Arndt**

Diagnosis. Non-photosynthetic, colourless, single celled, bacterivorous cryptophycean bi-flagellate from freshwater habitats. Cell body laterally compressed, clearly elongated, lean shape. Posterior end slightly rounded, proximal end truncated, obtuse and slightly angled. Cells carry two smooth flagella of subequal length, anterior flagellum slightly longer than posterior. Cell body covered with irregularly shaped, elongated periplast plates, four on the dorsal side and five on the ventral side, that have a dorso-ventral orientation.

Type species. *Limnogoniomonas fijiensis* n. g. et n. sp.

Etymology. The genus name is derived from the fact that the clade of *Limnogoniomonas* consists of sequences only originating from freshwater samples.

Species included. *Limnogoniomonas fijiensis* n. g. et n. sp.

Limnogoniomonas fijiensis **n. sp. Sachs et Arndt**

Diagnosis. Free living, colourless, bi-flagellated cryptophyte from freshwater environment. Laterally compressed cells, elongated cell body, rounded towards posterior end, angularly truncated and obtuse anterior end, with 7.6–11.5 µm with (average 9.3 ± 1.1 µm, n = 20) length and 3.9–7.3 µm (average 5.4 ± 0.9 µm, n = 20) width of cell body. Two flagella of subequal length arise from an anterior groove and are directed anterior-laterally with the anterior flagellum being slightly longer with 6.0–7.7 µm (average 6.8 ± 1.1 µm, n = 14) length than the posterior flagellum 4.5–6.6 µm (average 5.8 ± 0.5 µm, n = 13) long. Cells carry four irregularly

shaped periplast plates on the dorsal side and five on the ventral side. A band of ejectisomes
following the angle of the anterior margin is visible with 1.7–2.7 µm (average 2.2 ± 0.4 µm, n
=7) in length.

Type location. Rainforest pond, Suva, Island of Fiji, 18°08'29"S, 178°26'29"E

Type material. The type culture is currently deposited in the Heterotrophic Flagellate
Collection Cologne, as strain HFCC 162, specimen shown in Figure 3, B

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the
Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. The species name refers to the geographical origin of the type species, the Fiji
Islands in the Pacific Ocean and is dedicated to the friendly people of the Fiji archipelago.

Neptunogoniomonas **Sachs et Arndt n. g.**

Diagnosis. A non-photosynthetic, colourless, single celled, bacterivorous cryptophycean bi-
flagellate from marine or brackish habitats. Cell-body rounded, ovoid to obovoid in shape,
rounded posterior end, truncated anterior end. Cells have four rectangular periplast plates on
the dorsal side and three on the ventral side. Two flagella of subequal length, slightly longer
than cell length and with different orientation. During movement, anterior flagellum is directed
anterior-laterally, posterior flagellum is directed towards the cell. Anterior flagellum equipped
with mastigonemes/hairs, posterior flagellum smooth.

Type species. *Neptunogoniomonas avonlea* (Kim & Archibald, 2013) n. comb. Sachs & Arndt

Etymology. The prefix "*Neptuno*" refers to the roman god of the oceans, Neptune, as the type
for this genus was isolated from the Atlantic Ocean.

Neptunogoniomonas juergensii **n. sp. Sachs et Arndt**

Diagnosis. Free living, colourless, bi-flagellated cryptophyte from brackish environment.
Laterally compressed, ovoid cells with a truncated anterior end, 5.9–9.3 µm (average 7.7 ± 0.9
µm, n = 20) long and 4.8–7.1 µm (average 6.1 ± 0.6 µm, n = 20) wide. Flagella are of unequal
length, the anterior flagellum is directed anterior-laterally and 4.9–7.6 µm (average 6.7 µm,

SD=0.6, n = 20) long and is equipped with mastigonemes, posterior flagellum is directed
towards the cell body and oftentimes carried laying over the cell and of 5.5–9.0 µm (average
7.2 ± 0.9 µm, n = 20) length. Cells carry four rectangular periplast plates on the dorsal side
and three on the ventral side.

Type location. Island Hiddensee, Germany, Baltic Sea, 54°35'37"N 13°06'34"E

Type material. The type culture is currently deposited in the Heterotrophic Flagellate
Collection Cologne, as strain HFCC 22, specimen shown in Figure 4, D.

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the
Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. The species is dedicated to Klaus Jürgens for his major contributions to the
research on microbes of the Baltic Sea.

Marigoniomomonas **n. g. Sachs et Arndt**

Diagnosis. Non-photosynthetic, colourless, single celled, bacterivorous chryptophycean bi-
flagellate from marine environments. Cells laterally compressed, roundish to d-shaped due to
rounded posterior end and truncated, angled anterior end, angle pointing upwards from the
attachment point of the two unequally long flagella. Cells carry five rectangular periplast plates
on the dorsal side and four on the ventral side (no data for *M. madeiriensis* n. sp.)

Type species. *Marigoniomonas chilensis* n. g. et n. sp.

Etymology. The genus name refers to the marine environment as an origin for all included taxa

Marigoniomonas chilensis **n. sp. Sachs, Nitsche et Arndt**

Diagnosis. Free-living, colourless, bi-flagellated cryptophyte from marine environment.
Laterally compressed, roundish, slightly d-shaped cells with a truncated anterior end, 3.6–4.9
µm (average 4.4 ± 0.4 µm, n = 20) long and 2.9–4.6 µm (average 3.7 ± 0.5 µm, n = 20) wide.
Flagella are of unequal length and arise from an anterior depression, anterior flagellum is
approx. 1/3 shorter and directed anterior-laterally with 4.2–6.3 µm (average 4.9 ± 0.6 µm, n =
21) length and covered with mastigonemes, posterior flagellum is smooth and directed towards

the cell body and is often carried laying over the cell, 5.1–8.5 µm long. Cells carry five

rectangular periplast plates on the dorsal side and four on the ventral side. A band of

ejectisomes can be observed at the anterior margin of the cell with 1.5–1.9 µm (average 1.7 ±

0.1 µm, n=15) in length.

Type location. Pacific, coast of Mejillones, Chile, 23°3'13,72"S 70°32'58,63"W

Type material. The type culture is currently deposited in the Heterotrophic Flagellate

Collection Cologne, as strain HFCC 841, specimen shown in Figure 4, E

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the

Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

ZooBank registration: XXX

Marigoniomonas madeiriensis **n. sp. Sachs, Dünn et Arndt**

Diagnosis. Free-living, colourless, bi-flagellated cryptophyte from marine environment.

Laterally compressed cells, almost round, slightly pointed towards posterior end, truncated

anterior end, with 3.3–5.0 µm (average 4.2 ± 0.5 µm, n = 19) length and 3.1–5.0 µm (average

3.8 ± 0.5 µm, n = 20) width of cell body. Two comparably thick, smooth and short flagella of

subequal length arise from an anterior depression connected to a longitudinal groove. The

anterior flagellum is slightly longer and directed anterior-laterally with 4.1–5.7 µm 5.0 ± 0.5

µm (4.1–5.7 µm, n = 17) length, posterior flagellum is directed towards the cell body, 3.2–4.8

µm (4.0 ± 0.5 µm , n = 18) long. A prominent band of ejectisomes is visible on the anterior

margin of the cell with 1.2–1.9 µm (1.7 ± 0.2 µm, n = 16) length.

Type location. Atlantic Ocean**,** Madeira Island, Ribeira Brava, 32°36`33"N 16°53`16"W from

a depth of 950 m.

Type material. The type culture is currently deposited in the Heterotrophic Flagellate

Collection Cologne, as strain HFCC 272, specimen shown in Figure 4, M

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the

Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. The species name refers to the sampling site and is dedicated to the friendly people of Madeira Island, Portugal.

Remarks. The periplast plates could not be counted due to difficulties with SEM preparation.

Thalassogoniomonas **n. g. Sachs et Arndt**

Diagnosis. Non-photosynthetic, colourless, single celled, bacterivorous chryptophycean bi-flagellate from marine environments. Cells cup shaped, tapered posterior end and truncated, yet rounded anterior end. Two unequally long flagella arise from anterior margin of the cell, the anterior flagellum anterior-laterally directed, posterior flagellum carried over the cell in an almost curled manner. A prominent band of ejectisomes is visible transversally at the anterior margin of the cell.

Type species. *Thalassogoniomonas helgolandica* n. g. et n. sp.

Etymology. The prefix "Thalasso" (Greek: thalassa – ocean) refers to the ocean as the environment of origin.

Thalassogoniomonas helgolandica **n. sp. Sachs et Arndt**

Diagnosis. Free-living, colourless, bi-flagellated cryptophyte from marine environment. Laterally compressed cells, cup shaped, slightly elongated, with tapered posterior and truncated, obtuse anterior end, cells of 4.3–5.7 µm (average 4.2 ± 0.5 µm, n = 19) length and 3.1–4.6 µm (average 3.8 ± 0.5 µm, n = 19) width of cell body. Two flagella of unequal length, anterior flagellum with 3.3-6.3 µm (average 5.1 ± 0.7 µm, n =16) and directed anterior-laterally, posterior flagellum generally directed towards the cell body, 5.2–8.6 µm (average 7.4 ± 0.8 µm, n = 18) long. A prominent band of ejectisomes is visible on the anterior margin of the cell with 1.7–2.3 µm (average 1.9 ± 0.2 µm, n = 12) length.

Type location. North Sea, shore of the Island Helgoland (Germany), 54°11`01"N 7°53`27"W from a depth of 950 m.

Type material. The type culture is currently deposited in the Heterotrophic Flagellate Collection Cologne, as strain HFCC 1666, specimen shown in Figure 4, N

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. The species refers to the geographical origin, the Island Helgoland (Germany) in the North Sea close to the Biologische Anstalt Helgoland.

Poseidogoniomonas **n. g. Sachs et Arndt**

Diagnosis. Non-photosynthetic, colourless, single celled, bacterivorous cryptophycean bi-flagellate from marine environments. Cells laterally compressed, roundish, slightly elongated cell body, sides almost parallel to each other. Cells carry five rectangular periplast plates on the dorsal side and four on the ventral side. Two unequal flagella, with different directions, smooth shorter anterior flagellum directed anterior laterally and longer posterior flagellum directed towards or curved over the cell equipped with mastigonemes.

Type species. *Poseidogoniomonas azorensis* n. g. et n. sp.

Etymology. The addition "*Poseido*" refers to the Greek word for god of the ocean, Poseidon, and points to the marine origin of the type strain.

Poseidogoniomonas azorensis **n. sp. Sachs et Arndt**

Diagnosis. Free-living, colourless, bi-flagellated cryptophyte from marine environment. Laterally compressed cells, roundish, slightly elongated cell body, rounded towards posterior end, truncated anterior end, sides almost parallel to each other with 3.7–5.8 µm (average 4.7 ± 0.5 µm, n = 18) length and 2.7–4.8 µm (average 3.5 ± 0.5 µm, n =18) width of cell body. Two flagella of unequal length arise from an anterior depression. The anterior flagellum is directed anterior-laterally with 3.9–6.2 µm (average 5.0 ± 0.8 µm, n = 15) length and carries several mastigonemes, the posterior flagellum is directed towards the cell body often curved over the cell body, 5.0–8.8 µm (average 6.8 ± 1.1 µm, n = 14) long. Cells carry five rectangular periplast plates on the dorsal side and four on the ventral side. A band of ejectisomes is visible on the anterior margin of the cell with 1.4–1.8 µm (average 1.6 ± 0.1 µm, n = 5) in length.

Type location. Atlantic Ocean, Island of Faial, Azores (Portugal), 38°28'25"N 28°27'16"W.

Type material. The type culture is currently deposited in the Heterotrophic Flagellate
Collection Cologne, as strain HFCC 157, specimen shown in Figure 4, S.

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the
Accession Number xxx

ZooBank ID. XXX (will be provided prior to publication)

Etymology. Species name refers to its geographic origin and is dedicated to the friendly people
of the Azores islands, Portugal.

Cosmogoniomonas **n. g. Sachs et Arndt**

Diagnosis. Non-photosynthetic, colourless, single celled, bacterivorous cryptophycean bi-
flagellate from marine environments. Cells ovate with a truncated anterior and rounded
posterior end, dorso-ventrally flattended. Two flagella of similar length, one pointing forward
(can be equipped with appendages), one bent ventrally, trailing. Cells are covered with periplast
plates (exact number not stated, see remarks). Between the plates, lines of ejectisomes can be
found in the ridges. An additional transversal band of ejectisomes can be seen at the anterior
end of the cell.

Type species: *Cosmogoniomonas amphinema* (Larsen & Patterson 1990) n. comb. Sachs et
Arndt

Etymology: "*Cosmo*" refers to the supposedly cosmopolitan distribution of this genus.

Remarks: Species within the genus *Cosmogoniomonas* n. g. are morphologically relatively
similar, but genetically relatively diverse within a significantly separated clade (Figure 5).
Since morphological features overlap, we here define the 18S rDNA genotype EU047707
(Martin-Cereceda et al., 2010) as the neotype sequence for *Cosmogoniomonas amphinema* n.
comb. The detailed description by Martin-Cereceda et al. (2010) gives the best reference for
this species as the study only finds minor differences to the description of Larsen and Patterson
(1990) for *G. amphinema*. We further assume that those differences likely result from the
comparison of light microscopic (Larsen and Patterson, 1990) and electron microscopic
images. It is likely that structures like mastigonemes as flagellar appendages were simply not
visible under the light microscope and therefore not described by Larsen and Patterson (1990).
Nevertheless, features like periplast plates (or the ridges between them) are recognized by both

authors, unfortunately without further information about the number, making this specific

feature unusable for the genus description. For "*G. pacifica*" light microscopic studies ~~also~~

exist (Larsen and Patterson, 1990). In GenBank, there are only two sequences at the moment

that were assigned to "*G. pacifica*". While one sequence (LC647566) only has a minor (0.12

%) pairwise difference to *Cosmogoniomonas amphinema* n. comb. (EU047707), it is plausible

that it has been wrongly assigned and rather belongs to the latter. Hence, we have decided to

define the other existing sequence (AF508277, Deane et al., 2002) as the neotype sequence for

Cosmogoniomonas pacifica n. comb. as it includes the description of Larsen and Patterson

(1990) in the description for this species. Our choice is underpinned by the fact that AF208277

originates from the Australian Pacific.

Cosmogoniomonas amphinema **Sachs et Arndt n. comb.**

Basionym: *Goniomonas amphinema* Larsen & Patterson, 1990

Emended diagnosis: Free-living, colourless, bi-flagellated cryptophyte from marine
environment. Cells have an anterior truncate oval shape and are 5.6 ± 0.51 µm long and 3.5 ±
0.57 µm wide (n = 90). One straight shorter flagellum beating jerkily in all directions (5.18 ±
0.44 µm long), more rapidly than the trailing, longer one (6.32 ± 0.45 µm long, n = 45). The
short flagellum carries flagellar appendages. The flagellar vestibulum is about one fourth of
the cell length with a vacuole located on the opposite side to it. A line of ejectisomes appearing
transversally in the anterior end of the cell. A cytopharynx structure is often visible along the
longitudinal axis of the cell. Between the periplast plates, rows of emerging ejectisomes are
visible. The central part of the anterior part of the cell appears raised. On the ventral anterior
side, a granular area is visible. According to Martin-Cereceda et al. (2010).

Type location. Menai Strait, North Wales, UK, 53.231N, 4.164 W

Type material. Culture collection of Algae and Protozoa, CCAP 980/2, Figure 5 in Martin-
Cereceda et al. (2010).

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the
Accession Number EU047707 (Martin-Cereda et al., 2010).

Cosmogoniomonas pacifica **n. comb. Sachs et Arndt**

Basionym: *Goniomonas pacifica* Larsen and Patterson (1990)

Diagnosis of the original description by Larsen and Patterson (1990): "Cell outline ovate,
truncate anteriorly, rounded posteriorly, 8-10 µm long, 6-8 m wide, laterally compressed. With
a transverse band of ejectisomes near the anterior end. The left-hand side of the cell with 2-3
longitudinal very delicate ridges, the right-hand side with 3 ridges which are usually more
distinct. Nucleus situated in the middle of the cell near the dorsal side. Two, equal or slightly
unequal (dorsal longer) flagella emerge from a small depression anteriorly near the dorsal
surface, about 3/4 the cell length. At their insertion the bases lie parallel to the axis of the cell.
When swimming the flagella diverge; the one situated closest to the dorsal surface points
forwards and bends only slightly in dorsal direction, while the other flagellum usually bends
towards the ventral side. The cell skids parallel to the substratum when moving."

Type location. Port Phillip Bay, Melbourne, Victoria, Australia

Type material. Provasoli-Guillard Center for Culture of Marine Phytoplankton, strain
CCMP1869, Figure 1g in Larson and Patterson (1990).

Gene sequence data. The 18S rDNA sequence has been deposited at GenBank with the
Accession Number AF508277 (Deane et al., 2002).

Description in Larsen and Patterson (1990). "Cell ovate, truncate anteriorly, rounded
posteriorly, 8-10µm long, 6-8µm wide. With 2-3 longitudinal ridges on the left-hand side of
the cell, 3 on the right. Two, equal or slightly unequal flagella emerge from a small depression,
insertion along cell axis."

CRediT authorship contribution statement

Maria Sachs: Data curation, Formal analysis, Investigation, Visualization, Writing – original
draft. Frank Nitsche: Investigation, Supervision, Writing – review & editing. Hartmut Arndt:
Conceptualization, Investigation, Supervision, Funding acquisition, Project administration,
Writing – review & editing.

Acknowledgements

Special thanks go to Rosita Bieg, Brigitte Gräfe and Anke Pyschny for their valuable technical support. Many thanks go also to Sabine Schiwitza, who provided strain HFCC 1525 to the study. We are also very thankful to many students helping in field work in the Atacama Desert. This research was supported by grants of the German Federal Ministry for Education and Research (Bundesministerium für Bildung und Forschung, BMBF) within its DAM pilot mission project MGF Baltic Sea (MGF-Ostsee;grant number 03F0848A) to H.A., by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) regarding the Collaborative Research Centre 1211 "Evolution at the dry limit" (project number 268236062 (SFB 1211; B03, B02) and regarding the Project AR288/23,24 and MerMet 17-11 to H.A. and to F.N. (SPP 1991, project NI 1097/4).

Data availability

Data are published or will be made available on request.

References

Abramoff, M.D., Magalhaes, P.J. & Ram, S.J. (2004) Image processing with ImageJ. *Biophotonics International*, 11(7), 36–42.

Al-Qassab, S., Lee, W.J., Murray, S., Simpson, A.G.B. & Patterson, D.J. (2002) Flagellates from stromatolites and surrounding sediments in Shark Bay, Western Australia. *Acta Protozoologica*. 41, 91 – 144.

Archibald, J.M. (2020) Cryptomonads. *Current Biology*, 30(19), R1114–R1116. Available from: https://doi.org/10.1016/j.cub.2020.08.101

Arndt, H., Dietrich, D., Auer, B., Cleven, E., Gräfenhan, T., Weitere, M. & Mylnikov, A.
(2000) Functional Diversity of Heterotrophic Flagellates in Aquatic Ecosystems. In:
Leadbeater, B.S.C. & Green, J.C. (Eds.) *The Flagellates: Unity, Diversity and Evolution*.
Taylor & Francis, CRC Press, London, pp. 240–268. Available from:
https://doi.org/10.1201/9781482268225

Arndt, H., Ritter, B., Rybarski, A., Schiwitza, S., Dunai, T. & Nitsche, F. (2020) Mirroring the
effect of geological evolution: Protist divergence in the Atacama Desert. *Global and*
Planetary Change, 190, 103193. Available from:
https://doi.org/10.1016/j.gloplacha.2020.103193

Bernard, C., Simpson, A.G.B. & Patterson, D.J. (2000) Some free-living flagellates (Protista)
from anoxic habitats. *Ophelia*, 52(2), 113–142. Available from:
https://doi.org/10.1080/00785236.1999.10409422

Carr, M., Richter, D.J., Fozouni, P., Smith, T.J., Jeuck, A., Leadbeater, B.S.C. & Nitsche, F.
(2017) A six-gene phylogeny provides new insights into choanoflagellate evolution.
Molecular Phylogenetics and Evolution, 107, 166–178. Available from:
https://doi.org/10.1016/j.ympev.2016.10.011

Cenci, U., Sibbald, S.J., Curtis, B.A., Kamikawa, R., Eme, L., Moog, D., Henrissat, B.,
Maréchal, E., Chabi, M., Djemiel, C., Roger, A.J., Kim, E. & Archibald, J.M. (2018)
Nuclear genome sequence of the plastid-lacking cryptomonad *Goniomonas avonlea*
provides insights into the evolution of secondary plastids. *BMC Biology*, 16(1). Available
from: https://doi.org/10.1186/s12915-018-0593-5

Deane, J.A., Strachan, I.M., Saunders, G.W., Hill, D.R.A. & McFadden, G.I. (2002)
Cryptomonad evolution: nuclear 18S rDNA phylogeny versus cell morphology and
pigmentation. *Journal of Phycology*, 38, 1236–1244. Available from:
https://doi.org/10.1046/j.1529-8817.2002.01250.x

Ekelund, F. & Patterson, D.J. (1997) Some heterotrophic flagellates from a cultivated garden
soil in Australia. *Archiv für Protistenkunde*, 148(4), 461–478. Available from:
https://doi.org/10.1016/S0003-9365(97)80022-X

Fresenius, G. (1858) Beiträge zur Kenntnis mikroskopischer Organismen. Abhandlungen der
Senckenbergischen Naturforschenden Gesellschaft, 211–242. Brönner. Available from:
https://books.googleusercontent.com/books/

Guillard, R.L. & Lorenzen, C.J. (1972) Yellow-green algea with Chlorophyllide C1,2. *Journal of Phycology*, 8, 10–14.

Hall, T.A. (1999) BioEdit: a user-frienldy biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41(2), 95–98. Available from: https://doi.org/10.14601/phytopathol_mediterr-14998u1.29

Hernandez-Hernandez, T., Miller, E.C., Roman-Palacios, C. & Wiens, J.J. (2021) Speciation across the tree of life. *Biological Reviews*, 96, 1205–1242. Available from: https://doi.org/doi: 10.1111/brv.12698

Hill, D.R. (1991) Diversity of heterotrophic cryptomonads, In: Patterson, D.J., Larsen, J. (Eds.), The Biology of Free-Living Heterotrophic Flagellates. Clarendon Press, Oxford, England, pp. 235–240.

Hohlfeld, M., Meyer, C., Schoenle, A., Nitsche, F. & Arndt, H. (2022) Biogeography, autecology and phylogeny of percolomonads based on newly described species. *Journal of Eukaryotic Microbiology*, 70(1), e12930. Available from: https://doi.org/10.1111/jeu.12930

Huelsenbeck, J.P. & Ronquist, F. (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17(8), 754–755. Available from: https://doi.org/10.1093/bioinformatics/17.8.754

Jarreta de Castro, A.A. & Bicudo, C.E. (2023) *Goniomonas brasiliensis* n. sp., In: Flora Ficológica do Estado de São Paulo: Cryptophyceae. Rima Editora, Sao Carlos, pp. 73–74.

Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software Version 7: Improvements in performance and usability. *Molecular Biology and Evolution*, 30(4), 772–780. Available from: https://doi.org/10.1093/molbev/mst010

Keeling, P.J. (2004) Diversity and evolutionary history of plastids and their hosts. *American Journal of Botany*, 91(10), 1481–1493. Available from: https://doi.org/10.3732/ajb.91.10.1481

Kim, E. & Archibald, J.M. (2013) Ultrastructure and molecular phylogeny of the cryptomonad *Goniomonas avonlea* n. sp. *Protist*, 164(2), 160–182. Available from: https://doi.org/10.1016/j.protis.2012.10.002

Kugrens, P. & Lee, R.E. (1991) Organization of cryptomonads, in: Patterson, D.J., Larsen, J.
(Eds.), The Biology of Free-Living Heterotrophic Flagellates. Clarendon Press, Oxford,
England, pp. 219–233.

Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. (2018) MEGA X: Molecular
evolutionary genetics analysis across computing platforms. *Molecular Biology and*
Evolution, 35(6), 1547–1549. Available from: https://doi.org/10.1093/molbev/msy096

Larsen, J. & Patterson, D.J. (1990) Some flagellates (Protista) from tropical marine sediments.
Journal of Natural History, 24(4), 801–937. Available from:
https://doi.org/10.1080/00222939000770571

López-García, P., Rodríguez-Valera, F. & Moreira, D. (2002) Toward the monophyly of
Haeckel's Radiolaria: 18S rRNA environmental data support the sisterhood of
Polycystinea and Acantharea. *Molecular Biology and Evolution*, 19(1), 118–121.
Available from: https://doi.org/10.1093/oxfordjournals.molbev.a003976

Martin-Cereceda, M., Roberts, E.C., Wootton, E.C., Bonaccorso, E., Dyal, P., Guinea, A.,
Rogers, D., Wright, C.J. & Novarino, G. (2010) Morphology, ultrastructure, and small
subunit rDNA phylogeny of the marine heterotrophic flagellate *Goniomonas* aff.
amphinema. Journal of Eukaryotic Microbiology, 57(2), 159–170. Available from:
https://doi.org/10.1111/j.1550-7408.2009.00449.x

McFadden, G.I. (2018) Genome of tiny predator with big appetite. *BMC Biology*, 16(1), 140.
Available from: https://doi.org/10.1186/s12915-018-0610-8

McFadden, G.I. (2017) The cryptomonad nucleomorph. *Protoplasma*, 254(5), 1903–1907.
Available from: https://doi.org/10.1007/s00709-017-1153-5

Medlin, L., Elwood, H.J., Stickel, S. & Sogin, M. (1988) The characterization of enzymatically
amplified eukaryotic 16S-like rRNA-coding regions. *Gene*, 71(2), 491–499. Available
from: https://doi.org/10.1016/0378-1119(88)90066-2

Novarino, G. (2003) A companion to the identification of cryptomonad flagellates
(Cryptophyceae = Cryptomonadea). *Hydrobiologia*, 502(1), 225–270. Available from:
https://doi.org/10.1023/B:HYDR.0000004284.12535.25

Patterson, D.J. & Lee, W.J. (2000) Geographic distribution and diversity of free-living
heterotrophic flagellates, in: Leadbeater, B.S.C., Green, J.C. (Eds.), In: The Flagellates -
Unity, Diversity and Evolution. Taylor & Francis Ltd, London, pp. 269–287.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
Liu, L., Suchard, M.A. & Huelsenbeck, J.P. (2012) MrBayes 3.2: Efficient Bayesian
Phylogenetic Inference and Model Choice Across a Large Model Space. *Systematic*
Biology, 61(3), 539–542. Available from: https://doi.org/10.1093/sysbio/sys029

Schoenle, A., Hohlfeld, M., Rosse, M., Filz, P., Wylezich, C., Nitsche, F. & Arndt, H. (2020)
Global comparison of bicosoecid *Cafeteria*-like flagellates from the deep ocean and
surface waters, with reorganization of the family Cafeteriaceae. *European Journal of*
Protistology, 73, 125665. Available from:
https://doi.org/10.1016/j.ejop.2019.125665

Shiratori, T. & Ishida, K. (2016) A new heterotrophic cryptomonad: *Hemiarme marina* n.g., n.
sp. *Journal of Eukaryotic Microbiology*, 63, 804-812. Available from:
doi:10.1111/jeu.12327

Shalchian-Tabrizi, K., Bråte, J., Logares, R., Klaveness, D., Berney, C. & Jakobsen, K. (2008)
Diversification in unicellular eukaryotes: cryptomonad colonizations of marine and
freshwaters inferred from revised 18S rRNA phylogeny. *Environmental Microbiology,*
10(10), 2635–2644. Available from https://doi.org/10.1111/j.1462-2920.2008.01685.x

Simek, K., Mukherjee, I., Szoke-Nagy, T., Haber, M., Salcher, M. M., & Ghai, R. (2023)
Cryptic and ubiquitous aplastidic cryptophytes are key freshwater flagellated
bacterivores. *ISME Journal*, 17(1), 84-94. Available from:
https://doi.org/10.1038/s41396-022-01326-4

Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, 30(9), 1312–1313. Available from:
https://doi.org/10.1093/bioinformatics/btu033

Stein, F. (1878) Der Organismus der Flagellaten I., In: Der Organismus der Infusionstiere III.
Wilhelm Engelmann, Leipzig.

Von Der Heyden, S., Chao, E. & Cavalier-Smith, T. (2004) Genetic diversity of goniomonads:
an ancient divergence between marine and freshwater species. *European Journal of*

Phycology, 39(4), 343–350. Available from:
https://doi.org/10.1080/09670260400005567

Vørs, N. (1993) Marine heterotrophic amoebae, flagellates and heliozoa from Belize (Central
America) and Tenerife (Canary Islands), with descriptions of new species, *Luffisphaera*
bulbochaete n. sp., *L. longihastis* n. sp., *L. turrifor*mis n. sp. and *Paulinella intermedia*
n. sp. *Journal of Eukaryotic Microbiology*, 40, 272–287. Available from:
https://doi.org/10.1111/j.1550-7408.1993.tb04917.x

Wylezich, C., Meisterfeld, R., Meisterfeld, S. & Schlegel, M. (2002) Phylogenetic analyses of
small subunit ribosomal RNA coding regions reveal a monophyletic lineage of euglyphid
testate amoebae (Order Euglyphida). *Journal of Eukaryotic Microbiology*, 49(2), 108–
118. Available from: https://doi.org/10.1111/j.1550-7408.2002.tb00352.x

1015

Figure 1 Light and electron micrographs of the genera *Goniomonas* and *Aquagoniomonas* n. g. et n. sp. A-E *Goniomonas truncata* (bar indicates: A, D-5 µm, E-10 µm), F drawing of *Goniomonas truncata* (from Stein, 1878), drawing of *Monas truncata* (from Fresenius, 1858), H-K *Goniomonas rhenensis* n. sp. (bar indicates: H- 2µm, K-10 µm), L-P *Aquagoniomonas mylnikovii* n. g et. n. sp. (bar indicates: L, M, P-5 µm), Q-T *Aquagoniomonas atacamiensis* n. g. et n. sp. (bar indicates: Q, R-2 µm, S, T-5 µm).

Figure 2 Boxplots showing the comparison of morphological measurements of the described goniomonads. Letters in the box plots indicate the different species. a: *Limnogoniomonas fijiensis* n. g. et sp. n.; b: *Goniomonas rhenensis* n. sp.; c: *Goniomonas truncata*, d: *Aquagoniomonas mylnikovii* n. g. et n. sp., e: *Aquagoniomonas atacamiensis* n. g. et n. sp.; f: *Neptunogoniomonas juergensii* n. g. et n. sp.; g: *Marigoniomonas chilensis* n. g. et n. sp.; h: *Marigoniomonas madeiriensis* n. g. et n. sp.; i: *Thalassogoniomonas helgolandica* n. g. et n. sp.; j: *Poseidogoniomonas azorensis* n. g. et n. sp.

Figure 3 Light and electron micrographs showing *Limnogoniomonas fijiensis* gen et. n. sp. (A-E; bar indicates: A-2 µm, D-5 µm, E-3 µm)

Figure 4 Light and electron micrographs showing: A-D *Neptunogoniomonas juergensii* n. g. et n. sp. (bar indicates A, D- 5 µm), E-I *Marigoniomonas chilensis* n. g. et n. sp.(bar indicates: E, F, I- 5 µm), J-M *Marigoniomonas madeiriensis* n. g. et n. sp.(bar indicates: J - 5 µm, L,M-2 µm) , N-P *Thalassogoniomonas helgolandica* n. g. et n. sp. (bar indicates: N – 5µm), Q-U *Poseidogoniomonas azorensis* n. g. et n. sp. (bar indicates: Q, U - 2µm, S - 4 µm, T - 5 µm)

1041 **Figure 5** Maximum likelihood (ML) phylogenetic tree based on sequences of 18S rDNA.

1042 Investigated strains are marked in bold. Numbers at the nodes indicate bootstrap values of

1043 Maximum Likelihood analysis as well as Bayesian analysis, "- "indicates a differing topology

1044 for Bayesian analysis.

1045

1046 **Figure 1**

1047

1048 **Figure 2**

1049

1050

a   *Limnogoniomonas fijiensis*
(HFCC 162)

b   *Goniomonas rhenensis*
(HFCC 251)

c   *Goniomonas truncata*
(HFCC 235)

d   *Aquagoniomonas mylnikovii*
(HFCC 23)

e   *Aquagoniomonas atacamiensis*
(HFCC 5007)

f   *Neptunogoniomonas juergensii*
(HFCC 22)

g   *Marigoniomonas chilensis*
(HFCC 841)

h   *Marigoniomonas madeiriensis*
(HFCC 272)

i   *Thalassogoniomonas helgolandica*
(HFCC 1666)

j   *Poseidogoniomonas azorensis*
(HFCC 157)

1051

1052

1053   **Figure 3**

1054

1055

Figure 4

Figure 5