1

5
6
Cell division protein A (CdpA) assembles and anchors the two FtsZ rings at midcell in haloarchaea

9
Yan Liao[1,#], Vinaya Devidas Shinde[1], Dalong Hu[1,2], Zhuang Xu[3], Bill Söderström[1], Katharine A. Michie[4],
10
Iain G. Duggin[1,#]

14
1. Australian Institute for Microbiology & Infection, University of Technology Sydney, Australia

15
2. Saw SweeHock School of Public Health, National University of Singapore, Singapore

16
3. School of Mathematics and Statistics, The University of New South Wales, Australia

17
18
4. Structural Biology Facility, Mark Wainwright Analytical Centre, The University of New South Wales, Australia

20
Address correspondence to Yan Liao (yan.liao@uts.edu.au) and Iain Duggin (iain.duggin@uts.edu.au)

Abstract

Many archaea appear to divide through the coordinated activities of two distinct FtsZ homologues,
FtsZ1 and FtsZ2, and another bacterial cell division homologue, SepF, which are part of the midcell
division ring. Here, we discovered a novel cell division protein (HVO_0739, renamed CdpA) in
Haloferax volcanii. CdpA represents the prototypical characterized member of a protein family
widespread in Haloarchaea. Midcell division site localization of CdpA was dependent on the presence
of the ring-assembly protein FtsZ1, and the division constriction protein FtsZ2 influenced the proper
midcell assembly and structure of CdpA. In the absence of CdpA, cells frequently failed to divide
properly and FtsZ1 and FtsZ2 were uncoupled from midcell, with FtsZ1 forming poorly condensed
pseudo-helical structures, and FtsZ2 showing mispositioned foci and filaments that were often
disorientated and perpendicular to the division ring along the cell edge. The dynamic movements of
FtsZ1 and FtsZ2 clusters around the division ring were minimally affected by loss of CdpA, leading to
dynamic repositioning of the aberrant FtsZ structures. In contrast, CdpA formed relatively immobile
distinct foci around the ring. CdpA domain function studies, protein pull-down assays, and multimer
structure predictions suggested that CdpA forms a central part of a membrane anchor complex that
interacts with FtsZ2. CdpA has a transmembrane domain and a cytoplasmic disordered linker and small
C-terminal domain, which appear to tether multiple division proteins to the midcell membrane. Our
discovery of an archaeal FtsZ assembly and midcell anchor protein offers new insights into cell division
mechanisms that are similar across the tree of life.

Introduction

Cell division is one of the most fundamental processes in all organisms. It is universally carried out through the localized assembly of cytoskeletal proteins into a ring-like structure linked to the cell membrane and envelope that initiates a constriction process leading to division. In most bacteria, the cytoskeletal protein FtsZ and other cytoskeletal and accessory proteins form the division ring that dynamically drives cell envelope constriction[1]. Bacterial FtsZ appears to perform multiple functions, including acting as scaffolding for protein recruitment to the division site, triggering membrane constriction[2, 3], and dynamically guiding cell wall synthesis to drive its inward growth[4, 5]. The relative contributions of these mechanisms to division *in vivo* are yet to be precisely defined. The eukaryotic homolog of FtsZ is tubulin, which greatly elaborates on the core polymerization function common to this superfamily of proteins, and forms microtubules that provide structural support and play essential roles in diverse cellular processes including intracellular transport and division, although tubulin is not considered a main component of the eukaryotic division ring. Multiple tubulin proteins exist in each species, with diversified and specialized functions to support the complex cellular events and differentiation processes that they underpin.

Most bacterial species contain only one FtsZ, but archaea, which are phylogenetically more closely related to eukaryotes, most commonly have two FtsZ homologues belonging to distinct families[6-8]. The dual-FtsZ system might resemble an early stage in the evolutionary diversification and specialization of cytoskeletal functions, as seen in eukaryotes. By using the archaeal model organism, *Haloferax volcanii*, previous work showed that they divide through the coordinated activities of FtsZ1 and FtsZ2[9, 10]. The role of archaeal FtsZ1 was primarily ascribed to promoting assembly of the midcell ring, whereas FtsZ2 was more directly implicated in the ring constriction process[10]. Thus, some of the multiple functions of bacterial FtsZ appear to be at least partially separated into specialized homologs in the archaeal dual-FtsZ system. Furthermore, FtsZ2 was found to be specifically absent in certain methanogenic archaea that—rarely in archaea—have a peptidoglycan (PG) cell wall, suggesting that FtsZ2 is needed for division when a PG wall is not present to support the inward growth of the cell envelope[10].

Current knowledge of FtsZ-based cell division in archaea is limited. There are only a few proteins in this system that have been identified and characterized, despite the many different components present in bacterial and eukaryotic cell division systems. In addition to the two FtsZ proteins, an archaeal homologue of the bacterial division protein SepF[11, 12] and the very recently reported PRC barrel domain proteins (CdpB1/2)[13, 14] were found to have roles in division. *H. volcanii* SepF associates with FtsZ2 at the division site and potentially the cell membrane[11], and CdpB1 also associates with SepF[13, 14]. Crystal structures and AlphaFold2 multimer predictions suggested a tripartite complex formed by CdpB1, CdpB2 and SepF, with CdpB1 and CdpB2 forming a polymer of alternating heterodimers[13]. Currently, there is very little information on what functions these proteins play in the division mechanism or how they are linked to the envelope at midcell.

Other expected components of the FtsZ-based archaeal divisome are yet to be identified, though candidate genes linked to cell division were identified on the basis that their expression or promoter regions were influenced by a transcriptional regulator that appears to control cell division, CdrS, which is encoded by a small ORF in an operon with *ftsZ2*[15, 16]. In *H. volcanii*, CdrS-regulated genes included *ftsZ1*, *ftsZ2*, *sepF*, *cdpB1* (Hvo_1691), and the diadenylate cyclase gene (*dacZ*) implicated in signal transduction and regulation of division[16, 17]. Another previously uncharacterized gene, HVO_0739, was, with *sepF*, amongst the most strongly regulated by CdrS[16]. Here we have determined the function HVO_0739, renaming the protein CdpA (Cell division protein A), and we propose the Cdp- prefix may be given to other haloarchaeal proteins when they are discovered to participate directly in the division mechanism. We applied genetic and protein structure-function analyses combined with high-resolution fluorescence microscopy to demonstrate the functions of CdpA in coupling the two FtsZ rings and anchoring them at the midcell for division.

Results

CdpA is conserved in haloarchaea and required for normal cell division in *H. volcanii*

We identified CdpA (HVO_0739) homologs encoded in 506 published archaeal genomes and performed a phylogenetic analysis of the amino acid sequences. The *cdpA* gene is typically present as a single gene in Haloarchaea, in which it co-exists with the two FtsZs (Fig. 1a) in 96.5% of the species analyzed. Based on protein topology and structure predictions (Fig. 1b), and a multiple sequence alignment (Supplementary Fig. 1), CdpA is comprised of three main regions: a conserved N-terminal transmembrane domain, a less conserved cytoplasmic disordered linker region, and a small cytoplasmic C-terminal domain. CdpA does not have a direct homolog in Bacteria or some of the more distantly related archaeal taxa, although we observed homologous CdpA-like proteins, which showed significant homology but differed in length (± 20%) or number of N-terminal transmembrane segments (≠ 4) compared to the reference CdpA (HVO_0739) in a diverse range of archaea (Fig. 1a). Some Halobacteriota species contained multiple homologs within the genome (Fig. 1a). For example, in *H. volcanii*, a CdpA-like protein (HVO_2016) with four predicted transmembrane segments lies adjacent to HVO_2015, which encodes an actin superfamily protein (volactin)[18] hinting of a potential common function associated with diverse cytoskeletal systems.

To identify the role of CdpA, we investigated the consequences of deleting the *cdpA* gene in *H. volcanii*. Microscopy revealed a cell division defect (Fig. 1c), where the Δ*cdpA* mutant showed a mixture of some greatly enlarged cells that had failed to divide and others with almost normal size. This great heterogeneity suggests an incomplete division defect where failure of cell division in some cells leads to enlarged cells that have a lower probability of division than the smaller cells, which further exaggerates their size defect. Consistent with this, time-lapse microscopy of growing Δ*cdpA* cells showed that some divided at mid-cell and others initially developed division planes but were then unable to complete division (Fig. 1d, Supplementary Video 1). Interestingly, some of the greatly enlarged Δ*cdpA* cells appeared to lyse suddenly but produced cell fragments that were sometimes capable of ongoing growth and division (Fig. 1d, arrows; Supplementary Video 1); in others, blebbing from cellular lobes was seen. These appear to be typical features of *H. volcanii* cell division mutants that are accompanied by ongoing chromosome replication and metabolism to maintain culture viability[10].

To verify the cell division phenotype attributed to *cdpA*, we reintroduced the *cdpA* gene under the control of the tryptophan-regulated promoter (*p.tna*) on a plasmid in the Δ*cdpA* background. This rescued the cell division defect in the presence of moderate (0.2 mM) or high (2 mM) concentrations of tryptophan (Trp) (Supplementary Fig. 2), confirming that the *cdpA* open reading frame (ORF) is responsible for the observed cell division phenotype. Overexpression of *cdpA* in a wild-type *H. volcanii* background caused no detected cell division or morphological changes (Supplementary Fig. 2a).

CdpA localizes at the mid-cell division ring but differs in sub-ring localization compared to FtsZ1 and FtsZ2

We investigated the subcellular localization of CdpA by expressing GFP-tagged CdpA from a plasmid in wild-type and Δ*cdpA* strain backgrounds. Fluorescence microscopy showed that CdpA-GFP in both strains localized as a band at mid-cell in almost all cells (Fig. 1e, Supplementary Fig. 3). With relatively high concentrations of Trp (2 mM), small extracellular fluorescent particles were also seen (Supplementary Fig. 3b), suggesting that excess CdpA-GFP promotes the generation of such particles from cells. CdpA-GFP fully complemented the cell division defect of the Δ*cdpA* strain in the presence of 0.2 mM Trp induction or greater, showing that the tagged protein is functional in division (Supplementary Fig. 3).

We then investigated the localization of CdpA together with FtsZ1 or FtsZ2 by production of pairs of fusion proteins from a plasmid in the wild-type background. The degree of co-localization observed

between CdpA-GFP and FtsZ1-mCherry was strong, with a correlation coefficient of > 0.7 in 79% of
cells (Fig. 2a). CdpA-GFP showed a somewhat higher background throughout the rest of the cell than
FtsZ1-mCherry, relative to the midcell bands (Fig. 2a, normalized intensity graphs). The co-production
of CdpA-GFP and FtsZ1-GFP did not cause a substantial change in cell shape and size, similar to when
FtsZ1-mCherry or FtsZ2-mCherry are moderately produced[10]. The co-production of CdpA-GFP and
FtsZ2-mCherry resulted in enlarged or filamentous cells (Fig. 2a), suggestive of a functional
interference between these two tagged proteins. Nevertheless, CdpA-GFP and FtsZ2-mCherry still
showed strong co-localization, with 84% of cells having a correlation coefficient of > 0.8 (Fig. 2a,
righthand panels). In many of these enlarged cells, CdpA-GFP and FtsZ2-mCherry were co-localized
around mid-cell, whereas in some cells they were found near the quarter positions of the long axis of
the cell (Fig. 2a).

To investigate protein localization at higher resolution within the division ring, Structured-Illumination
Microscopy (SIM) was used to visualize cells co-producing CdpA- and FtsZ- fusion proteins. Live cell
2D-SIM revealed that CdpA-GFP and FtsZ1/2-mCherry had differing localization patterns within the
mid-cell ring (Fig. 2b). Quantification of fluorescence intensities revealed that CdpA-GFP and FtsZ1- or
FtsZ2-mCherry foci were colocalized in some regions but were clearly separated in others (Fig. 2b);
similar results were seen with cells immobilised in vertical agarose micro-holes to view the cells end-
on using confocal microscopy (Supplementary Fig. 4). We then used 3D-SIM with the normal mounting
of cells on agarose pads, which further resolved the individual ring sub-structures, and we saw that
CdpA-GFP was highly discontinuous within the ring, appearing as distinct small foci with gaps in
fluorescence intensity between them (Fig. 2c). FtsZ1-GFP and FtsZ2-GFP rings appeared more
continuous but still showed substantial variations in structure and intensity around the ring (Fig. 2c,
Supplementary Fig. 5, Supplementary Video 2-4).

CdpA has different effects on the localization of FtsZ1 and FtsZ2, and contributes strongly to their coupling at mid-cell

The division ring localization of CdpA, FtsZ1 and FtsZ2, and their different spatial patterns within the
ring, prompted us to investigate the dependency of CdpA and FtsZ1/FtsZ2 on each other for proper
assembly in the division ring. We assessed the localization of CdpA in the two *ftsZ* knockout strains
and vice versa. In the absence of FtsZ1, CdpA-GFP failed to localize as rings and infrequently appeared
as a faint midcell focus, and in an *ftsZ1/ftsZ2* double knock-out it showed a diffuse, non-localized signal
(Fig. 3a). On the other hand, FtsZ2 deletion did not prevent CdpA-GFP localization although it often
appeared as very irregularly structured, poorly condensed or indistinct rings (Fig. 3a). Thus, the proper
localization and midcell assembly of CdpA is strongly dependent on the presence of FtsZ1 and FtsZ2.

Deletion of *cdpA* had strikingly different effects on FtsZ1 and FtsZ2 function and localization patterns.
FtsZ1-mCh localizations in Δ*cdpA* appeared as rings but with broader zones compared to FtsZ1-mCh
ring structure in wild-type cells and occasional foci seen throughout the cells (Fig. 3b). Measurements
of the ratio of localized to non-localized FtsZ1-mCh per cell and each individual localization's thickness,
confirmed that the fluorescence in localized structures was similar in both strains, but the localizations
were relatively poorly condensed in many of the Δ*cdpA* cells compared to wild-type (Fig. 3b, graphs).
In these strains, the *cdpA* knockout resulted in ~30% of the total cells that we classified as filamentous
(cell area > 5 μm^2, circularity < 0.3) compared with ~5% in the wild-type (Supplementary Fig. 6). These
findings suggest that CdpA contributes to proper FtsZ1 ring condensation.

FtsZ2-GFP also showed poorly formed ring structures in the Δ*cdpA* background, but in contrast to
FtsZ1, localizations often appeared as foci, foci clusters, or short mispositioned filaments located along
the cell edge in a perpendicular orientation compared to normal rings (Fig. 3c). Consistent with a
previous observation[10], we saw that expression of FtsZ2-GFP caused a sensitive division defect,
resulting in somewhat enlarged cells in the wild-type and Δ*cdpA* backgrounds. Some cells showed
multiple localizations; the number of normal perpendicular orientated localizations of FtsZ2-GFP and

their frequency in relation to total cell length appeared to be reduced in Δc*dpA* (Fig. 3c, graphs;
Supplementary Fig. 7).

Highly informative results were obtained when producing both FtsZ1-mCh and FtsZ2-GFP in the Δ*cdpA*
background. The absence of CdpA resulted in obvious uncoupling of FtsZ1 and FtsZ2 filaments, with a
wide range of correlation coefficients (0.3-1) in most cells (Fig. 3d). FtsZ1-mCh appeared as rings with
additional poorly condensed foci throughout the cells, whilst FtsZ2-GFP was seen as foci or
mispositioned filaments, as seen during the single expression experiments above (Fig. 3d). We further
examined these strains with a relatively low concentration of tryptophan induction (0.1 mM Trp),
reflecting lower expression level of FtsZ1-mCh and FtsZ2-GFP[10], and observed broader zones of FtsZ1-
mCh localization and foci or mispositioned FtsZ2-GFP, demonstrating strong uncoupling of the two
FtsZs (Supplementary Fig. 8). These data suggest that the uncoupling is not caused by the enlarged
cell sizes and expression levels of FtsZ-FP proteins. We conclude that CdpA plays an important role in
assembly of FtsZ1 and FtsZ2 at midcell by promoting FtsZ1 ring condensation and the correct
orientation and ring-coupling of FtsZ2 structures.

CdpA is a relatively immobile midcell ring 'anchor' that has minimal effects on the dynamics of FtsZ1-
GFP and FtsZ2-GFP localization

To investigate CdpA and FtsZ movement at midcell, and whether CdpA influences FtsZ dynamics within
the division ring, we monitored the fluorescence of CdpA-GFP and FtsZ1/2-GFP in live cells over time.
We initially recorded 3D-SIM images with 1 min intervals and saw that FtsZ1-GFP was dynamic within
the ring, with changes in the intensity and location of clusters of fluorescence intensity over time
(Supplementary Fig. 9, Supplementary Video 5), consistent with previous lower-resolution
observations[10]. In contrast, CdpA-GFP showed no such movement in the ring over the same period
(Supplementary Fig. 10, Supplementary Video 6).

To better enable cluster tracking and reduce bleaching during image acquisition, we used total internal
reflection fluorescence (TIRF) microscopy to record protein movement near the cell surface closest to
the coverslip. Visual inspection of kymographs generated from the TIRF data recorded at the division
ring using 4 s frame intervals over 15 min, suggested that clusters of FtsZ1-GFP fluorescence were
moving in a directional manner (Fig. 4a). Further analysis revealed that these clusters exhibited
complex zigzag and crossover patterns, reflecting FtsZ1-GFP cluster movement in both directions
around the division plane (Fig. 4a, Supplementary Video 7-8). FtsZ2-GFP showed similar zigzag-like
patterns, but they were generally slower than FtsZ1-GFP (imaging required 10 s intervals over 40 min,
Fig. 4b, Supplementary Fig. 11, Supplementary Video 9-10). In contrast, visualization of CdpA-GFP with
TIRF over the 15- and 40-min periods showed no substantial directional movement and greatly
reduced fluctuations of fluorescence intensity over time compared to the FtsZs (Fig. 4c, 4d,
Supplementary Fig. 11-12, Supplementary Video 11-14).

It is important to note that the function of FtsZ2-GFP in cell division is more severely affected by the
GFP tag than FtsZ1-GFP[10, 19], whereas CdpA-GFP is fully functional (Fig. 1f). The dynamics of FtsZ2-GFP
or its response to other proteins might therefore be significantly affected by the tag, compared to the
untagged native FtsZ2. Here, our aim instead was to determine whether the relative dynamics of each
of the existing FtsZ-GFPs is altered by the presence or absence of CdpA. In the absence of CdpA, FtsZ1-
GFP still showed clusters with apparently similar movement rates as the wild-type background, and as
expected, they often spanned a relatively uncondensed zone of localization around midcell (Fig. 4e,
Supplementary Video 15-17). We then measured the FtsZ1-GFP fluorescence intensity over time in a
square region near the centre of a localization, and this appeared to show irregular, non-periodic
fluctuations (Fig. 4f, Supplementary Fig. 13). To identify any periodicity or dominant frequencies, we
plotted the power spectrum density (PSD)[5] in 21 cells of each strain (Fig. 4g, Supplementary Fig. 14).
In the wild-type background, there was an elevated FtsZ1-GFP periodicity at ~0.003 Hz on average
compared to Δ*cdpA* in which the average fluctuations showed a simpler decay-like curve consistent

with a lack of periodicity detected within the 15 min. The wild-type signal around 0.003 Hz was
dominant in ~24% of cells, and weak or not evident in ~76%. This heterogeneity might reflect complex
patterns of polymerization within the ring, or cells at different stages of midcell maturation and
activity (Supplementary Fig. 14). In the absence of CdpA, FtsZ1 polymerization rates or types might be
affected, or the loss of ring structural integrity could also lead to the loss of the localized periodic
fluctuations.

With FtsZ2-GFP in Δ*cdpA*, we saw striking dynamic shifts in the cellular position and orientation of
FtsZ2-GFP localizations over time (Fig. 4h-i, Supplementary Video 18-23), consistent with the strong
uncoupling and mispositioning of FtsZ2-GFP in the static images (Fig. 3c). However, Like FtsZ1-GFP,
FtsZ2-GFP clusters clearly displayed movement consistent with dynamic polymerization within the
localized structures in Δ*cdpA* that was comparable to the wild-type background (Fig. 4h-i,
Supplementary Video 9-10, 18-23). We conclude that CdpA is relatively immobile within the ring but
strongly influences the stable assembly and positioning of the two FtsZ proteins at midcell whilst
having minimal or moderate influence on FtsZ-GFP dynamics within the localized structures.

The three domains of CdpA have differing roles

CdpA contains a strongly predicted N-terminal transmembrane domain (amino acids 17-149), a
cytoplasmic unstructured linker region (150-284) and a smaller C-terminal domain (285-328) (Fig. 1b,
5a). Together with the results above, this expected topology suggests a model in which CdpA tethers
and/or helps assemble and orientate cytoplasmic FtsZ filaments and potentially other components at
the midcell membrane. We further investigated this by assessing the contributions of the three main
regions to CdpA localization and function.

We initially found that an isolated N-terminal domain (CdpA$_{NTD}$-GFP) localized weakly at mid-cell, a
NTD deletion construct (CdpA$_{\Delta NTD}$-GFP) failed to localize, and a CTD deletion construct (CdpA$_{\Delta CTD}$-GFP)
efficiently localized at mid-cell (Fig. 5b). The CdpA$_{NTD}$ is therefore essential for midcell ring localization,
but insufficient without the CdpA$_{Linker}$ for the normal degree of localization. It is possible that the linker
contributes to interactions promoting CdpA's assembly into the ring, and/or it alleviates potential
steric interference by the GFP when attached directly to the CdpA$_{NTD}$.

To investigate the influence of the three CdpA domains on cell division and FtsZ localization, we
generated in-frame deletions (scar-less) of each of the three regions in the endogenous chromosomal
copy of *cdpA*. In the absence of the NTD (*cdpA$_{\Delta NTD}$*), the cells became filamentous (Fig. 5c,
Supplementary Fig. 15a); the *cdpA$_{\Delta Linker}$* strain also showed a division defect of enlarged plate cells,
and *cdpA$_{\Delta CTD}$* showed a heterogeneous mixture of enlarged plates and filamentous cells. Thus, division
defects were clear in all three strains, but they appear to have differing cell morphologies. However,
these shape differences were neutralized upon transformation with plasmids (containing various FtsZ-
FPs), which resulted in a greater frequency of highly filamentous cells (circularity < 0.2) and no
enlarged plates (Fig. 5c, Supplementary Fig. 15a). This is consistent with the known influence of *H.*
volcanii plasmids on cell shapes[19].

In all three CdpA domain deletion backgrounds, FtsZ1-mCh appeared as poorly condensed rings, broad
zones of patchy localization or complex helicoid structures, but this was accentuated with *cdpA$_{\Delta Linker}$*
(Fig. 5c, Supplementary Fig. 15b), suggesting that this deletion causes additional interference to FtsZ1
localization. Cells producing FtsZ2-GFP in all three backgrounds were filamentous, with infrequent
rings or localizations usually near one or both poles (Fig. 5c). The *cdpA$_{\Delta Linker}$* background also caused
additional interference to FtsZ2-GFP localization because rings were rare or even not seen and foci
were often seen near the poles instead, whereas the *cdpA$_{\Delta CTD}$* background gave the mildest FtsZ2-GFP
defect in which rings more frequent in than the other mutants but less than wild-type (Fig. 5c,
Supplementary Fig. 15c).

Interestingly, when we co-produced FtsZ1-mCh and FtsZ2-GFP in the various *cdpA* deletion strains (Fig.
5d, Supplementary Fig. 16), the patterns for each protein were similar but some appeared more
accentuated than the corresponding single-FtsZ-FP strains. In *cdpA*$_{\Delta NTD}$, FtsZ1-mCh localized as rings,
poorly condensed rings or foci, while FtsZ2-GFP formed patches, foci, or rare poor rings and they
showed a weaker correlation of localization than the wild-type (Fig. 5e). In *cdpA*$_{\Delta Linker}$, FtsZ1-mCh
showed striking irregular bands scattered along the length of the cells, and FtsZ2-GFP showed a
combination of diffuse signal with additional scattered foci (Fig. 5d). This strain showed the weakest
co-localization amongst the three *cdpA* domain deletions (Fig. 5e) and the FtsZ1-mCh localizations also
occupied a much greater area of the cells than the FtsZ2-GFP localizations (Fig. 5f). In *cdpA*$_{\Delta CTD}$, we saw
strongly co-localized rings of FtsZ1-mCh and FtsZ2-GFP, but these still appeared less frequently than
in the wild type and were often near one pole (Fig. 5d-f).

The combined results suggest that: 1) The N-terminal transmembrane domain and possibly the linker
region of CdpA are important for CdpA midcell localization, FtsZ1 ring condensation, and FtsZ2 ring
assembly and positioning; 2) The linker region of CdpA is necessary to prevent strong localization
defects caused by direct fusion of the NTD and CTD, possibly by preventing steric hindrance; 3) The C-
terminal domain of CdpA does not play a major role in the assembly and coupling of FtsZ1 and FtsZ2
rings, but has at least one other significant yet currently unknown role in cell division.

297 CdpA is part of an anchor complex linked to FtsZ2 and other cell division proteins

Given the above results, we searched for CdpA interactions with other *H. volcanii* proteins by isolating
affinity-tagged CdpA (CdpA-HA) from cross-linked *H. volcanii* cell lysates (in the Δ*cdpA* background)
using anti-HA magnetic beads. Bound proteins were eluted and analysed by western blots probed with
antibodies recognising HA, FtsZ1 and FtsZ2 (Fig. 6a). Both FtsZ1 and FtsZ2 were detected in the lysate
(L) and unbound (FT) samples, however, only FtsZ2 was clearly detected in the bound protein elution
(E) sample (Fig. 6a). Neither protein was detected in the elution from the control cells containing the
HA vector only (Fig. 6a). These results suggest that CdpA can exist in a complex with FtsZ2.

We carried out additional pull-down experiments with CdpA-HA and FtsZ2-HA in their corresponding
knock-out backgrounds, compared to the wildtype background with HA vector only as the control, and
we identified bound proteins by mass spectrometry (MS). FtsZ1 was detected in the bound-protein
elution of both the CdpA-HA and FtsZ2-HA pull-downs, albeit with only moderate statistical support,
as assessed by mean MS peak area quantification (Log$_2$ fold-change (FC)=2.7, p=0.10 for the CdpA-HA
pull-down, and Log$_2$FC=3.0, p=0.03 for the FtsZ2-HA pull-down, n=6 independent experiments each).
FtsZ2 was identified significantly in the CdpA-HA pulldown (Log$_2$FC=5.7, p=0.002), and as expected in
the FtsZ2-HA pull-down (Log$_2$FC=10.6, p=0.08). The FtsZ2-HA pull-down only identified one other
protein with statistical support (i.e., CetZ2, which was not detected in the HA controls; Supplementary
Table 3). The low detection rate in the FtsZ2-HA pull-downs might be due to the relatively low
expression of the FtsZ2-HA fusion (Supplementary Fig. 17) or interference by the HA tag (as seen with
the function of FtsZ2-GFP). However, the CdpA-HA construct was well expressed and complemented
the division defect of the Δ*cdpA* background used in the pull-downs (Supplementary Fig. 17). The
CdpA-HA pull-down MS data included other proteins previously implicated in cell division, including
SepF (HVO_0392, Log$_2$FC=4.3, p=0.01), CdpB1 (HVO_1691, Log$_2$FC=3.4, p=0.07) and CdpB2 (HVO_1964,
Log$_2$FC=3.1, p=0.02) (Supplementary Table 3). Other identified proteins recently linked to cell
envelope-related functions included actin-like protein volactin (HVO_2015, Log$_2$FC=2.5, p=0.06) and
small GTPase ArvA (OapA, HVO_3014, Log$_2$FC=2.8, p=0.04)[18, 20].

To predict how the cell division proteins interact with CdpA, we used Alphafold2 (AF2) Multimer with
two molecules each of CdpA, CdpB1, CdpB2, FtsZ2, and SepF, as preliminary AF2 runs and existing
homolog structures[13, 14] had indicated that each protein can dimerise or polymerise. This revealed a
strikingly coherent predicted structure, consistent with the current experimental results, that we refer
to as the anchor complex (Fig. 6b). Most of the visible interactions within the predicted anchor

complex are supported by low pairwise alignment error scores between the corresponding regions
(Fig. 6c). The main exception to these relatively strong predictions was that the association between
the anchor complex and FtsZ2 was quite weak, hinting that interactions between the anchor complex
and FtsZ2 may occur via other unknown adaptors. The strongly predicted interactions notably include
a direct association of the cytoplasmic short N-terminal helix of CdpA (residues 1-10) with the interface
region of each CdpB1/2 heterodimer, which would bring CdpB1/2 close to the inner surface of the
membrane (Fig. 6b, 6c). Furthermore, CdpA is predicted to participate in a ternary association with
CdpB1 and SepF's N-terminal extension and membrane targeting sequence (MTS) that places the SepF
MTS in an ideal position to penetrate the inner leaflet of the cytoplasmic membrane whilst also bound
to CdpA at the interface of cytoplasm and membrane (Fig. 6b, 6c).

These experimental results and structural analysis have indicated that the CdpA NTD acts as a primary
membrane anchor located at discrete positions around the division ring, where SepF is expected to
play an adaptor-like role (braced by CdpB1/2 multimers) to connect the anchor complex with
cytoplasmic ring components including FtsZ2. We found that CdpA has multiple functions, as the inter-
domain disordered linker was critical for the proper condensation and positioning of FtsZ1 and FtsZ2
filaments, respectively, whereas the CTD was also important for division in other ways, likely through
unknown interaction partners.

Discussion

In bacteria and eukaryotes, many proteins have been identified to contribute to division ring assembly and anchoring. Here we identify and characterize the function of a ring assembly and anchor protein in haloarchaea, which we named Cell division protein A (CdpA). We found clear roles for CdpA in orchestrating the midcell assembly and co-positioning of FtsZ1 and FtsZ2 filaments to form the division ring in the model archaeon *H. volcanii*. Several other proteins involved in archaeal dual-FtsZ division have recently been identified, including SepF, CdpB1, and CdpB2[11, 13, 14], though their functions in the haloarchaeal division mechanism are unclear. Based on the current findings, we developed a structural model in which CdpA acts as the membrane-embedded component of a midcell anchor complex (including SepF, CdpB1, and CdpB2 linked to the membrane though multiple direct interactions with CdpA; Fig. 6). CdpA also functions as a tether and organizing center at relatively immobile and distinct locations around the division ring to promote FtsZ1 and FtsZ2 ring formation and maintenance, and likely other midcell ring proteins, via its cytoplasmic linker and C-terminal domains.

In bacteria, one of the best characterized FtsZ membrane linkage proteins is the widespread FtsA actin-superfamily protein. The essentiality of FtsA varies among different species. For example, in *Escherichia coli*, FtsA is essential[21], whereas in *Bacillus subtilis*, the *ftsA* gene can be deleted, although this affects Z-ring formation and causes cells to become enlongated[22]. Here, we were able to delete the *cdpA* gene in *H. volcanii*, which caused a division defect resulting in mixed populations of near-normal and greatly enlarged cells. We directly observed some of the greatly enlarged cells forming cellular fragments upon lysis of the main cell body, some of which were viable and capable of ongoing growth and division. This likely explains the survival of many *H. volcanii* cell division mutant strains in culture. As has been observed in bacteria, we anticipate that other currently uncharacterized haloarchaeal proteins, and potentially CdpA-like proteins, are also involved in midcell ring assembly or cytoskeletal protein anchoring. These may somewhat compensate for the loss of CdpA and enable some of the cells to proliferate.

Super-resolution imaging revealed that CdpA forms a more discontinuous or patchy ring-like structure at mid-cell compared to FtsZ1 or FtsZ2 rings and differing organization within the ring. FtsZ1 rarely formed normal Z rings and instead formed broad and ill-defined—often helicoid—structures, that still retained dynamic localization patterns. Although typical-looking FtsZ2-GFP rings were sometimes observed in the absence of CdpA, FtsZ2-GFP foci or filaments frequently appeared strongly mispositioned, including perpendicular to the normal ring along the cell edge. FtsZ2 also retained its dynamic properties, resulting in frequent repositioning of the misplaced FtsZ2 foci and filaments. This resulted in FtsZ1 and FtsZ2 becoming uncoupled in their subcellular localization patterns, supporting the notion that they can polymerize independently of one another[10].

Our protein pull-down experiments detected CdpA in complex with FtsZ2 but not clearly with FtsZ1, despite our finding that the proper localization of CdpA depends on the presence of both FtsZ2 and especially FtsZ1 (and vice versa). Current observations would suggest that FtsZ1 associates only indirectly with the CdpA-anchor complex, via another anchor, or in a manner that is not well maintained during the pull-down assay procedure. The CdpA pull-downs also identified SepF and CdpB1/2 amongst the interaction partners. The observed co-dependencies of localization, and the similar poorly-condensed Z-ring structures observed in the absence of CdpA, SepF[11], or CdpB1/2[13, 14], suggest a hypothesis in which these proteins assemble into a complex that functions in Z-ring assembly.

Previous studies reported that *H. volcanii* SepF is part of a complex with FtsZ2[11], and with CdpB1/2[13, 14]. Furthermore, SepF appears to localize at mid-cell via a short N-terminal amphipathic sequence, which was expected to act as a membrane targeting sequence (MTS)[11]. The finding that the SepF MTS

alone was sufficient for midcell ring localization[11] suggests that this short sequence does more than
just potential membrane targeting. Our combined results and structural modelling (Fig. 6) support the
existence of a midcell anchor complex that forms through multiple intimate associations, including
the N-terminal extensions of CdpA and SepF interacting in different ways with CdpB1/2 heterodimers,
and SepF's MTS (at the very N-terminus) simultaneously associating with the CdpA NTD and partly
embedded in the cytoplasmic membrane. The SepF MTS may thereby be sufficient for midcell
localization[11] due to its likely association with CdpA. The main membrane anchoring component
appears to be the CdpA NTD, linked to and braced by multimers of SepF and CdpB1/2.

Consistent with the structural model (Fig. 6), our domain localization studies indicated that the CdpA
NTD is essential for its midcell localization, whereas the CTD was dispensable for midcell localization
but required for normal division. The NTD and linker regions appear to be critical for assembly and
anchoring of FtsZ1 and FtsZ2 rings, whereas the CTD seems to not strongly affect FtsZ1 and FtsZ2.
These observations suggest a multi-functional role of CdpA in *H. volcanii* division, where the long,
disordered linker and C-terminal domain also play important but currently undefined roles in tethering
or assembling other proteins at midcell.

CdpA shows some noteworthy similarities and differences to other division ring anchor proteins. In
bacteria, the FtsZ ring is typically anchored to the midcell membrane by FtsA, ZipA or SepF, whereas
in non-FtsZ archaeal division based on the ESCRT-III (CdvB) cytoskeletal proteins, CdvA is considered
to play a different anchor-like role. FtsA is broadly conserved in most bacteria and contains a short C-
terminal MTS and also directly interacts with FtsZ, recruiting it to the inner membrane[23, 24]. ZipA is an
integral inner membrane protein found in Gram-negative bacteria, and is comprised of a large C-
terminal globular cytoplasmic domain linked to a single N-terminal transmembrane domain via an
extended linker[25, 26]. FtsA and ZipA have overlapping functions but likely serve differing roles in
modulating division ring assembly[27, 28] Both FtsA and ZipA interact directly with FtsZ through an
interaction hub located at its C-terminal tail[29]. SepF is conserved in Gram-positive bacteria and
contains an N-terminal MTS and forms functional dimers, which interact with the C-terminal domain
of FtsZ[30, 31]. In the Crenarchaeota, *Sulfolobus* CdvA links ESCRT-III (CdvB) proteins to the cell membrane
via direct interactions and is thought to assemble on the membrane surface[32].

Compared to these proteins, CdpA exhibits highest topological similarity to ZipA, suggesting an
analogous midcell anchor role. However, ZipA-GFP in *E. coli* showed clear movement around the
division ring[33], yet we found that CdpA-GFP remains relatively stable at mid-cell in *H. volcanii*,
suggesting they may operate via differing mechanisms within the ring. SepF homologues from
different bacterial species can polymerize into large protein rings associated with cell wall
remodelling[34], whereas there is no evidence to show that archaeal SepF is capable of polymerization[11,
12]. Our discovery of an archaeal FtsZ-assembly and transmembrane anchor protein confirms this as a
feature important in both major branches of microbial life.

Methods

Identification, phylogeny, and structural analysis of CdpA and CdpA-like sequences

The reference amino acid sequence of HVO_0739 (CdpA) was first input into JackHMMER[35] to search against UniProtKB database[36] with three iterations restricted by taxonomy in Archaea (taxid:2157). The output HMM profile was used to scan the amino-acid sequences of all proteins in 1,587 high quality diverse archaeal genomes from NCBI RefSeq database[37] with a significance threshold of an E-value of 10^{-5}. The identified sequences were then confirmed using a BLASTP v2.13.0[38] search against the reference CdpA sequence with an E-value cut-off of 10^{-5}; sequences giving no hit in the BLASTP search result were removed. The remaining 1,336 sequences were taken as CdpA candidates. Of these, proteins with a length between 80% to 120% of HVO_0739 (i.e., 260-400 residues) and with four transmembrane segments (predicted using TMHMM v2.0)[39] within the N-terminal 75% of the sequences were taken as CdpA family proteins (470 sequences), and the remaining 866 were designated CdpA-like proteins.

Multiple alignment of the 470 CdpA sequences was then performed using MUSCLE v3.8.31[40]. A neighbour-joining tree based on the amino-acid alignment was constructed using PHYLIP v3.698[41] with the default model and 1,000 bootstrap replicates. HVO_2016 (*H. volcanii* CdpA-like protein) was included as the outgroup to root the tree. The alignment was then reordered by tree structure. A reduced version of the alignment was generated by removing columns that had a gap in the *H. volcanii* CdpA reference sequence. This alignment was displayed using UGENE v46.0[42]. Taxonomy annotation was based on the NCBI Taxonomy database. PsiPred v4.02[43] was used to estimate the secondary structure of CdpA along the alignment.

Protein 3D structure predictions were downloaded in monomer form from the Alphafold structure database (https://www.alphafold.ebi.ac.uk) or generated as complexes using Alphafold v2.3.1 (Multimer)[44] using default settings with the full BFD reference database. Structures were visualised and rendered with PyMOL[45] v2.5.1. Transmembrane topology (Fig. 1b, 6b) was predicted with Phobius[46].

Growth of *H. volcanii* strains

The *H. volcanii* cultures were grown in Hv-Cab medium[9, 47]. Where necessary as auxotrophic requirements, media were supplemented with uracil (50 μg/mL) for Δ*pyrE2* strains. To control gene expression via the *p.tnaA* promoter, the indicated concentration of L-tryptophan (Sigma–Aldrich) was supplemented in these cultures. For microscopic imaging, the cultures were incubated at 42°C with rotary shaking (200 rpm) and were maintained in log growth (OD_{600} < 0.8) for at least 2 days before sampling for analysis of mid-log cultures. Representative data displayed in this paper are representative of at least two independent biological replicate experiments.

Construction of plasmids for gene deletion and genomic modification in *H. volcanii*

The plasmids, oligonucleotide sequences, and strains are given in the Supplementary Tables 1 and 2. The plasmid pTA131 was used as the basis to construct the plasmids for gene deletion[48]. To make the plasmid for *cdpA* deletion, the upstream and downstream flanks for homologous recombination on either side of *cdpA* were PCR amplified from *H. volcanii* H26 genomic DNA, using the primers given in Supplementary Table 1. The upstream flank fragment and downstream fragments were digested with HindIII and BamHI, respectively, and cloned into pTA131 (at HindIII-BamHI) using Gibson assembly reagents (NEB), resulting in pTA131_up_down_0739. The plasmids pTA131_CdpA_delNTD for N-terminal domain (17-149 aa) deletion and pTA131_CdpA_delLinker for linker region deletion (150-284 aa) were obtained synthesised and cloned (Genscript). To make the plasmid pTA131_CdpA_delCTD for C-terminal domain (285-328 aa) deletion, the upstream flank and downstream flanking fragments of the CdpA C-terminal domain were amplified by using the primers in Supplementary Table 1 and cloned into pTA131 (at HindIII-BamHI) using Gibson assembly (NEB) as described above.

Demethylated plasmids pTA131_up_down_0739, pTA131_CdpA_delNTD, pTA131_CdpA_delLinker,
and pTA131_CdpA_delCTD were used to transform *H. volcanni* H26 separately by using the pop-in-
pop-out method as described previously to delete the corresponding gene fragments[48]. Allele-
specific PCR was used to verify the expected deletion chromosomal structure, resulting in H26-base
strains carrying the *cdpA* mutations: Δ*cdpA*, *cdpA*$_{\Delta NTD}$, *cdpA*$_{\Delta Linker}$, and *cdpA*$_{\Delta CTD}$ (Supplementary Table
2).

Construction of plasmids for gene expression in *H. volcanii*

The plasmids pTA962[49] or pIDJL40[9] (containing GFP) were used as the basis to construct plasmids for
controlled expression of the *cdpA* genes or modified versions. To construct pTA962-*cdpA*, the *cdpA*
ORF (HVO_0739) was amplified from *H. volcanii* H26 genomic DNA using primers HVO_0739_r and
HVO_0739_r, then the product was cloned between the NdeI and BamHI sites of pTA962. The *cdpA*
ORF was also amplified without its stop codon, then cloned between the NdeI and BamHI sites of
pIDJL40, to create the *cdpA*-gfp fusion. For dual expression of *cdpA*-gfp and *ftsZ1/2*-mCherry, PvuII
fragment of pTA962-*ftsZ1*-mCherry or *ftsZ2*-mCherry[10] (containing a *p.tnaA-ftsZ*-mCherry gene fusion)
were ligated into HindIII-cut (Klenow blunt-ended) pIDJL40_HVO_0739 containing (*cdpA*-GFP),
creating plasmid with two independent copies of the CdpA-GFP and FtsZ1/2-mCherry.

To construct the CdpA$_{NTD}$-GFP fusion, the N-terminal domain of CdpA (1-149 aa) was amplified from
H. volcanii H26 genomic DNA using primers HVO_0739_f and CdpA_NTD_R, then the product was
cloned between NdeI and BamHI sites of pIDJL40. The mutant CdpA without the C-terminal domain
and without the N-terminal sequences were amplified from *H. volcanii* H26 and mutant *cdpA*$_{\Delta NTD}$
genomic DNA, respectively, by using the primers in Supplementary Table 1, and then cloned between
NdeI and BamHI sites of pIDJL40, to create CdpA$_{\Delta CTD}$-GFP and CdpA$_{\Delta NTD}$-GFP fusions. To construct
pHVID1-HA, synthetic DNA fragments encoding the HA tag with EcoRI and NotI overhangs were
obtained from Integrated DNA Technologies (IDT). These fragments were subsequently cloned into
the pHVID1 vector. To generate CdpA and FtsZ2 HA tagged plasmids, the respective genes amplified
from pIDJL40_*cdpA* and pIDJL40-*ftsZ2* were cloned between NdeI and BamHI sites of pHVID1-HA
plasmid, respectively. All plasmids were demethylated by passage through *E. coli* C2925 and re-
purified before transfer to *H. volcanii* H26 by polyethylene glycol-mediated spheroplast
transformation[48].

Microscopy

Phase-contrast and epifluorescence microscopy: a 2 μl sample of mid-log culture was loaded on a 1%
agarose pad containing 18% buffered saltwater (BSW) on a glass slide at room temperature, and a
#1.5 glass coverslip was placed on top[10]. The images were acquired on a GE DeltaVision Elite
microscope with a pco.edge 5.5 sCMOS camara and an Olympus 100X Plan Apo/NA 1.4 objective using
a FITC filter (ex 464-492 nm: em 500-549 nm) and mCherry filter set (ex 557-590, em 598-617 nm) for
visualizing GFP-tagged and mCherry-tagged proteins, respectively.

Time-lapse microscopy: the sample was prepared by using submerged sandwich technique as
previously described[10]. A 2 μl sample of mid-log culture was placed on a 0.3% wt/vol agarose pad
(containing the medium for the growth) prepared on an 8-mm glass-based (#1.5) FluoroDish (WPI). A
4 mL volume of medium was added to the FluoroDish to cover the agarose pad, and the lid was placed
on the FluoroDish to avoid evaporation. The images were captured at 42°C every 10 mins for 24 hrs
on a Nikon TiE2 inverted Fluorescence deconvolution microscope fitted with a 100 X oil-immersion
phase-contrast NA 1.45 objective.

2D- and 3D-Structured Illumination Microscopy (SIM): Samples were prepared on agarose pads as
described above for phase-contrast and epifluorescence microscopy. Images were captured on a Delta
Vision OMX SR 3D Structured Illumination Microscope with a 60 X Plan 1.42 NA Oil Immersion
Objective at room temperature for visualizing fluorescent fusion proteins, or at 42°C for 3D-SIM time-

lapse imaging. GFP and mCherry signals were collected through a FITC and a TRITC filter, respectively.
3D-SIM images were sectioned using a 125 nm z-step. The reconstructed images-stacks were 3D
rendered and visualized by using IMARIS x64 9.6.0 (Bitplane Scientific). 3D-SIM time series (acquired
every 30s – 2 min as required) were compiled from single time points using Imaris to create a time-
lapse series, followed by bleach correction (Histogram Match) and registration (using 'translation' in
TurboReg) in Image J 1.53v. The 3D intensity plots were generated by using '3D surface plot" in Image
J 1.53 v. Two regions of interest were monitored overtime show how FtsZ- or CdpA-GFP fluorescence
fluctuates in the Z ring.

Confocal imaging for standing cells: Production of microholes in agarose pads for standing cell
trapping were performed as previously described for bacterial cells[4, 50], with some required
modification for imaging archaeal cells. Liquefied agarose (4% w/v) in Hv-Cab media was dispersed on
glass slides and the silica mold was placed on top. Once the agarose solidified, the mold was removed
and ~ 4 µl of live cell mid-log culture was loaded on the top followed by placing the glass slides on a
microcentrifuge for a brief centrifuging to trap the cells in a vertical position in micron-sized holes in
agarose pads. The excess liquid was then removed, and the agarose pad was covered with a pre-
cleaned #1.5 cover glass. The confocal images of cells expressing CdpA-GFP and FtsZ-mCherry were
acquired on Leica STELLARIS SP8 confocal microscope equipped with a 63X oil immersion objective.
The Lightning module was used to generate deconvolved images.

Total Internal Reflection fluorescence (TIRF) microscopy: Samples were prepared as described for time-
lapse microscopy. The TIRF imaging was performed using a Nikon TiE-2 N-STORM microscope
operated in TIRF mode on a stage top incubator (OKO-lab) set to 42°C, equipped with a 100 X 1.49 NA
TIRF objective. The TIRF angle was set optimize the signal to noise ratio at a fixed angle of ~100°. The
imaging settings were chosen to minimize photobleaching with the power intensity setting as 5% and
the exposure time as ~ 150 – 200 ms. The dynamics of FtsZ1-GFP proteins were imaged at 4 second
intervals for 15 mins, and FtsZ2-GFP were imaged at 10 second intervals for 40 min. CdpA-GFP were
imaged either at 4 second intervals for 15 mins or 10 second intervals for 40 mins.

Image analysis

The analysis for cell shape (area vs circularity), FtsZ1/2 and CdpA medial localization along the cell long
axis, and FtsZ1-mCh intensity ratio and localization thickness were performed as described
previously[10]. The number of correct positioned FtsZ2-GFP rings per cell was manually counted and
plotted against cell length determined by FIJI.

For the dynamic analysis from TIRF data[5], fluorescence images were stabilized, and registered
(StackReg, Translation), and corrected for photobleaching (simple ratio), interpolated to 20 nm/pixel
via the bicubic method in Image J, and moving averaged over a 12-frame window of time. The
fluorescence intensity of an FtsZ polymer along the direction of its movement in each frame was
determined from the intensity along a line with a width of 25 interpolated pixels (~ 500 nm) manually
drawn across the full length of the path of the FtsZ polymer. This line was then used to plot the
kymograph (line width = 25) then background subtraction rolling ball =25, and smooth (for display
purpose). The speeds of clusters were calculated by manually measuring the slopes of leading and
trailing edges of clear fluorescence zigzags in the kymograph. The distributions of speed of FtsZ-FP
clusters were plotted in OrginPro (2021). For analysis of Z-ring intensity fluctuations during TIRF time-
lapse imaging, briefly, a cell of interest was cropped out from both fluorescence and bright-field
images. Second, the cell was rotated to align the Z-ring horizontally. Fluorescence background within
the cell was calculated by averaging the total integrated intensity over the area defined by the cell
contour. A square region of interest (ROI) with 3×3 pixels in the center of the ring was then chosen
manually. The corrected intensity was calculated as the integrated intensity within the ROI after
background subtraction and smoothed by a moving-average filter with a window size of 5 time points
in OriginPro (2021).

The Power Spectral Density (PSD) analysis was conducted using custom Matlab scripts. Firstly, a cell
of interest was cropped out from both fluorescence and bright-field images. The fluorescence images
were then registered to the bright-field image and corrected for sample/stage drift and
photobleaching. Following this, the cell was rotated manually to align the Z-ring horizontally. To
determine fluorescence background within the cell, the total integrated intensity over the area
defined by the cell outline was averaged. A square region of interest (ROI) with 3×3 pixels at the center
of the ring was then chosen manually. The local FtsZ-GFP (I) was computed as the integrated intensity
within the ROI after background subtraction.

Using fluorescence intensity time trace, $I(t)$, from the center of the division ring, the PSD curves were
calculated. This involved subtracting the mean value $\langle I(t) \rangle$ from $I(t)$ and subsequently applying the
Fast-Fourier Transform (FFT) to the shifted time trace (fft function in Matlab). For each cell, the PSD
curve across different frequencies f was then calculated as follows:

583
$$PSD(f) = \frac{|FFT(I(t) - \langle I(t) \rangle)|}{n\Delta f \cdot Var(I(t))},$$

where n is the total number of frames of the trace and Δf is the frame rate ($1/4 \ s^{-1}$). Note that the
PSD was normalized by the squared magnitude of the intensity trace (variance of $I(t)$). The mean PSD
curves were generated by averaging the PSD curves of individual cells in each strain.

The FtsZ1/FtsZ2 localization patterns were analyzed in Matlab 2019b (MathWorks). The phase-
contrast images were first preprocessed using a bottom-hat transform (imbothat function) with a disk-
shaped structuring element of a radius of 30 pixels for the kernel. These were binarized and individual
cells were segmented (bwconncomp function). Cell debris were excluded by removing objects with
area less than 1 μm^2 (~237 pixels). The filtered cell outline was parametrically smoothed by fitting
each component of cell outline (i.e. x- and y-components in the Cartesian coordinate system) with two
Fourier expansions up to the 8th order mode.

To quantify the localization patterns of FtsZ1 and FtsZ2, fluorescent images were pre-treated by
subtracting the average intensity of the background (calculated from the fluorescence images,
excluding regions of the detected cells). Additionally, a top-hat transform (imtophat function) was
applied with a disk-shaped structuring element of a radius of 10 pixels. FtsZ1 and FtsZ2 clusters within
each cell were identified and segmented by converting the processed fluorescence images to 8-bit
and applying the extended-maxima transform (imextendedmax function) with a threshold value of 1.
Small fluctuations in FtsZ1 and FtsZ2 fluorescence signals were neglected by removing clusters less
than 4 pixels. The localization area of FtsZ1/FtsZ2 in each cell is then defined as the total area of all
FtsZ1/FtsZ2 clusters found within the cell outline. The ratio of FtsZ2 localization area to FtsZ1
localization area was then determined for each cell. Cells were excluded from the analysis if either
FtsZ1 or FtsZ2 clusters were undetectable within the cell or if the cell exhibited a signal-to-noise ratio
of less than 2.

Immunoprecipitation of HA-tagged CdpA

The method was employed as describe by previous work[11] but with modifications. H26-HA and CdpA-
HA strains were grown in 300 mL of Hv-cab medium supplemented with 2 mM Trp until their OD600
reached 0.5. The cells were harvested by centrifugation at 6,000 x g for 15 min. The pelleted cells were
resuspended in 30 mL of 18% buffered salt water supplemented with 10 mM HEPES. For crosslinking
1.5% (w/v) formaldehyde was added to the cell suspension, and the mixture was incubated at room
temperature for 10 min with constant shaking. The crosslinking reaction was stopped by adding 20 mL

of 18% buffered salt-water containing 1.25 M glycine. The cells were centrifuged again for 15 min at
6,000 x g. The cell pellet was washed twice with 18% buffered salt water with 1.25 M glycine. After
the final wash, the cell pellet was resuspended in 3 mL of lysis buffer (1X PBS, 10 mM MgCl2, 10 µg/ml
DNase I and protease inhibitor cocktail). The cells where lysed by sonication for 5 min, with 5 sec on
and 5 sec off cycle at 35% amplitude. To remove the cell debris lysed cells were centrifuged at 8,000
rpm for 15 min at 4 °C in a microcentrifuge. The supernatant was used for immunoprecipitation with
washed anti-HA magnetic beads (Thermo Scientific) as described in the manufacturer's instructions.
The supernatant (1 mL) was added to 0.5 mg of anti-HA beads and incubated at room temperature for
30 min with gentle mixing. The beads were washed twice with buffer (Tis–buffered saline with 0.05%
Tween –20 Detergent) and eluted with 0.1 M glycine pH 2.0.

Western blotting

Samples were first run on a sodium dodecyl-sulphate polyacrylamide gel electrophoresis (SDS-PAGE)
and then electro transferred to a nitrocellulose membrane using the BioRad trans-blot turbo transfer
system. The membrane was blocked overnight in 5% (w/v) skim milk in Tris-buffered saline with 0.05%
Tween-20 (TBST). A primary antibody recognizing heamagglutanin (HA) (Thermo Fisher Scientific, Cat#
26183) (1:10,000), FtsZ1[10] (1:1,000) and FtsZ2[10] (1:2,000) were incubated with membrane for 2 h with
constant shaking at RT. Following that the membrane was incubated with the secondary antibody
against mouse (Goat anti-mouse IgG-HRP; AbCam 6789) for 1 h in a 1:5,000 dilution. Bands were
detected with enhanced chemiluminescence reagents (Thermo-Fisher) using an Amersham Imager
600 system.

Mass spectrometry (MS)

Fifty microgram of eluted proteins from the pull-down assay were subjected to trypsin digestion using
Promega Trypsin Gold, mass spectrometer grade (1 µg/µL) in 50 mM acetic acid. The digested samples
were processed for clean-up using solid-phase extraction (SPE). An equivalent amount of protein (15
µg) for each sample was loaded onto the SPE columns, which were then washed, and peptides eluted.
Columns were manually prepared using Empore SDS-RPS (Styrenedivinylbenzene reverse phase
sulfonate) with a pipette tip. The SDB-RPS based tips were used for peptide clean up.

Eluted samples were dried in a vacuum centrifuge, and peptides were reconstituted in the injection
solvent suitable for the LC/MS system. Analysis was performed on a Q Exactive Plus mass
spectrometer (Thermo Fisher Scientific), and the MS/MS data files were analyzed using PEAKS Studio
against *Haloferax volcanii* predicted proteome database using Uniprot. The MS peak areas obtained
for the peptides were then used for statistical analyses.

References

1. Adams, D.W. & Errington, J. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. *Nature Reviews Microbiology* **7**, 642-653 (2009).
2. Osawa, M. & Erickson, H.P. Liposome division by a simple bacterial division machinery. *Proceedings of the National Academy of Sciences* **110**, 11000-11004 (2013).
3. Szwedziak, P., Wang, Q., Bharat, T.A., Tsim, M. & Löwe, J. Architecture of the ring formed by the tubulin homologue FtsZ in bacterial cell division. *Elife* **3**, e04601 (2014).
4. Bisson-Filho, A.W. et al. Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division. *Science* **355**, 739-743 (2017).
5. Yang, X. et al. GTPase activity–coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis. *Science* **355**, 744-747 (2017).
6. Aylett, C.H. & Duggin, I.G. The tubulin superfamily in archaea. *Prokaryotic Cytoskeletons: Filamentous Protein Polymers Active in the Cytoplasm of Bacterial and Archaeal Cells*, 393-417 (2017).
7. Liao, Y., Ithurbide, S., de Silva, R.T., Erdmann, S. & Duggin, I.G. Archaeal cell biology: diverse functions of tubulin-like cytoskeletal proteins at the cell envelope. *Emerging Topics in Life Sciences* **2**, 547-559 (2018).
8. Vaughan, S., Wickstead, B., Gull, K. & Addinall, S.G. Molecular evolution of FtsZ protein sequences encoded within the genomes of archaea, bacteria, and eukaryota. *Journal of molecular evolution* **58**, 19-29 (2004).
9. Duggin, I.G. et al. CetZ tubulin-like proteins control archaeal cell shape. *Nature* **519**, 362-365 (2015).
10. Liao, Y., Ithurbide, S., Evenhuis, C., Löwe, J. & Duggin, I.G. Cell division in the archaeon Haloferax volcanii relies on two FtsZ proteins with distinct functions in division ring assembly and constriction. *Nature microbiology* **6**, 594-605 (2021).
11. Nussbaum, P., Gerstner, M., Dingethal, M., Erb, C. & Albers, S.V. The archaeal protein SepF is essential for cell division in Haloferax volcanii. *Nat Commun* **12**, 3469 (2021).
12. Pende, N. et al. SepF is the FtsZ anchor in archaea, with features of an ancestral cell division system. *Nature Communications* **12**, 3214 (2021).
13. Nussbaum, P. et al. Proteins containing photosynthetic reaction centre domains modulate FtsZ-based archaeal cell division. *Nat Microbiol* **9**, 698-711 (2024).
14. Zhao, S. et al. Widespread photosynthesis reaction centre barrel proteins are necessary for haloarchaeal cell division. *Nat Microbiol* **9**, 712-726 (2024).
15. Darnell, C.L. et al. The ribbon-helix-helix domain protein CdrS regulates the tubulin homolog ftsZ2 to control cell division in archaea. *MBio* **11**, 10.1128/mbio. 01007-01020 (2020).
16. Liao, Y. et al. CdrS is a global transcriptional regulator influencing cell division in Haloferax volcanii. *Mbio* **12**, 10.1128/mbio. 01416-01421 (2021).
17. Braun, F. et al. Cyclic nucleotides in archaea: Cyclic di-AMP in the archaeon Haloferax volcanii and its putative role. *Microbiologyopen* **8**, e00829 (2019).
18. Schiller, H. et al. Identification of structural and regulatory cell-shape determinants in Haloferax volcanii. *Nat Commun* **15**, 1414 (2024).
19. Ithurbide, S., de Silva, R.T., Brown, H.J., Shinde, V. & Duggin, I.G. A vector system for single and tandem expression of cloned genes and multi-colour fluorescent tagging in Haloferax volcanii. *Microbiology (Reading)* **170** (2024).
20. Mills, J. et al. Extracellular vesicle formation in Euryarchaeota is driven by a small GTPase. *Proc Natl Acad Sci U S A* **121**, e2311321121 (2024).
21. Addinall, S.G. & Lutkenhaus, J. FtsA is localized to the septum in an FtsZ-dependent manner. *Journal of bacteriology* **178**, 7167-7172 (1996).
22. Beall, B. & Lutkenhaus, J. Impaired cell division and sporulation of a Bacillus subtilis strain with the ftsA gene deleted. *Journal of bacteriology* **174**, 2398-2403 (1992).

23.   Pichoff, S. & Lutkenhaus, J. Tethering the Z ring to the membrane through a conserved
membrane targeting sequence in FtsA. *Molecular microbiology* **55**, 1722-1734 (2005).
24.   Pichoff, S. & Lutkenhaus, J. Identification of a region of FtsA required for interaction with FtsZ.
Molecular microbiology **64**, 1129-1138 (2007).
25.   Mosyak, L. et al. The bacterial cell-division protein ZipA and its interaction with an FtsZ
fragment revealed by X-ray crystallography. *The EMBO journal* **19**, 3179-3191 (2000).
26.   Moy, F.J., Glasfeld, E., Mosyak, L. & Powers, R. Solution structure of ZipA, a crucial component
of Escherichia coli cell division. *Biochemistry* **39**, 9146-9156 (2000).
27.   Cameron, T.A., Vega, D.E., Yu, C., Xiao, H. & Margolin, W. ZipA uses a two-pronged FtsZ-binding
mechanism necessary for cell division. *Mbio* **12**, e02529-02521 (2021).
28.   Pichoff, S. & Lutkenhaus, J. Unique and overlapping roles for ZipA and FtsA in septal ring
assembly in Escherichia coli. *The EMBO journal* (2002).
29.   Ortiz, C., Natale, P., Cueto, L. & Vicente, M. The keepers of the ring: regulators of FtsZ assembly.
FEMS microbiology reviews **40**, 57-67 (2016).
30.   Hamoen, L.W., Meile, J.C., De Jong, W., Noirot, P. & Errington, J. SepF, a novel FtsZ‐interacting
protein required for a late step in cell division. *Molecular microbiology* **59**, 989-999 (2006).
31.   Sogues, A. et al. Essential dynamic interdependence of FtsZ and SepF for Z-ring and septum
formation in Corynebacterium glutamicum. *Nature communications* **11**, 1641 (2020).
32.   Samson, R.Y. et al. Molecular and structural basis of ESCRT-III recruitment to membranes
during archaeal cell division. *Mol Cell* **41**, 186-196 (2011).
33.   Söderström, B., Chan, H., Shilling, P.J., Skoglund, U. & Daley, D.O. Spatial separation of FtsZ
and FtsN during cell division. *Molecular microbiology* **107**, 387-401 (2018).
34.   Wenzel, M. et al. Control of septum thickness by the curvature of SepF polymers. *Proceedings
of the National Academy of Sciences* **118**, e2002635118 (2021).
35.   Johnson, L.S., Eddy, S.R. & Portugaly, E. Hidden Markov model speed heuristic and iterative
HMM search procedure. *BMC bioinformatics* **11**, 1-8 (2010).
36.   Coudert, E. et al. Annotation of biologically relevant ligands in UniProtKB using ChEBI.
Bioinformatics **39**, btac793 (2023).
37.   O'Leary, N.A. et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic
expansion, and functional annotation. *Nucleic acids research* **44**, D733-D745 (2016).
38.   Camacho, C. et al. BLAST+: architecture and applications. *BMC bioinformatics* **10**, 1-9 (2009).
39.   Krogh, A., Larsson, B., Von Heijne, G. & Sonnhammer, E.L. Predicting transmembrane protein
topology with a hidden Markov model: application to complete genomes. *Journal of molecular
biology* **305**, 567-580 (2001).
40.   Edgar, R.C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic acids research **32**, 1792-1797 (2004).
41.   Felsenstein, J. PHYLIP (phylogeny inference package), version 3.5 c. (Joseph Felsenstein., 1993).
42.   Okonechnikov, K., Golosova, O., Fursov, M. & Team, U. Unipro UGENE: a unified
bioinformatics toolkit. *Bioinformatics* **28**, 1166-1167 (2012).
43.   Buchan, D.W. & Jones, D.T. The PSIPRED protein analysis workbench: 20 years on. *Nucleic
acids research* **47**, W402-W407 (2019).
44.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *nature* **596**, 583-
589 (2021).
45.   Schrodinger, L. The PyMOL molecular graphics system. *Version* **1**, 8 (2015).
46.   Kall, L., Krogh, A. & Sonnhammer, E.L. Advantages of combined transmembrane topology and
signal peptide prediction--the Phobius web server. *Nucleic Acids Res* **35**, W429-432 (2007).
47.   de Silva, R.T. et al. Improved growth and morphological plasticity of Haloferax volcanii.
Microbiology (Reading) **167** (2021).
48.   Allers, T., Ngo, H.-P., Mevarech, M. & Lloyd, R.G. Development of additional selectable
markers for the halophilic archaeon Haloferax volcanii based on the leuB and trpA genes.
Applied and environmental microbiology **70**, 943-953 (2004).

49. Allers, T., Barak, S., Liddell, S., Wardell, K. & Mevarech, M. Improved strains and plasmid
vectors for conditional overexpression of His-tagged proteins in Haloferax volcanii. *Applied*
and environmental microbiology **76**, 1759-1769 (2010).
50. Söderström, B., Badrutdinov, A., Chan, H. & Skoglund, U. Cell shape-independent FtsZ
dynamics in synthetically remodeled bacterial cells. *Nature Communications* **9**, 4323 (2018).

Acknowledgments

This study was supported by the University of Technology Sydney Chancellor's Research Fellowship
(PR021-13161 to Y.L.) and by the Australian Research Council (FT160100010 to I.G.D, and
DP220101143 to B.S.). For technical and facilities management support, we thank Matt Padula from
the Proteomics Core Facility, UTS, and Amy Bottomley, Christian Evenhuis and Louise Cole from the
Microbial Imaging Facility, UTS.

Author Contributions

Y.L and I.G.D. designed the research. Y.L. constructed most of the plasmids, strains and conducted the
primary microscopy imaging and analyses. V.D.S constructed plasmids and strains for pull-down
experiments and performed immunoprecipitation, western blot and mass-spectrometry analyses. D.H
performed phylogeny analysis of CdpA and CdpA-like sequences and contributed the production of
some image analysis figures using R programming. Z.X. performed PSD analysis and quantified the
localization area of FtsZ1/FtsZ2 in Matlab. B.S. contributed to the TIRF imaging and confocal imaging
of standing cells. K.A.M, and I.G.D. performed Alphafold2 Multimer prediction and analysis. Y.L. and
I.G.D. interpreted results and wrote the original draft. All authors reviewed and gave scientific input
to revise the manuscript. Y.L and I.G.D managed and supervised the project and acquired funding.

Competing interests statement

The authors declare no competing interests.

Fig. 1. CdpA is cell division protein in haloarchaea. a, Phylogenetic distribution of CdpA (orange circles), CdpA-like (pink), CetZ (blue), FtsZ1 (light green), and FtsZ2 (dark green) on a schematic reference phylogeny of the Archaea (Genome Taxonomy Database, release 207), expanding on Halobacteriota families. **b,** Alphafold2 prediction of CdpA monomer structure coloured by model confidence (blue - pLDDT > 90%, orange - pLDDT < 50%) and predicted overall membrane topology. **c,**

Phase-contrast microscopy of wild-type (WT) and Δ*cdpA* strains in steady mid-log cultures and the
corresponding cell shape analysis. n= number of cells examined over one biologically independent
experiment. The data shown is representative of at least two independent experiments. **d,** Live-cell
time-lapse images of Δ*cdpA* growing on an agarose media pad and showing cell lysis, fragmentation,
and division processes (arrows). **e**, **f,** Phase-contrast and fluorescence micrographs of wild-type cells
expressing CdpA-GFP (**e**) and Δ*cdpA* expressing CdpA-GFP (**f**) under control of the *p.tnaA* promoter
with 0.2 mM Trp induction. Demographic analysis (righthand panels) showing fluorescence intensity
along the long axis ordered by cell lengths, top to bottom. n= number of cells examined over one
biologically independent experiment. The data shown in panel d, e, f is representative of at least two
independent experiments. All scale bars, 5 μm.

Fig. 2. Sub-cellular and sub-ring localization of CdpA with FtsZ1 or FtsZ2. a, Phase-contrast and
fluorescence images of *H. volcanii* strains YL6 (H26 + CdpA-GFP + FtsZ1-mCherry dual expression) and
YL7 (H26 + CdpA-GFP + FtsZ2-mCherry dual expression) cultured with 0.2 mM Trp during mid-log
growth. The middle graphs show the intensity profile of the fluorescence signal (magenta for mCherry,
and green for GFP) by plotting the normalized median intensity perpendicular to the long axis of cells.
The right graphs show correlation coefficients plotted as a frequency distribution. n = number of cells
examined over one independent experiment. The data shown are representative of at least two
independent experiments. Scale bars, 5 μm. **b,** Dual-colour of 2D-SIM images of mid-log *H. volcanii*
YL6 and YL7 grown with 0.2 mM Trp. mCh (magenta) and GFP (green) fluorescence was quantified by
plotting normalized intensity in the respective channel along the ring band. Scale bars, 1 μm. **c,** 3D-
SIM images (xy-tilted) of mid-log *H. volcanii* YL29 (H26 + CdpA-GFP), YL24 (H26 + FtsZ1-GFP), and YL26
(H26 + FtsZ2-GFP) grown with 0.2 mM Trp. Scale bars, 0.5 μm. The data shown in panel b and c are
representative of at least two independent experiments.

Fig. 3. CdpA, FtsZ1 and FtsZ2 show different localization interdependencies and strongly influence each other's midcell structures. All strains were grown in Hv-Cab medium with 0.2 mM Trp, and mid-log cultures were sampled for microscopy. **a,** Phase-contrast and fluorescence composite images of CdpA-GFP in the absence of FtsZ1, FtsZ2, or both. Scale bars, 5 µm. **b,** Left: localization of FtsZ1-mCh in wild-type H26 and Δ*cdpA* backgrounds. The arrowhead indicates a poorly condensed FtsZ1-mCh

structure in Δ*cdpA*. Scale bars, 5 μm. The dotplots show quantification of the degree of FtsZ1-mCh
localization versus individual localization thickness as a measurement of protein condensation along
the long axis in the indicated backgrounds. The data points are coloured with a proximity heatmap. **c,**
localization of FtsZ2-GFP in H26 (scale bar, 5 μm) and Δ*cdpA* backgrounds (scale bars, 2 μm). The plots
show the number of normal localizations of FtsZ2-GFP versus cell length in the indicated backgrounds.
Coloured lines indicate the lengths of cells that have the indicated relative number of localizations per
unit length (one ring per ~ 4 um). **d,** Upper: localization of FtsZ1-mCherry and FtsZ2 GFP in H26 and
Δ*cdpA* strains. Scale bars, 2 μm, Bottom: correlation coefficient between FtsZ1-mCh and FtsZ2-GFP
plotted as a frequency distribution. The majority (97%) of H26 cells had a correlation coefficient of
≥0.8, while only 32% of Δ*cdpA* cells had a correlation coefficient of ≥0.8. For panel b, c, and d, n=
number of cells examined over one biologically independent experiment, and the data shown is
representative of at least two independent experiments.

Fig. 4. Division protein localization dynamics and the effect of *cdpA* deletion on FtsZ1 and FtsZ2 movement. Cells were grown in Hv-Cab without tryptophan induction to minimize the expression level of fluorescent fusion proteins. Mid-log cultures were sampled for TRIF imaging. The data shown is representative of at least two independent experiments. **a,** Maximum intensity projections (MIPs) of two representative wild-type H26 cells expressing FtsZ1-GFP (i and ii) are shown on the left, with the corresponding kymographs at positions indicated by red arrows in the MIPs shown on the right. **b,** MIPs of two representative wild-type H26 cells expressing FtsZ2-GFP (i and ii) and their corresponding

kymographs. **c-d,** MIPs of wild-type H26 cells expressing CdpA-GFP and their corresponding
kymographs with different imaging time-window (4s intervals for 15 mins, panel c; 10s intervals for 40
mins, panel d). **e,** MIP of one representative Δ*cdpA* expressing FtsZ1-GFP and its kymograph computed
from the intensity along the long axis of cell (between the two red arrows). The relative position (0-1)
along the long axis of cell is labelled below the kymograph. **f,** Montages of FtsZ1-GFP dynamics in
Δ*cdpA* and the integrated fluorescence time trace for the ROI region (red box) around the centre of
ring shown at the start of imaging. The read curve is the moving average (every 5 points) of the raw
intensity (grey dots). **g,** The mean power spectral density (PSD) curves (± SEM) of FtsZ1-GFP in wild-
type and Δ*cdpA* cells (n=21). **h-i,** Montages of representative FtsZ2-GFP dynamics in two Δ*cdpA* cells
and their corresponding kymographs indicate the changes in ring position (h) and orientation (i). All
scale bars are 1 μm.

Fig. 5. The three domains of CdpA have differing roles in subcellular localization and assembly of FtsZ1 and FtsZ2. a, Alphafold2 models of CdpA showing domain boundaries of the GFP tagged deletion constructs, including the N-terminal domain (NTD), disordered linker, and C-terminal domain (CTD). **b,** Fluorescence microscopy of H26 wild-type strains expressing the indicated constructs during mid-log growth in Hv-Cab medium with 0.2 mM Trp. Arrows indicate regions of weak mid-cell localization. Scale bars, 5 μm. The lower panels show demographic analyses of fluorescence along the long axis, ordered from shortest to longest cells (left to right). n=number of cells examined over one

representative experiment. **c,** Phase-contrast and fluorescence micrographs of the indicated *cdpA*
deletion strains producing either FtsZ1-mCh or FtsZ2-GFP, sampled during mid-log phase growth in
Hv-Cab + 0.2 mM Trp. Arrows indicate some faint foci of FtsZ2-GFP. **d,** Fluorescence images of the
indicated *cdpA* deletion strains co-producing FtsZ1-mCh and FtsZ2-GFP. **e,** Distributions of cellular
correlation coefficients and (**f**) ratio of localization area for FtsZ1-mCh:FtsZ2-GFP in the background
strains shown below the lower panel. The data shown are representative from one of two
independent experiments.

Fig. 6. CdpA is associated with FtsZ2 and other division proteins in a putative membrane anchor complex. a, Western blots of one of three replicate co-immunoprecipitation experiments. The cell lysate (L), unbound protein flow through (FT) and bound protein elution fraction (E) from control (H26 + HA vector only) and Δ*cdpA* + CdpA-HA were analysed by 15% SDS-PAGE and western blotting. The total protein stain (Ponceau S) is shown for the same membrane used for the anti-HA probe, and replicate membranes were probed with anti-FtsZ1 or anti-FtsZ2 as indicated. **b,** Alphafold2 model of the mid-cell anchor complex generated from a sequence input of 2 CdpA, 2 CdpB1, 2 CdpB2, 2 FtsZ2, and 2 SepF. The C-terminal regions of CdpA and the N- and C-terminal extensions of FtsZ2 that are largely disordered are not shown for clarity. **c,** The Alphafold2 predicted aligned error (%) heatmap plot of all input sequences (concatenated), indicating the degree of confidence in pairwise interactions between residues. **d,** The same structure and orientation as in panel b shown with surface electrostatics, indicating the approximate CdpA transmembrane domain (TMD) and SepF membrane targeting sequence (MTS).