Recording butterfly movements from a UAV

Emmanuel de Margerie*[1], Kyra Monmasson[1]

[1] CNRS, Univ. Rennes, UniCaen, UMR 6552, Rennes, France

* Corresponding Author
Correspondence: emmanuel.demargerie@univ-rennes.fr

Abstract

Tracking and understanding the movements of animals in the wild is a fast-growing area of research, known as *movement ecology*. However, tracking small animals such as flying insects, which cannot easily carry an electronic tag, remains challenging as existing field methods are costly either in terms of equipment or tracking effort (e.g. VHF radio-tracking, scanning harmonic radar). Here we attempted to record the movements of free-flying butterflies from an unmanned aerial vehicle (UAV), maintaining a static position in the sky and recording video vertically downwards. With an appropriate flight height and image filtering algorithm, we recorded 166 flight tracks of *Pieris* butterflies (*P. brassicae* and *P. rapae*), with a median tracking length of 40 m (median flight duration 13 s), and a high temporal resolution of 30 positions per second. Average flight direction varied significantly over the course of the flying season, from a northern azimuth in June and early July, to a southern azimuth in September, a trans-generational migratory behaviour that had previously been documented by field observations or experiments in flight cages. In addition, UAV imagery unlocks the possibility to measure high-resolution flight movement patterns (e.g. path tortuosity and transverse oscillations), which will possibly help understand perceptual and locomotor mechanisms underlying spatial behaviour. We explore the technical details associated with UAV tracking methodology, and discuss its limitations, in particular the biases associated with a 2D projection of 3D flight movements, and the difficulty to distinguish between visually similar species, such as *P. brassicae* and *P. rapae*.

Keywords: UAV image-based tracking, insect flight, insect migration, movement ecology.

                                          Introduction

Tracking the movements of an animal in its natural environment provides insights on its
ecology, such as habitat use (e.g. Da Silveira et al 2016), dispersal and migration behaviours
(Whitfield et al. 2024, Rotics et al. 2016). When the movement data are detailed enough (spatial
and temporal resolution), it is also possible to extract biological information on the animal's
locomotor biomechanics (Sherub et al. 2016, Hedrick et al. 2018, Ruaux et al. 2023), its spatial
search strategies (Shepard et al. 2011, Hernandez-Pilego et al. 2017, de Margerie et al. 2018),
and more broadly the perceptual and cognitive processes involved in movement (Kashetsky et
al. 2021). All of these biological and ecological aspects of animal movement have been
advantageously integrated into a synthetic "movement ecology" framework (Nathan et al.
2008, Abrahms et al. 2017, Joo et al. 2022).

However, not all animal species are easy to track. Large-enough animals (approx. > 100 g)
can carry GPS receivers (Cagnacci et al 2010), which are now often enhanced with additional
sensors (accelerometers, barometers, cameras) to better infer the animal's behaviour along its
route (Joo et al. 2022). To track the movements of smaller species such as flying insects,
tracking tags need to be much lighter, and alternative technologies to GPS are employed.
Beacons emitting simple VHF radio beeps have been used successfully for decades to track the
movements of flying and non-flying insects (Kissling et al. 2014). Yet, VHF radio-tracking
involves following the animal with one or multiple antennae (or deploying a fixed antennae
array; e.g. Knight et al. 2019), and the spatio-temporal resolution is inferior to GPS tracking.
Even lighter passive tags (of a few mg, without batteries) can be used to track the movements
of flying insects (Ovaskainen et al. 2008, Lihoreau et al. 2012, Maggiora et al. 2019). The
downside of these passive tags is that a scanning harmonic radar (SHR) - a heavy and expensive
device - has to be positioned in the field. The tracking range of SHR is near 1 km, and the
temporal resolution of the data is 0.33 Hz (i.e. sampling one position every 3 s, Ovaskainen et
al. 2008). Other shorter-range, passive-tag tracking systems also exist (portable harmonic radar,
RFID tags; see Kissling et al. 2014, Batsleer et al. 2020, Rhodes et al. 2022 for reviews).

For all animal-borne tracking systems, whether GPS receivers or other types of tags, the
impact of the tag on the animal's movements is a matter of concern. The carried mass, but also
the drag, the position, or the method of attachment of the tag must be carefully considered.
Impact studies are often necessary, both from an ethical point of view and in terms of the
reliability of the collected movement data (Wilson & McMahon 2006, Batsleer et al. 2020).

There are other routes to track flying insects, without having to place tags on animals. First,
the classic mark-recapture method allows to demonstrate the movement of an individual from
one point to another, and was thus used for studying insect migrations (e.g. butterflies;
Chowdhury et al. 2021, 2022). Trapping techniques along migratory routes, sometimes
combined with isotope analyses to determine the geographical origin of individuals, can also
provide information on migrations (e.g. dragonflies; Knoblauch et al. 2021, Oelman et al. 2023).
At a finer, local scale, if the insect flies fairly slowly in an open environment and remains visible,
it can be followed by foot and its passed positions can be drawn on a map (Brussard & Ehrlich
1970) or recorded by a GPS receiver carried by the observer (Delattre et al. 2013, Fernandez et
al. 2016). Tracking butterflies crossing a body of water from a boat is another possible technique
(Srygley & Oliveira 2001). Finally, when flying insects move in swarms of many individuals, radar

can be used to detect and measure the direction of these flights (e.g. Stefanescu et al. 2013),
but in this case individual movement is not accessible.

To complement these "tagless" tracking approaches, also called "passive sensing" (Rhodes
et al. 2022), we wondered whether flying insects could be tracked in video images filmed from
a UAV (Unmanned Air Vehicle, or drone) positioned above them in the sky.

Image-based tracking (from ground-based cameras) is a known technique to reconstruct
the 2D and 3D trajectories of animals, and in particular flying insects, with a high sampling
frequency (> 1Hz). It can be used in laboratory (e.g. Lihoreau et al. 2016), semi-natural outdoor
insectaries (Kitamura & Imafuku 2015, Le Roy et al. 2021) or even natural environments (Stürzl
et al. 2016, Jackson et al. 2016). Most often used to address biological questions related to
perception, cognition or locomotion, image-based tracking techniques, which are non-
invasive, are also attracting growing interest in ecology (Dell et al. 2014). Hybrid techniques
tracking tagged animals in videos are also being developed (Crall et al. 2015, Walter et al. 2021).

On the other hand, in the last decade, commercial UAVs have greatly improved in terms of
lightness, compactness, stability and image resolution, while decreasing in cost, making them
valuable tools for wildlife inventory and conservation (Wang et al. 2019, Charbonneau &
Lemaître 2021). Tracking animal movements using images recorded from a UAV is a next logical
step. It has recently been achieved for large species (e.g. reef shark: Rieucau et al. 2018, wild
dog: Haalck et al. 2020), and was even proposed for flying insects (Ivosevic et al. 2017).

Here we explore the validity of a UAV as a platform for remote, passive observation of the
movements of insects in flight. The motivation for this exploration is the many potential
benefits of such an approach, namely (1) non-invasiveness, i.e. tracking of animals moving
freely, without tags or capture, (2) in an open natural environment, over distances greater than
in an insectary, (3) with a spatio-temporal resolution superior to VHF or SHR tracking. If this type
of fine movement data in natural conditions can be extracted easily and cheaply, it could be
very useful for bridging the gap between laboratory studies on perceptual, cognitive and
locomotor mechanisms, and large-scale movement patterns observed in the natural
environment (e.g. migrations).

To begin this methodological exploration with a relatively simple case, we have focused on
Pieridae butterfly species (large white *Pieris brassicae* and small white *Pieris rapae*) because of
their relatively slow flight, good visibility and abundance in the field. We also limited the present
study to the simple case of a static UAV in the sky, filming vertically downwards (Fig. 1A) to
reconstruct flight trajectories in only 2 horizontal dimensions.

Material and Methods

112   This section describes the general methods we used to film butterflies in the field,
113 reconstruct their flight trajectories and describe their movements. We later explored and
114 validated these data using several specific methods and statistics, which are detailed at the
115 start of each results sub-section, for ease of reading.

Study site

We recorded butterfly flights from June to September 2021, in an agricultural area near the
city of Rennes, France (coordinates 48.105444, -1.560029). The local landscape is covered with

cultivated fields, few tree patches, small roads and farms, with the closest urbanized area
situated at 1.7 km. We filmed butterflies flying above 2 fields situated on each side of a narrow
road (Fig.1B). The western field (WF) contained an organic mixed crop (mainly wheat, fava and
lucerne) which was harvested in August, whereas the eastern field (EF) contained forage grass,
which was cut in June and August. We chose to record butterflies above two different fields
(filmed in alternance) to test our image filtering method above various backgrounds, and also
control whether the flight trajectories could be influenced by ground nature.
On the central road were the UAV takeoff/landing area, the pilot (EDM), the technical
assistant (KM), and an ultrasonic anemometer (Gill Maximet 501) recording wind speed and
direction every second, at 2 m height. We painted two permanent red crosses 20 m apart along
the central road (Fig.1D, E), as a reference line segment for positioning the UAV, and later for
scaling the video frames. The ground slope in the recorded area was less than 2°.

Figure 1 – UAV field of view (FOV). **(A)** Video recording geometry, showing the influence
of UAV flight height (*H*) and camera angular horizontal FOV (*hFOV*) on ground horizontal
FOV (*ghFOV*) and ground pixel pitch (*gPP*). **(B)** Aerial view of the study site, showing FOV
over western (WF) and eastern field (EF). *Source: geoportail.gouv.fr* **(C)** Variation of *ghFOV*
and *gPP*, as functions of *H*. These linear relationships depend on camera angular *hFOV*
and recorded image width *IW* (see equations 1-2). **(D-E)** Examples of video frames
recorded on July 8th 2021, over WF and EF, respectively. White arrows identify the 2
reference crosses painted 20 m apart on the road, and used for frame scaling.

UAV video recording

We used a Mavic Air 2 UAV (DJI, Nanshan, Shenzhen, China), which is a small commercial
quadcopter (takeoff weight 570 g, retail price ~1000 € in 2021). This UAV has a CMOS sensor of
3840 × 2160 pixel resolution (i.e. "4K" images, with 16:9 aspect ratio). According to the UAV
manual, the camera lens has an f/2.8 aperture and an 84° field of view. As most recent UAVs can

record with various zoom/resolution levels (which may involve sensor cropping), we measured
FOV in the lab, by placing the UAV camera at a known distance from a wall, and measuring the
horizontal distance along the wall that is effectively recorded within the UAV camera image. The
"horizontal" FOV ($hFOV$, i.e. along image width), was measured at 68.3° in the default "4K wide"
recording mode, that we used throughout the study. This angular $hFOV$ value was used to
choose a flight height. We computed the horizontal field of view on the ground ($ghFOV$, in
meters) when the UAV camera aims vertically downwards:

(1) $$ghFOV = 2H \tan\left(\frac{hFOV}{2}\right)$$

where H is the UAV height (m), and $hFOV$ is the horizontal angular FOV (°).
154
155 Proportional to $ghFOV$ is the corresponding pixel pitch on the ground, i.e. the distance on
156 the ground covered by a single pixel side (gPP, m):

157 (2) $$gPP = ghFOV/IW$$

158 where IW is the video image width, in pixels.
159
160 The area recorded on the ground ($gaFOV$, m²), can also be of interest:

161 (3) $$gaFOV = \frac{IH}{IW} ghFOV^2$$

162 where IH is the video image height, in pixels.
163
Based on these relations, we chose a 45 m UAV flight height, which covers a $ghFOV$ of 61 m
($gaFOV$ = 2096 m²), and a gPP of 16 mm (Fig. 1C). This pixel pitch value was voluntarily chosen
at a fraction of the body size of the *Pieris* species we wanted to track, which have a forewing
length around 30 mm.
We performed preliminary tests in the field that confirmed that at H = 45 m, *Pieris* butterflies
flying near the ground were projected in recorded images as pixel "blobs" with an area around
10 pixels, which is large enough to be reliably tracked from frame to frame (see Fig. 2A, C).
Higher camera height would allow larger FOV area on the ground, and hence longer tracking
durations, but would make automatic blob tracking less reliable.
Note that at H = 45m, the butterflies are not visible to the UAV pilot through the live video
feedback on the UAV controller screen. The video feedback has low resolution (720p/1080p),
and the controler screen (Ipad Mini 5, Apple, Cupertino, USA) has limited magnification and
contrast in outdoor conditions. Butterflies in UAV footage were only detectable *a posteriori*,
when playing recorded videos at full resolution on a computer screen in the lab.
All videos were recorded at 30 fps (29.97 frames per second), which in 4K resolution
produced video data at a rate of 13 MB/s. The UAV firmware automatically cut video files lasting
more than 5 min in multiple 4 GB bouts, which can complicate the post-processing workflow.
For this reason, we decided to keep each video duration below 5 min. Recording at 60 fps was
another possible option, but with a 15 MB/s data rate, this implied less data collected per video

frame. We sticked to 30 fps with the idea to obtain the best possible image quality per video
frame.
Manual exposure was tested initially, but proved unpractical as the ground luminance could
vary by several exposure values (Ev) when cloud shadows crossed the FOV. Hence, we used
auto-exposure with an exposure compensation of -0.7 to -1.3 Ev, as it improved the contrast
between the white butterflies and the background.

Time distribution of UAV flights

In order to cover the whole *Pieris* flight season, we chose to collect videos once a week, over
the months of June, July and September 2021. Each week, we chose a day that was favorable
for butterfly flight, i.e. with a weather forecast as warm, sunny and not very windy as possible.
We went on site in the afternoon (14:00-17:00). After takeoff, the UAV was positioned at $H = 45$
m above the EF. The UAV camera was tilted to a vertically-downward position, and the pilot
used the video feedback to place the scale points on the road to the left margin of the FOV (see
Fig. 1E). The pilot started the video recording, and the UAV then relied on its own sensors (GPS,
altimeter, etc.) to maintain its position without any pilot input, for about 4 min 30 s. Then the
video recording was stopped, the UAV was translated to the WF, height and scale alignment
(now on the right margin of the FOV, see Fig. 1D) were checked, and another 4 min 30 s video
was recorded. Repeating these steps, we could record a second EF video and a second WF video
before the UAV battery dropped to 20-30% of capacity, inciting landing the UAV and swapping
its battery. With this sequence, each UAV battery (rated at 11.6 V, 3500 mAh, 40.4 Wh) allowed
to record about 18 min of video. As we used 3 fully charged battery per field session, we were
able to collect about 1 hour of video per field session. Weather allowing, 3 field sessions could
take place in June (1st, 9th, 15th), 4 field sessions in July (1st, 8th, 15th, 22nd) and 4 in September
(6th, 16th, 22nd, 29th).

Video processing

In order to automate the tracking of butterflies in videos, pixels corresponding to the
butterfly should be easily distinguished from the background. In the present case, *Pieris* appear
in the raw video as blobs of bright pixels, but the background formed by the crops also has
multiple bright areas (Fig. 2A), which rules out a simple detection of the butterfly by
thresholding the raw video luminance levels. In addition, as the UAV is not perfectly static, and
the wind can cause plants on the ground to move slowly, the background is moving, which does
not favour background subtraction approaches (Piccardi 2004). We found a solution to this
issue by designing a custom filter that selects the pixels that blink in the video: when a butterfly
passes over an area, the pixels in that area become bright for only 1 frame, and then revert to
the background luminance. To apply this filter, we first transformed the RGB video frames into
greyscale. Each pixel then has a single luminance value (*v*) in the range [0, 255]. Then the "blink"
filter script performs the following calculations:
220
Pixel value variation from current frame (t) to next (t+1) and previous (t-1) video frames:

(4) $\Delta v_n = v_{t+1} - v_t$
(5) $\Delta v_p = v_{t-1} - v_t$

If both variations have the same sign, there was a luminance peak (positive or negative), and
a blink value (b) is computed. On the contrary, if variations have different signs, there was no
blinking, only monotonous pixel value variation.

(6)   if $sgn(\Delta v_n) = sgn(\Delta v_p)$ $\rightarrow$ $b_t = -sgn(\Delta v_n) * \min(|\Delta v_n|, |\Delta v_p|)$
(7)   else $\rightarrow$ $b_t = 0$

Note that b can be positive (bright blob passing over darker background) or negative (dark
blob passing over brighter background).
231
Interestingly, when going through a series of successive video frames, keeping memory of
extreme b values for each pixel (notated B) can be used to reveal flight trajectories as a series
of blobs, in a synthetic, diachronous image.

(8)   if $|b_{t+1}| > |b_t|$ $\rightarrow$ $B = b_{t+1}$
(9)   else $\rightarrow$ $B = b_t$

For easy display of b frames (and B synthetic images), pixel blink values must be rescaled
from [-255, 255] to [0, 255].

(10) $b = (b + 255)/2$

As a result, background appears as medium gray ($b = 128$), with blinking pixels as lighter or
darker gray blobs. Fig. 2 shows example results of the blink filter.
We used the blink filter in two ways. For the first exploration of our videos, we cut each video
into 20 s bouts, and generated a single B image summarising each bout. This enabled us to
quickly detect which video bouts containing butterfly tracks. On this basis, the FOV entry and
exit times of each butterfly were noted by replaying the original videos on a large screen. Based
on these time limits, we then generated a new greyscale video file, containing the series of b
frames corresponding to each butterfly track. This filtered greyscale video was then used to
automatically track the blobs.
We used DLTdv8 (Hedrick 2008) to extract blob centroid coordinates (in pixels) in successive
video frames. The blob was manually digitized (i.e. clicked) in the first few frames, and then
DLTdv8 uses a Kalman filter and 2D cross-correlation to find the blob in following frames
(without thresholding). As blob size and contrast can vary, and filtered b frames still contain
noise in the background, the automatic tracking process needed human supervision and
regular manual corrections, but with a convenient user interface to navigate through video
frames, DLTdv8 offered vast time savings compared to a fully manual digitization.
For scaling the butterfly track to real-world coordinates, we measured pixel coordinates of
the two painted reference crosses in one video frame (at mid-duration of the track), and fitted
a geometrical transformation (combining rotation, scaling and translation; *fitgeotrans* function
in Matlab) that resulted in (0, 0) and (0, -20) coordinates in real-world meters. This same
transformation was then applied to the whole butterfly coordinate series, transforming
coordinates in pixels to meters. We had measured with a compass in the field that the central
road   had   a   6°E   azimuth.   We   checked   and   refined   this   value   in   Geoportail

Figure 2 – Video frame filtering. **(A)** Magnified view of an original video frame, containing a *Pieris brassicae* image. **(B)** Same view after applying the blink filter (i.e. *b* frame). **(C)** A hypothetical 30 mm forewing length *Pieris*, with wings fully stretched, projected onto a pixel grid with a 16 mm *gPP*. If all partially covered pixels become brighter, the pixel blob area would be around 12 pixels. **(D)** Merging successive *b* frames reveals *P. brassicae* flight trace in a synthetic, diachronous image (i.e. *B* image). **(E)** Example trace of *Pieris rapae*. **(F)** Example trace of a dark species, either *Vanessa atalanta* or *Aglais io* (contrast ×2.0). **(G)** Example trace of *Melanargia galathea*. **(H)** Example trace where the shadow of a *Pieris* butterfly is also visible (contrast ×1.5).

Track selection

When we explored the *B* images synthesizing our videos, we mainly found flight traces of
Pieris butterflies (large white *P. brassicae*, small white *P. rapae*), appearing as clearly visible
white dotted traces (fig. 2D, E). These two species were also the most easily observed from the
ground during the field sessions. We also found a few flight traces of other species, that we had
observed in the field, such as the marbled white (*Melanargia galathea*), the red admiral
(*Vanessa atalanta*) or the peacock (*Aglais io*). However, as these traces were usually barely
visible and discontinuous (Fig. 2F, G), we did not analyse them further.

For some *Pieris* trajectories, the shadow of the butterfly formed a second, dark trace (Fig.
2H), which may potentially be used to determine the height of the butterfly relative to the
ground (knowing the current sun elevation). These shadows were mainly visible on almost bare
ground (September videos, after harvest), and only concerned a small number of *Pieris* traces.
We have therefore not used this additional source of 3D data in the present study.

Among the *Pieris* tracks, we applied the following exclusion criteria:
• A 10 m wide strip, containing the road and the adjacent flowered ditches, was excluded
from the analysis, so that each field (EF and WF) included a homogeneous area in terms
of ground nature.
• Very brief tracks (< 5 s), crossing just one corner of the FOV, were excluded.
• As we were mainly interested in migratory flights, defined as "directional flights in
which individuals appear undistracted by appetitive cues" (Chowdhury et al. 2021), we
focused on butterflies in continuous movement: Tracks marking one or more stops
while crossing the FOV were excluded (a stop being defined as remaining for at least 1
s within a radius of 0.1 m).
• Tracks in which the butterfly interacted with another individual (e.g. flight inflection
towards another butterfly, chases) were also excluded.

In the end, our sample comprised *N* = 166 *Pieris* flight tracks. Despite our constant sampling
efforts, these were not evenly distributed through the season: we recorded 12 tracks in June,
125 in July and 29 in September. Taking all these tracks together, 70524 butterfly positions were
recorded, representing a total of 39 min of flight time, and a flight distance covered of 7.4 km.

When the tracks contained missing positions (caused by the absence of blob on some *b*
frames), the (x, y) coordinates were linearly interpolated. These interpolated positions
represented 2.2 % of the dataset (1544 positions). The interpolated positions were rarely
contiguous, and the longest interpolated segment represented 7 successive positions (i.e. 0.23
s).

Track descriptive variables

312   For this pilot study, we computed a small set of basic descriptive variables for each 2D track.
313   • **Track duration** is the time the butterfly remained within the FOV (road zone
314   excluded).
315   • The change in 2D position from a frame to the next frame is named a step vector. **Track**
316   **length** is computed as the sum of step vector norms.

- 317 · Step speed is equal to step vector norm divided by the elapsed time (i.e. 1/29.97 s).
- 318 **Average speed** is the arithmetic mean of the series of step speed values. It indicates
- 319 how fast, on average, the butterfly flew along its 2D track. Average speed is also equal
- 320 to track length divided by track duration. Note that average speed is a ground speed,
- 321 not an air speed.
- 322 · A "beeline" vector is defined as the vector from the first to the last recorded position
- 323 of the butterfly. The beeline vector is equal to the vectorial sum of step vectors, and
- 324 hence represents the butterfly's resultant, directed movement across the FOV. **Beeline**
- 325 **azimuth** was the direction of the beeline vector, a circular variable in the interval [0°,
- 326 360°[, 0° corresponding to a northern azimuth.
- 327 · Track **straightness** is computed as the ratio of the beeline vector norm to the track
- 328 length. Straightness value is in the interval [0, 1]: 0 indicates that the butterfly
- 329 performed a loop (i.e. had the same entry and exit positions), while 1 indicates a
- 330 perfectly straight flight path. Straightness is inversely related to path tortuosity.
- 331 · **Beeline speed** is the norm of the beeline vector divided by track duration. It reflects
- 332 how fast the butterfly, on average, progressed in its directed movement. Beeline speed
- 333 is also equal to average speed multiplied by straightness. Note that beeline speed is
- 334 also a ground speed, not an air speed.

Processing and statistics software

Pixel blob tracking in videos was performed using DLTdv8 (Hedrick 2008;
https://biomech.web.unc.edu/dltdv/). All other analyses, from video processing to statistics,
were performed using Matlab R2018b (The MathWorks, Natick, MA, USA). We used the
CircStat2012a toolbox for circular statistics (Berens 2009). For a small number of captured
butterflies, forewing length measurement from field photographs were performed with ImageJ
V1.54g (http://imagej.org).

Results

Track general description

Fig. 3 shows the reconstructed 2D tracks, and associated variable distributions. We collected
66 Pierid flight tracks over WF, and 100 tracks over EF. Tracks had a median duration of 12.8 s
(range 5.1 to 56.5 s), for a median track length of 40.0 m (13.4 to 134.9 m). The median value for
average speed was 3.3 m.s^{-1} (1.9 to 9.2 m.s^{-1}), and 2.9 m.s^{-1} for beeline speed (0.2 to 9.1 m.s^{-1}).
Straightness distribution was strongly skewed towards straight tracks, with a median value of
0.93 (range 0.10 to 0.99). High straightness values cause similar values for average speed and
beeline speed in most tracks.

Figure 3 – *Pieris* flight tracks description. (A) Reconstructed 2D tracks over WF and EF. Black dots indicate last position of each track. **(B-F)** Distributions of descriptive variables: (B) track duration, (C) track length, (D) average speed, (E) track straightness and (F) beeline speed.

We used two-sample Kolmogorov-Smirnov (KS) tests to assess distribution differences between WF and EF tracks, and found no significant difference for track duration ($D_{(66,100)} = 0.10$, $p = 0.78$), track length ($D_{(66,100)} = 0.09$, $p = 0.86$), average speed ($D_{(66,100)} = 0.09$, $p = 0.86$), or beeline speed ($D_{(66,100)} = 0.18$, $p = 0.15$). However, there was a significant difference between straightness distribution over WF and EF ($D_{(66,100)} = 0.26$, $p = 0.009$), with straightness for EF tracks being even more skewed towards 1 (median 0.94) than WF tracks (median 0.89; see Fig. 3E).

Beyond average speed, direction and straightness, the 30 fps butterfly flight tracks extracted from UAV videos also contained fine-scale instantaneous information about flight speed and azimuth variation along the tracks. For example, Fig. 4 shows a track segment that depicts interesting movement patterns, in the form of meter-scale, sub-second transverse oscillations along the flight path.

Figure 4 – Benefit of 30 Hz positional data. The upper line is a 12 s flight track segment, reconstructed with the present method. The lower track is a copy (with a -3 m y offset for visualization) showing all recorded 2D positions (361 in total). The 5 circles represent the positions that could have been recorded with a SHR tracking system, that has a 90-fold lower sampling frequency (1 position every 3s.). Note that on the other hand, SHR benefits from a larger FOV, and hence longer tracking durations (see discussion).

Controlling UAV stability

During video recording, the UAV was not perfectly static, and whether this could have an effect on the reconstructed flight tracks was an important question. For a small random subset of tracking videos ($N = 10$), we digitized the two reference points painted on the road, describing a 20 m reference segment (RS), not only at mid-duration, but on every video frame, which allowed to monitor how the RS was transformed throughout the duration of tracking, due to UAV movements. We assumed that a combination of rotation, scaling and translation could affect the RS projection. Using the *fitgeotrans* function in Matlab, we obtained the geometric transformation matrix from the first frame's RS to each following frame, which allowed to monitor rotation, scaling and translation movement components separately.

Fig. 5 shows that the RS projected image was indeed affected by a combination of geometric transformations through time: over the investigated tracking durations (13 to 40 s), rotation of the RS could attain 0.25° (Fig. 5A), scaling variation could exceed ± 1 % (Fig. 5C), and horizontal (x) or vertical (y) translation of the RS image (Fig. 5B) could reach ± 10 pixels (i.e. 0.3 % of image width, or 0.16 m when projected on the ground). The amounts of transformation usually did not grow monotonically through time, reflecting that in the absence of pilot input, the UAV does not drift away from its initial position, but uses servoing based on its onboard sensors to try and maintain position and azimuth. Digitizing the RS only once at track mid-duration - i.e. assuming that the UAV is fully static during each track - can result in a ~ 0.2 m drift error on the butterfly reconstructed position (Fig. 5D). For the present work, we considered these levels of error to be acceptable, which is why we used the single RS digitizing method for all the remaining tracks.

Controlling butterfly flight height

With the present recording geometry, the flight tracks of all butterflies passing through the
pyramidal FOV (Fig. 1A), regardless of their flight height, are projected on a single sensor plane.
We reconstructed these tracks in 2D by assuming that the butterflies moved in the same plane
as our reference 20 m segment, i.e. flew at ground level. How this simplification departed from
the 3D reality needed investigation. We were especially interested in the possibility that some
butterflies might have crossed the FOV at a significant height (10-40 m), which would result in
overestimated flight speeds in our ground-projected reconstructions.

During field work, we visually monitored the field under the UAV, and noted each time we
saw a butterfly flying at a low height (defined as below the observer visual horizon, i.e. less than
2 m above the ground). By comparing our field notes with the timestamps of the reconstructed
tracks, N = 45 out of 166 tracks (27 %) corresponded to visually-detected butterflies flying at low
height (LH group). The remaining tracks were qualified as "unknown height" (UH group, N =
121), and may correspond either to high-flying butterflies (visually undetectable because of low
contrast against the sky), or to low-flying butterflies that remained undetected (because the
observer's attention was regularly directed at the UAV rather than at the ground).

We compared the beeline speed of the low-height (LH) and unknown-height (UH) groups.
Both groups had very similar speed distributions (Fig. 6A), as confirmed by statistical tests : LH
(2.9 ± 1.0 m.s^{-1}, mean ± SD) and UH tracks (3.1 ± 1.3 m.s^{-1}) did not significantly differ for mean
speed in a t-test ($t_{(164)}$ = 0.83, p = 0.4), and were not drawn from significantly different
distributions according to a KS test ($D_{(45,121)}$ = 0.11, p = 0.79).

Figure 6 – Speed distribution and flight height. (A) Beeline speed distribution observed for low height (LH) and unknown height (UH) butterfly tracks. **(B)** Kernel probability density estimation for LH and UH tracks, and for simulated tracks with uniform height distribution (see text).

In order to visualize what speed distribution would be obtained for butterflies flying far from the ground, we simulated 10^5 straight horizontal tracks, with flight speeds sampled from a normal distribution copied form the LH group (2.9 ± 1.0 m.s^{-1}), but crossing the FOV at heights uniformly distributed between 0 and 45 m. Simulated tracks crossing the FOV, and with a track duration of at least 5 s (to simulate comparable track selection), had a ground-projected speed distribution that was flatter and shifted towards high (i.e. overestimated) speeds when compared to both LH and UH tracks (see Fig. 6B). This further suggests that the UH group in our track sample mainly comprises low-flying butterflies that went visually undetected in the field. Still, we cannot exclude that a small number of tracks in the UH group (on the right tail of speed distribution, i.e. with reconstructed speed > 6 m/s) might have been high-flying butterflies.

Exploring blob area as a specific signature

We explored the possibility to discriminate *P. brassicae* from *P. rapae* tracks, based on pixel blob area in the videos. During field work we captured a small number (*N* = 8) of *Pieris sp.* individuals that had just passed through the UAV's FOV. Each individual, captured with a butterfly net, was briefly placed in a thin transparent box with a grid-patterned back, photographed with its wings stretched for later identification and size measurement, and then immediately released in the field. Back in the lab, we measured each individual's forewing length using ImageJ. The UAV video recordings corresponding to these individuals were filtered and digitized with the same methods as previously described, but were later re-analysed to measure the blob size in each filtered video frame. As a simple approach, the blob was defined as connected pixels with grey level *b* > 138, i.e. departing by more than 10 grey level from the mean background grey of 128. Sometimes the simple thresholding approach detected no blob,

but this method still allowed to obtain many (269 to 654) blob area values per individual, that could be compared to the animal's real size. Note that 4 out of the 8 tracks used for the present blob area analysis were not part of the final flight track sample ($N = 166$), because these captured individuals had flown near the road, or performed stops along their flight (see Track selection section).

Five captured individuals were identified as *P. rapae* (2 females + 3 males), with forewing length ranging from 22.8 to 28.0 mm. The 3 other captured individuals were *P. brassicae* (2 females + 1 male), with longer forewing, ranging from 31.2 to 36.7 mm. These values were in line with the literature, with the largest *P. rapae* individuals being close in size to the smallest *P. brassicae* specimens (Cook et al. 2022)

Figure 7 - Forewing length vs. pixel blob area distribution for 8 tracked and captured *Pieris* butterflies. Blue: *P. brassicae* ($N = 3$); Green: *P. rapae* ($N = 5$). Distributions are displayed as Kernel probability density estimates. Circles indicate the median blob area value for each butterfly.

Fig. 7 shows the blob area distributions observed for all 8 captured individuals. When considering only the median blob area for each individual, it was positively correlated to the forewing length of the animal (Spearman rank correlation, $r_{(6)} = 0.73$, $p = 0.047$). However, the relationship was not monotonically increasing ($r < 1$), and there was extensive overlap between blob area distributions. In other words, a smaller butterfly could often project as a larger pixel blob than a larger butterfly, depending on the compared video frames. As a result, we considered unreliable to use recorded pixel blob areas as a direct mean to discriminate *P. rapae* and *P. brassicae* flight tracks in the present work.

Proof of concept: Effect of season on migratory flight azimuth.

By observing the tracks at different stages of the flight period, it appeared visually that the flight azimuths have varied over the season (Fig. 8).

Figure 8 – *Pieris* flight tracks broken down by flight period (vertically) and field (horizontally). Black dots indicate last position of each track (N = 166). 30 m diameter circles represent the area considered for azimuth statistical comparisons (N = 119).

To further quantify seasonal variations in azimuth distributions, we focused on tracks
crossing a 30 m diameter disc located at the centre of the camera's FOV (Fig. 8). Indeed, the
rectangular shape of the camera FOV is less likely to record trajectories parallel to the longer
side of the FOV, and this bias must be corrected by considering only a circular surface inside the
FOV. When a track did not cross this central disc, it was therefore removed from the sample for
circular statistics. When a track crossed this disc more than once, only the longest segment
inside the disc was considered. This restriction of the FOV to a central disc had the effect of
reducing our sample from $N = 166$ to $N = 119$ (for this section only).
We tested the effects of period (June to September) and field (WF vs. EF) on beeline azimuth,
using a two-factor ANOVA for circular data (Harrison & Kanji 1988, in Berens 2009). As we
observed many more trajectories in July, we subdivided the month of July: *early July* (field
sessions on July 1st and 8th) and *late July* (July 15th and 21st). We then tested the uniformity of
the azimuth distributions for each period, using Rayleigh tests (Fisher 1995, in Berens 2009).

Figure 9 – Circular distributions of butterfly beeline azimuth, broken down by period
(vertically), and field (horizontally). Arrows represent resultant vectors. The left column
shows azimuth distributions for both fields pooled (WF + EF), with red arrows
representing significant resultant vectors after a Rayleigh test.

The ANOVA for circular data detected a significant effect of period on azimuth ($X^2_{(6)}$ = 43, p = 1.2 × 10^{-7}), but no effect of field side ($X^2_{(2)}$ = 0.51, p = 0.77), and no interaction between the two factors ($X^2_{(3)}$ = 5.5, p = 0.14). The distribution of azimuths during the 4 periods is shown in Fig. 9. In June, the butterflies flew most often to the north-east (Rayleigh test, N = 7, R = 0.65, p = 0.04; mean azimuth 24°). In early July, they flew most often to the north-west (N = 51, R = 0.56, p = 3.8×10^{-8}; mean azimuth 317°). In late July, tracks in many directions were observed, without any dominant azimuth, so that the azimuth distribution was not significantly different from a homogeneous distribution (N = 42, R = 0.15, p = 0.41). Finally, in September, tracks were most often oriented to the south (N = 19, R = 0.59, p = 0.001; mean azimuth 173°).

To verify if the above results were influenced by wind direction, we first tested whether butterfly beeline azimuth was correlated with wind direction, using a circular-circular correlation test (Jammalamadaka & Sengupta 2001, in Berens 2009). The test returned no significant correlation (c = - 0.01, p = 0.95; Fig. 10A). We also assessed whether butterfly preferentially flew downwind, crosswind or upwind, by computing the angular difference between beeline azimuth and wind direction, and testing whether this "angle to wind" variable departed from uniform distribution. Results suggested that butterflies did not preferentially fly at a specific angle to wind (Rayleigh test, N = 119, R = 0.15, p = 0.06; Fig. 10B), but the test was close to statistical significance, despite a small resultant vector (i.e. small effect size).

As a supplementary verification, we focused on tracks recorded during stronger winds (> 2 m.s^{-1}, N = 64), as these butterflies were expected to be the first to be affected by a possible wind influence. Both circular-circular correlation (c = - 0.04, p = 0.75) and Rayleigh test on angle to wind (N = 64, R = 0.08, p = 0.67) returned non-significant results (Fig. 10C, D). This comforted the conclusion that wind direction did not significantly bias butterfly flight azimuth in our data.

Figure 10 – Wind and flight azimuth. (A) Beeline azimuth vs. wind direction for N = 119 tracks. **(B)** Angle to wind for N = 119 tracks. 0° corresponds to upwind flight, 180° to downwind flight. **(C-D)** Same graphics for a N = 64 subset of tracks where wind speed exceeded 2 m.s^{-1}.

Discussion

Method validation, strengths and limits

We show that UAV-image-based tracking can be used to reconstruct free-flying butterfly
paths. The application of UAVs for butterfly movement study was proposed previously (Ivosevic
et al. 2017), but remained to be explored in reality. With relatively light, affordable and easily-
deployed gear in the field, we were able to track numerous wild, untagged *Pieris* butterflies,
over distances averaging 40 m. Given an average flight speed of about 3 m.s^{-1}, this translates to
tracking durations most often near 10-20 s, depending on each butterfly's flight speed and
straightness. This scale of recorded movement is larger than what can be achieved in most
insectaries, but remains modest compared to other open field methods (e.g. approximately 500
m for SHR; Ovaskainen et al. 2008, Lihoreau et al. 2012), and is directly constrained by the UAV
camera FOV.

In the present study, the FOV was voluntarily limited to 61 × 34 m (from a 45 m UAV height),
in order to maintain a butterfly blob area around 10 pixels, which was useful for reliable blob
tracking throughout video frame series (see Methods section). A first possible way to enlarge
FOV would be to use an UAV with higher camera resolution (e.g. 6K or 8K sensor, with matching
quality optics), which could potentially fly higher and record a wider ground area, while
maintaining a centimetric pixel pitch on the ground. Moreover, a higher signal-to-noise ratio in
video frames (e.g. from larger sensors and/or less compressed video file formats) might allow
to reliably detect smaller blobs throughout frame series, which in turn would allow larger pixel
pitch values, and even higher UAV flight height. A FOV exceeding 100 m in side length is probably
already possible with current high-end commercial UAVs, but with a heavier, less affordable
UAV, and proportional amounts of video data to process.

An alternative way to greatly increase recorded movement length would be to try and follow
butterflies with the UAV. This would necessitate (1) that the video feedback to the UAV pilot is
of sufficient magnification for a live view of individual butterflies and (2) a different approach to
video frame filtering, but these are interesting perspectives for future methodological
developments.

On the other hand, the main strength of the movement data we collected is the 30 Hz
temporal resolution, which is 2-3 orders of magnitude higher than non-image-based field
tracking methods applicable to flying insects: 0.33 Hz for SHR (Ovaskainen et al. 2008) and lower
(usually < 0.01 Hz) for automated radio-tracking (Kays et al. 2011). Here, with one location every
~10 cm along the flight path, our reconstructed tracks reveal fine-scale movement patterns (see
Fig. 2, 4, 8) that may provide interesting insights on biomechanical and/or navigational
processes at work during butterfly flight. An analysis focused on the oscillations along *Pieris*
flight paths is envisaged, but was beyond the scope of the present methodological study.

Another limitation of the present method is the projection of the FOV, a pyramidal 3D air
volume that butterflies can cross at various height, onto a theoretical 2D surface at ground
level. In the present dataset, we were able to verify by direct observation that at least 27 % of
tracked butterflies flew at less than 2 m above ground level. For these tracks, the 2D ground
projection implies only a small error on flight speed (i.e. an overestimation of less than 2 / (45 -

2) = 4.7 %). For other butterflies, for which we were unable to confirm flight height, we showed
that their projected flight speeds are no higher than confirmed "low flyers", and thus conclude
that they probably also crossed the FOV at low height (Fig. 6). This remains indirect evidence,
and we cannot exclude that for some rare tracks, we may have significantly overestimated flight
speed because the butterfly crossed the FOV at a higher height.

Worse, whatever the average flight height, all the vertical (z) movements of the butterflies,
even on a smaller scale, are lost when projected onto a (x, y) horizontal plane. For following
studies, it is therefore advisable to observe the 3D flight behaviour of butterflies beforehand,
depending on the species under consideration and the relief of the terrain, and to estimate
whether a 2D projection might lose important information about the animals' movements.

Still, in an hypothetical case where butterflies move mainly in horizontal planes, but at
various heights, it can be noted that the reconstruction of flight speeds will be heterogeneously
overestimated, but that angular and temporal variables (e.g. variation of azimuth through time)
remain unaffected by flight height.

If the 3D flight trajectory and/or elevation relative to the ground are really necessary for a
study (e.g. biomechanics), one can either try to find a study zone where the butterfly's shadow
is also visible in the image (Fig. 2H), or use other natively 3D optical methods, based on multiple
views of the flight volume (Theriault et al. 2014, de Margerie et al. 2015).

When the UAV receives no command from the pilot, its flight algorithm seeks to maintain its
horizontal position, height and azimuth, using sensory-motor regulation loops based on on-
board sensors (GPS, barometric altimeter, magnetic compass, accelerometers and
gyroscopes). It is probably on this point that commercial UAVs have made the most spectacular
progress in the last decade, and we were able to verify that, at least over the tracking times used
here (< 1 min), the movements of our UAV were indeed very limited (Fig. 5). This allowed us to
assume that the drone was static in the sky, at the cost of a drift of the order of 10-20 cm in the
reconstructed position of the butterfly, which we found acceptable for the question posed here
(i.e. the measurement of the beeline azimuth over several tens of meters). However, note that
most UAVs use barometric sensors for regulating flight height, and that atmospheric pressure
might vary significantly across flight times longer than a few minutes.

For studies requiring lower error caused by the drone's position, it remains possible to
repeat the measurement of the reference scale on every video frame, or even to perform a more
refined calibration of the projected image by continuously tracking multiple points on the
ground. If the drone's movements really need to be more closely controlled, more advanced
commercial drones are already available, using positioning technologies such as Differential
GPS (DGPS) or Real Time Kinematic (RTK), which can monitor the UAV position with centimeter
accuracy.

Our video tests suggest that many more butterfly species are potentially detectable using
the present method, whether they appear light against a dark background or dark against a
light background (Fig. 2). The blink filter proved efficient for erasing background textures, while
also tolerating a fair amount of movement in the background, caused either by the slow
movements of the UAV, or crop plants being blown over by the wind. Hence, we hope to see
following studies tracking other butterfly species in various landscapes. Unfortunately,

depending on the location and the flight season, it is possible to run into the problem that two morphologically similar species get filmed simultaneously, and it is precisely this problem that we encountered here with two *Pieris* species of similar aspects. We were not able to discriminate *P. brassicae* from *P. rapae* based on pixel blob area (Fig. 7). The wide, overlapping distributions of blob areas for each individual butterfly is not surprising: Butterflies can have any posture, from fully stretched to fully closed wings on different video frames, as a result of (1) flapping wing movement and (2) variable roll and pitch angles of the body along the flight path. Moreover, pixel blob area can also be affected by other factors such as (3) contrast with the local background, (4) "blob-clipping" (when the distance covered between two consecutive frames is less than the body length, which interferes with the filtering method), and (5) flight height. Note also that the relationship between projected wing area and blob area is expected to be subject to discrete steps, especially when the pixel pitch is close to animal body size (Fig. 2C). Provided that some of these sources of blob area variation can be better controlled, we do not rule out that pixel blob area could help discriminate similar butterfly species in future studies, but this would need more refined blob classification processes than what we could implement here.

Another route for reducing possible confusion between species could be a greater effort to identify individuals by systematically capturing them after they crossed the FOV, but this would however be limited to low-flying butterflies. This would also imply greater human presence and movements in the field, and probably additional precautions to avoid affecting the butterflies' movement behaviour.

P. brassicae and *P. rapae* migratory behaviours

Migratory behaviour has been documented in about 600 butterfly species (Chowdhury 2021, 2022). In butterflies, migration often occur over several generations, with successive generations following different flight azimuths to achieve a round-trip annual travel that can cover thousands of kilometers in some Nymphalidae species (e.g. monarch *Danaus Plexippus*: Brower 1995; painted lady *Vanessa cardui*: Stefanescu et al. 2013, Suchan et al. 2024).

Within Pieridae, another butterfly family with numerous migratory species (Chowdhury 2021), *P. brassicae* and *P. rapae* migratory behaviours have long been documented from observation of mass migrations (e.g. Vepsalainen 1968, now rarer due to the use of pesticides, but see John et al. 2008), and visually measured flight azimuth in the field (e.g. Baker 1968). More recently, these species' migratory behaviours have been studied by more experimental methods. Jones et al (1980) used a mark-recapture method: they fed *P. rapae* larvae with a special diet causing the adult females to later lay colored eggs, which allowed to study the movements of individual females over a 1 km square grid of cabbage plants. They reported that female *P. rapae* have a directed flight, but that the preferred azimuth varied from day to day. Gilbert & Raworth (2005) completed this work with field observation in the Pyrenees, and confirmed that both male and female *P. rapae* can migrate northward in spring and southward in the autumn, although some individuals and local populations did not appear to migrate. Spieth & Cordes (2012) collected eggs of *P. brassicae* from several Western Europe regions, and later measured the spontaneous flight azimuth of female individuals, in a 2 m diameter flight cage. They showed that the preferred flight azimuth depended on the season and the geographic origin. In regions where *P. brassicae* has a cycle of 2 generations per year, the first

generation followed a northern azimuth (modulated by the precise geographic origin), whereas
the last generation flew southward. In regions where *P. brassicae* has 3 generations per year,
both the first and second generation flew northward, and only the last generation had a reverse,
southward flight azimuth.

Compared with these pre-existing data on migration in *P. brassicae* and *P. rapae*, flight
tracks recorded from a UAV allow more accurate measurement of flight azimuth, without the
need to capture or mark the butterflies (which might influence spontaneous flight behaviour),
at an intermediate scale between a flight cage (or insectary) and mark-recapture data. For a
relatively large sample of individuals from both species, we measured flights that were mainly
directed northwards in the early season (June, early July), and a return southward azimuth in
the late season (September). Our azimuth results under natural conditions are therefore
congruent with the existing literature for both *P. brassicae* (Spieth & Cordes 2012) and *P. rapae*
(Gilbert & Raworth 2005).

As a bonus, our high sampling frequency trajectories appear to contain previously
unavailable fine-scale information on *Pieris* flight speed, tortuosity of the flight path, and
patterns of rapid azimuth variation (transverse oscillations). This refined data may be studied
in greater detail and perhaps reveal information on locomotor behaviour and the perceptual
mechanisms underlying *Pieris* spatial behaviour, whether migratory or not. A major limitation
of our approach at this stage remains the confusion between *P. rapae* and *P. brassicae*, because
of their similar morphologies and colors, and their concurrent flight in our geographical area.

Conclusion

Our results reveal that video tracking of butterflies from a UAV is possible, and capable of
providing movement data in fully natural conditions, at an unprecedented spatio-temporal
resolution, and at a modest cost. This fine-scale data could prove invaluable for understanding
the spatial behaviour of many butterfly species in the wild, which has so far been only partially
documented (Chowdhury 2021, 2022) as it remains a technical challenge to track to such small
species in the field. We hope that the present methodology exploration can serve as a starting
point and motivate following studies on spatial behaviour and movement ecology in flying
insects.

Acknowledgments

We thank the owner of the crop fields for authorizing our presence and UAV flights over his
property. We thank Tristan Lafranchis for discussions on the behaviour and biology of *Pieris*
brassicae and *Pieris rapae*. We thank Ty Hedrick for discussions on the use of DLTdv. We thank
Virginie Durier and Simon Benhamou for discussions on circular statistics.

Funding

The authors declare that they have received no specific funding for this study

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Links to be specified in the final version of the manuscript

References

Abrahms B, Seidel DP, Dougherty E, Hazen EL, Bograd SJ, Wilson AM, Weldon McNutt J, Costa DP, Blake S, Brashares JS, Getz WM (2017) Suite of simple metrics reveals common movement syndromes across vertebrate taxa. *Movement Ecology*, **5**, 12. https://doi.org/10.1186/s40462-017-0104-2

Baker R (1968) A possible method of evolution of the migratory habit in butterflies. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **253**, 309–341. https://doi.org/10.1098/rstb.1968.0003

Batsleer F, Bonte D, Dekeukeleire D, Goossens S, Poelmans W, Van der Cruyssen E, Maes D, Vandegehuchte ML (2020) The neglected impact of tracking devices on terrestrial arthropods. *Methods in Ecology and Evolution*, **11**, 350–361. https://doi.org/10.1111/2041-210X.13356

Berens P (2009) CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software*, **31**, 1–21. https://doi.org/10.18637/jss.v031.i10

Brower LP (1995) Understanding and misunderstanding the migration of the monarch butterfly (Nymphalidae) in North America: 1857-1995. *Journal of the Lepidopterists' Society*, **49**, 304--385.

Brussard PF, Ehrlich PR (1970) Adult Behavior and Population Structure in *Erebia Epipsodea* (Lepidoptera: Satyrinae). *Ecology*, **51**, 880–885. https://doi.org/10.2307/1933981

Cagnacci F, Boitani L, Powell RA, Boyce MS (2010) Animal ecology meets GPS-based radiotelemetry: a perfect storm of opportunities and challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 2157–2162. https://doi.org/10.1098/rstb.2010.0107

Charbonneau P, Lemaître J (2021) Revue des applications et de l'utilité des drones en conservation de la faune. *Le Naturaliste canadien*, **145**, 3–34. https://doi.org/10.7202/1080620ar

Chowdhury S, Fuller RA, Dingle H, Chapman JW, Zalucki MP (2021) Migration in butterflies: a global overview. *Biological Reviews*, **96**, 1462–1483. https://doi.org/10.1111/brv.12714

Chowdhury S, Zalucki MP, Amano T, Poch TJ, Lin M-M, Ohwaki A, Lin D-L, Yang L, Choi S-W, Jennions MD, Fuller RA (2022) Trends and progress in studying butterfly migration. *Integrative Conservation*, **1**, 8–24. https://doi.org/10.1002/inc3.13

Cook PM, Tordoff GM, Davis TM, Parsons MS, Dennis EB, Fox R, Botham MS, Bourn NAD (2022) Traits data for the butterflies and macro-moths of Great Britain and Ireland. *Ecology*, **103**, e3670. https://doi.org/10.1002/ecy.3670

Crall JD, Gravish N, Mountcastle AM, Combes SA (2015) BEEtag: A Low-Cost, Image-Based
Tracking System for the Study of Animal Behavior and Locomotion. *PLOS ONE*, **10**, e0136487.
https://doi.org/10.1371/journal.pone.0136487
Da Silveira NS, Niebuhr BBS, Muylaert R de L, Ribeiro MC, Pizo MA (2016) Effects of Land Cover
on the Movement of Frugivorous Birds in a Heterogeneous Landscape. *PLOS ONE*, **11**,
e0156688. https://doi.org/10.1371/journal.pone.0156688
Delattre T, Vernon P, Burel F (2013) An agri-environmental scheme enhances butterfly dispersal
in European agricultural landscapes. *Agriculture, ecosystems & environment*, **166**, 102–109.
Dell AI, Bender JA, Branson K, Couzin ID, Polavieja GG de, Noldus LPJJ, Pérez-Escudero A,
Perona P, Straw AD, Wikelski M, Brose U (2014) Automated image-based tracking and its
application in ecology. *Trends in Ecology & Evolution*, **29**, 417–428.
https://doi.org/10.1016/j.tree.2014.05.004
Fernández P, Rodríguez A, Obregón R, de Haro S, Jordano D, Fernández-Haeger J (2016) Fine
Scale Movements of the Butterfly *Plebejus argus* in a Heterogeneous Natural Landscape as
Revealed by GPS Tracking. *Journal of Insect Behavior*, **29**, 80–98.
https://doi.org/10.1007/s10905-016-9543-7
Gilbert N, Raworth DA (2005) Movement and migration patterns in *Pieris rapae* (Pieridae).
Journal of the Lepidopterists' Society, **59**, 10–18.
Haalck L, Mangan M, Webb B, Risse B (2020) Towards image-based animal tracking in natural
environments using a freely moving camera. *Journal of neuroscience methods*, **330**, 108455.
Hedrick TL (2008) Software techniques for two- and three-dimensional kinematic
measurements of biological and biomimetic systems. *Bioinspiration & Biomimetics*, **3**,
034001. https://doi.org/10.1088/1748-3182/3/3/034001
Hedrick TL, Pichot C, de Margerie E (2018) Gliding for a free lunch: biomechanics of foraging
flight in common swifts (*Apus apus*). *Journal of Experimental Biology*, **221**, jeb186270.
https://doi.org/10.1242/jeb.186270
Hernández-Pliego J, Rodríguez C, Dell'Omo G, Bustamante J (2017) Combined use of tri-axial
accelerometers and GPS reveals the flexible foraging strategy of a bird in relation to weather
conditions. *PLOS ONE*, **12**, e0177892. https://doi.org/10.1371/journal.pone.0177892
Ivosevic B, Han Y-G, Kwon O (2017) Monitoring butterflies with an unmanned aerial vehicle:
current possibilities and future potentials. *Journal of Ecology and Environment*, **41**, 12.
https://doi.org/10.1186/s41610-017-0028-1
Jackson BE, Evangelista DJ, Ray DD, Hedrick TL (2016) 3D for the people: multi-camera motion
capture in the field with consumer-grade cameras and open source software. *Biology Open*,
5, 1334–1342. https://doi.org/10.1242/bio.018713
John E, Cottle N, McArthur A, Makris C (2008) Eastern Mediterranean migrations of the Small
White *Pieris rapae* (Linnaeus, 1758) (Lepidoptera: Pieridae): observations from Cyprus 2001
and 2007. *Entomologist's Gazette*, **59**, 71–78.
Jones RE, Gilbert N, Guppy M, Nealis V (1980) Long-Distance Movement of *Pieris rapae*. *Journal
of Animal Ecology*, **49**, 629–642. https://doi.org/10.2307/4268
Joo R, Picardi S, Boone ME, Clay TA, Patrick SC, Romero-Romero VS, Basille M (2022) Recent
trends in movement ecology of animals and human mobility. *Movement Ecology*, **10**, 26.
https://doi.org/10.1186/s40462-022-00322-9

Kashetsky T, Avgar T, Dukas R (2021) The Cognitive Ecology of Animal Movement: Evidence
From Birds and Mammals. *Frontiers in Ecology and Evolution*, **9**.
https://doi.org/10.3389/fevo.2021.724887

Kays R, Tilak S, Crofoot M, Fountain T, Obando D, Ortega A, Kuemmeth F, Mandel J, Swenson G,
Lambert T, Hirsch B, Wikelski M (2011) Tracking Animal Location and Activity with an
Automated Radio Telemetry System in a Tropical Rainforest. *The Computer Journal*, **54**,
1931–1948. https://doi.org/10.1093/comjnl/bxr072

Kissling WD, Pattemore DE, Hagen M (2014) Challenges and prospects in the telemetry of
insects. *Biological Reviews*, **89**, 511–530. https://doi.org/10.1111/brv.12065

Kitamura T, Imafuku M (2015) Behavioural mimicry in flight path of Batesian intraspecific
polymorphic butterfly *Papilio polytes*. *Proceedings of the Royal Society B: Biological Sciences*,
282, 20150483. https://doi.org/10.1098/rspb.2015.0483

Knight SM, Pitman GM, Flockhart DTT, Norris DR (2019) Radio-tracking reveals how wind and
temperature influence the pace of daytime insect migration. *Biology Letters*, **15**, 20190327.
https://doi.org/10.1098/rsbl.2019.0327

Knoblauch A, Thoma M, Menz MH (2021) Autumn southward migration of dragonflies along the
Baltic coast and the influence of weather on flight behaviour. *Animal Behaviour*, **176**, 99–109.

Le Roy C, Amadori D, Charberet S, Windt J, Muijres FT, Llaurens V, Debat V (2021) Adaptive
evolution of flight in *Morpho* butterflies. *Science*, **374**, 1158–1162.
https://doi.org/10.1126/science.abh2620

Lihoreau M, Ings TC, Chittka L, Reynolds AM (2016) Signatures of a globally optimal searching
strategy in the three-dimensional foraging flights of bumblebees. *Scientific Reports*, **6**, 30401.
https://doi.org/10.1038/srep30401

Lihoreau M, Raine NE, Reynolds AM, Stelzer RJ, Lim KS, Smith AD, Osborne JL, Chittka L (2012)
Radar Tracking and Motion-Sensitive Cameras on Flowers Reveal the Development of
Pollinator Multi-Destination Routes over Large Spatial Scales. *PLOS Biology*, **10**, e1001392.
https://doi.org/10.1371/journal.pbio.1001392

Maggiora R, Saccani M, Milanesio D, Porporato M (2019) An Innovative Harmonic Radar to Track
Flying Insects: the Case of *Vespa velutina*. *Scientific Reports*, **9**, 11964.
https://doi.org/10.1038/s41598-019-48511-8

de Margerie E, Pichot C, Benhamou S (2018) Volume-concentrated searching by an aerial
insectivore, the common swift, *Apus apus*. *Animal Behaviour*, **136**, 159–172.
https://doi.org/10.1016/j.anbehav.2017.11.026

de Margerie E, Simonneau M, Caudal J-P, Houdelier C, Lumineau S (2015) 3D tracking of animals
in the field using rotational stereo videography. *The Journal of Experimental Biology*, **218**,
2496–2504. https://doi.org/10.1242/jeb.118422

Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE (2008) A movement
ecology paradigm for unifying organismal movement research. *Proceedings of the National*
Academy of Sciences, **105**, 19052–19059. https://doi.org/10.1073/pnas.0800375105

Oelmann Y, Fiedler D, Michaelis R, Leivits M, Braun A, Gschwind P, Neidhardt H, Willigalla C
(2023) Autumn migration of the migrant hawker (*Aeshna mixta*) at the Baltic coast. *Movement*
Ecology, **11**, 52. https://doi.org/10.1186/s40462-023-00415-z

Ovaskainen O, Smith AD, Osborne JL, Reynolds DR, Carreck NL, Martin AP, Niitepõld K, Hanski I
(2008) Tracking butterfly movements with harmonic radar reveals an effect of population

age on movement distance. *Proceedings of the National Academy of Sciences*, **105**, 19090–19095. https://doi.org/10.1073/pnas.0802066105

Piccardi M (2004) Background subtraction techniques: a review. In: *2004 IEEE International Conference on Systems, Man and Cybernetics (IEEE Cat. No.04CH37583)* , pp. 3099–3104 vol.4. https://doi.org/10.1109/ICSMC.2004.1400815

Rhodes MW, Bennie JJ, Spalding A, ffrench-Constant RH, Maclean IMD (2022) Recent advances in the remote sensing of insects. *Biological Reviews*, **97**, 343–360. https://doi.org/10.1111/brv.12802

Rieucau G, Kiszka JJ, Castillo JC, Mourier J, Boswell KM, Heithaus MR (2018) Using unmanned aerial vehicle (UAV) surveys and image analysis in the study of large surface-associated marine species: a case study on reef sharks *Carcharhinus melanopterus* shoaling behaviour. *Journal of Fish Biology*, **93**, 119–127. https://doi.org/10.1111/jfb.13645

Rotics S, Kaatz M, Resheff YS, Turjeman SF, Zurell D, Sapir N, Eggers U, Flack A, Fiedler W, Jeltsch F, Wikelski M, Nathan R (2016) The challenges of the first migration: movement and behaviour of juvenile vs. adult white storks with insights regarding juvenile mortality. *Journal of Animal Ecology*, **85**, 938–947. https://doi.org/10.1111/1365-2656.12525

Ruaux G, Monmasson K, Hedrick TL, Lumineau S, de Margerie E (2023) Flight behaviours and energy savings in adult and juvenile house martins (*Delichon urbicum*) foraging near their breeding colony. *Behavioral Ecology and Sociobiology*, **77**, 63. https://doi.org/10.1007/s00265-023-03332-8

Shepard ELC, Lambertucci SA, Vallmitjana D, Wilson RP (2011) Energy Beyond Food: Foraging Theory Informs Time Spent in Thermals by a Large Soaring Bird. *PLOS ONE*, **6**, e27375. https://doi.org/10.1371/journal.pone.0027375

Sherub S, Bohrer G, Wikelski M, Weinzierl R (2016) Behavioural adaptations to flight into thin air. *Biology Letters*, **12**, 20160432. https://doi.org/10.1098/rsbl.2016.0432

Spieth H, Cordes R (2012) Geographic comparison of seasonal migration events of the large white butterfly, *Pieris brassicae*. *Ecological Entomology*, **37**. https://doi.org/10.1111/j.1365-2311.2012.01385.x

Srygley RB, Oliveira EG (2001) Sun Compass and Wind Drift Compensation in Migrating Butterflies. *The Journal of Navigation*, **54**, 405–417. https://doi.org/10.1017/S0373463301001448

Stefanescu C, Páramo F, Åkesson S, Alarcón M, Ávila A, Brereton T, Carnicer J, Cassar LF, Fox R, Heliölä J, Hill JK, Hirneisen N, Kjellén N, Kühn E, Kuussaari M, Leskinen M, Liechti F, Musche M, Regan EC, Reynolds DR, Roy DB, Ryrholm N, Schmaljohann H, Settele J, Thomas CD, van Swaay C, Chapman JW (2013) Multi-generational long-distance migration of insects: studying the painted lady butterfly in the Western Palaearctic. *Ecography*, **36**, 474–486. https://doi.org/10.1111/j.1600-0587.2012.07738.x

Stürzl W, Zeil J, Boeddeker N, Hemmi JM (2016) How wasps acquire and use views for homing. *Current Biology*, **26**, 470–482.

Suchan T, Bataille CP, Reich MS, Toro-Delgado E, Vila R, Pierce NE, Talavera G (2024) A trans-oceanic flight of over 4,200 km by painted lady butterflies. *Nature Communications*, **15**, 5205. https://doi.org/10.1038/s41467-024-49079-2

Theriault DH, Fuller NW, Jackson BE, Bluhm E, Evangelista D, Wu Z, Betke M, Hedrick TL (2014)
A protocol and calibration method for accurate multi-camera field videography. *The Journal*
of Experimental Biology, **217**, 1843–1848. https://doi.org/10.1242/jeb.100529
Vepsäläinen K (1968) The immigration of Pieris brassicae L. (Lep., Pieridae) into Finland in 1966,
with a general discussion on insect migration. *Annales Entomologici Fennici*, **34**, 223–243.
Walter T, Degen J, Pfeiffer K, Stöckl A, Montenegro S, Degen T (2021) A new innovative real-time
tracking method for flying insects applicable under natural conditions. *BMC Zoology*, **6**, 35.
https://doi.org/10.1186/s40850-021-00097-3
Wang D, Shao Q, Yue H (2019) Surveying Wild Animals from Satellites, Manned Aircraft and
Unmanned Aerial Systems (UASs): A Review. *Remote Sensing*, **11**, 1308.
https://doi.org/10.3390/rs11111308
Whitfield DP, Fielding AH, Anderson D, Benn S, Reid R, Tingay R, Weston E (2024) Sex difference
in natal dispersal distances of Golden Eagles *Aquila chrysaetos* in Scotland. *Ibis*, **166**, 146–
155. https://doi.org/10.1111/ibi.13225
Wilson RP, McMahon CR (2006) Measuring devices on wild animals: what constitutes acceptable
practice? *Frontiers in Ecology and the Environment*, **4**, 147–154. https://doi.org/10.1890/1540-
9295(2006)004[0147:MDOWAW]2.0.CO;2