Butterfly movements recorded from a UAV depict migratory flight behaviour in the wild.

Emmanuel de Margerie*[1], Kyra Monmasson[1]

[1] CNRS, Univ. Rennes, UniCaen, UMR 6552, Rennes, France

* Corresponding Author
Correspondence: emmanuel.demargerie@univ-rennes.fr

Abstract

Tracking and understanding the movements of animals in the wild is a fast-growing area of research, known as *movement ecology*. However, tracking small animals such as flying insects, which cannot easily carry an electronic tag, remains challenging as existing field methods are costly either in terms of equipment or tracking effort (e.g. VHF radio-tracking, scanning harmonic radar). Here we attempted to record the movements of free-flying butterflies from an unmanned aerial vehicle (UAV), maintaining a static position in the sky and recording video vertically downwards. With an appropriate flight height and image filtering algorithm, we recorded 166 flight tracks of *Pieris* butterflies (*P. brassicae* and *P. rapae*), with a median tracking length of 40 m (median flight duration 13 s), and a high temporal resolution of 30 positions per second. Average flight direction varied significantly over the course of the flying season, from a northward azimuth in June and early July, to a southward azimuth in September, a transgenerational migratory behaviour that had previously been documented by field observations or experiments in flight cages. In addition, UAV imagery unlocks the possibility to measure high-resolution flight movement patterns (e.g. path tortuosity and transverse oscillations), which will possibly help understand perceptual and locomotor mechanisms underlying spatial behaviour. We explore the technical details associated with UAV tracking methodology, and discuss its limitations, in particular the biases associated with a 2D projection of 3D flight movements, and the difficulty to distinguish between visually similar species, such as *P. brassicae* and *P. rapae*.

Keywords: UAV image-based tracking, insect flight, insect migration, movement ecology.

Introduction

Tracking the movements of an animal in the wild provides insights on its ecology, such as
habitat use (e.g. Da Silveira et al 2016), dispersal and migration behaviours (Whitfield et al. 2024,
Rotics et al. 2016). When the movement data are detailed enough, both spatially and
temporally, it is also possible to extract biological information on the animal's locomotor
biomechanics (Sherub et al. 2016, Hedrick et al. 2018, Ruaux et al. 2023), its spatial search
strategies (Shepard et al. 2011, Hernandez-Pilego et al. 2017, de Margerie et al. 2018), and more
broadly the perceptual and cognitive processes involved in movement (Kashetsky et al. 2021).
All of these biological and ecological aspects of animal movement have been advantageously
integrated into a synthetic "movement ecology" framework (Nathan et al. 2008, Abrahms et al.
2017, Joo et al. 2022).

However, not all animal species are easy to track. Large-enough animals (approx. > 100 g)
can carry GPS receivers (Cagnacci et al 2010), which are now often enhanced with additional
sensors (accelerometers, barometers, cameras) to better infer the animal's behaviour along its
route (Joo et al. 2022). To track the movements of smaller species such as flying insects,
tracking tags need to be much lighter, and alternative technologies to GPS are employed.
Beacons emitting simple VHF radio beeps have been used successfully for decades to track the
movements of flying and non-flying insects (Kissling et al. 2014). Yet, VHF radio-tracking
involves following the animal with one or multiple antennae (or deploying a fixed antennae
array; e.g. Knight et al. 2019), and the spatio-temporal resolution is inferior to GPS tracking.
Even lighter, passive tags (of a few mg, without any battery) can be used to track the movements
of flying insects (Ovaskainen et al. 2008, Lihoreau et al. 2012, Maggiora et al. 2019). The
downside of these passive tags is that a scanning harmonic radar (SHR) - a heavy and expensive
device - has to be deployed in the field. The tracking range of SHR is near 1 km, and the temporal
resolution of the data is one position every 3 s (Ovaskainen et al. 2008). Other shorter-range,
passive-tag tracking systems also exist (portable harmonic radar, RFID tags; see Kissling et al.
2014, Batsleer et al. 2020, Rhodes et al. 2022 for reviews).

For all animal-borne tracking systems, whether GPS receivers or other types of tags, the
impact of the tag on the animal's movements is a matter of concern. The carried mass, but also
the drag, the position, or the method of attachment of the tag must be carefully considered.
Impact studies are often necessary, both from an ethics point of view and in terms of the
reliability of the collected movement data (Wilson & McMahon 2006, Batsleer et al. 2020).

There are other routes to track flying insects, without having to place tags on animals. First,
the classic mark-recapture method allows to demonstrate the movement of an individual from
one point to another, and was thus used for studying insect migrations (e.g. butterflies;
Chowdhury et al. 2021, 2022). Trapping techniques along migratory routes, sometimes
combined with isotope analyses to determine the geographical origin of individuals, can also
provide information on migrations (e.g. dragonflies; Knoblauch et al. 2021, Oelman et al. 2023).
At a finer, local scale, if the insect flies slowly enough, in an open environment where it remains
visible, it can be followed by foot and its passed positions can be drawn on a map (Brussard &
Ehrlich 1970) or recorded by a GPS receiver carried by the observer (Delattre et al. 2013,
Fernandez et al. 2016). Tracking butterflies crossing a body of water from a small boat is another
possible technique (Srygley & Oliveira 2001). Also, when flying insects move in swarms of many

individuals, radar can be used to detect and measure the direction of these flights (e.g.
Stefanescu et al. 2013), but in this case individual movement is not directly accessible.

To complement these "tagless" tracking approaches, sometimes called "passive sensing"
(Rhodes et al. 2022), we wondered whether flying insects could be tracked in video images
filmed from a UAV (Unmanned Air Vehicle, or drone) positioned above them in the sky.

Image-based tracking from ground-based cameras is a known technique to reconstruct the
2D and 3D trajectories of animals, and in particular flying insects, with a high sampling
frequency (> 1 Hz). It can be used in laboratory settings (e.g. Lihoreau et al. 2016), semi-natural
outdoor insectaries (Kitamura & Imafuku 2015, Le Roy et al. 2021, Kleckova et al. 2024) or even
natural environments (Stürzl et al. 2016, Jackson et al. 2016). Most often used to address
biological questions related to perception, cognition or locomotion, image-based tracking
techniques, which are less invasive, are also attracting growing interest in ecology (Dell et al.
2014). Hybrid techniques tracking tagged animals in videos are also being developed (Crall et
al. 2015, Walter et al. 2021).

On the other hand, in the last decade, commercial UAVs have greatly improved in terms of
compactness, stability and image resolution, while decreasing in cost, making them valuable
tools for wildlife inventory and conservation (Wang et al. 2019, Charbonneau & Lemaître 2021).
Tracking animal movements using videos recorded from a UAV is a next logical step, and it has
already been achieved for a variety of large species (e.g. reef shark: Rieucau et al. 2018, wild
dog: Haalck et al. 2020, zebras and geladas: Koger et al. 2023). For flying insects, UAV-image-
based tracking has been proposed previously (Ivosevic et al. 2017), but remains to be tested
and validated. Most recently, Vo-Doan et al. (2024) successfully tracked a honey bee from a
special UAV-borne optical system (Fast Lock-On), but this technique requires that the insect
carries a reflective marker.

Here we explore the validity of a UAV as a platform for remote, passive observation of the
movements of untagged insects in flight. The motivation for this exploration is the many
potential benefits of such an approach, namely (1) non-invasiveness, i.e. tracking of animals
moving freely, without tags and associated capture procedures, (2) in an open natural
environment, over distances greater than in an insectary, (3) with a spatio-temporal resolution
superior to VHF or SHR tracking. If this type of fine movement data in natural conditions can be
collected easily and cheaply, it could be very useful for bridging the gap between laboratory
studies on perceptual, cognitive and locomotor mechanisms, and large-scale movement
patterns observed in the natural environment (e.g. insect migrations).

To begin this methodological exploration with a relatively simple case, we have focused on
Pieris butterfly species (large white *Pieris brassicae* and small white *Pieris rapae*) because of
their relatively slow flight, good visibility and abundance in the field. We also limited the present
study to the simple situation of a single, static UAV in the sky, filming vertically downwards (Fig.
1A) to reconstruct flight trajectories in only 2 horizontal dimensions.

Material and Methods

117   This section describes the general methods we used to film butterflies in the field,
118 reconstruct their flight trajectories and describe their movements. We later explored and
119 validated these data using diverse specific methods and statistics, which are detailed at the
120 start of each results sub-section, for ease of reading.

Study site

We recorded butterfly flights from June to September 2021, in an agricultural area near the city of Rennes, France (coordinates 48.105444, -1.560029). The local landscape is covered with cultivated fields, few tree patches, small roads and farms, with the closest urbanized area situated at 1.7 km. We filmed butterflies flying above 2 fields situated on each side of a narrow road (Fig.1B). The western field (WF) contained an organic mixed crop (mainly wheat, fava and lucerne) which was harvested in August, whereas the eastern field (EF) contained forage grass, which was cut in June and August. We chose to record butterflies above two different fields (filmed in alternance) to test our image filtering method above various backgrounds, and also control whether the flight trajectories could be influenced by ground vegetation.

On the central road were the UAV takeoff/landing area, the UAV pilot (EDM), the technical assistant (KM), and an ultrasonic anemometer (Gill Maximet 501) recording wind speed and the direction from which the wind originates every second, at 2 m height. We painted two permanent red crosses 20 m apart along the central road (Fig.1D, E), as a reference line segment for positioning the UAV, and later for scaling the video frames. The ground slope in the recorded area was less than 2°.

Figure 1 – UAV field of view (FOV). **(A)** Video recording geometry, showing the influence of UAV flight height (*H*) and camera angular horizontal FOV (*hFOV*) on ground horizontal FOV (*ghFOV*) and ground pixel pitch (*gPP*). **(B)** Aerial view of the study site, showing FOV over western (WF) and eastern field (EF). *Source for aerial photography: geoportail.gouv.fr* **(C)** Variation of *ghFOV* and *gPP*, as functions of *H*. These linear relationships depend on camera angular *hFOV* and recorded image width *IW* (see equations 1-2). **(D-E)** Examples of video frames recorded on July 8[th] 2021, over WF and EF, respectively. White arrows identify 2 reference crosses painted 20 m apart on the road, and used for image scaling.

UAV video recording

We used a Mavic Air 2 UAV (DJI, Nanshan, Shenzhen, China), which is a small commercial
quadricopter (takeoff weight 570 g, retail price ~1000 € in 2021). This UAV has a CMOS sensor
(6.4 × 4.8 mm) which can record 3840 × 2160 pixel videos (i.e. "4K" images, with 16:9 aspect
ratio). According to the UAV manual, the camera lens has an f/2.8 aperture and an 84° field of
view. As most recent UAVs can record with various aspect ratios and resolution levels (which
may involve sensor cropping, i.e. digital zoom), we preferred to measure FOV in the lab, by
placing the UAV camera at a known distance from a wall, and measuring the horizontal distance
along the wall that is effectively included in the UAV camera image. The "horizontal" FOV (*hFOV*,
i.e. along image width), was measured at 68.3° in the default "4K wide" recording mode, that
we used throughout the present study. This angular *hFOV* value was used to choose a flight
height. We computed the horizontal field of view on the ground (*ghFOV*, in meters) when the
UAV camera aims vertically downwards:

(1) $ghFOV = 2H \tan\left(\frac{hFOV}{2}\right)$

where *H* is the UAV height (m), and *hFOV* is the horizontal angular FOV (°).
161
162   Proportional to *ghFOV* is the corresponding pixel pitch on the ground, i.e. the distance on
163   the ground covered by a single pixel side (*gPP*, m):

164   (2) $gPP = ghFOV / IW$

165   where *IW* is the video image width, in pixels.
166
167   The total recorded ground area (*gaFOV*, m²), can also be of interest:

168   (3) $gaFOV = \frac{IH}{IW} ghFOV^2$

169   where *IH* is the video image height, in pixels.
170
Based on these relations, we chose a 45 m UAV flight height, which covers a *ghFOV* of 61 m
(*gaFOV* = 2096 m²), and corresponds to a *gPP* of 16 mm (Fig. 1C). This pixel pitch value was
voluntarily chosen at a fraction of the body size of the *Pieris* species we wanted to track, which
have a forewing length around 30 mm.
We performed preliminary tests in the field that confirmed that at *H* = 45 m, *Pieris* butterflies
flying near the ground were projected in recorded images as pixel "blobs" with an area around
10 pixels, which is large enough to be reliably tracked from frame to frame (see Fig. 2A, C).
Higher camera height would allow larger FOV area on the ground, and hence longer tracking
durations, but automatic tracking of smaller blobs would become less reliable.
Note that at *H* = 45m, the butterflies are not visible to the UAV pilot through the live video
feedback on the UAV controller screen. The video feedback has lower resolution (*IH* = 720 or
1080 pixels) than the recorded video, and the controler screen (Ipad Mini 5, Apple, Cupertino,
USA) also has limited magnification and contrast in outdoor conditions. Butterflies in UAV

footage were only detectable *a posteriori*, when playing recorded videos at full resolution on a
computer screen in the lab.

All videos were recorded at "30 fps" (29.97 video frames per second), which in 4K resolution
produced video data at a rate of 13 MB/s. The UAV firmware automatically cut video files lasting
more than 5 min in multiple 4 GB bouts, which can complicate the post-processing workflow.
For this reason, we decided to keep each video duration below 5 min. Recording at 60 fps was
another possible option, but with a 15 MB/s data rate, this implied less data collected per video
frame. We sticked to 30 fps, aiming to obtain the best possible image quality per video frame.

Constant, manual exposure was tested initially, but proved unpractical as the ground
luminance could vary by several exposure values (Ev) when cloud shadows crossed the FOV.
Hence, we used auto-exposure with an exposure compensation of -0.7 to -1.3 Ev, as it improved
the contrast between the white butterflies and the background.

Time distribution of UAV flights

In order to distribute observations accross the *Pieris* flight season (which in France can span
from April to October; Lafranchis et al. 2015), we chose to collect videos once a week, over the
months of June, July and September 2021. Each week, we chose a day that was favorable for
butterfly flight, i.e. with a weather forecast as warm, sunny and not very windy as possible. We
went on site in the afternoon (14:00-16:30). After takeoff, the UAV was positioned at H = 45 m
above the EF. The UAV camera was tilted to a vertically-downward position, and the pilot used
the video feedback to place the scale points on the road to the left margin of the FOV (see Fig.
1E). The pilot started the video recording, and the UAV then relied on its own sensors (GPS,
altimeter, etc.) to maintain its position without any pilot input, for about 4 min 30 s. Then the
video recording was stopped, the UAV was translated to the WF, height and scale alignment
(now on the right margin of the FOV, see Fig. 1D) were checked, and another 4 min 30 s video
was recorded. Repeating these steps, we could record a second EF video and a second WF video
before the UAV battery dropped to 20-30% of capacity, inciting landing the UAV and swapping
its battery. With this sequence, each UAV battery (rated at 11.6 V, 3500 mAh, 40.4 Wh) allowed
to record about 18 min of video. As we used 3 fully charged battery per field session, we were
able to collect about 1 hour of video per field session. Weather allowing, 3 field sessions could
take place in June (1st, 9th, 15th), 4 field sessions in July (1st, 8th, 15th, 22nd) and 4 in September
(6th, 16th, 22nd, 29th), for a total video duration of 10.7 hours (48.8% over WF, 51.2% over EF).

Video processing

In order to automate the tracking of butterflies in videos, pixels corresponding to the
butterfly should be easily distinguished from the background. In the present case, *Pieris*
butterflies appear in the raw video as blobs of bright pixels, but the background formed by the
vegetation also has multiple bright areas (Fig. 2A), which rules out a simple detection of the
butterfly by thresholding the raw video luminance levels. In addition, as the UAV is not perfectly
static, and the wind can cause vegetation on the ground to move slowly, the background is
moving, which does not favour background subtraction approaches (Piccardi 2004). We found
a solution to this issue by designing a custom filter that selects the pixels that blink in the video:
when a butterfly passes over an area, the pixels in that area become brighter for only 1 frame,
and then revert to the background luminance. To apply this filter, we first transformed the RGB

video frames into greyscale. Each pixel then has a single luminance value (v) in the range [0,
255]. Then the "blink" filter script performs the following calculations:
228
Pixel value variations from current frame (t) to next (t+1) and previous (t-1) video frames:

(4) $\Delta v_n = v_{t+1} - v_t$
(5) $\Delta v_p = v_{t-1} - v_t$

If both variations have the same sign, there was a luminance peak (positive or negative), and
a blink value (b) is computed. On the contrary, if variations have different signs, there was no
blinking, only monotonous pixel value variation.

(6)   if $sgn(\Delta v_n) = sgn(\Delta v_p)$ $\rightarrow$ $b_t = -sgn(\Delta v_n) * \min(|\Delta v_n|, |\Delta v_p|)$
(7)   else $\rightarrow$ $b_t = 0$

Note that b can be positive (bright blob passing over darker background) or negative (dark
blob passing over brighter background).
239
Interestingly, when going through a series of successive video frames, keeping memory of
extreme b values for each pixel (notated B) can be used to reveal flight trajectories as a series
of blobs, in a synthetic, diachronous image.

(8)   if $|b_{t+1}| > |b_t|$ $\rightarrow$ $B = b_{t+1}$
(9)   else $\rightarrow$ $B = b_t$

For easy display of b frames (and B synthetic images), pixel blink values are rescaled from [-
255, 255] to [0, 255].

(10) $b = (b + 255)/2$

As a result, background appears as medium grey (b = 128), with blinking pixels as lighter or
darker gray blobs. Fig. 2 shows example results of the blink filter.
We used the blink filter in two processing steps. For the initial exploration of our videos, we
cut each video into 20 s bouts, and generated a single B image summarising each bout. This
enabled us to quickly detect which video bouts contained butterfly tracks. On this basis, the
FOV entry and exit times of each butterfly were precisely noted by playing the original videos
on a large screen. Based on these time limits, we then generated for each butterfly track a new
greyscale, uncompressed video file, containing the series of b frames. This filtered greyscale
video was then used to automatically track the blobs.
We used DLTdv8 (Hedrick 2008) to extract blob centroid coordinates (in pixels) in successive
video frames. The blob was manually digitized (i.e. mouse clicked) in the first few frames, and
then DLTdv8 uses a Kalman filter and 2D cross-correlation to find the blob in following frames
(without thresholding). As blob size and contrast can vary, and filtered b frames still contain
some noise in the background, the automatic tracking process needed human supervision and
frequent manual corrections, but with a convenient user interface to navigate through video
frames, DLTdv8 offered vast time savings compared to a fully manual digitization.

Figure 2 – Video frame filtering. **(A)** Magnified view of an original video frame, containing a *Pieris brassicae* image (as identified after butterfly capture). **(B)** Same view after applying the blink filter (i.e. *b* frame). **(C)** A hypothetical 30 mm forewing length *Pieris*, with wings fully stretched, projected onto a pixel grid with a 16 mm *gPP*. If all partially covered pixels appear brighter, the pixel blob area would be around 12 pixels. **(D)** Merging successive *b* frames reveals *P. brassicae* flight trace in a synthetic, diachronous image (i.e. *B* image). **(E)** Example trace of *Pieris rapae*. **(F)** Example trace of a dark-coloured species, either *Vanessa atalanta* or *Aglais io* (contrast ×2.0). **(G)** Example trace of *Melanargia galathea.* **(H)** Example trace where the shadow of a *Pieris* butterfly is also visible (contrast ×1.5).

For scaling the butterfly track to real-world coordinates, we measured pixel coordinates of the two painted reference crosses in one video frame (at mid-duration of the track), and fitted a geometrical transformation (combining rotation, scaling and translation; *fitgeotrans* function in Matlab) that resulted in (0, 0) and (0, -20) coordinates in real-world meters. This same transformation was then applied to the whole butterfly coordinate series, transforming coordinates in pixels to meters. We had measured with a compass in the field that the central road had a 6°E azimuth. We checked and refined this value in a GIS software (https://www.geoportail.gouv.fr), and thus applied a 5.83° CW rotation to all butterfly coordinates, so that in all graphical representations, y axis has a 0°, northward azimuth.

Track selection

When we explored the *B* images synthesizing our videos, we mainly found flight traces of *Pieris* butterflies (large white *P. brassicae*, small white *P. rapae*), appearing as clearly visible white dotted traces (fig. 2D, E). These two species were also the most easily observed from the ground during the field sessions. We also found a few flight traces of other species, that we had observed in the field, such as the marbled white (*Melanargia galathea*), the red admiral (*Vanessa atalanta*) or the peacock (*Aglais io*). However, as these traces were less frequent, and usually barely visible and discontinuous (Fig. 2F, G), we did not analyse them further.

For some *Pieris* traces, the shadow of the butterfly formed a second, dark trace (Fig. 2H), which may potentially be used to determine the height of the butterfly relative to the ground (knowing the associated sun elevation angle). These shadows were only visible on almost bare ground (September videos, after crop harvest), and thus concerned a small number of *Pieris* traces. We have therefore not used this additional source of 3D data in the present study.

Among the *Pieris* tracks, we applied the following exclusion criteria:
- A 10 m wide strip, containing the road and the adjacent flowered ditches, was excluded from the analysis, so that each field (WF and EF) included a homogeneous area in terms of ground vegetation.
- Many very brief tracks (< 5 s), often crossing just one corner of the FOV, were considered less informative and excluded from analysis.
- As we were mainly interested in migratory flights, defined as "directional flights in which individuals appear undistracted by appetitive cues" (Chowdhury et al. 2021), we chose to focus on butterflies in continuous movement: Tracks marking one or more stops while crossing the FOV were excluded (a stop being defined as remaining for at least 1 s within a radius of 0.1 m). These tracks with stops were in the minority (~ 1 out of 6 tracks).
- Tracks in which the butterfly interacted with another individual (e.g. flight inflection towards another butterfly, chases) were also excluded (~ 1 out of 13 tracks).

In the end, our sample comprised $N = 166$ *Pieris* flight tracks. Despite our constant sampling efforts, these were not evenly distributed through the season: we recorded 12 tracks in June, 125 in July and 29 in September. Taking all these tracks together, 70524 butterfly positions were recorded, representing a total of 39 min of flight time, and a flight distance of 7.4 km.

When the tracks contained missing positions (caused by the absence of blob in some *b*
frames), the (x, y) coordinates were linearly interpolated. These interpolated positions
represented 2.2 % of the dataset (1544 positions). The interpolated positions were rarely
contiguous, and the longest interpolated segment represented 7 successive positions (i.e. 0.23
s).

Track descriptive variables

For this pilot study, we computed a small set of basic descriptive variables for each 2D track.
- **Track duration** is the time the butterfly remained within the FOV (road zone
excluded).
- The change in 2D position from a frame to the next frame is named a step vector. **Track
length** is computed as the sum of step vector norms.
- Step speed is equal to step vector norm divided by the elapsed time (i.e. 1/29.97 s).
Average speed is the arithmetic mean of the series of step speed values. It indicates
how fast, on average, the butterfly flew along its 2D track. Average speed is also equal
to track length divided by track duration. Note that average speed is a ground speed,
not an air speed.
- A "beeline" vector is defined as the vector from the first to the last recorded position
of the butterfly. The beeline vector is equal to the vectorial sum of step vectors, and
hence represents the butterfly's resultant, directed movement across the FOV. **Beeline
azimuth** was the direction of the beeline vector, a circular variable in the interval [0°,
360°[, 0° corresponding to a northward azimuth.
- Track **straightness** is computed as the ratio of the beeline vector norm to the track
length. Straightness value is in the interval [0, 1]: 0 indicates that the butterfly
performed a loop (i.e. had the same entry and exit positions), while 1 indicates a
perfectly straight flight path. Straightness is inversely related to path tortuosity.
- **Beeline speed** is the norm of the beeline vector divided by track duration. It reflects
how fast the butterfly, on average, progressed in its directed movement. Beeline speed
is also equal to average speed multiplied by straightness. Note that beeline speed is
also a ground speed, not an air speed.
- **Wind speed** and **direction** for each track were computed from the vectorial sum of the
n wind vectors recorded during track duration, divided by n.

Processing and statistics software

Pixel blob tracking in videos was performed using DLTdv8 (Hedrick 2008;
https://biomech.web.unc.edu/dltdv/). Other analyses, from video processing to statistics, were
performed using Matlab R2018b (The MathWorks, Natick, MA, USA). We used the CircStat2012a
toolbox for circular statistics (Berens 2009). For a small number of captured butterflies,
forewing length measurement from field photographs were performed with ImageJ V1.54g
(http://imagej.org).

Results

Track general description

Fig. 3 shows the reconstructed 2D tracks, and associated variable distributions. We collected 66 *Pieris* flight tracks over WF, and 100 tracks over EF. Tracks had a median duration of 12.8 s (range 5.1 to 56.5 s, Fig. 3B), for a median track length of 40.0 m (13.4 to 134.9 m, Fig. 3C). The median value for average speed was 3.3 m.s^{-1} (1.9 to 9.2 m.s^{-1}, Fig. 3D), and 2.9 m.s^{-1} for beeline speed (0.2 to 9.1 m.s^{-1}, Fig. 3F). Straightness distribution was strongly skewed towards straight tracks, with a median value of 0.93 (range 0.10 to 0.99, Fig. 3E). High straightness values imply similar values for average speed and beeline speed in most tracks.

Figure 3 – *Pieris* flight tracks description. (A) Reconstructed 2D tracks over WF and EF. Black dots indicate last position of each track. **(B-F)** Distributions of descriptive variables: (B) track duration, (C) track length, (D) average speed, (E) track straightness and (F) beeline speed.

We used two-sample Kolmogorov-Smirnov (KS) tests to assess distribution differences between WF and EF tracks, and found no significant difference for track duration ($D_{(66,100)} = 0.10$, $p = 0.78$), track length ($D_{(66,100)} = 0.09$, $p = 0.86$), average speed ($D_{(66,100)} = 0.09$, $p = 0.86$), or beeline

speed ($D_{(66,100)} = 0.18$, $p = 0.15$). However, there was a significant difference between straightness
distribution over WF and EF ($D_{(66,100)} = 0.26$, $p = 0.009$), with straightness for EF tracks being even
more skewed towards 1 (median 0.94) than WF tracks (median 0.90; see Fig. 3E).

Beyond average speed and straightness, the 30 fps butterfly flight tracks extracted from UAV
videos also contained fine-scale instantaneous information about flight speed and azimuth
variation along the tracks. For example, Fig. 4 shows a track segment that depicts interesting
movement patterns, in the form of meter-scale, sub-second transverse oscillations along the
flight path.

Figure 4 – Benefit of 30 Hz positional data. The upper line is a 12 s flight track segment,
reconstructed with the present method. The lower track is a copy (with a -3 m y offset for
visualization) showing all recorded 2D positions (361 in total). The 5 circles represent the
positions that could have been recorded with a SHR tracking system, that has a 90-fold
lower sampling frequency (1 position every 3s.). Note that on the other hand, SHR
benefits from a larger FOV, and hence longer tracking durations (see discussion).

Controlling UAV stability

During video recording, the UAV was not perfectly static, and whether this could have an
effect on the reconstructed flight tracks was an important issue. For a small random subset of
tracking videos ($N = 10$), we digitized the two reference points painted on the road, describing
a 20 m reference segment (RS), not only at mid-duration, but on every video frame, which
allowed to monitor how the RS was transformed throughout the duration of tracking, due to
UAV movements. We assumed that a combination of rotation, scaling and translation could
affect the RS projection. Using the *fitgeotrans* function in Matlab, we obtained the geometric
transformation matrix from the first frame's RS to each following frame, which allowed to
monitor rotation, scaling and translation movement components separately.

Fig. 5 shows that the RS projected image was indeed affected by a combination of geometric
transformations through time. Over the investigated tracking durations (13 to 40 s), rotation of
the RS could attain 0.26° (Fig. 5A), and was on average 0.07° (root mean square, RMS). Scaling
variation could attain ± 1.13 % (0.36 % RMS; Fig. 5C). Horizontal (x) or vertical (y) translation of
the RS image (Fig. 5B) could reach ± 11.2 pixels (3.6 pixels RMS), which represents less than 0.3
% of image width, or 0.18 m when projected on the ground. The amounts of transformation
usually did not grow monotonically through time, reflecting that in the absence of pilot input,

the UAV does not simply drift away from its initial position, but uses servoing based on its
onboard sensors to try and maintain position and azimuth. Digitizing the RS only once at track
mid-duration - i.e. assuming that the UAV is fully static during each track - resulted in a maximal
0.28 m error (0.08 m RMS) on the butterfly reconstructed 2D position (Fig. 5D). For the present
work, we considered these levels of error to be acceptable, which is why we used the single RS
digitizing method for all the remaining tracks.

413   **Figure 5 – UAV movements.** Reference segment transformation through time for 10
414   tracking videos, decomposed into **(A)** rotation, **(B)** translation and **(C)** scaling
415   components. **(D)** Resulting 2D error on the butterfly 2D position, when only a single
416   digitization of the RS is performed at mid-duration.

Controlling butterfly flight height

With the present recording geometry, the flight tracks of all butterflies passing through the
pyramidal FOV (Fig. 1A), regardless of their flight height, are projected on a single sensor plane.
We reconstructed these tracks in 2D by assuming that the butterflies moved in the same plane
as our reference 20 m segment, i.e. flew at ground level. How this simplification departed from
the 3D reality needed investigation. We were especially interested in the possibility that some
butterflies might have crossed the FOV at a significant height (10-40 m), which would result in
vastly overestimated flight speeds in our ground-projected reconstructions.
During field work, we visually monitored the fields under the UAV, and noted each time we
saw a *Pieris* butterfly passing at a low height (defined as below the observer visual horizon, i.e.
less than ~2 m above the ground). By comparing our field notes with the timestamps of the
reconstructed tracks, N = 45 out of 166 tracks (i.e. 27 %) corresponded to visually-detected
butterflies flying at low height (LH group). The remaining tracks were qualified as "unknown
height" (UH group, N = 121), and may correspond either to high-flying butterflies (visually
undetected because of low contrast against the sky), or to low-flying butterflies that remained

undetected (because the observer's attention was regularly directed at the UAV rather than at
the ground).

We compared the beeline speeds of the low-height (LH) and unknown-height (UH) groups.
Both groups had very similar speed distributions (Fig. 6A), as confirmed by statistical tests : LH
(2.9 ± 1.0 m.s^{-1}, mean $\pm$ SD) and UH tracks (3.1 ± 1.3 m.s^{-1}) did not significantly differ for mean
speed in a t-test ($t_{(164)} = 0.84$, $p = 0.40$), and were not drawn from different distributions according
to a KS test ($D_{(45,121)} = 0.11$, $p = 0.79$).

Figure 6 – Speed distribution and flight height. (A) Ground beeline speed distribution
observed for low height (LH) and unknown height (UH) butterfly tracks. **(B)** Kernel
probability density estimation for LH and UH tracks, and for simulated tracks with
uniform height distribution (see text).

In order to visualize what speed distribution would be obtained for butterflies flying far from
the ground, we simulated 10^5 straight horizontal tracks, with flight speeds sampled from a
normal distribution copied form the LH group (2.9 ± 1.0 m.s^{-1}), but crossing the FOV at heights
uniformly distributed between 0 and 45 m. Simulated tracks crossing the FOV, and with a track
duration of at least 5 s (to simulate comparable track selection), had a ground-projected speed
distribution that was flatter and shifted towards high (i.e. overestimated) speeds when
compared to both LH and UH tracks (see Fig. 6B). This further suggests that the UH group in our
track sample mainly comprises low-flying butterflies that went visually undetected in the field.
Still, we cannot exclude that a small number of tracks in the UH group (on the right tail of speed
distribution, e.g. with reconstructed speed > 6 m/s) might correspond to high-flying butterflies.

Exploring blob area as a specific signature

We explored the possibility to discriminate *P. brassicae* from *P. rapae* tracks, based on pixel
blob area in the videos. During field work we captured a small number (*N* = 8) of *Pieris*
individuals that had just passed through the UAV's FOV. Each individual, captured with a
butterfly net, was briefly placed in a thin transparent box with a grid-patterned back,

photographed with its wings stretched for later identification and size measurement, and then
immediately released. Back in the lab, we measured each individual's forewing length using
ImageJ. The UAV video recordings corresponding to these individuals were filtered and
digitized with the same methods as previously described, but were later re-analysed to measure
the blob size in each filtered video frame. As a simple approach, the blob was defined as
connected pixels with grey level $b > 138$, i.e. departing by more than 10 grey level from the mean
background grey of 128. Sometimes the simple thresholding approach detected no blob, but
this method still allowed to obtain many (269 to 654) blob area values per individual, that could
be compared to the animal's real size. Note that 4 out of the 8 tracks used for the present blob
area analysis were not part of the final flight track sample ($N = 166$), because these captured
individuals had flown near the road, or performed stops along their flight (see Track selection
section).

Five captured individuals were identified as *P. rapae* (2 females + 3 males), with forewing
length ranging from 22.8 to 28.0 mm. The 3 other captured individuals were *P. brassicae* (2
females + 1 male), with longer forewing, ranging from 31.2 to 36.7 mm. These values were in line
with the literature, with the largest *P. rapae* individuals being close in size to the smallest *P.*
brassicae specimens (Cook et al. 2022)

Figure 7 - Forewing length vs. pixel blob area distribution for 8 tracked and captured
Pieris butterflies. Blue: *P. brassicae* ($N = 3$); Green: *P. rapae* ($N = 5$). Distributions are
displayed as Kernel probability density estimates. Circles indicate the median blob area
value for each butterfly.

Fig. 7 shows the blob area distributions observed for all 8 captured individuals. When
considering only the median blob area for each individual, it was positively correlated to the
forewing length of the animal (Spearman rank correlation, $r_{(6)} = 0.73$, $p = 0.047$). However, the
relationship was not monotonically increasing ($r < 1$), and there was extensive overlap between

blob area distributions. In other words, a smaller butterfly could often project as a larger pixel
blob than a larger butterfly, depending on the compared video frames. As a result, we
considered unreliable to use recorded pixel blob areas as a direct mean to discriminate *P. rapae*
and *P. brassicae* flight tracks in the present work.

Effect of advancing season on flight azimuth.

By observing the tracks at different times in the flight period, it appeared visually that the
flight azimuths have varied over the season (Fig. 8).

Figure 8 – *Pieris* flight tracks broken down by flight period (vertically) and field
(horizontally). Black dots indicate last position of each track (*N* = 166). 30 m diameter
circles represent the area considered for azimuth statistical comparisons (*N* = 119).

To further quantify seasonal variations in azimuth distributions, we computed the beeline
azimuth of tracks crossing a 30 m diameter disc located at the centre of the camera's FOV (Fig.
8). Indeed, the rectangular shape of the camera FOV is less likely to record trajectories parallel
to the longer side of the FOV (see Fig. S1), and this bias can be corrected by considering only an
area enclosed by a circle inside the FOV. When a track did not cross this central disc, it was
therefore removed from the sample for circular statistics. When a tortuous track crossed this
disc more than once, only the longest segment inside the disc was considered. This restriction
of the FOV to a central disc had the effect of reducing our sample from $N = 166$ to $N = 119$ (for
this section only).

We tested the effects of period (June to September) and field (WF vs. EF) on beeline azimuth,
using a two-factor ANOVA for circular data (Harrison & Kanji 1988, in Berens 2009). As we
observed many more trajectories in July, we subdivided the month of July: *early July* (field
sessions on July 1st and 8th) and *late July* (July 15th and 21st). We then tested the uniformity of
the azimuth distributions for each period, using Rayleigh tests (Fisher 1995, in Berens 2009).

Figure 9 – Circular distributions of *Pieris* beeline azimuth, broken down by period
(vertically), and field (horizontally). Arrows represent resultant vectors. The right column
shows azimuth distributions for both fields pooled (WF + EF), with red arrows
representing significant resultant vectors according to a Rayleigh test.

The ANOVA for circular data detected a significant effect of period on azimuth ($X^2_{(6)}$ = 43, p =
1.2 × 10^{-7}), but no effect of field side ($X^2_{(2)}$ = 0.51, p = 0.77), and no interaction between the two
factors ($X^2_{(3)}$ = 5.5, p = 0.14). The distribution of azimuths during the 4 periods is shown in Fig. 9.
In June, the butterflies flew most often to the north-east (Rayleigh test, N = 7, R = 0.65, p = 0.04;
mean azimuth 24°). In early July, they flew most often to the north-west (N = 51, R = 0.56, p = 4
× 10^{-8}; mean azimuth 317°). In late July, tracks in all directions were observed, without any
dominant azimuth, so that the azimuth distribution was not significantly different from a
homogeneous distribution (N = 42, R = 0.15, p = 0.41). Finally, in September, tracks were most
often oriented to the south (N = 19, R = 0.59, p = 0.001; mean azimuth 173°).

To verify if the above results were influenced by wind direction, we first tested whether
butterfly beeline azimuth was correlated with wind direction, using a circular-circular
correlation test (Jammalamadaka & Sengupta 2001, in Berens 2009). The test returned no
significant correlation (c = - 0.01, p = 0.95; Fig. 10A). We also assessed whether butterfly
preferentially flew downwind, crosswind or upwind, by computing the angular difference
between beeline azimuth and wind direction, and testing whether this "angle to wind" variable
departed from a uniform distribution. Results suggested that butterflies did not preferentially
fly at a specific angle to wind (Rayleigh test, N = 119, R = 0.15, p = 0.06; Fig. 10B), but the test was
close to statistical significance, despite a small resultant vector (i.e. small effect size).

As a supplementary verification, we focused on tracks recorded during stronger winds (> 2
$m.s^{-1}$, N = 64), as these butterflies were expected to be the most affected by a possible wind
influence. Both circular-circular correlation (c = - 0.04, p = 0.75) and Rayleigh test on angle to
wind (N = 64, R = 0.08, p = 0.67) returned non-significant results (Fig. 10C, D). This comforted the
conclusion that wind direction did not significantly bias butterfly flight azimuth in our data.

Figure 10 – Wind and *Pieris* flight azimuth. (A) Beeline azimuth vs. wind direction for N
= 119 tracks. **(B)** Angle to wind for N = 119 tracks. 0° corresponds to upwind flight, 180° to
downwind flight. **(C-D)** Same graphics for the N = 64 tracks where wind speed exceeded
2 $m.s^{-1}$.

Discussion

Butterfly tracking method validation, strengths and limits

We show that UAV-image-based tracking can be used to reconstruct free-flying butterfly
paths. With light, affordable and easily-deployed gear in the field, we were able to track
numerous wild, untagged *Pieris* butterflies, over distances averaging 40 m. Given an average
flight speed of about 3 m.s^{-1}, this translates to tracking durations most often near 10-20 s (Fig.
3B-D), depending on each butterfly's flight speed and straightness. This scale of recorded
movement is larger than what can be achieved in most insectaries, but remains modest
compared to some other open field methods (e.g. approximately 500 m for SHR; Ovaskainen et
al. 2008, Lihoreau et al. 2012), and is directly constrained by the UAV camera FOV.

In the present study, the FOV was voluntarily limited to 61 × 34 m (from a 45 m UAV height),
in order to maintain a butterfly blob area around 10 pixels, which was useful for reliable blob
tracking throughout video frame series (see Methods section). A first possible way to enlarge
FOV would be to use an UAV with higher camera resolution (e.g. 6K or 8K sensor, with matching
optics quality), which could potentially fly higher and record a wider ground area, while
maintaining a centimetric pixel pitch on the ground. Moreover, a higher signal-to-noise ratio in
video frames (e.g. from larger sensors and/or less compressed video file formats) might allow
to reliably detect smaller blobs throughout frame series, which in turn would allow larger pixel
pitch values, and even higher UAV flight height. A FOV exceeding 100 m in side length is probably
already possible with current high-end commercial UAVs, but with a heavier, less affordable
UAV, and larger amounts of video data to process.

An alternative way to greatly increase recorded movement length would be to try and follow
butterflies with the UAV. This would necessitate (1) that the video feedback to the UAV pilot is
of sufficient magnification for a live view of individual butterfly blobs and (2) a different
approach to video frame filtering (accounting for quickly moving image background), but these
are interesting perspectives for future methodological developments.

On the other hand, the main strength of the movement data we collected is the 30 Hz
temporal resolution, which is 2-3 orders of magnitude higher than non-image-based field
tracking methods applicable to flying insects: 0.33 Hz for SHR (Ovaskainen et al. 2008) and lower
(usually < 0.01 Hz) for automated radio-tracking (Kays et al. 2011). Here, with one location every
~10 cm along the flight path, the reconstructed tracks reveal fine-scale movement patterns (see
Fig. 2, 4, 8), and offer access to flight speed and tortuosity in the wild, with improved accuracy.
Such refined movement data may provide interesting insights on biomechanical and/or
orientation processes at work during butterfly flight. More detailed analyses focused on the
effect of wind on flight speeds, and the oscillation patterns along *Pieris* flight paths are
envisaged, but were beyond the scope of the present methodology presentation.

Another limitation of the present method is the projection of the FOV, a pyramidal 3D air
volume that butterflies can cross at various height, onto a virtual 2D surface at ground level. In
the present dataset, we were able to verify by direct observation that at least 27 % of tracked
butterflies flew at less than 2 m above ground level. For these tracks, the 2D ground projection
implies only a small error on flight speed (i.e. an overestimation of less than 2 / (45 - 2) = 4.7 %).

For other butterflies, for which we were unable to confirm flight height, we showed that their
projected flight speeds are no higher than confirmed "low flyers", and thus conclude that they
probably also crossed the FOV at low height (Fig. 6). This remains indirect evidence, and we
cannot exclude that for a minority of tracks, we might have significantly overestimated flight
speed because the butterfly crossed the FOV at a higher height.
Note that, in the hypothetical situation where butterflies move mainly in horizontal planes,
but at various heights, the reconstruction of flight speeds will be heterogeneously
overestimated, but the angular and temporal variables (e.g. variation of azimuth through time)
remain unaffected by flight height.
Beyond the average flight height, all the vertical (z) movements of the butterflies are lost
when projected onto a (x, y) horizontal plane. For following studies, it is therefore advisable to
observe the 3D flight behaviour of butterflies beforehand, depending on the species under
consideration and the relief of the terrain, and to estimate whether a 2D projection might
overlook important information about the animals' movements in the local lanscape. If the 3D
flight trajectory and/or elevation relative to the ground are really necessary for a study (e.g.
flight biomechanics), one can either try to find a study zone where the butterfly's shadow is also
visible in the image (Fig. 2H), or opt for other natively 3D optical methods, based on multiple
views of the flight volume (Theriault et al. 2014, de Margerie et al. 2015).
608
When the UAV receives no command from the pilot, its flight control algorithm seeks to
maintain its horizontal position, height and azimuth, using sensory-motor regulation loops
based on many on-board sensors (GPS, barometric altimeter, magnetic compass,
accelerometers, gyroscopes, downward vision system). Commercial UAVs have made
spectacular progress on stability in the last decade, and we were able to verify that, at least over
the tracking durations used here (< 1 min), the movements of our UAV were indeed very limited
(Fig. 5). This allowed us to assume that the drone was static in the sky, at the cost of a drift under
30 cm in the reconstructed position of the butterfly, which we found acceptable for the question
posed here (i.e. the measurement of the beeline azimuth over several tens of meters). Still, note
that most UAVs use barometric sensors for regulating flight height, and that atmospheric
pressure might vary significantly across flight times longer than a few minutes.
For studies requiring lower error caused by the drone's position, it is possible to repeat the
measurement of the reference scale on every video frame, or to perform a more refined
calibration of the projected image by continuously tracking multiple points on the ground. If
the drone's movements really need to be more closely controlled, more advanced commercial
drones are available, using positioning technologies such as Differential GPS (DGPS) or Real
Time Kinematic (RTK), which can monitor the UAV position with centimeter accuracy.
626
Our video tests suggest that many more butterfly species are potentially detectable using
the present method, whether they appear light against a dark background or dark against a
light background (Fig. 2). The blink filter proved efficient for erasing background textures, while
also tolerating a fair amount of background movement, caused either by the slow drift of the
UAV, or vegetation being blown over by the wind. Hence, we hope to see following studies
tracking other butterfly species in various landscapes. Unfortunately, depending on the
location and the flight season, it is possible to run into the issue that two (or more) species with

similar sizes and colors get filmed simultaneously, and this is the problem we encountered here with two *Pieris* species. We were not able to discriminate *P. brassicae* from *P. rapae* based on pixel blob area (Fig. 7). The wide, overlapping distributions of blob areas for each individual butterfly is not surprising: Flying butterflies can have any posture, from fully stretched to fully closed wings on different video frames, as a result of (1) flapping wing movement and (2) variable roll and pitch angles of the body along the flight path. Moreover, pixel blob area can also be affected by other factors such as (3) contrast with the local background, (4) "blob-clipping" (i.e. when the distance covered between two consecutive frames is less than the body length, which interferes with the filtering method), and (5) flight height. Note also that the relationship between butterfly size and blob area is expected to follow discrete steps (pixels), especially when the pixel pitch is close to animal body size (Fig. 2C).

Provided that some of these sources of blob area variation can be better controlled, we do not rule out that pixel blob area could help discriminate butterfly species in future studies, but this would need more refined blob classification processes than what we implemented here.

Another option could be to discriminate species based on flight speed or flight behaviour (e.g. tortuosity). A first issue with this approach is that in studies aiming at describing the flight behaviour, it could lead to circular inference. Moreover, Kleckova et al. (2024) reported that different species (*P. rapae* and *P. napi*) can have smaller differences in their flight parameters (measured in an insectary) than spring and summer generations of the same species.

Another route for reducing possible confusion between species could be a greater effort to identify each individual's species (and sex) in the field, either by remote visual/photographic identification, or by systematically capturing butterflies after they crossed the UAV's FOV. This would be limited to low-flying butterflies, and also implies greater human presence and movements in the field, which might affect the butterflies' movement patterns. As well, capturing the butterflies before they cross the FOV, and tracking their movements once released was not retained as a valid option, as released butterflies do not immediately display their normal flight behaviour (Nikoleav 1974, Dudley & Srygley 1994).

Observed flight speed and straightness distributions

Ground-based multi-camera settings can be used to measure butterfly flight speed in 3D, most often inside insectaries (e.g. Kitamura & Imafuku 2015, Le Roy et al. 2021, Kleckova et al. 2024). Unfortunately, many butterflies species tend to fly slower in insectaries than in the wild (Dudley & Srygley 1994), and captivity can also affect other flight parameters (e.g. glide duration: Le Roy et al. 2021). The present data offers a nice opportunity to measure *Pieris* flight speed in undisturbed, natural conditions, although in 2D only. The median average ground speed we report, at 3.3 m.s^{-1}, comes out lower than some earlier measurements on smaller samples of migrating *Pieris* in the field (e.g. 3.6 m.s^{-1} in *P. brassicae*, Nikolaev 1974; 4.4 m.s^{-1} in *P. rapae*, Dunn 2024). More interestingly, across our relatively large sample from multiple field sessions, we observed a wide speed range (from 1.9 to 9.2 m.s^{-1} for track average ground speed, Fig. 3D), and this does not even account for flight speed variations within each track. This again calls for a detailed (upcoming) analysis of instantaneaous ground and air speeds, taking wind into account, that might provide novel insights on the flight behaviour of *Pieris* (e.g. wind drift compensation; Gilbert & Singer 1975, Srygley & Oliveira 2001).

Flight speeds over WF (mixed crop) and EF (grass) did not differ, and most tracks had a very
high straightness value, compatible with a "directional, undistracted" migratory flight
behaviour (Chowdhury et al. 2021). Still, we measured that tracks over WF were not quite as
straight as over EF (Fig. 3E), suggesting that the more diverse vegetation in WF might have
"distracted" butterflies to some extent. Also, we note that some butterflies in our sample
exhibited tortuous rather than directional flight (see Fig. S2 for tracks with straightness < 0.6).
Moreover, note that the speed and straightness distributions we report would probably be
different if we had included butterflies marking stops along their flights. Thus it would be
incorrect to assume that all the *Pieris* butterflies we observed in the field exhibited a migratory
behaviour.
The higher number of butterflies passing over EF (*N* = 100 vs. *N* = 66 for WF) was intriguing.
After a close examination of sample sizes for each field session (see table S1), it appears that EF
and WF butterfly numbers only differed significantly for 1 out of 11 field sessions, thus we do
not conclude that EF consistently attracted more butterflies than WF.

P. brassicae and *P. rapae* migratory behaviours

Migratory behaviour has been documented in about 600 butterfly species (Chowdhury 2021,
2022). In butterflies, migration often occur over several generations, with successive
generations following different flight azimuths to achieve a round-trip annual travel that can
cover thousands of kilometers in some Nymphalidae species (e.g. monarch *Danaus Plexippus*:
Brower 1995; painted lady *Vanessa cardui*: Stefanescu et al. 2013, Suchan et al. 2024).
Within Pieridae, another butterfly family with numerous migratory species (Chowdhury
2021), *P. brassicae* and *P. rapae* migratory behaviours have long been documented from the
observation of mass migrations (Williams et al. 1942, Vepsalainen 1968). These mass migrations
are now rarer due to the use of pesticides (Spieth & Cordes 2012), but see John et al. (2008) and
Dunn (2024) for recent reports of group migrations in *P. rapae*. Early attempts at quantifying the
directions and distance of *Pieris* migratory flights used methods such as visually estimating the
flight azimuth in the field (e.g. Baker 1968), or difficult mark-recapture experiments (Roer 1959,
1961). More recently, these species' movements have been studied by more advanced
experiments. Jones et al (1980) used an egg marking method: they fed *P. rapae* larvae with a
special diet causing the adult females to later lay colored eggs, which allowed to study the
movements of individual females over a 1 km square grid of cabbage plants. They reported that
australian female *P. rapae* have a directed flight, but that the preferred azimuth varied from day
to day (with some northward bias). Gilbert & Raworth (2005) performed complementary
experiments in Australia that showed similar daily directionality in male *P. rapae*. The same
authors report field observations in the Pyrenees mountains in Europe: there, a portion of the
P. rapae population migrated northward in spring and southward in autumn, in line with earlier
observations in England (Baker, 1968). For *P. brassicae*, Spieth & Cordes (2012) collected eggs
from several Western Europe regions, and later measured the spontaneous flight azimuth of
adult female individuals, in a 2 m octogonal flight cage. They showed that the preferred flight
azimuth depended on the season and the geographic origin. In regions where *P. brassicae* has
a cycle of 2 generations per year, the first generation usually followed a northward azimuth
(modulated by the precise geographic origin), whereas the last generation flew southward. In
regions where *P. brassicae* has 3 generations per year, both the first and second generations

flew northward, and only the last generation had a reverse, southward flight azimuth. Using a similar flight cage, Larranaga et al. (2013) confirmed a mean northward azimuth in both females and males *P. brassicae* of the first generation.

Compared with these pre-existing data on migration in *P. brassicae* and *P. rapae*, flight tracks recorded from a UAV allow more accurate measurement of flight azimuth, without the need to capture or mark the butterflies (which affects spontaneous flight behaviour: Nikoleav 1974, Dudley & Srygley 1994), at an intermediate scale between a flight cage (or insectary) and mark-recapture experiments. For a relatively large (N = 119), mixed sample of individuals from both species, we measured flights that were mainly directed northwards in the early season (June, early July), and a return southward azimuth in the late season (September). Our azimuth data in natural conditions (Fig. 8-9) are therefore congruent with the existing literature for both *P. brassicae* (Spieth & Cordes 2012) and *P. rapae* (Gilbert & Raworth 2005). Note that the absence of a dominant azimuth for late July might be the result of *P. brassicae* already shifting to southward flight, with *P. rapae* still flying predominantly northward at this period (R. Baker, personal communication).

As a plus, our high sampling frequency trajectories appear to contain previously unavailable fine-scale information on *Pieris* flight speeds, tortuosity of the flight path, and patterns of rapid azimuth variation (transverse oscillations). This refined data may be studied in greater detail and perhaps reveal information on locomotor behaviour and the perceptual mechanisms underlying *Pieris* spatial behaviour, whether migratory or not. A major limitation of our approach at this stage remains the non-distinction between *P. rapae* and *P. brassicae*, because of their similar sizes and colors, and their concurrent flight in our geographical area.

Conclusion

Our results reveal that video tracking of butterflies from a UAV is possible, and capable of providing movement data in fully natural conditions, at an unprecedented spatio-temporal resolution, and at a modest cost. This fine-scale data could prove precious for understanding the spatial behaviour of many butterfly species in the wild, which has so far been only partially documented and understood (Spieth & Cordes 2012, Chowdhury 2021, 2022) as it remains a technical challenge to track to such small species in the field. We hope that the present UAV-based methodology exploration can serve as a starting point and motivate following studies on spatial behaviour and movement ecology in flying insects.

Acknowledgments

We thank the owner of the crop fields for authorizing our presence and UAV flights over his property. We thank Robin Baker and Tristan Lafranchis for discussions on the behaviour and biology of *Pieris brassicae* and *Pieris rapae*. We thank Ty Hedrick for discussions on the use of DLTdv. We thank Virginie Durier and Simon Benhamou for discussions on circular statistics. We thank François Cornu and Ignace Cacciaguerra (Pôle drones, CNRS) for their assistance with the UAV pilot certification, and regulations on the use of UAVs in the field .

Funding

The authors declare that they have received no specific grant for this study, funded solely by the basic allowances from the CNRS and the University of Rennes.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code and supplementary information availability

Data and scripts needed to replicate the statistical results presented here will be available at https://dx.doi.org/10.6084/m9.figshare.28043333, as well as supplementary figures and table.

References

Abrahms B, Seidel DP, Dougherty E, Hazen EL, Bograd SJ, Wilson AM, Weldon McNutt J, Costa DP, Blake S, Brashares JS, Getz WM (2017) Suite of simple metrics reveals common movement syndromes across vertebrate taxa. *Movement Ecology*, **5**, 12. https://doi.org/10.1186/s40462-017-0104-2

Baker R (1968) A possible method of evolution of the migratory habit in butterflies. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **253**, 309–341. https://doi.org/10.1098/rstb.1968.0003

Batsleer F, Bonte D, Dekeukeleire D, Goossens S, Poelmans W, Van der Cruyssen E, Maes D, Vandegehuchte ML (2020) The neglected impact of tracking devices on terrestrial arthropods. *Methods in Ecology and Evolution*, **11**, 350–361. https://doi.org/10.1111/2041-210X.13356

Berens P (2009) CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software*, **31**, 1–21. https://doi.org/10.18637/jss.v031.i10

Brower LP (1995) Understanding and misunderstanding the migration of the monarch butterfly (Nymphalidae) in North America: 1857-1995. *Journal of the Lepidopterists' Society*, **49**, 304--385.

Brussard PF, Ehrlich PR (1970) Adult Behavior and Population Structure in *Erebia epipsodea* (Lepidoptera: Satyrinae). *Ecology*, **51**, 880–885. https://doi.org/10.2307/1933981

Cagnacci F, Boitani L, Powell RA, Boyce MS (2010) Animal ecology meets GPS-based radiotelemetry: a perfect storm of opportunities and challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 2157–2162. https://doi.org/10.1098/rstb.2010.0107

Charbonneau P, Lemaître J (2021) Revue des applications et de l'utilité des drones en conservation de la faune. *Le Naturaliste canadien*, **145**, 3–34. https://doi.org/10.7202/1080620ar

Chowdhury S, Fuller RA, Dingle H, Chapman JW, Zalucki MP (2021) Migration in butterflies: a global overview. *Biological Reviews*, **96**, 1462–1483. https://doi.org/10.1111/brv.12714

Chowdhury S, Zalucki MP, Amano T, Poch TJ, Lin M-M, Ohwaki A, Lin D-L, Yang L, Choi S-W,
Jennions MD, Fuller RA (2022) Trends and progress in studying butterfly migration.
Integrative Conservation, **1**, 8–24. https://doi.org/10.1002/inc3.13
Cook PM, Tordoff GM, Davis TM, Parsons MS, Dennis EB, Fox R, Botham MS, Bourn NAD (2022)
Traits data for the butterflies and macro-moths of Great Britain and Ireland. *Ecology*, **103**,
e3670. https://doi.org/10.1002/ecy.3670
Crall JD, Gravish N, Mountcastle AM, Combes SA (2015) BEEtag: A Low-Cost, Image-Based
Tracking System for the Study of Animal Behavior and Locomotion. *PLOS ONE*, **10**, e0136487.
https://doi.org/10.1371/journal.pone.0136487
Da Silveira NS, Niebuhr BBS, Muylaert R de L, Ribeiro MC, Pizo MA (2016) Effects of Land Cover
on the Movement of Frugivorous Birds in a Heterogeneous Landscape. *PLOS ONE*, **11**,
e0156688. https://doi.org/10.1371/journal.pone.0156688
Delattre T, Vernon P, Burel F (2013) An agri-environmental scheme enhances butterfly dispersal
in European agricultural landscapes. *Agriculture, ecosystems & environment*, **166**, 102–109.
Dell AI, Bender JA, Branson K, Couzin ID, Polavieja GG de, Noldus LPJJ, Pérez-Escudero A,
Perona P, Straw AD, Wikelski M, Brose U (2014) Automated image-based tracking and its
application in ecology. *Trends in Ecology & Evolution*, **29**, 417–428.
https://doi.org/10.1016/j.tree.2014.05.004
de Margerie E, Pichot C, Benhamou S (2018) Volume-concentrated searching by an aerial
insectivore, the common swift, *Apus apus*. *Animal Behaviour*, **136**, 159–172.
https://doi.org/10.1016/j.anbehav.2017.11.026
de Margerie E, Simonneau M, Caudal J-P, Houdelier C, Lumineau S (2015) 3D tracking of animals
in the field using rotational stereo videography. *The Journal of Experimental Biology*, **218**,
2496–2504. https://doi.org/10.1242/jeb.118422
Dudley R, Srygley RB (1994) Flight Physiology of Neotropical Butterflies: Allometry of Airspeeds
During Natural Free Flight. *Journal of Experimental Biology*, **191**, 125–139.
https://doi.org/10.1242/jeb.191.1.125
Dunn K A (2024) remarkable migration of the Small Cabbage White Pieris rapae (L.)
(Lepidoptera: Pieridae) in the Melbourne region of Victoria, Australia that spanned four days
in November 2023, with evidence of post-migratory clustering. *Calodema*, **1040**, 1–44.
Fernández P, Rodríguez A, Obregón R, de Haro S, Jordano D, Fernández-Haeger J (2016) Fine
Scale Movements of the Butterfly *Plebejus argus* in a Heterogeneous Natural Landscape as
Revealed by GPS Tracking. *Journal of Insect Behavior*, **29**, 80–98.
https://doi.org/10.1007/s10905-016-9543-7
Gilbert N, Raworth DA (2005) Movement and migration patterns in *Pieris rapae* (Pieridae).
Journal of the Lepidopterists' Society, **59**, 10–18.
Haalck L, Mangan M, Webb B, Risse B (2020) Towards image-based animal tracking in natural
environments using a freely moving camera. *Journal of neuroscience methods*, **330**, 108455.
Hedrick TL (2008) Software techniques for two- and three-dimensional kinematic
measurements of biological and biomimetic systems. *Bioinspiration & Biomimetics*, **3**,
034001. https://doi.org/10.1088/1748-3182/3/3/034001
Hedrick TL, Pichot C, de Margerie E (2018) Gliding for a free lunch: biomechanics of foraging
flight in common swifts (*Apus apus*). *Journal of Experimental Biology*, **221**, jeb186270.
https://doi.org/10.1242/jeb.186270

Hernández-Pliego J, Rodríguez C, Dell'Omo G, Bustamante J (2017) Combined use of tri-axial
accelerometers and GPS reveals the flexible foraging strategy of a bird in relation to weather
conditions. *PLOS ONE*, **12**, e0177892. https://doi.org/10.1371/journal.pone.0177892

Ivosevic B, Han Y-G, Kwon O (2017) Monitoring butterflies with an unmanned aerial vehicle:
current possibilities and future potentials. *Journal of Ecology and Environment*, **41**, 12.
https://doi.org/10.1186/s41610-017-0028-1

Jackson BE, Evangelista DJ, Ray DD, Hedrick TL (2016) 3D for the people: multi-camera motion
capture in the field with consumer-grade cameras and open source software. *Biology Open*,
5, 1334–1342. https://doi.org/10.1242/bio.018713

John E, Cottle N, McArthur A, Makris C (2008) Eastern Mediterranean migrations of the Small
White *Pieris rapae* (Linnaeus, 1758) (Lepidoptera: Pieridae): observations from Cyprus 2001
and 2007. *Entomologist's Gazette*, **59**, 71–78.

Jones RE, Gilbert N, Guppy M, Nealis V (1980) Long-Distance Movement of *Pieris rapae*. *Journal
of Animal Ecology*, **49**, 629–642. https://doi.org/10.2307/4268

Joo R, Picardi S, Boone ME, Clay TA, Patrick SC, Romero-Romero VS, Basille M (2022) Recent
trends in movement ecology of animals and human mobility. *Movement Ecology*, **10**, 26.
https://doi.org/10.1186/s40462-022-00322-9

Kashetsky T, Avgar T, Dukas R (2021) The Cognitive Ecology of Animal Movement: Evidence
From Birds and Mammals. *Frontiers in Ecology and Evolution*, **9**.
https://doi.org/10.3389/fevo.2021.724887

Kays R, Tilak S, Crofoot M, Fountain T, Obando D, Ortega A, Kuemmeth F, Mandel J, Swenson G,
Lambert T, Hirsch B, Wikelski M (2011) Tracking Animal Location and Activity with an
Automated Radio Telemetry System in a Tropical Rainforest. *The Computer Journal*, **54**,
1931–1948. https://doi.org/10.1093/comjnl/bxr072

Kissling WD, Pattemore DE, Hagen M (2014) Challenges and prospects in the telemetry of
insects. *Biological Reviews*, **89**, 511–530. https://doi.org/10.1111/brv.12065

Kitamura T, Imafuku M (2015) Behavioural mimicry in flight path of Batesian intraspecific
polymorphic butterfly *Papilio polytes*. *Proceedings of the Royal Society B: Biological Sciences*,
282, 20150483. https://doi.org/10.1098/rspb.2015.0483

Kleckova I, Linke D, Rezende FDM, Rauscher L, Le Roy C, Matos-Maraví P (2024) Flight behaviour
diverges more between seasonal forms than between species in Pieris butterflies. *Ecology
and Evolution*, **14**, e70012. https://doi.org/10.1002/ece3.70012

Knight SM, Pitman GM, Flockhart DTT, Norris DR (2019) Radio-tracking reveals how wind and
temperature influence the pace of daytime insect migration. *Biology Letters*, **15**, 20190327.
https://doi.org/10.1098/rsbl.2019.0327

Knoblauch A, Thoma M, Menz MH (2021) Autumn southward migration of dragonflies along the
Baltic coast and the influence of weather on flight behaviour. *Animal Behaviour*, **176**, 99–109.

Koger B, Deshpande A, Kerby JT, Graving JM, Costelloe BR, Couzin ID (2023) Quantifying the
movement, behaviour and environmental context of group-living animals using drones and
computer vision. *Journal of Animal Ecology*, **92**, 1357–1371. https://doi.org/10.1111/1365-
2656.13904

Lafranchis T, Jutzeler D, Guillosson J-Y, Kan P, Kan B (2015) *La vie des papillons : écologie,
biologie et comportement des Rhopalocères de France*. Diatheo. Paris.

Larranaga N, Baguette M, Calvez O, Trochet A, Ducatez S, Legrand D (2013) Intra- and inter-
individual variation in flight direction in a migratory butterfly co-vary with individual
mobility. *Journal of Experimental Biology*, **216**, 3156–3163.
https://doi.org/10.1242/jeb.082883

Le Roy C, Amadori D, Charberet S, Windt J, Muijres FT, Llaurens V, Debat V (2021) Adaptive
evolution of flight in *Morpho* butterflies. *Science*, **374**, 1158–1162.
https://doi.org/10.1126/science.abh2620

Lihoreau M, Ings TC, Chittka L, Reynolds AM (2016) Signatures of a globally optimal searching
strategy in the three-dimensional foraging flights of bumblebees. *Scientific Reports*, **6**, 30401.
https://doi.org/10.1038/srep30401

Lihoreau M, Raine NE, Reynolds AM, Stelzer RJ, Lim KS, Smith AD, Osborne JL, Chittka L (2012)
Radar Tracking and Motion-Sensitive Cameras on Flowers Reveal the Development of
Pollinator Multi-Destination Routes over Large Spatial Scales. *PLOS Biology*, **10**, e1001392.
https://doi.org/10.1371/journal.pbio.1001392

Maggiora R, Saccani M, Milanesio D, Porporato M (2019) An Innovative Harmonic Radar to Track
Flying Insects: the Case of *Vespa velutina*. *Scientific Reports*, **9**, 11964.
https://doi.org/10.1038/s41598-019-48511-8

Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE (2008) A movement
ecology paradigm for unifying organismal movement research. *Proceedings of the National
Academy of Sciences*, **105**, 19052–19059. https://doi.org/10.1073/pnas.0800375105

Nikolaev NA (1974) Flight speed and types for certain Pieridae and Nyphalidae (Lepidoptera).
Moscow University Biological Bulletin, **29**, 17–20.

Oelmann Y, Fiedler D, Michaelis R, Leivits M, Braun A, Gschwind P, Neidhardt H, Willigalla C
(2023) Autumn migration of the migrant hawker (*Aeshna mixta*) at the Baltic coast. *Movement
Ecology*, **11**, 52. https://doi.org/10.1186/s40462-023-00415-z

Ovaskainen O, Smith AD, Osborne JL, Reynolds DR, Carreck NL, Martin AP, Niitepõld K, Hanski I
(2008) Tracking butterfly movements with harmonic radar reveals an effect of population
age on movement distance. *Proceedings of the National Academy of Sciences*, **105**, 19090–
19095. https://doi.org/10.1073/pnas.0802066105

Piccardi M (2004) Background subtraction techniques: a review. In: *2004 IEEE International
Conference on Systems, Man and Cybernetics (IEEE Cat. No.04CH37583)* , pp. 3099–3104 vol.4.
https://doi.org/10.1109/ICSMC.2004.1400815

Rhodes MW, Bennie JJ, Spalding A, ffrench-Constant RH, Maclean IMD (2022) Recent advances
in the remote sensing of insects. *Biological Reviews*, **97**, 343–360.
https://doi.org/10.1111/brv.12802

Rieucau G, Kiszka JJ, Castillo JC, Mourier J, Boswell KM, Heithaus MR (2018) Using unmanned
aerial vehicle (UAV) surveys and image analysis in the study of large surface-associated
marine species: a case study on reef sharks *Carcharhinus melanopterus* shoaling behaviour.
Journal of Fish Biology, **93**, 119–127. https://doi.org/10.1111/jfb.13645

Roer H (1959) Über Flug- und Wandergewohnheiten von Pieris brassicae L. *Zeitschrift für
Angewandte Entomologie*, **44**, 272–309. https://doi.org/10.1111/j.1439-0418.1959.tb00927.x

Roer H (1961) Ergebnisse mehrjahriger Markierungsversuche zur Erforschung der Flug-und
Wandergewohnheiten europäischer Schmetterlinge. *Zool. Anz*, **167**, 456–463.

Rotics S, Kaatz M, Resheff YS, Turjeman SF, Zurell D, Sapir N, Eggers U, Flack A, Fiedler W, Jeltsch
F, Wikelski M, Nathan R (2016) The challenges of the first migration: movement and
behaviour of juvenile vs. adult white storks with insights regarding juvenile mortality.
Journal of Animal Ecology, **85**, 938–947. https://doi.org/10.1111/1365-2656.12525

Ruaux G, Monmasson K, Hedrick TL, Lumineau S, de Margerie E (2023) Flight behaviours and
energy savings in adult and juvenile house martins (*Delichon urbicum*) foraging near their
breeding colony. *Behavioral Ecology and Sociobiology*, **77**, 63.
https://doi.org/10.1007/s00265-023-03332-8

Shepard ELC, Lambertucci SA, Vallmitjana D, Wilson RP (2011) Energy Beyond Food: Foraging
Theory Informs Time Spent in Thermals by a Large Soaring Bird. *PLOS ONE*, **6**, e27375.
https://doi.org/10.1371/journal.pone.0027375

Sherub S, Bohrer G, Wikelski M, Weinzierl R (2016) Behavioural adaptations to flight into thin
air. *Biology Letters*, **12**, 20160432. https://doi.org/10.1098/rsbl.2016.0432

Spieth H, Cordes R (2012) Geographic comparison of seasonal migration events of the large
white butterfly, *Pieris brassicae*. *Ecological Entomology*, **37**. https://doi.org/10.1111/j.1365-
2311.2012.01385.x

Srygley RB, Oliveira EG (2001) Sun Compass and Wind Drift Compensation in Migrating
Butterflies. *The Journal of Navigation*, **54**, 405–417.
https://doi.org/10.1017/S0373463301001448

Stefanescu C, Páramo F, Åkesson S, Alarcón M, Ávila A, Brereton T, Carnicer J, Cassar LF, Fox R,
Heliölä J, Hill JK, Hirneisen N, Kjellén N, Kühn E, Kuussaari M, Leskinen M, Liechti F, Musche
M, Regan EC, Reynolds DR, Roy DB, Ryrholm N, Schmaljohann H, Settele J, Thomas, van
Swaay C, Chapman JW (2013) Multi-generational long-distance migration of insects:
studying the painted lady butterfly in the Western Palaearctic. *Ecography*, **36**, 474–486.
https://doi.org/10.1111/j.1600-0587.2012.07738.x

Stürzl W, Zeil J, Boeddeker N, Hemmi JM (2016) How wasps acquire and use views for homing.
Current Biology, **26**, 470–482.

Suchan T, Bataille CP, Reich MS, Toro-Delgado E, Vila R, Pierce NE, Talavera G (2024) A trans-
oceanic flight of over 4,200 km by painted lady butterflies. *Nature Communications*, **15**, 5205.
https://doi.org/10.1038/s41467-024-49079-2

Theriault DH, Fuller NW, Jackson BE, Bluhm E, Evangelista D, Wu Z, Betke M, Hedrick TL (2014)
A protocol and calibration method for accurate multi-camera field videography. *The Journal*
of Experimental Biology, **217**, 1843–1848. https://doi.org/10.1242/jeb.100529

Vepsäläinen K (1968) The immigration of Pieris brassicae L. (Lep., Pieridae) into Finland in 1966,
with a general discussion on insect migration. *Annales Entomologici Fennici*, **34**, 223–243.

Vo-Doan TT, Titov VV, Harrap MJM, Lochner S, Straw AD (2024) High-resolution outdoor
videography of insects using Fast Lock-On tracking. *Science Robotics*, **9**, eadm7689.
https://doi.org/10.1126/scirobotics.adm7689

Walter T, Degen J, Pfeiffer K, Stöckl A, Montenegro S, Degen T (2021) A new innovative real-time
tracking method for flying insects applicable under natural conditions. *BMC Zoology*, **6**, 35.
https://doi.org/10.1186/s40850-021-00097-3

Wang D, Shao Q, Yue H (2019) Surveying Wild Animals from Satellites, Manned Aircraft and
Unmanned Aerial Systems (UASs): A Review. *Remote Sensing*, **11**, 1308.
https://doi.org/10.3390/rs11111308

Whitfield DP, Fielding AH, Anderson D, Benn S, Reid R, Tingay R, Weston E (2024) Sex difference
in natal dispersal distances of Golden Eagles *Aquila chrysaetos* in Scotland. *Ibis*, **166**, 146–
155. https://doi.org/10.1111/ibi.13225
Williams CB, Cockbill GF, Gibbs ME, Downes JA (1942) Studies in the Migration of Lepidoptera.
Transactions of the Royal Entomological Society of London, **92**, 101–283.
https://doi.org/10.1111/j.1365-2311.1942.tb03319.x
Wilson RP, McMahon CR (2006) Measuring devices on wild animals: what constitutes acceptable
practice? *Frontiers in Ecology and the Environment*, **4**, 147–154. https://doi.org/10.1890/1540-
9295(2006)004[0147:MDOWAW]2.0.CO;2