Effects of copepod chemical cues on intra- and extracellular

toxins in two species of *Dinophysis*

Milad Pourdanandeh[a*], Véronique Séchet[b], Liliane Carpentier[b], Damien Réveillon[c], Fabienne

Hervé[c], Clarisse Hubert[c], Philipp Hess[c], Erik Selander[a,d].

a: Department of Marine Sciences, University of Gothenburg, Medicinaregatan 7b, SE41390,

Gothenburg, Sweden

b: IFREMER, PHYTOX, PHYSALG, F-44000 Nantes, France

c: IFREMER, PHYTOX, METALG, F-44000 Nantes, France

d: Functional ecology section - Department of Biology, Lund University, Lund, Sweden

* Corresponding author: milad.pourdanandeh@gu.se

Abstract

Copepods may contribute to harmful algal bloom formation by selectively rejecting harmful cells. Additionally, copepods and the chemical cues they exude, copepodamides, have been shown to induce increased toxin production in paralytic and amnesic toxin producing microalgae. However, it is unknown if diarrhetic shellfish toxin (DST) producers such as *Dinophysis* respond to copepods or copepodamides in a similar fashion. Here we expose laboratory cultures of *Dinophysis sacculus* and *D. acuminata* to direct grazing by *Acartia* sp. copepods or copepodamides and measure their toxins after three days. Total *Dinophysis*-produced toxins (DPTs), okadaic acid, pectenotoxin-2, and C9-diol ester of okadaic acid, increased by 8 - 45% in *D. sacculus* but was significantly different from controls only in the highest (10 nM) copepodamide treatment whereas toxin content was not affected in *D. acuminata*. Growth rate was low across all groups and explained up to 91% of the variation in toxin content. DPTs were redistributed from internal compartments to the extracellular medium in the highest copepodamide treatments (5 - 10 nM), which were two to three times higher than controls and indicates an active release or passive leakage of toxins. Untargeted analysis of endometabolomes indicated significant changes in metabolite profiles for both species in response to the highest copepodamide treatments, independent of known toxins. However, it is not clear whether these are stress responses or caused by more complex mechanisms. The relatively small grazer-induced effect in *Dinophysis* observed here, compared to several species of *Alexandrium* and *Pseudo-nitzschia* reported previously, suggests that DPT production in *Dinophysis* is likely not induced by copepods, except perhaps in patches with high copepod densities. Thus, DPTs may, represent either a constitutive chemical defence for *Dinophysis*, or serve an altogether different purpose.

1. Introduction

Dinophysis is a highly specialised genus of non-constitutive mixotrophic dinoflagellates (Reguera et al., 2012; Mitra et al., 2016). They feed on ciliates of the genus *Mesodinium*, and retain the photosynthetically active chloroplasts (Hansen et al., 2013) that the ciliate, in turn, obtained from their own cryptomonad prey such as *Teleaulax* spp., *Plagioselmis* spp., and *Geminigera* spp. At least eight species of *Dinophysis* can be reared on a diet consisting of the ciliate *Mesodinium rubrum* (Park et al., 2006; Hansen et al., 2013; Séchet et al., 2021) which in turn are fed with cryptophytes. *Dinophysis* spp. grow at rates similar to other small dinoflagellates in fed cultures, but more slowly or not at all in the prolonged absence of prey (Reguera et al., 2012).

The link between diarrhetic shellfish toxins (DSTs: okadaic acid group toxins and dinophysistoxins (DTXs)) and the dinoflagellate *Dinophysis fortii* was established following multiple episodes of shellfish poisoning in Japan during the late 1970s (Yasumoto et al., 1980). The active compounds, identified as groups of polyethers, notably include okadaic acid (OA), DTXs, and their derivatives. They also produce pectenotoxins (PTXs), initially included as DSTs (Yasumoto et al., 1985; Ishige et al., 1988) but now concluded to be non-diarrhetic in humans (Terao et al., 1986; Matsushima et al., 2015). In sufficient amounts, the DSTs cause severe gastroenteritis in humans (Yasumoto et al., 1985) while pectenotoxins have been shown to negatively impact fertilisation success of oyster gametes (Gaillard et al., 2020),development of fish embryos (Gaillard et al., 2023), be hepatotoxic to some vertebrate hepatocytes (Terao et al., 1986; Fladmark et al., 1998), and are potently cytotoxic to several human cell lines (Jung et al., 1995).

Dinophysis is a cosmopolitan genus (Reguera et al., 2012) and the resulting syndrome, Diarrhetic Shellfish Poisoning (DSP), constitutes the largest share of harmful algal events in the North-East Atlantic (55%) reported in the Harmful Algae Event Database

(Karlson et al., 2021). *Dinophysis* blooms are hard to foresee due to their unpredictable

population dynamics and highly variable toxin production. The reason for their variable toxin

content is not fully understood (Karlson et al., 2021) but is negatively correlated to cell

density on the Swedish west coast. For example, Lindahl and colleagues (2007) reported

okadaic acid content higher than 5 pg cell^{-1} in blooms with less than 100 cells L^{-1} but, with

few exceptions, below 2 pg cell^{-1} in blooms with more than 250 cells L^{-1}. Recent findings

suggest that interactions with grazers may contribute to harmful algal bloom formation

(Selander et al., 2019; Trapp et al., 2021). Chemical alarm cues from copepods,

copepodamides (Selander et al., 2015), induce several-fold increases in cellular toxin content

for paralytic and amnesic shellfish toxin (PSTs and ASTs, respectively) producers such as

Alexandrium and *Pseudo-nitzschia*. (Selander et al., 2006; Tammilehto et al., 2015; Selander

et al., 2019) and vary in composition between copepod taxa (Arnoldt et al., 2024). Moreover,

copepods reject the more toxic cells and feed selectively on less defended cells (Xu and

Kiørboe, 2018; Ryderheim et al., 2021; Olesen et al., 2022). As a result, the relative

abundance of harmful cells will increase and to some extent contribute to harmful algal

bloom formation (Xu et al., 2018). Recently, Trapp and colleagues (2021) tested this

hypothesis by exploring the correlation between copepod biomass, or copepodamides in

mussels, with the amount of DSTs in mussels on the Swedish west coast. They found that

both copepod biomass and the concentration of copepodamides in blue mussels correlate to

time-lagged DST content and can be used together with *Dinophysis* spp. cell densities to

improve lead time and forecasting precision of DST concentrations in shellfish. It is,

however, not known if copepods, or their chemical cues, induce the production of toxins in

Dinophysis, or how their biosynthetic machinery responds to grazer presence.

Here, we address this knowledge gap by exposing laboratory cultures of

Dinophysis sacculus and *D. acuminata* to direct grazing from *Acartia* sp. copepods, and to

grazer-free copepodamide extracts. We measure growth rates, intra- and extracellular DPTs,

and perform untargeted metabolomic analyses to determine the effect of copepod grazers on

DPT production and metabolic state of *Dinophysis* spp. cells.

2. Materials and Methods

2.1 *Dinophysis* cultures

Dinophysis sacculus (IFR-DSa-01Lt, accession no. MT 37 1867) and *Dinophysis acuminata*

(IFR-DAu-01Ey, accession no. MT 365104), isolated from Eyrac Pier, Arcachon, France in

May 2015 and April 2020 respectively, were used. Both cultures were kept in diluted K (-Si)/

L1/20 (-Si) culture media (Guillard and Hargraves, 1993) prepared with filtered seawater (0.2

µm Steritop Corning, Corning, United States) at pH 8.2 and a salinity of 35. They were

maintained in a thermo-regulated room at 17 °C and ~55 µmol m^2 s^{-1} photosynthetically

active radiation on a 12h:12h light:dark cycle in 50 mL Erlenmeyer flasks. Irradiance was

delivered by Osram Fluora 36W (Munich, Germany) and Philips Daylight 36W (Amsterdam,

Netherlands). *Dinophysis* cultures were fed daily with 100 µL concentrated *Mesodinium*

rubrum culture (DK-2009), isolated from Helsingør harbour (Denmark) in 2009, which in

turn had been fed *Teleaulax amphioxeia* (strain AND-0710). Before the experiment, cultures

were checked under a microscope to make sure that all *Mesodinium* cells had been consumed.

2.2 Copepodamides

103 The copepodamides used here were extracted and purified from commercially fished *Calanus*

104 *finmarchicus*, as described by Selander and colleagues (2015). The copepodamide blend

105 from *Calanus* sp. (Fig. 1A) is more complex than that of the *Acartia* sp. copepods used in

106 experiments but also includes the copepodamides found in *Acartia* sp. (Fig. 1B,

107 Supplementary Table S1).

2.3 Experimental procedure

Cultures of *D. sacculus* and *D. acuminata*, with cell concentrations of 1728 ± 62 cells mL^{-1}

and 1254 ± 76 cells mL^{-1} (mean $\pm$ SEM) respectively, were divided between fifteen 20 mL

Erlenmeyer flasks. Nine flasks for each species were prepared by adding copepodamides

dissolved in methanol (10 µL) to achieve nominal copepodamide concentrations of 1, 5, or 10

nM, each in triplicate. Triplicate controls for each species were prepared in the same way, but

with 10 µL methanol without copepodamides. The methanol was allowed to evaporate before

adding the

Dinophysis cultures, leaving the copepodamides coated onto the culture flasks walls. The

resulting effective concentration in the culture media, due to a combination of dissolution and

degradation, typically averages between 0.1% and 1.5% of the nominal concentration

(Selander et al., 2019). To maintain copepodamide stimuli throughout the experimental

period, the experimental cultures were transferred to freshly coated Erlenmeyer flasks after

48 hours. The experiment was concluded after three days (68.5 h).

To control for the possibility that *Dinophysis* respond to other copepod cues

than copepodamides, we included a living copepod treatment as a positive control. The

copepods (*Acartia* sp.) were collected with a plankton net at Le Croisic (WGS84

decimal degrees: 47.299796, -2.517140) the day before the experimental start and

individually isolated with a Pasteur pipette under a dissecting microscope into a multi-well

dish. The water was carefully replaced by medium three times to avoid contaminating

organisms or chemical stimuli from the net haul. Finally, a single copepod was added to three

flasks (resulting in 50 ind L^{-1}) for both *Dinophysis* species together with a small volume of

medium (<100 µL). The copepods were inspected daily, and non-motile copepods were

replaced. In total, two copepods, one in each *Dinophysis* species, were replaced. All

experimental units received 100 µL *Mesodinium* on day 0, 1, and 2 (85, 53, and 89×10^3 cells

mL^{-1} respectively) to avoid starvation of the *Dinophysis* cells.

2.4 Toxin and metabolomic samples

Triplicate samples were taken from all treatments of both species at the start and end of the

experiment. Well-mixed samples of 15 mL were centrifuged (4300g, 20 min, 4 °C) and

supernatants carefully pipetted into a 50 mL centrifuge tube. The remaining pellets and

supernatants were stored frozen at -80 °C until extraction.

2.5 Growth rate

At the end of the experiment, a well-mixed sample (1.5 mL) from each replicate was

removed for cell enumeration. The sample was diluted to 15 mL and preserved with acid

Lugol's solution. Aliquots of 1 mL were counted under an inverted microscope until >500

cells per replicate had been counted. Specific growth rate (μ) was calculated as:

$$\mu = \frac{ln(N_1) - ln(N_0)}{t_1 - t_0}$$

where N_0 and N_1 are cell concentrations at the start (t_0) and end (t_1) of the experiment (Frost,

1972).

2.6 Copepodamide analysis

Five adult *Acartia* copepods, possibly including some non-conspicuous late-stage

copepodites (C5), were pipetted into an HPLC vial; the water removed with a Pasteur pipette,

and extracted in 1.5 mL methanol for at least 24 h. The methanol extract was transferred to a

glass tube and evaporated under a stream of nitrogen gas at 40 °C, resuspended in 50 μL

methanol and transferred to a low volume HPLC-vial insert. Copepodamides were analysed

on an Agilent 1200 series HPLC fitted with a 6470 triple quadrupole mass spectrometer

(Santa Clara, CA, USA). A multiple reaction monitoring experiment targeting the 10 known

copepodamides was performed to quantify both the copepodamide extract from *Calanus* sp.

used in experiments and the extract from *Acartia* sp. Grebner et al. (2019), for further details

see Selander et al. (2019).

2.7 Toxin analyses

Okadaic acid group toxins and pectenotoxins were extracted following Séchet and colleagues

(2021). Briefly, intracellular toxins were extracted from cell pellets twice with 0.75 mL

methanol, vortexed, and sonicated (25 kHz for 10 min). Samples were then centrifuged for 5

min (4300 g, 4 °C), the extracts were pooled, filtered (0.2 µm Nanosep MF filter, Pall), and

stored at -20 °C until analysis with liquid chromatography coupled to tandem mass

spectrometry (LC-MS/MS). Extracellular toxins were recovered from the supernatant after

two liquid-liquid partitioning procedures with 25 mL dichloromethane. After evaporation to

dryness, and resuspension in 1 mL methanol, the extracts were filtered and stored at -20 °C

until LC-MS/MS analysis.

167   Toxins were separated on an a ultra-fast liquid chromatography system

(Prominence UFLC-XR, Shimadzu, Japan), equipped with a reversed-phase C18 Kinetex

column (100 Å, 2.6 μm, 100 × 2.1 mm, Phenomenex, Le Pecq, France) and coupled to a

triple quadrupole/ion trap mass spectrometer (4000QTrap, Sciex, Framingham, MA, USA),

as described by Séchet and colleagues (2021). Authentic standards (OA, DTX-1, DTX-2, and

PTX-2) for quantification were obtained from the National Research Council of Canada

(Halifax, Nova Scotia, Canada) and the uncertified C9-diol ester of OA (C9) from Cifga

(through Novakits, Nantes, France).

2.8 Metabolomics

Metabolomic profiles of the extracts prepared for intracellular toxin analysis above were

acquired by liquid chromatography coupled to high resolution mass spectrometry (LC-

HRMS) as in Gémin et al. (2021). Briefly, an UHPLC (1290 Infinity II, Agilent technologies,

Santa Clara, CA, USA) coupled to a quadrupole time-of-flight mass spectrometer (QTOF

6550, Agilent technologies, Santa Clara, CA, USA) equipped with a Dual Jet Stream ESI

interface was used. Both positive (+MS) and negative (-MS) full scan modes were used over

a mass-to-charge ratio (m/z) ranging from 100 to 1700. The injection volume was 5 µL. The

batch was prepared as recommended by Broadhurst and colleagues (2018), by using a mix of

four phycotoxins (OA, DTX-1, DTX2, PTX-2) as the standard reference material and a

mixture of all samples for the qualitative control (QC) samples. The injection order was

randomised, and QCs were injected every 5 samples.

187   LC-HRMS raw data was converted to mzXML-format using MS-Convert in

ProteoWizard 3.0 (Chambers et al., 2012) and pre-processed with the

Workflow4Metabolomics 4.0 e-infrastructure (Guitton et al., 2017). Peak picking, grouping,

retention time correction, and peak filling were performed with its "CentWave",

"PeakDensity", "PeakGroups", and "FillchromPeaks" algorithms. Annotation (e.g. isotopes

and adducts) was conducted with the "CAMERA" algorithm (Kuhl et al., 2012). Intra-batch

signal intensity drift was corrected for, in the –MS data only, by fitting a locally quadratic

(LOESS) regression model to the QC values (van der Kloet et al., 2009; Dunn et al., 2011).

Three successive filtering steps based on signal-to-noise ratio with blanks, variability among

pools, and autocorrelation were applied using in-house scripts in R, as described by Georges

des Aulnois et al. (2020). Pre-processing of +MS data matrices for each species led

to 19427/20570 features for *D. acuminata* and *D. sacculus*, respectively, while -MS data

matrices led to 4406/4816. After filtrations 1788/695 and 422/324 features remained,

respectively.

201   The data matrix in +MS and -MS for each species was log transformed and

Pareto scaled, and a weighting factor based on the maximal cell number was applied for each

sample. Unsupervised principal component analysis (PCA) was used for data exploration,

while features significantly affected by copepodamides or copepods were selected by

ANOVA followed by a Tukey's post hoc test, with adjusted p-value cut-offs set to 0.001 or

0.01 depending on the species and mode of ionisation. Statistical analyses of metabolomics

were performed with MetaboAnalyst 5.0 (Pang et al., 2021)

208   In parallel, HRMS data were acquired by autoMS/MS as in Georges des

Aulnois et al. (2019) to attempt annotation of the significant features revealed by

metabolomics, using both GNPS (Wang et al., 2016) and SIRIUS (Dührkop et al., 2019). The

comparison of MS/MS spectra with these reference databases provided a level 2 of

annotation, as defined by Schymanski and colleagues (2014). In addition, diagnostic ions in

MS/MS spectra were considered, especially for the

(lyso)diacylglycerylcarboxyhydroxymethylcholine (lysoDGCC) and

(lyso)phosphatidylcholine (lysoPC) families (Roach et al., 2021; Rey et al., 2022). When the

fatty acyl chain could not be confirmed, the features were considered "lysoDGCC-like" or

"lysoPC-like" (.xlsx file; Supplementary spreadsheet 1).

2.9 Statistical analyses of growth rates and toxins

For analyses of toxin induction, the three toxin congeners (OA, PTX-2, C9) were pooled

(referred to as total toxins or total DPTs). Differences in specific growth rate (μ) and

intracellular toxins (% of total toxins) as a function of species (two levels: *D. sacculus* & *D.*

acuminata) and treatment (five levels: Control, Copepod, 1 nM, 5 nM & 10 nM) were tested

with two-way analyses of variance (ANOVA). Volume normalised total (intracellular and

extracellular) DPT content (ng mL^{-1}) was first analysed with a two-way analysis of

covariance (ANCOVA) with growth rate as covariate. Because there was no interaction effect

between treatment and species, but a significant difference between the species, they were

analysed with separate one-way ANCOVAs instead. The proportions (%) of individual toxin

congeners (OA, PTX-2, and C9) in *D. sacculus* were tested with separate one-way ANOVAs.

All proportions were logit (Ln(p/1-p)) transformed for better statistical properties (Warton

and Hui, 2011).

Assumptions of linearity were evaluated visually and with linear regressions.

Normality was visually inspected and formally tested with Shapiro-Wilk's test, and

homogeneity of variances was tested with Levene's test. Significant ANOVAs ($p \leq 0.05$)

were followed up with Tukey's honest significant difference (HSD) post-hoc procedure.

Estimates of covariate corrected mean toxin content, and their 95% confidence intervals (CIs

hereafter), were extracted from the linear models using the base R *predict* function at

covariate level (i.e. growth rate) set to zero. Unless otherwise stated, all summary statistics

values are given as mean (95% CI: lower limit - upper limit) or (mean, lower CI limit - upper

CI limit). Effect sizes are derived from log response ratios (LRR) and presented as the

percentage difference (lower 95% CI limit – upper CI confidence) or as (mean, lower CI

limit–upper CI limit) (Hedges et al., 1999). Partial eta-squared (η^2_p) was used as effect size

for linear models.

Statistical analyses of growth rates and toxins, and visualisations of growth rate,

toxins, and PCA scores derived from metabolomics data were performed in R v.4.4.1 (R Core

Team, 2024) using RStudio v.2024.4.1.748 (Posit team, 2024). The code (.html file;

Supplementary code & output) and the toxin analysis data it uses (.csv file; Supplementary

spreadsheet 2) are available as supplementary materials. These, Supplementary spreadsheet 1,

and the HRMS metabolomics data are also openly accessible from a public Zenodo

repository (Pourdanandeh et al., 2024). Supplementary tables and figures directly referred to

in the text is found in Supplementary information

3. Results

3.1 Copepodamides

The *Acartia* sp. copepods used as grazers in this experiment contained approximately ~0.5

pmol ind^{-1} dihydro-copepodamides (hereafter dhCAs). The most abundant compound was

18:4-dhCA, followed by 20:5-dhCA and 22:6-dhCA, along with trace amounts of the

deacylated molecular scaffold (Fig. 1B). The copepodamides extracted from *Calanus*

finmarchicus used in the experiment included all three dhCAs found in *Acartia* sp. as minor

components along with other copepodamides (Fig. 1A, Table S1).

3.2 Growth rate

261   Average growth rates were low and ranged from -0.10 to 0.05 day^{-1} (Fig. 2) but did not differ

262   between treatments (p = 0.22, Table 1) or the two species of *Dinophysis* (p = 0.94). However,

263   growth rate was positively correlated with total DSTs in both *D. sacculus* (p < 0.001) and *D.*

264   *acuminata* (p = 0.025) and accounted for 91% and 44% of the variance left unexplained by

265   treatment, respectively.

3.3 Toxins

Although total *Dinophysis*-produced toxins (DPTs) in *D. sacculus* generally increased (8 -

45%) with higher copepodamide concentrations and exposure to live copepods, only the

highest copepodamide treatment was significantly different from control conditions (p =

0.008, Fig. 3. Table 1). *D. sacculus* cells produced 45.1% (20.3 - 74.6) more DPTs in the 10

nM copepodamide treatment than in controls and 34% (12.2 - 50.4) more than in the live

copepod treatment (p = 0.027). *D. acuminata* okadaic acid (OA)content displayed a similar

trend, with 18.5 - 20.5% increases in all treatment groups compared to controls which,

however, were not significant (p = 0.35, Table 1). N.B. raw means and confidence intervals,

i.e. without correcting for growth rates, is available in Fig. S1.

276   Toxins were redistributed from the inside of cells to their surroundings in both

277   species (Fig. 4), and the effect was most pronounced for *D. acuminata* where the extracellular

278   fraction doubled from 46 - 54% to 90 - 94% (p < 0.001, Table 1). Intracellular toxins in the 5

279   and 10 nM copepodamide treatments were significantly lower than in all other treatments for

280   both species (Table. 1), ranging from 53 to 72% and 13 to 21% of control levels for *D.*

281   *sacculus* and *D. acuminata*, respectively.

282   PTX-2 accounted for almost all the increase in total DPTs produced by *D.*

sacculus (Fig. S2), but the proportions of individual toxin congeners (OA, PTX-2, C9) did

not significantly differ between any treatment groups (p-values: 0.062, 0.6, 0.16; Table 1).

On average, these congeners accounted for 4.7, 94, and 1.3% of total toxins, respectively

(Fig. 5). *D. acuminata* only produced OA.

3.4 Metabolomics

Metabolomic profiles of *Dinophysis* cells exposed to control conditions, live copepods, or 1

nM copepodamides formed clear clusters for both species (Fig. 6, Fig. S3). Meanwhile, the 5

and 10 nM treatments diverged from this cluster and, to a lesser extent, from each other. This

suggests a dose-dependent effect of copepodamides on the metabolomes of both *D. sacculus*

and *D. acuminata*. Indeed, 55 and 15 features (in +MS and -MS respectively) differed

significantly between treatments for *D. sacculus*, and 28 and 11 were significantly different

for *D. acuminata* (Supplementary spreadsheet 3).

295   Overall, two expression profiles were observed for the two *Dinophysis* species:

24/11 and 10/2 features (+MS/-MS) were co-upregulated with increasing copepodamide

concentration, while 30/4 and 18/9 features were co-downregulated in D. *sacculus* and D.

acuminata, respectively (Figs. S4-S7). One peculiar over-expressed feature was also noted in

the copepod treatment for *D. sacculus* only (M298T600, +MS). Proportionally, more features

were co-downregulated with copepodamide concentration for *D. acuminata* than for *D.*

sacculus.

4. Discussion

4.1 Toxin induction

We present the first experimental study that, to our knowledge, assesses the effects of copepod grazer cues (copepodamides) on *Dinophysis* toxin production and their metabolomes. Total *Dinophysis* produced toxins (DPTs) increased. slightly in *D. sacculus* cells that were exposed to grazing *Acartia* sp. copepods or copepodamides (Fig. 3) and displayed a positive trend as copepodamide concentration increased. However, this increase, ranging from 8 to 45%, was significantly different from non-induced cells only in the highest copepodamide treatment (10 nM). The effective copepodamide concentration in this treatment is one to two orders of magnitude higher than the average concentrations expected in natural conditions (Selander et al., 2019), but possibly on par with those experienced near copepods or in patches with high (>50 ind L^{-1}) copepod densities (Lindström et al., 2017). In addition, *D. acuminata* toxins did not differ significantly from non-exposed cells. It's therefore possible that the variation in cellular toxin content observed in *Dinophysis* cells in the field (Lindahl et al., 2007) are related to changes in strain composition as blooms develop, because variability in toxin content between strains of the same species or clade may be up to ten-fold (Séchet et al., 2021). In contrast to the present study, paralytic and amnesic shellfish toxins producers often respond to copepods and copepodamides by several fold increases in toxin production over similar time frames (Bergkvist et al., 2008; Lundholm et al., 2018; Griffin et al., 2019; Selander et al., 2019; Ryderheim et al., 2021). Hence, it seems unlikely that grazer-induced DPT production contributes significantly to the total of toxin content in natural populations of *Dinophysis* spp., at least based on the conditions and time frame covered here. It should, however, be noted that growth rate was low to absent in our experiment (Fig. 2), which may have constrained toxin production. Toxin production in *Alexandrium catenella* (previously *A. fundyense*) mainly takes place directly after division, in the G1 phase of the cell cycle (Taroncher-Oldenburg et al., 1997) and with limited growth

few cells would have passed the G1 phase during our exposure period. Regardless, the

modest induction of DPTs here raises question about the primary purpose of *Dinophysis*

toxins. Do they serve as a constitutive defence trait? Do they function as deterrents for

specific grazers not tested here? Are they used to immobilise prey? Or do they have a

different role altogether?

Grazer induced toxin production indicates that the toxin functions as a defence

against that grazer (Stahl, 1888), especially when the induction correlates to increase

resistance to grazers (Ryderheim et al., 2021; Olesen et al., 2022). Both inducible and

constitutional chemical defences are common in plant secondary metabolism (Stamp, 2003).

Inducible defences are favoured when there is a reliable (i.e., source specific and non-

persistent) cue associated with the level of threat, the threat level is variable, and the defence

metabolites are costly (Karban and Baldwin, 1997; Tollrian and Harvell, 1999).

Copepodamides are a reliable cue as they correlate to copepod density and rapidly degrade in

seawater (Selander et al., 2019). Daily vertical migration by both copepods and *Dinophysis*,

combined with differences in seasonal population dynamics of both predator and prey, indeed

results in a variable threat level across both time and space, which should favour the

evolution of inducible defences (Bergkvist et al., 2008; Ryderheim et al., 2021; Rigby et al.,

2022). However, the costs of producing DPTs are unknown. Unlike the nitrogen-rich PSTs

and AST alkaloids, DSTs and PTXs are cyclic polyethers and contain little to no nitrogen,

potentially making them less costly to produce in the often nitrogen-limited marine

ecosystem (Bristow et al., 2017). If metabolically cheap, there is less selective pressure on

Dinophysis to modulate toxin production according to grazing pressure and a high

constitutive chemical defence could instead be adaptive (Karban et al., 1997; Chen, 2008).

This is supported by the plant apparency hypothesis (Feeny, 1975, 1976; Rhoades and Cates,

1976), which predicts that large and/or more apparent (e.g., highly motile) organisms have a

higher degree of constitutive defence, unless this decreases their overall fitness (Harvell,

1990; Stamp, 2003). If DPTs function as chemical grazer deterrents it is possible that

Dinophysis could employ a combination of both strategies, permanently producing toxins as a

constitutive defence but able to increase this production when induced by increased grazing

risk. PTX-2 was the main contributor to the increase in total DPTs produced by *D. sacculus*

(Fig. S2) here. It could be that PTXs are used by *Dinophysis* as copepod specific deterrents,

whereas DSTs lack any such function. Despite likely being non-diarrhetic to humans and

differing from DSTs in both structure and effect, it remains a relevant inclusion as PTXs has

demonstrated toxicity to some invertebrates (Pease et al., 2022) and recently suggested to

reduce survival of the copepod *Acartia tonsa* (Ladds et al., 2024). Additionally, PTX-2 has

shown toxicity to several vertebrate (Terao et al., 1986; Fladmark et al., 1998) and human

cells (Jung et al., 1995), underscoring the potential ecological and health impacts of its

production and introduction to food webs.

It is also possible that the copepods used here do not represent the main grazers

of *Dinophysis* and that DPTs evolved to deter other grazers. Although many copepods co-

exist with *Dinophysis* and are known to feed on them in both field and experimental settings

(Carlsson et al., 1995; Frangoulis et al., 2022), the extent of this is unknown. Furthermore,

Acartia spp. and several other copepod species (e.g., *Euterpina acutifrons*) avoid feeding on

D. acuminata at bloom densities (Turner and Anderson, 1983; Maneiro et al., 2000).

Copepods are estimated to consume between 10 and 25% of pelagic primary production

(Calbet, 2001), whereas microzooplankton such as ciliates, heterotrophic flagellates. and

rotifers consume an estimated 67% (Calbet and Landry, 2004). Microzooplankton grazers

generally target prey as small or smaller than themselves (Hansen et al., 1994), and several

species of mixotrophic and heterotrophic dinoflagellates such as *Alexandrium ostenfeldii*

(Jacobson and Anderson, 1996), *Noctiluca scintillans* (Escalera et al., 2007), and *Fragilidium*

spp. (Jacobsen, 1999; Nézan and Chomérat, 2009) have been observed to feed on *Dinophysis*.

In fact, when offered a mixture of prey organisms (diatoms, a cryptophyte, a ciliate, a

raphidophyte, and dinoflagellates) *F. duplocampanaeforme* fed exclusively on *Dinophysis*

spp. (Park and Kim, 2010). Similarly, ciliates such as tintinnids (e.g., *Favella*) co-occur with

Dinophysis populations in the field (Santhanam and Srinivasap, 1996) and feed on

Dinophysis (Maneiro et al., 1998; Maneiro et al., 2000). Thus, microplankton could constitute

an important group of grazers for *Dinophysis*, and a possible target for chemical defences that

should be further explored.

4.2 Extracellular toxins – active release or passive leakage?

When exposed to high (5 – 10 nM) copepodamide concentrations, DPTs were redistributed

from intracellular compartments to the extracellular medium in both species, which increased

two to three-fold (Fig. 4). It is unknown if the redistribution was actively controlled or

resulted from increased leakage from injured or lysed cells. However, a notable fraction of

total *Dinophysis* toxins is commonly found in seawater (MacKenzie et al., 2004; Jørgensen

and Andersen, 2007; Johansen, 2008). Phycotoxins can serve to immobilise prey before

ingestion, which is experimentally supported for *Dinophysis* (Giménez Papiol et al., 2016;

Mafra Jr et al., 2016) and other microzooplankton (Skovgaard and Hansen, 2003; Sheng et

al., 2010; Blossom et al., 2012). It is, however, unclear whether DPTs serve this function for

Dinophysis or if other compounds are involved. Mafra Jr et al. (2016) reported that purified

solutions of OA, dinophysistoxin-1 (DTX1), and PTX-2 were ineffective against *Mesodinium*

rubrum prey. Cell-free culture medium imprinted by *Dinophysis* compounds, including free

polyunsaturated fatty acids such as eicosapentaenoic acid and docosahexaenoic acid,

decreased their motility and survival substantially. This suggests that *Dinophysis* likely use

some allelochemicals to aid in prey capture and may have actively released DPTs into the

extracellular medium for this purpose in our experiment. However, this does not account for

the consistent differential increase in extracellular DPTs in only the high copepodamide

treatments for both species and lacks sufficient experimental evidence to be conclusively

supported regardless.

Alternatively, DPTs released into the environment could act as a form of

chemical aposematism (Weldon, 2013; Rojas et al., 2017), increasing its concentration in the

phycosphere to discourage grazers capable of sensing them. It seems implausible, however,

given the high spatial dispersion of cells, even in moderate turbulence, for this kind of

allelopathic signal to confer an adaptive benefit to the individual cell (Jonsson et al., 2009).

Interestingly though, a recent experiment found that extracellular DPTs significantly

increased for *D. acuminata* cells when exposed to grazing *Acartia tonsa* copepods (Ladds et

al., 2024). Although this could simply be due to toxins leaking into the medium when cells

were fed on by the copepods, it tentatively supports our current findings and may suggest that

Dinophysis actively releases toxins into the extracellular environment as a grazer deterrent.

The redistribution of toxins could also be due to passive leaking from injured or

lysed cells. Cells exposed to high copepodamide concentrations grew more slowly than

control cells, although not significantly so, potentially indicating a toxic effect or a cost

associated with the copepodamide treatments on *Dinophysis*. However, all treatment groups

showed negligible growth, which suggests a more general cause, such as the physical

handling associated with the transfer of cultures to freshly coated culture vessels. *Dinophysis*

is indeed sensitive to handling and leak their cellular toxins when physically disturbed

(Jørgensen and Andersen, 2007; Johansen, 2008). This handling would, however, likely

affect the treatments similarly and does not sufficiently explain the differential increase in

only the two highest copepodamide groups for both species. Alternatively, the poor growth

and consequent lysis was caused by an increased allocation cost associated with DPT

production and/or other, unknown, grazer induced metabolites (Ryderheim et al., 2021;

Brown et al., 2022; Olesen et al., 2022; Park et al., 2023).

4.3 Metabolomics

We examined endometabolomic profiles for both *Dinophysis* species after exposure to

copepodamides or a living *Acartia* sp. copepod. Untargeted high resolution mass

spectrometry analyses revealed potential grazer deterrent candidates among features that co-

upregulated with increased copepodamide concentration. Alternatively, features that were

both co-upregulated and co-downregulated with copepodamide concentration may be

biomarkers indicating a stress-response or a potential growth cost associated with chemical

defences induced by the copepodamides. However, it remains unclear if, how, or to what

extent the expression pattern in the affected features is linked to growth rate. Overall, in all

but two annotated features choline was involved, either directly or in polar lipids.

Importantly, a series of five unsaturated lysoDGCC lipids and two lysoPC-like lipids, with

fatty acid chains from 16 to 28 carbons, were identified in *D. sacculus*.

442   Diacylglycerylcarboxyhydroxymethylcholine (DGCC) are a family of betaine

lipids that can be produced by dinoflagellates and may be their dominant representatives

(Flaim et al., 2014; Cañavate et al., 2016). Although the exact location and roles of these

DGCCs and their deacylated forms (i.e. lyso DGCCs) are unknown, they have been reported

as biomarkers of thermal tolerance, depending on their abundance or degrees of acylation and

unsaturation (Flaim et al., 2014; Rosset et al., 2019; Roach et al., 2021). Interestingly, the

only putatively annotated feature in *D. acuminata* was choline, which constitutes the polar

head of DGCC betaine lipids, but whether it is linked with DGCC or not remains unknown.

All the significantly affected features were upregulated with increasing copepodamide

concentrations but not affected by the live copepod, which is supported by the PCAs and

likely reflect a true physiological response (Fig. 6, Fig. S3). A more comprehensive analysis

of the total pool and nature of lysoPCs and lysoDGCCs in *Dinophysis* species could help

understanding their function and how they are affected by stressors. While these phospho-

and betaine lipids could mainly be localised in extra-chloroplastic membranes, they may be

primary acceptors of *de novo* formed fatty acid before being redistributed from the cytoplasm

to the chloroplast where they participate in membrane lipid synthesis (Eichenberger and

Gribi, 1997).

459   While extracellular fractions would have been interesting to analyse in this

context of chemically mediated induction of grazer deterrents, metabolomic analyses were

only performed on intracellular samples (i.e. endometabolomes) given the limited amount of

material, which is inherent to the highly challenging cultivation of *Dinophysis*. In addition,

this may have led to sensitivity issues (e.g. no detection of low intensity features) and could

have contributed to the low number of annotated features (e.g. poorly informative MS/MS

spectra). This is, however, common when trying to annotate marine microalgal metabolomes

(Zendong et al., 2016; Gémin et al., 2021; Yon et al., 2024). Regardless, knowledge about

grazer deterring metabolites in microalgae is generally poor, especially for effects induced by

chemical cues such as copepodamides, except for "known toxins" such as PSTs or ASTs.

4.4 Ecosystem implications for HAB formation

Harmful algal blooms (HABs) incur damages to human health and economic activities worth

billions of dollars annually (Hoagland and Scatasta, 2006; Kouakou and Poder, 2019).

Copepods are believed to contribute to HAB formation primarily by two mechanisms,

inducing upregulation of toxin production (Bergkvist et al., 2008; Selander et al., 2015;

Tammilehto et al., 2015) and/or by selectively feeding on non-toxic competitors (Selander et

al., 2006; Prevett et al., 2019; Ryderheim et al., 2021; Olesen et al., 2022). Dinoflagellates

grow slowly compared to e.g., many diatoms and are poor competitors for nutrients compared

with similarly sized organisms (Banse, 1982). Thus, harmful algae in general, and

dinoflagellates in particular, may be dependent on defensive traits (Fistarol et al., 2004;

Selander et al., 2019) and/or selective grazers (Irigoien et al., 2005; Selander et al., 2006;

Prevett et al., 2019). Indeed, when copepods are largely removed by the invasive ctenophore

Mnemiopsis leidyi, community structure changes in favour of faster growing diatoms

(Tiselius and Møller, 2017). Defended dinoflagellates are also positively correlated to

copepods in the ocean (Prevett et al., 2019), which is also true for DSTs in mussels on the

Swedish west coast (Trapp et al., 2021). If DSTs in mussels are indeed causally related to

copepods, our results suggest that it is not driven by grazer-induced production but rather by

selective feeding on less defended taxa, or other variables correlated to copepod density,

leading to increased relative abundance of *Dinophyis* spp (Carlsson et al., 1995; Xu and

Kiørboe, 2018). Future research should aim to determine the functional relationships between

Dinophysis and their predators, the mechanism(s) causing increased extracellular toxins, and

whether *Dinophysis* produced toxins result in any adaptive benefits for *Dinophysis* in

interactions with copepods and other grazers.

Fig. 1: Chromatogram of (**A**) the *Calanus finmarchicus* derived and purified copepodamide

mixture used in experiments and (**B**) copepodamides found in the *Acartia* sp. copepods used

as live grazers in this induction experiment. The trace amounts of 14:0 CA found in the

purified mixture were too low to be observed in this figure.

Fig. 2: Growth rate of *Dinophysis sacculus* (black) and *Dinophysis acuminata* (grey) after

68.5 hours of exposure to 1-10 nM concentrations of copepodamides, a living *Acartia* sp.

copepod, or control conditions without copepods or copepod cues. Transparent diamonds

represent individual growth rates. Circles denote mean growth rates (n = 3) and error bars

show their 95% confidence intervals.

Fig. 3: Total *Dinophysis* produced toxins, normalised by sample volume, for *Dinophysis*

sacculus (black) and *Dinophysis acuminata* (grey) after 68.5 hours of exposure to 1-10 nM

concentrations of copepodamides, a living *Acartia* sp. copepod, or control conditions without

copepods or copepod cues. Circles represent the covariate (growth rate) adjusted mean values

(n = 3), predicted at growth rate = 0, and error bars denote the 95% confidence intervals of

the predicted means. Transparent diamonds show the measured toxin content in each

replicate (i.e., not covariate adjusted). Letters denote statistically homogeneous subgroups for

D sacculus. There were no significantly different subgroups in *D acuminata.*

Fig. 4: Proportion of intracellular (grey) and extracellular (white) toxins for *Dinophysis*

sacculus (left) and *Dinophysis acuminata* (right) after 68.5 hours of exposure to 1-10 nM

concentrations of copepodamides, a living *Acartia* sp. copepod, or control conditions without

copepods or copepod cues. Bars represent the mean value of three replicates (n = 3) and error

bars show the 95% confidence intervals of the means. Letters denote statistically

homogeneous treatments. Note that only the lower confidence limits are shown for

extracellular bars.

Fig. 5: Composition of *Dinophysis* produced toxins (white = OA (Okadaic acid), light grey =

PTX-2 (Pectenotoxin-2) and dark grey = C9 (OA-C9-diol ester) for *Dinophysis sacculus* (*D.*

acuminata produced OA only) after 68.5 hours of exposure to 1-10 nM concentrations of

copepodamides, a living *Acartia* sp. copepod or control conditions without copepods or

copepod cues. Bars are the mean values of three replicates (n = 3) and error bars denote the

95% confidence intervals of the means. Note that only the lower confidence limits are shown

for C9.

Fig. 6: PCA score plot from LC-HRMS derived metabolomic profiles for *D. sacculus* (left,

+MS data) and *D. acuminata* (right, -MS data) after 68.5 hours of exposure to three

concentrations of chemical cues purified from copepods (copepodamides, triangles), a living

Acartia sp. copepod (circles), or control conditions (squares). Transparent symbols without

borders are individual replicate dimensionality reductions, and opaque symbols with black

border denote the centroids (averages) for each group.

534 Table 1. Results from linear models analysing growth rate (model I), total *Dinophysis*

535 produced toxins (DPTs; models II-IV), proportion of toxins inside vs. outside of cells (model

536 V), and individual toxin congener proportions of the total DPTs (models VI-VIII). Bold

537 values indicate statistically significant (p ≤ 0.05) model terms, IV = independent variable, Cv

538 = covariate.

Model type	Model number	Response variable	Term	SS	MS	df	F	p	Partial η^2
Two-way ANOVA	I	Growth rate	Treatment (IV)	0.082	0.0205	4	1.59	0.22	0.241
			Species (IV)	7.2E-05	7.2E-05	1	0.006	0.94	3E-04
			Treatment * Species	0.006	0.0015	4	0.11	0.98	0.022
			Residuals	0.26	0.0128	20			
Two-way ANCOVA	II	Total toxin content	Growth rate (Cv)	24738	24738	1	95.6	**< 0.001**	0.834
			Treatment (IV)	7218	1805	4	6.97	**0.00**	0.595
			Species (IV)	9446	9446	1	36.50	**< 0.001**	0.658
			Treatment * Species	1047	262	4	1.01	0.43	0.176
			Residuals	4918	259	19			
One-way ANCOVA	III	Total toxin content of *D. sacculus*	Growth rate (Cv)	23883.0	23883	1	89.3	**< 0.001**	0.91
			Treatment (IV)	6754	1689	4	6.3	**0.011**	0.74
			Residuals	2406	267	9			
One-way ANCOVA	IV	Total toxin content of *D. acuminata*	Growth rate (Cv)	1495	1495	1	7.19	**0.025**	0.444
			Treatment (IV)	1060	265	4	1.28	0.35	0.362
			Residuals	1871	208	9			
Two-way ANOVA	V	Intracellular toxin proportion	Treatment (IV)	29.9	7.48	4	92	**< 0.001**	0.949
			Species (IV)	37.4	37.45	1	461	**< 0.001**	0.958
			Treatment * Species	1.38	0.345	4	4.24	**0.012**	0.459
			Residuals	1.62	0.081	20			
One-way ANOVA	VI	OA proportion	Treatment (IV)	0.034	0.0085	4	3.195	0.062	0.561
				0.026	0.0026	10			
One-way ANOVA	VII	PTX-2 proportion	Treatment (IV)	0.01	0.0030	4	0.73	0.59	0.227
				0.04	0.0042	10			
One-way ANOVA	VIII	C9 proportion	Treatment (IV)	0.17	0.0413	4	0.855	0.162	0.255
				0.48	0.0484	10			

CRediT authorship contribution statement

Milad Pourdanandeh: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing – original draft, Writing – review & editing, Visualisation, Project administration

Véronique Séchet: Methodology, Resources, Investigation, Writing- review and editing

Liliane Carpentier: Methodology, Resources, Investigation

Damien Réveillon: Conceptualization, Validation, Formal analysis, Investigation, Data curation, Visualisation, Writing – original draft, Writing – review & editing

Fabienne Hervé: Formal analysis, Investigation, Methodology, Data curation, Writing- Review & Editing

Clarisse Hubert: Formal analysis, Investigation

Philipp Hess: Conceptualization, writing – review & editing, project administration

Erik Selander: Conceptualization, Methodology, Investigation, Formal analysis, Writing – original draft, Writing - review & editing, Visualisation, Project administration

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding sources

559   This work was funded by the Swedish Research Council grant to **E.S** (VR 2019-05238) and

560   the Ifremer/ODE/PHYTOX core budgets.

562   Data availability

563   Data and analysis code are available at Zenodo: https://doi.org/10.5281/zenodo.11091359

565   Acknowledgements

566   We thank two anonymous reviewers for their insightful comments and constructive

567   suggestions, which greatly improved the quality and clarity of this manuscript. We also thank

568   Jonathan N. Havenhand for valuable insight and discussions regarding the statistical analyses.

570   References

Arnoldt, S., Pourdanandeh, M., Spikkeland, I., Andersson, M.X., Selander, E., 2024. Mass

spectroscopy reveals compositional differences in copepodamides from limnic and marine

copepods. Scientific Reports 14(1), 3147. https://doi.org/10.1038/s41598-024-53247-1

Banse, K., 1982. Cell volumes, maximal growth rates of unicellular algae and ciliates, and the

role of ciliates in the marine pelagial. Limnology and Oceanography 27(6), 1059-1071.

https://doi.org/10.4319/lo.1982.27.6.1059

Bergkvist, J., Selander, E., Pavia, H., 2008. Induction of toxin production in dinoflagellates:

the grazer makes a difference. Oecologia 156, 147-154. https://doi.org/10.1007/s00442-008-

0981-6

Blossom, H.E., Daugbjerg, N., Hansen, P.J., 2012. Toxic mucus traps: A novel mechanism

that mediates prey uptake in the mixotrophic dinoflagellate *Alexandrium pseudogonyaulax*.

Harmful Algae 17, 40-53. https://doi.org/10.1016/j.hal.2012.02.010

Bristow, L.A., Mohr, W., Ahmerkamp, S., Kuypers, M.M.M., 2017. Nutrients that limit

growth in the ocean. Current Biology 27(11), R474-R478.

https://doi.org/10.1016/j.cub.2017.03.030

Broadhurst, D., Goodacre, R., Reinke, S.N., Kuligowski, J., Wilson, I.D., Lewis, M.R., Dunn,

W.B., 2018. Guidelines and considerations for the use of system suitability and quality

control samples in mass spectrometry assays applied in untargeted clinical metabolomic

studies. Metabolomics 14(6), 72. https://doi.org/10.1007/s11306-018-1367-3

Brown, E.R., Moore, S.G., Gaul, D.A., Kubanek, J., 2022. Predator cues target signaling

pathways in toxic algal metabolome. Limnology and Oceanography 67(6), 1227-1237.

https://doi.org/10.1002/lno.12069

Calbet, A., 2001. Mesozooplankton grazing effect on primary production: A global

comparative analysis in marine ecosystems. Limnology and Oceanography 46(7), 1824-1830.

https://doi.org/10.4319/lo.2001.46.7.1824

Calbet, A., Landry, M.R., 2004. Phytoplankton growth, microzooplankton grazing, and

carbon cycling in marine systems. Limnology and Oceanography 49(1), 51-57.

https://doi.org/10.4319/lo.2004.49.1.0051

Cañavate, J.P., Armada, I., Ríos, J.L., Hachero-Cruzado, I., 2016. Exploring occurrence and

molecular diversity of betaine lipids across taxonomy of marine microalgae. Phytochemistry

124, 68-78. https://doi.org/10.1016/j.phytochem.2016.02.007

Carlsson, P., Granéli, E., Finenko, G., Maestrini, S.Y., 1995. Copepod grazing on a

phytoplankton community containing the toxic dinoflagellate *Dinophysis acuminata*. Journal

of Plankton Research 17(10), 1925-1938. https://doi.org/10.1093/plankt/17.10.1925

Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto,

L., Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N.,

Frewen, B., Baker, T.A., Brusniak, M.-Y., Paulse, C., Creasy, D., Flashner, L., Kani, K.,

Moulding, C., Seymour, S.L., Nuwaysir, L.M., Lefebvre, B., Kuhlmann, F., Roark, J., Rainer,

P., Detlev, S., Hemenway, T., Huhmer, A., Langridge, J., Connolly, B., Chadick, T., Holly,

K., Eckels, J., Deutsch, E.W., Moritz, R.L., Katz, J.E., Agus, D.B., MacCoss, M., Tabb, D.L.,

Mallick, P., 2012. A cross-platform toolkit for mass spectrometry and proteomics. Nature

Biotechnology 30(10), 918-920. https://doi.org/10.1038/nbt.2377

Chen, M.S., 2008. Inducible direct plant defense against insect herbivores: a review. Insect

science 15(2), 101-114.

Dunn, W.B., Broadhurst, D., Begley, P., Zelena, E., Francis-McIntyre, S., Anderson, N.,

Brown, M., Knowles, J.D., Halsall, A., Haselden, J.N., Nicholls, A.W., Wilson, I.D., Kell,

D.B., Goodacre, R., The Human Serum Metabolome, C., 2011. Procedures for large-scale

metabolic profiling of serum and plasma using gas chromatography and liquid

chromatography coupled to mass spectrometry. Nature Protocols 6(7), 1060-1083.

https://doi.org/10.1038/nprot.2011.335

Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A.A., Melnik, A.V., Meusel, M.,

Dorrestein, P.C., Rousu, J., Böcker, S., 2019. SIRIUS 4: a rapid tool for turning tandem mass

spectra into metabolite structure information. Nature Methods 16(4), 299-302.

https://doi.org/10.1038/s41592-019-0344-8

Eichenberger, W., Gribi, C., 1997. Lipids of *Pavlova lutheri*: Cellular site and metabolic role

of DGCC. Phytochemistry 45(8), 1561-1567. https://doi.org/10.1016/S0031-9422(97)00201-

X

Escalera, L., Pazos, Y., Moroño, Á., Reguera, B., 2007. *Noctiluca scintillans* may act as a

vector of toxigenic microalgae. Harmful Algae 6(3), 317-320.

https://doi.org/10.1016/j.hal.2006.04.006

Feeny, P., 1975. Biochemical coevolution between plants and their insect herbivores,

Coevolution of Animals and Plants: Symposium V, First International Congress of

Systematic and Evolutionary Biology, 1973. University of Texas Press, pp. 1-19.

Feeny, P., 1976. Plant apparency and chemical defense, In: Wallace, J.W., Mansell, R.L.

(Eds.), Biochemical interaction between plants and insects. Springer US, Boston, MA, pp. 1-

40. https://doi.org/10.1007/978-1-4684-2646-5_1

Fistarol, G.O., Legrand, C., Selander, E., Hummert, C., Stolte, W., Granéli, E., 2004.

Allelopathy in *Alexandrium* spp.: effect on a natural plankton community and on algal

monocultures. Aquatic Microbial Ecology 35(1), 45-56. https://doi.org/10.3354/ame035045

Fladmark, K.E., Serres, M.H., Larsen, N.L., Yasumoto, T., Aune, T., DØskeland, S.O., 1998.

Sensitive detection of apoptogenic toxins in suspension cultures of rat and salmon

hepatocytes. Toxicon 36(8), 1101-1114. https://doi.org/10.1016/S0041-0101(98)00083-X

Flaim, G., Obertegger, U., Anesi, A., Guella, G., 2014. Temperature-induced changes in lipid

biomarkers and mycosporine-like amino acids in the psychrophilic dinoflagellate *Peridinium*

aciculiferum. Freshwater Biology 59(5), 985-997. https://doi.org/ 10.1111/fwb.12321

Frangoulis, C., Christou, E.D., Varkitzi, I., Zervoudaki, S., Maneiro, I., Svensen, C., Pagou,

K., Assimakopoulou, G., Hatzianestis, I., Granéli, E., 2022. Impact of a *Dinophysis*

acuminata bloom on the copepod *Acartia clausi*: First indications, Water.

Frost, B.W., 1972. Effects of size and concentration of food particles on the feeding behavior

of the marine planktonic copepod *Calanus Pacificus*. Limnology and Oceanography 17(6),

805-815. https://doi.org/10.4319/lo.1972.17.6.0805

Gaillard, S., Le Goïc, N., Malo, F., Boulais, M., Fabioux, C., Zaccagnini, L., Carpentier, L.,

Sibat, M., Réveillon, D., Séchet, V., Hess, P., Hégaret, H., 2020. Cultures of *Dinophysis*

sacculus, *D. acuminata* and pectenotoxin 2 affect gametes and fertilization success of the

Pacific oyster, *Crassostrea gigas*. Environmental Pollution 265, 114840.

https://doi.org/10.1016/j.envpol.2020.114840

Gaillard, S., Réveillon, D., Mason, P.L., Ayache, N., Sanderson, M., Smith, J.L., Giddings,

S., McCarron, P., Séchet, V., Hégaret, H., Hess, P., Vogelbein, W.K., 2023. Mortality and

histopathology in sheepshead minnow (*Cyprinodon variegatus*) larvae exposed to

pectenotoxin-2 and *Dinophysis acuminata*. Aquatic Toxicology 257, 106456.

https://doi.org/10.1016/j.aquatox.2023.106456

Gémin, M.-P., Bertrand, S., Séchet, V., Amzil, Z., Réveillon, D., 2021. Combined effects of

temperature and light intensity on growth, metabolome and ovatoxin content of a

mediterranean *Ostreopsis* cf. *ovata* strain. Harmful Algae 106, 102060.

https://doi.org/10.1016/j.hal.2021.102060

Georges des Aulnois, M., Réveillon, D., Robert, E., Caruana, A., Briand, E., Guljamow, A.,

Dittmann, E., Amzil, Z., Bormans, M., 2020. Salt shock responses of *Microcystis* revealed

through physiological, transcript, and metabolomic analyses, Toxins.

Georges des Aulnois, M., Roux, P., Caruana, A., Réveillon, D., Briand, E., Hervé, F., Savar,

V., Bormans, M., Amzil, Z., 2019. Physiological and metabolic responses of freshwater and

brackish-water strains of *Microcystis aeruginosa* acclimated to a salinity gradient: Insight

into salt tolerance. Applied and Environmental Microbiology 85(21), e01614-01619.

https://doi.org/10.1128/AEM.01614-19

Giménez Papiol, G., Beuzenberg, V., Selwood, A.I., MacKenzie, L., Packer, M.A., 2016. The

use of a mucus trap by *Dinophysis acuta* for the capture of *Mesodinium rubrum* prey under

culture conditions. Harmful Algae 58, 1-7. https://doi.org/10.1016/j.hal.2016.07.001

Grebner, W., Berglund, E.C., Berggren, F., Eklund, J., Harðadóttir, S., Andersson, M.X.,

Selander, E., 2019. Induction of defensive traits in marine plankton—new copepodamide

structures. Limnology and Oceanography 64(2), 820-831. https://doi.org/10.1002/lno.11077

Griffin, J.E., Park, G., Dam, H.G., 2019. Relative importance of nitrogen sources, algal alarm

cues and grazer exposure to toxin production of the marine dinoflagellate Alexandrium

catenella. Harmful Algae 84, 181-187. https://doi.org/10.1016/j.hal.2019.04.006

Guillard, R.R.L., Hargraves, P.E., 1993. *Stichochrysis immobilis* is a diatom, not a

chrysophyte. Phycologia 32(3), 234-236. https://doi.org/10.2216/i0031-8884-32-3-234.1

Guitton, Y., Tremblay-Franco, M., Le Corguillé, G., Martin, J.-F., Pétéra, M., Roger-Mele,

P., Delabrière, A., Goulitquer, S., Monsoor, M., Duperier, C., Canlet, C., Servien, R.,

Tardivel, P., Caron, C., Giacomoni, F., Thévenot, E.A., 2017. Create, run, share, publish, and

reference your LC–MS, FIA–MS, GC–MS, and NMR data analysis workflows with the

Workflow4Metabolomics 3.0 Galaxy online infrastructure for metabolomics. The

International Journal of Biochemistry & Cell Biology 93, 89-101.

https://doi.org/10.1016/j.biocel.2017.07.002

Hansen, B., Bjornsen, P.K., Hansen, P.J., 1994. The size ratio between planktonic predators

and their prey. Limnology and Oceanography 39(2), 395-403.

https://doi.org/10.4319/lo.1994.39.2.0395

Hansen, P.J., Nielsen, L.T., Johnson, M., Berge, T., Flynn, K.J., 2013. Acquired phototrophy

in *Mesodinium* and *Dinophysis* – A review of cellular organization, prey selectivity, nutrient

uptake and bioenergetics. Harmful Algae 28, 126-139.

https://doi.org/10.1016/j.hal.2013.06.004

Harvell, C.D., 1990. The ecology and evolution of inducible defenses. The Quarterly Review

of Biology 65(3), 323-340. https://doi.org/10.1086/416841

Hedges, L.V., Gurevitch, J., Curtis, P.S., 1999. The meta-analysis of response ratios in

experimental ecology. Ecology 80(4), 1150-1156. https://doi.org/10.1890/0012-

9658(1999)080[1150:TMAORR]2.0.CO;2

Hoagland, P., Scatasta, S., 2006. The economic effects of harmful algal blooms, In: Granéli,

E., Turner, J.T. (Eds.), Ecology of Harmful Algae. Springer Berlin Heidelberg, Berlin,

Heidelberg, pp. 391-402. https://doi.org/10.1007/978-3-540-32210-8_30

Irigoien, X., Flynn, K.J., Harris, R.P., 2005. Phytoplankton blooms: a 'loophole' in

microzooplankton grazing impact? Journal of Plankton Research 27(4), 313-321.

https://doi.org/10.1093/plankt/fbi011

Ishige, M., Satoh, N., Yasumoto, T., 1988. Pathological studies on the mice administrated

with the causative agent of diarrhetic shellfish poisoning (okadaic acid and pectenotoxin-2).

Bulletin of the Hokkaido Institute of Public Health 38, 15-18.

Jacobsen, D.M., 1999. A brief history of dinoflagellate feeding research. Journal of

Eukaryotic Microbiology 46(4), 376-381.

Jacobson, D.M., Anderson, D.M., 1996. Widespread phagocytosis of ciliates and other

protists by marine mixotrophic and heteotrophic dinoflagellates. Journal of Phycology 32(2),

279-285. https://doi.org/10.1111/j.0022-3646.1996.00279.x

Johansen, M., 2008. *PhD thesis.* On *Dinophysis* - Occurrence and toxin content, Department

of Marine Ecology. University of Gothenburg

Jonsson, P.R., Pavia, H., Toth, G., 2009. Formation of harmful algal blooms cannot be

explained by allelopathic interactions. Proceedings of the National Academy of Sciences

106(27), 11177-11182. https://doi.org/10.1073/pnas.0900964106

Jung, J.H., Sim, C.J., Lee, C.-O., 1995. Cytotoxic Compounds from a Two-Sponge

Association. Journal of Natural Products 58(11), 1722-1726.

https://doi.org/10.1021/np50125a012

Jørgensen, K., Andersen, P., 2007. Relation between the concentration of *Dinophysis*

acuminata and Diarrheic Shellfish Poisoning toxins in blue mussels (*Mytilus edulis*) during a

toxic episode in the Limfjord (Denmark), 2006. Journal of Shellfish Research 26(4), 1081-

1087. https://doi.org/10.2983/0730-8000(2007)26[1081:RBTCOD]2.0.CO;2

Karban, R., Agrawal, A.A., Mangel, M., 1997. The benefits of induced defenses against

herbivores. Ecology 78(5), 1351-1355. https://doi.org/10.2307/2266130

Karban, R., Baldwin, I.T., 1997. Induced responses to herbivory, 1 ed. University of Chicago

Press, Chicago, Illinois, USA.

Karlson, B., Andersen, P., Arneborg, L., Cembella, A., Eikrem, W., John, U., West, J.J.,

Klemm, K., Kobos, J., Lehtinen, S., Lundholm, N., Mazur-Marzec, H., Naustvoll, L.,

Poelman, M., Provoost, P., De Rijcke, M., Suikkanen, S., 2021. Harmful algal blooms and

their effects in coastal seas of Northern Europe. Harmful Algae 102, 101989.

https://doi.org/10.1016/j.hal.2021.101989

Kouakou, C.R.C., Poder, T.G., 2019. Economic impact of harmful algal blooms on human

health: a systematic review. Journal of Water and Health 17(4), 499-516.

https://doi.org/10.2166/wh.2019.064

Kuhl, C., Tautenhahn, R., Böttcher, C., Larson, T.R., Neumann, S., 2012. CAMERA: An

integrated strategy for compound spectra extraction and annotation of liquid

chromatography/mass spectrometry data sets. Analytical Chemistry 84(1), 283-289.

https://doi.org/10.1021/ac202450g

Ladds, M., Smith, J., Strohm, V., Gobler, C.J., 2024. Acartia tonsa grazing on the harmful

dinoflagellate Dinophysis acuminata reduces copepod survival and increases extracellular

toxin concentrations. Marine Ecology Progress Series 745, 25-40.

Lindahl, O., Lundve, B., Johansen, M., 2007. Toxicity of *Dinophysis* spp. in relation to

population density and environmental conditions on the Swedish west coast. Harmful Algae

6(2), 218-231. https://doi.org/10.1016/j.hal.2006.08.007

Lindström, J., Grebner, W., Rigby, K., Selander, E., 2017. Effects of predator lipids on

dinoflagellate defence mechanisms - increased bioluminescence capacity. Scientific Reports

7(1), 13104. https://doi.org/10.1038/s41598-017-13293-4

Lundholm, N., Krock, B., John, U., Skov, J., Cheng, J., Pančić, M., Wohlrab, S., Rigby, K.,

Nielsen, T.G., Selander, E., Harðardóttir, S., 2018. Induction of domoic acid production in

diatoms - Types of grazers and diatoms are important. Harmful Algae 79, 64-73.

https://doi.org/10.1016/j.hal.2018.06.005

MacKenzie, L., Beuzenberg, V., Holland, P., McNabb, P., Selwood, A., 2004. Solid phase

adsorption toxin tracking (SPATT): a new monitoring tool that simulates the biotoxin

contamination of filter feeding bivalves. Toxicon 44(8), 901-918.

https://doi.org/10.1016/j.toxicon.2004.08.020

Mafra Jr, L.L., Nagai, S., Uchida, H., Tavares, C.P.S., Escobar, B.P., Suzuki, T., 2016.

Harmful effects of *Dinophysis* to the ciliate *Mesodinium rubrum*: Implications for prey

capture. Harmful Algae 59, 82-90. https://doi.org/10.1016/j.hal.2016.09.009

Maneiro, I., D'Aleo, O., Guisande, C., Reguera, B., 1998. Interactions between the DSP agent

Dinophysis acuminata and the microzooplankton community, In: Reguera, B., Blanco, J.,

Fernández, M.L., Wyatt, T. (Eds.), Harmful Algae. Xunta de Galicia and Intergovernmental

Oceanographic Commission of UNESCO, pp. 386-389.

Maneiro, I., Frangópulos, M., Guisande, C., Fernández, M., Reguera, B., Riveiro, I., 2000.

Zooplankton as a potential vector of diarrhetic shellfish poisoning toxins through the food

web. Marine Ecology Progress Series 201, 155-163.

Matsushima, R., Uchida, H., Nagai, S., Watanabe, R., Kamio, M., Nagai, H., Kaneniwa, M.,

Suzuki, T., 2015. Assimilation, Accumulation, and Metabolism of Dinophysistoxins (DTXs)

and Pectenotoxins (PTXs) in the Several Tissues of Japanese Scallop Patinopecten

yessoensis. Toxins 7(12), 5141-5154. https://doi.org/10.3390/toxins7124870

Mitra, A., Flynn, K.J., Tillmann, U., Raven, J.A., Caron, D., Stoecker, D.K., Not, F., Hansen,

P.J., Hallegraeff, G., Sanders, R., Wilken, S., McManus, G., Johnson, M., Pitta, P., Våge, S.,

Berge, T., Calbet, A., Thingstad, F., Jeong, H.J., Burkholder, J., Glibert, P.M., Granéli, E.,

Lundgren, V., 2016. Defining planktonic protist functional groups on mechanisms for energy

and nutrient acquisition: Incorporation of diverse mixotrophic strategies. Protist 167(2), 106-

120. https://doi.org/10.1016/j.protis.2016.01.003

Nézan, E., Chomérat, N., 2009. *Fragilidium duplocampanaeforme* sp. nov. (Dinophyceae): A

new phagotrophic dinoflagellate from the French Atlantic coast. European Journal of

Protistology 45(1), 2-12. https://doi.org/10.1016/j.ejop.2008.04.002

Olesen, A.J., Ryderheim, F., Krock, B., Lundholm, N., Kiørboe, T., 2022. Costs and benefits

of predator-induced defence in a toxic diatom. Proceedings of the Royal Society B:

Biological Sciences 289(1972), 20212735. https://doi.org/10.1098/rspb.2021.2735

Pang, Z., Chong, J., Zhou, G., de Lima Morais, D.A., Chang, L., Barrette, M., Gauthier, C.,

Jacques, P.-É., Li, S., Xia, J., 2021. MetaboAnalyst 5.0: narrowing the gap between raw

spectra and functional insights. Nucleic Acids Research 49(W1), W388-W396.

https://doi.org/10.1093/nar/gkab382

Park, G., Norton, L., Avery, D., Dam, H.G., 2023. Grazers modify the dinoflagellate

relationship between toxin production and cell growth. Harmful Algae 126, 102439.

https://doi.org/10.1016/j.hal.2023.102439

Park, M.G., Kim, M., 2010. Prey specificity and feeding of the thecate mixotrophic

dinoflagellate *Fragilidium duplocampanaeforme*. Journal of Phycology 46(3), 424-432.

https://doi.org/10.1111/j.1529-8817.2010.00824.x

Park, M.G., Kim, S., Kim, H.S., Myung, G., Kang, Y.G., Yih, W., 2006. First successful

culture of the marine dinoflagellate *Dinophysis acuminata*. Aquatic Microbial Ecology 45(2),

101-106.

Pease, S.K.D., Brosnahan, M.L., Sanderson, M.P., Smith, J.L., 2022. Effects of Two Toxin-

Producing Harmful Algae, Alexandrium catenella and Dinophysis acuminata (Dinophyceae),

on Activity and Mortality of Larval Shellfish. Toxins 14(5).

https://doi.org/10.3390/toxins14050335

Posit team. RStudio: Integrated Development Environment for R v. 2024.4.1.748. *Posit*

Software, PBC, Boston, MA, USA. https://posit.co/. (2024).

Pourdanandeh, M., Séchet, V., Carpentier, L., Réveillon, D., Hervé, F., Hubert, C., Hess, P.,

Selander, E. Data and analysis code for a toxin induction study on two species of *Dinophysis*

dinoflagellates. 2024. *Zenodo*. https://doi.org/10.5281/zenodo.11091359.

Prevett, A., Lindström, J., Xu, J., Karlson, B., Selander, E., 2019. Grazer-induced

bioluminescence gives dinoflagellates a competitive edge. Current Biology 29(12), R564-

R565.

R Core Team. R: A Language and Environment for Statistical Computing v. 4.4.1. *R*

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/. (2024).

Reguera, B., Velo-Suárez, L., Raine, R., Park, M.G., 2012. Harmful *Dinophysis* species: A

review. Harmful Algae 14, 87-106. https://doi.org/10.1016/j.hal.2011.10.016

Rey, F., Melo, T., Lopes, D., Couto, D., Marques, F., Domingues, M.R., 2022. Applications

of lipidomics in marine organisms: progress, challenges and future perspectives. Molecular

Omics 18(5), 357-386. https://doi.org/10.1039/D2MO00012A

Rhoades, D.F., Cates, R.G., 1976. Toward a general theory of plant antiherbivore chemistry,

In: Wallace, J.W., Mansell, R.L. (Eds.), Biochemical Interaction Between Plants and Insects.

Springer US, Boston, MA, pp. 168-213. https://doi.org/10.1007/978-1-4684-2646-5_4

Rigby, K., Kinnby, A., Grønning, J., Ryderheim, F., Cervin, G., Berdan, E.L., Selander, E.,

2022. Species specific responses to grazer cues and acidification in phytoplankton - Winners

and losers in a changing world. Frontiers in Marine Science 9, 875858.

https://doi.org/10.3389/FMARS.2022.875858/BIBTEX

Roach, T.N.F., Dilworth, J., H, C.M., Jones, A.D., Quinn, R.A., Drury, C., 2021.

Metabolomic signatures of coral bleaching history. Nature Ecology & Evolution 5(4), 495-

503. https://doi.org/10.1038/s41559-020-01388-7

Rojas, B., Burdfield-Steel, E., Pakkanen, H., Suisto, K., Maczka, M., Schulz, S., Mappes, J.,

2017. How to fight multiple enemies: target-specific chemical defences in an aposematic

moth. Proceedings of the Royal Society B: Biological Sciences 284(1863), 20171424.

https://doi.org/10.1098/rspb.2017.1424

Rosset, S., Koster, G., Brandsma, J., Hunt, A.N., Postle, A.D., D'Angelo, C., 2019. Lipidome

analysis of Symbiodiniaceae reveals possible mechanisms of heat stress tolerance in reef

coral symbionts. Coral Reefs 38(6), 1241-1253. https://doi.org/10.1007/s00338-019-01865-x

Ryderheim, F., Selander, E., Kiørboe, T., 2021. Predator-induced defence in a dinoflagellate

generates benefits without direct costs. The ISME Journal 15(7), 2107-2116.

https://doi.org/10.1038/s41396-021-00908-y

Santhanam, R., Srinivasap, A., 1996. Impact of dinoflagellate *Dinophysis caudata* bloom on

the hydrography and fishery potentials of Tuticorin Bay, South India, In: Yasumoto, T.,

Oshima, Y., Fukuyo, Y. (Eds.), Harmful and Toxic Algal Blooms. Intergovernmental

Oceanographic Commission of UNESCO, Sendai, Japan, pp. 41-44.

Schymanski, E.L., Jeon, J., Gulde, R., Fenner, K., Ruff, M., Singer, H.P., Hollender, J., 2014.

Identifying small molecules via high resolution mass spectrometry: Communicating

confidence. Environmental Science & Technology 48(4), 2097-2098.

https://doi.org/10.1021/es5002105

Séchet, V., Sibat, M., Billien, G., Carpentier, L., Rovillon, G.-A., Raimbault, V., Malo, F.,

Gaillard, S., Perrière-Rumebe, M., Hess, P., 2021. Characterization of toxin-producing strains

of *Dinophysis* spp. (Dinophyceae) isolated from French coastal waters, with a particular focus

on the *D. acuminata*-complex. Harmful Algae 107, 101974.

Selander, E., Berglund, E.C., Engström, P., Berggren, F., Eklund, J., Harðardóttir, S.,

Lundholm, N., Grebner, W., Andersson, M.X., 2019. Copepods drive large-scale trait-

mediated effects in marine plankton. Science Advances 5(2).

https://doi.org/10.1126/sciadv.aat5096

Selander, E., Kubanek, J., Hamberg, M., Andersson, M.X., Cervin, G., Pavia, H., 2015.

Predator lipids induce paralytic shellfish toxins in bloom-forming algae. Proceedings of the

National Academy of Sciences 112(20), 6395-6400.

https://doi.org/10.1073/pnas.1420154112

Selander, E., Thor, P., Toth, G., Pavia, H., 2006. Copepods induce paralytic shellfish toxin

production in marine dinoflagellates. Proceedings of the Royal Society B: Biological

Sciences 273(1594), 1673-1680. https://doi.org/10.1098/rspb.2006.3502

Sheng, J., Malkiel, E., Katz, J., Adolf, J.E., Place, A.R., 2010. A dinoflagellate exploits toxins

to immobilize prey prior to ingestion. Proceedings of the National Academy of Sciences

107(5), 2082-2087. https://doi.org/10.1073/pnas.0912254107

Skovgaard, A., Hansen, P.J., 2003. Food uptake in the harmful alga *Prymnesium parvum*

mediated by excreted toxins. Limnology and Oceanography 48(3), 1161-1166.

https://doi.org/10.4319/lo.2003.48.3.1161

Stahl, E., 1888. Pflanzen und Schnecken: eine biologische Studie über die Schutzmittel der

Pflanzen gegen Schneckenfrass. Fischer. G.

Stamp, N., 2003. Out of the quagmire of plant defense hypotheses. The Quarterly Review of

Biology 78(1), 23-55. https://doi.org/10.1086/367580

Tammilehto, A., Nielsen, T.G., Krock, B., Møller, E.F., Lundholm, N., 2015. Induction of

domoic acid production in the toxic diatom *Pseudo-nitzschia seriata* by calanoid copepods.

Aquatic Toxicology 159, 52-61. https://doi.org/10.1016/j.aquatox.2014.11.026

Taroncher-Oldenburg, G., Kulis, D.M., Anderson, D.M., 1997. Toxin variability during the

cell cycle of the dinoflagellate *Alexandrium fundyense*. Limnology and Oceanography

42(5part2), 1178-1188. https://doi.org/10.4319/lo.1997.42.5

Terao, K., Ito, E., Yanagi, T., Yasumoto, T., 1986. Histopathological studies on experimental

marine toxin poisoning. I. Ultrastructural changes in the small intestine and liver of suckling

mice induced by dinophysistoxin-1 and pectenotoxin-1. Toxicon 24(11), 1141-1151.

https://doi.org/10.1016/0041-0101(86)90140-6

Tiselius, P., Møller, L.F., 2017. Community cascades in a marine pelagic food web controlled

by the non-visual apex predator *Mnemiopsis leidyi*. Journal of Plankton Research 39(2), 271-

279. https://doi.org/10.1093/plankt/fbw096

Tollrian, R., Harvell, C.D., 1999. The ecology and evolution of inducible defenses. Princeton

university press, Princeton, New Jersey, USA.

Trapp, A., Heuschele, J., Selander, E., 2021. Eavesdropping on plankton - Can zooplankton

monitoring improve forecasting of biotoxins from harmful algae blooms? Limnology and

Oceanography 66(9), 3455-3471. https://doi.org/10.1002/lno.11891

Turner, J.T., Anderson, D.M., 1983. Zooplankton grazing during dinoflagellate blooms in a

Cape Cod embayment, with observations of predation upon tintinnids by copepods. Marine

Ecology 4(4), 359-374.

van der Kloet, F.M., Bobeldijk, I., Verheij, E.R., Jellema, R.H., 2009. Analytical error

reduction using single point calibration for accurate and precise metabolomic phenotyping.

Journal of Proteome Research 8(11), 5132-5141. https://doi.org/10.1021/pr900499r

Wang, M., Carver, J.J., Phelan, V.V., Sanchez, L.M., Garg, N., Peng, Y., Nguyen, D.D.,

Watrous, J., Kapono, C.A., Luzzatto-Knaan, T., Porto, C., Bouslimani, A., Melnik, A.V.,

Meehan, M.J., Liu, W.-T., Crüsemann, M., Boudreau, P.D., Esquenazi, E., Sandoval-

Calderón, M., Kersten, R.D., Pace, L.A., Quinn, R.A., Duncan, K.R., Hsu, C.-C., Floros,

D.J., Gavilan, R.G., Kleigrewe, K., Northen, T., Dutton, R.J., Parrot, D., Carlson, E.E., Aigle,

B., Michelsen, C.F., Jelsbak, L., Sohlenkamp, C., Pevzner, P., Edlund, A., McLean, J., Piel,

J., Murphy, B.T., Gerwick, L., Liaw, C.-C., Yang, Y.-L., Humpf, H.-U., Maansson, M.,

Keyzers, R.A., Sims, A.C., Johnson, A.R., Sidebottom, A.M., Sedio, B.E., Klitgaard, A.,

Larson, C.B., Boya P, C.A., Torres-Mendoza, D., Gonzalez, D.J., Silva, D.B., Marques,

L.M., Demarque, D.P., Pociute, E., O'Neill, E.C., Briand, E., Helfrich, E.J.N., Granatosky,

E.A., Glukhov, E., Ryffel, F., Houson, H., Mohimani, H., Kharbush, J.J., Zeng, Y., Vorholt,

J.A., Kurita, K.L., Charusanti, P., McPhail, K.L., Nielsen, K.F., Vuong, L., Elfeki, M.,

Traxler, M.F., Engene, N., Koyama, N., Vining, O.B., Baric, R., Silva, R.R., Mascuch, S.J.,

Tomasi, S., Jenkins, S., Macherla, V., Hoffman, T., Agarwal, V., Williams, P.G., Dai, J.,

Neupane, R., Gurr, J., Rodríguez, A.M.C., Lamsa, A., Zhang, C., Dorrestein, K., Duggan,

B.M., Almaliti, J., Allard, P.-M., Phapale, P., Nothias, L.-F., Alexandrov, T., Litaudon, M.,

Wolfender, J.-L., Kyle, J.E., Metz, T.O., Peryea, T., Nguyen, D.-T., VanLeer, D., Shinn, P.,

Jadhav, A., Müller, R., Waters, K.M., Shi, W., Liu, X., Zhang, L., Knight, R., Jensen, P.R.,

Palsson, B.Ø., Pogliano, K., Linington, R.G., Gutiérrez, M., Lopes, N.P., Gerwick, W.H.,

Moore, B.S., Dorrestein, P.C., Bandeira, N., 2016. Sharing and community curation of mass

spectrometry data with Global Natural Products Social Molecular Networking. Nature

Biotechnology 34(8), 828-837. https://doi.org/10.1038/nbt.3597

Warton, D.I., Hui, F.K.C., 2011. The arcsine is asinine: the analysis of proportions in

ecology. Ecology 92(1), 3-10. https://doi.org/10.1890/10-0340.1

Weldon, P.J., 2013. Chemical aposematism. Chemoecology 23(4), 201-202.

https://doi.org/10.1007/s00049-013-0140-3

Xu, J., Kiørboe, T., 2018. Toxic dinoflagellates produce true grazer deterrents. Ecology

99(10), 2240-2249. https://doi.org/10.1002/ecy.2479

Xu, J., Nielsen, L.T., Kiørboe, T., 2018. Foraging response and acclimation of ambush

feeding and feeding-current feeding copepods to toxic dinoflagellates. Limnology and

Oceanography 63(4), 1449-1461. https://doi.org/10.1002/lno.10782

Yasumoto, T., Murata, M., Oshima, Y., Sano, M., Matsumoto, G.K., Clardy, J., 1985.

Diarrhetic shellfish toxins. Tetrahedron 41(6), 1019-1025. https://doi.org/10.1016/S0040-

4020(01)96469-5

Yasumoto, T., Oshima, Y., Sugawara, W., Fukuyo, Y., Oguri, H., Igarashi, T., Fujita, N.,

1980. Identification of *Dinophysis fortii* as the causative organism of Diarrhetic Shellfish

Poisoning. Nippon Suisan Gakkaishi 46(11), 1405-1411.

https://doi.org/10.2331/suisan.46.1405

Yon, T., Réveillon, D., Sibat, M., Holland, C., Litaker, R.W., Nascimento, S.M., Rossignoli,

A.E., Riobó, P., Hess, P., Bertrand, S., 2024. Targeted and non-targeted mass spectrometry to

explore the chemical diversity of the genus *Gambierdiscus* in the Atlantic Ocean.

Phytochemistry 222, 114095. https://doi.org/10.1016/j.phytochem.2024.114095

Zendong, Z., Bertrand, S., Herrenknecht, C., Abadie, E., Jauzein, C., Lemée, R., Gouriou, J.,

Amzil, Z., Hess, P., 2016. Passive sampling and high resolution mass spectrometry for

chemical profiling of French coastal areas with a focus on marine biotoxins. Environmental

Science & Technology 50(16), 8522-8529. https://doi.org/10.1021/acs.est.6b02081

944