Title: BCAS2 promotes primitive hematopoiesis by sequestering β-catenin within

the nucleus

Authors: Guozhu Ning[1,2#], Yu Lin[3#], Haixia Ma[4,5], Jiaqi Zhang[4,5], Liping Yang[1],

Zhengyu Liu[1], Lei Li[4,5]*, Xinyu He[1]*, Qiang Wang[1]*

Affiliations:

[1]Innovation Centre of Ministry of Education for Development and Diseases, the Sixth

Affiliated Hospital, School of Medicine, South China University of Technology,

Guangzhou 510006, China

[2]Affiliated Hospital of Guangdong Medical University & Key Laboratory of

Zebrafish Model for Development and Disease of Guangdong Medical University,

Zhanjiang 524001, China

[3]School of Pharmacy, Qiqihar Medical University, Qiqihar 161006, China

[4]Institute State Key Laboratory of Stem Cell and Reproductive Biology, Institute of

Stem Cell and Regeneration, Beijing Institute of Stem Cell and Regenerative

Medicine, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.

[5]University of Chinese Academy of Sciences, Beijing 100049, China.

[#]These authors contributed equally.

*To whom correspondence should be addressed: (Lei Li) lil@ioz.ac.cn; (Xinyu He)

hexyu@scut.edu.cn; (Qiang Wang) qiangwang@scut.edu.cn

Abstract

Breast carcinoma amplified sequence 2 (BCAS2), a core component of the hPrP19

complex, plays crucial roles in various physiological and pathological processes.

However, whether BCAS2 has functions other than being a key RNA-splicing

regulator within the nucleus remains unknown. Here, we show that BCAS2 is

essential for primitive hematopoiesis in zebrafish and mouse embryos. The activation

of Wnt/β-catenin signal, which is required for hematopoietic progenitor differentiation,

is significantly decreased upon depletion of *bcas2* in zebrafish embryos and mouse

embryonic fibroblasts (MEFs). Interestingly, BCAS2 deficiency has no obvious

impact on the splicing efficiency of β-catenin pre-mRNA, while significantly

attenuating β-catenin nuclear accumulation. Moreover, we find that BCAS2 directly

binds to β-catenin via its coiled-coil domains, thereby sequestering β-catenin within

the nucleus. Thus, our results uncover a previously unknown function of BCAS2 in

promoting Wnt signaling by enhancing β-catenin nuclear retention during primitive

hematopoiesis.

Introduction

Hematopoiesis refers to the lifelong process by which all blood cell lineages are generated. It begins at the early stage of embryonic development, providing the growing embryo with sufficient oxygen and nutrients.[1] Evolutionarily conserved across vertebrate species, hematopoiesis consists of two successive and partially overlapping waves: primitive and definitive. In mammals, the first wave of hematopoiesis occurs in the yolk-sac blood islands, producing primitive erythroid, megakaryocyte and macrophage progenitors, which can be observed in mouse embryos as early as embryonic day 7.25 (E7.25).[2-4] In zebrafish, primitive hematopoiesis initiates at around 11 hours post fertilization (hpf), when hemangioblasts emerge from the anterior lateral mesoderm (ALM) and posterior lateral mesoderm (PLM) and later differentiate into both hematopoietic and endothelial cells.[5-7]

Breast cancer amplified sequence 2 (BCAS2), also known as pre-mRNA splicing factor SPF27, is a 26-kDa nuclear protein containing two coiled-coil (CC) domains.[8] It was initially found to be overexpressed and amplified in human breast cancer cell lines.[9-11] Further studies have identified BCAS2 as a vital component of the human Prp19/CDC5L complex, which forms the catalytic ribonucleoprotein (RNP) core of spliceosome and is required for the activation of pre-mRNA splicing.[9,12,13] In Drosophila, the function of BCAS2 in RNA splicing is essential for cell viability.[14] In mouse, disruption of *Bcas2* in male germ cells impairs mRNA splicing and leads to a failure of spermatogenesis.[15] Additionally, BCAS2 has been shown to be a negative

regulator of p53 by directly interacting with p53 or modulating alternative splicing of

Mdm4, a major p53 inhibitor.[8,16] Zebrafish *bcas2* transcripts were enriched in the sites

of both primitive and definitive hematopoiesis during embryonic development.[16]

However, a previous study showed that p53 overactivation induced by zebrafish

bcas2 depletion did not affect primitive hematopoiesis, but impaired definitive

hematopoiesis.[16] In recent years, several studies have highlighted the importance of

regulating the expression and activity of p53 in primitive erythroid cell differentiation

in both mouse and zebrafish embryos.[17-19] Thus, it is necessary to reexamine the exact

function of BCAS2 in primitive hematopoiesis.

Wnt signaling, usually categorized into canonical and non-canonical pathways, is

involved in the process of hematopoiesis.[20-22] Notably, the canonical Wnt signaling

pathway, which is dependent on the nuclear accumulation of β-catenin to regulate

gene transcription, controls primitive hematopoietic progenitor formation and

promotes definitive hematopoietic stem cell (HSC) specification.[23-25] For instance, it

has been demonstrated in *Xenopus* that, Wnt4-mediated activation of Wnt/β-catenin

signaling plays a critical role in the induction and maintenance of primitive

hematopoiesis.[26] Moreover, transient inhibition of canonical Wnt signaling in

zebrafish embryos impairs embryonic blood formation.[27] However, previous studies

utilizing human pluripotent stem cells revealed an opposite role of Wnt/β-catenin

pathway in primitive progenitor generation.[25,28] Therefore, the impact of Wnt/β-

catenin signaling on primitive hematopoiesis remains elusive and even controversial.

Moreover, it has been suggested that BCAS2 is important for neural stem cell

proliferation and dendrite growth in mice by regulating β-catenin pre-mRNA

splicing.[29,30] As a nuclear protein, it is unclear whether BCAS2 can modulate Wnt/β-

catenin signaling in a splicing-independent manner.

In this study, we generated two zebrafish *bcas2* mutant lines, both of which

exhibited defects in male fertility and embryonic HSC formation, similar to what was

previously reported in mice and zebrafish.[15,16] More importantly, loss-of-function

experiments suggest that BCAS2 is necessary for primitive hematopoiesis in both

zebrafish and mouse embryos. We further find that *bcas2* is dispensable for the

survival and proliferation of hematopoietic cells, but plays a crucial role in the

differentiation of the hematopoietic lineage from hemangioblasts. Using a

comprehensive approach, we reveal that BCAS2 is a nuclear retention factor for β-

catenin during primitive hematopoiesis. Subsequent biochemical and functional

experiments demonstrate that BCAS2 directly binds to β-catenin and suppresses its

nuclear export to promote Wnt signal activation and hematopoietic progenitor

differentiation. Furthermore, the CC domains on BCAS2 and the Armadillo (ARM)

repeats on β-catenin are responsible for their interaction. Collectively, we have

uncovered a novel function of BCAS2 in regulating Wnt/β-catenin signaling by

sequestering β-catenin within the nucleus during primitive hematopoiesis.

Methods

Animal models

Our studies including animal maintenance and experiments were performed in

compliance with the guidelines of the Animal Care and Use Committee of the South

China University of Technology (Permission Number: 2023092). Seven strains of

zebrafish were used in this study, including Tübingen wild-type, *bcas2* mutant, *cloche*

mutant, *Tg(gata1:GFP)*, *Tg(coro1a:eGFP)*, *Tg(kdrl:GFP)*, and *Tg(hsp70l:dkk1b-*

GFP). *cloche* mutant, *Tg(gata1:GFP)* and *Tg(kdrl:GFP)* lines were provided by

Professor Feng Liu (Chinese Academy of Sciences). *Tg(coro1a:eGFP)* was provided

by Professor Yiyue Zhang (South China University of Technology). *Tg(hsp70l:dkk1b-*

GFP) strain was purchased from China Zebrafish Resource Center. *Bcas2^{Floxed/Floxed}*

(*Bcas2^{F/F}*) mouse line was generated as previously described.[15] Genotyping of

Bcas2^{F/F} mouse and *Flk1*-Cre mouse was performed using primers listed in

Supplemental Table 1. The mouse model with *Bcas2* specifically disrupted in the

hemangioblasts was derived from mating female *Bcas2^{F/F}* mice with *Flk1*-Cre

transgenic mice. All mouse lines were maintained on a mixed background

(129/C57BL/6).

Cell lines and transfection

HEK293T, HeLa, SW480, and L cell lines were obtained from ATCC. Conditional

Bcas2 knockout (*Bcas2*-cKO) mouse embryonic fibroblasts (MEFs) were prepared

from *Bcas2^{F/F}* embryos at E13.5. Cells were cultured in Dulbecco's modified eagle's

medium (DMEM, HyClone) supplemented with 10% fetal bovine serum (FBS,

HyClone) and 1% penicillin-streptomycin (HyClone) at 37 °C and 5% CO_2. L cell

lines expressing Wnt3a were maintained under similar conditions in the presence of

400 µg/ml G-418, from which Wnt3a conditioned medium (Wnt3a CM) was

generated. Culture medium prepared from L-cells was used as a control. To stimulate

Wnt signaling, cells were treated with Wnt3a CM in a 1:1 ratio with normal media. To

deplete *Bcas2* expression, *Bcas2*-cKO MEFs were cultured in medium containing 2

µM tamoxifen for 72 h and the knockout efficiency was evaluated using western blot

analysis. The same cells cultured without tamoxifen were used as a control. To silence

BCAS2 expression, shRNA constructs in pLL 3.7-GFP plasmid were generated to

target the following sequences: shRNA1, GAATGTGTAAACAATTCTA; shRNA2:

GAAGGAACTTCAGAAGTTA. Transfection was performed with Lipofectamine

2000 (11668019, Invitrogen) according to the manufacturer's instructions.

Generation of CRISPR-Cas9-mediated *bcas2* knockout zebrafish

The *bcas2* knockout zebrafish mutants were generated by CRISPR-Cas9 system as

previously described.[31] The guide RNA was designed to target the sequences 5′-

GGCGCAGCTGGAGCATCAGG-3′ within exon 4 of *bcas2*. Humanized Cas9

mRNA and gRNA were co-injected into wild-type embryos at the 1-cell stage.

Embryos or adult fin clips were collected to prepare genomic DNA. To screen for

mutant alleles, the genomic regions containing gRNA-targeted sequences were

amplified by polymerase chain reaction (PCR) with primers listed in Supplemental

Table 1. The PCR products were sequenced or digested with T7 endonuclease or

restriction enzyme FspI for genotyping.

RNA, morpholinos, and microinjection

Capped mRNAs for human *BCAS2*, *BCAS2 △CC1-2*, *BCAS2 CC1-2* and mouse *ΔN-β-catenin* mRNA were synthesized from the corresponding linearized plasmids using an mMESSAGE mMACHINE T7 transcription kit (AM1344, Ambion). Morpholinos (MOs) were designed and purchased from Gene Tools: mismatch MO (cMO 5'-AGCCACTCATCCTGCTCCTCCCATC-3'), and *bcas2* translation-blocking MO (tMO; 5'-AGCGACTGATGCTGGTCCTGCCATC-3'). The mRNAs and morpholinos were injected into embryos at the 1- to 2-cell stage.

Whole-mount *in situ* hybridization

Digoxigenin-labeled and fluorescein-labeled probes were synthesized using a RNA Labeling kit (11175025910, Roche). Whole-mount *in situ* hybridization (WISH) and double fluorescence *in situ* hybridization for zebrafish embryos were performed following previously published methods.[32,33] Anti-digoxigenin-POD (11633716001, Roche) and anti-fluorescein-POD (11426346910, Roche) were used to detect digoxigenin-labeled probes and fluorescein-labeled probes, respectively. After WISH, the stained embryos were embedded in OCT and sections were prepared with a LEICA CM1900. The mouse yolk sac layers were separated as previously described.[34]

***o*-Dianisidine staining**

To evaluate hemoglobin level, embryos were harvested at 36 hpf or 48 hpf, then stained with *o*-dianisidine as previously described.[35]

Proliferation and apoptosis assays

Embryos were incubated with 10 mM bromodeoxyuridine (BrdU) (B5002, Sigma) for

20 min. The incorporated BrdU was detected with anti-BrdU (B2531, Sigma)

antibody. TUNEL staining was performed using *In Situ* Cell Death Detection Kit,

TMR red (12156792910, Roche) according to the manufacturer's recommendation.

Heat shock treatment

To induce *dkk1* expression, *Tg(hsp70l:dkk1b-GFP)* embryos were subjected to heat

shock (42°C) for 10 min at 10 hpf, and then collected at the indicated stage for WISH.

Dual reporter assay

HEK293T cells or MEFs were seeded in 24-well plates and transfected with a Super-

TOPflash plasmid containing multimerized TCF-binding elements and a Renilla

luciferase plasmid, along with the indicated vectors. Then cells were treated with 100

ng/mL LiCl and/or Wnt3a CM for 12 h and assayed for luciferase activity using the

Dual luciferase system (E1910, Promega).

Immunoprecipitation, GST pulldown, and western blotting

For immunoprecipitation, HEK293T cells were transfected with the indicated

plasmids and collected 48 h after transfection. Subsequently, HEK293T cells were

lysed in a lysis buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 2 mM EDTA and

0.5% Nonidet P-40) containing protease inhibitors. Immunoprecipitation was

performed in accordance with the standard protocols.

For GST pulldown assay, GST, GST tagged β-catenin ARM 1-12 and His tagged

BCAS2 were expressed in *Escherichia coli* BL21, then purified using Glutathione-

Sepharose 4B beads (71024800-GE, GE Healthcare) and HisPur Ni-NTA beads

(88831, Thermo Fisher), respectively. GST and GST-β-catenin ARM 1-12 proteins

were immobilized onto Glutathione-Sepharose 4B beads and incubated with purified

His-BCAS2 at 4°C for 4 h. Beads were washed three times and analyzed using

western blotting.

Cytoplasmic and nuclear extracts were separated with nuclear and cytoplasmic

extraction kit (CW0199, CWBIO). Cell lysates were subjected to

immunoprecipitation with anti-Flag M2 affinity gel (A2220, Sigma) or anti-c-Myc

agarose affinity gel (A7470, Sigma) antibodies. Proteins were analyzed by western

blot using anti-Flag (F2555, Sigma), anti-HA (CW0092A, CWBIO), anti-β-catenin

(M24002, Abmart), anti-BCAS2 (10414-1-AP, Proteintech), anti- β -Tubulin

(CW0265A, CWBIO), anti-GFP (A11120, Thermo Fisher), anti-Histone H3 (ab1791,

Abcam), anti-GST (SAB4200237, Sigma), and anti-His Tag (AF5060, Beyotime)

antibodies.

Immunofluorescence staining

Cells on coverslips and embryos were processed for immunofluorescence staining as

previously described.[36,37] Before fixation, *bcas2*-deficient MEFs were treated with a

concentration of 20 μM MG132 or 20 nM leptomycin B (LMB) for 6 h, while

Tg(gata1:GFP) embryos were treated with 20 nM LMB from the bud stage to the 10-

somite stage. The prepared samples were stained with anti-BCAS2 (10414-1-AP,

Proteintech), anti-β-catenin (M24002, Abmart), and anti-GFP (A11122, Invitrogen).

Meanwhile, 4′,6-Diamidine-2′-phenylindole dihydrochloride (DAPI, 10236276001,

Sigma) was used to label nuclei. Fluorescence imaging was performed using a Nikon

A1R+ confocal microscope and all images were captured with the same settings. The

relative fluorescence intensity was calculated by dividing the fluorescence intensity of

nuclear β-catenin by the fluorescence intensity of DAPI.

Bimolecular fluorescence complementation assay

To construct the plasmids for Bimolecular fluorescence complementation assay

(BiFC), BCAS2 was fused to the N-terminal half of yellow fluorescent protein (YN-

BCAS2) and β-catenin to the C-terminal half (YC-β-catenin). YN-BCAS2 and YC-β-

catenin were either individually or collectively transfected into HeLa cells.

Fluorescence was detected 48 h after transfection using a Nikon A1R+confocal

microscope.

Fluorescence recovery after photobleaching

BCAS2 and GFP tagged S37A-β-catenin were co-transfected into HeLa cells.

Fluorescence recovery after photobleaching (FRAP) assay was performed according

to previously reported methods.[38] The cells were bleached by the 488 nm laser line of

the 20 mW argon laser at 100% power. About 90% of nuclear or cytoplasmic GFP

signal was bleached. Images were acquired with 35 frames at 25-s intervals by a

META Zeiss 510 confocal microscope.

RNA sequencing

Embryos were collected at the 10-somite stage and gently transferred into lysis buffer.

Reverse transcription was performed using a SMARTer Ultra Low RNA Kit

(Clontech, 634437) directly from the cell lysates. The cDNA library was prepared

using an Advantage 2 PCR Kit (Clontech，639206) and then sequenced via the

Illumina sequencing platform (NovaSeq 6000). The difference in the number of

alternative splicing events between groups was analyzed using rMATS (version 4.1.0).

Reverse transcription PCR

Total RNA was isolated from wild-type and *bcas2* mutant embryos at the 10-somite

stage with MicroElute Total RNA kit (OMEGA, R6831-01), followed by reverse

transcription using ReverTra Ace qPCR RT Kit (Toyobo, FsQ-101). The cDNA was

amplified with primers listed in Supplemental Table 2.

Quantification and statistical analysis

Images were quantified with Image J. Statistical data were analyzed using Graph Pad

Prism. Comparisons between experimental groups were done using the Student's *t-test*.

Data are presented as mean ± standard deviation. *P < 0.05, P < 0.01, P < 0.001*, and *P*

< 0.0001 were considered statistically significant and marked with *, **, ***, and

****, respectively (Student's *t-test*).

Results

BCAS2 is necessary for primitive hematopoiesis

To confirm that *bcas2* is expressed in the posterior intermediate cell mass (ICM)

where primitive hematopoiesis occurs in zebrafish, we first examined the

spatiotemporal expression pattern of *bcas2* during zebrafish embryo development by

performing whole mount *in situ* hybridization (WISH). The results showed that *bcas2*

was ubiquitously expressed from 1-cell stage to 10-somite stage (Supplemental Figure

1A). Its expression in the ICM became detectable at 18 hours post fertilization (hpf)

and was significantly elevated at 22 hpf (Figure 1A). We further observed that *bcas2*

was co-expressed with the primitive erythropoietic marker *gata1* in the ICM at 22 hpf

by fluorescence *in situ* hybridization (FISH) (Figure 1B). In contrast, *bcas2* was

hardly detectable in the ICM in *cloche*[-/-] mutants that lack both endothelial and

hematopoietic cells (Figure 1C). These results demonstrate a dynamic expression of

bcas2 in the ICM and imply a potential role of this gene in primitive hematopoiesis.

To gain insight into the developmental function of *bcas2*, we employed

CRISPR/Cas9 system to generate *bcas2* mutants. Two mutant lines were obtained,

designated *bcas2*[Δ7] (with a 7-base deletion) and *bcas2*[Δ14] (with a 14-base deletion).

These mutations led to premature translation termination, which resulted in truncated

Bcas2 proteins lacking the C-terminal CC domains (Supplemental Figure 2A-B).

bcas2$^{+/Δ7}$ and *bcas2$^{+/Δ14}$* mutants were identified by T7 endonuclease I assay or

restriction enzyme analysis (FspI) (Supplemental Figure 2C). We found that nearly

85% of the embryos derived from crossing *bcas*2$^{+/-}$ males with *bcas*2$^{+/-}$ females did

not develop to the cleavage stage (Supplemental Figure 2D). Only 3% of the living

embryos were homozygotes. In contrast, embryos obtained by crossing between wild-

type males and *bcas*2$^{+/-}$ females were viable and showed normal morphology, with a

heterozygosity rate consistent with Mendelian inheritance. This could be explained by

male infertility as previously documented in *Bcas2* knockout mice.[15] Combining the

above findings, we propose that Bcas2 may have an evolutionarily conserved role in

spermatogenesis.

Given the difficulty of obtaining homozygous mutants, embryos lacking one copy

of *bcas2* gene were produced from crosses between heterozygous females and wild-

type males. We observed a significant decrease of *bcas2* expression in the ICM region

in *bcas2$^{+/Δ7}$* or *bcas2$^{+/Δ14}$* mutants, likely resulting from nonsense-mediated RNA

decay (Figure 1D). In line with a previous report,[16] a marked reduction in the

expression of the HSC marker *cmyb* and T-cell marker *rag1* was found in *bcas2$^{+/Δ7}$* or

bcas2$^{+/Δ14}$ embryos at 5 dpf, indicating an essential role of *bcas2* in definitive

hematopoiesis (Supplemental Figure 3A-B). These findings suggest that *bcas2$^{+/Δ7}$* and

bcas2$^{+/Δ14}$ mutants can be used to examine the involvement of *bcas2* in primitive

hematopoiesis.

To explore whether *bcas2* is required for primitive hematopoiesis, we first

examined the expression of primitive erythropoietic markers *gata1* and *hbbe3* in

bcas2$^{+/\Delta7}$ and *bcas2*$^{+/\Delta14}$ embryos at 22 hpf, and observed a marked decrease in the

expression of these genes in the mutants (Figure 1E-F). Surprisingly, *o*-dianisidine

staining showed similar hemoglobin contents in the *bcas2*$^{+/\Delta7}$ and *bcas2*$^{+/\Delta14}$ embryos

at 48 hpf compared with control embryos, suggesting that the defect in primitive

hematopoiesis induced by haploinsufficiency of *bcas2* was alleviated at later

developmental stages. In order to further explore the role of BCAS2 in primitive

hematopoiesis, we identified several *bcas2*$^{\Delta14/\Delta14}$ mutants from about 100 embryos.

These homozygous mutants display a severe decrease in hemoglobin (Figure 1G).

Moreover, injection of a translation-blocking MO into wild-type embryos to

downregulate *bcas2* expression resulted in severe defects in erythropoiesis at 22 hpf

and 48 hpf (Supplemental Figure 4A-D). These results indicate that *bcas2* is

indispensable for primitive hematopoiesis in zebrafish. In addition, transgenic mice

expressing Cre recombinase under the control of the *Flk1* promoter were crossed to

Bcas2$^{F/F}$ animals to induce the deletion of *Bcas2* in endothelial/hematopoietic cells.

We found that red blood cells were eliminated in the yolk sac of *Bcas2*$^{F/F}$;*Flk1-Cre*

mice at E12.5 despite the presence of vessels (Figure 1H). Therefore, Bcas2 has a

conserved role in vertebrates to regulate primitive hematopoiesis.

bcas2 deficiency impairs hematopoietic progenitor differentiation

The decrease of primitive hematopoietic cells in *bcas2* deficient animals may be

attributed to a number of possible causes: excessive apoptosis, hampered proliferation

of hematopoietic cells, or impaired differentiation of hematopoietic progenitor cells.

To shed light on this issue, we first performed terminal deoxynucleotidyl transferase-

mediated dUTP nick end labeling (TUNEL) assay in *Tg(gata1:GFP)* embryos at the

10-somite stage to examine DNA fragmentations in apoptotic cells, and found no

obvious apoptotic signal in the *gata1*[+] hematopoietic cells in either *bcas2*[+/Δ14]

embryos or their wild-type siblings (Supplemental Figure 5A). Meanwhile, BrdU

incorporation assay revealed no significant difference in hematopoietic cell

proliferation between *bcas2*[+/Δ14] mutants and their corresponding wild-types

(Supplemental Figure 5B-C). These data suggest that *bcas2* is dispensable for the

survival and proliferation of hematopoietic cells.

In the developing embryo, hemangioblasts are derived from the ventral mesoderm

at early somite stage, and then differentiate into both hematopoietic and endothelial

lineages.[39,40] To test whether *bcas2* functions in cell fate decision during primitive

hematopoietic cell development, the expression of hemangioblast markers *npas4l*, *scl*,

and *gata2* in *bcas2*[+/Δ14] embryos was examined at the 1- to 2-somite stage. As shown

in Figure 2A, haploinsufficiency of *bcas2* did not affect the emergence of the

hemangioblast population. Then we extended our analysis to include the markers of

hematopoietic and endothelial progenitors. Consistent with the decrease in primitive

hematopoietic cells in *bcas2* deficient mutants, a marked reduced expression of

erythrocyte progenitor markers *gata1* and *hbbe3* was observed in the posterior lateral

mesoderm of *bcas2*[+/-] embryos at the 10-somite stage (Figure 2B). Interestingly, the

expression of myeloid progenitor marker *pu.1* was also dramatically decreased

(Figure 2C). Moreover, overexpression of human BCAS2 enhanced the expression of

gata1 in both wild-type and mutant embryos at the 10-somite stage (Figure 2D). In

contrast, the endothelial progenitor marker *fli1a* was expressed at a similar level in

bcas2[+/-] embryos as in wild-type animals (Figure 2E). Consistently, blood vessels

appeared normal in *bcas2*[+/-] mutants with *Tg(kdrl:GFP)* background at 54 hpf (Figure

2F). These data provide convincing evidence that *bcas2* is required for the

differentiation of the hematopoietic lineage from hemangioblasts during primitive

hematopoiesis.

BCAS2 functions in primitive hematopoiesis by activating Wnt signaling

Previous studies have shown that Wnt/β-catenin plays a key role in primitive

hematopoiesis.[26,27,41] As both BCAS2 and β-catenin-like 1 (CTNNBL1) are members

of the Prp19/CDC5L complex, which is a major building block of the spliceosome's

catalytic RNP core,[13] we speculate that BCAS2 may be a regulator of Wnt signaling

through interaction with β-catenin during hematopoiesis. To test our hypothesis,

human BCAS2 was overexpressed in HEK293T cells and MEFs. Ectopic expression

of BCAS2 enhanced the Wnt3a-induced expression of the TOPflash luciferase

reporter in a dose-dependent manner (Figure 3A-B). Importantly, Wnt3a-induced

luciferase activity in HEK293T cells could be effectively reduced by knockdown of

BCAS2 using two shRNAs targeting different regions of human *BCAS2* (Figure 3C

and Supplemental Figure 6). Similar results were also observed in *Bcas2*-cKO MEFs

in the presence of tamoxifen (Figure 3D). Furthermore, the expression of *cdx4* and

hoxa9a, which are targets of canonical Wnt signaling in the regulation of

hematopoietic development,[42,43] were downregulated in the lateral plate mesoderm of

bcas2[+/-] embryos at the 6-somite stage (Supplemental Figure 7). These findings

support that BCAS2 promotes Wnt signaling activation.

To confirm that Wnt signaling was required for zebrafish embryonic hematopoiesis,

we induced the expression of canonical Wnt inhibitor Dkk1 by heat-shocking

Tg(hsp70l:dkk1-GFP)[w32] embryos at the bud stage.[44] As expected, diminished

expression of *gata1* and *hbbe3* was detected in the resulting embryos at the 10-somite

stage (Figure 3E-F). In addition, treatment with a small molecule β-catenin antagonist

CCT036477 from 9 hpf did not affect the expression of hemangioblast markers

npas4l,[45] *scl*, and *gata2* or endothelial marker *fli1a* (Supplemental Figure 8A-B), yet

significantly reduced the expression of erythroid progenitor marker *gata1* in wild-type

embryos (Supplemental Figure 8C), suggesting that canonical Wnt signaling may not

be required for the generation of hemangioblasts or their endothelial differentiation,

but is pivotal for their hematopoietic differentiation. To further validate that *bcas2*

functions in primitive hematopoiesis via Wnt/β-catenin signaling, the expression

pattern of β-catenin was examined in *bcas2* morphants with *Tg(gata1:GFP)*

background at the 10-somite stage by immunofluorescent staining. The signal of

nuclear β-catenin was substantially decreased in hematopoietic progenitor cells

(Figure 3G-H) and primitive myeloid cells (Supplemental Figure 9A-B). Moreover,

overexpression of ΔN-β-catenin, a constitutively active form of β-catenin, effectively

restored the expression of *hbbe3* in *bcas2* morphants and mutants (Figure 3I-J). All

these data suggest that BCAS2 functions in primitive hematopoiesis by regulating

Wnt/β-catenin signaling.

BCAS2 promotes β-catenin nuclear accumulation independently of protein

stability regulation

To investigate how BCAS2 regulates Wnt/β-catenin signaling, HEK293T cells were

treated with LiCl, a canonical Wnt agonist that inhibits GSK-3β activity and stabilizes

cytosolic β-catenin.[46] The results showed that TOPflash activity was significantly

elevated in LiCl-treated cells (Figure 4A). BCAS2 overexpression further upregulated,

whereas shRNA-mediated knockdown of *BCAS2* downregulated LiCl-induced

TOPflash activity (Figure 4A-B). Likewise, HEK293T cells transfected with S37A-β-

catenin, a constitutively active form of β-catenin that is resistant to GSK-3β-mediated

degradation, [47] displayed a much higher level of TOPflash activity, which was reduced

by *BCAS2* knockdown (Figure 4C). These results strongly imply that BCAS2

regulates Wnt signaling downstream of β-catenin stability control.

To test the above hypothesis, we evaluated nuclear β-catenin level by performing

immunofluorescence staining and immunoblotting experiments. Upon tamoxifen

exposure, nuclear accumulation of β-catenin induced by LiCl was greatly inhibited in

Bcas2-cKO MEFs, while nuclear/cytoplasmic fractionation suggested that

cytoplasmic β-catenin level remained relatively unchanged (Figure 4D-E). Similarly,

silencing *BCAS2* with shRNA led to reduced nuclear β-catenin in the human colon

cancer cell line SW480, in which β-catenin was activated because of mutations in the

adenomatous polyposis coli protein (APC), an integral component of the β-catenin

destruction complex (Figure 4F).[48] Next, MG132, a proteasome inhibitor, was applied

to activate Wnt/β-catenin signaling in *Bcas2*-cKO MEFs by inhibiting β-catenin

degradation. In the absence of tamoxifen and MG132, endogenous β-catenin was

localized almost exclusively in the cytoplasm; MG132 treatment dramatically

triggered β-catenin accumulation in the nuclei (Figure 4G). However, in *Bcas2*-cKO

MEFs exposed to tamoxifen, MG132 treatment was not able to induce nuclear

accumulation of β-catenin (Figure 4G). These findings indicate that BCAS2 promotes

β-catenin nuclear accumulation in a manner that is independent of β-catenin stability

regulation.

BCAS2 sequesters β-catenin within the nucleus

In addition to be affected by protein stability, the nuclear level of β-catenin is also

fine-tuned by the opposing actions of nuclear export and import.[49-51] To examine the

effect of BCAS2 on the nuclear import and export of β-catenin, fluorescent recovery

after photobleaching (FRAP) experiments was carried out in HeLa cells expressing

GFP-tagged S37A-β-catenin. After photobleaching the nucleus, no significant

difference was found in the recovery of nuclear GFP signals between the cells with

and without overexpression of BCAS2 (Supplemental Figures 10A, A', and C),

suggesting that BCAS2 does not regulate β-catenin nuclear import. Conversely, after

photobleaching the cytoplasm, BCAS2 overexpressed cells showed a much slower

recovery of cytoplasmic fluorescence (Supplemental Figures 10B, B', and C),

indicating that BCAS2 inhibits β-catenin nuclear export.

It has been suggested that the nuclear exit of β-catenin can be either dependent or

independent on CRM1, a major nuclear export receptor.[52] To shed light on the

mechanism underlying BCAS2 mediated β-catenin nuclear retention, we treated

Bcas2-cKO MEFs with the CRM1-specific export inhibitor LMB.[53] Regardless of the

presence or absence of endogenous BCAS2, LMB treatment could effectively

increase the level of β-catenin in the nucleus (Figure 5A). Consistently, treatment of

LMB was able to rescue the impaired nuclear accumulation of β-catenin in BCAS2-

deficient SW480 cells (Figure 5B). Moreover, when *bcas2* morphants in *Tg*

(gata1:GFP) background were treated with LMB from bud stage to 10 somite stage,

the level of nuclear β-Catenin was partially recovered (Figure 5C-D). Importantly, the

expression of *gata1* was also restored in *bcas2* mutants upon LMB treatment (Figure

5E). We further tested if BCAS2 specifically regulates CRM1-mediated nuclear

export of β-catenin by analyzing the nucleocytoplasmic distribution of other known

CRM1 cargoes, such as ATG3 and CDC37L.[54] Intriguingly, BCAS2 overexpression in

HeLa cells only slightly enhanced the nuclear localization of CDC37L, and had no

significant impact on that of ATG3 (Supplemental Figure 11), indicating the

specificity of BCAS2-mediated inhibition of CRM1-dependent nuclear export of β-

catenin. Taken together, these findings suggest that BCAS2 negatively regulates

CRM1-mediated nuclear export of β-catenin.

BCAS2 directly interacts with β-catenin in the nucleus

To investigate whether BCAS2 inhibits the nuclear export of β-catenin through

physical binding, HEK293T cells were transfected with Flag-tagged β-catenin and

HA-tagged BCAS2 constructs. Co-immunoprecipitation (Co-IP) experiments showed

that Flag-β-catenin was precipitated with HA-BCAS2 as well as endogenous BCAS2,

indicating an interaction between these two proteins (Figure 6A-B). In addition, the

interaction was enhanced upon Wnt3a stimulation (Figure 6C). Given that Wnt ligand

stimulation ultimately induces β-catenin nuclear accumulation, this enhanced

interaction implies that BCAS2 associates with β-catenin within the nucleus.

Therefore, we performed the bimolecular fluorescence complementation (BiFC) assay

to visualize the interaction of BCAS2 and β-catenin in living cells. In this assay, the

N-terminal fragment of yellow fluorescent protein (YFP) was fused to BCAS2 (YN-

BCAS2), while the C-terminal fragment was fused to β-catenin (YC-β-catenin)

(Figure 6D). As expected, the YFP fluorescence was specifically observed in the

nucleus (Figure 6E).

Previous studies have divided the β-catenin protein into three distinct domains,

including the N-terminal domain (residues 1–133), the central domain with 12 ARM

repeats (residues 134–670), and the C-terminal domain (residues 671–781).[55] To

identify the BCAS2 binding site, constructs expressing various truncated forms of β-

catenin were generated and co-transfected with BCAS2 into HEK293T cells (Figure

6F). Co-IP assays revealed that deletion of the N-terminal or C-terminal domain of β-

catenin did not alter the interaction between β-catenin and BCAS2 (Figure 6G). In

contrast, when the ARM repeats 1-12 of β-catenin were deleted, the resulting deletion

mutant showed virtually no interaction with BCAS2 (Figure 6G). GST pull-down

assay also demonstrated a direct interaction between BCAS2 and the ARM repeats of

β-catenin (Figure 6H). These results indicate that BCAS2 physically binds to the

ARM repeats of β-catenin. Furthermore, we found that the ARM repeats 9-12, but not

1-8, bound to BCAS2 (Figure 6G).

Transcriptional activation of the canonical Wnt target genes depends on β-catenin

nuclear localization and its physical association with TCF/LEF family members. As

the binding sites for TCF have been located in the ARM repeats 3-10 of β-catenin,[56] it

is likely that BCAS2-mediated nuclear sequestration of β-catenin through interacting

with the ARM repeats 9-12 would be compatible with the initiation of gene

transcription by allowing for the association of β-catenin and TCF. To validate this

possibility, co-IP assays were performed and we found that β-catenin still bound with

TCF4 in the presence of BCAS2 (Supplemental Figure 12), confirming that the

binding of BCAS2 to β-catenin would not interfere with the formation of β-

catenin/TCF complex.

BCAS2 enhances β-catenin nuclear accumulation through its CC domains

To determine which domain of BCAS2 binds to β-catenin, we constructed a series of

deletion mutants of BCAS2 (Figure 7A). Notably, we observed that among these

truncated mutants, only the one lacking both CC1 and CC2 domains lost the ability to

interact with β-catenin (Figure 7B). Moreover, these two CC domains alone or

together could interact with β-catenin (Figure 7C). Therefore, we conclude that

BCAS2 binds to β-catenin via its CC domains.

We next examined whether the CC domains are required for BCAS2 to promote

Wnt/β-catenin signaling. As shown in Figure 7D, overexpression of BCAS2 without

the CC domains failed to increase LiCl-induced TOPflash activity in HEK-293T cells.

Likewise, overexpression of the full-length or the CC domains alone, but not BCAS2

lacking the CC domains, restored the nuclear accumulation of β-catenin in *bcas2*

morphants (Figure 7E-F and Supplemental Figure 13A-B). The expression of *gata1* in

bcas2 mutants was also recovered by overexpression of the full-length BCAS2, but

not the truncated form without the CC-domains (Figure 7G). Collectively, these

findings indicate that BCAS2 positively regulates Wnt signaling through sequestering

β-catenin within the nucleus via its CC domains during primitive hematopoiesis.

As BCAS2 is involved in the Prp19-CDC5L spliceosome complex that regulates

RNA splicing during spermiogenesis, neurogenesis, and definitive hematopoiesis,[15,16]

we wondered if this protein participates in primitive hematopoiesis via mRNA

alternative splicing. To this end, we performed RNA sequencing of 10-somite stage

embryos to identify abnormal events in alternative splicing in *bcas2*[+/Δ14] mutants.

However, upon haploinsufficiency of *bcas2*, neither the number of five major types of

alternative splicing events, nor the typical forms of alternative splicing were

significantly affected (Supplemental Figure 14A-B). Additionally, haploinsufficiency

of *bcas2* did not result in the alternative splicing of *mdm4* that predisposes cells to

undergo p53-mediated apoptosis in definitive hematopoiesis, as reported previously

by Yu *et al.* (Supplemental Figure 14C).[16,57] Furthermore, the splicing efficiency of β-

catenin pre-mRNA remained almost unchanged in *bcas2*[+/Δ14] mutants (Supplemental

Figure 14D). These results demonstrate that the defects in primitive hematopoiesis of

bcas2$^{+/Δ14}$ mutants are independent of the regulatory role of Bcas2 in pre-mRNA

splicing.

Discussion

BCAS2 is a 26-kDa nuclear protein involved in a multitude of developmental

processes, such as Drosophila wing development, dendritic growth, and

spermatogenesis.[8,14,15,58-60] In our study, we generated *bcas2* knockout zebrafish. The

heterozygotes not only showed male infertility, resembling the phenotype of *Bcas2*

germ cell-specific knockout mice reported previously,[15] but also exhibited impaired

definitive hematopoiesis, consistent with the earlier study.[16] Importantly, we found a

marked decrease in the expression of the primitive erythroid progenitor markers *gata1*

and *hbbe3* in these heterozygous mutants, which was rescued by overexpression of

BCAS2. Moreover, the defective primitive hematopoiesis in mutant zebrafish was

phenocopied in hemangioblast-specific *Bcas2* knockout mice. While the reason(s) for

the discrepancy between our data and the observations made by Yu *et al.* regarding the

role of *bcas2* in the development of primitive erythroid and myeloid cells remains to

be determined,[16] our findings in zebrafish and mouse embryos provide solid evidence

that BCAS2 plays a conserved role in primitive hematopoiesis.

As demonstrated in previous studies, BCAS2 is involved in various developmental

events by regulating pre-mRNA splicing.[14-16,29,30] However, our data showed that

haploinsufficiency of *bcas2* did not affect alternative splicing during primitive

hematopoiesis. These results imply that one copy of the *bcas2* gene is sufficient to

support mRNA splicing in zebrafish. Instead, we find that Bcas2 promotes primitive

hematopoiesis by sequestering β-catenin within the nucleus. It has been reported that

the *bcas2* deletion in zebrafish embryos induces alternative splicing of *Mdm4* that

predisposes cells to undergo p53-mediated apoptosis in HSPCs during definitive

hematopoiesis.[16] Intriguingly, we found that the loss of one copy of *bcas2* gene in

zebrafish also resulted in severe impairment of HSPCs and their derivatives. It is

possible that Bcas2 might also have a role in definitive hematopoiesis independent of

its splicing regulatory function.

For the past decades, given the contradictory conclusions obtained from various *in*

vitro and *in vivo* studies, the function of Wnt/β-catenin signaling in primitive

hematopoiesis remains elusive and controversial.[25-28] In the present study, we have

provided several lines of evidence supporting that Wnt/β-catenin signaling positively

regulates primitive hematopoiesis: (1) Inhibition of Wnt/β-catenin by overexpression

of the canonical Wnt inhibitor Dkk1 disrupts the formation of erythrocyte progenitors

at the 10-somite stage. (2) Defects in primitive hematopoiesis in *bcas2* morphants and

mutants are readily restored by overexpression of ΔN-β-catenin, a constitutively

active β-catenin. (3) Overexpression of the full-length BCAS2, but not the CC

domain-deleted BCAS2, restores the formation of the primitive erythroid progenitor

in *bcas2* mutants. (4) BCAS2 overexpression enhances the development of primitive

blood cells in wild-type embryos. All these data suggest that BCAS2-mediated Wnt/β-

catenin signal activation is necessary for primitive hematopoiesis.

In addition, Wnt/β-catenin signaling has been known as an important pathway

involved in the regulation of axis determination and neural patterning during

gastrulation.[61-64] In our study, neither the heterozygous *bcas2* mutant embryos nor the

very few homozygous ones, exhibited any morphological defects typically associated

with inhibition of Wnt signaling, such as ventralization or brain anteriorization. This

may be due to the presence of maternal Bcas2 in heterozygous and homozygous

mutant embryos which were derived from crossing *bcas2*[+/-] males with *bcas2*[+/-]

females.

CRM1 can facilitate β-catenin nuclear export in distinct ways.[65] For example,

CRM1 usually recognizes and binds with the nuclear export signal (NES) sequences

in chaperon proteins, such as APC, Axin and Chibby,[66-68] to mediate the nuclear

export of β-catenin. On the other hand, CRM1 can also bind directly to and function

as an efficient nuclear exporter for β-catenin.[69] Since BCAS2 has not been reported to

contain any recognizable NES sequences, it will be interesting to investigate whether

BCAS2 competitively inhibits β-catenin from associating with CRM1, or with the

chaperone proteins.

In summary, we uncover a novel role of BCAS2 in primitive hematopoiesis

through enhancing nuclear retention of β-catenin. Our study provides new insights

into the mechanism of BCAS2-mediated Wnt signal activation during primitive

hematopoiesis. Given that BCAS2 and Wnt signaling are well documented to

contribute to cancer development,[70-74] it is appealing to further explore whether our

findings can be applied to future cancer research.

Acknowledgements

We acknowledge the financial support of the National Natural Science Foundation of China (32025014 and 32330029) and the National Key Research and Development Program of China (2018YFA0800200 and 2020YFA0804000).

Author Contributions

Guozhu Ning: Writing - Original draft, Visualization, Software, Methodology, Investigation, Formal analysis, Data curation, Conceptualization**. Yu Lin:** Visualization, Software, Methodology, Investigation, Formal analysis, Data curation**. Haixia Ma, Jiaqi Zhang, Liping Yang, Zhengyu Liu:** Methodology, Validation**. Lei Li:** Resources**. Xinyu He:** Writing - Review & Editing**. Qiang Wang:** Project administration, Supervision, Funding acquisition**.**

Conflict-of-interest disclosure

The authors declare no competing financial interests.

Additional files

Supplemental Figures

Supplemental Table 1

Supplemental Table 2

References

1. Galloway JL, Zon LI. Ontogeny of hematopoiesis: examining the emergence of hematopoietic cells in the vertebrate embryo. *Curr Top Dev Biol*. 2003;53:139-158.

2. Murry CE, Keller G. Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development. *Cell*. 2008;132(4):661-680.

3. Ferkowicz MJ, Yoder MC. Blood island formation: longstanding observations and modern interpretations. *Exp Hematol*. 2005;33(9):1041-1047.

4. Palis J. Hematopoietic stem cell-independent hematopoiesis: emergence of erythroid, megakaryocyte, and myeloid potential in the mammalian embryo. *FEBS Lett*. 2016;590(22):3965-3974.

5. Paik EJ, Zon LI. Hematopoietic development in the zebrafish. *Int J Dev Biol*. 2010;54(6-7):1127-1137.

6. Detrich HW, 3rd, Kieran MW, Chan FY, et al. Intraembryonic hematopoietic cell migration during vertebrate development. *Proc Natl Acad Sci U S A*. 1995;92(23):10713-10717.

7. Leung AY, Mendenhall EM, Kwan TT, et al. Characterization of expanded intermediate cell mass in zebrafish chordin morphant embryos. *Dev Biol*. 2005;277(1):235-254.

8. Kuo PC, Tsao YP, Chang HW, et al. Breast cancer amplified sequence 2, a novel negative regulator of the p53 tumor suppressor. *Cancer Res*. 2009;69(23):8877-8885.

9. Neubauer G, King A, Rappsilber J, et al. Mass spectrometry and EST-database searching allows characterization of the multi-protein spliceosome complex. *Nature*

Genetics. 1998;20(1):46-50.

10. Nagasaki K, Maass N, Manabe T, et al. Identification of a novel gene, DAM1, amplified at chromosome 1p13.3-21 region in human breast cancer cell lines. *Cancer Letters*. 1999;140(1-2):219-226.

11. Qi C, Zhu YT, Chang J, Yeldandi AV, Rao MS, Zhu YJ. Potentiation of estrogen receptor transcriptional activity by breast cancer amplified sequence 2. *Biochem Biophys Res Commun*. 2005;328(2):393-398.

12. Ajuh P, Kuster B, Panov K, Zomerdijk JC, Mann M, Lamond AI. Functional analysis of the human CDC5L complex and identification of its components by mass spectrometry. *EMBO J*. 2000;19(23):6569-6581.

13. Grote M, Wolf E, Will CL, et al. Molecular architecture of the human Prp19/CDC5L complex. *Mol Cell Biol*. 2010;30(9):2105-2119.

14. Chen PH, Lee CI, Weng YT, et al. BCAS2 is essential for Drosophila viability and functions in pre-mRNA splicing. *RNA*. 2013;19(2):208-218.

15. Liu W, Wang F, Xu Q, et al. BCAS2 is involved in alternative mRNA splicing in spermatogonia and the transition to meiosis. *Nat Commun*. 2017;8:14182.

16. Yu S, Jiang T, Jia D, et al. BCAS2 is essential for hematopoietic stem and progenitor cell maintenance during zebrafish embryogenesis. *Blood*. 2019;133(8):805-815.

17. Bissinger R, Lang E, Gonzalez-Menendez I, et al. Genetic deficiency of the tumor suppressor protein p53 influences erythrocyte survival. *Apoptosis*. 2018;23(11-12):641-650.

18. Yang S, Cao S, Xu X, et al. adducin 1 is essential for the survival of erythroid

precursors via regulating p53 transcription in zebrafish. *iScience*. 2023;26(9):107516.

19. Stanic K, Reig G, Figueroa RJ, et al. The Reprimo gene family member, reprimo-

like (rprml), is required for blood development in embryonic zebrafish. *Sci Rep*.

2019;9(1):7131.

20. Richter J, Traver D, Willert K. The role of Wnt signaling in hematopoietic stem

cell development. *Critical Reviews in Biochemistry and Molecular Biology*.

2017;52(4):414-424.

21. Krimpenfort RA, Nethe M. Canonical Wnt: a safeguard and threat for

erythropoiesis. *Blood Advances*. 2021;5(18):3726-3735.

22. Kokolus K, Nemeth MJ. Non-canonical Wnt signaling pathways in hematopoiesis.

Immunologic Research. 2010;46(1-3):155-164.

23. Tarafdar A, Dobbin E, Corrigan P, Freeburn R, Wheadon H. Canonical Wnt

signaling promotes early hematopoietic progenitor formation and erythroid

specification during embryonic stem cell differentiation. *PLoS One*.

2013;8(11):e81030.

24. Nostro MC, Cheng X, Keller GM, Gadue P. Wnt, activin, and BMP signaling

regulate distinct stages in the developmental pathway from embryonic stem cells to

blood. *Cell Stem Cell*. 2008;2(1):60-71.

25. Sturgeon CM, Ditadi A, Awong G, Kennedy M, Keller G. Wnt signaling controls

the specification of definitive and primitive hematopoiesis from human pluripotent

stem cells. *Nature Biotechnology*. 2014;32(6):554-+.

26. Tran HT, Sekkali B, Van Imschoot G, Janssens S, Vleminckx K. Wnt/beta-catenin

signaling is involved in the induction and maintenance of primitive hematopoiesis in

the vertebrate embryo. *Proc Natl Acad Sci U S A*. 2010;107(37):16160-16165.

27. Lengerke C, Schmitt S, Bowman TV, et al. BMP and Wnt specify hematopoietic

fate by activation of the Cdx-Hox pathway. *Cell Stem Cell*. 2008;2(1):72-82.

28. Paluru P, Hudock KM, Cheng X, et al. The negative impact of Wnt signaling on

megakaryocyte and primitive erythroid progenitors derived from human embryonic

stem cells. *Stem Cell Research*. 2014;12(2):441-451.

29. Chen HH, Lu HY, Chang CH, et al. Breast carcinoma-amplified sequence 2

regulates adult neurogenesis via β-catenin. *Stem Cell Research & Therapy*.

2022;13(1).

30. Huang CW, Chen YW, Lin YR, et al. Conditional Knockout of Breast Carcinoma

Amplified Sequence 2 (BCAS2) in Mouse Forebrain Causes Dendritic Malformation

via β-catenin. *Scientific Reports*. 2016;6.

31. Chang N, Sun C, Gao L, et al. Genome editing with RNA-guided Cas9 nuclease

in zebrafish embryos. *Cell Res*. 2013;23(4):465-472.

32. Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 are required for

mesendoderm induction by transforming growth factor-beta/nodal signals in zebrafish.

J Biol Chem. 2008;283(4):2418-2426.

33. Welten MC, de Haan SB, van den Boogert N, et al. ZebraFISH: fluorescent in situ

hybridization protocol and three-dimensional imaging of gene expression patterns.

Zebrafish. 2006;3(4):465-476.

34. Wallingford MC, Giachelli CM. Loss of PiT-1 results in abnormal endocytosis in

the yolk sac visceral endoderm. *Mech Dev*. 2014;133:189-202.

35. Lieschke GJ, Oates AC, Crowhurst MO, Ward AC, Layton JE. Morphologic and

functional characterization of granulocytes and macrophages in embryonic and adult

zebrafish. *Blood*. 2001;98(10):3087-3096.

36. Wei S, Dai MM, Liu ZT, et al. The guanine nucleotide exchange factor Net1

facilitates the specification of dorsal cell fates in zebrafish embryos by promoting

maternal β-catenin activation. *Cell Research*. 2017;27(2):202-225.

37. Yang SY, Ning GZ, Hou YM, et al. Myoneurin regulates BMP signaling by

competing with Ppm1a for Smad binding. *Iscience*. 2022;25(6).

38. Schmierer B, Hill CS. Kinetic analysis of Smad nucleocytoplasmic shuttling

reveals a mechanism for transforming growth factor beta-dependent nuclear

accumulation of Smads. *Mol Cell Biol*. 2005;25(22):9845-9858.

39. Vogeli KM, Jin SW, Martin GR, Stainier DY. A common progenitor for

haematopoietic and endothelial lineages in the zebrafish gastrula. *Nature*.

2006;443(7109):337-339.

40. Reischauer S, Stone OA, Villasenor A, et al. Cloche is a bHLH-PAS transcription

factor that drives haemato-vascular specification. *Nature*. 2016;535(7611):294-298.

41. Sun H, Wang Y, Zhang J, et al. CFTR mutation enhances Dishevelled degradation

and results in impairment of Wnt-dependent hematopoiesis. *Cell Death Dis*.

2018;9(3):275.

42. Pilon N, Oh K, Sylvestre JR, Bouchard N, Savory J, Lohnes D. is a direct target

of the canonical Wnt pathway. *Developmental Biology*. 2006;289(1):55-63.

43. Sun HQ, Wang Y, Zhang JT, et al. CFTR mutation enhances Dishevelled degradation and results in impairment of Wnt-dependent hematopoiesis. *Cell Death & Disease*. 2018;9.

44. Stoick-Cooper CL, Weidinger G, Riehle KJ, et al. Distinct Wnt signaling pathways have opposing roles in appendage regeneration. *Development*. 2007;134(3):479-489.

45. Krause U, Ryan DM, Clough BH, Gregory CA. An unexpected role for a Wnt-inhibitor: Dickkopf-1 triggers a novel cancer survival mechanism through modulation of aldehyde-dehydrogenase-1 activity. *Cell Death & Disease*. 2014;5.

46. Meijer L, Skaltsounis AL, Magiatis P, et al. GSK-3-selective inhibitors derived from Tyrian purple indirubins. *Chem Biol*. 2003;10(12):1255-1266.

47. Easwaran V, Song V, Polakis P, Byers S. The ubiquitin-proteasome pathway and serine kinase activity modulate adenomatous polyposis coli protein-mediated regulation of beta-catenin-lymphocyte enhancer-binding factor signaling. *J Biol Chem*. 1999;274(23):16641-16645.

48. Rosin-Arbesfeld R, Cliffe A, Brabletz T, Bienz M. Nuclear export of the APC tumour suppressor controls beta-catenin function in transcription. *EMBO J*. 2003;22(5):1101-1113.

49. Lu Y, Xie S, Zhang W, et al. Twa1/Gid8 is a beta-catenin nuclear retention factor in Wnt signaling and colorectal tumorigenesis. *Cell Res*. 2017;27(12):1422-1440.

50. Henderson BR, Fagotto F. The ins and outs of APC and beta-catenin nuclear

transport. *EMBO Rep*. 2002;3(9):834-839.

51. Henderson BR. Nuclear-cytoplasmic shuttling of APC regulates beta-catenin

subcellular localization and turnover. *Nat Cell Biol*. 2000;2(9):653-660.

52. Xu L, Massague J. Nucleocytoplasmic shuttling of signal transducers. *Nat Rev*

Mol Cell Biol. 2004;5(3):209-219.

53. Wolff B, Sanglier JJ, Wang Y. Leptomycin B is an inhibitor of nuclear export:

inhibition of nucleo-cytoplasmic translocation of the human immunodeficiency virus

type 1 (HIV-1) Rev protein and Rev-dependent mRNA. *Chem Biol*. 1997;4(2):139-

147.

54. Kirli K, Karaca S, Dehne HJ, et al. A deep proteomics perspective on CRM1-

mediated nuclear export and nucleocytoplasmic partitioning. *Elife*. 2015;4.

55. Dimitrova YN, Li J, Lee YT, et al. Direct ubiquitination of beta-catenin by Siah-1

and regulation by the exchange factor TBL1. *J Biol Chem*. 2010;285(18):13507-

13516.

56. Graham TA, Weaver C, Mao F, Kimelman D, Xu WQ. Crystal structure of a β-

catenin/Tcf complex. *Cell*. 2000;103(6):885-896.

57. Rallapalli R, Strachan G, Cho B, Mercer WE, Hall DJ. A novel MDMX transcript

expressed in a variety of transformed cell lines encodes a truncated protein with

potent p53 repressive activity. *Journal of Biological Chemistry*. 1999;274(12):8299-

8308.

58. Huang CW, Chen YW, Lin YR, et al. Conditional Knockout of Breast Carcinoma

Amplified Sequence 2 (BCAS2) in Mouse Forebrain Causes Dendritic Malformation

via beta-catenin. *Sci Rep*. 2016;6:34927.

59. Xu Q, Wang F, Xiang Y, et al. Maternal BCAS2 protects genomic integrity in mouse early embryonic development. *Development*. 2015;142(22):3943-3953.

60. Zhang J, Liu W, Li G, et al. BCAS2 is involved in alternative splicing and mouse oocyte development. *FASEB J*. 2022;36(2):e22128.

61. Yamaguchi TP. Heads or tails: Wnts and anterior-posterior patterning. *Current Biology*. 2001;11(17):R713-R724.

62. Kozmikova I, Kozmik Z. Wnt/β-catenin signaling is an evolutionarily conserved determinant of chordate dorsal organizer. *Elife*. 2020;9.

63. Brafman D, Willert K. Wnt/β-Catenin Signaling during Early Vertebrate Neural Development. *Developmental Neurobiology*. 2017;77(11):1239-1259.

64. Lickert H, Cox B, Wehrle C, Taketo MM, Kemler R, Rossant J. Dissecting Wnt/β-catenin signaling during gastrulation using RNA interference in mouse embryos. *Development*. 2005;132(11):2599-2609.

65. Morgan RG, Ridsdale J, Tonks A, Darley RL. Factors Affecting the Nuclear Localization of β-Catenin in Normal and Malignant Tissue. *Journal of Cellular Biochemistry*. 2014;115(8):1351-1361.

66. Neufeld KL, Nix DA, Bogerd H, et al. Adenomatous polyposis coli protein contains two nuclear export signals and shuttles between the nucleus and cytoplasm. *Proceedings of the National Academy of Sciences of the United States of America*. 2000;97(22):12085-12090.

67. Li FQ, Mofunanya A, Harris K, Takemaru KI. Chibby cooperates with 14-3-3 to

regulate β-catenin subcellular distribution and signaling activity. *Journal of Cell*

Biology. 2008;181(7):1141-1154.

68. Cong F, Varmus H. Nuclear-cytoplasmic shuttling of Axin regulates subcellular

localization of β-catenin. *Proceedings of the National Academy of Sciences of the*

United States of America. 2004;101(9):2882-2887.

69. Ki H, Oh M, Chung SW, Kim K. β-Catenin can bind directly to CRM1

independently of adenomatous polyposis coli, which affects its nuclear localization

and LEF-1/β-catenin-dependent gene expression. *Cell Biology International*.

2008;32(4):394-400.

70. Murillo-Garzón V, Kypta R. WNT signalling in prostate cancer. *Nature Reviews*

Urology. 2017;14(11):683-696.

71. Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. *Oncogene*.

2017;36(11):1461-1473.

72. Yu FY, Yu CH, Li FF, et al. Wnt/β-catenin signaling in cancers and targeted

therapies. *Signal Transduction and Targeted Therapy*. 2021;6(1).

73. Salmerón-Hernández A, Noriega-Reyes MY, Jordan A, Baranda-Avila N, Langley

E. BCAS2 Enhances Carcinogenic Effects of Estrogen Receptor Alpha in Breast

Cancer Cells. *International Journal of Molecular Sciences*. 2019;20(4).

74. Wang LP, Chen TY, Kang CK, Huang HP, Chen SL. BCAS2, a protein enriched

in advanced prostate cancer, interacts with NBS1 to enhance DNA double-strand

break repair. *British Journal of Cancer*. 2020;123(12):1796-1807.

Figures and figure legends

Figure 1

Figure 1. *bcas2* is expressed in the ICM and required for primitive hematopoiesis.

(A) WISH assay showing *bcas2* expression in the ICM at the 18-somite stage and 22

hpf. The dotted lines represent the section position and the black arrowheads indicate

the ICM region. n, notochord. (B) Double FISH assay showing the expression pattern

of *bcas2* and *gata1* in the ICM at 22 hpf. Scale bar, 50 μm. (C-D) Comparison of

bcas2 expression in *cloche* mutants (C) or *bcas2* heterozygous mutants (D) along with

their corresponding siblings. (E-F) Expression analysis of *gata1* and *hbbe3* in

bcas2$^{+/\Delta 7}$ and *bcas2*$^{+/\Delta 14}$ embryos. (G) Hemoglobin detection using *o*-dianisidine

staining in *bcas2* homozygous mutant at 36 and 48 hpf. (H) Representative images of

yolk sac from the hemangioblast-specific *Bcas2* knockout mice and their siblings.

Bcas2$^{F/F}$ females were crossed with *Bcas2*$^{F/+}$;*Flk1-Cre* males to induce the deletion of

Bcas2 in hemangioblasts. Scale bar, 1 mm.

Figure 2

Figure 2. *bcas2* is required for hematopoietic progenitor differentiation. (A-C)

Expression analysis of hemangioblast markers *npas4l, scl, gata2* (A), erythroid

progenitor markers *gata1*, *hbbe3* (B), and myeloid marker *pu.1* (C) in *bcas2$^{+/\Delta14}$*

embryos and their wild-type siblings at indicated stages. (D) Expression changes of

gata1 in *bcas2$^{+/\Delta14}$* embryos overexpressing BCAS2 at the 10-somite stage. The

indicated embryos were injected with or without 300 pg of human *BCAS2* mRNA at

the one-cell stage. (E) Expression of endothelial marker *fli1a* in *bcas2$^{+/\Delta14}$* and sibling

embryos at the 10-somite stage. (G) Confocal imaging of *bcas2$^{+/\Delta14}$* and control

sibling *Tg(kdrl:GFP)* embryos at 54 hpf. Scale bar, 500 μm.

Figure 3

Figure 3. BCAS2 promotes primitive hematopoiesis via activating Wnt signaling.

(A-B) Overexpression of BCAS2 increases Wnt3a-induced TOPflash activity in

HEK293T cells (A) and MEFs (B). Different amounts of plasmid expressing BCAS2

(0, 80, 160, or 320 ng/well) were transfected into cells, together with the super-

TOPflash luciferase and Renilla luciferase vectors. After 36 h of transfection, cells

were treated with or without Wnt3a CM for 12 h and harvested for luciferase assays.

*P < 0.05; **P < 0.01 (Student's *t-test*). (C) The Wnt3a-induced TOPflash activity is

decreased in BCAS2-deficient cells. HEK293T Cells were transfected with shRNA

plasmids, along with indicated plasmids, and harvested for luciferase reporter assay.

*P < 0.05 (Student's *t-test*) (D) *Bcas2*-cKO MEFs prepared from *Bcas2^{F/F}* mouse

embryos were incubated in medium containing 100 μM tamoxifen for 72 h and then

subjected to western blotting and luciferase reporter assay. **P < 0.01 (Student's *t-*

test). (E-F) Expression analysis of *gata1* (E) and *hbbe3* (F) in *Tg(hsp70l:dkk1b-GFP)*

embryos after heat shock at 16 hpf. (G-H) Immunofluorescence staining of β-catenin

in *Tg(gata1:GFP)* embryos at 16 hpf. The embryos were injected with 8 ng of the

indicated MO at the one-cell stage. The dotted lines show the GFP-positive

hematopoietic progenitor cells. Scale bars, 5 μm. The relative fluorescence intensity

of nuclear β-catenin was quantified in (H). **P < 0.01 (Student's *t-test*). (I-J)

Expression of *hbbe3* in *bcas2* morphants (I) and *bcas2^{+/Δ14}* mutants (J) overexpressing

ΔN-β-catenin. Embryos were injected with the indicated MO together with *ΔN-β-*

*cateni*n mRNA at the one-cell stage and harvested at the 10-somite stage for *in situ*

hybridization.

Figure 4

Figure 4. BCAS2 is essential for β-catenin nuclear accumulation. (A-C) BCAS2

enhances LiCl-induced TOPflash activity in HEK293T cells. Cells were transfected

with BCAS2 expression plasmids (A), shRNA plasmids (B), or S37A-β-catenin

expression plasmids (C), together with the TOPflash luciferase and Renilla luciferase

vectors. After transfection, cells were subsequently treated with or without 100 ng/mL

LiCl for 12 h and assayed for luciferase activity. *P < 0.05; **P < 0.01 (Student's *t*-

test). (D-E) *Bcas2*-cKO MEFs were incubated with tamoxifen for 24 h and then

treated with or without 100 ng/mL LiCl. The nuclear accumulation of β-catenin was

analyzed using immunofluorescence (D) and western blotting (n=3) (E) Scale bars, 10

μm. (F) SW480 cells were transfected with the indicated shRNA constructs, and the

endogenous β-catenin protein was detected using immunofluorescence 48 h after

transfection. The expression of GFP served as a transfection control. The arrowheads

indicate the cells transfected with indicated shRNA constructs. Scale bars, 10 μm. (G)

Bcas2-cKO MEFs were cultured in the presence of tamoxifen for 24 h and then

treated with 20 μM MG132 for 6 h. The expression of BCAS2 and β-catenin was

measured by immunofluorescence. Scale bars, 10 μm.

Figure 5

Figure 5. BCAS2 functions in CRM1-mediated nuclear export of β-catenin. (A)

Tamoxifen-treated *Bcas2*-cKO MEFs were incubated with 20 nM LMB for 3 h. The

expression of Bcas2 and β-catenin was analyzed using immunofluorescence. The

arrowheads show the cells with nuclear β-catenin accumulation. Scale bars, 10 μm. (B)

SW480 cells were transfected with the indicated shRNA constructs and then treated

with LMB for 3 h before immunostaining. GFP was regarded as a transfection control.

The arrowheads indicate the transfected cells. Scale bars, 10 μm. (C-D)

Immunofluorescence staining of β-catenin in *bcas2* morphants with *Tg(gata1:GFP)*

background at 16 hpf. Embryos were exposed to 20 nM LMB from the bud stage. The

dotted lines indicate the GFP-positive hematopoietic progenitor cells. Scale bars, 5

μm. The relative fluorescence intensity of nuclear β-catenin was quantified in (D). ns,

not significant; **$P < 0.01$ (Student's *t-test*). (E) *bcas2$^{+/Δ14}$* embryos were treated with

20 nM LMB for 6 h and then subjected to WISH assay to analyze the expression of

gata1 at the indicated stages.

Figure 6

Figure 6. BCAS2 interacts with β-catenin. (A-C) Flag-tagged β-catenin was co-transfected with or without HA-tagged BCAS2 into HEK293T cells. Cell lysates were immunoprecipitated using anti-Flag antibody. Eluted proteins were analyzed by western blotting using indicated antibodies. In panel C, for Wnt signaling activation, cells were treated with Wnt3a CM for 5 h before harvest. (D-E) YN-BCAS2 and YC-β-catenin were either individually or collectively transfected into HeLa cells. The expression of YN-BCAS2 and YC-β-catenin was analyzed with anti-GFP antibody (D). The reconstituted YFP fluorescence in living cells was detected by confocal laser scanning microscopy with excitation at 488 nm (E). Scale bars, 10 μm. (F) Schematics of full-length and deletion mutants of β-catenin. (G) HEK293T cells were transfected with HA-tagged BCAS2 and Flag-tagged deletion mutants of β-catenin. Cell lysates were then immunoprecipitated using anti-Flag antibody followed by western blot analysis. (H) GST pull-down assays were performed using bacterially expressed GST, GST-ARM1-12, and His-BCAS2.

Figure 7

Figure 7. BCAS2 sequesters β-catenin in the nucleus via its CC domains. (A)

Schematics of full length and deletion mutants of BCAS2. (B-C) HEK293T cells were

transfected with Flag-β-catenin and indicated deletion mutants of BCAS2. Cell lysates

were subjected to immunoprecipitation with anti-Flag antibody. Eluted proteins were

immunoblotted using anti-HA (B) or anti-GFP antibodies (C) for BCAS2 detection.

(D) HEK293T cells transfected with the indicated plasmids were treated with 100

ng/mL LiCl for 12 h, and then subjected to luciferase assay. Note that overexpression

of BCAS2 without the CC domains failed to increase LiCl-induced TOPflash activity.

Data represent the mean ± SD of three independent experiments. ns, not significant;

**$P < 0.01$ (Student's *t-test*). (E-F) Immunofluorescence staining of β-catenin in

Tg(gata1:GFP) embryos at 16 hpf. The embryos were injected with 8 ng *bcas2* MO

and 300 pg of full length *BCAS2* mRNA or *ΔCC1-2* mRNA at the one-cell stage.

Scale bars, 5 μm. The relative fluorescence intensity of nuclear β-catenin was

quantified in (F). ns, not significant; ****$P < 0.01$; ****$P < 0.0001$** (Student's *t-test*). (G)

Transcripts of *gata1* were evaluated by WISH in *bcas2$^{+/Δ14}$* embryos injected with 300

pg of *BCAS2* mRNA or *ΔCC1-2* mRNA.

Supplemental figures

Figure S1. Expression patterns of *bcas2* in wild-type embryos during

development. Expression of *bcas2* in wild-type embryos at the indicated

developmental stages was analyzed using whole-mount *in situ* hybridization.

Figure S2. Zebrafish *bcas2* mutants are generated by using CRISPR/Cas9 system.

(A) Schematic showing generation of *bcas2* mutants. Two mutant lines were obtained

with mutations that resulted in premature translation termination, resulting in

truncated Bcas2 proteins lacking the C-terminal CC domains. (B) Identification of

bcas2 mutations using DNA sequencing. (C) The *bcas2*[+/Δ7] and *bcas2*[+/Δ14] mutants

were identified via T7 endonuclease (the upper panel) or FspI restriction enzyme (the

lower panel) digestions. (D) Bright field images of embryos derived from crossing

indicated females with heterozygous male mutants. Black arrows refer to the embryos

that exhibited an abnormal cleavage. The ratio of viable embryos was indicated.

Figure S3. *bcas2* is essential for definitive hematopoiesis. (A-B) Expression

changes of *cymb* and *rag1* in *bcas2$^{+/\Delta 7}$* (A) and *bcas2$^{+/\Delta 14}$* (B) embryos compared with

their siblings. Arrowheads on the left panels indicate the caudal hematopoietic tissue,

and those on the right panels indicate the thymus.

969 **Figure S4. Knockdown of *bcas2* impairs primitive hematopoiesis.** (A) Western blot

970 analysis showing the expression changes of Bcas2 protein in wild-type embryos and

971 embryos injected with 8 ng control MO (cMO) or *bcas2* MO. (B-C) Expression of

972 *gata1* (B) and *hbbe3* (C) in *bcas2* morphants and control embryos. (D) Detection of

973 hemoglobin levels by *o*-dianisidine staining in *bcas2* morphants and control embryos.

Figure S5. *bcas2* **is dispensable for the survival and proliferation of**

hematopoietic cells. (A) Cell apoptosis was detected by TUNEL assay in *bcas2$^{+/\Delta14}$*

embryos and their siblings with *Tg(gata1:GFP)* background. Scale bar, 50 μm. (B-C)

Cell proliferation was detected by BrdU staining in *bcas2$^{+/\Delta14}$* embryos and their

siblings with *Tg(gata1:GFP)* background. Scale bars, 20 μm. The percentage of

BrdU$^+$/GFP$^+$ cells (% over the total number of GFP$^+$ cells) was quantified from six

embryos in each group (C). Data represent the mean ± SD of three independent

experiments. ns, not significant (Student's *t-test*).

Figure S6. Western blot analysis of HEK293T Cells transfected with corresponding shRNA constructs.

999 **Figure S7. Expression patterns of *cdx4* and *hoxa9a* in *bcas2*^{+/Δ14} embryos and**

1000 **their siblings at the 6-somite stage.** Black arrows indicate the posterior lateral

1001 mesoderm.

1002

1003

Figure S8. Inhibition of Wnt signaling does not affect the generation of hemangioblasts or their endothelial differentiation, but impairs their hematopoietic differentiation. Expression of hemangioblast markers *npas4l, scl, gata2* (A), endothelial marker *fli1a* (B) and erythroid progenitor marker *gata1* (C) at the indicated stages. Wild-type embryos were treated with 10 μM CCT036477 from 9 hpf and then collected for whole-mount *in situ* hybridizations.

1011

Figure S9. Knockdown of *bcas2* significantly reduces nuclear β-catenin in the primitive myeloid cells. Immunofluorescence staining of β-catenin in *Tg(coro1a:GFP)* embryos at 17 hpf. The embryos were injected with 8 ng of the indicated MOs at the one-cell stage and then collected for immunofluorescence staining. The dotted lines refer to the GFP-positive primitive myeloid cells. Scale bars, 10 μm. The relative fluorescence intensity of nuclear β-catenin was quantified in (B). **$P < 0.01$ (Student's *t-test*).

1020

Figure S10. BCAS2 inhibits the nuclear export of β-catenin. (A-B') GFP-tagged

S37A-β-catenin was co-expressed with or without BCAS2 in HeLa cells. The entire

nucleus (A) or entire cytoplasm (B) was bleached. (A-A') Representative images at

the indicated timepoints showing the kinetics of nuclear import of S37A-β-catenin are

shown in (A) and the related fluorescence recovery curves are shown in (A'). Time-

lapse imaging (B) and recovery curves (B') from cytoplasmic photobleaching

experiments showed the kinetics of nuclear export of S37A-β-catenin. Scale bars, 10

μm. (C) Quantitative analysis of fluorescence recoveries after photobleaching (n=7).

Data represent the mean ± SD of three independent experiments. Student's t-test was

used. ns, not significant, **$P < 0.01$ (Student's *t-test*).

1032

Figure S11. Overexpression of BCAS2 slightly enhances the nuclear accumulation of CDC37L, and has no influence on the distribution of ATG3. HeLa cells were transfected with the indicated plasmids and then subjected to immunostaining using anti-Flag and anti-HA antibodies. The nuclei were labeled with DAPI. Scale bar, 10 μm.

1033

1034

1035

1036

1037

1038

1039

1040 **Figure S12. The interaction between β-catenin and TCF4 remains unaffected in**

1041 **the presence of BCAS2.** HEK293T cells were transfected with the indicated

1042 constructs. Cell lysates were immunoprecipitated using anti-c-Myc antibody and the

1043 eluted proteins were analyzed by western blotting.

1044

1045

Figure S13. Overexpression of the CC domains of BCAS2 restores nuclear β-catenin accumulation in *bcas2* morphants. Immunofluorescence staining of β-catenin in *Tg(gata1:GFP)* embryos at 16 hpf. The embryos were injected with 8 ng *bcas2* MO and 300 pg of *BCAS2 CC1-2* mRNA at the one-cell stage. The dotted lines show the GFP-positive hematopoietic progenitor cells. Scale bars, 10 μm. The relative fluorescence of nuclear β-catenin was quantified in (B). *$P < 0.05$ (Student's *t-test*).

1052

1053

Figure S14. Haploinsufficiency of *bcas2* does not affect pre-mRNA splicing during primitive hematopoiesis. (A) The number of five major types of alternative splicing events was analyzed from RNA sequencing data. Embryos were lysed at the 10-somite stage and subjected to reverse transcription. The cDNA library was prepared, sequenced and then analyzed using rMATS. The five main alternative splicing types refer to exon skipping (SE), alternative 3' splicing site (A3SS), alternative 5' splicing site (A5SS), intron retention (RI), and mutually exclusive exons (MEX). (B) Examples of different types of alternative splicing were analyzed by reverse transcription PCR using total RNA from sibling and *bcas2^{+/Δ14}* embryos at the 10-somite stage. (C) Reverse transcription PCR analysis of total *mdm4*-FL and *mdm4*-S isoforms in *bcas2^{+/Δ14}* and sibling embryos at the 10-somite stage. (D) Analysis the pre-mRNA and mature mRNA of *β-catenin* in *bcas2^{+/Δ14}* and sibling embryos at the

1066   10-somite stage.

1067

1068 **Supplemental Table 1**

1069
<div align="center">

Primers Used for Genotyping

</div>

Strain	Directions	Sequence (5' to 3')
Flk1-Cre mouse	Forward	CGGTTATTCAACTTGCACCAC
	Reverse	CAGGACTGAAAGCCCAGACT
Bcas2^{Flox/Flox} mouse	Forward	ATTCCAGCAGTTGGTGTGGG
	Reverse	CATTGCTGGACAGAAGGTGAG
Flk1-Cre;*Bcas2*^{Flox/Flox} mouse	Forward	AGGTGTATGAATGCCTGAACAAG
	Reverse	CATTGCTGGACAGAAGGTGAG
bcas2 knockout zebrafish	Forward	TGCACATACAGTAATAGGCTTACCC
	Reverse	GTCTGATTTGCATCAAAAGATGTGA

1070

1071 **Supplemental Table 2**

1072

	Primers Used for Reverse Transcription-PCR	
Symbol	**Directions**	**Sequence (5' to 3')**
ddb2	Forward	CGCTCACTTAAATCTTACAAGCTGC
	Reverse	AGTCTGGGGTACGAGACAATATCT
	Reverse	TTGGCAAAGACTTGTATAACGGATC
acox	Forward	ATCCAGATTTCCAACATGAAGACCT
	Reverse	AGGTGCCTAACATATGGAATGACTC
	Forward	TTCTCTCGAAGTGAGCGATATGA
	Reverse	TCTGTGCGTAGGTGCCAATAATCTC
ubap2l	Forward	GAATGTTGGTGTCAATGCATCAGCT
	Reverse	CTGCGTCTTTGTGTAAACAGAGCCT
cdk16	Forward	ATGAGGAAGATCAAACGGCAGCTGT
	Reverse	AAATTTTGCGACCGGGGTTGTTTCT
rbbp4	Forward	TGGGACACTCGTTCCAATAACACAT
	Reverse	CTGGAAGATTTCATCTTTGTGCGAC
her6	Forward	GGAGAAAGAAGAAGAGCGAGAATC
	Reverse	GCTGCATGTTTCTGAGATGTTTCAC
cirbpb	Forward	TCCTACAGAGACGGTTACGACAGTT
	Reverse	CCTCACCAGGCTTGAAACTCTTACA
cdkn2aip	Forward	TCGCTAATGAGGACTTGTCTTTGGA
	Reverse	ATTTATCTTGTCCATCACACGCTGC
mapk1	Forward	GTCTGTTGGTTGTATTCTGGCTGAG
	Reverse	AGGATCATAGTACTGCTCCAGGTAC
add1	Forward	AGGATGCTGGATAATCTGGGCTACA
	Reverse	GTTGCGCATCTCTTGGACTTCTTTC
Mdm4-FL	Forward	GGTCAGGTGTCCAGTGAGTCAATAA
	Reverse	AAACCATCTGAGGAGTCTTCATCTG
Mdm4-S	Forward	GAAGATCCTGGTCAGACTCCTCAGA
	Reverse	CGGGAGAGAGTTGATTGGTGTGAAT
β-catenin	Forward	GATGACGATGTGGATAATCAGGTGC
	Reverse	GAACTGTGTGGAAGGTATCTGCATG
	Forward	GGAGACAATTTCTCATTATTCTGGC
	Reverse	TCAGTAGTTTGGTGAGCTCTGGAAT
β-actin	Forward	ATGGATGATGAAATTGCCGCAC
	Reverse	ACCATCACCAGAGTCCATCACG

1073