Title: BCAS2 promotes primitive hematopoiesis by sequestering β-catenin within

the nucleus

Authors: Guozhu Ning[1,2#], Yu Lin[3#], Haixia Ma[4,5], Jiaqi Zhang[4,5], Liping Yang[1],

Zhengyu Liu[1], Lei Li[4,5]*, Xinyu He[1]*, Qiang Wang[1]*

Affiliations:

[1]Innovation Centre of Ministry of Education for Development and Diseases, the Sixth

Affiliated Hospital, School of Medicine, South China University of Technology,

Guangzhou 510006, China

[2]Affiliated Hospital of Guangdong Medical University & Key Laboratory of Zebrafish

Model for Development and Disease of Guangdong Medical University, Zhanjiang

524001, China

[3]School of Pharmacy, Qiqihar Medical University, Qiqihar 161006, China

[4]Institute State Key Laboratory of Stem Cell and Reproductive Biology, Institute of

Stem Cell and Regeneration, Beijing Institute of Stem Cell and Regenerative Medicine,

Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.

[5]University of Chinese Academy of Sciences, Beijing 100049, China.

[#]These authors contributed equally.

*To whom correspondence should be addressed: (Lei Li) lil@ioz.ac.cn; (Xinyu He)

hexyu@scut.edu.cn; (Qiang Wang) qiangwang@scut.edu.cn

Running title: BCAS2 sequesters β-catenin within the nucleus

Abstract

Breast carcinoma amplified sequence 2 (BCAS2), a core component of the hPrP19 complex, plays an important role in RNA-splicing and DNA damage. However, whether BCAS2 has other functions within the nucleus remains largely unknown. Here, we show that BCAS2 is essential for primitive hematopoiesis in zebrafish and mouse embryos. The activation of Wnt/β-catenin signal, which is required for hematopoietic progenitor differentiation, is significantly decreased upon depletion of *bcas2* in zebrafish embryos and mouse embryonic fibroblasts. Interestingly, haploinsufficiency of *bcas2* has no obvious impact on the splicing efficiency of β-catenin pre-mRNA, while significantly attenuating β-catenin nuclear accumulation. Moreover, we find that BCAS2 directly binds to β-catenin via its coiled-coil domains, thereby sequestering β-catenin within the nucleus. Thus, our results uncover a previously unknown function of BCAS2 in promoting Wnt signaling by enhancing β-catenin nuclear retention during primitive hematopoiesis.

Key words: BCAS2, Wnt/β-catenin, coiled-coil domain, nuclear retention, primitive hematopoiesis

Introduction

Hematopoiesis refers to the lifelong process by which all blood cell lineages are generated. It begins at the early stage of embryonic development, providing the growing embryo with sufficient oxygen and nutrients.[1] Evolutionarily conserved across vertebrate species, hematopoiesis consists of two successive and partially overlapping waves: primitive and definitive. In mammals, the first wave of hematopoiesis occurs in the yolk-sac blood islands, producing primitive erythroid, megakaryocyte and macrophage progenitors, which can be observed in mouse embryos as early as embryonic day 7.25 (E7.25).[2-4] In zebrafish, primitive hematopoiesis initiates at around 11 hours post fertilization (hpf), when hemangioblasts emerge from the anterior lateral mesoderm (ALM) and posterior lateral mesoderm (PLM) and later differentiate into both hematopoietic and endothelial cells.[5-7]

Breast cancer amplified sequence 2 (BCAS2), also known as pre-mRNA splicing factor SPF27, is a 26-kDa nuclear protein containing two coiled-coil (CC) domains.[8] It was initially found to be overexpressed and amplified in human breast cancer cell lines.[9-11] Further studies have identified BCAS2 as a vital component of the human Prp19/CDC5L complex, which forms the catalytic ribonucleoprotein (RNP) core of spliceosome and is required for the activation of pre-mRNA splicing.[9,12,13] In Drosophila, the function of BCAS2 in RNA splicing is essential for cell viability.[14] In mouse, disruption of *Bcas2* in male germ cells impairs mRNA splicing and leads to a failure of spermatogenesis.[15] Additionally, BCAS2 also plays an important role in the

DNA damage response through the replication protein A (RPA) complex.[8,16]

Interestingly, it has been shown that BCAS2 is a negative regulator of p53 by directly

interacting with p53 or modulating alternative splicing of Mdm4, a major p53

inhibitor.[8,17] Zebrafish *bcas2* transcripts were enriched in the sites of both primitive and

definitive hematopoiesis during embryonic development.[17] However, a previous study

showed that p53 overactivation induced by zebrafish *bcas2* depletion did not affect

primitive hematopoiesis, but impaired definitive hematopoiesis.[17] In recent years,

several studies have highlighted the importance of regulating the expression and

activity of p53 in primitive erythroid cell differentiation in both mouse and zebrafish

embryos.[18-20] Thus, it is necessary to reexamine the exact function of BCAS2 in

primitive hematopoiesis.

Wnt signaling, usually categorized into canonical and non-canonical pathways, is

involved in the process of hematopoiesis.[21-23] Notably, the canonical Wnt signaling

pathway, which is dependent on the nuclear accumulation of β-catenin to regulate gene

transcription, controls primitive hematopoietic progenitor formation and promotes

definitive hematopoietic stem cell (HSC) specification.[24-26] For instance, it has been

demonstrated in *Xenopus* that, Wnt4-mediated activation of Wnt/β-catenin signaling

plays a critical role in the induction and maintenance of primitive hematopoiesis.[27]

Moreover, transient inhibition of canonical Wnt signaling in zebrafish embryos impairs

embryonic blood formation.[28] However, previous studies utilizing human pluripotent

stem cells revealed an opposite role of Wnt/β-catenin pathway in primitive progenitor

generation.[26,29] Therefore, the impact of Wnt/β-catenin signaling on primitive

hematopoiesis remains elusive and even controversial. Moreover, it has been suggested

that BCAS2 is important for neural stem cell proliferation and dendrite growth in mice

by regulating β-catenin pre-mRNA splicing.[30,31] As a nuclear protein, it is unclear

whether BCAS2 can modulate Wnt/β-catenin signaling in a splicing-independent

manner.

In this study, we generated two zebrafish *bcas2* mutant lines, both of which

exhibited defects in male fertility and embryonic HSC formation, similar to what was

previously reported in mice and zebrafish.[15,17] More importantly, loss-of-function

experiments suggest that BCAS2 is necessary for primitive hematopoiesis in both

zebrafish and mouse embryos. We further find that *bcas2* is dispensable for the survival

and proliferation of hematopoietic cells, but plays a crucial role in the differentiation of

the hematopoietic lineage from hemangioblasts. Using a comprehensive approach, we

reveal that BCAS2 is a nuclear retention factor for β-catenin during primitive

hematopoiesis. Subsequent biochemical and functional experiments demonstrate that

BCAS2 directly binds to β-catenin and suppresses its nuclear export to promote Wnt

signal activation and hematopoietic progenitor differentiation. Furthermore, the CC

domains on BCAS2 and the Armadillo (ARM) repeats on β-catenin are responsible for

their interaction. Collectively, we have uncovered a novel function of BCAS2 in

regulating Wnt/β-catenin signaling by sequestering β-catenin within the nucleus during

primitive hematopoiesis.

Methods

Animal models

Our studies including animal maintenance and experiments were performed in

compliance with the guidelines of the Animal Care and Use Committee of the South

China University of Technology (Permission Number: 2023092). Six strains of

zebrafish were used in this study, including Tübingen wild-type, *bcas2* mutant, *Cloche*

mutant, *Tg(gata1a:GFP), Tg(kdrl:GFP),* and *Tg(hsp70l:dkk1b-GFP)*. *Cloche* mutant,

Tg(gata1a:GFP) and *Tg(kdrl:GFP)* lines were provided by Professor Feng Liu

(Chinese Academy of Sciences). *Tg(hsp70l:dkk1b-GFP)* strain was purchased from

China Zebrafish Resource Center. $Bcas2^{Floxed/Floxed}$ ($Bcas2^{F/F}$) mouse line was generated

as previously described.[15] Genotyping of $Bcas2^{F/F}$ mouse and *Flk1*-Cre mouse was

performed using primers listed in Supplemental Table 1. The mouse model with *Bcas2*

specifically disrupted in the hemangioblasts was derived from mating female $Bcas2^{F/F}$

mice with *Flk1*-Cre transgenic mice. All mouse lines were maintained on a mixed

background (129/C57BL/6).

Cell lines and transfection

HEK293T, HeLa, SW480, and L cell lines were obtained from ATCC. Conditional

Bcas2 knockout (*Bcas2*-cKO) mouse embryonic fibroblasts (MEFs) were prepared

from $Bcas2^{F/F}$ embryos at E13.5. Cells were cultured in Dulbecco's modified eagle's

medium (DMEM, HyClone) supplemented with 10% fetal bovine serum (FBS,

HyClone) and 1% penicillin-streptomycin (HyClone) at 37 °C and 5% CO_2. L cell lines

expressing Wnt3a were maintained under similar conditions in the presence of 400

µg/ml G-418, from which Wnt3a conditioned medium (Wnt3a CM) was generated.

Culture medium prepared from L-cells was used as a control. To stimulate Wnt

signaling, cells were treated with Wnt3a CM in a 1:1 ratio with normal media. To

deplete *Bcas2* expression, *Bcas2*-cKO MEFs were cultured in medium containing 2 µM

tamoxifen for 72 h and the knockout efficiency was evaluated using western blot

analysis. The same cells cultured without tamoxifen were used as a control. To silence

BCAS2 expression, shRNA constructs in pLL 3.7-GFP plasmid were generated to target

the following sequences: shRNA1, GAATGTGTAAACAATTCTA; shRNA2:

GAAGGAACTTCAGAAGTTA. Transfection was performed with Lipofectamine

2000 (11668019, Invitrogen) according to the manufacturer's instructions.

Generation of CRISPR-Cas9-mediated *bcas2* knockout zebrafish

The *bcas2* knockout zebrafish mutants were generated by CRISPR-Cas9 system as

previously described.[32] The guide RNA was designed to target the sequences 5′-

GGCGCAGCTGGAGCATCAGG-3′ within exon 4 of *bcas2*. Humanized Cas9 mRNA

and gRNA were co-injected into wild-type embryos at the 1-cell stage. Embryos or

adult fin clips were collected to prepare genomic DNA. To screen for mutant alleles,

the genomic regions containing gRNA-targeted sequences were amplified by

polymerase chain reaction (PCR) with primers listed in Supplemental Table 1. The PCR

products were sequenced or digested with T7 endonuclease or restriction enzyme FspI

for genotyping.

RNA, morpholinos, and microinjection

Capped mRNAs for zebrafish *bcas2*, human *BCAS2*, *BCAS2 △CC1-2*, and mouse *ΔN-*

β-catenin mRNA were synthesized from the corresponding linearized plasmids using

an mMESSAGE mMACHINE T7 transcription kit (AM1344, Ambion). Morpholinos

(MOs) were designed and purchased from Gene Tools: mismatch MO (cMO 5'-

AGCCACTCATCCTGCTCCTCCCATC-3'), and *bcas2* translation-blocking MO

(tMO; 5'-AGCGACTGATGCTGGTCCTGCCATC-3'). The mRNAs and morpholinos

were injected into embryos at the 1- to 2-cell stage.

Whole-mount *in situ* hybridization

Digoxigenin-labeled and fluorescein-labeled probes were synthesized using a RNA

Labeling kit (11175025910, Roche). Whole-mount *in situ* hybridization (WISH) and

double fluorescence *in situ* hybridization for zebrafish embryos were performed

following previously published methods.[33,34] Anti-digoxigenin-POD (11633716001,

Roche) and anti-fluorescein-POD (11426346910, Roche) were used to detect

digoxigenin-labeled probes and fluorescein-labeled probes, respectively. After WISH,

the stained embryos were embedded in OCT and sections were prepared with a LEICA

CM1900. The mouse yolk sac layers were separated as previously described.[35]

o-Dianisidine staining

171 To evaluate hemoglobin level, embryos were harvested at 36 hpf or 48 hpf, then stained

172 with *o*-dianisidine as previously described.[36]

Proliferation and apoptosis assays

175 Embryos were incubated with 10 mM bromodeoxyuridine (BrdU) (B5002, Sigma) for

176 20 min. The incorporated BrdU was detected with anti-BrdU (B2531, Sigma) antibody.

177 TUNEL staining was performed using *In Situ* Cell Death Detection Kit, TMR red

178 (12156792910, Roche) according to the manufacturer's recommendation.

Heat shock treatment

181 To induce *dkk1* expression, *Tg(hsp70l:dkk1b-GFP)* embryos were subjected to heat

182 shock (42°C) for 10 min at 10 hpf, and then collected at the indicated stage for WISH.

Dual reporter assay

185 HEK293T cells or MEFs were seeded in 24-well plates and transfected with a Super-

186 TOPflash plasmid containing multimerized TCF-binding elements and a Renilla

187 luciferase plasmid, along with the indicated vectors. Then cells were treated with 100

188 ng/mL LiCl and/or Wnt3a CM for 12 h and assayed for luciferase activity using the

189 Dual luciferase system (E1910, Promega).

Immunoprecipitation, GST pulldown, and western blotting

For immunoprecipitation, HEK293T cells were transfected with the indicated plasmids and collected 48 h after transfection. Subsequently, HEK293T cells were lysed in a lysis buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 2 mM EDTA and 0.5% Nonidet P-40) containing protease inhibitors. Immunoprecipitation was performed in accordance with the standard protocols.

For GST pulldown assay, GST, GST tagged β-catenin ARM 1-12 and His tagged BCAS2 were expressed in *Escherichia coli* BL21, then purified using Glutathione-Sepharose 4B beads (71024800-GE, GE Healthcare) and HisPur Ni-NTA beads (88831, Thermo Fisher), respectively. GST and GST-β-catenin ARM 1-12 proteins were immobilized onto Glutathione-Sepharose 4B beads and incubated with purified His-BCAS2 at 4°C for 4 h. Beads were washed three times and analyzed using western blotting.

Cytoplasmic and nuclear extracts were separated with nuclear and cytoplasmic extraction kit (CW0199, CWBIO). Proteins were analyzed by western blot using affinity-purified anti-Flag (F2555, Sigma), anti-HA (CW0092A, CWBIO), anti-β-catenin (M24002, Abmart), anti-BCAS2 (10414-1-AP, Proteintech), anti-Tubulin (CW0265A, CWBIO), anti-GFP (A11120, Thermo Fisher), anti-Histone H3 (ab1791, Abcam), anti-GST (SAB4200237, Sigma), and anti-His Tag (AF5060, Beyotime) antibodies.

Immunofluorescence staining

Cells on coverslips and embryos were processed for immunofluorescence staining as previously described.[37,38] Before fixation, *bcas2*-deficient MEFs were treated with a concentration of 20 μM MG132 or 20 nM leptomycin B (LMB) for 6 h, while *Tg(gata1a:GFP)* embryos were treated with 20 nM LMB from the bud stage to the 10-somite stage. Then the prepared samples were stained with anti-BCAS2 (10414-1-AP, Proteintech), anti-β-catenin (M24002, Abmart), and anti-GFP (A11122, Invitrogen). Meanwhile, 4′,6-Diamidine-2′-phenylindole dihydrochloride (DAPI, 10236276001, Sigma) was used to label nuclei. Fluorescence imaging was performed using a Nikon A1R+ confocal microscope and all images were captured with the same settings.

Bimolecular fluorescence complementation assay

To construct the plasmids for Bimolecular fluorescence complementation assay (BiFC), BCAS2 was fused to the N-terminal half of yellow fluorescent protein (YN-BCAS2) and β-catenin to the C-terminal half (YC-β-catenin). YN-BCAS2 and YC-β-catenin were either individually or collectively transfected into HeLa cells. Fluorescence was detected 48 h after transfection using a Nikon A1R+confocal microscope.

Fluorescence recovery after photobleaching

BCAS2 and GFP tagged S37A-β-catenin were co-transfected into HeLa cells.

Fluorescence recovery after photobleaching (FRAP) assay was performed according to

previously reported methods.[39] Images were acquired by a META Zeiss 510 confocal

microscope.

RNA sequencing

Embryos were collected at the 10-somite stage and gently transferred into lysis buffer.

Reverse transcription was performed using a SMARTer Ultra Low RNA Kit (Clontech,

634437) directly from the cell lysates. The cDNA library was prepared using an

Advantage 2 PCR Kit (Clontech，639206) and then sequenced via the Illumina

sequencing platform (NovaSeq 6000). The difference in the number of alternative

splicing events between groups was analyzed using rMATS (version 4.1.0).

Reverse transcription PCR

Total RNA was isolated from wild-type and *bcas2* mutant embryos at the 10-somite

stage with MicroElute Total RNA kit (OMEGA, R6831-01), followed by reverse

transcription using ReverTra Ace qPCR RT Kit (Toyobo, FsQ-101). The cDNA was

amplified with primers listed in Supplemental Table 2.

Quantification and statistical analysis

Images were quantified with Image J. Statistical data were analyzed using Graph Pad

Prism. Comparisons between experimental groups were done using the Student's *t-test*.

Data are presented as mean ± standard deviation. Differences were considered

significant at *P < 0.05* and marked with "*", very significant at *P < 0.01* and marked

with "**".

Results

BCAS2 is necessary for primitive hematopoiesis

To confirm that *bcas2* is expressed in the posterior intermediate cell mass (ICM) where

primitive hematopoiesis occurs in zebrafish, we first examined the spatiotemporal

expression pattern of *bcas2* during zebrafish embryo development by performing whole

mount *in situ* hybridization (WISH). The results showed that *bcas2* was ubiquitously

expressed from 1-cell stage to 10-somite stage (Supplemental Figure 1A). Its

expression in the ICM became detectable at 18 hours post fertilization (hpf) and was

significantly elevated at 22 hpf (Figure 1A). We further observed that *bcas2* was co-

expressed with the primitive erythropoietic marker *gata1* in the ICM at 22 hpf by

fluorescence *in situ* hybridization (FISH) (Figure 1B). In contrast, *bcas2* was hardly

detectable in the ICM in *cloche*[-/-] mutants that lack both endothelial and hematopoietic

cells (Figure 1C). These results demonstrate a dynamic expression of *bcas2* in the ICM

and imply a potential role of this gene in primitive hematopoiesis.

To gain insight into the developmental function of *bcas2*, we employed

CRISPR/Cas9 system to generate *bcas2* mutants. Two mutant lines were obtained,

designated *bcas2Δ7* (with a 7-base deletion) and *bcas2Δ14* (with a 14-base deletion).

These mutations led to premature translation termination, which resulted in truncated

Bcas2 proteins lacking the C-terminal CC domains (Supplemental Figure 2A-B).

bcas2Δ7$^{+/-}$ and *bcas2Δ14$^{+/-}$* mutants were identified by T7 endonuclease I assay or

restriction enzyme analysis (FspI) (Supplemental Figure 2C). We found that nearly 85%

of the embryos derived from crossing *bcas2$^{+/-}$* males with *bcas2$^{+/-}$* females did not

develop to the cleavage stage (Supplemental Figure 2D). Only 3% of the living embryos

were homozygotes. In contrast, embryos obtained by crossing between wild-type males

and *bcas2$^{+/-}$* females were viable and showed normal morphology, with a

heterozygosity rate consistent with Mendelian inheritance. This could be explained by

male infertility as previously documented in *Bcas2* knockout mice.[15] Combining the

above findings, we propose that Bcas2 may have an evolutionarily conserved role in

spermatogenesis.

Given the difficulty of obtaining homozygous mutants, embryos lacking one copy

of *bcas2* gene were produced from crosses between heterozygous females and wild-

type males. We observed a significant decrease of *bcas2* expression in the ICM region

in *bcas2Δ7$^{+/-}$* or *Δ14$^{+/-}$* mutants, likely resulting from nonsense-mediated RNA decay

(Figure 1D). In line with a previous report,[17] a marked reduction in the expression of

the HSC marker *cmyb* and T-cell marker *rag1* was found in *bcas2Δ7$^{+/-}$* or *Δ14$^{+/-}$*

embryos at 5 hpf, indicating an essential role of *bcas2* in definitive hematopoiesis

(Supplemental Figure 3A-B).

To explore whether *bcas2* is required for primitive hematopoiesis, we first examined

the expression of primitive erythropoietic markers *gata1* and *hbbe3* in *bcas2Δ7*$^{+/-}$ and

Δ14$^{+/-}$ embryos at 22 hpf, and observed a marked decrease in the expression of these

genes in the mutants (Figure 1E-F). Surprisingly, *o*-dianisidine staining showed similar

hemoglobin contents in the *bcas2Δ7*$^{+/-}$ and *Δ14*$^{+/-}$ embryos at 48 hpf compared with

control embryos, suggesting that the defect in primitive hematopoiesis induced by

haploinsufficiency of *bcas2* was alleviated at later developmental stages. In order to

further explore the role of BCAS2 in primary hematopoiesis, we identified several

bcas2Δ14$^{-/-}$ mutants from about 100 embryos from the mating of *bcas2Δ14*$^{+/-}$ male and

female. These homozygous mutants display a severe decrease in hemoglobin (Figure

1G). Similarly, injection of a translation-blocking MO into wild-type embryos to

downregulate *bcas2* expression resulted in severe defects in erythropoiesis at 22 hpf

and 48 hpf (Supplemental Figure 4A-D). These results indicate that *bcas2* is

indispensable for primitive hematopoiesis in zebrafish. In addition, to induce the

deletion of Bcas2 in mammalian endothelial/hematopoietic cells, transgenic mice

expressing Cre recombinase under the control of the *Flk1* promoter were crossed to

Bcas2$^{F/F}$ animals. We found that red blood cells were eliminated in the yolk sac of

Bcas2$^{F/F}$*;Flk1-Cre* mice at E12.5 despite the presence of vessels (Figure 1H). Therefore,

Bcas2 has a conserved role in vertebrates to regulate primitive hematopoiesis.

bcas2 deficiency impairs hematopoietic progenitor differentiation

The decrease of primitive hematopoietic cells in *bcas2* deficient animals may be

attributed to a number of possible causes: excessive apoptosis, hampered proliferation

of hematopoietic cells, or impaired differentiation of hematopoietic progenitor cells. To

shed light on this issue, we first performed terminal deoxynucleotidyl transferase-

mediated dUTP nick end labeling (TUNEL) assay in *Tg(gata1:GFP)* embryos at the

10-somite stage to examine DNA fragmentations in apoptotic cells, and found no

obvious apoptotic signal in the *gata1*$^+$ hematopoietic cells in either *bcas2Δ14*$^{+/-}$

embryos or their wild-type siblings (Supplemental Figure 5A). Meanwhile, BrdU

incorporation assay revealed no significant difference in hematopoietic cell

proliferation between *bcas2Δ14*$^{+/-}$ mutants and their corresponding wild-types

(Supplemental Figure 5B-C). These data suggest that *bcas2* is dispensable for the

survival and proliferation of hematopoietic cells.

In the developing embryo, hemangioblasts are derived from the ventral mesoderm

at early somite stage, and then differentiate into both hematopoietic and endothelial

lineages.[40,41] To test whether *bcas2* functions in cell fate decision during primitive

hematopoietic cell development, the expression of hemangioblast markers *npas4l*, *scl*,

and *gata2* in *bcas2Δ14*$^{+/-}$ embryos was examined at the 1- to 2-somite stage. As shown

in Figure 2A, haploinsufficiency of *bcas2* did not affect the emergence of the

hemangioblast population. Then we extended our analysis to include the markers of

hematopoietic and endothelial progenitors. Consistent with the decrease in primitive

hematopoietic cells in *bcas2* deficient mutants, a marked reduced expression of

erythrocyte progenitor markers *gata1* and *hbbe3* was observed in the posterior lateral

mesoderm of *bcas2*[+/-] embryos at the 10-somite stage (Figure 2B). Interestingly, the

expression of myeloid progenitor marker *pu.1* was also dramatically decreased (Figure

2C). Moreover, overexpression of human BCAS2 enhanced the expression of *gata1* in

both wild-type and mutant embryos at the 10-somite stage (Figure 2D). In contrast, the

endothelial progenitor marker *fli1a* was expressed at a similar level in *bcas2*[+/-] embryos

as in wild-type animals (Figure 2E). Consistently, blood vessels appeared normal in

bcas2[+/-] mutants with *Tg(kdrl:GFP)* background at 54 hpf (Figure 2F). These data

provide convincing evidence that *bcas2* is required for the differentiation of the

hematopoietic lineage from hemangioblasts during primitive hematopoiesis.

BCAS2 functions in primitive hematopoiesis by activating Wnt signaling

Previous studies have shown that Wnt/β-catenin plays a key role in primitive

hematopoiesis.[27,28,42] As both BCAS2 and β-catenin-like 1 (CTNNBL1) are members

of the Prp19/CDC5L complex, which is a major building block of the spliceosome's

catalytic RNP core,[13] we speculate that BCAS2 may be a regulator of Wnt signaling

through interaction with β-catenin during hematopoiesis. To test our hypothesis, human

BCAS2 was overexpressed in HEK293T cells and MEFs. Ectopic expression of

BCAS2 enhanced the Wnt3a-induced expression of the TOPflash luciferase reporter in

a dose-dependent manner (Figure 3A-B). Importantly, Wnt3a-induced luciferase

activity in HEK293T cells could be effectively reduced by knockdown of *BCAS2* using

two shRNAs targeting different regions of human *BCAS2* (Figure 3C-D). Similar results

were also observed in *Bcas2*-cKO MEFs in the presence of tamoxifen (Figure 3E).

These findings support that BCAS2 promotes Wnt signaling activation.

To confirm that Wnt signaling was required for zebrafish embryonic hematopoiesis,

we induced the expression of canonical Wnt inhibitor Dkk1 by heat-shocking

Tg(hsp70l:dkk1-GFP)^(w32) embryos at the bud stage.[43] As expected, diminished

expression of *gata1* and *hbbe3* was detected in the resulting embryos at the 10-somite

stage (Figure 3F-G). To validate that *bcas2* functions in primitive hematopoiesis via

Wnt/β-catenin signaling, the expression pattern of β-catenin was examined in *bcas2*

morphants with *Tg(gata1:GFP)* background at the 10-somite stage by

immunofluorescent staining. The signals of both cytoplasmic and nuclear β-catenin

were substantially decreased in hematopoietic progenitor cells (Figure 3H). Moreover,

overexpression of ΔN-β-catenin, a constitutively active form of β-catenin, effectively

restored the expression of *hbbe3* in *bcas2* morphants and mutants (Figure 3I-J). All

these data suggest that BCAS2 functions in primitive hematopoiesis by regulating

Wnt/β-catenin signaling.

BCAS2 promotes β-catenin nuclear accumulation independently of protein

stability regulation

To investigate how BCAS2 regulates Wnt/β-catenin signaling, HEK293T cells were

treated with LiCl, a canonical Wnt agonist that inhibits GSK-3β activity and stabilizes

cytosolic β-catenin.[44] The results showed that TOPflash activity was significantly

elevated in LiCl-treated cells (Figure 4A). BCAS2 overexpression further upregulated,

whereas shRNA-mediated knockdown of *BCAS2* downregulated LiCl-induced

TOPflash activity (Figure 4A-B). Likewise, HEK293T cells transfected with S37A-β-

catenin, a constitutively active form of β-catenin that is resistant to GSK-3β-mediated

degradation, [45] displayed a much higher level of TOPflash activity, which was reduced

by *BCAS2* knockdown (Figure 4C). These results strongly imply that BCAS2 regulates

Wnt signaling downstream of β-catenin stability control.

To test the above hypothesis, we evaluated nuclear β-catenin level by performing

immunofluorescence staining and immunoblotting experiments. Upon tamoxifen

exposure, nuclear accumulation of β-catenin induced by LiCl was greatly inhibited in

Bcas2-cKO MEFs, while nuclear/cytoplasmic fractionation suggested that cytoplasmic

β-catenin level remained relatively unchanged (Figure 4D-E). Similarly, silencing

BCAS2 with shRNA led to reduced nuclear β-catenin in the human colon cancer cell

line SW480, in which β-catenin was activated because of mutations in the adenomatous

polyposis coli protein (APC), an integral component of the β-catenin destruction

complex (Figure 4F).[46] Next, MG132, a proteasome inhibitor, was applied to activate

Wnt/β-catenin signaling in *Bcas2*-cKO MEFs by inhibiting β-catenin degradation. In

the absence of tamoxifen and MG132, endogenous β-catenin was localized almost

exclusively in the cytoplasm; MG132 treatment dramatically triggered β-catenin

accumulation in the nuclei (Figure 4G). However, in *Bcas2*-cKO MEFs exposed to

tamoxifen, MG132 treatment was not able to induce nuclear accumulation of β-catenin

(Figure 4G). These findings indicate that BCAS2 promotes β-catenin nuclear accumulation in a manner that is independent of β-catenin stability regulation.

BCAS2 sequesters β-catenin within the nucleus

In addition to be affected by protein stability, the nuclear level of β-catenin is also fine-tuned by the opposing actions of nuclear export and import.[47-49] To examine the effect of BCAS2 on the nuclear import and export of β-catenin, fluorescent recovery after photobleaching (FRAP) experiments was carried out in HeLa cells expressing GFP-tagged S37A-β-catenin. After photobleaching the nucleus, no significant difference was found in the recovery of nuclear GFP signals between the cells with and without overexpression of BCAS2 (Supplemental Figures 6A, A', and C), suggesting that BCAS2 does not regulate β-catenin nuclear import. Conversely, after photobleaching the cytoplasm, BCAS2 overexpressed cells showed a much slower recovery of cytoplasmic fluorescence (Supplemental Figures 6B, B', and C), indicating that BCAS2 inhibits β-catenin nuclear export.

It has been suggested that the nuclear exit of β-catenin can be either dependent or independent on CRM1, a major nuclear export receptor.[50] To shed light on the mechanism underlying BCAS2 mediated β-catenin nuclear retention, we treated *Bcas2*-cKO MEFs with the CRM1-specific export inhibitor leptomycin B (LMB).[51] Regardless of the presence or absence of endogenous BCAS2, LMB treatment could effectively increase the level of β-catenin in the nucleus (Figure 5A). Consistently,

treatment of LMB was able to rescue the impaired nuclear accumulation of β-catenin

in BCAS2-deficient SW480 cells (Figure 5B). Moreover, when *bcas2* morphants in *Tg*

(gata1:GFP) background were treated with LMB from bud stage to 10 somite stage,

the level of nuclear β-Catenin was partially recovered compared with that of control

embryos (Figure 5C). Importantly, the expression of *gata1* was also restored in *bcas2*

mutants upon LMB treatment (Figure 5D). Taken together, these findings suggest that

BCAS2 negatively regulates CRM1-mediated nuclear export of β-catenin.

BCAS2 directly interacts with β-catenin in the nucleus

To investigate whether BCAS2 inhibits the nuclear export of β-catenin through physical

binding, HEK293T cells were co-transfected with Flag-tagged β-catenin and HA-

tagged BCAS2 constructs. Co-immunoprecipitation (Co-IP) experiments showed that

Flag-β-catenin was precipitated with HA-BCAS2 as well as endogenous BCAS2,

indicating an interaction between these two proteins (Figure 6A-B). In addition, the

interaction was enhanced upon Wnt3a stimulation (Figure 6C). Given that Wnt ligand

stimulation ultimately induces β-catenin nuclear accumulation, this enhanced

interaction implies that BCAS2 associates with β-catenin within the nucleus. Therefore,

we performed the bimolecular fluorescence complementation (BiFC) assay to visualize

the interaction of BCAS2 and β-catenin in living cells. In this assay, the N-terminal

fragment of yellow fluorescent protein (YFP) was fused to BCAS2 (YN-BCAS2),

while the C-terminal fragment was fused to β-catenin (YC-β-catenin) (Figure 6D). As

expected, the YFP fluorescence was specifically observed in the nucleus (Figure 6E).

Previous studies have divided the β-catenin protein into three distinct domains,

including the N-terminal domain (residues 1–133), the central domain with 12 ARM

repeats (residues 134–670), and the C-terminal domain (residues 671–781).[52] To

identify the BCAS2 binding site, constructs expressing various truncated forms of β-

catenin were generated and co-transfected with BCAS2 into HEK293T cells (Figure

6F). Co-IP assays revealed that deletion of the N-terminal or C-terminal domain of β-

catenin did not alter the interaction between β-catenin and BCAS2 (Figure 6G). In

contrast, when the ARM repeats 1-12 of β-catenin were deleted, the resulting deletion

mutant showed virtually no interaction with BCAS2 (Figure 6G). GST pull-down assay

also demonstrated a direct interaction between BCAS2 and the ARM repeats of β-

catenin (Figure 6H). These results indicate that BCAS2 physically binds to the ARM

repeats of β-catenin. Furthermore, we found that the ARM repeats 9-12, but not 1-8,

bound to BCAS2 (Figure 6G).

BCAS2 enhances β-catenin nuclear accumulation through its CC domains

To determine which domain of BCAS2 binds to β-catenin, we constructed a series of

deletion mutants of BCAS2 (Figure 7A). Notably, we observed that among these

truncated mutants, only the one lacking both CC1 and CC2 domains lost the ability to

interact with β-catenin (Figure 7B). Moreover, these two CC domains alone or together

could interact with β-catenin (Figure 7C). Therefore, we conclude that BCAS2 binds to

β-catenin via its CC domains.

We next examined whether the CC domains is required for BCAS2 to promote

Wnt/β-catenin signaling. As shown in Figure 7D, overexpression of BCAS2 without

the CC domains failed to increase LiCl-induced TOPflash activity in HEK-293T cells.

Likewise, overexpression of the full-length BCAS2, but not BCAS2 lacking the CC

domains, could restore the nuclear accumulation of β-catenin in *bcas2* morphants and

gata1 expression in *bcas2* mutants (Figure 7E-F). Collectively, these findings indicate

that BCAS2 positively regulates Wnt signaling through sequestering β-catenin within

the nucleus via its CC domains during primitive hematopoiesis.

As BCAS2 is involved in the Prp19-CDC5L spliceosome complex that regulates

RNA splicing during spermiogenesis, neurogenesis, and definitive hematopoiesis,[15,17]

we wondered if this protein participates in primitive hematopoiesis via mRNA

alternative splicing. To this end, we performed RNA sequencing of 10-somite stage

embryos to identify abnormal events in alternative splicing in *bcas2Δ14$^{+/-}$* mutants.

However, upon haploinsufficiency of *bcas2*, neither the number of five major types of

alternative splicing events, nor the typical forms of alternative splicing were

significantly affected (Supplemental Figure 7A-B). Additionally, haploinsufficiency of

bcas2 did not result in the alternative splicing of *mdm4* that predisposes cells to undergo

p53-mediated apoptosis in definitive hematopoiesis, as reported previously by Yu *et al.*

(Supplemental Figure 7C).[17,53] Furthermore, the splicing efficiency of β-catenin pre-

mRNA remained almost unchanged in *bcas2Δ14$^{+/-}$* mutants (Supplemental Figure 7C).

These results demonstrate that the defects in primitive hematopoiesis of *bcas2Δ14*[+/-]

mutants are independent of the regulatory role of Bcas2 in pre-mRNA splicing.

Discussion

BCAS2 is a 26-kDa nuclear protein involved in a multitude of developmental processes,

such as Drosophila wing development, dendritic growth, and spermatogenesis.[8,14-]

[16,54,55] In our study, we generated *bcas2* knockout zebrafish. The heterozygotes not only

showed male infertility, resembling the phenotype of *Bcas2* germ cell-specific knockout

mice reported previously,[15] but also exhibited impaired definitive hematopoiesis,

consistent with the earlier study.[17] Importantly, we found a marked decrease in the

expression of the primitive erythroid progenitor markers *gata1* and *hbbe3* in these

heterozygous mutants, which was rescued by overexpression of BCAS2. Moreover, the

defective primitive hematopoiesis in zebrafish mutants was phenocopied in

hemangioblast-specific *Bcas2* knockout mice. While the reason(s) for the discrepancy

between our data and the observations made by Yu *et al.* regarding the role of *bcas2* in

the development of primitive erythroid and myeloid cells remains to be determined,[17]

our findings in zebrafish and mouse embryos provide solid evidence that BCAS2 plays

a conserved role in primitive hematopoiesis.

As demonstrated in previous studies, BCAS2 is involved in various developmental

events by regulating pre-mRNA splicing.[14,15,17,30,31] However, our data showed that

haploinsufficiency of *bcas2* did not affect alternative splicing during primitive

hematopoiesis. These results imply that one copy of the *bcas2* gene is sufficient to

support mRNA splicing in zebrafish. Instead, we find that Bcas2 promotes primitive

hematopoiesis by sequestering β-catenin within the nucleus. It has been reported that

the *bcas2* deletion in zebrafish embryos induces alternative splicing of *mdm4* that

predisposes cells to undergo p53-mediated apoptosis in HSPCs during definitive

hematopoiesis.[17] Intriguingly, we found that the loss of one copy of *bcas2* gene in

zebrafish also resulted in severe impairment of HSPCs and their derivatives. It is

possible that Bcas2 might also have a role in definitive hematopoiesis independent of

its splicing regulatory function.

For the past decades, given the contradictory conclusions obtained from various *in*

vitro and *in vivo* studies, the function of Wnt/β-catenin signaling in primitive

hematopoiesis remains elusive and controversial.[26-29] In the present study, we have

provided several lines of evidence supporting that Wnt/β-catenin signaling positively

regulates primitive hematopoiesis: (1) Inhibition of Wnt/β-catenin by overexpression

of the canonical Wnt inhibitor Dkk1 disrupts the formation of erythrocyte progenitors

at the 10-somite stage. (2) Defects in primitive hematopoiesis in *bcas2* morphants and

mutants are readily restored by overexpression of ΔN-β-catenin, a constitutively active

β-catenin. (3) Overexpression of the full-length BCAS2, but not the CC domain-deleted

BCAS2, restores the formation of the primitive erythroid progenitor in *bcas2* mutants.

(4) BCAS2 overexpression enhances the development of primitive blood cells in wild-

type embryos. All these data suggest that BCAS2-mediated Wnt/β-catenin signal

activation is necessary for primitive hematopoiesis.

The nuclear translocation of β-catenin is a key event in canonical Wnt signal

activation. However, the precise mechanism for the nuclear-cytoplasmic shuttling of β-

catenin is not completely understood.[50] It has been demonstrated that β-catenin can

transit into and out of nucleus using distinct molecular interactions.[56-58] In some

physiological contexts, β-catenin binds to APC and/or Axin proteins, which are actively

co-exported from the nucleus by the nuclear export factor CRM1.[59-61] Furthermore,

CRM1 can bind directly to and function as an efficient nuclear exporter for β-catenin

independently of APC.[62] Our FRAP experiments reveal that BCAS2 is not involved in

β-catenin nuclear import, but inhibits the nuclear export of β-catenin. Treatment of

LMB, a CRM1-specific export inhibitor, could rescue the defect in β-catenin nuclear

accumulation in BCAS2-deficient cells, implying that BCAS2 has an inhibitory role in

CRM1-mediated nuclear export of β-catenin. We further show that BCAS2 interacts

directly with β-catenin and enhances β-catenin nuclear accumulation through its CC

domains. In the future, it will be interesting to investigate whether BCAS2 competes

with APC, Axin, and/or CRM1 for binding to β-catenin.

In summary, we uncover a novel role of BCAS2 in primitive hematopoiesis through

enhancing the nuclear retention of β-catenin. Our study provides new insights into the

mechanism of BCAS2-mediated Wnt signal activation during primitive hematopoiesis.

Given that BCAS2 and Wnt signaling are well documented to contribute to cancer

development,[63-67] it is appealing to further explore whether our findings can be applied

to future cancer research.

Acknowledgements

We acknowledge the financial support of the National Natural Science Foundation of

China (32025014 and 32330029) and the National Key Research and Development

Program of China (2018YFA0800200 and 2020YFA0804000).

Conflict-of-interest disclosure

The authors declare no competing financial interests.

References

1.   Galloway JL, Zon LI. Ontogeny of hematopoiesis: examining the emergence of

hematopoietic cells in the vertebrate embryo. *Curr Top Dev Biol*. 2003;53:139-158.

2.   Murry CE, Keller G. Differentiation of embryonic stem cells to clinically relevant

populations: lessons from embryonic development. *Cell*. 2008;132(4):661-680.

3.   Ferkowicz MJ, Yoder MC. Blood island formation: longstanding observations and

modern interpretations. *Exp Hematol*. 2005;33(9):1041-1047.

4.   Palis J. Hematopoietic stem cell-independent hematopoiesis: emergence of

erythroid, megakaryocyte, and myeloid potential in the mammalian embryo. *FEBS Lett*.

2016;590(22):3965-3974.

5.   Paik EJ, Zon LI. Hematopoietic development in the zebrafish. *Int J Dev Biol*.

2010;54(6-7):1127-1137.

6.   Detrich HW, 3rd, Kieran MW, Chan FY, et al. Intraembryonic hematopoietic cell

migration during vertebrate development. *Proc Natl Acad Sci U S A*.

1995;92(23):10713-10717.

7.   Leung AY, Mendenhall EM, Kwan TT, et al. Characterization of expanded

intermediate cell mass in zebrafish chordin morphant embryos. *Dev Biol*.

2005;277(1):235-254.

8.   Kuo PC, Tsao YP, Chang HW, et al. Breast cancer amplified sequence 2, a novel

negative regulator of the p53 tumor suppressor. *Cancer Res*. 2009;69(23):8877-8885.

9.   Neubauer G, King A, Rappsilber J, et al. Mass spectrometry and EST-database

searching allows characterization of the multi-protein spliceosome complex. *Nature*

Genetics. 1998;20(1):46-50.

10. Nagasaki K, Maass N, Manabe T, et al. Identification of a novel gene, DAM1,

amplified at chromosome 1p13.3-21 region in human breast cancer cell lines. *Cancer*

Letters. 1999;140(1-2):219-226.

11. Qi C, Zhu YT, Chang J, Yeldandi AV, Rao MS, Zhu YJ. Potentiation of estrogen

receptor transcriptional activity by breast cancer amplified sequence 2. *Biochem*

Biophys Res Commun. 2005;328(2):393-398.

12. Ajuh P, Kuster B, Panov K, Zomerdijk JC, Mann M, Lamond AI. Functional

analysis of the human CDC5L complex and identification of its components by mass

spectrometry. *EMBO J*. 2000;19(23):6569-6581.

13. Grote M, Wolf E, Will CL, et al. Molecular architecture of the human

Prp19/CDC5L complex. *Mol Cell Biol*. 2010;30(9):2105-2119.

14. Chen PH, Lee CI, Weng YT, et al. BCAS2 is essential for Drosophila viability and

functions in pre-mRNA splicing. *RNA*. 2013;19(2):208-218.

15. Liu W, Wang F, Xu Q, et al. BCAS2 is involved in alternative mRNA splicing in

spermatogonia and the transition to meiosis. *Nat Commun*. 2017;8:14182.

16. Xu Q, Wang F, Xiang Y, et al. Maternal BCAS2 protects genomic integrity in

mouse early embryonic development. *Development*. 2015;142(22):3943-3953.

17. Yu S, Jiang T, Jia D, et al. BCAS2 is essential for hematopoietic stem and

progenitor cell maintenance during zebrafish embryogenesis. *Blood*. 2019;133(8):805-

815.

18. Bissinger R, Lang E, Gonzalez-Menendez I, et al. Genetic deficiency of the tumor

suppressor protein p53 influences erythrocyte survival. *Apoptosis*. 2018;23(11-12):641-

650.

19. Yang S, Cao S, Xu X, et al. adducin 1 is essential for the survival of erythroid

precursors via regulating p53 transcription in zebrafish. *iScience*. 2023;26(9):107516.

20. Stanic K, Reig G, Figueroa RJ, et al. The Reprimo gene family member, reprimo-

like (rprml), is required for blood development in embryonic zebrafish. *Sci Rep*.

2019;9(1):7131.

21. Richter J, Traver D, Willert K. The role of Wnt signaling in hematopoietic stem

cell development. *Critical Reviews in Biochemistry and Molecular Biology*.

2017;52(4):414-424.

22. Krimpenfort RA, Nethe M. Canonical Wnt: a safeguard and threat for

erythropoiesis. *Blood Advances*. 2021;5(18):3726-3735.

23. Kokolus K, Nemeth MJ. Non-canonical Wnt signaling pathways in hematopoiesis.

Immunologic Research. 2010;46(1-3):155-164.

24. Tarafdar A, Dobbin E, Corrigan P, Freeburn R, Wheadon H. Canonical Wnt

signaling promotes early hematopoietic progenitor formation and erythroid

specification during embryonic stem cell differentiation. *PLoS One*. 2013;8(11):e81030.

25. Nostro MC, Cheng X, Keller GM, Gadue P. Wnt, activin, and BMP signaling

regulate distinct stages in the developmental pathway from embryonic stem cells to

blood. *Cell Stem Cell*. 2008;2(1):60-71.

26. Sturgeon CM, Ditadi A, Awong G, Kennedy M, Keller G. Wnt signaling controls

the specification of definitive and primitive hematopoiesis from human pluripotent

stem cells. *Nature Biotechnology*. 2014;32(6):554-+.

27. Tran HT, Sekkali B, Van Imschoot G, Janssens S, Vleminckx K. Wnt/beta-catenin

signaling is involved in the induction and maintenance of primitive hematopoiesis in

the vertebrate embryo. *Proc Natl Acad Sci U S A*. 2010;107(37):16160-16165.

28. Lengerke C, Schmitt S, Bowman TV, et al. BMP and Wnt specify hematopoietic

fate by activation of the Cdx-Hox pathway. *Cell Stem Cell*. 2008;2(1):72-82.

29. Paluru P, Hudock KM, Cheng X, et al. The negative impact of Wnt signaling on

megakaryocyte and primitive erythroid progenitors derived from human embryonic

stem cells. *Stem Cell Research*. 2014;12(2):441-451.

30. Chen HH, Lu HY, Chang CH, et al. Breast carcinoma-amplified sequence 2

regulates adult neurogenesis via β-catenin. *Stem Cell Research & Therapy*. 2022;13(1).

31. Huang CW, Chen YW, Lin YR, et al. Conditional Knockout of Breast Carcinoma

Amplified Sequence 2 (BCAS2) in Mouse Forebrain Causes Dendritic Malformation

via β-catenin. *Scientific Reports*. 2016;6.

32. Chang N, Sun C, Gao L, et al. Genome editing with RNA-guided Cas9 nuclease in

zebrafish embryos. *Cell Res*. 2013;23(4):465-472.

33. Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 are required for

mesendoderm induction by transforming growth factor-beta/nodal signals in zebrafish.

J Biol Chem. 2008;283(4):2418-2426.

34. Welten MC, de Haan SB, van den Boogert N, et al. ZebraFISH: fluorescent in situ

hybridization protocol and three-dimensional imaging of gene expression patterns.

Zebrafish. 2006;3(4):465-476.

35. Wallingford MC, Giachelli CM. Loss of PiT-1 results in abnormal endocytosis in

the yolk sac visceral endoderm. *Mech Dev*. 2014;133:189-202.

36. Lieschke GJ, Oates AC, Crowhurst MO, Ward AC, Layton JE. Morphologic and

functional characterization of granulocytes and macrophages in embryonic and adult

zebrafish. *Blood*. 2001;98(10):3087-3096.

37. Wei S, Dai MM, Liu ZT, et al. The guanine nucleotide exchange factor Net1

facilitates the specification of dorsal cell fates in zebrafish embryos by promoting

maternal β-catenin activation. *Cell Research*. 2017;27(2):202-225.

38. Yang SY, Ning GZ, Hou YM, et al. Myoneurin regulates BMP signaling by

competing with Ppm1a for Smad binding. *Iscience*. 2022;25(6).

39. Schmierer B, Hill CS. Kinetic analysis of Smad nucleocytoplasmic shuttling

reveals a mechanism for transforming growth factor beta-dependent nuclear

accumulation of Smads. *Mol Cell Biol*. 2005;25(22):9845-9858.

40. Vogeli KM, Jin SW, Martin GR, Stainier DY. A common progenitor for

haematopoietic and endothelial lineages in the zebrafish gastrula. *Nature*.

2006;443(7109):337-339.

41. Reischauer S, Stone OA, Villasenor A, et al. Cloche is a bHLH-PAS transcription

factor that drives haemato-vascular specification. *Nature*. 2016;535(7611):294-298.

42. Sun H, Wang Y, Zhang J, et al. CFTR mutation enhances Dishevelled degradation

and results in impairment of Wnt-dependent hematopoiesis. *Cell Death Dis*.

2018;9(3):275.

43. Stoick-Cooper CL, Weidinger G, Riehle KJ, et al. Distinct Wnt signaling pathways

have opposing roles in appendage regeneration. *Development*. 2007;134(3):479-489.

44. Meijer L, Skaltsounis AL, Magiatis P, et al. GSK-3-selective inhibitors derived

from Tyrian purple indirubins. *Chem Biol*. 2003;10(12):1255-1266.

45. Easwaran V, Song V, Polakis P, Byers S. The ubiquitin-proteasome pathway and

serine kinase activity modulate adenomatous polyposis coli protein-mediated

regulation of beta-catenin-lymphocyte enhancer-binding factor signaling. *J Biol Chem*.

1999;274(23):16641-16645.

46. Rosin-Arbesfeld R, Cliffe A, Brabletz T, Bienz M. Nuclear export of the APC

tumour suppressor controls beta-catenin function in transcription. *EMBO J*.

2003;22(5):1101-1113.

47. Lu Y, Xie S, Zhang W, et al. Twa1/Gid8 is a beta-catenin nuclear retention factor

in Wnt signaling and colorectal tumorigenesis. *Cell Res*. 2017;27(12):1422-1440.

48. Henderson BR, Fagotto F. The ins and outs of APC and beta-catenin nuclear

transport. *EMBO Rep*. 2002;3(9):834-839.

49. Henderson BR. Nuclear-cytoplasmic shuttling of APC regulates beta-catenin

subcellular localization and turnover. *Nat Cell Biol*. 2000;2(9):653-660.

50. Xu L, Massague J. Nucleocytoplasmic shuttling of signal transducers. *Nat Rev Mol*

Cell Biol. 2004;5(3):209-219.

51. Wolff B, Sanglier JJ, Wang Y. Leptomycin B is an inhibitor of nuclear export:

inhibition of nucleo-cytoplasmic translocation of the human immunodeficiency virus

type 1 (HIV-1) Rev protein and Rev-dependent mRNA. *Chem Biol*. 1997;4(2):139-147.

52. Dimitrova YN, Li J, Lee YT, et al. Direct ubiquitination of beta-catenin by Siah-1

and regulation by the exchange factor TBL1. *J Biol Chem*. 2010;285(18):13507-13516.

53. Rallapalli R, Strachan G, Cho B, Mercer WE, Hall DJ. A novel MDMX transcript

expressed in a variety of transformed cell lines encodes a truncated protein with potent

p53 repressive activity. *Journal of Biological Chemistry*. 1999;274(12):8299-8308.

54. Huang CW, Chen YW, Lin YR, et al. Conditional Knockout of Breast Carcinoma

Amplified Sequence 2 (BCAS2) in Mouse Forebrain Causes Dendritic Malformation via beta-catenin. *Sci Rep*. 2016;6:34927.

55. Zhang J, Liu W, Li G, et al. BCAS2 is involved in alternative splicing and mouse oocyte development. *FASEB J*. 2022;36(2):e22128.

56. Koike M, Kose S, Furuta M, et al. β-catenin shows an overlapping sequence requirement but distinct molecular interactions for its bidirectional passage through nuclear pores. *Journal of Biological Chemistry*. 2004;279(32):34038-34047.

57. Mis M, O'Brien S, Steinhart Z, et al. IPO11 mediates βcatenin nuclear import in a subset of colorectal cancers. *Journal of Cell Biology*. 2020;219(2).

58. Hendriksen J, Fagotto F, van der Velde H, van Schie M, Noordermeer J, Fornerod M. RanBP3 enhances nuclear export of active β-catenin independently of CRM1. *Journal of Cell Biology*. 2005;171(5):785-797.

59. Henderson BR. Nuclear-cytoplasmic shuttling of APC regulates β-catenin subcellular localization and turnover. *Nature Cell Biology*. 2000;2(9):653-660.

60. Rosin-Arbesfeld R, Townsley F, Bienz M. The APC tumour suppressor has a nuclear export function. *Nature*. 2000;406(6799):1009-1012.

61. Cong F, Varmus H. Nuclear-cytoplasmic shuttling of Axin regulates subcellular localization of β-catenin. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;101(9):2882-2887.

62. Ki H, Oh M, Chung SW, Kim K. β-Catenin can bind directly to CRM1 independently of adenomatous polyposis coli, which affects its nuclear localization and

LEF-1/β-catenin-dependent gene expression. *Cell Biology International*. 2008;32(4):394-400.

63. Murillo-Garzón V, Kypta R. WNT signalling in prostate cancer. *Nature Reviews Urology*. 2017;14(11):683-696.

64. Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. *Oncogene*. 2017;36(11):1461-1473.

65. Yu FY, Yu CH, Li FF, et al. Wnt/β-catenin signaling in cancers and targeted therapies. *Signal Transduction and Targeted Therapy*. 2021;6(1).

66. Salmerón-Hernández A, Noriega-Reyes MY, Jordan A, Baranda-Avila N, Langley E. BCAS2 Enhances Carcinogenic Effects of Estrogen Receptor Alpha in Breast Cancer Cells. *International Journal of Molecular Sciences*. 2019;20(4).

67. Wang LP, Chen TY, Kang CK, Huang HP, Chen SL. BCAS2, a protein enriched in advanced prostate cancer, interacts with NBS1 to enhance DNA double-strand break repair. *British Journal of Cancer*. 2020;123(12):1796-1807.

736   **Figures and figure legends**

737   Figure 1

Figure 1. *bcas2* is expressed in the ICM and required for primitive hematopoiesis.

(A) WISH assay showing *bcas2* expression in the ICM at the 18-somite stage and 22

hpf. The dotted lines represent the section position and the black arrowheads indicate

the ICM region. n, notochord. (B) Double FISH assay showing the expression pattern

of *bcas2* and *gata1* in the ICM at 22 hpf. Scale bar, 50 μm. (C-D) Comparison of *bcas2*

expression in *cloche* mutants (C) or *bcas2* heterozygous mutants (D) along with their

corresponding siblings. (E-F) Expression analysis of *gata1* and *hbbe3* in *bcas2Δ7*$^{+/-}$ and

bcas2Δ14$^{+/-}$ embryos. (G) Hemoglobin detection using *o*-dianisidine staining in *bcas2*

homozygous mutant at 36 and 48 hpf. (H) Representative images of yolk sac from the

hemangioblast-specific *Bcas2* knockout mice and their siblings. *Bcas2*$^{F/F}$ females were

crossed with *Bcas2*$^{F/+}$*;Flk1-Cre* males to induce the deletion of *Bcas2* in

hemangioblasts. Scale bars, 1 mm.

765   Figure 2

Figure 2. *bcas2* is required for hematopoietic progenitor differentiation. (A-C)

Expression analysis of hemangioblast markers *npas4l*, *scl, gata2* (A), erythroid

progenitor markers *gata1*, *hbbe3* (B), and myeloid marker *pu.1* (C) in *bcas2Δ14*$^{+/-}$

embryos and their wild-type siblings at indicated stages. (D) Expression changes of

gata1 in *bcas2Δ14*$^{+/-}$ embryos overexpressing BCAS2 at the 10-somite stage. The

indicated embryos were injected with or without 300 pg of human BCAS2 mRNA at

the one-cell stage. (E) Expression of endothelial marker *fli1a* in *bcas2Δ14*$^{+/-}$ and sibling

embryos at the 10-somite stage. (G) Confocal imaging of *bcas2Δ14*$^{+/-}$ and control

sibling *Tg(kdrl:GFP)* embryos at 54 hpf. Scale bars, 500 μm.

798   Figure 3

Figure 3. BCAS2 promotes primitive hematopoiesis via activating Wnt signaling.

(A-B) Overexpression of BCAS2 increases Wnt3a-induced TOPflash activity in

HEK293T cells (A) and MEFs (B). Different amounts of plasmid expressing BCAS2

(0, 80, 160, or 320 ng/well) were transfected into cells, together with the super-

TOPflash luciferase and Renilla luciferase vectors. After 36 h of transfection, cells were

treated with or without Wnt3a CM for 12 h and harvested for luciferase assays. *$P <$

0.05; **$P < 0.01$ (Student's *t*-test). (C-E) The Wnt3a-induced TOPflash activity is

decreased in BCAS2-deficient cells. HEK293T Cells were co-transfected with shRNA

plasmids, along with indicated plasmids, and harvested for western blot analysis (C) or

luciferase reporter assay (D). *Bcas2*-cKO MEFs prepared from *Bcas2$^{F/F}$* mouse

embryos were incubated in medium containing 100 μM tamoxifen for 72 h and then

subjected to western blotting and luciferase reporter assay (E). *$P < 0.05$; **$P < 0.01$

(Student's *t*-test). (F-G) Expression analysis of *gata1* (F) and *hbbe3* (G) in

Tg(hsp70l:dkk1b-GFP) embryos after heat shock at 16 hpf. (H) Immunofluorescence

staining of β-catenin in *Tg(gata1:GFP)* embryos at 16 hpf. The embryos were injected

with 8 ng of the indicated MO at the one-cell stage and collected at the 10-somite stage.

The dotted lines show the GFP-positive hematopoietic progenitor cells. Scale bars, 5

μm. (I-J) Expression of *hbbe3* in *bcas2* morphants (I) and *bcas2Δ14$^{+/-}$* mutants (J)

overexpressing ΔN-β-catenin. Embryos were injected with the indicated MO together

with *ΔN-β-cateni*n mRNA at the one-cell stage and harvested at the 10-somite stage for

in situ hybridization.

Figure 4

Figure 4. BCAS2 is essential for β-catenin nuclear accumulation. (A-C) BCAS2

enhances LiCl-induced TOPflash activity in HEK293T cells. Cells were transfected

with BCAS2 expression plasmids (A), shRNA plasmids (B), or S37A-β-catenin

expression plasmids (C), together with the TOPflash luciferase and Renilla luciferase

vectors. After transfection, cells were subsequently treated with or without 100 ng/ml

LiCl for 12 h and assayed for luciferase activity. *P < 0.05; **P < 0.01 (Student's *t-*

test). (D-E) *Bcas2*-cKO MEFs were incubated with tamoxifen for 24 h and then treated

with or without 100 ng/ml LiCl. The nuclear accumulation of β-catenin was analyzed

using immunofluorescence (D) and western blotting (n=3) (E) Scale bars, 10 μm. (F)

SW480 cells were transfected with the indicated shRNA constructs, and the endogenous

β-catenin protein was detected using immunofluorescence 48 h after transfection. The

expression of GFP served as a transfection control. The arrowheads indicate the cells

transfected with indicated shRNA constructs. Scale bars, 10 μm. (G) *Bcas2*-cKO MEFs

were cultured in the presence of tamoxifen for 24 h and then treated with 20 μM MG132

for 6 h. The expression of BCAS2 and β-catenin was measured by immunofluorescence.

Scale bars, 10 μm.

854   Figure 5

Figure 5. BCAS2 functions in CRM1-mediated nuclear export of β-catenin. (A)

Tamoxifen-treated *Bcas2*-cKO MEFs were incubated with 20 nM LMB for 3 h. The

expression of Bcas2 and β-catenin was analyzed using immunofluorescence. The

arrowheads show the cells with nuclear β-catenin accumulation. Scale bars, 10 μm. (B)

SW480 cells were transfected with the indicated shRNA constructs and then treated

with LMB for 3 h before immunostaining. GFP was regarded as a transfection control.

The arrowheads indicate the transfected cells. Scale bars, 10 μm. (C)

Immunofluorescence staining of β-catenin in *bcas2* morphants with *Tg(gata1:GFP)*

background. Embryos were exposed to 20 nM LMB from the bud stage. The dotted

lines indicate the GFP-positive hematopoietic progenitor cells. Scale bars, 5 μm. (D)

bcas2Δ14$^{+/-}$ embryos were treated with 20 nM LMB for 6 h and then subjected to WISH

assay to analyze the expression of *gata1* at the indicated stages.

884   Figure 6

Figure 6. BCAS2 interacts with β-catenin. (A-C) Flag-tagged β-catenin was co-

transfected with or without HA-tagged BCAS2 into HEK293T cells. Cell lysates were

immunoprecipitated using anti-Flag antibody. Eluted proteins were analyzed by

western blotting using indicated antibodies. In panel C, for Wnt signaling activation,

cells were treated with Wnt3a CM for 5 h before harvest. (D-E) YN-BCAS2 and YC-

β-catenin were either individually or collectively transfected into HeLa cells. The

expression of YN-BCAS2 and YC-β-catenin was analyzed with anti-GFP antibody (D).

The reconstituted YFP fluorescence in living cells was detected by confocal laser

scanning microscopy with excitation at 488 nm (E). Scale bars, 10 μm. (F) Schematics

of full-length and deletion mutants of β-catenin. (G) HEK293T cells were transfected

with HA-tagged BCAS2 and Flag-tagged deletion mutants of β-catenin. Cell lysates

were then immunoprecipitated using anti-Flag antibody followed by western blot

analysis. (H) GST pull-down assays were performed using bacterially expressed GST,

GST-ARM1-12, and His-BCAS2.

911 Figure 7

921 **Figure 7. BCAS2 sequesters β-catenin in the nucleus via its CC domains.** (A)

922 Schematics of full length and deletion mutants of BCAS2. (B-C) HEK293T cells were

923 transfected with Flag-β-catenin and indicated deletion mutants of BCAS2. Cell lysates

924 were subjected to immunoprecipitation with anti-Flag antibody. Eluted proteins were

925 immunoblotted using anti-HA (B) or anti-GFP antibodies (C) for BCAS2 detection. (D)

926 HEK293T cells transfected with the indicated plasmids were treated with 100 ng/ml

927 LiCl for 12 h, and then subjected to luciferase assay. Note that overexpression of

928 BCAS2 without the CC domains failed to increase LiCl-induced TOPflash activity.

929 Data represent the mean ± SD of three independent experiments. ns, not significant;

930 **P < 0.01 (Student's *t-test*). (E) Immunofluorescence staining of β-catenin in

931 *Tg(gata1:GFP)* embryos injected with 8 ng *bcas2* MO and 300 pg of full length *BCAS2*

932 mRNA or *ΔCC1-2* mRNA at the one-cell stage. Scale bars, 5 μm. (F) Transcripts of

933 *gata1* were evaluated by WISH in *bcas2Δ14$^{+/-}$* embryos injected with 300 pg of *BCAS2*

934 mRNA or *ΔCC1-2* mRNA.