Reward history guides focal attention in whisker somatosensory cortex

Deepa L. Ramamurthy[1], Lucia Rodriguez[1,2], Celine Cen[1], Siqian Li[1], Andrew Chen[1], Daniel E. Feldman[1,3,*]

1. Department of Neuroscience and Helen Wills Neuroscience Institute, UC Berkeley
2. Neuroscience PhD Program, UC Berkeley
3. Lead Contact

*Corresponding author. Correspondence should be addressed to:

     Daniel E. Feldman
     dfeldman@berkeley.edu
     142 Weill Hall #3200
     Dept. of Neuroscience
     Univ. of California, Berkeley
     Berkeley, CA 94720-3200

Pages: 55
Figures: 7
Supplementary Figures: 7
Main text: 5187 words

ABSTRACT

Prior reward is a potent cue for attentional capture, but the underlying neurobiology is largely unknown. In a novel whisker touch detection task, we show that mice flexibly shift attention between specific whiskers on a trial-by-trial timescale, guided by the recent history of stimulus-reward association. Two-photon calcium imaging and spike recordings revealed a robust neurobiological correlate of attention in the somatosensory cortex (S1), boosting sensory responses to the attended whisker in L2/3 and L5, but not L4. Attentional boosting in L2/3 pyramidal cells was topographically precise and whisker-specific, and shifted receptive fields toward the attended whisker. L2/3 VIP interneurons were broadly activated by whisker stimuli, motion, and arousal but did not carry a whisker-specific attentional signal, and thus did not mediate spatially focused tactile attention. Together, these findings establish a new model of focal attention in the mouse whisker tactile system, showing that the history of stimuli and rewards in the recent past can dynamically engage local modulation in cortical sensory maps to guide flexible shifts in ongoing behavior.

INTRODUCTION

Humans and other animals engage attention to prioritize processing of behaviorally relevant stimuli in complex environments, including for vision[1], audition[2] and touch[3]. Past experiences across different timescales play an important role in guiding attention[4]. In humans, prior stimulus-reward association is a highly robust cue for attentional capture[5-17], such that perceptual detection or discrimination is selectively enhanced for previously rewarded stimuli, even when those stimuli are no longer important to current goals[14]. The effects of reward history on allocation of attentional priority have been termed experience-driven attention[15], value-driven attention[12-14], memory-guided attention[16], or attentional bias by previous reward history[17].

The neurobiology of attention has been studied extensively in non-human primates[1,18], and has been shown to involve boosting of signal-to-noise ratio for neural encoding of attended sensory features across the cortical sensory hierarchy, including primary sensory cortex. However, the fine-scale organization of attentional boosting in sensory cortex, and the neural circuits that control it, remain unclear. Mice provide powerful cell-type specific tools to identify precise neural circuit mechanisms underlying attentional processing. Here, we developed a new model of focal attention based on reward history in the mouse whisker tactile system, and used it to investigate the neurobiological basis of attention in sensory cortex.

We studied attentional behavior in head-fixed mice during a Go/NoGo whisker touch detection task which included Go stimuli on many different whiskers. When the whisker location of a touch stimulus was unpredictable from trial to trial, mice naturally used the history of stimulus-reward association as a cue to guide attention to specific, recently rewarded whiskers. This task generates a rich set of trial histories to probe which stimulus and reward contingencies guide attention, and allows us to track the shifting locus of attention on a trial-by-trial time scale. Using this platform, we identified a robust, spatially focused neural correlate of attention in whisker somatosensory cortex (S1), and tested a major circuit model of attention involving VIP interneurons in sensory cortex. More broadly, our study shows that in addition to the well-known role of learned stimulus-reward associations in slowly driving cortical map plasticity, the recent

history of stimuli and rewards acts on a fast time scale to dynamically and locally modify cortical
sensory maps with corresponding stimulus-specific shifts in perceptual sensitivity.

RESULTS

To study attention, we developed a head-fixed whisker detection task. Mice had nine whiskers
inserted in a piezo array, and on each trial, one randomly selected whisker (Go trials) or no whisker
(NoGo trials) was deflected in a brief train. Mice were rewarded for licking during a response
window on Go trials (Hits) but not on NoGo trials (False Alarms) (**Fig. 1A-B**). The task was
performed in darkness and incorporated a delay period of 0, 0.5, or 1-sec in different mice to
separate whisker stimuli from response licks. Go and NoGo trials were randomly intermixed with
a variable inter-trial interval (ITI), and whisker identity on Go trials was randomly selected. Thus,
mice could not anticipate the upcoming whisker or precise trial timing. Sequential trials could be
one Go and one NoGo, two NoGo trials, two Go trials on different whiskers, or two Go trials on
the same whisker (**Fig. 1C**).

Expert mice effectively distinguished Go from NoGo trials, quantified by d' from signal
detection theory (**Fig. S1A**). We analyzed behavior in each session during a continuous task-
engaged phase that excluded early and late low-performance epochs ($d'<0.5$) that reflect
motivational effects[19]. In expert mice, overall mean d' was 0.923 ± 0.051 (n = 476 sessions, 22
mice), but local fluctuations in d' regularly occurred over the time course of several trials,
suggesting that the recent history of stimuli or rewards may dynamically alter sensory detection
behavior (**Fig. S1A**). To examine this, we classified each current Go trial based on the history of
whisker stimuli and reward on immediately preceding trials. We used trial history categories of:
i) prior NoGo, ii) prior Hit to the same whisker as the current trial, iii) prior Hit to a different whisker,
iv) prior Miss to the same whisker, and v) prior Miss to a different whisker. Current NoGo trials
were classified into categories of prior NoGo, prior Hit (to any whisker), or prior Miss (to any
whisker). We separately tracked trials preceded by a single Hit trial from those preceded by
multiple sequential prior Hits ("prior >1 Hit"; **Fig. S1B**).
97
Reward history cues focal attention in the whisker system
Prior trial history strongly influenced detection on the current trial. When the prior trial was a
NoGo, mice detected the current Go whisker with $d' = 1.13 ± 0.02$ (termed d'_{NoGo}, mean ± SEM
across 476 sessions, 22 mice), which we consider baseline detection sensitivity. d' for detecting
a given Go whisker was elevated following 1 prior Hit to the same whisker, and even more so
following multiple consecutive Hits to the same whisker ($d'_{>1HitSame} = 2.45 ± 0.11$, p = 1.0e-4 vs
d'_{NoGo}, permutation test). In contrast, d' was reduced following one or multiple prior Hits to a
different whisker than the current trial ($d'_{>1HitDiff} = 0.82 ± 0.06$, p = 1.0e-4 vs d'_{NoGo}, permutation
test) (**Fig. 1D-E**). d' after multiple prior Hits to the same whisker ($d'_{>1HitSame}$) was substantially
greater than after multiple prior Hits to a different whisker ($d'_{>1HitDiff}$) (**Fig. 1D**). Thus, recent Hits
engage a whisker-specific boost in detection, evident as a whisker-specific increase in d', termed
Δ d' (**Fig. 1D**). This effect was found across mice with 0, 0.5, or 1-sec delay period (p = 1.0e-4
for $d'_{>1HitSame}$ vs $d'_{>1HitDiff}$ in each case), so these data were combined for behavioral analyses (**Fig.
S1D**).

Trial history-dependent boosting of detection required the conjunction of prior stimulus
plus reward, because d' did not increase after a prior Miss to the same whisker ($d'_{MissSame}$ = 0.90
± 0.09, p = 4.2e-3 relative to d'_{NoGo}). Boosting also failed to occur if the mouse licked to the prior
Go but received no reward (i.e., unrewarded hits) or a very small reward (<4% of maximal reward
volume ($d'_{>1HitSame}$ vs $d'_{>1HitDiff}$, p = 0.49, permutation test) (**Fig. 1F**). Thus, boosting was not driven
by prior whisker deflection alone, or stimulus-evoked licking, but by whisker-reward association
on recent trials.

We interpret this effect as whisker-specific attention, because it shares characteristic
features of stimulus-specific attention documented in primates during detection tasks[20-24]. This
involves both increased sensitivity (d') and a shift in decision criterion (c, also referred to as
response bias), with the whisker-specific Δd' reflecting increased Hit rate following >1 prior Hit to
the same whisker relative to >1 prior Hit to a different whisker (p = 1.0e-4, permutation test; **Fig.
1G**). The whisker-specific increase in d' was observed across mice (**Fig. 1H**; p = 1.0e-4,
permutation test). We also observed shifts in criterion[20,22-23] (Δ c) which with a more modest
whisker-specific component ($c_{>1HitSame}$ vs $c_{>1HitDiff}$, p = 0.002, permutation test) (**Fig. 1I**). On
average, the whisker-specific shift in sensitivity was larger than the whisker-specific shift in
criterion (**Fig. 1J**; Δ $d'_{>1HitSame}$ = 1.4 ± 0.20, Δ $d'_{>1HitDiff}$ = -0.47 ± 0.11, p = 1.0e-4, permutation test;
Δ $c_{>1HitSame}$ = -2.03 ± 0.18, Δ $c_{>1HitDiff}$ = -1.34 ± 0.08, p = 7.0e-4, permutation test).

Mice trained on all 3 delay periods showed the whisker-specific boost in d' (**Fig. S1C-D**).
Reaction times on Go trials were reduced by prior Hit trials, as expected for attention (assessed
in mice without a delay period), but this effect was not whisker-specific, which may reflect a ceiling
effect (**Fig. S1E**). The task interleaved whisker deflections of varying amplitude, and whisker-
specific boosting of d' was greatest for current Go trials with low-amplitude (weak) deflections, as
expected for attention (**Fig. S1F**). Trial history effects were consistently observed across different
variations of the task (used in different mice) in which we manipulated the stimulus probability of
each whisker in blocks, or manipulated the probability of sequential same-whisker Go trials to
make prior same Hit histories more likely (**Fig. S1G-H**). Trial history effects were driven by
stimulus-reward association, not by stimulus salience or reward probability, because whisker
deflections were physically identical and reward probability was always 100% for each whisker.

Behavioral shifts in d' exhibited the hallmark effects of focused attention: spatial specificity,
temporal specificity, and flexible targeting. Spatially, d' was boosted most strongly by prior Hits to
the same whisker, more weakly by prior Hits to an immediate same-row or same-arc neighbor,
and not at all by prior Hits to a diagonally adjacent or more distant whisker (**Fig. 1K**). Thus,
boosting is somatotopically organized. Temporally, attentional boosting fell off with the time
interval between Go trials (which varied due to variable ITI and intervening NoGo trials), and
subsided after ~10s (**Fig. 1L**). Enhancement of d' was flexibly shifted to different whiskers in an
interleaved manner, and had similar magnitude when cued by trial history to any of the 9 whiskers
in rows B-D or arcs 1-3 (**Fig. 1M**). Thus, mice use recent history of stimulus-reward association
to dynamically boost sensory detection of spatially specific whiskers on a rapid trial-by-trial
timescale, consistent with attentional enhancement[1,18,20-23, 25-26]. These properties strongly
resemble attentional capture guided by reward history in humans and non-human primates[4-17,24].

Whisker-specific attention is not mediated by whisker or body movement

To test whether trial history effects involve whisker or body movement, we extracted these movements, plus pupillary dilations related to arousal[27-28], from behavioral videos of 9 mice (74 sessions) using DeepLabCut[29] (**Fig. S2A**). Reward retrieval at the end of Hit trials was associated with whisker movement, body movement (detected from platform motion), and pupil dilation that slowly subsided during the ITI before the next trial (**Fig. 2A**). The magnitude of movement and pupil dilation during the ITI was greater after 1 or >1 prior Hits, relative to prior NoGo, but was identical for prior same and prior different conditions (**Fig. 2A, Fig. S2B**). Thus, mice exhibited increased motion and arousal following prior Hits. During the subsequent trial, whisker stimulation evoked modest whisker and body motion during the stimulus period, and these were also heightened after prior Hits (p = 1e-4, permutation test), indicating that behavioral arousal and motion effects from prior Hits persisted into subsequent trials, but did not differ between prior same whisker and prior different conditions (p = 0.39, permutation test **Fig. 2A-B**).

To test whether active whisker movement contributed to the whisker-specific $\Delta\,d'$ effect, we paralyzed whisker movements with Botulinum toxin B (Botox) injection in the vibrissal pad in 4 task-expert mice. Paralysis was maintained over 7-8 days, and history effects were compared between standard sessions prior to Botox and the Botox sessions. The average whisker-specific shift in behavioral d' and c following >1 prior hit did not differ between Botox and non-Botox sessions (**Fig. 2C-D**) demonstrating that whisker-specific attentional effects do not require active whisker movement. Thus, although prior hits also engage increases in whisker motion, body motion and arousal, whisker-specific attentional effects were independent of these effects on global behavior state.

Neural correlates of focal attention in L2/3 PYR cells in S1

The somatotopic precision of attentional effects on behavior (**Fig. 1K**) suggests a neural basis in a somatotopically organized brain area like S1. We performed 2-photon imaging in S1, using Drd3-Cre;Ai162D mice that transgenically express GCaMP6s in L2/3 PYR cells. Mice performed the task with a delay period that separated whisker-evoked responses (analyzed 0-799 ms after stimulus onset) from later licks and rewards, and any trials with early licks were excluded. Imaging fields were localized in the S1 whisker map by post-hoc cytochrome oxidase staining for whisker barrel boundaries in L4 (**Fig. 3A-B**). 61% of L2/3 PYR neurons were whisker responsive, and trial history modulated whisker responses for many individual neurons. For example, in **Fig. 3C**, neurons increased their response to the C3 whisker after >1 prior hit to that same whisker, but not after >1 prior hit to a different whisker.

On average, L2/3 PYR cells responded to a given Go whisker when the prior trial was a NoGo, responded more strongly following 1 prior Hit to the same whisker (p = 1e-4), and even more after >1 prior Hit to the same whisker (p = 1e-4). This boosting of sensory responses did not occur when the prior trial was a Miss to the same whisker, or >1 Hit to a different whisker (n = 6 mice, 70 sessions, 6906 cells, **Fig. 3D-E**). Thus, recent stimulus-reward association modulated whisker-evoked $\Delta F/F$ in L2/3 PYR cells in a way that closely resembled the behavioral attention effect (**Fig. 3E** vs **Fig. 1D**). On current NoGo trials, no whisker stimulus was presented and $\Delta F/F$ traces were largely flat, except for NoGo trials following prior Hit trials, which exhibited a surprising rising $\Delta F/F$ signal. We interpret this as an expectation or arousal effect, which parallels the

increased FA rate on these trials (**Fig. 1G**). These effects were evident in single example fields
(**Fig. S3A**). History-dependent boosting of whisker responses was most evident in whisker-
responsive cells (defined from trials after a prior NoGo), and did not occur in cells that were non-
responsive after a prior NoGo (**Fig. S3B-C**). Boosting after prior Hits was whisker-specific in all
6/6 mice, but its magnitude varied across mice (**Fig. 3F**) and correlated with the magnitude of the
behavioral attention effect measured by $\Delta d'$ in each mouse (**Fig. 3G**).

We quantified the attention effect in individual cells using three attention modulation
indices (AMI). $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ quantify the change in whisker-evoked
response observed after >1 prior Hit to the same (or different) whisker vs after a prior NoGo. Most
cells showed positive $AMI_{>1HitSame-NoGo}$ values and negative $AMI_{>1HitDiff-NoGo}$ values, indicating up-
and down-modulation of response magnitude by the identity of the prior whisker Hit. $AMI_{>1HitSame-}$
$_{>1HitDiff}$ compares response magnitude after >1 prior hit to the same whisker vs >1 prior hit to a
different whisker. This was shifted to positive values for most L2/3 PYR cells, indicating whisker-
specific attentional modulation (**Fig. 3H, I**). This was reproducible across individual mice (**Fig.
S3D**). Modulation of whisker-evoked responses for all cells sorted by $AMI_{>1HitSame->1HitDiff}$ is shown
in **Fig. S3E**.

History-dependent boosting was somatotopically restricted in S1. After >1 prior hits to a
given whisker, responses to an immediate same-row adjacent neighboring whisker were boosted
strongly, those to an immediate same-arc neighbor were boosted less, and those to diagonal
adjacent neighbors or further whiskers were boosted the least or not at all. This somatotopic
profile of $\Delta F/F$ boosting strongly resembled the somatotopy of behavioral d' boosting (**Fig. 3J**).
Spatially within S1, >1 prior hits to a reference whisker boosted whisker-evoked $\Delta F/F$ to that
whisker most strongly for L2/3 PYR cells within the reference whisker column and in the near half
of the neighboring column, and weakly or not at all beyond that. This somatotopically constrained
boosting was not observed for >1 prior hits to a different whisker, or following a miss to the
reference whisker (**Fig. 3K**). Analysis of individual cell AMI confirmed somatotopically precise
boosting (**Fig. S4A**). This defines the spatial profile of the trial history-based 'attentional spotlight'
in S1 as boosting responses to the cued whisker within a region of 1.5 columns width in L2/3. At
the center of the spotlight, the representation of the attended whisker is boosted within its own
column (**Fig. S4B-C**).
226

Attention involves shifts in receptive fields toward the attended whisker
In primates, attention not only increases sensory response magnitude and signal-to-noise ratio in
sensory cortex, but can also shift neural tuning toward attended stimuli[30-32]. We tested whether
attention involves receptive field shifts by L2/3 PYR cells in S1. We analyzed all whisker-
responsive cells relative to the boundaries of the nearest column (defined in the Prior NoGo
condition). We calculated the mean receptive field across 9 whiskers centered on the columnar
whisker (CW) when the prior trial was a NoGo (**Fig. 4A**, black traces in center panel). We then
recalculated the receptive field, for these same cells, for >1 Prior Hit to each of the surround
whiskers (purple traces in outside panels).   For many whiskers, recent stimulus-reward
association shifted the mean receptive field toward the attended whisker or nearby whiskers (**Fig.
4A**). To quantify this effect, we calculated the tuning center of mass (CoM) across the 3 by 3
whisker array for these mean receptive fields. History-based cueing to an attentional target

whisker generally shifted tuning CoM towards that whisker (**Fig. 4B**). An exception was when
attention was cued upward, which caused little upward CoM shift. This was not explained by
known experimental factors, but could reflect a spatial asymmetry in attentional effects[33] on
whisker touch (**Fig. 4B**). Attentional cueing to the CW elevated responses to that whisker within
its S1 column but caused very little tuning change (**Fig. S5A**).

To measure receptive field shifts in individual neurons, we quantified the shift in CoM for
each responsive cell along an attentional axis from the CW to the attentional target whisker, which
was one of the 8 surrounding whiskers (**Fig. S5B-C**). In the >1 prior Hit condition, the mean CoM
shift along the attentional axis was 0.12 ± 0.04 (n = 173 cells with sufficient whisker sampling, p
= 3.6e-3, one-sample permutation test vs. mean of 0), indicating a tuning shift toward the attended
whisker. A range of tuning shifts were observed, with significantly more cells showing CoM shifts
towards the attended whisker than away from it (62% vs. 38%, p = 2.3e-3, binomial exact test for
difference from 0.5). Thus, attentional cueing involves receptive field shifts as well as modulation
of whisker response magnitude.
253
Attention boosts population decoding of recently rewarded whiskers
Is the magnitude of attentional boosting of L2/3 PYR responses sufficient to improve neural coding
of attended whisker stimuli on single trials? To test this, we built a simple neural decoder that
uses logistic regression to predict the presence of any whisker stimulus from single-trial
population mean ΔF/F calculated across all whisker-responsive cells in a single imaging field.
Each field spanned ~1-1.5 columns within the 9-whisker region of S1, typically centered over the
location of column corresponding to the center whisker on the piezo array. A separate decoder
was trained on each session (n = 70 sessions, 6 mice), and performance was assessed from
held-out trials (**Fig. 5A**). Because S1 is somatotopically organized, we observed modest, above-
chance performance for detecting any of the 9 whiskers from mean activity in a single field
(relative to NoGo trials or shuffled data), strong performance for detecting the field best whisker
(fBW) that is topographically matched to the field location, and no ability to detect non-
topographically aligned whiskers (non-fBWs) (**Fig. 5B**).

Population mean ΔF/F and decoder performance were related to current trial outcome,
with greater ΔF/F and stimulus prediction on current Hit trials than Miss trials, and on current false
alarm trials than correct rejection trials. This is consistent with the known representation of
sensory decision in S1[34-35] (**Fig. 5C-D**). Next, we examined decoding as a function of prior trial
history. Detection of any whisker stimulus from Go trials was improved following >1 Hit to the
same whisker, relative to prior NoGo (p = 1e-0.4, permutation test) or to >1 prior Hit to a different
whisker (p = 0.001, permutation test) (**Fig. 5E**). This boost in decoding performance did not occur
when the decoder was tested only on fBW trials but did occur for non-fBW trials (**Fig. 5F-G**). This
same effect was found when decoding was evaluated on current Hit trials only, so this is an effect
of trial history, not current sensory decision (**Fig. 5E-F**).

Thus, attention to a whisker strengthens its encoding on single trials in S1. The lack of
improvement in decoding attended fBWs likely reflects the strong coding of these whiskers under
baseline conditions, such that attentional boosting of CW responses (**Fig. S4**) does not further
improve detection.

Attentional modulation of neural coding with Neuropixels spike recordings

To examine attentional modulation of S1 neural coding at finer temporal resolution and across layers, we recorded extracellular spiking in S1 using Neuropixels probes[36] that spanned L1-6 (**Fig. 6A**). Mice performed a modified version of the task in which Go stimuli were distributed over only 4 or 5 whiskers, rather than 9, to enable adequate sampling of each history condition per session. Behaviorally, mice performing the task with 4-5 whiskers showed the whisker-specific Δ d' attention effect, but at lower magnitude due to the smaller number of whiskers (**Fig. S6A**). All sessions included the CW for the recording site plus 3 nearby whiskers. We spike sorted to identify single units, classified units as regular-spiking (RS) or fast-spiking (FS), and assigned laminar identity based on CSD analysis of local field potentials (**Fig. S6B-F**). Many single units showed history-dependent modulation of whisker-evoked spiking (**Fig. 6B**).

S1 units responded to each deflection in the stimulus train. On average for L2/3 RS units, whisker-evoked spiking was boosted in Go trials after >1 prior Hit to the same whisker, relative to >1 prior Hit to a different whisker (**Fig. 6C-D**). This effect did not occur after prior Miss trials, and firing on NoGo trials was not significantly regulated (**Fig. 6D**). L4 RS units showed only a slight trend for history-dependent modulation that did not reach significance (**Fig. 6E-F**). L5a/b RS units (grouped together) showed a whisker-specific attentional boost similar to L2/3 (**Fig. 6G-H**).

To examine heterogeneity across units we calculated AMI for each unit. Most L2/3 RS units responded more strongly after prior Hits to the same whisker relative to prior NoGo (as measured by $AMI_{>1HitSame-NoGo}$) and more weakly after prior Hits to a different whisker relative to prior NoGo (as measured by $AMI_{>1HitDiff-NoGo}$). This whisker-specific attentional effect, evident as the separation between $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ distributions, was not present in L4, and was weaker in L5a/b (**Fig. 6I**). These laminar trends were also apparent in $AMI_{>1HitSame->1HitDiff}$, which was shifted positively in L2/3 relative to L5a/b and L4 units (**Fig. 6J**). Calculating the mean AMI across neurons confirmed whisker-specific attentional shifts in L2/3 and L5, but not in L4 (**Fig. 6K;** ($AMI_{>1HitSame-NoGo}$ vs $AMI_{>1HitDiff-NoGo,}$ L2/3: p = 0.03, L4 p = 0.65, L5a/b: p = 0.01; $AMI_{>1HitSame->1HitDiff}$, L2/3: p = 0.031, L4 p = 0.51, L5a/b: p = 0.06, permutation test). This suggests that history-dependent attentional modulation is not simply inherited from the thalamus, but has a cortical component.

VIP interneurons do not carry a simple "attend here" signal

We used reward history-based attention in S1 to investigate the candidate involvement of VIP interneurons in attentional control. VIP cells are known to disinhibit PYR cells to increase PYR sensory gain during arousal, locomotion, and whisking[37-41]. For attention, this same VIP circuit has been hypothesized to be activated by long-range (e.g., top-down) inputs, and to act to amplify local PYR responses to selected sensory features[40-41] (**Fig. 7A**). Whether VIP cells mediate goal-directed attention is still unclear[42-43], and their involvement in history-based attention has not been tested. We used 2-photon imaging from L2/3 VIP cells in VIP-Cre;Ai162 mice to ask whether VIP cells are activated when mice direct attention to a particular whisker column in S1 (**Fig. 7B**).

VIP cells in S1 are activated by arousal (indexed by pupil size), whisker and body movement, and goal-directed licking during this whisker detection task[44]. These behaviors all peak at the end of Hit trials, as mice retrieve rewards, and then systematically decline during the ITI, which ends with a 3-sec lick-free period that is required to initiate the next trial. As a result,

VIP cell ΔF/F falls systematically during the ITI after Hit trials, correlated with these behavioral
variables, and falls less after NoGo or Miss trials (**Fig. 7C**). VIP cells in S1 also show robust
whisker stimulus-evoked ΔF/F transients, which ride on this declining baseline[44]. To test whether
whisker-evoked VIP responses are greater when reward history cues attentional capture, we
calculated mean ΔF/F traces (n = 7 mice, 103 sessions, 1843 VIP cells) as a function of trial
history. Baseline (prestimulus) ΔF/F declined more steeply on trials following prior Hits than
following prior NoGo or Miss, as expected. Superimposed on this, and clearest after detrending
the baseline, whisker-evoked Δ F/F was also increased after prior Hit trials. However, this was
not whisker-specific (**Fig. 7D-F**). AMI analysis confirmed increased responsiveness for most VIP
cells in both prior >1 Hit same and prior >1 Hit different conditions, but no whisker-specific
attention effect (the $\text{AMI}_{>1HitSame->1HitDiff}$ index was peaked at 0; **Fig. 7G**, for all cells individually see
Fig. S7).

Together, these findings indicate that VIP cells as a population do not carry a whisker-
specific attention signal, but do exhibit a general increase in activity with multiple prior Hits to any
whisker that is consistent with global arousal and motion effects. Although we cannot rule out
that a subpopulation of VIP cells may carry a whisker-specific "attend here" signal, the VIP
population as a whole does not.

DISCUSSION

Attention captured by recent history cues[4-17], including stimulus-, reward-, and choice-history,
provides a powerful model to study mechanisms of attention. In our paradigm, enhanced
detection (D *d'*) of whisker stimuli was driven by recent whisker stimulus-reward association, had
the defining features of selective attention (spatially focused, flexibly allocated, and temporally
constrained)[4,20-23]. Behavioral and neural effects in our study were whisker-specific, and thus did
not correspond to a global arousal or motion effect[45]. They were not explained by priming, which
occurs in response to stimulus presentation without reward association, does not require
detection of the priming stimulus, and typically has short (<100 ms) duration[46].

Attentional boosting in our study was not bottom-up attention, because it was not driven
by physical stimulus salience (whisker stimuli had the same stimulus strength), and it was
not top-down attention, because it was automatically engaged and not goal-directed (i.e.,
whisker stimuli were all equally rewarded, so it did not increase overall reward rate in the
task). These results align well with automatic attentional capture by reward history in humans[4-
17], which is theorized to represent a category of attentional processes often called "selection
history"[8-10] distinct from classical top-down and bottom-up attention. Rodents are well known to
exhibit history-dependent response biases (i.e., shifts in decision criterion for behavioral choices)
in perceptual decision-making tasks[47-51], including serial dependence[47,51-52], contraction bias[47-
48,51], adaptation aftereffects[53], win-stay/lose-switch strategies[54-56] and choice alternation[54]. Our
findings show that mice also use prior reward history to prioritize sensory processing, through
stimulus-specific shifts in perceptual sensitivity (*d'*), in addition to shifts in decision criterion (c).

The neural correlates of attention have been primarily studied in non-human primates, and
include increased sensory-evoked spike rate[18,57-58], reduced variability[18], neural synchrony
modulation[59], and changes in receptive fields[18,30-32] including in receptive size, boosting of peak

responses, and receptive field shifts. In primates, these effects are greatest at higher levels of the
sensory hierarchy but also occur in primary sensory cortex[18,60]. The precise spatial organization
of these coding effects in sensory cortex has been unknown, and has important implications for
identifying the neural control circuits for attention.   On the macroscopic scale, human brain
imaging and focal pharmacological inactivation studies in non-human primates indicate that
spatial attention in vision is retinotopically organized within visual cortical areas[1,18,61-64]. But the
precise spatial organization of attentional modulation in sensory cortex (i.e., the spatial profile of
the spotlight of attention) has not been known. Importantly, our task design (in which we track
spontaneous behaviors, and separate stimulus and lick response windows with a delay period)
allowed us to distinguish attentional signals from global arousal and motion signals, which
dominate neural activity during behavior and are widespread across cortical areas[45].

We took advantage of S1 whisker map topography[65] to quantitatively define the precise
spatial structure of the attentional spotlight relative to anatomical cortical columns in S1.
Attentional capture boosted sensory responses to the attended whisker in a region comprising
that whisker's column plus the near half of surrounding columns. In this region, whisker-evoked
spike rate and receptive field peak increased (in the central attended column) and receptive fields
shifted toward the attended whisker (for cells in surrounding columns). Together, this increased
total neural activity evoked by the attended whisker, both by increasing the number of PYR cells
responding to that whisker, and by elevating the number of spikes per cell. This somatotopically
restricted boosting[61-66] is distinct from the spatially broad modulation of sensory responses that
occurs across entire cortical areas (or multiple areas) in response to global behavioral state (e.g.,
arousal indexed by pupil size, active whisker movement for S1, or locomotion for V1)[45]. Our results
show that attentional boosting can be flexibly targeted with a precision of ~300 μm in cortical
space for stimulus-specific modulation of the neural code.   Thus, neural control circuits for
attention (which may involve feedforward, local, feedback, or neuromodulatory circuits) must
operate with this spatial precision.

History signals in mouse cortex have not previously been described for attention, but have
been identified in posterior parietal cortex (PPC) and orbitofrontal cortex (OFC) during decision
making[67-69] and in reversal learning[70-73]. S1 receives instructional signals from OFC that are
necessary for reversal learning[72], but whether this pathway plays a role in attention cued by recent
reward history is unknown.

The neural mechanisms and control circuits for attention remain poorly understood, and
likely differ between different forms of attention. We found that attention to touch modulates PYR
sensory responses in L2/3 and L5a/b but not L4, suggesting either an intracortical origin, or a
thalamic origin in secondary thalamic nuclei like the posterior medial nucleus (POm), which
projects to L2/3 and L5a. Thalamic control of attention has been implicated in visual and tactile
cross-modal attention tasks in mice[74], as well as some non-human primate studies[75]. We tested
one major circuit model[39-43] for attentional boosting in sensory cortex, that long-range inputs
amplify pyramidal (PYR) cell sensory responses by activating local VIP interneurons in sensory
cortex[39,76]. L2/3 VIP interneurons receive local, feedforward, and feedback glutamatergic input,
as well as by neuromodulatory input, and inhibit other cortical interneurons to disinhibit PYR cells,
thus boosting PYR sensory responses[39,76]. VIP interneurons are activated by global behavioral
state (e.g., locomotion and whisking), by spontaneous arousal during quiet wakefulness[27,37-38],

and by top-down contextual signals[40]. Top-down input from anterior cingulate cortex (ACC) to VIP
cells in sensory cortex has been suggested to mediate top-down attentional effects on sensory
processing[39]. However, recent studies have questioned this model[42,77], finding that L2/3 VIP
modulation of PYR activity is orthogonal to attentional effects in a cross-modal attention task[42].
We tested the potential involvement of VIP cells in focal attention by asking whether VIP cell
activity is enhanced in attended columns, as required if these cells contribute to boosting of PYR
cell responsiveness. We found that L2/3 VIP cells, at least as a full population, do not carry this
whisker-specific "attend-here" signal, but instead show general, non-whisker-specific activation in
response to any prior Hit, consistent with an arousal- or global behavior-related signal. This
suggests VIP cells are more engaged in global modulation of whisker sensory responsiveness
during arousal and motion, and not whisker-specific attention cued by reward history.

Attention to whisker touch cued by recent reward history in mice is a novel paradigm for
studying the neurobiological mechanisms of focal attention. This model complements recent
visual tasks that aim to study top-down[79-82] and bottom-up[83-84] attention in head-fixed mice.
Together, these paradigms can reveal the extent to which common vs. distinct neurobiological
mechanisms are engaged in different forms of attention, and across different sensory modalities.
425

METHODS

Animals
All methods followed NIH guidelines and were approved by the UC Berkeley Animal Care and
Use Committee. The study used 22 mice. These included 7 Drd3-Cre;Ai162D mice and 10 VIP-
Cre;Ai162D mice (used for behavior and 2-photon imaging), and 5 offspring from Drd3-Cre x
Ai162D crosses (genotype not determined) used in extracellular recording experiments. VIP-Cre
(JAX # 10908) and Ai162D mice (JAX # 031562) were from The Jackson Laboratory. Drd3-Cre
mice were from Gensat MMRRC (strain number 034610).

Mice were kept in a reverse 12:12 light cycle, and were housed with littermates before
surgery and individually after cranial window surgery. Mice were roughly evenly divided between
male and female, and no sex differences were found for the results reported here. Behavioral,
imaging, and analysis methods were as described in Ramamurthy et al., 2023[44], and are here
described more briefly. Results reported for 7 VIP-Cre;Ai162D mice and 3 Drd3-Cre;Ai162D mice
are new analyses which include data from the dataset reported in Ramamurthy et al., 2023.
440

Surgery for behavioral training and 2-photon imaging
Mice (2-3 months of age) were anesthetized with isoflurane (1-3%) and maintained at 37°C.
Dexamethasone (2 mg/kg) was given to minimize inflammation, meloxicam (5-10 mg/kg) for
analgesia, and enrofloxacin (10 mg/kg) to prevent infection. Using sterile technique, a lightweight
(<3 g) metal head plate containing a 6 mm aperture was affixed to the skull using cyanoacrylate
glue and Metabond (C&B Metabond, Parkell). The headplate allowed both head fixation and 2-
photon imaging through the aperture. Intrinsic signal optical imaging (ISOI) was used to localize
either C-row (C1, C2, C3) or D-row (D1, D2, D3) barrel columns in S1[85], and a 3 mm craniotomy
was made within the aperture using a biopsy punch over either the C2 or D2 column. The
craniotomy was covered with a 3 mm diameter glass coverslip (#1 thickness, CS-3R, Warner
Instruments) over the dura, and sealed with Metabond to form a chronic cranial window. Mice

were monitored on a heating pad until sternal recumbency was restored, given subcutaneous
buprenorphine (0.05 mg/kg) to relieve post-operative pain and then returned to their cages. After
the mice recovered for a week, behavioral training began.
455
Behavioral task
To motivate behavioral training, each mouse received 0.8-1.5 mL of water daily, calibrated to
maintain 85% of pre-training body weight. Mice were weighed and observed daily. Behavioral
training sessions took place 5-7 days per week. For behavioral training, the mouse was head-
fixed and rested on a spring-mounted stage[44,86]. Nine whiskers were inserted in a 3 x 3 piezo
array, typically centered on a D-row or C-row whisker. Piezo tips were located ~5 mm from
the face, and each whisker was held in place by a small amount of rubber cement. A tenth
piezo was present near the 3x3 array but did not hold any whisker ("dummy piezo"). A
capacitive lick sensor (for imaging experiments) or an infrared (IR) lick sensor (for extracellular
recording experiments) detected licks, and water reward (mean 4 µl) was delivered via a
solenoid valve. Mice were transiently anesthetized with isoflurane (0.5-2.0%) at the start of
each session to enable head-fixation and whisker insertion, after which isoflurane was
discontinued and behavioral testing began after the effects of anesthesia had fully recovered.
Behavior was performed in the dark with 850 nm IR illumination for video monitoring. Masking
noise was presented from nearby speakers to mask piezo actuator sounds. Task control, user
input and task monitoring were performed using custom Igor Pro (WaveMetrics) routines and an
Arduino Mega 2560 microcontroller board.
473
<u>Training stages</u>
A series of training stages (1-5 days each) were used to shape behavior on the Go/NoGo
detection task. In Stage 1, mice were habituated to the experimental rig and to handling. In Stage
2 mice were head-fixed and conditioned to lick for water reward at the port. In Stage 3, mice
received a reward (cued by a blue light) for suppressing licks for at least ~3 seconds, termed the
Interlick Interval (ILI) threshold. Stage 4 introduced whisker stimulation for the first time, with 50%
Go trials (whisker stimulation) and 50% NoGo trials (no whisker stimulus), with automatic reward
delivery in the response window on Go trials (i.e., classical conditioning). The dummy piezo was
actuated on all NoGo trials, so that any unmasked piezo sounds did not provide cues for task
performance. The Go whisker randomly chosen from among 9 possible whiskers. This stage
ended when mice shifted licks in time to occur before reward delivery. In Stage 5, training
switched to operant conditioning mode, and mice were required to lick in the response window (0
- 300 ms after whisker stimulus onset) to receive a reward. There was no delay period at this
stage. Learning progress was tracked by divergence of Go/NoGo lick probability. In Stage 6, the
delay period was introduced. To do this, we introduced a trial abort window in which licking during
the stimulus presentation period caused a trial to be canceled without reward, to discourage
licking during the stimulus presentation. Simultaneously, we implemented a ramp-plateau reward
gradient within the response window, so that later licks resulted in a larger reward. Over the course
of this stage, the trial abort window was gradually lengthened, and the time of reward plateau was
gradually increased. Learning progress was tracked by the gradual increase in median trial first
lick time. Stage 7 represented the final whisker detection task, which included completely
randomized Go/NoGo trials, fixed trial abort window and reward plateau parameters, and

eliminating the blue light that signaled reward delivery. Mice were deemed task experts when
they exhibited stable performance at d' > 1 (mean running d-prime) for three consecutive
sessions. Expert mice performed 500 - 1000 trials daily.
499
Task structure
Task structure was identical to Ramamurthy et al., 2023[44] . Briefly, on each trial, a whisker
stimulus was applied to either one randomly selected whisker (Go trials, 50-60% of trials) or no
whisker (NoGo trials, 40-50% of trials). The whisker stimulus consisted of a train of five deflections
separated by 100 ms each. Every deflection was a 300 μm amplitude (6º angular deflection)
rostrocaudal ramp-and-return movement with 5 ms rise/fall time and 10 ms duration. NoGo trials
presented the same stimulus on a dummy piezo that did not contact a whisker, so that any
unmasked auditory cue from piezo movement was matched between Go and NoGo trials. Trial
onset was irregular with an ITI of 3 ± 2 s. Mice had to restrict licking to greater than 3-sec interlick
interval (ILI) to initiate the next trial. On Go trials, mice were rewarded for licking within the 2.0 s
response window with ILI of <300 ms. Licking was not rewarded on NoGo trials. Each trial
outcome was recorded as a Hit, Miss, False Alarm, or Correct Rejection.

Different delay periods were used in different mice (**Fig. S3C-D**). Mice used in imaging or
extracellular recording experiments had a delay period of 500 ms (for all spike recording mice and
7/14 2p imaging mice) or 1000 ms (for the other seven 2p imaging mice). This was used to
separate sensory-driven neural activity from action- and reward-related activity. Three mice
used only for behavioral data collection were tested without a delay period, which enabled
testing of attention effects on lick response latency (**Fig. S3E**).

Because reward size varied with lick time in the response window, reward volume varied
across trials. In addition, for mice with the 1000 ms delay period, mice sometimes licked on Go
trials after stimulus presentation but before the response window opened, and thus earned no
reward. These represent unrewarded Hits, so that attention effects could be quantified based on
absence of reward and reward size on prior Hit (**Fig. 1F**).
523
Task variations
All mice in this study were trained on an "equal probability" (EqP) version of the task in which
whisker identity on each Go trial was randomly selected from nine possible whiskers with equal
1/9 probability (EqP). Two other task variations manipulated either the global or local probability
of each specific whisker, while still randomly selecting whisker identity on each trial. In high
probability (HiP) sessions, we manipulated the global stimulus probability of each whisker by
presenting one whisker with higher probability (80% of Go trials) than the others. This was done
in 300-400 trial blocks, interleaved with standard EqP blocks (13 mice) on the same day. In high
probability of same whisker (HiPSame) sessions (run on separate days from EqP sessions) we
manipulated the local probability of repeating the same whisker stimulus on consecutive trials,
while maintaining the overall probability of each whisker at 1/9. This was done in blocks
interleaved with EqP blocks (7 mice). Whisker-specific attentional cueing was observed in all 3
task variants (**Fig. 1H**), so data from all variations were combined for the rest of the analyses. All
22 mice were trained and tested on the EqP version and either the HiP or HiPSame version of
the task, but EqP blocks/sessions contributed trials to history analysis only in a subset of animals,

since multiple hits to a given whisker in a single session were adequately sampled only in sessions
with a higher number of total trials.

A modified task version was used for extracellular recording experiments, in order to
adequately sample trial history conditions when only 2-4 days of acute recording were possible
per mouse. To do this, we reduced the number of whiskers sampled during Go trials from 9
whiskers to either 4 or 5 whiskers 5 (4-5-whisker task; **Fig. 6A**). The 4-5-whisker task was used
in 3 mice for extracellular recordings, and was also applied in 2 mice that were used in PYR cell
imaging, where performance could be compared to the standard 9-whisker task (**Fig. S6A**).

Behavioral movies & DeepLabCut tracking
Behavioral movies were acquired at 15-30 frames/sec using either a Logitech HD Pro Webcam
C920 (modified for IR detection) or FLIR Blackfly S (BFS-U3-63S4M; used in video analyses).
DeepLabCut[29] was used to track spontaneous face and body movements. Movies were manually
labeled to generate training datasets for tracking facial motion (snout tip, whisker pad and 2-3
whiskers), body motion (corner of the mouse stage, whose motion reflects limb and postural
movements), pupil size (8 labels on the circumference of the pupil), eyelids (8 labels on the
circumference of the eyelid), and licking (tongue and lickport). Three separate networks were
trained (100,000-200,000 iterations) such that a good fit to training data was achieved (loss <
0.005). One network each was trained for face/body motion for the two camera setups (version
1 network: 1110 labeled frames from 37 video clips across 6 mice) and another for pupillometry
(version 2 network: 2463 labeled frames from 27 video clips across 2 mice).

Behavioral movies from 74 sessions in 9 mice were analyzed for whisker motion (average
across all whisker-related labels), body motion (stage corner) and pupil size (ellipse fit to the pupil
markers). Blinking artifacts were removed using a one-dimensional moving median filter (40 frame
window) applied to the trace of pupil size as a function of time. Pupil size measured on each frame
was normalized to the mean pupil size over each individual session.

Whisker paralysis by Botox injection
Four mice (2 VIP-Cre;Ai162D mice and 2 Drd3-Cre;Ai162D mice) that were used for imaging
experiments also underwent Botox injection to induce paralysis of whisking[34,44,86-88]. Both whisker
pads were injected with Botox (Botulinum Neurotoxin Type A from Clostridium botulinum, List Labs
#130B). A stock solution of 40 ng/µl Botox was prepared with 1mg/ml bovine serum albumin in
distilled water. Each whisker pad was injected with 1 µl of a 10 pg/µl final dilution using a microliter
syringe (Hamilton). Whisking stopped within 1 day, and gradually recovered in ~1 week. Following
the initial dose, a 50% Botox supplement was injected once per week, as needed. Imaging was
performed > 24 hours after any Botox injection.

Behavioral Analysis
Behavioral performance was assessed using the signal detection theory measures[89] of
detection sensitivity (d') and criterion (c), calculated from Hit rates (HR) and False Alarm rates
(FA), as per their standard definitions:

$$d' = Z_{HR} - Z_{FA}$$

$$c = \frac{1}{2}\left(Z_{HR} + Z_{FA}\right)$$

where Z is the inverse cumulative of the normal distribution.

To assess the overall behavioral performance of mice on the whisker detection task, *d'*
was computed across trials over the entire behavioral session. For each session, a sliding *d'*
cutoff (calculated over a 50 trial sliding window) was applied to the start and end of the session,
and analysis was restricted between the first and last trial that met the threshold, to minimize
satiety effects. A standard sliding *d'* cutoff of 0.5 was used for all behavioral and imaging
analyses. We tested *d'* cutoffs 0.5, 0.7 and 1 to ensure that choice of *d'* cutoff did not affect key
results. A sliding *d'* cutoff of 1.2 was used for extracellular recording analyses.
590

591   ***Definition of trial histories***
592   For each *current trial*, trial history conditions were defined based on outcome and stimulus on
593   *prior trials*, as follows. These definitions are illustrated in **Fig. S1B**.

594   History conditions were defined for *current Go trials* as follows:
595   1. Prior Miss
596   a. On the *same* whisker: The current trial is a Go and the outcome on the previous
597   trial was a Miss to the same whisker.
598   b. On a *different* whisker: The current trial is a Go and the outcome on the previous
599   trial was a Miss to a different whisker.
600   2. Prior NoGo: The previous trial was a NoGo (any outcome).
601   3. Prior 1 Hit
602   a. On the *same* whisker: The current trial is a Go and the outcome on the previous
603   trial was a Hit to the same whisker.
604   b. On a *different* whisker: The current trial is a Go and the outcome on the previous
605   trial was a Hit to a different whisker.
606   4. Prior >1 Hit:
607   a. On the *same* whisker: The current trial is a Go and the outcomes on the previous
608   two or more trials were Hits to the same whisker as the whisker presented on
609   the current trial.
610   b. On a *different* whisker: The current trial is a Go and the outcomes on the previous
611   two or more trials were Hits to a single consistent whisker that was a different
612   identity than the whisker presented on the current trial (i.e., two or more Hits in
613   a row to the same whisker that differed from the current Go trial).

614   For each history condition defined above for *current Go trials*, a matched condition was
615   defined for *current NoGo trials*. This allowed us to compute behavioral *d'* and *c* values within
616   each history condition, and to compare neural signals on Go and NoGo trials within each history
617   condition. History conditions for *current NoGo trials* were defined as follows:
618   1. Prior Miss: The current trial is a NoGo and the outcome on the previous trial was a miss
619   to any whisker. NoGo trials in this category were used for comparison with Go trials in
620   both categories 1a and 1b above.
621   2. Prior NoGo: The previous trial was a NoGo (any outcome).
622   3. Prior 1 Hit: The current trial is a NoGo and the outcome on the previous trial was a hit
623   to any whisker. NoGo trials in this category were used for comparison with Go trials in
624   both categories 3a and 3b above.

4. Prior >1 Hit: The current trial is a NoGo and the outcomes on the previous two or more
trials were Hits to any single repeated whisker. NoGo trials in this category were used
for comparison with Go trials in categories 4a and 4b, above.

The history conditions above were defined based on sequences of consecutive trials,
including both Go and NoGo. The history-dependent effects on detection behavior were
maintained when history conditions were defined by ignoring NoGo trials, and categorizing
history based solely on Go trials (data not shown). NoGo trials were ignored when
characterizing the temporal profile of attention as a function of time since last Go trial (**Fig. 1L**).

2-photon calcium imaging
A Sutter Moveable-Objective Microscope with resonant-galvo scanning (RESSCAN-MOM, Sutter)
was used to perform 2p imaging in expert mice. A Ti-Sapphire femtosecond pulsed laser
(Coherent Chameleon Ultra II) tuned to 920 nm, or an ALCOR 920 nm fixed wavelength
femtosecond fiber laser (Spark Lasers), was used for GCaMP6s excitation. A water-dipping
objective (16x, 0.8 NA, Nikon) was used, and emission was band-pass filtered (HQ 575/50 filter,
Chroma) and detected by GaAsP photomultiplier tubes (H10770PA-40, Hamamatsu). Single Z-
plane images (512 x 512 pixels) were acquired serially at 7.5 Hz (30 Hz averaged every 4 frames)
using ScanImage 5 software (Vidrio Technologies). Laser power measured at the objective was
60 - 90 mW. On average, 14 imaging fields (305 µm x 305 µm) were obtained per mouse at depths
of 110 - 250 µm below the cortical surface[90]. If there was >25% XY overlap, imaging fields were
required to be at least 20 µm apart in depth to avoid repeated imaging of the same cells. After
completion of all imaging experiments, the mouse was euthanized and the brain was collected to
perform histology.
647

Histological localization of imaging fields
The brain was extracted and fixed overnight in 4% paraformaldehyde. After flattening, the cortex
was sunk in 30% sucrose and sectioned at 50-60 µm parallel to the surface. Cytochrome oxidase
(CO) staining showed surface vasculature in the most superficial tangential section as well as
boundaries of barrels in L4. Histological sections were manually aligned using Fiji[91], and imaging
fields were localized in the whisker map aided by the surface blood vessels imaged at the
beginning of each session. The centroid of each of the nine anatomical barrels corresponding to
whiskers stimulated in each session and the XY coordinates of all imaged cells were localized
relative to barrel boundaries. A cell was located within a specific barrel column if >50% of its pixels
were within its boundaries, and cells outside barrel boundaries were classified as septal cells.
Major and minor axes of all barrels were averaged to calculate the mean barrel width.
659

Image processing, ROI selection, and ΔF/F calculation
Custom MATLAB pipeline code (Ramamurthy et al, 2023[44]; adapted from LeMessurier, 2019[92])
was used for image processing. Correction for slow XY drift was performed using dftregistration[93].
Regions-of-interest (ROIs) were manually drawn as ellipsoid regions over the somata of neurons
visible in the average projection across the full imaging movie after registration. Mean
fluorescence of the pixels in each ROI was calculated to obtain the raw fluorescence time series.
For PYR cell imaging, neuropil masks were created as 10 pixel-wide rings beginning two pixels
from the somatic ROI, excluding any pixels correlated with any somatic ROI (r>0.2). Mean

fluorescence of neuropil masks was scaled by 0.3 and subtracted from raw somatic ROI
fluorescence. Neuropil subtraction was not performed for VIP cells, which were spatially well-
separated. The mean fluorescence time series was converted to ΔF/F for each ROI, defined as
$(F_t-F_0)/F_0$, where F_0 is the 20th percentile of fluorescence across the entire imaging movie and F_t
is the fluorescence on each frame.
673
Quantification of whisker-evoked responses
Whisker-evoked ΔF/F signal on Go trials was quantified in a post-stimulus analysis window (7
frames, 0.799 s), relative to pre-event baseline window (2 frames, 0.270 s). Whisker responses
were measured as (mean ΔF/F in the post-stimulus window – mean ΔF/F in the baseline window)
for each ROI. On NoGo trials, the ΔF/F analysis was aligned to the NoGo stimulus (dummy piezo
deflection). Whisker responses for each cell were normalized by z-scoring to prestimulus baseline
activity. For some analyses, each ROI's Go-NoGo response magnitude to every whisker was also
calculated as (median whisker-evoked ΔF/F signal across Go trials – median ΔF/F signal across
NoGo trials). Trials aborted due to licks occurring during the post-stimulus window (0 - 0.799 s)
were excluded from analyses. If at least one whisker produced a significant response above
baseline activity (permutation test), the cell was considered to be whisker-responsive. This was
done by combining the whisker-evoked ΔF/F signal distribution on Go trials with the ΔF/F signal
distribution on NoGo trials, randomly splitting the combined distribution into two groups and
comparing the difference in their means to the true Go-NoGo distribution difference (10,000
iterations). Differences greater than the 95th percentile of the permuted distribution were
assessed as significant. The nine whisker response p-values were corrected for multiple
comparisons (False Discovery Rate correction[94]). Cells without a positive ΔF/F response to at
least one whisker were considered non-responsive.

The standard method for assessing whisker-responsiveness used all trials belonging to
each session (combining trials across all history conditions). In the analysis of attentional
modulation of receptive fields, the significance of whisker responses was separately assessed
using only trials in the Prior NoGo category and compared to trials in Prior >1 Hit condition, which
allowed us to test whether there was history-dependent acquisition of whisker-evoked responses
by previously non-responsive cells.
698
Definition of each cell's columnar whisker (CW) and best whisker (BW)
Each cell's anatomical home column was determined by histological localization of the cell relative
to barrel column boundaries. For cells located within column boundaries, the CW was the whisker
corresponding to its anatomical home location. For septa-related cells (i.e., those outside of
column boundaries and above a L4 septum), the CW was the whisker corresponding to the
nearest barrel column. The best whisker (BW) was defined for each cell as the whisker that
evoked the numerically highest magnitude response.
706
Attention Modulation Index (AMI)
Multiple AMI metrics were used to quantify attentional modulation of whisker response magnitude
in individual cells. The definitions were:

AMI (>1Hit-NoGo):

$$AMI_{>1HitSame-NoGo} = \frac{Go_{Prior>1HitSame} - Go_{PriorNoGo}}{|Go_{Prior>1HitSame} + Go_{PriorNoGo}|}$$

$$AMI_{>1HitDiff-NoGo} = \frac{Go_{Prior>1HitDiff} - Go_{PriorNoGo}}{|Go_{Prior>1HitDiff} + Go_{PriorNoGo}|}$$

AMI(>1HitSame->1HitDiff):

$$AMI_{>1HitSame->1HitDiff} = \frac{Go_{Prior>1HitSame} - Go_{Prior>1HitDiff}}{|Go_{Prior>1HitSame} + Go_{Prior>1HitDiff}|}$$

where Go = mean whisker-evoked ΔF/F for current Go trials (on any whisker) with the specified
trial history.

Attentional modulation of receptive fields
Population average 9-whisker receptive fields were constructed centered on the CW, and included
the CW plus the 8 immediately adjacent whiskers. A separate population average receptive field
was calculated for each trial history (**Fig. 4A**). These represent the average tuning of cells within
each whisker column, following each trial history. To test for shifts in receptive fields by attention
to specific whiskers, we first computed the center of mass (CoM) of the population average
receptive fields for each history condition. CoM was calculated in a Cartesian CW-centered
whisker space, as defined in **Fig. S5B**. The CW position is considered the origin in this space.
Receptive field shifts associated with prior trial history were visualized as vectors from CoM
measured after NoGo trials, to CoM measured after >1 prior hit to specific whiskers.

To quantify the receptive field shift (ΔRF CoM) for individual cells, we defined the *attention*
axis as the axis connecting the CW position to the attended whisker position in the Cartesian CoM
space. We projected the CoM$_{PriorNoGo}$ and CoM$_{Prior>1Hit}$ onto this axis, and computed the receptive
field shift (ΔRF CoM) as the distance between these projected positions normalized to the
distance from CW to attended whisker along the attention axis (**Fig. S5C**). Since not all whisker
positions could be sampled for all cells across history conditions, ΔRF CoM was quantified only
for the subset of cells for which at least 6 of the 9 whisker positions were sampled in both Prior
>1 Hit and Prior NoGo conditions. Only response magnitudes at whisker positions sampled in
both Prior >1 Hit and Prior NoGo conditions for any given cell contributed to the CoMs computed
for that cell. The RF shift was computed separately for each attended whisker position that was
sampled for a given cell, and then averaged across these attended whisker positions to generate
a single RF shift metric for that cell.

Somatotopic profile of attentional modulation
To quantify the somatotopic profile of attentional modulation in S1 (**Fig. 3K**), we considered each
of the 9 tested whiskers separately. For each whisker (termed the reference whisker), every cell
was placed in a spatial bin representing its distance to the center of the reference whisker column
in S1. Both columnar and septal-related cells were included. Mean whisker response magnitude
was calculated, separated by trial history, in each bin. This was repeated for all 9 reference
whiskers, and **Fig. 3K** shows the average response. Thus, the Prior NoGo trace reflects normal
somatotopy, i.e., the normal point representation of an average whisker. The somatotopic profile

of attentional modulation is evident as the difference between other history conditions and the
Prior NoGo condition.

A similar binning procedure was used to calculate the somatotopic profile of AMI
modulation across S1 columns (**Fig. S4A**).

Imaging analysis for VIP cells
Analysis of VIP cell responses was performed similarly to PYR cells, except that neuropil
subtraction was not performed. To separate whisker-evoked VIP responses from slow trends in
VIP baseline activity related to whisker motion, body motion and arousal in the ITI (Ramamurthy
et al., 2023[44] and **Fig. 2**) we applied linear baseline detrending. For detrending, the median pre-
stimulus baseline trace (in a 1.07-sec window) was calculated across Go trials (aligned to stimulus
onset time) and NoGo trials (aligned to dummy piezo onset time). A line was fit to this median
trace. This line was extrapolated and subtracted from each individual trial to yield the full peri-
stimulus trace. This linear detrending was done separately for each history condition, due to the
differences in pre-stimulus slopes for each condition. Note that linear detrending for prior same
and prior different categories in each reward condition was identical. Analysis of history effects on
VIP whisker responses (**Fig. 7F-G**) was performed after linear detrending. While VIP cells did not
show whisker-specific attentional modulation (**Fig. 7F-G**), we verified that PYR cells still showed
whisker-specific attentional boosting after detrending with the same methods (data not shown).

Neural decoding from population activity on single trials
We used a generalized linear model (GLM, Matlab 'glmfit') to predict the presence of a whisker
stimulus from the trial-by-trial mean population activity of whisker-responsive L2/3 PYR cells in
2p imaging experiments. For each session, mean ΔF/F in the post-stimulus window (0 - 0.799 s)
for each trial was calculated across all whisker-responsive ROIs that were simultaneously imaged
in a single behavioral session. This single-trial population activity was used as the predictor of
stimulus presence (current Go trial) or stimulus absence (current NoGo trial).

A separate decoder was fit from the data for each imaging session. We fitted a logistic
regression model using leave-one-out cross-validation to predict the presence of any of the 9
whiskers, and tested it on hold-out data. Training/testing datasets were randomly re-sampled
(majority class undersampled to match the minority class) to have identical numbers of trials within
each response category (Hit, Miss, CR, FA), in order to remove bias. Decoder performance was
measured as the average fraction of trials classified as containing a whisker stimulus (assessed
over 25-50 iterations) and compared to performance for a decoder trained with shuffled trial labels.

For each field, we defined the fBW (field best whisker) as the whisker which evoked the
numerically highest mean population ΔF/F. A single decoder was trained for each session to
predict any whisker from training data containing Go trials from all whiskers, as well as NoGo
trials. Decoder performance was assessed either for detecting any whisker, or just the fBW, or
just non-fBW trials.

Extracellular recordings
For surgical preparation for mice used in extracellular recording experiments, methods were
similar to that described above, except a lightweight chronic head post was affixed to the skull

using cyanoacrylate glue and Metabond, and ISOI was performed to localize D-row (D1, D2, D3)
barrel columns in S1. A 5-mm diameter glass coverslip (#1 thickness, CS-3R, Warner Instruments)
was placed over the skull, sealed with Kwik-Cast silicone adhesive (World Precision Instruments)
and dental cement. Mice recovered for 7 days prior to the start of behavioral training (~4 weeks
before recording).

The day before recording, mice were anaesthetized with isoflurane, the protective
coverslip was removed, and a craniotomy (~ 1.2 x 1.2 mm) was made over S1, centered over the
D-row (D1, D2, D3) barrel columns localized by ISOI. A plastic ring was cemented around the
craniotomy to create a recording chamber. During recording sessions, mice were anesthetized
and positioned on the rig. A reference ground was attached inside the chamber. The craniotomy
and reference wire were covered in a saline bath. Recordings were made with Neuropixels 1.0
probes using SpikeGLX software release v.20201024 (http://billkarsh.github.io/SpikeGLX/), Imec
phase30 v3.31. Acute recordings were made in external reference mode with action potentials
(AP) sampled at 30 kHz at 500x gain, and local field potential (LFP) sampled at 2.5 kHz at 250x
gain. The AP band was common average referenced and band-pass filtered from 0.3 kHz to
6 kHz. The Neuropixels probe was mounted on a motorized stereotaxic micromanipulator (MP-
285, Sutter Instruments) and advanced through the dura mater (except in cases where the dura
had detached during the craniotomy). To reduce insertion-related mechanical tissue damage and
to increase the single unit yield, the probe was lowered with a slow insertion speed of 1-2 µm/sec.
The probe was first lowered to 700 µm, and a short 10-minute recording was conducted to map
its location in S1. After identifying the columnar whisker for the recording penetration location,
probe insertion continued until the final depth was reached. The probe was then left untouched
for ~ 20 minutes. The craniotomy was sealed after probe insertion with silicone sealant (Kwik-
Cast, World Precision Instruments) to prevent drying. Anesthesia was then discontinued, and
mice were allowed to fully wake up before recording.

After recording was complete for the day, the probe was removed, the craniotomy was
sealed with silicone sealant (Kwik-Cast, World Precision Instruments), and the recording chamber
was sealed with a cover glass and a thin layer of dental cement. 3-4 sequential days of recording
were performed in each mouse, with the probe located in a different whisker column in S1 on
each day. On the final recording day, a Neuropixels probe was coated with red-fluorescent DiI
(1,1′-Dioctadecyl-3,3,3′,3′-tetramethylindocarbocyanine perchlorate; Sigma-Aldrich) dissolved in
100% ethanol, 1-2mg/mL, which was allowed to partially dry on probe before probe insertion. The
probe was briefly inserted to deposit DiI at recording and several fiducial sites. The mouse was
euthanized and the brain was extracted, sectioned tangentially to the pial surface, and processed
to stain for cytochrome oxidase (CO). DiI deposition sites in L2/3 were localized relative to column
boundaries in CO from L4 (**Fig. S6B**).

The columnar location of each recording penetration was determined from DiI marks on
the last recording day, plus relative locations of other penetrations based on reference images of
surface vasculature and microdrive coordinates. The laminar depth of each recording was
determined from current source density (CSD) and LFP power spectrum analysis, as described
below.
836

Analysis of extracellular recording data

<u>Spike Sorting</u>

Spike sorting was performed by automatic clustering using Kilosort3 followed by manual curation using the 'phy' GUI (https://github.com/kwikteam/phy). Isolated units were manually inspected for mean spike waveform, stability over time, and inter-spike interval refractory period violations (we required that < 2% of intervals < 1.5 ms). Only well-isolated single units were analyzed. Single units were classified as regular-spiking or fast-spiking based on trough-to-peak duration of the spike waveform at the highest-amplitude recording channel, with a separation criterion of 0.45 ms[95] (**Fig. S6C**). Only data from regular spiking units was analyzed here.

<u>Layer assignment using CSD and LFP power spectrum analysis</u>

To calculate the CSD for each recording penetration, the whisker stimulus-evoked local field potential (LFP, 500Hz low-pass) was calculated for each Neuropixels channel. LFP traces were normalized to correct for variations in channel impedance and were interpolated between channels (20 um site spacing, 1.6-2x interpolation) prior to calculating the second spatial derivative, which defines the CSD[96]. For visualization, CSDs were convolved with a 2D (depth x time) Gaussian, which revealed depth-restricted regions of current sources and sinks in response to each whisker stimulus. The L4-L5A boundary was defined from CSD as the zero-crossing between the most negative current sink (putative L4) and the next deeper current source (putative L5A) (**Fig. S6F**). To estimate the brain surface location (defining the top of L1), we computed the LFP power spectrum as a function of channel depth. A sharp increase in low-frequency LFP power marked the brain surface, which was used to verify correct selection of the L4-L5A boundary from CSD. Each cortical layer was then assigned boundary depths based on layer thicknesses reported in Lefort et al., 2009[97].

Whisker response quantification and attentional modulation for spike recordings

Firing rates for Go and NoGo trials were quantified in a 0.5 s window after stimulus onset (lick-free window). Units were classified as whisker-responsive or non-responsive by testing for greater firing rate on Go vs NoGo trials, using a permutation test, as for the 2p imaging data. To quantify whisker-evoked response magnitude, firing rate for each unit was z-scored relative to pre-stimulus baseline. Peristimulus time histograms (PSTHs) were constructed using 10-ms bins aligned to stimulus onset. AMI metrics were used to quantify attentional modulation of whisker-evoked spiking, and were calculated exactly as for 2p imaging data, but from z-scored whisker-evoked firing rate in the post-stimulus window.

Statistical analysis of summary data

Statistics were performed in Matlab. Sample size (n) and p-value for each analysis are reported in the figure panel, with the statistical test reported in the figure legend, or sometimes in the *Results* section. Permutation tests for the difference in means were used to assess differences in mean between two groups (referred to in brief as 'permutation test') or differences in the mean of a distribution relative to zero. Binomial exact test was used to assess the statistical significance of deviations from the expected distribution of observations into two categories. Summary data are reported as mean ± SEM, unless otherwise specified. Population means were compared using permutation tests, corrected for multiple comparisons (False Discovery Rate correction[94])

as needed. To compare population means with multiple subgroups, we first assessed whether a main effect was present using a permutation test, and then performed post hoc tests for pairwise means, correcting for multiple comparisons (with False Discovery Rate correction). We used a significance level (alpha) of 0.05. All statistical tests reported in the *Results* use n of cells, but we also verified that all major behavioral and neural effects were consistent across individual mice. We show individual mouse data for comparison with population data summaries. To account for inter-individual variability, we also verified all key results using a linear mixed effects model with the formula:

$$Response\ Variable \sim Fixed\ Effect + (1|Mouse\ Sex) + (1|Mouse\ ID)$$

where mouse sex and mouse ID are modeled as random effects.

DATA AVAILABILITY

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

ACKNOWLEDGMENTS

This work was supported by NIH Grant 2 R01 NS092367 (D.E.F.) and NIH Grant 5 K99 NS129753-02 (D.L.R.).

AUTHOR CONTRIBUTIONS

D.L.R. and D.E.F. designed the study. D.L.R., L.R., C.C., S.L., and A.C. performed the experiments. D.L.R., L.R., C.C., and S.L. analyzed the data. D.L.R., L.R., and D.E.F. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Maunsell, J.H.R. Neuronal Mechanisms of Visual Attention. Annu Rev Vis Sci 1, 373-391 (2015).
2. Hromádka, T. & Zador, A.M. Toward the mechanisms of auditory attention. Hear Res 229, 180-185 (2007).
3. Gallace, A. & Spence, C. 147 Tactile attention. in In Touch with the Future: The sense of touch from cognitive neuroscience to virtual reality 0 (Oxford University Press, 2014).
4. Nobre, A.C. & Stokes, M.G. Premembering Experience: A Hierarchy of Time-Scales for Proactive Attention. Neuron 104, 132-146 (2019).
5. Awh, E., Belopolsky, A.V. & Theeuwes, J. Top-down versus bottom-up attentional control: a failed theoretical dichotomy. Trends in cognitive sciences 16, 437-443 (2012).
6. Theeuwes, J. & Failing, M. Attentional Selection: Top-Down, Bottom-Up and History-Based Biases. (Cambridge University Press, Cambridge, 2020).

7. Meyer, K.N., Sheridan, M.A. & Hopfinger, J.B. Reward history impacts attentional orienting and inhibitory control on untrained tasks. Atten Percept Psychophys 82, 3842-3862 (2020).

8. Failing, M. & Theeuwes, J. Selection history: How reward modulates selectivity of visual attention. Psychonomic Bulletin & Review 25, 514-538 (2018).

9. Anderson, B.A. & Britton, M.K. Selection history in context: Evidence for the role of reinforcement learning in biasing attention. Atten Percept Psychophys 81, 2666-2672 (2019).

10. Anderson, B.A., Kim, H., Kim, A.J., Liao, M.-R., Mrkonja, L., Clement, A., and Grégoire, L. The past, present, and future of selection history. Neuroscience & Biobehavioral Reviews 130, 326-350. https://doi.org/10.1016/j.neubiorev.2021.09.004. (2021).

11. Chelazzi, L., Perlato, A., Santandrea, E. & Della Libera, C. Rewards teach visual selective attention. Vision Research 85, 58-72 (2013).

12. Anderson, B.A. A value-driven mechanism of attentional selection. J Vis 13 (2013).

13. Anderson, B.A. Neurobiology of value-driven attention. Current Opinion in Psychology 29, 27-33 (2019).

14. Anderson, B.A., Laurent, P.A. & Yantis, S. Value-driven attentional capture. Proceedings of the National Academy of Sciences 108, 10367 (2011).

15. Addleman, D.A. & Jiang, Y.V. Experience-Driven Auditory Attention. Trends in Cognitive Sciences 23, 927-937 (2019).

16. Chen, D. & Hutchinson, J.B. What Is Memory-Guided Attention? How Past Experiences Shape Selective Visuospatial Attention in the Present. Curr Top Behav Neurosci 41, 185-212 (2019).

17. Lynn, J., and Shin, M. Strategic top-down control versus attentional bias by previous reward history. Atten Percept Psychophys 77, 2207-2216. 10.3758/s13414-015-0939-9. (2015).

18. Noudoost, B., Chang, M.H., Steinmetz, N.A. & Moore, T. Top-down control of visual attention. Curr Opin Neurobiol 20, 183-190 (2010).

19. Matteucci, G., et al. Cortical sensory processing across motivational states during goal-directed behavior. Neuron 110, 4176-4193.e4110 (2022).

20. Luo, T.Z. & Maunsell, J.H.R. Attention can be subdivided into neurobiological components corresponding to distinct behavioral effects. Proceedings of the National Academy of Sciences 116, 26187-26194 (2019).

21. Krauzlis, R.J., Wang, L., Yu, G. & Katz, L.N. What is attention? Wiley Interdiscip Rev Cogn Sci 14, e1570 (2023).

22. Luo, T.Z. & Maunsell, J.H. Neuronal Modulations in Visual Cortex Are Associated with Only One of Multiple Components of Attention. Neuron 86, 1182-1188 (2015).

23. Luo, T.Z. & Maunsell, J.H.R. Attentional Changes in Either Criterion or Sensitivity Are Associated with Robust Modulations in Lateral Prefrontal Cortex. Neuron 97, 1382-1393.e1387 (2018).

24. Peck, C.J., Jangraw, D.C., Suzuki, M., Efem, R. & Gottlieb, J. Reward modulates attention independently of action value in posterior parietal cortex. J Neurosci 29, 11182-11191 (2009).

25. Desimone, R. & Duncan, J. Neural mechanisms of selective visual attention. Annu Rev Neurosci 18, 193-222 (1995).

26. Cohen, M.R. & Maunsell, J.H.R. A Neuronal Population Measure of Attention Predicts Behavioral Performance on Individual Trials. The Journal of Neuroscience 30, 15241 (2010).

27. Reimer, J., et al. Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. Neuron 84, 355-362 (2014).

28. Reimer, J., et al. Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nature Communications 7, 13289 (2016).

29. Mathis, A., et al. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nature Neuroscience 21, 1281-1289 (2018).

30. Womelsdorf, T., Anton-Erxleben, K., Pieper, F. & Treue, S. Dynamic shifts of visual receptive fields in cortical area MT by spatial attention. Nature Neuroscience 9, 1156-1160 (2006).

31. Womelsdorf, T., Anton-Erxleben, K. & Treue, S. Receptive field shift and shrinkage in macaque middle temporal area through attentional gain modulation. J Neurosci 28, 8934-8944 (2008).

32. Anton-Erxleben, K., Stephan, V.M. & Treue, S. Attention reshapes center-surround receptive field structure in macaque cortical area MT. Cereb Cortex 19, 2466-2478 (2009).

33. He, S., Cavanagh, P. & Intriligator, J. Attentional resolution and the locus of visual awareness. Nature 383, 334-337 (1996).

34. Yang, H., Kwon, S.E., Severson, K.S. & O'Connor, D.H. Origins of choice-related activity in mouse somatosensory cortex. Nat Neurosci 19, 127-134 (2016).

35. Yamashita, T. & Petersen, C. Target-specific membrane potential dynamics of neocortical projection neurons during goal-directed behavior. Elife 5 (2016).

36. Jun, J.J., et al. Fully integrated silicon probes for high-density recording of neural activity. Nature 551, 232-236 (2017).

37. Lee, S., Kruglikov, I., Huang, Z.J., Fishell, G. & Rudy, B. A disinhibitory circuit mediates motor integration in the somatosensory cortex. Nat Neurosci 16, 1662-1670 (2013).

38. Fu, Y., et al. A cortical circuit for gain control by behavioral state. Cell 156, 1139-1152 (2014).

39. Zhang, S., et al. Selective attention. Long-range and local circuits for top-down modulation of visual cortex processing. Science 345, 660-665 (2014).

40. Batista-Brito, R., Zagha, E., Ratliff, J.M. & Vinck, M. Modulation of cortical circuits by top-down processing and arousal state in health and disease. Current opinion in neurobiology 52, 172-181 (2018).

41. Xia, R., Chen, X., Engel, T.A. & Moore, T. Common and distinct neural mechanisms of attention. Trends Cogn Sci 28, 554-567 (2024).

42. Myers-Joseph, D., Wilmes, K.A., Fernandez-Otero, M., Clopath, C. & Khan, A.G. Disinhibition by VIP interneurons is orthogonal to cross-modal attentional modulation in primary visual cortex. Neuron 112, 628-645.e627 (2024).

43. Rahmatullah, N., et al. Hypersensitivity to Distractors in Fragile X Syndrome from Loss of Modulation of Cortical VIP Interneurons. The Journal of Neuroscience 43, 8172 (2023).

44. Ramamurthy, D.L., et al. VIP interneurons in sensory cortex encode sensory and action signals but not direct reward signals. Curr Biol 33, 3398-3408.e3397 (2023).

45. Musall, S., Kaufman, M.T., Juavinett, A.L., Gluf, S. & Churchland, A.K. Single-trial neural dynamics are dominated by richly varied movements. Nature Neuroscience 22, 1677-1686 (2019).

46. Schmidt, F., Haberkamp, A. & Schmidt, T. Dos and don'ts in response priming research. Adv Cogn Psychol 7, 120-131 (2011).

47. Akrami, A., Kopec, C.D., Diamond, M.E. & Brody, C.D. Posterior parietal cortex represents sensory history and mediates its effects on behaviour. Nature 554, 368-372 (2018).

48. Hachen, I., Reinartz, S., Brasselet, R., Stroligo, A. & Diamond, M.E. Dynamics of history-dependent perceptual judgment. Nature Communications 12, 6036 (2021).

49. Waiblinger, C., McDonnell, M.E., Borden, P.Y. & Stanley, G.B. Emerging experience-dependent dynamics in primary somatosensory cortex reflect behavioral adaptation. bioRxiv, 2021.2001.2029.428886 (2021).

50. Lak, A., et al. Reinforcement biases subsequent perceptual decisions when confidence is low, a widespread behavioral phenomenon. eLife 9, e49834 (2020).

51. Boboeva, V., Pezzotta, A., Clopath, C. & Akrami, A. Unifying network model links recency and central tendency biases in working memory. eLife 12, RP86725 (2024).

52. Cicchini, G.M., Mikellidou, K. & Burr, D.C. Serial Dependence in Perception. Annual Review of Psychology 75, 129-154 (2024).

53. Samonds, J.M., Lieberman, S. & Priebe, N.J. Motion Discrimination and the Motion Aftereffect in Mouse Vision. eNeuro 5 (2018).

54. Urai, A.E., de Gee, J.W., Tsetsos, K. & Donner, T.H. Choice history biases subsequent evidence accumulation. eLife 8, e46331 (2019).

55. Ashwood, Z.C., et al. Mice alternate between discrete strategies during perceptual decision-making. Nat Neurosci 25, 201-212 (2022).

56. Roy, N.A., Bak, J.H., Akrami, A., Brody, C.D. & Pillow, J.W. Extracting the dynamics of behavior in sensory decision-making experiments. Neuron 109, 597-610.e596 (2021).

57. Reynolds, J.H., Pasternak, T. & Desimone, R. Attention increases sensitivity of V4 neurons. Neuron 26, 703-714 (2000).

58. Briggs, F., Mangun, G.R. & Usrey, W.M. Attention enhances synaptic efficacy and the signal-to-noise ratio in neural circuits. Nature 499, 476-480 (2013).

59. Cohen, M.R. & Maunsell, J.H. Attention improves performance primarily by reducing interneuronal correlations. Nature neuroscience 12, 1594-1600 (2009).

60. Luck, S.J., Chelazzi, L., Hillyard, S.A. & Desimone, R. Neural Mechanisms of Spatial Selective Attention in Areas V1, V2, and V4 of Macaque Visual Cortex. Journal of Neurophysiology 77, 24-42 (1997).

61. Silver, M.A., Ress, D. & Heeger, D.J. Topographic maps of visual spatial attention in human parietal cortex. J Neurophysiol 94, 1358-1371 (2005).

62. Tootell, R.B., et al. The retinotopy of visual spatial attention. Neuron 21, 1409-1422 (1998).

63. Woldorff, M.G., et al. Retinotopic organization of early visual spatial attention effects as revealed by PET and ERPs. Hum Brain Mapp 5, 280-286 (1997).

64. Bollimunta, A., Bogadhi, A.R. & Krauzlis, R.J. Comparing frontal eye field and superior colliculus contributions to covert spatial attention. Nat Commun 9, 3553 (2018).

65. Staiger, J.F. & Petersen, C.C.H. Neuronal Circuits in Barrel Cortex for Whisker Sensory
Perception. Physiol Rev 101, 353-415 (2021).

66. Engel, T.A., et al. Selective modulation of cortical state during spatial attention. Science
354, 1140-1144 (2016).

67. Hattori, R., Danskin, B., Babic, Z., Mlynaryk, N. & Komiyama, T. Area-Specificity and
Plasticity of History-Dependent Value Coding During Learning. Cell 177, 1858-1872.e1815
(2019).

68. Hwang, E.J., Dahlen, J.E., Mukundan, M. & Komiyama, T. History-based action selection
bias in posterior parietal cortex. Nature Communications 8, 1242 (2017).

69. Findling, C., et al. Brain-wide representations of prior information in mouse decision-
making. bioRxiv, 2023.2007.2004.547684 (2023).

70. Miller, K.J., Botvinick, M.M. & Brody, C.D. Value representations in the rodent orbitofrontal
cortex drive learning, not choice. eLife 11, e64575 (2022).

71. Marmor, O., Pollak, Y., Doron, C., Helmchen, F. & Gilad, A. History information emerges
in the cortex during learning. eLife 12, e83702 (2023).

72. Banerjee, A., et al. Value-guided remapping of sensory cortex by lateral orbitofrontal
cortex. Nature 585, 245-250 (2020).

73. Chéreau, R., et al. Dynamic perceptual feature selectivity in primary somatosensory cortex
upon reversal learning. Nature Communications 11, 3245 (2020).

74. Petty, G.H. & Bruno, R.M. Attentional modulation of secondary somatosensory and visual
thalamus of mice. eLife. 13:RP97188 (2024).

75. Kastner, S. & Arcaro, M.J. The Thalamus in Attention. in The Thalamus (ed. M.M. Halassa)
324-339 (Cambridge University Press, Cambridge, 2022).

76. Pi, H.J., et al. Cortical interneurons that specialize in disinhibitory control. Nature 503, 521-
524 (2013).

77. Hu, F. & Dan, Y. An inferior-superior colliculus circuit controls auditory cue-directed visual
spatial attention. Neuron 110, 109-119.e103 (2022).

78. Speed, A. & Haider, B. Probing mechanisms of visual spatial attention in mice. Trends in
Neurosciences (2021).

79. Speed, A., Del Rosario, J., Mikail, N. & Haider, B. Spatial attention enhances network,
cellular and subthreshold responses in mouse visual cortex. Nature Communications 11,
505 (2020).

80. Kanamori, T. & Mrsic-Flogel, T.D. Independent response modulation of visual cortical
neurons by attentional and behavioral states. Neuron 110, 3907-3918.e3906 (2022).

81. McBride, E.G., Lee, S.J. & Callaway, E.M. Local and Global Influences of Visual Spatial
Selection and Locomotion in Mouse Primary Visual Cortex. Curr Biol 29, 1592-1605.e1595
(2019).

82. Lehnert, J., et al. Visual attention to features and space in mice using reverse correlation.
Current Biology 33, 3690-3701.e3694 (2023).

83. You, W.-K. & Mysore, S.P. Endogenous and exogenous control of visuospatial selective
attention in freely behaving mice. Nature Communications 11, 1986 (2020).

84. Goldstein, S., Wang, L., McAlonan, K., Torres-Cruz, M. & Krauzlis, R.J. Stimulus-driven
visual attention in mice. Journal of Vision 22, 11-11 (2022).

85. Drew, P.J. & Feldman, D.E. Intrinsic Signal Imaging of Deprivation-Induced Contraction of
Whisker Representations in Rat Somatosensory Cortex. Cerebral Cortex 19, 331-348
(2009).
86. Wang, H.C., LeMessurier, A.M. & Feldman, D.E. Tuning instability of non-columnar
neurons in the salt-and-pepper whisker map in somatosensory cortex. Nature
Communications 13, 6611 (2022).
87. Landers, M., Pytte, C. & Zeigler, H.P. Reversible blockade of rodent whisking: Botulinum
toxin as a tool for developmental studies. Somatosensory & Motor Research 19, 358-363
(2002).
88. Yang, H., Kwon, S.E., Severson, K.S. & O'Connor, D.H. Origins of choice-related activity
in mouse somatosensory cortex. Nature Neuroscience 19, 127-134 (2016).
89. Green, D.M. & Swets, J.A. Signal detection theory and psychophysics (John Wiley,
Oxford, England, 1966).
90. Pronneke, A., et al. Characterizing VIP Neurons in the Barrel Cortex of VIPcre/
tdTomato Mice Reveals Layer-Specific Differences. Cereb Cortex 25, 4854-4868 (2015).
91. Schindelin, J., et al. Fiji: an open-source platform for biological-image analysis. Nature
methods 9, 676-682 (2012).
92. LeMessurier, A.M. (2019). Imaging_analysis_pipeline. GitHub 5a21c9f.
https://github.com/alemessurier/imaging_analysis_pipeline
93. Guizar-Sicairos, M., Thurman, S.T. & Fienup, J.R. Efficient subpixel image registration
algorithms. Opt. Lett. 33, 156-158 (2008).
94. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological) 57, 289-300 (1995).
95. Barthó, P., et al. Characterization of neocortical principal cells and interneurons by network
interactions and extracellular features. J Neurophysiol 92, 600-608 (2004).
96. Freeman, J.A. & Nicholson, C. Experimental optimization of current source-density
technique for anuran cerebellum. J Neurophysiol 38, 369-382 (1975).
97. Lefort, S., Tomm, C., Floyd Sarria, J.C. & Petersen, C.C.H. The Excitatory Neuronal
Network of the C2 Barrel Column in Mouse Primary Somatosensory Cortex. Neuron 61,
301-316 (2009).

FIGURES AND FIGURE LEGENDS

Figure 1. Recent reward history cues spatially specific attention for whisker touch. A. Whisker detection task in head-fixed mice. **B.** Trial structure. Delay period was 0, 500, or 1000 ms in different mice (see **Figure S1**). Bottom, Trial types and trial outcomes. **C.** Example trial sequence showing interleaved Go and NoGo trials, with the identity of the deflected whisker on each Go trial chosen randomly. Water drop indicates successful Hit and reward. **D.** Mean effect of trial history on detection sensitivity (d-prime) on the current trial. Error bars are SEM across sessions. One or more prior Hits to the same whisker increased d' for detecting that whisker on the current trial ('attend toward'), while prior Hits to a different whisker decreased d' (attend away). P-values: (1) from permutation test (prior same 1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior different >1 Hit: p = 1e-4), (2) from linear mixed effects model with fixed effect of prior history class (p = 8.7e-27). **E.** Same effect in a single example mouse. Left, the underlying effect on hit rate for 5 example sessions (thin lines) and mean ± SEM for all 34 sessions. Right, d' for this mouse from 34 sessions. **F.** d' did not increase when the prior trial was an unrewarded hit to the same whisker or earned very small reward (<4%: p = 0.49, >4%: p = 1e-4). P-values are for difference in d' between prior same and prior different conditions (permutation test). **G.** Effect of >1 prior hits on Hit Rate and FA rate on the current trial. Connected symbols are the three history conditions for one mouse. Gray lines, receiver-operating characteristic (ROC) curves for different d' levels. Prior hits to the same whisker drove a whisker-specific increase in hit rate over FA rate that improved d'. **H.** Behavioral shift in d' for each individual mouse. **I.** Behavioral shifts in criterion (c) were also observed, but were less whisker-specific. **J.** Whisker-specific changes in d' ($\Delta d'$) and c (Δc) for each mouse (number is mouse identity). We focus on $\Delta d'$ to index whisker-specific attention. **K.** Spatial gradient of the attention effect. The facial position of the prior Hit whisker is plotted on the x-axis, relative to the current trial whisker. P-values are for difference from prior same (same row: p = 0.27, same arc: 1e-3, diagonal: 1e-4, further: 1e-4, permutation test). Bottom, each position shown schematically. **L.** Temporal profile of the whisker-specific attention effect (3-5 s: 5e-3, 5-7 s: 1e-4, 7-9 s: 1e-4, 9-11 s: 0.17, >11s: 0.15, permutation test), P-values are from permutation test. Bottom, Inter-Go-trial intervals sampled in the task. **M.** Flexible targeting of attention. x-axis represents identity of whisker in current Go trial, grouped into C, D, or E-row whiskers (left), or arcs 1, 2 or 3 (right). For all of these whiskers, prior reward history boosts detection in a whisker-specific way. Bottom, each row or arc shown schematically. P-values are for prior same vs prior different > 1 hit (p = 1e-4 for all whisker positions, permutation test). See also **Figure S1**.

Figure 1. Recent reward history cues spatially specific attention for whisker touch.

Figure 2. Whisker motion, body motion, and arousal do not account for whisker-specific behavioral effects. A. Mean whisker motion, platform motion (proxy for body motion), and pupil size traces, obtained using DeepLabCut from behavioral videos, across 43,647 trials, 74 sessions in 9 mice. Each panel shows the last 3 seconds of the intertrial interval (ITI) period after the prior trial, plus the stimulus period of the current trial. Traces and shading are mean ± SEM across all trials. Dashed line, stimulus onset (Go trials) or dummy piezo onset (NoGo trials). Prior trial identity is indicated by the Prior trial history. Δ whisker position and Δ platform position traces were zeroed to the mean of positions in a 0.25-sec window prior to stimulus onset (t = 0). Pupil size was normalized within each session to the mean pupil size across the whole session. Bottom row, lick histogram. **B.** Mean whisker movement, body movement, and pupil size change during the stimulus period. Prior Hits increased stimulus-evoked whisker and body motion on subsequent trials (Δ whisker motion, prior >1 hit same vs prior NoGo: p = 1e-4, prior >1 hit same vs >1 hit different: p = 0.39; Δ body motion, prior >1 hit same vs prior NoGo: p = 1e-4, prior >1 hit same vs >1 hit different: p = 0.39; Δ pupil area, prior >1 hit same vs prior NoGo: p = 0.49, prior >1 hit same vs >1 hit different: p = 0.53). p-values are for >1 Prior Hit vs Prior NoGo (top), and >1 Prior Hit Same vs > 1 Prior Hit Different (right) (permutation test). **C.** Design of Botox experiment. Behavior was assayed on an average of 12 sessions prior to Botox injection, and 7 sessions after Botox whisker paralysis. **D.** Reward history-dependent attention effect in each of the 4 mice tested, for standard sessions (before Botox, open symbols) and Botox sessions (filled symbols). M, Mouse numbers as in **Fig. 1**. Large points are mean ± SEM across mice. Conventions as in **Fig. 1J**. Whisker paralysis did not alter the mean whisker-specific d-prime effect or criterion effect (p = 0.88, comparing same vs. different shift in Δ d-prime for standard and Botox sessions). See also **Figure S2**.

Figure 2. Whisker motion, body motion, and arousal do not account for whisker-specific behavioral effects.

Figure 3. Neural correlates of attentional capture in L2/3 pyramidal cells in S1. A-B. Example imaging field in S1 centered on the D3 column. L2/3 PYR cells are expressing GCaMP6s. Scale bar = 100 μm. **C.** Example trials showing responses to the columnar whisker D3, but only weak responses to the surround whisker C3 that were strongly modulated by prior trial history. C3 responses increased following multiple prior hits to the C3 whisker (left), but not following other history conditions (right). **D.** Mean ΔF/F traces across all neurons in 70 sessions, separated by trial history. Each trace is the mean response to any Go whisker (solid) or on NoGo trials (dash). Shading shows SEM, which is often thinner than the mean trace. **E.** Quantification of whisker-evoked ΔF/F by current trial type (Go or NoGo) and by prior trial condition (prior same 1 Hit vs prior NoGo: $p = 1e-4$, prior same >1 Hit vs prior NoGo: $p = 1e-4$, prior different >1 Hit vs prior NoGo: $p = 0.1$, prior same >1 Hit vs prior different >1 Hit: $p = 1e-4$, permutation test). Bars are SEM. **F.** Mean modulation of ΔF/F response magnitude calculated by individual mouse. Bars are SEM. **G.** Correlation between history-based modulation of PYR whisker responses and behavioral d-prime, by mouse ($r = 0.837$, $p = 6.9e-4$). **H.** $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ for each cell. Positive values denote greater response compared to the prior NoGo condition. **I.** $AMI_{>1HitSame->1HitDiff}$ for each cell. **J.** Boosting of whisker-evoked ΔF/F responses as a function of somatotopic offset between prior >1Hit whisker and the current trial whisker. Bars show d-prime (data from **Fig. 1K**, for the 6 mice used in PYR imaging) for comparison. **K.** Somatotopic organization of attentional capture. Left, the y-axis shows mean ΔF/F evoked by a reference whisker, as a function of cell position in S1 relative to the center of the reference whisker column. When calculated from Prior NoGo trials (thick black trace), this defines the classic point representation of a single whisker. This is boosted in prior same >1 hit trials, but not in prior different >1 hit trials, or prior same miss trials. Right, same data presented as difference from the Prior NoGo condition, to plot magnitude of modulation by trial history. See also **Figure S3** and **Figure S4**.

Figure 3. Neural correlates of attentional capture in L2/3 pyramidal cells in S1.

Figure 4. Attentional cueing involves receptive field shifts toward attended whiskers. A. Mean whisker-evoked ΔF/F traces for all whisker-responsive cells in all imaged columns. Center, mean ± SEM (across N = 5399 cells) for trials when prior trial was NoGo, separated by the identity of the current trial whisker. This reports the average whisker tuning curve for these neurons, in the absence of attentional cueing. Outer flanks, the whisker responses measured when prior trial history was >1Hit to the indicated attentional target whisker (thick purple trace is mean, thin traces show ± SEM). Purple fill is drawn between mean traces to aid visualization. r, u, c, d denote rostral, up, caudal, or down from CW. **B.** Mean tuning center-of-mass (CoM) when the prior trial was NoGo (black circle) vs after prior >1 Hit to each of the indicated whiskers as defined in panel **A**. CoM coordinate system is shown in **Figure S4B**. Vectors are color-coded for whether the target whisker was rostral, caudal, up, or down from the CW. **C.** Magnitude of CoM shift along the attention axis, as defined in **Figure S4C**. Negative values are shifts away from the attended whisker. The mean CoM shift was significantly greater than zero (p = 3.6e-3, permutation test). See also **Figure S5**.

Figure 4. Attentional cueing involves receptive field shifts toward attended whiskers.

Figure 5. Attentional cueing improves neural decoding of attended whiskers on single trials. A. Neural decoder design. Left, example field showing L2/3 PYR cells tuned to different best whiskers intermixed in each column, consistent with prior studies[86]. Right, on each trial, population mean ΔF/F was calculated across all responsive cells. Leave-one-out cross-validation was used to fit a logistic regression predicting stimulus presence (any whisker Go trial) or absence (NoGo trial). **B.** Decoder performance determined from held-out trials. Each dot is one session. Decoder performance is low for any whisker, because many trials are Go trials for whiskers that are not strongly represented in the imaging field. Decoder performance for fBW trials is high, because this whisker is strongly represented in the imaging field. **C-D.** Mean ΔF/F **(C)** and fraction of trials decoded as a stimulus (Go) trial **(D)**, separated by current trial type and outcome. Each dot is a session. All prior history conditions are combined in **C-D** panels (ΔF/F, HR vs Miss: $p = 1e-4$, FA vs CR: $p = 1e-4$; decoder, HR vs Miss: $p = 1e-4$, FA vs CR: $p = 9e-4$, permutation test). **E.** Mean decoder performance separated by trial history type, when current trial is any Go whisker (maroon or yellow) or a NoGo trial (gray dash). Solid lines with circle markers show decoder performance across all current trial outcomes, and dotted lines with asterisk markers show decoder performance on current hit trials only. (All trials, prior same >1 Hit vs prior NoGo: $p = 1e-4$, prior different >1 Hit vs prior NoGo: $p = 0.21$, prior same >1 Hit vs prior different >1 Hit: $p = 1e-4$; Hit trials only, prior same >1 Hit vs prior NoGo: $p = 7e-3$, prior different >1 Hit vs prior NoGo: $p = 0.05$, prior same >1 Hit vs prior different >1 Hit: $p = 2e-3$, permutation test). **F.** Same as **(E)**, but for decoder performance when current trial is an fBW Go trial or a NoGo trial. **G.** Same as **(E)**, but for decoder performance when current trial is a non-fBW Go trial or a NoGo trial. **H.** Summary of attentional modulation of decoding accuracy for the prior >1 Hit trial history condition. >1 prior Hit to a whisker improves single-trial decoding for non-fBW whiskers, but not for the fBW, in each field (non-fBW: $p = 1e-4$, fBW: $p = 0.648$). P-values are for prior same vs prior different > 1 hit (permutation test).

Figure 5. Attentional cueing improves neural decoding of attended whiskers on single trials.

Figure 6. Attentional effects on extracellular single-unit spiking in S1. **A.** Neuropixels recording during the whisker detection task. **B.** Mean PSTH for an example L2/3 RS unit recorded in the D1 whisker column, across all Go trials (D1, C2, delta, and gamma whiskers, top) and NoGo trials (bottom). Whisker-evoked responses were boosted when prior trial history was 1 or >1 Hit to the same whisker, and reduced when prior trial was >1 Hit to a different whisker. **C, E, G.** Layer-specific analysis for RS units of population mean PSTH (10 ms bins) for Go trials, by trial history. **D, F, H.** For the same units, population average whisker-evoked response on current Go trials (500-ms window) by trial history, or for current NoGo trials in an equivalent window (prior same >1 Hit vs prior different >1 Hit, L2/3: p = 6.9e-4, L4: 0.08, L5a/b: 1e-4, permutation test). **I.** Cumulative histogram of $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ values for RS units by layer. **J.** Cumulative histogram of $AMI_{>1HitSame->1HitDiff}$ by layer. **K.** Mean AMI values by layer ($AMI_{>1HitSame-NoGo}$ vs $AMI_{>1HitDiff-NoGo}$, L2/3: 0.03, L4: 0.65, L5a/b: 0.01; $AMI_{>1HitSame->1HitDiff}$, L2/3: 0.01, L4:0.51, L5a/b: 0.06)). See also **Figure S6**.

Figure 6. Attentional effects on extracellular single-unit spiking in S1.

Figure 7. L2/3 VIP cells carry a general arousal signal, but not a whisker-specific attentional signal. A. Circuit model for potential VIP cell role in arousal, movement, and attentional modulation of sensory responses in sensory cortex. **B.** Example imaging field in L2/3 in a VIP-Cre;Ai162D mouse. Scale bar = 50 µm. **C.** Mean ΔF/F traces from L2/3 VIP cells across all mice and sessions during the last 3 seconds of the intertrial interval (ITI) period after the prior trial for prior rewarded (prior 1 Hit or prior >1 Hit trials) and prior unrewarded trials (prior NoGo or prior Miss). Traces are zeroed to the last 2 frames prior to stimulus onset. The declining baseline, evident on both prior rewarded (blue) and unrewarded (gray) trials, is due to VIP cell encoding of arousal, whisker motion, and body motion which all decline in the inter-trial interval as described in a previous study[44], and shown in **Fig. 2A**. Prior rewarded trials show a steeper peak and steeper decline due to licking and reward consumption at the end of the prior trial. **D.** Mean ΔF/F traces across all VIP neurons in 103 sessions, separated by trial history. Each trace is the mean response to any Go whisker (solid) or on NoGo trials (dash). Shading shows SEM. Conventions as in **Fig. 3D**. **E.** Same data as in **(D)** after subtracting an extrapolation of the pre-stimulus baseline trend (see Methods). Whisker-evoked responses are modulated by prior hits on either the same or different whiskers, which is consistent with a global arousal effect, but not whisker-specific attentional capture. **F.** Quantification of trial history effects for L2/3 VIP cells (prior same 1 Hit vs prior NoGo: p = 0.11, prior same >1 Hit vs prior NoGo: p = 1e-3, prior different >1 Hit vs prior NoGo: p = 1e-3, prior same >1 Hit vs prior different >1 Hit: p = 0.54, permutation test). Conventions as in **Fig. 3E**. **G.** $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ for each cell. Positive values denote greater response compared to the prior NoGo condition. **H.** $AMI_{>1HitSame->1HitDiff}$ for each cell. See also **Figure S7**.

Figure 7. L2/3 VIP cells carry a general arousal signal, but not a whisker-specific attentional signal.

SUPPLEMENTARY INFORMATION

Figure S1. Further characterization of attentional capture by recent reward history. A. Behavioral performance in a single example session. Sliding *d'* was computed from Hit rate and FA rate using a 50-trial sliding window. A *d'* cutoff (black dashed line) was applied to exclude trials at the start and end of the session. **B.** Definition of trial history classes for current Go trials and current NoGo trials, using current D2 whisker trials as an example. When calculating *d'* and *c*, Go and NoGo trials with matched trial histories were selected, as illustrated. **C.** Delay periods used in different groups of mice. 0 ms (no delay period) was used only for behavioral analyses. For 500 ms delay periods (used for 2p imaging and extracellular spike recordings), and for 1000 ms delay periods (used for 2p imaging), reward volume was ramped during the delay period to encourage delayed licking (see Methods). **D.** Whisker-specific attentional cueing based on reward history was observed for mice trained with all 3 delay periods. (p = 1e-4 for prior same vs prior different for all delays, permutation test). **E.** Reaction times on Hit trials for different trial histories, for mice trained without a delay period. Prior hits decreased reaction time relative to prior NoGo (prior same >1: p = 1e-4, prior different >1 hit: p = 1e-4), but there was no significant difference between prior same and prior different conditions (prior same >1 hit vs prior different >1 hit: p = 0.21, permutation test). **F.** Trial history effect as a function of whisker stimulus amplitude (i.e., stimulus strength) on the current trial. This shows that history-dependent boosting in d' is strongest for low amplitude whisker stimuli (<50 μm: p = 0.03, 50-100 μm: p = 1e-4, 100-150 μm: p = 1e-4, 150-200 μm: p = 1e-4, 200-300 μm: p = 1e-4). P-values are for prior-same vs prior-different (permutation test). **G.** The three task designs used to set different whisker stimulus probabilities. In all designs, reward probability for a Hit on any whisker, P(reward), was 100%. In the equal probability (EqP) design, each of the 9 whiskers was presented with equal probability. In the high-probability (HiP) design, sessions were divided into one or more blocks of a few hundred trials, and within each block a single whisker (example, gray box) had a much higher probability (0.8) than the others (0.025). In the HiPSame design, all whiskers were presented at an equal overall probability, but sequential presentation was biased so that there was a higher probability (0.25-0.4) of the same whisker being repeated on the next trial, for all whisker identities. In all 3 designs, whisker identity was randomly selected on each Go trial, following the intended probability structure. **H.** Whisker-specific attentional effects cued by reward history were observed in all three task designs. 13 mice were tested with EqP and HiP designs (top), and 7 mice were tested with EqP and HiPSame designs (bottom). Related to **Figure 1**.

Figure S1. Further characterization of attentional capture by recent reward history.

Figure S2. Behavioral imaging and additional analysis of history-dependent effects on spontaneous behaviors. A. Example behavioral movie frame showing DeepLabCut tracking of whisker motion (pad + 3 whiskers), platform motion (proxy for body motion), and pupil size measurement. **B.** Mean whisker motion, body motion, and pupil size changes during the ITI period of the current trial, as a function of trial history. Conventions and p-values as in **Fig. 2B**. Prior Hits increased average whisker and body motion during the ITI period (Δ whisker motion, prior >1 hit same vs prior NoGo: p = 1e-4, prior >1 hit same vs >1 hit different: p = 1e-4; Δ body motion, prior >1 hit same vs prior NoGo: p = 1e-4, prior >1 hit same vs >1 hit different: p = 0.55; Δ pupil area, prior >1 hit same vs prior NoGo: p = 0.07, prior >1 hit same vs >1 hit different: p = 0.16). Related to **Figure 2**.

Figure S2. Behavioral imaging and additional analysis of history-dependent effects on spontaneous behaviors.

Figure S3. Additional analysis of history cueing effects on PYR activity. **A.** Example imaging field showing modulation of whisker-evoked response magnitude (color scale) by trial history. Top 2 rows, the average response to all Go whiskers. Response magnitude was increased for many cells in the prior Same >1 Hit condition, and not in the prior Different > 1 Hit condition. Bottom row, activity on NoGo trials, showing expectation or global arousal effect after > 1 prior Hit to any whisker. **B.** Mean whisker-evoked ΔF/F traces, as in **Fig. 3D**, calculated separately for whisker-responsive or non-responsive neurons (identified based on prior NoGo trials). Conventions as in **Fig 3D**. **C.** Quantification of the data in (b). Conventions as in **Fig. 3E**. Prior hits to the same whisker increase whisker-evoked responses in responsive cells (prior same 1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 0.09, prior same >1 Hit vs prior >1 Hit current NoGo: p = 1e-4, permutation test), but do not cause the appearance of whisker responses in non-responsive cells (prior same 1 Hit vs prior NoGo: p = 0.02, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 0.60, prior same >1 Hit vs prior >1 Hit current NoGo: p = 0.34, permutation test). **D.** CDFs of AMI for each cell, separated by mouse. The whisker-specific effect on response magnitude was observed in 6/6 mice. **E.** Mean ΔF/F trace for each L2/3 PYR cell, sorted by $AMI_{>1HitSame->1HitDiff}$ for each cell. Related to **Figure 2** and **Figure S4**.

Figure S3. Additional analysis of history cueing effects on PYR activity.

Figure S4. Additional analysis of somatotopic profile of attentional boosting in S1. **A.** Mean AMI for neuronal responses to a reference whisker, binned by cell location relative to the center of the reference whisker column. Left, $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$. Right, $AMI_{>1HitSame->1HitDiff}$. Conventions as in **Fig. 3K**, except that finer spatial bins are used within the reference whisker column. Whisker-specific attentional boosting was evident in a region <200 um from the reference column center ($AMI_{>1HitSame-NoGo}$ vs $AMI_{>1HitDiff-NoGo}$, 0 -50 µm: 1e-4, 50-140 µm: 4e-4,140-280 µm: 0.20, 280- 420 µm: 0.48, 420-560 µm: 0.03, 560-700 µm: 0.92, permutation test; $AMI_{>1HitSame->1HitDiff}$, 0 -50 µm: 1e-4, 50-140 µm: 1e-4,140-280 µm: 0.46, 280- 420 µm: 0.23, 420-560 µm: 0.01, 560-700 µm: 0.88, one-sample permutation test vs mean of 0). **B-C.** Trial history-dependent modulation of neural responses to the CW. For this analysis, only cells located within columnar boundaries were analyzed (i.e., septal cells were excluded). **B.** >1 prior Hit to CW increases CW-evoked responses. Conventions as in **Fig. 3D**. **C.** Quantification of trial history effects on CW responses (prior same 1 Hit vs prior NoGo: p = 0.55, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 0.76, prior same >1 Hit vs prior different >1 Hit: p = 1e-4, permutation test). Conventions as in **Fig. 3E**. Related to **Figure 2** and **Figure S3**.

Figure S4. Additional analysis of somatotopic profile of attentional boosting in S1.

Figure S5. Additional analysis of receptive field shifts. A. Modulation of mean whisker receptive field when prior trial was >1 hit to the CW (purple), relative to when prior trial was NoGo (black). Format as in **Figure 4A**. **B.** Coordinate system for calculating tuning CoM. The CW is at (0,0), and whiskers are separated by 1 unit in row or arc dimensions. **C.** Calculation of CoM tuning shift along the attention axis. The attention axis was defined as the line connecting the CW coordinate (0,0) with the coordinate of the attended whisker. The CoM measured after prior NoGo trials and after prior >1 hit to the attended whisker were projected onto this axis, and the CoM shift was calculated as distance along this axis, with positive values indicating shift toward the attended whisker. Related to **Figure 4**.

Figure S5. Additional analysis of receptive field shifts.

Figure S6. **Neuropixels recording methods. A.** Behavioral shifts in d' and c for mice that were initially trained on the 9-whisker task version, and then were tested either using 9 whiskers (open) or 4-5 whiskers only (filled). When performing the task with 4-5 whiskers, reward history still drives a whisker-specific Δ d' effect, but this is smaller than when performing with 9 whiskers (p = 1e-4, comparing same vs. different shift in Δ d-prime for 9-whisker task and 4-5 whisker task). Numbered points are individual mice. Larger symbols are mean ± SEM across mice. **B.** Example histological reconstruction of a Neuropixels recording site to the D1 whisker column. Dashed circles are probe tracks marked with DiI, outlines are from L4 barrels, white circles are reconstructed penetration sites. Asterisk marks the recording site for the example cell in **Fig. 6B** and in **S6** panel **E-G**. Scale bar = 100 μm. **C.** Classification of FS and RS units by trough-to-peak time in extracellular spike waveform. Bottom, example RS and FS spike waveform. **D.** Left, example current source density (CSD) analysis to identify laminar boundaries. See Methods for details. Right, power spectrum analysis to identify the top of L1, defined as the transition from high to low LFP power across a wide range of frequency bands. **E.** Mean spike waveform for the example L2/3 RS unit from **Fig. 6B**. Blue traces, individual spikes. **F.** Whisker-evoked spike rasters separated by current trial Go whisker (gamma, delta, D1, C2) or no whisker stimulus (NoGo trials), for the example L2/3 RS unit from **Fig. 6B**. All trial histories are combined in this plot. **G.** Whisker-evoked PSTH for the same example unit as in panel **E**. Plotting conventions as in **Fig. 6C**. Related to **Figure 6**.

Figure S6. Neuropixels recording methods.

Figure S7. Additional analysis of attentional signals in VIP cells. A. Mean raw ΔF/F trace for each L2/3 VIP cell, sorted by $AMI_{>1HitSame->1HitDiff}$ for each cell. **B.** Mean detrended ΔF/F raw trace for each L2/3 VIP cell, sorted by $AMI_{>1HitSame->1HitDiff}$ for each cell. Related to **Figure 7**.

Figure S7. Additional analysis of attentional signals in VIP cells.