Reward history guides attentional selection in whisker somatosensory (S1) cortex

Deepa L. Ramamurthy[1], Lucia Rodriguez[1,2], Celine Cen[1], Siqian Li[1], Andrew Chen[1], Daniel E. Feldman[1]

1. Department of Neuroscience and Helen Wills Neuroscience Institute, UC Berkeley
2. Neuroscience PhD Program, UC Berkeley

Corresponding author. Correspondence should be addressed to:
   Daniel E. Feldman
   142 Weill Hall #3200
   Dept. of Neuroscience
   Univ. of California, Berkeley
   Berkeley, CA 94720-3200

Pages: 53
Figures: 7
Supplementary Figures: 7
Main text: 4970 words

Abstract

The history of stimuli and rewards in the recent past drives an automatic form of attention in animals and humans in which attentional priority is given to previously rewarded stimuli. The neurobiological basis for this form of attention is unknown. In a novel whisker touch detection task, we show that mice flexibly shift attention between specific whiskers, based on this recent history of stimulus-reward association. 2-photon calcium imaging and spike recordings revealed a robust neurobiological correlate in somatosensory cortex (S1), involving topographically precise, whisker-specific boosting of L2/3 pyramidal (PYR) cell sensory responses to attend whiskers, and receptive fields shifts towards attended whiskers. L2/3 VIP interneurons were activated by whisker stimuli and arousal but did not carry a whisker-specific attentional signal, so do not mediate this form of attention. Thus, reward history drives attentional capture that is associated with dynamic, topographically precise modulation of sensory evoked activity in S1.

Introduction

Humans and animals engage attention to prioritize processing of behaviorally relevant stimuli in complex environments, including for vision[1], audition[2] and touch[3]. Multiple forms of attention exist, including classical top-down attention (voluntary attention to goal-relevant stimuli) and bottom-up attention (attentional capture by physically salient stimuli), and additional forms including the automatic capture of attention in response to recent stimulus-reward association[4-16]. In humans, this reward history-dependent attention is highly robust, such that perceptual performance is selectively enhanced for previously rewarded stimuli, even when those stimuli are no longer important to current goals[12]. This phenomenon represents a category of attentional processes often called "selection history"[7,13,16] which is distinct from traditional top-down and bottom-up attention. The effects of reward history have been termed experience-driven attention[14], value-driven attention[10], memory-guided attention[15], or attentional bias by previous reward history[16].

The neurobiology of top-down and bottom-up attention have been studied extensively in non-human primates[1,17], and have been shown to involve boosting of signal-to-noise ratio for neural encoding of attended sensory features across the cortical sensory hierarchy, including primary sensory cortex. However, the neural circuits that control attentional boosting of sensory signals remain unclear. Identifying these circuit mechanisms would be experimentally most efficient in mice, which provide powerful cell-type specific tools for investigating attentional processing. Here, we developed a new model of focal attention based on reward history in the mouse whisker tactile system, and used it to investigate the neurobiological basis of attention in sensory cortex.

We studied a simple Go-NoGo whisker touch detection task in head-fixed mice. When the whisker location of a touch stimulus was unpredictable from trial to trial, mice naturally used the history of stimulus-reward association to guide attention to specific, recently rewarded whiskers. This task generates a rich set of trial histories to probe what stimulus and reward contingencies guide attention and to track the shifting locus of attention on a trial-by-trial time scale. Using this platform, we identified a robust, spatially focused neural correlate of attention in whisker somatosensory cortex (S1), and tested a major circuit model of attention involving VIP interneurons in sensory cortex.

Results

To study attention, we developed a head-fixed whisker detection task. Mice had nine whiskers inserted in a piezo array, and on each trial, one randomly selected whisker (Go trials) or no whisker

(NoGo trials) was deflected in a brief train. Mice were rewarded for licking during a response
window on Go trials (Hits) but not on NoGo trials (False Alarms) (**Fig. 1a-b**). The task was performed
in darkness and incorporated a delay period of 0, 0.5, or 1 sec in different mice to separate whisker
stimuli from response licks. Go and NoGo trials were randomly intermixed with a variable inter-trial
interval, and whisker identity on Go trials was randomly selected. Thus, mice could not anticipate
the upcoming whisker or precise trial timing. Sequential trials could be one Go and one NoGo, two
NoGo trials, two Go trials on different whiskers, or two Go trials on the same whisker (**Fig. 1c**).
80   Expert mice effectively distinguished Go from NoGo trials, quantified by d' from signal
detection theory (**Fig. S1a**). We analyzed behavior in each session during a continuous task-
engaged phase that excluded early and late low-performance epochs ($d'<0.5$) that reflect
motivational effects[18]. In expert mice, overall mean d' was 0.923 ± 0.051 (n=476 sessions, 22 mice),
but local fluctuations in d' regularly occurred over the time course of several trials, suggesting that
the recent history of stimuli or rewards may dynamically alter sensory detection behavior
(**Extended Data Fig. 1a**). To examine this, we classified each current Go trial based on the history
of whisker stimuli and reward on immediately preceding trials. We used trial history categories of:
i) prior NoGo, ii) prior Hit to the same whisker as the current trial, iii) prior Hit to a different whisker,
iv) prior Miss to the same whisker, and v) prior Miss to a different whisker. Current NoGo trials were
classified into categories of prior NoGo, prior Hit (to any whisker), or prior Miss (to any whisker). We
separately tracked trials preceded by a single Hit trial from those preceded by multiple sequential
prior Hits ("prior > 1 Hit") (**Extended Data Fig. 1b**).
93
Reward history cues focal attention in the whisker system
Prior trial history strongly influenced detection on the current trial. When the prior trial was a NoGo,
mice detected the current Go whisker with d' = 1.13 ± 0.02 (termed d'_{NoGo}, mean ± SEM across 476
sessions, 22 mice), which we consider baseline detection sensitivity. d' for detecting a given Go
whisker was elevated following 1 prior Hit to the same whisker, and even more so following multiple
consecutive Hits to the same whisker ($d'_{>1HitSame}$ = 2.45 ± 0.11, p = 1.0e-04 vs. d'_{NoGo}, permutation
test). In contrast, d' was reduced following one or multiple prior Hits to a different whisker ($d'_{>1HitDiff}$ =
0.82 ± 0.06, p= 1.0e-04 vs. d'_{NoGo}, permutation test) (**Fig. 1d-e**). d' after multiple prior Hits to the
same whisker ($d'_{>1HitSame}$) was substantially greater than after multiple prior Hits to a different
whisker ($d'_{>1HitDiff}$) (**Fig. 1d**). Thus, recent Hits engage a whisker-specific boost in detection, evident
as a whisker-specific increase in d', termed Δ d' (**Fig. 1d**). This effect was found across mice with 0,
0.5, or 1 sec delay period (p = 1.0e-04 for $d'_{>1HitSame}$ vs. $d'_{>1HitDiff}$ in each case), so these data were
combined for behavioral analyses (**Fig. S1d**).
Trial history-dependent boosting of detection required the conjunction of prior stimulus
plus reward, because d' did not increase after a prior Miss to the same whisker ($d'_{MissSame}$ = 0.90 ±
0.09, p = 4.2e-3 relative to d'_{NoGo}). Boosting also failed to occur if the mouse licked to the prior Go
but received no reward (i.e., unrewarded hits) or a very small reward, (<4% of maximal reward
volume ($d'_{>1HitSame}$ vs. $d'_{>1HitDiff}$, p = 0.49, permutation test) (**Fig. 1f**). Thus, boosting was not driven by
prior whisker deflection alone, or stimulus-evoked licking, but by whisker-reward association on
recent trials.
We interpret this effect as whisker-specific attention, because it shares the properties of
attention documented in primates during detection tasks[19-23]. It involves both increased sensitivity
(d') and a shift in criterion (c), with the whisker-specific Δd' reflecting increased Hit rate following >1
prior Hit to the same whisker relative to >1 prior Hit to a different whisker (p = 1.0e-04, , permutation
test; **Fig. 1g**). The whisker-specific increase in d' was observed across mice (**Fig. 1h**; p = 1.0e-04,
permutation test). We also observed shifts in criterion[19,21-22] (Δ c) which with a more modest
whisker-specific component ($c_{>1HitSame}$ vs. $c_{>1HitDiff}$, p = 0.002 , permutation test) (**Fig. 1i**). On average,

the whisker-specific shift in sensitivity was larger than the whisker-specific shift in criterion (**Fig. 1j**; $\Delta d'_{>1HitSame}$ = 1.4 ± 0.20, $\Delta d'_{>1HitDiff}$ =-0.47 ± 0.11, p = 1.0e-04, permutation test; $\Delta c_{>1HitSame}$ = -2.03 ± 0.18, $\Delta c_{>1HitDiff}$ =-1.34 ± 0.08 , p = 7.0e-04, permutation test).

Mice trained on all 3 delay periods showed the whisker-specific boost in d' (**Extended Data Fig. 1c-d**). Reaction times on Go trials were reduced by prior Hit trials, as expected for attention (assessed in mice without a delay period), but this effect was not whisker-specific, which may reflect a ceiling effect (**Extended Data Fig. 1e**). The task interleaved whisker deflections of varying amplitude, and whisker-specific boosting of d' was greatest for current Go trials with low-amplitude (weak) deflections, as expected for attention (**Fig. S1f**). Trial history effects were consistently observed across different variations of the task (used in different mice) in which we manipulated the stimulus probability of each whisker in blocks, or manipulated the probability of sequential same-whisker Go trials to make prior same Hit histories more likely (**Fig. S1g-h**). Trial history effects were driven by stimulus-reward association, not by stimulus salience or reward probability, because all whisker deflections were, on average, physically identical and reward probability was always 100% for each whisker.

136 Behavioral shifts in d' exhibited the hallmark effects of attentional selection: spatial specificity, temporal specificity, and flexible targeting . Spatially, d' was boosted most strongly by prior Hits to the same whisker, more weakly by prior Hits to an immediate same-row or same-arc neighbor, and not at all by prior Hits to a diagonally adjacent or more distant whisker (**Fig. 1k**). Thus, boosting is somatotopically organized. Temporally, attentional boosting fell off with the time interval between Go trials (which varied due to variable ITI and intervening NoGo trials), and subsided after ~10s (**Fig. 1l**). Enhancement of d' was flexibly shifted to different whiskers in an interleaved manner, and had similar magnitude when cued by trial history to any of the 9 whiskers in rows B-D or arcs 1-3 (**Fig. 1m**). Thus, mice use recent history of stimulus-reward association to dynamically boost sensory detection on spatially specific whiskers on a rapid trial-by-trial timescale, consistent with attentional selection[1,17,19-20, 24-25]. These properties are identical to attentional capture guided by reward history in humans and non-human primates[4-16].

Whisker-specific attention is not mediated by whisker or body movement

150 To test whether trial history effects involve whisker or body movement, we extracted these movements, plus pupillary dilations related to arousal[26-27] , from behavioral videos of 9 mice (74 sessions) using DeepLabCut[28] (**Extended Data Fig. 2a**). Reward retrieval at the end of Hit trials was associated with whisker movement, body movement (detected from platform motion), and pupil dilation that slowly subsided during the ITI before the next trial (**Fig. 2a**). The magnitude of movement and pupil dilation during the ITI was greater after 1 or >1 prior Hits, relative to prior NoGo, but was identical for prior same and prior different conditions (**Fig. 2a, Extended Data Fig. 2b**). Thus, mice exhibited increased general arousal following prior Hits. During the subsequent trial, whisker stimulation evoked modest whisker and body motion during the stimulus period, and these were also heightened after prior Hits (p = 1e-04, permutation test), indicating that behavioral arousal from prior Hits persisted into subsequent trials, but did not differ between prior same whisker and prior different conditions (p=0.39, permutation test **Fig. 2a-b**).

162 To test whether active whisker movement contributed to the whisker-specific $\Delta d'$ effect, we paralyzed whisker movements with Botulinum toxin B (Botox) injection in the vibrissal pad in 4 task-expert mice. Paralysis was maintained over 7 days, and history effects were compared between standard sessions prior to Botox and the Botox sessions. The average whisker-specific shift in behavioral d' and c following >1 prior hit did not differ between Botox and non-Botox sessions (**Fig. 2c-d**). Thus, prior hits engage general arousal as well as whisker-specific attention, and whisker-specific attentional effects do not require active whisker movement.

Neural correlates of reward history-guided attention in L2/3 PYR cells in S1

The somatotopic precision of attention selection (**Fig. 1k**) suggests a neural basis in a somatotopically organized brain area like S1. We performed 2-photon imaging in S1, using Drd3-Cre;Ai162D mice that transgenically express GCaMP6s in L2/3 PYR cells. Mice performed the task with a delay period that separated whisker-evoked responses (analyzed 0-799 ms after stimulus onset) from later licks and rewards, and any trials with early licks were excluded. Imaging fields were localized in the S1 whisker map by post-hoc cytochrome oxidase staining for whisker barrel boundaries in L4 **(Fig. 3a-b)**. 61% of L2/3 PYR neurons were whisker responsive, and trial history modulated whisker responses for many individual neurons. For example, in **Fig. 3c**, neurons increased their response to the C3 whisker after >1 prior hit to that same whisker, but not after >1 prior hit to a different whisker.

On average, L2/3 PYR cells responded to a given Go whisker when the prior trial was a NoGo, responded more strongly following 1 prior Hit to the same whisker (p=1e-04), and even more after > 1 prior Hit to the same whisker (p=1e-04). This boosting of sensory responses did not occur when the prior trial was a Miss to same whisker, or >=1 Hit to a different whisker (n = 6 mice, 70 sessions, 6906 cells, **Fig. 3d-e**). Thus, recent stimulus-reward association modulated whisker-evoked ΔF/F in L2/3 PYR cells in a way that closely resembled the behavioral attention effect (**Fig. 3e** vs. **Fig. 1d**). On current NoGo trials, no whisker stimulus was presented and ΔF/F traces were largely flat, except for NoGo trials following prior Hit trials, which exhibited a surprising rising ΔF/F signal. We interpret this as an expectation or arousal effect, which parallels the increased FA rate on these trials (**Fig. 1g**). These effects were evident in single example fields (**Extended Data Fig. 3a**). History-dependent boosting of whisker responses was only evident in whisker-responsive cells (defined from prior NoGo trials), and not in non-responsive cells, which only showed the arousal effect (**Extended Data Fig. 3b-c**). Boosting after prior Hits was whisker specific in all 6/6 mice, but its magnitude varied across mice (**Fig. 3f**) and correlated with the magnitude of the behavioral attention effect measured by $\Delta d'$ in each mouse (**Fig. 3g**).

We quantified the attention effect in individual cells using three attention modulation indices (AMI). $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ quantify the change in whisker-evoked response observed after >1 prior Hit to the same (or different) whisker vs. after a prior NoGo. Most cells showed positive $AMI_{>1HitSame-NoGo}$ values and negative $AMI_{>1HitDiff-NoGo}$ values, indicating up- and down-modulation of response magnitude by identity of the prior whisker Hit. $AMI_{>1HitSame->1HitDiff}$ compares response magnitude after >1 prior hit to the same whisker vs. >1 prior hit to a different whisker. This was shifted to positive values for most L2/3 PYR cells, indicating whisker-specific attentional modulation (**Fig. 3h, i**). This was reproducible across individual mice (**Extended Data Fig. 3d**). Modulation of whisker-evoked responses for all cells sorted by $AMI_{>1HitSame->1HitDiff}$ is shown in **Extended Data Fig. 3e**.

History-dependent boosting in S1. After >1 prior hits to a given whisker, responses to an immediate same-row adjacent neighboring whisker were boosted strongly, those to an immediate same-arc neighbor were boosted less, and those to diagonal adjacent neighbors or further whiskers were boosted the least or not at all. This somatotopic profile of ΔF/F boosting strongly resembled the somatotopy of behavioral d' boosting (**Fig. 3j**). Spatially within S1, >1 prior hits to a reference whisker boosted whisker-evoked ΔF/F to that whisker most strongly for L2/3 PYR cells within the reference whisker column and in the near half of the neighboring column, and weakly or not at all beyond that. This somatotopically constrained boosting was not observed for >1 prior hits to a different whisker, or following a miss to the reference whisker (**Fig. 3k**). Analysis of individual cell AMI confirmed somatotopically precise boosting (**Extended Data Fig. 4a**). This defines the spatial profile of the trial history-based 'attentional spotlight' in S1 as boosting responses to the cued

217   whisker within a region of 1.5 columns width in L2/3. At the center of the spotlight, the
218   representation of the history-cued whisker is boosted within its own column (**Extended Data Fig.**
219   **4b-c**).
220

Attention involves shifts in receptive fields toward the attended whisker
In primates, attention not only increases sensory response magnitude and signal-to-noise ratio in
sensory cortex, but can also shift neural tuning toward attended stimuli[29-31]. We tested whether
reward history-cued attention involves receptive field shifts by L2/3 PYR cells in S1. We analyzed all
whisker-responsive cells relative to the boundaries of the nearest column (defined in the Prior
NoGo condition). We calculated the mean receptive field across 9 whiskers centered on the
columnar whisker (CW) when the prior trial was a NoGo (**Fig. 4a**, black traces in center panel). We
then recalculated the receptive field, for these same cells, for >1 Prior Hit to each of the surround
whiskers (purple traces in outside panels). For many whiskers, recent stimulus-reward association
shifted the mean receptive field toward the history-cued whisker or nearby whiskers (**Fig. 4a**). To
quantify this effect, we calculated the tuning center of mass (CoM) across the 3 by 3 whisker array
for these mean receptive fields. History-based cueing to an attentional target whisker generally
shifted tuning CoM towards that whisker, except when attention was cued upward, which caused
little upward CoM shift, which could reflect a spatial asymmetry in attentional effects[32] on whisker
touch (**Fig. 4b**). Attentional cueing to the CW elevated responses to that whisker within its S1
column but caused very little tuning change (**Extended Data Fig. 5a**).
To measure receptive field shifts in individual neurons, we quantified the shift in CoM for
each responsive cell along an attentional axis from the CW to the attentional target whisker, which
was one of the 8 surrounding whiskers (**Extended Data Fig. 5b-c**). In the >1 prior Hit condition, the
mean CoM shift along the attentional axis was 0.12 ± 0.04 (n=173 cells with sufficient whisker
sampling, p=3.6e-3, one-sample permutation test vs. mean of 0), indicating a tuning shift toward
the attended whisker. A range of tuning shifts were observed, with 62% of cells showing CoM shifts
towards the attended whisker, and 38% showing shifts in the opposite direction (p=2.3e-3, binomial
exact test for difference from 0.5). Thus, attentional cueing involves receptive field shifts as well as
modulation of whisker response magnitude.
246

Attention boosts population decoding of recently rewarded whiskers
Is the magnitude of attentional boosting of L2/3 PYR responses sufficient to improve neural coding
of attended whisker stimuli on single trials? To test this, we built a simple neural decoder that uses
logistic regression to predict the presence of any whisker stimulus from single-trial population
mean ΔF/F calculated across all whisker-responsive cells in a single imaging field. Each field
spanned ~1-1.5 columns within the 9-whisker region of S1, typically centered over the location of
column corresponding to the center whisker on the piezo array. A separate decoder was trained on
each session (n=70 sessions, 6 mice), and performance was assessed from held-out trials (**Fig. 5a**).
Because S1 is somatotopically organized, we observed modest, above-chance performance for
detecting any of the 9 whiskers from mean activity in a single field (relative to NoGo trials or
shuffled data), strong performance for detecting the field best whisker (fBW) that is topographically
matched to the field location, and no ability to detect non-topographically aligned whiskers (non-
fBWs) (**Fig. 5b**).
Population mean ΔF/F and decoder performance were related to current trial outcome, with
greater ΔF/F and stimulus prediction on current Hit trials than Miss trials, and on current false
alarm trials than correct rejection trials. This is consistent with the known representation of
sensory decision in S1[33-34] (**Fig. 5c-d**). Next, we examined decoding as a function of prior trial

history. Detection of any whisker stimulus from Go trials was improved following >1 Hit to the same
whisker, relative to prior NoGo (p=1e-0.4, test) or to >1 prior Hit to a different whisker (p=0.001, test)
(**Fig. 5e**). This boost in decoding performance did not occur when the decoder was tested only on
fBW trials but did occur for non-fBW trials (**Fig. 5f-g**). This same effect was found when decoding
was evaluated on current Hit trials only, so this is an effect of trial history, not current sensory
decision (**Fig. 5e-f**).
Thus, history-cued attention to a whisker strengthens its encoding on single trials in S1. The
lack of improvement in decoding attended fBWs likely reflects the strong coding of these whiskers
under baseline conditions, so that attentional boosting of CW responses (**Extended Data Fig. 4**)
does not further improve detection.
274
Attentional modulation of neural coding with Neuropixels spike recordings
To examine attentional modulation of S1 neural coding at finer temporal resolution and across
layers, we recorded extracellular spiking in S1 using Neuropixels probes[35] that spanned L1-6 (**Fig.**
6a). Mice performed a modified version of the task in which Go stimuli were distributed over only 4
or 5 whiskers, rather than 9, to enable adequate sampling of each history condition per session.
Behaviorally, mice performing the task with 4-5 whiskers showed the whisker-specific Δ*d'* attention
effect, but at lower magnitude due to the smaller number of whiskers (**Extended Data Fig. 6a**). All
sessions included the CW for the recording site plus 3 nearby whiskers. We spike sorted to identify
single units, classified units as regular-spiking (RS) or fast-spiking (FS), and assigned laminar
identity based on CSD analysis of local field potentials (**Extended Data Fig. 6b-f**). Many single
units showed history-dependent modulation of whisker-evoked spiking (**Fig. 6b**).
S1 units responded to each deflection in the stimulus train. On average for L2/3 RS units,
whisker-evoked spiking was boosted in Go trials after >1 prior Hit to the same whisker, relative to >1
prior Hit to a different whisker (**Fig. 6c-d**). This effect did not occur after prior Miss trials, and firing
on NoGo trials was not significantly regulated (**Fig. 6d**). L4 RS units showed only a slight trend for
history-dependent modulation that did not reach significance (**Fig. 6e-f**). L5A-B RS units (grouped
together) showed a whisker-specific attentional boost similar to L2/3 (**Fig. 6g-h**).
To examine heterogeneity across units we calculated AMI for each unit. Most L2/3 RS units
responded more strongly after prior Hits to the same whisker relative to prior NoGo (as measured
by $AMI_{>1HitSame-NoGo}$) and more weakly after prior Hits to a different whisker relative to prior NoGo (as
measured by $AMI_{>1HitDiff-NoGo}$). This whisker-specific attentional effect, evident as the separation
between $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ distributions, was not present in L4, and was weaker in
L5A-B (**Fig. 6i**). These laminar trends were also apparent in $AMI_{>1HitSame->1HitDiff}$, which was shifted
positively in L2/3 relative to L5A-B and L4 units (**Fig. 6j**). Calculating the mean AMI across neurons
confirmed whisker-specific attentional shifts in L2/3 and L5, but not in L4 (**Fig. 6k;** ($AMI_{>1HitSame-NoGo}$
vs. $AMI_{>1HitDiff-NoGo}$, L2/3: p = 0.03, L4 p=0.65, L5a/b: p = 0.01; $AMI_{>1HitSame->1HitDiff}$, L2/3: p = 0.031, L4
p=0.51, L5a/b: p = 0.06, permutation test). This suggests that history-dependent attentional
modulation is not simply inherited from the thalamus, but has a cortical component.
303
VIP interneurons do not carry a simple "attend here" signal
We used reward history-based attention in S1 to investigate the candidate involvement of VIP
interneurons in attentional selection. VIP cells are known to disinhibit PYR cells to increase PYR
sensory gain during arousal, locomotion, and whisking[36-40]. For attention, this same VIP circuit has
been hypothesized to be activated by long-range (e.g., top-down) inputs, and act to amplify local
PYR responses to selected sensory features[39-40] (**Fig. 7a**). Whether VIP cells mediate goal-directed
attention is still unclear[41-42], and their involvement in history-based attention has not been tested.

We used 2-photon imaging from L2/3 VIP cells in VIP-Cre;Ai162 mice to ask whether VIP cells are
activated when mice direct attention to a particular whisker column in S1 (**Fig. 7b**).
VIP cells in S1 are activated by arousal (indexed by pupil size), whisker and body movement,
and goal-directed licking during this whisker detection task[43] . These behaviors all peak at the end
of Hit trials, as mice retrieve rewards, and then systematically decline during the ITI, which ends
with a 3-sec lick-free period that is required to initiate the next trial. As a result, VIP cell ΔF/F falls
systematically during the ITI after Hit trials, correlated with these behavioral variables, and falls less
after NoGo or Miss trials (**Fig. 7c**). VIP cells in S1 also show robust whisker stimulus-evoked ΔF/F
transients, which ride on this declining baseline (Ramamurthy et al., 2023). To test whether
whisker-evoked VIP responses are greater when reward history cues attentional selection, we
calculated mean Δ F/F traces (n=7 mice, 103 sessions, 1843 VIP cells) as a function of trial history.
Baseline (prestimulus) ΔF/F declined more steeply on trials following prior Hits than following prior
NoGo or Miss, as expected. Superimposed on this, and clearest after detrending the baseline,
whisker-evoked Δ F/F was also increased after prior Hit trials. However, this was not whisker-
specific (**Fig. 7d-f**). AMI analysis confirmed increased responsiveness for most VIP cells in both
prior >1 Hit same and prior >1 Hit different conditions, but no whisker-specific attention effect (the
$AMI_{>1HitSame->1HitDiff}$ index is peaked at 0) (**Fig. 7g**, for all cells individually see **Extended Data Fig. 7**).
Together, these findings indicate that VIP cells as a population do not carry a whisker-
specific attention signal, but do exhibit a general increase in activity with any prior Hit that
resembles a general arousal effect. A subpopulation of VIP cells may carry a whisker-specific
"attend here" signal, but not the general VIP population as a whole.
332

Discussion

Attention captured by recent history cues[4-16], including stimulus-, reward-, and choice-history, is
distinct from classical top-down and bottom-up forms of attention. The neural mechanisms and
control circuits for attention remain poorly understood. History-guided attentional capture
provides a useful model for identifying such mechanisms, because it is spatially focused, flexibly
allocated, and temporally constrained (characteristic features of selective attention), but is
automatic and thus does not have to be explicitly trained. Our demonstration of reward history-
guided attention for whisker touch in mice establishes a novel model for studying the
neurobiological mechanisms of attentional capture. This model complements recent visual tasks
to study top-down[44-49] and bottom-up[49-50] attention in head-fixed mice, and thus may reveal the
extent to which common neurobiological mechanisms are engaged in different forms of attention
and different sensory modalities.
345
The recent history effects on sensory detection in our study match well with attentional
capture by reward history in humans[4-16]. Enhanced detection (quantified by d-prime) was driven by
recent whisker-reward association and was whisker-specific, and thus was distinct from global
arousal or motion effects[51]. It was not short-term or early priming, which occurs in response to
stimulus presentation only (not reward), and occurs even when the priming stimulus is
undetectable and typically operates on a very short (<100 ms) time scale[52]. It was distinct from
oddball paradigms that generate stimulus-specific adaptation and enhanced behavioral
responses to novel stimuli[53]. Unlike bottom-up attention, it was not driven by physical
stimulus salience. Unlike top-down attention, it was automatic and not goal-directed (since
allocating attention to a prior Hit whisker did not increase the overall probability of obtaining
reward in the task). Rodents are well known to exhibit history-dependent choice biases in
decision-making tasks[54-57], including serial dependence[58-59], contraction bias[54,60], adaptation

aftereffects[61], win-stay/lose-switch strategies[62-64] and choice alternation[62,65]. Our findings show that mice use prior reward history not only to bias their subsequent choices (as in these choice biases), but also to prioritize sensory processing.

The neural correlates of attention have been primarily studied in non-human primates, and include increased sensory-evoked spike rate[17,66-67], reduced variability[17], neural synchrony modulation[68], and changes in receptive fields[17,29-31] including boosting of the peak, changes in receptive size and receptive field shifts. In primates, these effects are greatest at higher levels of the sensory hierarchy but also occur in primary sensory cortex[17,70]. The precise spatial organization of these coding effects in sensory cortex has been unknown, but has important implications for understanding neural control circuits for attention. We took advantage of S1 whisker map topography[71] to define the precise topography of the attentional spotlight for reward history-guided attention. We found that attentional capture modulated sensory responses in a S1 region spanning ~1.5 cortical columns, consisting of increases in whisker-evoked spike rate and RF peak in the central attended column, and shifting of RFs toward the attended whisker for cells in surrounding columns. The net effect was an increase in total population activity evoked by the attended whisker, due to both increasing the number of PYR cells responding to that whisker, and elevating the number of spikes per cell. This somatotopically restricted boosting[72-75] is distinct from global behavioral state-related modulation in sensory cortex (e.g., related to arousal indexed by pupil size, or active whisker movement, or locomotion for V1 responses), in which strong, enhancement of sensory activity occurs that is area-wide or spans multiple cortical areas[51].

History signals in mouse cortex have not previously been described in the context of attention but have been studied in the context of choice biases during decision making[76-78] and in reversal learning[79-82]. Higher order cortical areas including posterior parietal cortex (PPC)[54] and orbitofrontal cortex (OFC)[81] of mice these history signals were found to have a causal role in the representation and use of these forms of history information. Additionally, S1 cortex receives instructional signals from OFC, and was shown to be necessary for reversal learning, a history-dependent behavior[81].

The circuit origins of attentional modulation are unknown. We found that reward history-guided attention modulated PYR sensory responses in L2/3 and L5 but not L4, suggesting either an intracortical origin, or a thalamic origin in secondary thalamic nuclei like the POm that projects to L2/3 and L5a. Thalamic control of attention has been implicated in visual and tactile cross-modal attention tasks in mice[83], as well as some NHP studies[84]. We tested one major circuit model[39-40] for attentional boosting in sensory cortex, that long-range inputs amplify pyramidal (PYR) cell activity by activation of local VIP interneurons in sensory cortex[84]. L2/3 VIP interneurons are activated by both neuromodulatory and glutamatergic inputs, and inhibit other cortical inhibitory interneurons to disinhibit PYR cells, thus boosting PYR sensory responses[38,85]. VIP interneurons are activated by global behavioral state such as locomotion and whisking, by spontaneous arousal during quiet wakefulness[26,36,37,], and by top-down contextual signals[39-40]. VIP cells in sensory cortex receive top-down input from anterior cingulate cortex (ACC), which has been suggested to mediate top-down attentional effects on sensory processing[38]. However, recent studies have suggested that L2/3 VIP modulation of PYR activity is orthogonal to attentional effects measured in a cross-modal attention task[41], and that cingulate cortex is not engaged when attention is endogenously deployed[86]. We tested the potential involvement of VIP cells in reward history-cued attention by testing whether attention activates VIP cells in attended columns to boost PYR cell responsiveness. We found that

405 L2/3 VIP cells, at least as a full population, do not carry this attend-here signal (however, we cannot
406 rule out that a subpopulation of VIP cells does carry this signal).
407
408   Overall, this study establishes reward history-guided attention in the mouse whisker system
409 as a promising platform to examine other circuit mechanisms for attention.
410

ONLINE METHODS

412
Animals
All methods followed NIH guidelines and were approved by the UC Berkeley Animal Care
and Use Committee. The study used 22 mice. These included 7 Drd3-Cre;Ai162D mice and 10 VIP-
Cre;Ai162D mice (used for behavior and 2-photon imaging), and 5 offspring from Drd3-Cre x Ai162D
crosses (genotype not determined) used in extracellular recording experiments. VIP-Cre (JAX #
10908) and Ai162D mice (JAX # 031562) were from The Jackson Laboratory. Drd3-Cre mice were
from Gensat MMRRC (strain number 034610).
Mice were kept in a reverse 12:12 light cycle, and were housed with littermates before
surgery and individually after cranial window surgery. Mice were roughly evenly divided between
male and female, and no sex differences were found for the results reported here. Behavioral,
imaging, and analysis methods were as described in Ramamurthy et al., 2023[43], and are here
described more briefly.
425
Surgery for behavioral training and 2-photon imaging
Mice (2-3 months of age) were anesthetized with isoflurane (1-3%) and maintained at 37°C.
Dexamethasone (2 mg/kg) was given to minimize inflammation, meloxicam (5-10 mg/kg) for
analgesia, and enrofloxacin (10 mg/kg) to prevent infection. Using sterile technique, a lightweight
(<3 g) metal head plate containing a 6 mm aperture was affixed to the skull using cyanoacrylate
glue and Metabond (C&B Metabond, Parkell). The headplate allowed both head fixation and 2-
photon imaging through the aperture. Intrinsic signal optical imaging (ISOI) was used to localize
either C-row (C1, C2, C3) or D-row (D1, D2, D3) barrel columns in S1[87], and a 3 mm craniotomy was
made within the aperture using a biopsy punch over either the C2 or D2 column. The craniotomy
was covered with a 3 mm diameter glass coverslip (#1 thickness, CS-3R, Warner Instruments) over
the dura, and sealed with Metabond to form a chronic cranial window. Mice were monitored on a
heating pad until sternal recumbency was restored, given subcutaneous buprenorphine (0.05
mg/kg) to relieve post-operative pain and then returned to their cages. After the mice recovered for
a week, behavioral training began.
440
Behavioral task
To motivate behavioral training, each mouse received 0.8-1.5 mL of water daily,
calibrated to maintain 85% of pre-training body weight. Mice were weighed and observed daily.
Behavioral training sessions took place 5-7 days per week. For behavioral training, the mouse
was head-fixed and rested on a spring-mounted stage[43,88]. Nine whiskers were inserted in a 3 x 3
piezo array, typically centered on a D-row or C-row whisker. Piezo tips were located ~5 mm from
the face, and each whisker was held in place by a small amount of rubber cement. A tenth piezo
was present near the 3x3 array but did not hold any whisker ("dummy piezo"). A capacitive lick
sensor (for imaging experiments) or an infrared (IR) lick sensor (for extracellular recording
experiments) detected licks, and water reward (mean 4 μl) was delivered via a solenoid valve.
Mice were transiently anesthetized with isoflurane (0.5-2.0%) at the start of each session to

enable head-fixation and whisker insertion, after which isoflurane was discontinued and
behavioral testing began after the effects of anesthesia had fully recovered. Behavior was
performed in the dark with 850 nm IR illumination for video monitoring. Masking noise was
presented from nearby speakers to mask piezo actuator sounds. Task control, user input and
task monitoring were performed using custom Igor Pro (WaveMetrics) routines and an Arduino
Mega 2560 microcontroller board.
458
Training stages
A series of training stages (1-5 days each) were used to shape behavior on the Go/NoGo
detection task. In Stage 1, mice were habituated to the experimental rig and to handling. In Stage 2
mice were head-fixed and conditioned to lick for water reward at the port. In Stage 3, mice received
a reward (cued by a blue light) for suppressing licks for at least ~3 seconds, termed the Interlick
Interval (ILI) threshold. Stage 4 introduced whisker stimulation for the first time, with 50% Go trials
(whisker stimulation) and 50% NoGo trials (no whisker stimulus), with automatic reward delivery in
the response window on Go trials (i.e., classical conditioning). The dummy piezo was actuated on
all NoGo trials, so that any unmasked piezo sounds did not provide cues for task performance. The
Go whisker randomly chosen from among 9 possible whiskers. This stage ended when mice shifted
licks in time to occur before reward delivery. In Stage 5, training switched to operant conditioning
mode, and mice were required to lick in the response window (0 - 300 ms after whisker stimulus
onset) to receive a reward. There was no delay period in this stage. Learning progress was tracked
by divergence of Go/NoGo lick probability. In Stage 6, the delay period was introduced. To do this,
we introduced a trial abort window in which licking during the stimulus presentation period caused
a trial to be canceled without reward, to discourage licking during the stimulus presentation.
Simultaneously, we implemented a ramp-plateau reward gradient within the response window, so
that later licks resulted in a larger reward. Over the course of this stage, the trial abort window was
gradually lengthened, and the time of reward plateau was gradually increased. Learning progress
was tracked by the gradual increase in median trial first lick time. Stage 7 represented the final
whisker detection task, which included completely randomized Go/NoGo trials, fixed trial abort
window and reward plateau parameters, and eliminating the blue light that signaled reward
delivery. Mice were deemed task experts when they exhibited stable performance at $d' > 1$ (mean
running d-prime) for three consecutive sessions. Expert mice performed 500 - 1000 trials daily.
483
Task structure
Task structure was identical to Ramamurthy et al., 2023[43] . Briefly, on each trial, a whisker
stimulus was applied to either one randomly selected whisker (Go trials, 50-60% of trials) or no
whisker (NoGo trials, 40-50% of trials). The whisker stimulus consisted of a train of five deflections
separated by 100 ms each. Every deflection was a 300 µm amplitude (6^0 angular deflection)
rostrocaudal ramp-and-return movement with 5 ms rise/fall time and 10 ms duration. NoGo trials
presented the same stimulus on a dummy piezo that did not contact a whisker, so that any
unmasked auditory cue from piezo movement was matched between Go and NoGo trials. Trial
onset was irregular with an ITI of 3 ± 2 s. Mice had to restrict licking to greater than 3 sec interlick
interval (ILI) to initiate the next trial. On Go trials, mice were rewarded for licking within the 2.0 s
response window with ILI of <300 ms. Licking was not rewarded on NoGo trials. Each trial outcome
was recorded as a Hit, Miss, False Alarm, or Correct Rejection.
Different delay periods were used in different mice (**Extended Data Fig. 3c-d**). Mice used in
imaging or extracellular recording experiments had a delay period of 500 ms (for all spike recording
mice and 7/14 2p imaging mice) or 1000 ms (for the other seven 2p imaging mice). This was used to
separate sensory-driven neural activity from action- and reward-related activity. Three mice used

only for behavioral data collection were tested without a delay period, which enabled testing of
attention effects on lick response latency (**Extended Data Fig. 3e**).
Because reward size varied with lick time in the response window, reward volume varied
across trials. In addition, for mice with the 1000 ms delay period, mice sometimes licked on Go
trials after stimulus presentation but before the response window opened, and thus earned no
reward. These represent unrewarded Hits, so that attention effects could be quantified based on
absence of reward and reward size on prior Hit (**Fig. 1f**).
507
Task variations
All mice in this study were trained on an "equal probability" (EqP) version of the task in
which whisker identity on each Go trial was randomly selected from nine possible whiskers with
equal 1/9 probability (EqP). Two other task variations manipulated either the global or local
probability of each specific whisker, while still randomly selecting whisker identity on each trial. In
HiP sessions, we manipulated the global stimulus probability of each whisker by presenting one
whisker with higher probability (80% of Go trials) than the others. This was done in 300-400 trial
blocks, interleaved with standard EqP blocks (13 mice) on the same day. In HiPsame sessions (run
on separate days from EqP sessions) we manipulated the local probability of repeating the same
whisker stimulus on consecutive trials, while maintaining the overall probability of each whisker at
1/9. This was done in blocks interleaved with EqP blocks (7 mice). Whisker-specific attentional
cueing was observed in all 3 task variants (**Fig. 1h**), so data from all variations were combined for
the rest of analysis. All 22 mice were trained and tested on the EqP version and either the HiP or
HiPSame version of the task, but EqP blocks/sessions contributed trials to history analysis only in a
subset of animals, since multiple hits to a given whisker in a single session were adequately
sampled only in sessions with a higher number of total trials.
A modified task version was used for extracellular recording experiments, in order to
adequately sample trial history conditions when only 2-4 days of acute recording were possible per
mouse. To do this, we reduced the number of whiskers sampled during Go trials from 9 whiskers to
either 4 or 5 whiskers 5 (4-5-whisker task; **Fig. 6a**). The 4-5-whisker task was used in 3 mice for
extracellular recordings, and was also applied in 2 mice that were used in PYR cell imaging, where
performance could be compared to the standard 9-whisker task (**Extended Data Fig 6a**).
530
Behavioral movies and DeepLabCut tracking
Behavioral movies were acquired at 15-30 frames/sec using either a Logitech HD Pro
Webcam C920 (modified for IR detection) or FLIR Blackfly S (BFS-U3-63S4M; used in video
analyses). DeepLabCut[28] was used to track spontaneous face and body movements. Movies were
manually labeled to generate training datasets for tracking facial motion (snout tip, whisker pad and
2-3 whiskers), body motion (corner of the mouse stage, whose motion reflects limb and postural
movements), pupil size (8 labels on the circumference of the pupil), eyelids (8 labels on the
circumference of the eyelid), and licking (tongue and lickport). Three separate networks were
trained (100,000-200,000 iterations) such that a good fit to training data was achieved (loss <
0.005). One network each was trained for face/body motion for the two camera setups (version 1
network: 1110 labeled frames from 37 video clips across 6 mice) and another for pupillometry
(version 2 network: 2463 labeled frames from 27 video clips across 2 mice).
Behavioral movies from 74 sessions in 9 mice were analyzed for whisker motion (average
across all whisker-related labels), body motion (stage corner) and pupil size (ellipse fit to the pupil
markers). Blinking artifacts were removed using a one-dimensional moving median filter (40 frame
window) applied to the trace of pupil size as a function of time. Pupil size measured on each frame
was normalized to the mean pupil size over each individual session.

548
Whisker paralysis by Botox injection
Four mice (2 VIP-Cre;Ai162D mice and 2 Drd3-Cre;Ai162D mice) that were used for imaging
experiments also underwent Botox injection to induce paralysis of whisking[43,88-90]. Both whisker
pads were injected with Botox (Botulinum Neurotoxin Type A from Clostridium botulinum, List Labs
#130B). A stock solution of 40 ng/µl Botox was prepared with 1mg/ml bovine serum albumin in
distilled water. Each whisker pad was injected with 1 µl of a 10 pg/µl final dilution using a microliter
syringe (Hamilton). Whisking stopped within 1 day, and gradually recovered in ~1 week. Following
the initial dose, a 50% Botox supplement was injected once per week, as needed. Imaging was
performed > 24 hours after any Botox injection.

Behavioral Analysis
560   Behavioral performance was assessed using the signal detection theory measures[91] of
561   detection sensitivity (*d'*) and criterion (*c*), calculated from Hit rates (HR) and False Alarm rates
562   (FA), as per their standard definitions:

$$d' = Z_{HR} - Z_{FA}$$

$$c = \frac{1}{2}(Z_{HR} + Z_{FA})$$

where Z is the inverse cumulative of the normal distribution.
To assess the overall behavioral performance of mice on the whisker detection task, *d'*
was computed across trials over the entire behavioral session. For each session, a sliding *d'*
cutoff (calculated over a 50 trial sliding window) was applied to the start and end of the session,
and analysis was restricted between the first and last trial that met the threshold, to minimize
satiety effects. A standard sliding *d'* cutoff of 0.5 was used for all behavioral and imaging
analyses. We tested *d'* cutoffs 0.5, 0.7 and 1 to ensure that choice of *d'* cutoff did not affect key
results. A sliding d' cutoff of 1.2 was used for extracellular recording analyses.

Definition of trial histories
575   For each *current trial*, trial history conditions were defined based on outcome and stimulus on
576   *prior trials*, as follows. These definitions are illustrated in **Extended Data Fig. 1b**.
577   History conditions were defined for *current Go trials* as follows:
578   1. Prior Miss
579   a. On the *same* whisker: The current trial is a Go and the outcome on the previous
580   trial was a Miss to the same whisker.
581   b. On a *different* whisker: The current trial is a Go and the outcome on the previous
582   trial was a Miss to a different whisker.
583   2. Prior NoGo: The previous trial was a NoGo (any outcome).
584   3. Prior 1 Hit
585   a. On the *same* whisker: The current trial is a Go and the outcome on the previous
586   trial was a Hit to the same whisker.
587   b. On a *different* whisker: The current trial is a Go and the outcome on the previous
588   trial was a Hit to a different whisker.
589   4. Prior >1 Hit:
590   a. On the *same* whisker: The current trial is a Go and the outcomes on the previous
591   two or more trials were Hits to the same whisker as the whisker presented on the
592   current trial.
593   b. On a *different* whisker: The current trial is a Go and the outcomes on the previous
594   two or more trials were Hits to a single consistent whisker that was a different

595   identity than the whisker presented on the current trial (i.e., two or more Hits in a
596   row to the same whisker that differed from the current Go trial).
597
598   For each history condition defined above for *current Go trials*, a matched condition was
599   defined for *current NoGo trials*. This allowed us to compute behavioral *d'* and *c* values within
600   each history condition, and to compare neural signals on Go and NoGo trials within each history
601   condition. History conditions for *current NoGo trials* were defined as follows:
602   1. Prior Miss: The current trial is a NoGo and the outcome on the previous trial was a miss to
603   any whisker. NoGo trials in this category were used for comparison with Go trials in both
604   categories 1a and 1b above.
605   2. Prior NoGo: The previous trial was a NoGo (any outcome).
606   3. Prior 1 Hit: The current trial is a NoGo and the outcome on the previous trial was a hit to
607   any whisker. NoGo trials in this category were used for comparison with Go trials in both
608   categories 3a and 3b above.
609   4. Prior >1 Hit: The current trial is a NoGo and the outcomes on the previous two or more
610   trials were Hits to any single repeated whisker. NoGo trials in this category were used for
611   comparison with Go trials in categories 4a and 4b, above.
612
613   The history conditions above were defined based on sequences of consecutive trials,
614   including both Go and NoGo. The history-dependent effects on detection behavior were
615   maintained when history conditions were defined by ignoring NoGo trials, and categorizing
616   history based solely on Go trials (data not shown). NoGo trial were ignored when characterizing
617   the temporal profile of attention as a function of time since last Go trial (**Fig. 1l**).
618
2-photon calcium imaging
A Sutter Moveable-Objective Microscope with resonant-galvo scanning (RESSCAN-MOM,
Sutter) was used to perform 2p imaging in expert mice. A Ti-Sapphire femtosecond pulsed laser
(Coherent Chameleon Ultra II) tuned to 920 nm, or an ALCOR 920 nm fixed wavelength
femtosecond fiber laser (Spark Lasers), was used for GCaMP6s excitation. A water-dipping
objective (16x, 0.8 NA, Nikon) was used, and emission was band-pass filtered (HQ 575/50 filter,
Chroma) and detected by GaAsP photomultiplier tubes (H10770PA-40, Hamamatsu). Single Z-
plane images (512 x 512 pixels) were acquired serially at 7.5 Hz (30 Hz averaged every 4 frames)
using ScanImage 5 software (Vidrio Technologies). Laser power measured at the objective was 60 -
90 mW. On average, 14 imaging fields (305 µm x 305 µm) were obtained per mouse at depths of 110
- 250 µm below the cortical surface[92]. If there was >25% XY overlap, imaging fields were required to
be at least 20 µm apart in depth to avoid repeated imaging of the same cells. After completion of all
imaging experiments, the mouse was euthanized and the brain was collected to perform histology.
632
Histological localization of imaging fields
The brain was extracted and fixed overnight in 4% paraformaldehyde. After flattening, the
cortex was sunk in 30% sucrose and sectioned at 50-60 µm parallel to the surface. Cytochrome
oxidase (CO) staining showed surface vasculature in the most superficial tangential section as well
as boundaries of barrels in L4. Histological sections were manually aligned using Fiji[93], and imaging
fields were localized in the whisker map aided by the surface blood vessels imaged at the beginning
of each session. The centroid of each of the nine anatomical barrels corresponding to whiskers
stimulated in each session and the XY coordinates of all imaged cells were localized relative to
barrel boundaries. A cell was located within a specific barrel column if >50% of its pixels were

within its boundaries, and cells outside barrel boundaries were classified as septal cells. Major and
minor axes of all barrels were averaged to calculate the mean barrel width.
644
Image processing, ROI selection, and ΔF/F calculation
Custom MATLAB pipeline code (Ramamurthy et al, 2023[43]; adapted from LeMessurier,
2019[94]) was used for image processing. Correction for slow XY drift was performed using
dftregistration[95]. Regions-of-interest (ROIs) were manually drawn as ellipsoid regions over the
somata of neurons visible in the average projection across the full imaging movie after registration.
Mean fluorescence of the pixels in each ROI was calculated to obtain the raw fluorescence time
series. For PYR cell imaging, neuropil masks were created as 10 pixel-wide rings beginning two
pixels from the somatic ROI, excluding any pixels correlated with any somatic ROI (r>0.2). Mean
fluorescence of neuropil masks was scaled by 0.3 and subtracted from raw somatic ROI
fluorescence. Neuropil subtraction was not performed for VIP cells, which were spatially well-
separated. The mean fluorescence time series was converted to ΔF/F for each ROI, defined as $(F_t -
F_0)/F_0$, where F_0 is the 20th percentile of fluorescence across the entire imaging movie and F_t is the
fluorescence on each frame.
658
Quantification of whisker-evoked responses.
Whisker-evoked ΔF/F signal on Go trials was quantified in a post-stimulus analysis window (7
frames, 0.799 s), relative to pre-event baseline window (2 frames, 0.270 s). Whisker responses were
measured as (mean ΔF/F in the post-stimulus window – mean ΔF/F in the baseline window) for each
ROI. On NoGo trials, the ΔF/F analysis was aligned to the NoGo stimulus (dummy piezo deflection).
Whisker responses for each cell were normalized by z-scoring to prestimulus baseline activity. For
some analyses, each ROI's Go-NoGo response magnitude to every whisker was also calculated as
(median whisker-evoked ΔF/F signal across Go trials – median ΔF/F signal across NoGo trials). Trials
aborted due to licks occurring during the post-stimulus window (0 - 0.799 s) were excluded from
analyses. If at least one whisker produced a significant response above baseline activity
(permutation test), the cell was considered to be whisker-responsive. This was done by combining
the whisker-evoked ΔF/F signal distribution on Go trials with the ΔF/F signal distribution on NoGo
trials, randomly splitting the combined distribution into two groups and comparing the difference in
their means to the true Go-NoGo distribution difference (10,000 iterations). Differences greater
than the 95th percentile of the permuted distribution were assessed as significant. The nine
whisker response p-values were corrected for multiple comparisons (False Discovery Rate
correction[96]). Cells without a positive ΔF/F response to at least one whisker were considered non-
responsive.
The standard method for assessing whisker-responsiveness used all trials belonging to
each session (combining trials across all history conditions). In the analysis of attentional
modulation of receptive fields, the significance of whisker responses was separately assessed
using only trials in the Prior NoGo category and compared to trials in Prior >1 Hit condition, which
allowed us to test whether there was history-dependent acquisition of whisker-evoked responses
by previously non-responsive cells.
683
Definition of each cell's columnar whisker (CW) and best whisker (BW)
Each cell's anatomical home column was determined by histological localization of the cell
relative to barrel column boundaries. For cells located within column boundaries, the CW was the
whisker corresponding to its anatomical home location. For septa-related cells (i.e., those outside
of column boundaries and above a L4 septum), the CW was the whisker corresponding to the

nearest barrel column. The best whisker (BW) was defined for each cell as the whisker that evoked
the numerically highest magnitude response.

Attention Modulation Index (AMI)
Multiple AMI metrics were used to quantify attentional modulation of whisker response
magnitude in individual cells. The definitions were:

AMI (>1Hit-NoGo):

697
$$AMI_{>1HitSame-NoGo} = \frac{Go_{Prior>1HitSame} - Go_{PriorNoGo}}{|Go_{Prior>1HitSame} + Go_{PriorNoGo}|}$$

699
$$AMI_{>1HitDiff-NoGo} = \frac{Go_{Prior>1HitDiff} - Go_{PriorNoGo}}{|Go_{Prior>1HitDiff} + Go_{PriorNoGo}|}$$

701   AMI(>1HitSame->1HitDiff):

702
$$AMI_{>1HitSame->1HitDiff} = \frac{Go_{Prior>1HitSame} - Go_{Prior>1HitDiff}}{|Go_{Prior>1HitSame} + Go_{Prior>1HitDiff}|}$$

where Go = mean whisker-evoked ΔF/F for current Go trials (on any whisker) with the specified trial
history.

Attentional modulation of receptive fields
Population average 9-whisker receptive fields were constructed centered on the CW, and
included the CW plus the 8 immediately adjacent whiskers. A separate population average
receptive field was calculated for each trial history (**Fig. 4a**). These represent the average tuning of
cells within each whisker column, following each trial history. To test for shifts in receptive fields by
attention to specific whiskers, we first computed the center of mass (CoM) of the population
average receptive fields for each history condition. CoM was calculated in a Cartesian CW centered
whisker space, as defined in **Extended Data Fig. 5b**. The CW position is considered the origin in
this space. Receptive field shifts associated with prior trial history were visualized as vectors from
CoM measured after NoGo trials, to CoM measured after >1 prior hit to specific whiskers.
To quantify the receptive field shift (ΔRF CoM) for individual cells, we defined the *attention*
axis as the axis connecting the CW position to the attended whisker position in the Cartesian CoM
space. We projected the CoM$_{PriorNoGo}$ and CoM$_{Prior>1Hit}$ onto this axis, and computed the receptive
field shift (ΔRF CoM) as the distance between these projected positions normalized to the distance
from CW to attended whisker along the attention axis (**Extended Data Fig. 5c**). Since not all
whisker positions could be sampled for all cells across history conditions, ΔRF CoM was quantified
only for the subset of cells for which at least 6 of the 9 whisker positions were sampled in both Prior
>1 Hit and Prior NoGo conditions. Only response magnitudes at whisker positions sampled in both
Prior >1 Hit and Prior NoGo conditions for any given cell contributed to the CoMs computed for that
cell. The RF shift was computed separately for each attended whisker position that was sampled
for a given cell, and then averaged across these attended whisker positions to generate a single RF
shift metric for that cell.

Somatotopic profile of attentional modulation
To quantify the somatotopic profile of attentional modulation in S1 (**Fig. 3k**), we considered
each of the 9 tested whiskers separately. For each whisker (termed the reference whisker), every
cell was placed in a spatial bin representing its distance to the center of the reference whisker
column in S1. Both columnar and septal-related cells were included. Mean whisker response

magnitude was calculated, separated by trial history, in each bin. This was repeated for all 9
reference whiskers, and **Fig. 3k** shows the average response. Thus, the Prior NoGo trace reflects
normal somatotopy, i.e., the normal point representation of an average whisker. The somatotopic
profile of attentional modulation is evident as the difference between other history conditions and
the Prior NoGo condition.
A similar binning procedure was used to calculate the somatotopic profile of AMI
modulation across S1 columns (**Extended Data Fig. 4a**).

Imaging analysis for VIP cells
Analysis of VIP cell responses was performed similarly to PYR cells, except that neuropil
subtraction was not performed. To separate whisker-evoked VIP responses from slow trends in VIP
baseline activity related to whisker motion, body motion and arousal in the ITI (Ramamurthy et al.,
2023) (and **Fig. 2**) we applied linear baseline detrending. For detrending, the median pre-stimulus
baseline trace (in a 1.07 sec window) was calculated across Go trials (aligned to stimulus onset
time) and NoGo trials (aligned to dummy piezo onset time). A line was fit to this median trace. This
line was extrapolated and subtracted from each individual trial to yield the full peri-stimulus trace.
This linear detrending was done separately for each history condition, due to the differences in pre-
stimulus slopes for each condition. Note that linear detrending for prior same and prior different
categories in each reward condition was identical. Analysis of history effects on VIP whisker
responses (**Fig. 7f-g**) was performed after linear detrending. While VIP cells did not show whisker-
specific attentional modulation (**Fig. 7f-g**), we verified that PYR cells still showed whisker-specific
attentional boosting after detrending with the same methods (data not shown).

Neural decoding from population activity on single trials
We used a generalized linear model (GLM, Matlab 'glmfit') to predict the presence of a
whisker stimulus from the trial-by-trial mean population activity of whisker-responsive L2/3 PYR
cells in 2p imaging experiments. For each session, mean ΔF/F in the post-stimulus window (0 -
0.799 s) for each trial was calculated across all whisker-responsive ROIs that were simultaneously
imaged in a single behavioral session. This single-trial population activity was used as the predictor
of stimulus presence (current Go trial) or stimulus absence (current NoGo trial).
A separate decoder was fit from the data for each imaging session. We fit a logistic
regression model using leave-one-out cross-validation to predict the presence of any of the 9
whiskers, and tested it on hold-out data. Training/testing datasets were randomly re-sampled
(majority class undersampled to match the minority class) to have identical numbers of trials within
each response category (Hit, Miss, CR, FA), in order to remove bias. Decoder performance was
measured as the average fraction of trials classified as containing a whisker stimulus (assessed
over 25-50 iterations) and compared to performance for a decoder trained with shuffled trial labels.
For each field, we defined the fBW (field best whisker) as the whisker which evoked the
numerically highest mean population ΔF/F. A single decoder was trained for each session to
predict any whisker from training data containing Go trials from all whiskers, as well as NoGo trials.
Decoder performance was assessed either for detecting any whisker, or just the fBW, or just non-
fBW trials.

Extracellular recordings
For surgical preparation for mice used in extracellular recording experiments, methods were
similar to that described above, except a lightweight chronic head post was affixed to the skull
using cyanoacrylate glue and Metabond, and ISOI was performed to localize D-row (D1, D2, D3)
barrel columns in S1. A 5-mm diameter glass coverslip (#1 thickness, CS-3R, Warner Instruments)

was placed over the skull, sealed with Kwik-Cast silicone adhesive (World Precision Instruments)
and dental cement. Mice recovered for 7 days prior to the start of behavioral training (~4 weeks
before recording).
The day before recording, mice were anaesthetized with isoflurane, the protective coverslip
was removed, and a craniotomy (~ 1.2 x 1.2 mm) was made over S1, centered over the D-row (D1,
D2, D3) barrel columns localized by ISOI. A plastic ring was cemented around the craniotomy to
create a recording chamber. During recording sessions, mice were anesthetized and positioned on
the rig. A reference ground was attached inside the chamber. The craniotomy and reference wire
were covered in a saline bath. Recordings were made with Neuropixels 1.0 probes using SpikeGLX
software release v.20201024 (http://billkarsh.github.io/SpikeGLX/), Imec phase30 v3.31. Acute
recordings were made in external reference mode with action potentials (AP) sampled at 30 kHz at
500x gain, and local field potential (LFP) sampled at 2.5 kHz at 250x gain. The AP band common
average referenced and band-pass filtered from 0.3 kHz to 6 kHz. The Neuropixels probe was
mounted on a motorized stereotaxic micromanipulator (MP-285, Sutter Instruments) and advanced
through the dura mater (except in cases where the dura had detached during the craniotomy). To
reduce insertion-related mechanical tissue damage and to increase the single unit yield, the probe
was lowered with a slow insertion speed of 1-2 μm/sec. The probe was first lowered to 700 μm, and
a short 10-minute recording was conducted to map its location in S1. After identifying the columnar
whisker for the recording penetration location, probe insertion continued until the final depth was
reached. The probe was then left untouched for ~ 20 minutes. The craniotomy was sealed after
probe insertion with silicone sealant (Kwik-Cast, World Precision Instruments) to prevent drying.
Anesthesia was then discontinued, and mice were allowed to fully wake up before recording.
After recording was complete for the day, the probe was removed, the craniotomy was
sealed with silicone sealant (Kwik-Cast, World Precision Instruments), and the recording chamber
was sealed with a cover glass and a thin layer of dental cement. 3-4 sequential days of recording
were performed in each mouse, with the probe located in a different whisker column in S1 on each
day. On the final recording day, a Neuropixels probe was coated with red-fluorescent DiI (1,1'-
Dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate; Sigma-Aldrich) dissolved in 100%
ethanol, 1-2mg/mL, which was allowed to partially dry on probe before probe insertion. The probe
was briefly inserted to deposit DiI at recording and several fiducial sites. The mouse was
euthanized and the brain extracted, sectioned tangentially to the pial surface, and processed to
stain for cytochrome oxidase (CO). DiI deposition sites in L2/3 were localized relative to column
boundaries in CO from L4 (**Extended Data Fig. 6b**).
The columnar location of each recording penetration was determined from DiI marks on the
last recording day, plus relative locations of other penetrations based on reference images of
surface vasculature and microdrive coordinates. The laminar depth of each recording was
determined from current source density (CSD) and LFP power spectrum analysis, as described
below.
821
Analysis of extracellular recording data
<u>Spike Sorting</u>
Spike sorting was performed by automatic clustering using Kilosort3 followed by manual
curation using the 'phy' GUI (https://github.com/kwikteam/phy). Isolated units were manually
inspected for mean spike waveform, stability over time, and inter-spike interval refractory period
violations (we required that < 2% of intervals < 1.5 ms). Only well-isolated single units were
analyzed. Single units were classified as regular-spiking or fast-spiking based on trough-to-peak
duration of the spike waveform at the highest-amplitude recording channel, with a separation

criterion of 0.45ms[97] (**Extended Data Fig. 6c**). Only data from regular spiking units was analyzed
here.
832
<u>Layer assignment using CSD and LFP power spectrum analysis</u>
To calculate the CSD for each recording penetration, the whisker stimulus-evoked local
field potential (LFP, 500Hz low-pass) was calculated for each Neuropixels channel. LFP traces were
normalized to account for differences in channel impedance and interpolated between channels
(20 um site spacing, 0.6x interpolation) before calculating the second spatial derivative, which
defines the CSD[98] . For visualization, CSDs were convolved with a 2D (depth x time) Gaussian,
which revealed depth-restricted regions of current sources and sinks in response to each whisker
stimulus. The L4-L5A boundary was defined from CSD as the zero-crossing between the most
negative current sink (putative L4) and the next deeper current source (putative L5A) (**Extended**
Data Fig. 6f). To estimate the brain surface location (defining the top of L1), we computed the LFP
power spectrum as a function of channel depth. A sharp increase in low-frequency LFP power
marked the brain surface, which was used to verify correct selection of the L4-L5A boundary from
CSD. Each cortical layer was then assigned boundary depths based on layer thicknesses reported
in Lefort et al., 2009[99].
847
Whisker response quantification and attentional modulation for spike recordings
Firing rates for Go and NoGo trials were quantified in a 0.5 s window after stimulus onset
(lick-free window p. Units were classified as whisker-responsive or non-responsive by testing for
greater firing rate on Go vs. NoGo trials, using a permutation test, as for the 2p imaging data. To
quantify whisker evoked response magnitude, firing rate for each unit was z-scored relative to pre-
stimulus baseline. Peristimulus time histograms (PSTHs) were constructed using 10-ms bins
aligned to stimulus onset. AMI metrics were used to quantify attentional modulation of whisker-
evoked spiking, and were calculated exactly as for 2p imaging data, but from z-scored whisker-
evoked firing rate in the post-stimulus window.
857
Statistical analysis of summary data
Statistics were performed in Matlab. Sample size (n) and p-value for each analysis is
reported in the figure panel, with the statistical test reported in the figure legend, or sometimes in
the *Results* section. Permutation tests for the difference in means were used to assess differences
in mean between two groups (referred to in brief as 'permutation test') or differences in the mean of
a distribution relative to zero. Binomial exact test was used to assess the statistical significance of
deviations from the expected distribution of observations into two categories. Summary data are
reported as mean ± SEM, unless otherwise specified. Population means were compared using
permutation tests, corrected for multiple comparisons (False Discovery Rate correction[48]) as
needed. To compare population means with multiple subgroups, we first assessed whether a main
effect was present using a permutation test, and then performed post hoc tests for pairwise means,
correcting for multiple comparisons (with False Discovery Rate correction). We used a significance
level (alpha) of 0.05. All statistical tests reported in the *Results* use n of cells, but we also verified
that all major behavioral and neural effects were consistent across individual mice. We show
individual mouse data for comparison with population data summaries. To account for inter-
individual variability, we also verified all key results using a linear mixed effects model with the
formula:
875
876         $Response\ Variable \sim Fixed\ Effect + (1|Mouse\ Sex) + (1|Mouse\ ID)$
877

878 where mouse sex and mouse ID are modeled as random effects.
879

References

1. Maunsell, J.H.R. Neuronal Mechanisms of Visual Attention. Annu Rev Vis Sci 1, 373-391 (2015).
2. Hromádka, T. & Zador, A.M. Toward the mechanisms of auditory attention. Hear Res 229, 180-185 (2007).
3. Gallace, A. & Spence, C. 147 Tactile attention. in In Touch with the Future: The sense of touch from cognitive neuroscience to virtual reality 0 (Oxford University Press, 2014).
4. Awh, E., Belopolsky, A.V. & Theeuwes, J. Top-down versus bottom-up attentional control: a failed theoretical dichotomy. Trends in cognitive sciences 16, 437-443 (2012).
5. Theeuwes, J. & Failing, M. Attentional Selection: Top-Down, Bottom-Up and History-Based Biases. (Cambridge University Press, Cambridge, 2020).
6. Meyer, K.N., Sheridan, M.A. & Hopfinger, J.B. Reward history impacts attentional orienting and inhibitory control on untrained tasks. Atten Percept Psychophys 82, 3842-3862 (2020).
7. Failing, M. & Theeuwes, J. Selection history: How reward modulates selectivity of visual attention. Psychonomic Bulletin & Review 25, 514-538 (2018).
8. Chelazzi, L., Perlato, A., Santandrea, E. & Della Libera, C. Rewards teach visual selective attention. Vision Research 85, 58-72 (2013).
9. Anderson, B.A. A value-driven mechanism of attentional selection. J Vis 13 (2013).
10. Anderson, B.A. Neurobiology of value-driven attention. Current Opinion in Psychology 29, 27-33 (2019).
11. Nobre, A.C. & Stokes, M.G. Premembering Experience: A Hierarchy of Time-Scales for Proactive Attention. Neuron 104, 132-146 (2019).
12. Anderson, B.A., Laurent, P.A. & Yantis, S. Value-driven attentional capture. Proceedings of the National Academy of Sciences 108, 10367 (2011).
13. Anderson, B.A., et al. The past, present, and future of selection history. Neuroscience & Biobehavioral Reviews 130, 326-350 (2021).
14. Addleman, D.A. & Jiang, Y.V. Experience-Driven Auditory Attention. Trends in Cognitive Sciences 23, 927-937 (2019).
15. Chen, D. & Hutchinson, J.B. What Is Memory-Guided Attention? How Past Experiences Shape Selective Visuospatial Attention in the Present. Curr Top Behav Neurosci 41, 185-212 (2019).
16. Anderson, B.A. & Britton, M.K. Selection history in context: Evidence for the role of reinforcement learning in biasing attention. Atten Percept Psychophys 81, 2666-2672 (2019).
17. Noudoost, B., Chang, M.H., Steinmetz, N.A. & Moore, T. Top-down control of visual attention. Curr Opin Neurobiol 20, 183-190 (2010).
18. Matteucci, G., et al. Cortical sensory processing across motivational states during goal-directed behavior. Neuron 110, 4176-4193.e4110 (2022).
19. Luo, T.Z. & Maunsell, J.H.R. Attention can be subdivided into neurobiological components corresponding to distinct behavioral effects. Proceedings of the National Academy of Sciences 116, 26187-26194 (2019).

20. Krauzlis, R.J., Wang, L., Yu, G. & Katz, L.N. What is attention? Wiley Interdiscip Rev Cogn Sci
14, e1570 (2023).

21. Luo, T.Z. & Maunsell, J.H. Neuronal Modulations in Visual Cortex Are Associated with Only
One of Multiple Components of Attention. Neuron 86, 1182-1188 (2015).

22. Luo, T.Z. & Maunsell, J.H.R. Attentional Changes in Either Criterion or Sensitivity Are
Associated with Robust Modulations in Lateral Prefrontal Cortex. Neuron 97, 1382-
1393.e1387 (2018).

23. Peck, C.J., Jangraw, D.C., Suzuki, M., Efem, R. & Gottlieb, J. Reward modulates attention
independently of action value in posterior parietal cortex. J Neurosci 29, 11182-11191
(2009).

24. Desimone, R. & Duncan, J. Neural mechanisms of selective visual attention. Annu Rev
Neurosci 18, 193-222 (1995).

25. Cohen, M.R. & Maunsell, J.H.R. A Neuronal Population Measure of Attention Predicts
Behavioral Performance on Individual Trials. The Journal of Neuroscience 30, 15241 (2010).

26. Reimer, J., et al. Pupil Fluctuations Track Fast Switching of Cortical States during Quiet
Wakefulness. Neuron 84, 355-362 (2014).

27. Reimer, J., et al. Pupil fluctuations track rapid changes in adrenergic and cholinergic activity
in cortex. Nature Communications 7, 13289 (2016).

28. Mathis, A., et al. DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. Nature Neuroscience 21, 1281-1289 (2018).

29. Womelsdorf, T., Anton-Erxleben, K., Pieper, F. & Treue, S. Dynamic shifts of visual receptive
fields in cortical area MT by spatial attention. Nature Neuroscience 9, 1156-1160 (2006).

30. Womelsdorf, T., Anton-Erxleben, K. & Treue, S. Receptive field shift and shrinkage in
macaque middle temporal area through attentional gain modulation. J Neurosci 28, 8934-
8944 (2008).

31. Anton-Erxleben, K., Stephan, V.M. & Treue, S. Attention reshapes center-surround receptive
field structure in macaque cortical area MT. Cereb Cortex 19, 2466-2478 (2009).

32. He, S., Cavanagh, P. & Intriligator, J. Attentional resolution and the locus of visual
awareness. Nature 383, 334-337 (1996).

33. Yang, H., Kwon, S.E., Severson, K.S. & O'Connor, D.H. Origins of choice-related activity in
mouse somatosensory cortex. Nat Neurosci 19, 127-134 (2016).

34. Yamashita, T. & Petersen, C. Target-specific membrane potential dynamics of neocortical
projection neurons during goal-directed behavior. Elife 5 (2016).

35. Jun, J.J., et al. Fully integrated silicon probes for high-density recording of neural activity.
Nature 551, 232-236 (2017).

36. Lee, S., Kruglikov, I., Huang, Z.J., Fishell, G. & Rudy, B. A disinhibitory circuit mediates motor
integration in the somatosensory cortex. Nat Neurosci 16, 1662-1670 (2013).

37. Fu, Y., et al. A cortical circuit for gain control by behavioral state. Cell 156, 1139-1152
(2014).

38. Zhang, S., et al. Selective attention. Long-range and local circuits for top-down modulation
of visual cortex processing. Science 345, 660-665 (2014).

39. Batista-Brito, R., Zagha, E., Ratliff, J.M. & Vinck, M. Modulation of cortical circuits by top-
down processing and arousal state in health and disease. Current opinion in neurobiology
52, 172-181 (2018).

40. Makino, H., Hwang, E.J., Hedrick, N.G. & Komiyama, T. Circuit Mechanisms of Sensorimotor
Learning. Neuron 92, 705-721 (2016).
41. Myers-Joseph, D., Wilmes, K.A., Fernandez-Otero, M., Clopath, C. & Khan, A.G. Disinhibition
by VIP interneurons is orthogonal to cross-modal attentional modulation in primary visual
cortex. Neuron 112, 628-645.e627 (2024).
42. Rahmatullah, N., et al. Hypersensitivity to Distractors in Fragile X Syndrome from Loss of
Modulation of Cortical VIP Interneurons. The Journal of Neuroscience 43, 8172 (2023).
43. Ramamurthy, D.L., et al. VIP interneurons in sensory cortex encode sensory and action
signals but not direct reward signals. Curr Biol 33, 3398-3408.e3397 (2023).
44. Speed, A. & Haider, B. Probing mechanisms of visual spatial attention in mice. Trends in
Neurosciences (2021).
45. Speed, A., Del Rosario, J., Mikail, N. & Haider, B. Spatial attention enhances network,
cellular and subthreshold responses in mouse visual cortex. Nature Communications 11,
505 (2020).
46. Kanamori, T. & Mrsic-Flogel, T.D. Independent response modulation of visual cortical
neurons by attentional and behavioral states. Neuron 110, 3907-3918.e3906 (2022).
47. McBride, E.G., Lee, S.J. & Callaway, E.M. Local and Global Influences of Visual Spatial
Selection and Locomotion in Mouse Primary Visual Cortex. Curr Biol 29, 1592-1605.e1595
(2019).
48. Lehnert, J., et al. Visual attention to features and space in mice using reverse correlation.
Current Biology 33, 3690-3701.e3694 (2023).
49. You, W.-K. & Mysore, S.P. Endogenous and exogenous control of visuospatial selective
attention in freely behaving mice. Nature Communications 11, 1986 (2020).
50. Goldstein, S., Wang, L., McAlonan, K., Torres-Cruz, M. & Krauzlis, R.J. Stimulus-driven visual
attention in mice. Journal of Vision 22, 11-11 (2022).
51. Musall, S., Kaufman, M.T., Juavinett, A.L., Gluf, S. & Churchland, A.K. Single-trial neural
dynamics are dominated by richly varied movements. Nature Neuroscience 22, 1677-1686
(2019).
52. Schmidt, F., Haberkamp, A. & Schmidt, T. Dos and don'ts in response priming research. Adv
Cogn Psychol 7, 120-131 (2011).
53. Musall, S., Haiss, F., Weber, B. & von der Behrens, W. Deviant Processing in the Primary
Somatosensory Cortex. Cerebral Cortex 27, 863-876 (2017).
54. Akrami, A., Kopec, C.D., Diamond, M.E. & Brody, C.D. Posterior parietal cortex represents
sensory history and mediates its effects on behaviour. Nature 554, 368-372 (2018).
55. Hachen, I., Reinartz, S., Brasselet, R., Stroligo, A. & Diamond, M.E. Dynamics of history-
dependent perceptual judgment. Nature Communications 12, 6036 (2021).
56. Waiblinger, C., McDonnell, M.E., Borden, P.Y. & Stanley, G.B. Emerging experience-
dependent dynamics in primary somatosensory cortex reflect behavioral adaptation.
bioRxiv, 2021.2001.2029.428886 (2021).
57. Lak, A., et al. Reinforcement biases subsequent perceptual decisions when confidence is
low, a widespread behavioral phenomenon. eLife 9, e49834 (2020).
58. Fischer, J. & Whitney, D. Serial dependence in visual perception. Nature Neuroscience 17,
738-743 (2014).

59. Sheehan, T.C. & Serences, J.T. Attractive serial dependence overcomes repulsive neuronal
adaptation. PLOS Biology 20, e3001711 (2022).

60. Tal-Perry, N. & Yuval-Greenberg, S. Contraction bias in temporal estimation. Cognition 229,
105234 (2022).

61. Webster, M.A. 655 Adaptation Aftereffects in the Perception of Faces. in The Oxford
Compendium of Visual Illusions (ed. A.G. Shapiro & D. Todorovic) 0 (Oxford University Press,
2017).

62. Urai, A.E., de Gee, J.W., Tsetsos, K. & Donner, T.H. Choice history biases subsequent
evidence accumulation. eLife 8, e46331 (2019).

63. Ashwood, Z.C., et al. Mice alternate between discrete strategies during perceptual
decision-making. Nat Neurosci 25, 201-212 (2022).

64. Roy, N.A., Bak, J.H., Akrami, A., Brody, C.D. & Pillow, J.W. Extracting the dynamics of
behavior in sensory decision-making experiments. Neuron 109, 597-610.e596 (2021).

65. Urai, A.E., Braun, A. & Donner, T.H. Pupil-linked arousal is driven by decision uncertainty
and alters serial choice bias. Nature Communications 8, 14637 (2017).

66. Reynolds, J.H., Pasternak, T. & Desimone, R. Attention increases sensitivity of V4 neurons.
Neuron 26, 703-714 (2000).

67. Briggs, F., Mangun, G.R. & Usrey, W.M. Attention enhances synaptic efficacy and the signal-
to-noise ratio in neural circuits. Nature 499, 476-480 (2013).

68. Chalk, M., et al. Attention reduces stimulus-driven gamma frequency oscillations and spike
field coherence in V1. Neuron 66, 114-125 (2010).

69. Cohen, M.R. & Maunsell, J.H. Attention improves performance primarily by reducing
interneuronal correlations. Nature neuroscience 12, 1594-1600 (2009).

70. Luck, S.J., Chelazzi, L., Hillyard, S.A. & Desimone, R. Neural Mechanisms of Spatial
Selective Attention in Areas V1, V2, and V4 of Macaque Visual Cortex. Journal of
Neurophysiology 77, 24-42 (1997).

71. Staiger, J.F. & Petersen, C.C.H. Neuronal Circuits in Barrel Cortex for Whisker Sensory
Perception. Physiol Rev 101, 353-415 (2021).

72. Engel, T.A., et al. Selective modulation of cortical state during spatial attention. Science
354, 1140-1144 (2016).

73. Silver, M.A., Ress, D. & Heeger, D.J. Topographic maps of visual spatial attention in human
parietal cortex. J Neurophysiol 94, 1358-1371 (2005).

74. Tootell, R.B., et al. The retinotopy of visual spatial attention. Neuron 21, 1409-1422 (1998).

75. Woldorff, M.G., et al. Retinotopic organization of early visual spatial attention effects as
revealed by PET and ERPs. Hum Brain Mapp 5, 280-286 (1997).

76. Hattori, R., Danskin, B., Babic, Z., Mlynaryk, N. & Komiyama, T. Area-Specificity and
Plasticity of History-Dependent Value Coding During Learning. Cell 177, 1858-1872.e1815
(2019).

77. Hwang, E.J., Dahlen, J.E., Mukundan, M. & Komiyama, T. History-based action selection bias
in posterior parietal cortex. Nature Communications 8, 1242 (2017).

78. Findling, C., et al. Brain-wide representations of prior information in mouse decision-
making. bioRxiv, 2023.2007.2004.547684 (2023).

79. Miller, K.J., Botvinick, M.M. & Brody, C.D. Value representations in the rodent orbitofrontal
cortex drive learning, not choice. eLife 11, e64575 (2022).

80. Marmor, O., Pollak, Y., Doron, C., Helmchen, F. & Gilad, A. History information emerges in
the cortex during learning. eLife 12, e83702 (2023).

81. Banerjee, A., et al. Value-guided remapping of sensory cortex by lateral orbitofrontal cortex.
Nature 585, 245-250 (2020).

82. Chéreau, R., et al. Dynamic perceptual feature selectivity in primary somatosensory cortex
upon reversal learning. Nature Communications 11, 3245 (2020).

83. Petty, G.H. & Bruno, R.M. Attentional modulation of secondary somatosensory and visual
thalamus of mice. eLife. 13:RP97188 (2024).

84. Kastner, S. & Arcaro, M.J. The Thalamus in Attention. in The Thalamus (ed. M.M. Halassa)
324-339 (Cambridge University Press, Cambridge, 2022).

85. Pi, H.J., et al. Cortical interneurons that specialize in disinhibitory control. Nature 503, 521-
524 (2013).

86. Hu, F. & Dan, Y. An inferior-superior colliculus circuit controls auditory cue-directed visual
spatial attention. Neuron 110, 109-119.e103 (2022).

87. Drew, P.J. & Feldman, D.E. Intrinsic Signal Imaging of Deprivation-Induced Contraction of
Whisker Representations in Rat Somatosensory Cortex. Cerebral Cortex 19, 331-348 (2009).

88. Wang, H.C., LeMessurier, A.M. & Feldman, D.E. Tuning instability of non-columnar neurons
in the salt-and-pepper whisker map in somatosensory cortex. Nature Communications 13,
6611 (2022).

89. Landers, M., Pytte, C. & Zeigler, H.P. Reversible blockade of rodent whisking: Botulinum
toxin as a tool for developmental studies. Somatosensory & Motor Research 19, 358-363
(2002).

90. Yang, H., Kwon, S.E., Severson, K.S. & O'Connor, D.H. Origins of choice-related activity in
mouse somatosensory cortex. Nature Neuroscience 19, 127-134 (2016).

91. Green, D.M. & Swets, J.A. Signal detection theory and psychophysics (John Wiley, Oxford,
England, 1966).

92. Pronneke, A., et al. Characterizing VIP Neurons in the Barrel Cortex of VIPcre/
tdTomato Mice Reveals Layer-Specific Differences. Cereb Cortex 25, 4854-4868 (2015).

93. Schindelin, J., et al. Fiji: an open-source platform for biological-image analysis. Nature
methods 9, 676-682 (2012).

94. LeMessurier, A.M. (2019). Imaging_analysis_pipeline. GitHub 5a21c9f.
https://github.com/alemessurier/imaging_analysis_pipeline

95. Guizar-Sicairos, M., Thurman, S.T. & Fienup, J.R. Efficient subpixel image registration
algorithms. Opt. Lett. 33, 156-158 (2008).

96. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological) 57, 289-300 (1995).

97. Barthó, P., et al. Characterization of neocortical principal cells and interneurons by network
interactions and extracellular features. J Neurophysiol 92, 600-608 (2004).

98. Freeman, J.A. & Nicholson, C. Experimental optimization of current source-density
technique for anuran cerebellum. J Neurophysiol 38, 369-382 (1975).

99. Lefort, S., Tomm, C., Floyd Sarria, J.C. & Petersen, C.C.H. The Excitatory Neuronal Network
of the C2 Barrel Column in Mouse Primary Somatosensory Cortex. Neuron 61, 301-316
(2009).

Acknowledgments

This work was supported by NIH Grant 2 R01 NS092367 (D.E.F.) and NIH Grant 5 K99 NS129753-02 (D.L.R.).

Author Contributions

D.L.R. and D.E.F. designed the study. D.L.R., L.R., C.C., S.L., and A.C. performed the experiments. D.L.R., L.R., C.C., and S.L. analyzed the data. D.L.R., L.R., and D.E.F. wrote the manuscript.

Competing Interests

The authors declare no competing interests.

Figures and Figure legends

Figure 1. Recent reward history cues spatially specific attention for whisker touch. a, Whisker detection task in head-fixed mice. **b,** Trial structure. Delay period was 0, 500, or 1000 ms in different mice (see **Figure S1**). Bottom, Trial types and trial outcomes. **c,** Example trial sequence showing interleaved Go and NoGo trials, with the identity of the deflected whisker on each Go trial chosen randomly. Water drop indicates successful Hit and reward. **d,** Mean effect of trial history on detection sensitivity (d-prime) on the current trial. Error bars are SEM across sessions. One or more prior Hits to the same whisker increased d' for detecting that whisker on the current trial ('attend toward'), while prior Hits to a different whisker decreased d' (attend away). P-values: (1) from permutation test (prior same 1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior different >1 Hit: p = 1e-4), (2) from linear mixed effects model with fixed effect of prior history class (p = 8.7e-27). **e,** Same effect in a single example mouse. Left, the underlying effect on hit rate for 5 example sessions (thin lines) and mean ± SEM for all 34 sessions. Right, d' for this mouse from 34 sessions. **f,** d' did not increase when the prior trial was an unrewarded hit to the same whisker or earned very small reward (<4%: p = 0.49, >4%: p = 1e-4). P-values are for difference in d' between prior same and prior different conditions (permutation test). **g,** Effect of >1 prior hits on Hit Rate and FA rate on the current trial. Connected symbols are the three history conditions for one mouse. Gray lines, receiver-operating characteristic (ROC) curves for different d' levels. Prior hits to the same whisker drove a whisker-specific increase in hit rate over FA rate that improved d'. **h,** Behavioral shift in d' for each individual mouse. **i,** Behavioral shifts in criterion (*c*) were also observed, but were less whisker-specific. **j,** Whisker-specific changes in d' (Δ d') and c (Δ *c*) for each mouse (number is mouse identity). We focus on Δ d' to index whisker-specific attention. **k,** Spatial gradient of the attention effect. The facial position of the prior Hit whisker is plotted on the x-axis, relative to the current trial whisker. P-values are for difference from prior same (same row: p =0.27, same arc: 1e-3, diagonal: 1e-4, further: 1e-4, permutation test). Bottom, each position shown schematically. **l,** Temporal profile of the whisker-specific attention effect (3-5 s: 5e-3, 5-7 s: 1e-4, 7-9 s: 1e-4, 9-11 s: 0.17, >11s: 0.15, permutation test), P-values are from permutation test. Bottom, Inter-Go-trial intervals sampled in the task. **m,** Flexible targeting of attention. x-axis represents identity of whisker in current Go trial, grouped into C, D, or E-row whiskers (left), or arcs 1, 2 or 3 (right). For all of these whiskers, prior reward history boosts detection in a whisker-specific way. Bottom, each row or arc shown schematically. P-values are for prior same vs. prior different > 1hit (p = 1e-4 for all whisker positions, permutation test). See also **Extended Figure 1**.

Figure 1. Recent reward history cues spatially specific attention for whisker touch.

Figure 2. Whisker motion, body motion, and arousal do not account for whisker-specific behavioral effects. a, Mean whisker motion, platform motion (proxy for body motion), and pupil size traces, obtained using DeepLabCut from behavioral videos, across 43,647 trials, 74 sessions in 9 mice. Each panel shows the last 3 seconds of the intertrial interval (ITI) period after the prior trial, plus the stimulus period of the current trial. Traces and shading are mean ± SEM across all trials. Dashed line, stimulus onset (Go trials) or dummy piezo onset (NoGo trials). Prior trial identity is indicated by the Prior trial history. Δ whisker position and Δ platform position traces were zeroed to the mean of positions in a 0.25 sec window prior to stimulus onset (t= 0). Pupil size was normalized within each session to the mean pupil size across the whole session. Bottom row, lick histogram. **b,** Mean whisker movement, body movement, and pupil size change during the stimulus period. Prior Hits increased stimulus-evoked whisker and body motion on subsequent trials (Δ whisker motion, prior >1 hit same vs. prior NoGo: p = 1e-4, prior >1 hit same vs. >1 hit different: p = 0.39; Δ body motion, prior >1 hit same vs. prior NoGo: p = 1e-4, prior >1 hit same vs. >1 hit different: p = 0.39; Δ pupil area, prior >1 hit same vs. prior NoGo: p = 0.49, prior >1 hit same vs. >1 hit different: p = 0.53). p-values are for >1 Prior Hit vs. Prior NoGo (top), and >1 Prior Hit Same vs. > 1 Prior Hit Different (right) (permutation test). **c,** Design of Botox experiment. Behavior was assayed on an average of 12 sessions prior to Botox injection, and 7 sessions after Botox whisker paralysis. **d,** Reward history-dependent attention effect in each of the 4 mice tested, for standard sessions (before Botox, open symbols) and Botox sessions (filled symbols). M, Mouse numbers as in **Fig. 1**. Large points are mean ± SEM across mice. Conventions as in **Fig. 1j**. Whisker paralysis did not alter the mean whisker-specific d-prime effect or criterion effect.

Figure 2. Body motion, whisker motion, or arousal does not account for whisker-specific behavioral effects.

Figure 3. Neural correlates of attentional capture in L2/3 pyramidal cells in S1. a-b, Example imaging field in S1 centered on the D3 column. L2/3 PYR cells are expressing GCaMP6s. Scale bar = 100 µm. **c,** Example trials showing responses to the columnar whisker D3, but only weak responses to the surround whisker C3 that were strongly modulated by prior trial history. C3 responses increased following multiple prior hits to the C3 whisker (left), but not following other history conditions (right). **d,** Mean ΔF/F traces across all neurons in 70 sessions, separated by trial history. Each trace is the mean response to any Go whisker (solid) or on NoGo trials (dash). Shading shows SEM, which is often thinner than the mean trace. **e,** Quantification of whisker-evoked ΔF/F by current trial type (Go or NoGo) and by prior trial condition (prior same 1 Hit vs prior NoGo: p = 1e-4, prior same >1 Hit vs prior NoGo: p = 1e-4, prior different >1 Hit vs prior NoGo: p = 0.1, prior same >1 Hit vs prior different >1 Hit: p = 1e-4, permutation test). Bars are SEM. **f,** Mean modulation of ΔF/F response magnitude calculated by individual mouse. Bars are SEM. **g,** Correlation between history-based modulation of PYR whisker responses and behavioral d-prime, by mouse (r = 0.837, p = 6.9e-4). **h,** $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ for each cell. Positive values denote greater response compared to the prior NoGo condition. **i,** $AMI_{>1HitSame->1HitDiff}$ for each cell. **j,** Boosting of whisker-evoked ΔF/F responses as a function of somatotopic offset between prior >1Hit whisker and the current trial whisker. Bars show d-prime (data from **Fig. 1k**, for the 6 mice used in PYR imaging) for comparison. **k,** Somatotopic organization of attentional capture. Left, the y-axis shows mean ΔF/F evoked by a reference whisker, as a function of cell position in S1 relative to the center of the reference whisker column. When calculated from Prior NoGo trials (thick black trace), this defines the classic point representation of a single whisker. This is boosted in prior same >1 hit trials, but not in prior different >1 hit trials, or prior same miss trials. Right, same data presented as difference from the Prior NoGo condition, to plot magnitude of modulation by trial history.

Figure 3. Neural correlates of attentional capture in L2/3 pyramidal cells in S1.

Figure 4. Attentional cueing involves receptive field shifts toward attended whiskers. a, Mean whisker-evoked ΔF/F traces for all whisker-responsive cells in all imaged columns. . Center, mean ± SEM (across N=5399 cells) for trials when prior trial was NoGo, separated by the identity of the current trial whisker. This reports the average whisker tuning curve for these neurons, in the absence of attentional cueing. Outer flanks, the whisker responses measured when prior trial history was >1Hit to the indicated attentional target whisker (thick purple trace is mean, thin traces show ± SEM). Purple fill is drawn between mean traces to aid visualization. r, u, c, d denote rostral, up, caudal, or down from CW. **b,** Mean tuning center-of-mass (CoM) when the prior trial was NoGo (black circle) vs. after prior >1 Hit to each of the indicated whiskers as defined in panel **a**. CoM coordinate system is shown in **Extended Data Figure 4b**. Vectors are color-coded for whether the target whisker was rostral, caudal, up, or down from the CW. **c,** Magnitude of CoM shift along the attention axis, as defined in **Extended Data Figure 4c**. Negative values are shifts away from the attended whisker. The mean CoM shift was significantly greater than zero (p=3.6e-3, permutation test).

a Attentional target whisker (cued by prior >1 hit)

(r,u) (-,u) (c,u)

Whisker-responsive cells
70 sessions, 6 mice, 5399 cells

Up

Rostral CW Caudal

(r,-) (c,-)

Down

2 ΔF/F
1 s

Whisker row

(r,d) (-,d) (c,d)

Whisker arc

b

(r,-)

Rostral Caudal
(c,-)
(r,u) (-,-) (c,u)
(c,d)
(r,d)

0.1 whisker units

70 sessions, 6 mice, 5399 cells

Up
(r,u) (-,-) (c,u)
(-,u)
(-,d) Down (c,d)
(r,d)

c

no shift p= 3.6e-3 complete shift

173 cells

cells

-1.5 -1 -0.5 0 0.5 1 1.5

Δ RF CoM
CoM shift along
attention axis

away from attended whisker   towards attended whisker

Figure 4. Receptive fields shift toward the attended whisker during attentional capture.

Figure 5. Attentional cueing improves neural decoding of attended whiskers on single trials. a, Neural decoder design. Left, example field showing L2/3 PYR cells tuned to different best whiskers intermixed in each column, consistent with prior studies[88]. Right, on each trial, population mean ΔF/F was calculated across all responsive cells. Leave-one-out cross-validation was used to fit a logistic regression predicting stimulus presence (any whisker Go trial) or absence (NoGo trial). **b,** Decoder performance determined from held-out trials. Each dot is one session. Decoder performance is low for any whisker, because many trials are Go trials for whiskers that are not strongly represented in the imaging field. Decoder performance for fBW trials is high, because this whisker is strongly represented in the imaging field. **c-d,** Mean ΔF/F **(c)** and fraction of trials decoded as a stimulus (Go) trial **(d)**, separated by current trial type and outcome. Each dot is a session. All prior history conditions are combined in **c-d** panels (ΔF/F, HR vs. Miss: p=1e-4, FA vs. CR: p=1e-4; decoder, HR vs. Miss: p=1e-4, FA vs. CR: p= 9e-4, permutation test). **e,** Mean decoder performance separated by trial history type, when current trial is any Go whisker (maroon or yellow) or a NoGo trial (gray dash). Solid lines with circle markers show decoder performance across all current trial outcomes, and dotted lines with asterisk markers show decoder performance on current hit trials only. (All trials, prior same >1 Hit vs prior NoGo: p =1e-4, prior different >1 Hit vs prior NoGo: p =0.21, prior same >1 Hit vs prior different >1 Hit: p =1e-4; Hit trials only, prior same >1 Hit vs prior NoGo: p =7e-3, prior different >1 Hit vs prior NoGo: p =0.05, prior same >1 Hit vs prior different >1 Hit: p =2e-3, permutation test). **f,** Same as **(e)**, but for decoder performance when current trial is an fBW Go trial or a NoGo trial. **g,** Same as **(e)**, but for decoder performance when current trial is a non-fBW Go trial or a NoGo trial. **h,** Summary of attentional modulation of decoding accuracy for the prior >1 Hit trial history condition. >1 prior Hit to a whisker improves single-trial decoding for non-fBW whiskers, but not for the fBW, in each field (non-fBW: p=1e-4, fBW: p=0.648). P-values are for prior same vs. prior different > 1 hit (permutation test).

Figure 5. Attentional cueing improves neural decoding of attended whiskers on single trials.

Figure 6. Attentional effects on extracellular single-unit spiking in S1. a, Neuropixels recording during the whisker detection task. **b,** Mean PSTH for an example L2/3 RS unit recorded in the D1 whisker column, across all Go trials (D1, C2, delta, and gamma whiskers, top) and NoGo trials (bottom). Whisker-evoked responses were boosted when prior trial history was 1 or >1 Hit to the same whisker, and reduced when prior trial was >1 Hit to a different whisker. **c, e, g,** Layer-specific analysis for RS units of population mean PSTH (10 ms bins) for Go trials, by trial history. **d, f, h,** For the same units, population average whisker-evoked response on current Go trials (500-ms window) by trial history, or for current NoGo trials in an equivalent window (prior same >1 Hit vs prior different >1 Hit, L2/3: p=6.9e-4, L4: 0.08, L5a/b: 1e-4, permutation test). **i,** Cumulative histogram of $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ values for RS units by layer. **j,** Cumulative histogram of $AMI_{>1HitSame->1HitDiff}$ by layer. **k,** Mean AMI values by layer ($AMI_{>1HitSame-NoGo}$ vs. $AMI_{>1HitDiff-NoGo}$, L2/3: 0.03, L4: 0.65, L5a/b: 0.01; $AMI_{>1HitSame->1HitDiff}$, L2/3: 0.01, L4:0.51, L5a/b: 0.06).

Figure 6. Attentional effects on extracellular single-unit spiking in S1.

Figure 7. L2/3 VIP cells carry a general arousal signal, but not a whisker-specific attentional signal. a, Circuit model for potential VIP cell role in arousal, movement, and attentional modulation of sensory responses in sensory cortex. **b,** Example imaging field in L2/3 in a VIP-Cre;Ai162D mouse. Scale bar = 50 μm. **c,** Mean ΔF/F traces from L2/3 VIP cells across all mice and sessions during the last 3 seconds of the intertrial interval (ITI) period after the prior trial for prior rewarded (prior 1 Hit or prior >1 Hit trials) and prior unrewarded trials (prior NoGo or prior Miss). Traces are zeroed to the last 2 frames prior to stimulus onset. The declining baseline, evident on both prior rewarded (blue) and unrewarded (gray) trials, is due to VIP cell encoding of arousal, whisker motion, and body motion which all decline in the inter-trial interval as described in a previous study[43], and shown in **Fig. 2a**. Prior rewarded trials show a steeper peak and steeper decline due to licking and reward consumption at the end of the prior trial. **d,** Mean ΔF/F traces across all VIP neurons in 103 sessions, separated by trial history. Each trace is the mean response to any Go whisker (solid) or on NoGo trials (dash). Shading shows SEM. Conventions as in **Fig. 3d**. **e,** Same data as in **(d)** after subtracting an extrapolation of the pre-stimulus baseline trend (see Methods). Whisker-evoked responses are modulated by prior hits on either the same or different whiskers, which is consistent with a general arousal effect, but not whisker-specific attentional capture. **f,** Quantification of trial history effects for L2/3 VIP cells (prior same 1 Hit vs prior NoGo: p =0.11, prior same >1 Hit vs prior NoGo: p =1e-3, prior different >1 Hit vs prior NoGo: p =1e-3, prior same >1 Hit vs prior different >1 Hit: p =0.54, permutation test). Conventions as in **Fig. 3e**. **f,** $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$ for each cell. Positive values denote greater response compared to the prior NoGo condition. **g,** $AMI_{>1HitSame->1HitDiff}$ for each cell.

Figure 7. L2/3 VIP cells carry an arousal signal, but not a whisker-specific attention signal.

Extended Data

Extended Data Figure 1. Further characterization of reward history-cued attention. a, Behavioral performance in a single example session. Sliding *d'* was computed from Hit rate and FA rate using a 50-trial sliding window. A *d'* cutoff (black dashed line) was applied to exclude trials at the start and end of the session. **b,** Definition of trial history classes for current Go trials and current NoGo trials, using current D2 whisker trials as an example. When calculating *d'* and *c*, Go and NoGo trials with matched trial histories were selected, as illustrated. **c,** Delay periods used in different groups of mice. 0 ms (no delay period) was used only for behavioral analyses. For 500 ms delay periods (used for 2p imaging and extracellular spike recordings), and for 1000 ms delay periods (used for 2p imaging), reward volume was ramped during the delay period to encourage delayed licking (see Methods). **d,** Whisker-specific attentional cueing based on reward history was observed for mice trained with all 3 delay periods. (p=1e-4 for prior same vs. prior different for all delays, permutation test). **e,** Reaction times on Hit trials for different trial histories, for mice trained without a delay period. Prior hits decreased reaction time relative to prior NoGo (prior same >1: p=1e-4, prior different >1 hit: p=1e-4), but there was no significant difference between prior same and prior different conditions (prior same >1 hit vs. prior different >1 hit: p=0.21). , permutation test). **f,** Trial history effect as a function of whisker stimulus amplitude (i.e., stimulus strength) on the current trial. This shows that history-dependent boosting in d' is strongest for low amplitude whisker stimuli (<50 μm: p = 0.03, 50-100 μm: p = 1e-4, 100-150 μm: p = 1e-4, 150-200 μm: p = 1e-4, 200-300 μm: p = 1e-4). P-values are for prior-same vs. prior-different (permutation test).**g,** The three task designs used to set different whisker stimulus probabilities. In all designs, reward probability for a Hit on any whisker, P(reward), was 100%. In the equal probability (EqP) design, each of the 9 whiskers was presented with equal probability. In the high-probability (HiP) design, sessions were divided into one or more blocks of a few hundred trials, and within each block a single whisker (example, gray box) had a much higher probability (0.8) than the others (0.025). In the HiPsame design, all whiskers were presented at an equal overall probability, but sequential presentation was biased so that there was a higher probability (0.25-0.4) of the same whisker being repeated on the next trial, for all whisker identities. In all 3 designs, whisker identity was randomly selected on each Go trial, following the intended probability structure. **h,** Whisker-specific attentional effects cued by reward history were observed in all three task designs. 13 mice were tested with EqP and HiP designs (top), and 7 mice were tested with EqP and HiPSame designs (bottom).

Extended Data Figure 1. Further characterization of reward history-cued attention.

Extended Data Figure 2. Behavioral imaging and additional analysis of history-dependent effects on spontaneous behaviors. **a,** Example behavioral movie frame showing DeepLabCut tracking of whisker motion (pad + 3 whiskers), platform motion (proxy for body motion), and pupil size measurement. **b,** Mean whisker motion, body motion, and pupil size changes during the ITI period of the current trial, as a function of trial history. Conventions and p-values as in **Fig. 2b**. Prior Hits increased average whisker and body motion during the ITI period (Δ whisker motion, prior >1 hit same vs. prior NoGo: p = 1e-4, prior >1 hit same vs. >1 hit different: p = 1e-4; Δ body motion, prior >1 hit same vs. prior NoGo: p = 1e-4, prior >1 hit same vs. >1 hit different: p = 0.55; Δ pupil area, prior >1 hit same vs. prior NoGo: p = 0.07, prior >1 hit same vs. >1 hit different: p = 0.16).

Extended Data Figure 2. Behavioral imaging and additional analysis of history-dependent behavioral effects.

Extended Data Figure 3. Additional analysis of history cueing effects on PYR activity. a, Example imaging field showing modulation of whisker-evoked response magnitude (color scale) by trial history. Top 2 rows, the average response to all Go whiskers. Response magnitude was increased for many cells in the prior Same >1 Hit condition, and not in the prior Different > 1 Hit condition. Bottom row, activity on NoGo trials, showing expectation or general arousal effect after > 1 prior Hit to any whisker. **b,** Mean whisker-evoked ΔF/F traces, as in **Fig. 3d**, calculated separately for whisker-responsive or non-responsive neurons (identified based on prior NoGo trials). Conventions as in **Fig 3d. c,** Quantification of the data in (b). Conventions as in **Fig. 3e**. Prior hits to the same whisker increase whisker-evoked responses in responsive cells (prior same 1 Hit vs prior NoGo: p =1e-4, prior same >1 Hit vs prior NoGo: p =1e-4, prior different >1 Hit vs prior NoGo: p =0.09, prior same >1 Hit vs prior >1 Hit current NoGo: p =1e-4, permutation test), but do not cause the appearance of whisker responses in non-responsive cells (prior same 1 Hit vs prior NoGo: p =0.02, prior same >1 Hit vs prior NoGo: p =1e-4, prior different >1 Hit vs prior NoGo: p =0.60, prior same >1 Hit vs prior >1 Hit current NoGo: p =0.34, permutation test). **d,** CDFs of AMI for each cell, separated by mouse. The whisker-specific effect on response magnitude was observed in 6/6 mice. **e,** Mean ΔF/F trace for each L2/3 PYR cell, sorted by $AMI_{>1HitSame->1HitDiff}$ for each cell.

Extended Data Figure 3. Additional analysis of history cueing effects on PYR cell activity.

Extended Data Figure 4. Additional analysis of somatotopic profile of attentional boosting in S1. a, Mean AMI for neuronal responses to a reference whisker, binned by cell location relative to the center of the reference whisker column. Left, $AMI_{>1HitSame-NoGo}$ and $AMI_{>1HitDiff-NoGo}$. Right, $AMI_{>1HitSame->1HitDiff}$. Conventions as in **Fig. 3k**, except that finer spatial bins are used within the reference whisker column. Whisker-specific attentional boosting was evident in a region <200 um from the reference column center ($AMI_{>1HitSame-NoGo}$ vs. $AMI_{>1HitDiff-NoGo}$, 0 -50 µm: 1e-4, 50-140 µm: 4e-4,140-280 µm: 0.20, 280- 420 µm: 0.48, 420-560 µm: 0.03, 560-700 µm: 0.92, permutation test; $AMI_{>1HitSame->1HitDiff}$, 0 -50 µm: 1e-4, 50-140 µm: 1e-4,140-280 µm: 0.46, 280- 420 µm: 0.23, 420-560 µm: 0.01, 560-700 µm: 0.88, permutation test). **b-c,** Trial history-dependent modulation of neural responses to the CW. For this analysis, only cells located within columnar boundaries were analyzed (i.e., septal cells were excluded). **b,** >1 prior Hit to CW increases CW-evoked responses. Conventions as in Fig. 3d. **c,** Quantification of trial history effects on CW responses (prior same 1 Hit vs prior NoGo: p =0.55, prior same >1 Hit vs prior NoGo: p =1e-4, prior different >1 Hit vs prior NoGo: p =0.76, prior same >1 Hit vs prior different >1 Hit: p =1e-4, permutation test). Conventions as in **Fig. 3e**.

Extended Data Figure 4. Additional analysis of somatotopic profile of attentional boosting in S1.

Extended Data Figure 5. Additional analysis of receptive field shifts. a, Modulation of mean whisker receptive field when prior trial was >1 hit to the CW (purple), relative to when prior trial was NoGo (black). Format as in **Figure 4a**. **b,** Coordinate system for calculating tuning CoM. The CW is at (0,0), and whiskers are separated by 1 unit in row or arc dimensions. **c,** Calculation of CoM tuning shift along the attention axis. The attention axis was defined as the line connecting the CW coordinate (0,0) with the coordinate of the attended whisker. The CoM measured after prior NoGo trials and after prior >1 hit to the attended whisker were projected onto this axis, and the CoM shift was calculated as distance along this axis, with positive values indicating shift toward the attended whisker.

Extended Data Figure 5. Additional analysis of receptive field shifts.

Extended Data Figure 6. Neuropixels recording methods. **a,** Behavioral shifts in d' and c for mice that were initially trained on the 9-whisker task version, and then were tested either using 9 whiskers (open) or 4-5 whiskers only (filled). When performing the task with 4-5 whiskers, reward history still drives a whisker-specific Δ d' effect, but this is smaller than when performing with 9 whiskers. Numbered points are individual mice. Larger symbols are mean ± SEM across mice. **b,** Example histological reconstruction of a Neuropixels recording site to the D1 whisker column. Dashed circles are probe tracks marked with DiI, outlines are from L4 barrels, white circles are reconstructed penetration sites. Asterisk marks the recording site for the example cell in **Fig. 6b** and in **S6-1** panel **e-g**. Scale bar = 100 µm. **c,** Classification of FS and RS units by trough-to-peak time in extracellular spike waveform. Bottom, example RS and FS spike waveform. **d,** Left, example current source density (CSD) analysis to identify laminar boundaries. See Methods for details. Right, power spectrum analysis to identify the top of L1, defined as the transition from high to low LFP power across a wide range of frequency bands. **e,** Mean spike waveform for the example L2/3 RS unit from **Fig. 6b**. Blue traces, individual spikes. **f,** Whisker-evoked spike rasters separated by current trial Go whisker (gamma, delta, D1, C2) or no whisker stimulus (NoGo trials), for the example L2/3 RS unit from **Fig. 6b**. All trial histories are combined in this plot. **g,** Whisker-evoked PSTH for the same example unit as in panel **e**. Plotting conventions as in **Fig. 6c**.

Extended Data Figure 6. Neuropixels recording methods.

Extended Data Figure 7. Additional analysis of attentional signals in VIP cells. a, Mean raw ΔF/F trace for each L2/3 VIP cell, sorted by $\text{AMI}_{>1\text{HitSame}\text{->}1\text{HitDiff}}$ for each cell. **b,** Mean detrended ΔF/F raw trace for each L2/3 VIP cell, sorted by $\text{AMI}_{>1\text{HitSame}\text{->}1\text{HitDiff}}$ for each cell.

Extended Data Figure 7. Additional analysis of attentional signals in VIP cells.