Clostridioides difficile colonization is not mediated by bile salts and utilizes

Stickland fermentation of proline in an *in vitro* model

Xiaoyun Huang[1], April E. Johnson[1], Joshua N. Brehm[2], Thi Van Thanh Do[1], Thomas A.

Auchtung[1], Hugh C. McCullough[1], Armando I. Lerma[1], Sigmund J. Haidacher[3,4], Kathleen

M. Hoch[3,4], Thomas D. Horvath[3,4,5], Joseph A. Sorg[2], Anthony M. Haag[3,4], and Jennifer M.

Auchtung[1]*

*Address correspondence to Jennifer M. Auchtung, jauchtung2@unl.edu

[1]Department of Food Science and Technology and Nebraska Food for Health Center, University

of Nebraska-Lincoln, Lincoln, NE 68588 USA

[2]Department of Biology, Texas A&M University, College Station, TX USA

[3]Department of Pathology and Immunology, Baylor College of Medicine, Houston, TX USA

[4]Texas Children's Microbiome Center, Department of Pathology, Texas Children's Hospital,

Houston, TX USA

[5]Department of Pharmacy Practice & Translational Research, University of Houston, Houston,

TX USA

Present address: Xiaoyun Huang, Fermenich Incorporated, Princeton, NJ USA; Thomas

Auchtung, Synbiotic Health, Lincoln, NE USA; Hugh McCullough, Department of Biological

Systems Engineering, Lincoln, NE USA

Xiaoyun Huang and April Johnson contributed equally to this work. Author order determined by

majority of ten coin flips.

Keywords: Stickland fermentation, bile metabolism, microbiota, antibiotic disruption

ABSTRACT

Treatment with antibiotics is a major risk factor for *Clostridioides difficile* infection, likely due to depletion of the gastrointestinal microbiota. Two microbiota-mediated mechanisms thought to limit *C. difficile* colonization include conversion of conjugated primary bile salts into secondary bile salts toxic to *C. difficile* growth, and competition between the microbiota and *C. difficile* for limiting nutrients. Using a continuous flow model that simulates the nutrient conditions of the distal colon, we investigated how treatment with six clinically-used antibiotics influenced susceptibility to *C. difficile* infection in 12 different microbial communities cultivated from healthy individuals. Antibiotic treatment reduced microbial richness; disruption varied by antibiotic class and microbiota composition, but did not correlate with *C. difficile* susceptibility. Antibiotic treatment also disrupted microbial bile salt metabolism, increasing levels of the primary bile salt, cholate. However, changes in bile salt did not correlate with increased *C. difficile* susceptibility. Further, bile salts were not required to inhibit *C. difficile* colonization. We tested whether amino acid fermentation contributed to persistence of *C. difficile* in antibiotic-treated communities. *C. difficile* mutants unable to use proline as an electron acceptor in Stickland fermentation due to disruption of proline reductase (*prdB-*) had significantly lower levels of colonization than wild-type strains in four of six antibiotic-treated communities tested. Inability to ferment glycine or leucine as electron acceptors, however, was not sufficient to limit colonization in any communities. This data provides further support for the importance of bile salt-independent mechanisms in regulating colonization of *C. difficile.*

IMPORTANCE

C. difficile is one of the leading causes of hospital-acquired infections and antibiotic-associated

diarrhea. Several potential mechanisms through which the microbiota can limit *C. difficile*

infection have been identified and are potential targets for new therapeutics. However, it is

unclear which mechanisms of *C. difficile* inhibition represent the best targets for development of

new therapeutics. These studies demonstrate that in a complex *in vitro* model of *C. difficile*

infection, colonization resistance is independent of microbial bile salt metabolism. Instead, the

ability of *C. difficile* to colonize is dependent upon its ability to metabolize proline, although

proline-dependent colonization is context-dependent and is not observed in all disrupted

communities. Altogether, these studies support the need for further work to understand how bile-

independent mechanisms regulate *C. difficile* colonization.

INTRODUCTION

Clostridioides difficile is one of the leading causes of nosocomial infections due to its

transmissibility as environmentally-resistant spores and its ability to infect patients who have

been treated with antibiotics (1, 2). Many classes of antibiotics can disrupt the colonic microbiota

(3–5); disruption can decrease production of inhibitory metabolites (6–9) and reduce competition

for limiting nutrients (10–13), providing favorable conditions for *C. difficile* infection. Although

multiple mechanisms for colonization resistance have been identified, an understanding of the

hierarchical importance of these mechanisms in *C. difficile* colonization and disease is just

beginning to emerge.

Microbial bile salt metabolism has been extensively studied for its role in limiting *C. difficile*

colonization. The primary bile salts cholate and chenodeoxycholate are synthesized in the liver

and are conjugated to glycine or taurine to improve solubility (14). Once secreted into the

intestine, microbial enzymes begin modifying these bile salts, removing conjugated amino acids

and dehydroxylating primary bile salts into secondary bile salts (14). *C. difficile* spore

germination is stimulated by cholate-family bile salts (cholate, taurocholate, glycocholate,

deoxycholate) and inhibited by chenodeoxycholate-family bile salts (15, 16), although there is

variation in these responses between strains (17). Germination is enhanced by amino acid (15,

18) and calcium co-germinants, which act synergistically with bile salts to enhance germination

(19). Secondary bile salts (deoxycholate, lithocholate) inhibit growth of vegetative *C. difficile* in

vitro (7, 15, 20), and low levels of secondary bile salts correlate with *C. difficile* infection in

humans (21, 22) and in mouse models (7, 23, 24).

However, recent studies have demonstrated that our understanding of the role of bile salts in

C. difficile colonization resistance may be incomplete. *C. difficile* spore colonization and

fulminant disease was observed in *Cypb8b1*[-/-] mice unable to make cholate-family bile salts (25),

and resistance to *C. difficile* colonization was observed even though these mice do not produce

the secondary bile salts deoxycholate or lithocholate. *Clostridium scindens*, a microbe inferred to

inhibit *C. difficile* growth through dehydroxylation of primary bile salts to secondary bile salts

(24), was also shown to inhibit *C. difficile in vitro* through production of tryptophan-derived

antibiotics (9) and other uncharacterized metabolites (26) and to potentially compete with *C.*

difficile for nutrients required for growth in *Cypb8b1*[-/-] monocolonized mice (25). In addition,

both cholate and deoxycholate were shown to induce similar *C. difficile* stress responses,

although 10X higher concentrations of cholate were used to observe these effects (27).

Competition for nutrients between commensal microbes and *C. difficile* has long been

postulated as a mechanism for colonization resistance, with Wilson and colleagues utilizing

continuous-flow culture systems to demonstrate competition between the microbiota and *C.*

difficile in vitro (28, 29). *C. difficile* can metabolize mucin monosaccharides *in vitro* (30), and

preferentially expresses pathways for degradation of mucin monosaccharides in mouse models

(31, 32), indicating that metabolism of mucin monosaccharides may be a niche open to *C.*

difficile during infection. Similarly, *C. difficile* metabolism of carbohydrates and sugar alcohols

from the host and its diet may be another niche open to *C. difficile* during infection, as

expression of genes in these metabolic pathways increase during infection in germ-free and

antibiotic-treated mice (11, 31, 32). In addition to carbohydrate metabolism, *C. difficile* also

efficiently utilizes amino acids through Stickland fermentation (33), which is a metabolic

pathway limited primarily to proteolytic Clostridial species (34), providing a unique metabolic

niche within the GI tract. Increasing evidence from human (10, 35) and mouse (11, 13, 25, 31,

32, 36) studies points to Stickland fermentation with proline as an electron acceptor as a

preferred nutritional niche for *C. difficile* in the GI tract. *C. difficile* can also use glycine (33) and

leucine (37) as electron acceptors in Stickland fermentation, with a glycine reductase mutant

recently shown to delay morbidity in a hamster model of disease (38). Multiple regulatory

pathways converge to coordinately regulate the expression of proline, glycine, and leucine

reductase pathways (34), and more work is needed to understand how the integration of these

pathways contributes to disease in different environmental and nutritional contexts.

Previously, we described an *in vitro* minibioreactor array (MBRA) model for characterization

of *C. difficile* colonization resistance in the presence of human fecal microbial communities

under nutritional conditions that simulate the distal colon (39), that has subsequently been used

to characterize microbes that inhibit *C. difficile* colonization and/or toxin production (40–42). In

this study, we used the MBRA model to investigate how treatment with six clinically-used

antibiotics differentially impacted susceptibility to *C. difficile* infection in microbial communities

cultivated from 12 healthy individuals. As expected, we observed that antibiotic treatment

reduced microbial richness and altered microbial diversity, although the extent of antibiotic-

mediated disruption varied across individual communities and was not well correlated with

susceptibility to *C. difficile* infection. Antibiotic treatment also reduced microbial metabolism of

cholate to deoxycholate, although there was no correlation between deoxycholate levels and

susceptibility to *C. difficile* infection. To further test whether bile salts contributed to *C. difficile*

colonization resistance, we cultivated fecal communities in the presence and absence of bovine

bile. Similar to what was observed by Aguirre et al. (25) in *Cypb8b1*$^{-/-}$ mice, we observed bile

salts were not required for *C. difficile* colonization resistance and that the low level of *C. difficile*

spore germination that occurred in the absence of bile was sufficient to allow colonization of

antibiotic-treated communities by *C. difficile* spores. Using *C. difficile* mutants unable to utilize

proline, glycine, or leucine as electron acceptors during Stickland fermentation due to a mutation

in proline reductase (*prdB* (43)) glycine reductase (*grdA*), or 2-hydroxyisocaproate

dehydrogenase (*ldhA*), we demonstrated that proline reduction was required for *C. difficile* to

colonize a subset of antibiotic-disrupted microbial communities, whereas glycine and leucine

reduction was not required for *C. difficile* colonization in the same communities. These results

can serve as a foundation for further mechanistic characterization of the hierarchical importance

of different nutritional environments in limiting *C. difficile* colonization.

RESULTS

Loss of microbial richness is not sufficient to promote *C. difficile* colonization. Previously,

we described an *in vitro* model for culturing microbial communities from fecal samples under

conditions that simulated the nutritional environment of the distal colon (44). Communities were

shown to maintain a subset of microbial richness and diversity found in the starting fecal

samples, similar to other *in vitro* models (44) and germ-free animal models colonized with

human feces (45, 46). Further, communities cultured from most healthy individuals demonstrated

resistance to colonization by *C. difficile*, which could be disrupted by treatment with clindamycin

(39, 41). To better understand how this model can be used to measure effects of antibiotics on

microbiota disruption and *C. difficile* colonization, we compared treatment with six individual

antibiotics (Fig. 1A) on microbial community composition and *C. difficile* susceptibility of

communities cultured from twelve different healthy individuals. These antibiotics were selected

because they are all used clinically, but vary in their therapeutic spectrum of antimicrobial

activity.

Fourteen independent bioreactors were inoculated with each fecal sample and allowed to

stabilize in continuous culture for seven days. Reactors were treated in duplicate with one of six

antibiotics twice daily for five days (Figure 1B); two reactors served as untreated controls. Two

days after the end of antibiotic treatment, communities were challenged with vegetative cells of

C. difficile strain 2015 (CD2015), a ribotype 027 clinical isolate (39). Samples were collected for

analysis of microbial community composition by 16S rRNA gene sequencing from three

replicates of the initial fecal sample prior to inoculation, and from cultured communities prior to

administration of antibiotics (Day -7), at the end of antibiotic treatment (Day -2), and two days

later (Day 0), just prior to *C. difficile* challenge (Figure 1A). Samples were collected from

cultured communities for analysis of *C. difficile* levels on days 3 and 7 post-infection.

Prior to antibiotic treatment (Day -7), there were no differences in richness (Observed OTUs

with ≥99% average nucleotide identity, Figure S1A) or microbial diversity (Inverse Simpson,

Figure S1B) between treatment groups, although there was a loss in richness and diversity

compared to the fecal sample inocula as reported previously ((44); Figures S1A and S1B).

Communities were composed primarily of taxa from the Bacteroidetes, Firmicutes,

Proteobacteria, and Verrucomicrobia phyla, with members of Fusobacteria, Synergistetes, and

Actinobacteria phyla found in some communities (Figure S2).

Treatment with antibiotics significantly reduced microbiota richness at the end of antibiotic

treatment (Day -2) relative to the baseline sample collected from the same reactor at Day -7

(Figure 1C) and led to significantly lower levels of richness compared to untreated controls on

Day -2 (Figure S1A); decreases ranged from 1.1-fold (cefaclor) to 1.5-fold (clindamycin).

Microbial diversity declined significantly from baseline to the end of antibiotic treatment with

just two antibiotics – clindamycin and ceftriaxone (Day -2, Figure 1D). These antibiotics also

showed significant lower levels of diversity compared to untreated samples at Day -2, as did the

Augmentin-treated communities (Figure S1B).

We also measured changes in shared community composition (Figure 1D) and structure

(Figures 1E, F) by calculating Jaccard and Bray-Curtis dissimilarity measures from communities

before antibiotic treatment (Day -7) to communities after antibiotic treatment (Day -2). (Jaccard

and Bray-Curtis dissimilarity measures compare the proportion of taxa that are shared between

two communities, with Jaccard providing an unweighted measure of dissimilarity between

communities and Bray-Curtis providing a measure of dissimilarity that is weighted based on taxa

abundance. In both cases, values closer to 0 are more similar.) As had been reported previously

(44), cultivation led to significant shifts in microbial composition and structure from initial fecal

samples (Figures 1E, F) to Day -7. Following this initial reorganization in community

composition and structure, continued cultivation led to lower levels of change in community

composition and structure (Figure 1E and 1F, see untreated samples on Day -2 and Day 0).

Treatment with all antibiotics led to larger changes in community composition from Day -7 to

Day -2 compared to untreated communities (Figure 1E), whereas only Augmentin and

clindamycin also led to significantly larger changes in community structure compared to

untreated communities (Figure 1F).

Two days following the end of antibiotic treatment (Day 0), reduced richness compared to

baseline samples persisted for all antibiotic-treated communities with the exception of those

treated with cefaclor (Figure 1C), although Augmentin-treated communities exhibited small, but

statistically significant increases in microbial richness (Figure 1C). Augmentin-treated

communities also exhibited small, but statistically significant increases in microbial diversity

(Figure 1D) and decreases in Jaccard (Figure S1E) and Bray-Curtis (Figure S1F) dissimilarity,

indicating a potential return towards baseline following cessation of this antibiotic. Clindamycin-

treated communities continued to exhibit decreased microbial diversity (Figure 1D) and less

similar community composition to baseline (Figure 1E).

C. difficile susceptibility, measured as levels of *C. difficile* detected on Day 7 following

challenge with *C. difficile* cells, varied across different antibiotic treatments (Figure 1E) and

fecal donors (Figure S3). While *C. difficile* levels on Day 3 and 7 were collected, we used levels

of *C. difficile* on Day 7 rather than Day 3 as a marker of colonization to provide sufficient time

for loss of non-replicating cells and spores under continuous flow culture conditions. Of the

twelve fecal samples tested, untreated communities from six fecal samples were resistant to

colonization, defined as *C. difficile* colony forming units (CFU)/mL undetectable in both

replicates, and untreated communities from one fecal sample were colonized at high levels, with

both replicates colonized with *C. difficile* levels in the highest quartile. The remaining five

untreated communities exhibited variable susceptibility to colonization (Figure S3A), with two

fecal communities exhibiting low (1st quartile) or undetectable levels of colonization in both

replicates. Clindamycin was the only antibiotic that significantly increased median levels of *C.*

difficile colonization above levels observed in untreated communities (Figure 2A); ten of twelve

fecal sample communities were colonized following treatment with clindamycin (Figure S3A).

Susceptibility to colonization varied across communities from different fecal samples that were

treated with other antibiotics (Figure S3A).

We tested whether there were correlations between levels of richness or microbial diversity

on Day -2 or Day 0 and found that there were no significant correlations with *C. difficile*

colonization levels (Figure S4). We also assessed whether changes in community composition

(Jaccard dissimilarity) or structure (Bray-Curtis dissimilarity) from Day -7 to Day -2 or Day 0

correlated with *C. difficile* colonization levels. We observed weak correlations with changes in

microbial composition from baseline (Day -7) to Day -2 (Figure 2B) and Day 0 (Figure 2C).

There were no significant correlations with changes in community structure from baseline to Day

-2 or Day 0 (Figure 2E). Similar results were observed for *C. difficile* levels on Day 3 (Figure

S5). We used LEfSe (47) to identify specific OTUs whose abundance on Day -2 or Day 0

correlated with *C. difficile* colonization. We found a small number of OTUs whose abundance

on Day -2 or Day 0 significantly correlated with *C. difficile* levels (Figure 2F), with four OTUs

more abundant in non-colonized communities on Day -2 and/or Day 0 and five OTUs more

abundant in colonized communities on Day -2 and/or Day 0.

Antibiotic treatment alters bile salt metabolism. Bioreactor medium (BRM3) contains bovine

bile as a complex source of bile salts. Analysis of bioreactor medium indicated that cholate

family bile salts predominated, with approximately 36% taurocholate (198 µM), 34%

glycocholate (190 µM), 10% cholate (56 µM), and 2% deoxycholate (13 µM) of total bile salts

measured (Figure S6A-B). To understand how microbiota cultivation and antibiotic treatment

altered bile salt pools in bioreactor cultures, we focused on quantification of the amino acid

conjugated primary bile salt, taurocholate, the primary bile salt, cholate, and the secondary bile

salt, deoxycholate across the communities described in Figure 1. We observed that microbiota

cultivation led to a >6,500-fold decrease in median levels of taurocholate and a 45-fold increase

in median levels of deoxycholate in spent culture supernatant in untreated communities on Day -

2 compared to fresh medium (Figure S6C). Treatment with four antibiotics – Augmentin,

ceftriaxone, clindamycin, and fidaxomicin – led to significantly higher levels of cholate

compared to untreated communities at the end of antibiotic treatment (Day -2; Figure 3A).

Deoxycholate and taurocholate levels were not significantly different between untreated

communities and communities treated with antibiotics on Day -2 (Figure 3B-C). There were no

significant differences in bile salt levels between untreated and antibiotic-treated communities on

Day 0 (Figure 3D-F). There was no correlation between levels of *C. difficile* on Day 7 and

cholate levels on Day -2 (Figure 3G) or Day 0 (R^2=8.5X10^{-8}, p=0.997).

Previous studies have found that deoxycholate inhibits the growth of vegetative *C. difficile*

cells (7, 15, 20), with inhibitory concentrations ranging from 0.01 – 0.1% (250 - 2500 µM). The

median concentration of deoxycholate measured in untreated communities on Day 0, the day of

C. difficile challenge, was 602 µM (Figure 3B); median concentrations in antibiotic-treated

communities ranged from 408 µM (Augmentin) to 608 µM (clindamycin), but were not

significantly different than those observed in untreated communities (Figure 3E). Because both

C. difficile susceptibility and deoxycholate levels varied by fecal donor and antibiotic treatment

(Figure S7), we tested whether deoxycholate levels on Day 0 correlated with *C. difficile*

colonization. We observed that there were no significant differences in levels of deoxycholate

between colonized and uncolonized communities (Figure 3H), nor were there significant

correlations between deoxycholate levels and *C. difficile* levels on Day 7 (Figure 3I). In addition,

we observed that 21.4% of colonized communities (15/70) had deoxycholate levels $\geq$1000 $\square$M

on Day 0 (Figure 3I, symbols colored red).

Bile was not required for *C. difficile* colonization resistance *in vitro*. While the previous

results were consistent with the hypothesis that secondary bile salts do not mediate colonization

resistance in communities cultured in bioreactors, we could not rule out effects of other bile salts

that were not measured. To test whether bile salts were required, we monitored colonization

resistance of microbial communities established from five fecal samples cultured in parallel in

media with and without added bile. Specifically, replicate bioreactors were established for each

fecal sample in media either containing or lacking bovine bile. Triplicate communities were

treated with clindamycin as previously described (48), while the remaining three replicates for

each type of media were left untreated (Figure 4A). Communities were challenged with

vegetative cells of CD2015 and *C. difficile* levels were monitored over time. We found *C.*

difficile colonization was similarly suppressed in communities cultured in the presence or

absence of bile and that treatment with clindamycin led to similar levels of *C. difficile*

colonization (Figure 4B).

Bile was not required for *C. difficile* spores to colonize *in vitro*. Cholate family bile salts are

known to greatly enhance *C. difficile* spore germination during *in vitro* culture (15, 49, 50),

although they are not required for spore germination in a mouse model of infection (25).

Previously, we reported that we were unable to colonize a clindamycin-treated microbial

community cultured in an MBRA model with *C. difficile* spores (39). Because these studies were

performed using a microbial community formed from pooling twelve fecal donors, we tested the

ability of the spores to colonize clindamycin-treated communities established from fecal samples

collected from five individuals. We found that the majority of clindamycin-treated communities

could be colonized following challenge with *C. difficile* spores (Figure 4C), although

communities from one fecal sample were colonized at a lower frequency by spores than by

vegetative cells (Figure S8). We then tested whether bile salts were required for *C. difficile*

spores to colonize clindamycin-treated fecal microbial communities cultured in MBRAs. We

observed that colonization by spores was not significantly different between communities

cultured in media with and without bile (Figure 4D).

Because cholate family bile salts were previously reported to enhance germination, we

compared the ability of media with and without bile to enhance colony formation of spores in

pure culture. Consistent with previous observations (15, 49), incubation of CD2015 spores in

bioreactor medium containing bile for one hour enhanced colony formation by germinated *C.*

difficile spores by >400-fold compared to incubation in the presence of bioreactor medium

without bile (Figure 5). We next tested how spent culture medium collected from clindamycin-

treated communities cultured in the presence or absence of bile would impact spore germination.

Low levels of spore germination were observed in spent culture medium from fecal communities

cultured in the presence and absence of bile (Figure 5). Altogether, this data indicates that the

levels of bile salts present in spent culture medium from communities cultured in the presence of

bile do not have significant impacts on germination. Further, while the levels of spore

germination and colony formation in spent culture medium are low, these low levels of

germination would result in ~10^3 vegetative cells, a level previously shown to infect antibiotic

disrupted communities (39).

Proline metabolism was required for *C. difficile* to colonize a subset of fecal samples *in*

***vitro*.** Proteolytic metabolism through Stickland fermentation has previously been reported as an

important nutritional niche for *C. difficile* in human (10, 35) and animal models of infection (11,

13, 25, 31, 32, 36, 38). As bioreactor medium contains low levels of fermentable carbohydrates

and does not contain mucin-associated monosaccharides, we hypothesized that Stickland

fermentation with proline as an electron acceptor could be important for colonization of

antibiotic-disrupted fecal communities. To test this hypothesis, we compared the colonization of

mutants defective in the ability to metabolize proline (*prdB*::CT (43)) to a wild type (wt) strain;

both wild type and *prdB*::CT were in the CD196 background, which is a non-epidemic ribotype

027 isolate (51) . Fecal samples from six healthy individuals were cultured in replicate

bioreactors, treated with clindamycin, and challenged with wt or *prdB*::CT strains using the

approach described in Fig. 4A.

Overall, we observed that fecal communities challenged with *prdB* mutants showed decreasing

levels of colonization over time, with significantly lower levels observed on day 3 and 5 of

colonization compared to wild type (Figure 6A). However, the ability of wt and *prdB*::CT strains

to colonize was dependent upon the fecal community tested (Figure 6B). Wild-type strains

colonized five of the six clindamycin-disrupted communities (FS515, FS228, FS235, FS685,

FS133) whereas *prdB*::CT strains failed to colonize two of these susceptible communities

(FS228, FS235). In the other three susceptible communities, there was either a lack of *prdB*::CT

colonization in the majority of replicates tested (FS515: 10/12 replicates; FS685; 8/13

replicates), or there was no difference between levels of wt and *prdB*::CT colonization levels

(FS133). Based on these results, we conclude that Stickland fermentation with proline as an

electron acceptor can play an important role in the ability of *C. difficile* to persist in complex

communities *in vitro*, but this is dependent upon the composition of the fecal community and its

ability to limit *C. difficile* colonization through other mechanisms.

As both glycine and leucine can also serve as electron acceptors for Stickland

fermentation, we hypothesized that fermentation of glycine or leucine may also contribute to

persistence in clindamycin-treated communities. We tested the ability of mutants defective in

Stickland fermentation of glycine and leucine due to mutations in glycine reductase ($\Delta grdA$) and

2-hydroxyisocaproate dehydrogenase ($\Delta ldhA$), respectively to persist in clindamycin-treated

communities. Persistence of these mutants, generated in the hypervirulent ribotype 027 R20291

background, were compared to R20291 wild type strains in four of five fecal samples tested in

Figure 6A (FS515, FS228, FS235, and FS133). In these studies, we saw no significant

differences in colonization levels between wild type and mutant strains over time, indicating that

defects in glycine or leucine fermentation may not be sufficient to limit *C. difficile* colonization

in this model.

DISCUSSION

The factors that govern whether or not an individual exposed to *C. difficile* will go on to develop

symptomatic disease are complex. While it is clear that disruption of the GI microbiota is a key

risk factor for infection and disease, the extent to which different microbes interact with each

other and the host to limit *C. difficile* infection and disease progression are not completely

understood. Developing tools that allow microbe and metabolite interactions to be investigated in

the absence of a host can help to provide insights into whether mechanisms that inhibit or

promote *C. difficile* colonization are more likely to be causative or correlative. This level of

understanding is necessary as new therapeutic approaches that more narrowly target *C. difficile*,

such as defined microbial consortia are developed.

To gain greater insights into factors that govern *C. difficile* colonization resistance in a fecal

MBRA model, we investigated how six different clinically-used antibiotics impacted the ability

of the microbiota to resist *C. difficile* colonization. As expected, we observed that antibiotic

treatment led to loss of microbial richness, although the extent of microbiota disruption varied

both by the class of antibiotic tested and by the composition of the microbial community.

Consistent with previous meta-analyses of human antibiotic exposure and risk for *C. difficile*

infection in non-hospitalized patients (52, 53), clindamycin was most strongly associated with

susceptibility to colonization in antibiotic-treated communities, with ten of twelve communities

tested showing susceptibility following treatment. However, the magnitude of microbiota

disruption across antibiotic treatments did not correlate with susceptibility to *C. difficile*

colonization, with some communities with significant losses in richness resisting *C. difficile*

colonization and other communities with modest richness losses exhibiting susceptibility to

colonization. This data provides further support for the hypothesis that *C. difficile* colonization

resistance is likely due to the presence of specific microbes and their functions that limit *C.*

difficile colonization rather than overall microbiota diversity.

Due to the compositional diversity between fecal communities tested, there were relatively

few conserved taxa whose presence or absence correlated with susceptibility to *C. difficile*

colonization, which is consistent with the model that differences in microbiota function rather

than composition are important for understanding *C. difficile* susceptibility. Similar to previous studies (41, 54), however, levels of multiple *Lachnospiraceae* OTUs were higher in resistant communities.

To better understand which microbiota functions may be important for colonization resistance in this model, we investigated two mechanisms previously proposed to regulate *C. difficile* colonization: microbial metabolism of bile acids and Stickland fermentation with proline. glycine, or leucine as an electron acceptor. We observed that microbial communities cultured in bioreactors were able to convert the taurocholate in bovine bile into deoxycholate. Treatment with antibiotics had no significant impacts on levels of taurocholate. Microbes encoding bile salt hydrolases, which remove taurine and glycine from primary bile salts (14), are broadly distributed amongst members of the GI microbiota (55), which may have contributed to the ability of this function to persist during antibiotic treatment.

In contrast, enzymes necessary to perform $7\square$-dehydroxylation of cholate to deoxycholate are restricted to a smaller subset of the GI microbiota, primarily members of the Clostridiales family and the *Clostridium* genus (56). Treatment with Augmentin, ceftriaxone, clindamycin, and fidaxomicin significantly increased levels of cholate at the end of antibiotic treatment on Day -2, suggesting that microbes capable of $7\square$-dehydroxylation of cholate to deoxycholate may be lost in some communities, although levels of deoxycholate were not significantly lower at these time points. This lack of significant differences in deoxycholate levels across all fecal sample communities may partly be due to differences in how different fecal communities responded to antibiotic treatment (Figure S7), as some communities (e.g., FS3 and FS9) exhibited much larger decreases in deoxycholate than others (e.g., FS6 and FS12). There was also no correlation between levels of deoxycholate and susceptibility to *C. difficile* colonization,

indicating that secondary bile salt mediated inhibition was unlikely to be a mechanism

preventing colonization in this model. This was further supported by experiments investigating

colonization resistance in the presence and absence of bile, which demonstrated bile was not

required for *C. difficile* colonization resistance.

Lack of inhibition by secondary bile salts was surprising, as levels of deoxycholate measured

in spent culture medium (431 - 608 μM) were within the range shown to inhibit *C. difficile*

growth in pure culture *in vitro* (250 - 2500 μM; (7, 15, 20)). Nevertheless, >20% of communities

colonized with *C. difficile* had deoxycholate levels $\geq$ 1000 ▢M on Day 0 of colonization,

suggesting inhibitory effects observed in pure culture may not be the same in complex culture in

the presence of other microbes. Further studies are needed to understand whether the presence of

other microbes could mitigate some of the toxic effects of deoxycholate on *C. difficile* growth.

We also observed that spores were capable of infecting communities treated with

clindamycin and this was independent of the presence of bile. Consistent with what has been

previously reported, low levels of germination were observed in the absence of bile and in spent

culture medium cultured in the presence or absence of bile. However, these low levels of

germination would result in ~10^3 vegetative cells, a level previously shown to infect antibiotic

disrupted communities cultured in bioreactors (39). This data provides further support to prior

studies that have proposed that control of *C. difficile* germination and growth (12, 25) through

restoration of microbial bile salt metabolism may not be sufficient to limit *C. difficile*

colonization.

While levels of bile salts did not correlate with susceptibility to infection, the ability of *C.*

difficile to metabolize proline as an electron acceptor for Stickland fermentation was required for

persistence in the majority of disrupted communities tested. These results are consistent with

evidence from human (10, 35) and mouse (11, 13, 25, 31, 32, 36) studies that point to Stickland

fermentation with proline as an electron acceptor as a preferred nutritional niche for *C. difficile*

in the GI tract. Stickland fermentation is a metabolic pathway limited primarily to proteolytic

Clostridial species (34). This includes microbes such as *C. scindens*, which can compete with *C.*

difficile for proline (25), dehydroxylate primary bile salts to secondary bile salts (24), and can

produce a bacteriocin that limits *C. difficile* growth (9). Identifying microbes that effectively

compete with *C. difficile* for proline and other amino acids following antibiotic treatment may be

one approach to limit *C. difficile* colonization.

It is unclear which metabolites were supporting *C. difficile* growth in antibiotic-treated

communities where *prdB* mutants persisted at higher levels over time. Stickland fermentation

with glycine or leucine as an electron acceptor is one potential niche that could have been

utilized by *prdB* mutants, although we observed the mutants in either pathway alone (Δ*grdA* and

Δ*ldhA*) was not sufficient to limit *C. difficile* colonization. Recent studies in a hamster model of

infection have shown that hamsters infected with Δ*grdAB* mutants in strain CD630 show delayed

morbidity compared to wild type CD630, although there were no significant differences between

wt and Δ*grdAB* mutants in levels of *C. difficile* or toxin recovered from the cecum at necropsy

(38), indicating the role of glycine fermentation *in vivo* may be relatively modest. Alternatively,

C. difficile may have utilized carbohydrate fermentation pathways. Further studies are needed to

more fully understand how *C. difficile* colonizes disrupted communities.

Altogether, these studies highlight the importance of improving our understanding of bile-

independent mechanisms regulating *C. difficile* colonization. While it is clear that Stickland

fermentation of amino acids with proline as an electron acceptor is important for *C. difficile*

persistence in many disrupted communities, other nutritional niches should also be explored,

including whether Stickland fermentation with both glycine and leucine may contribute to

reduced persistence. Studies similar to those described here, examining variations in fecal

microbial communities and environmental conditions, will further clarify the hierarchical

importance of different nutritional environments in limiting *C. difficile* colonization.

MATERIALS AND METHODS

Strains used in this study. *C. difficile* strain 2015 (39), a fully-sequenced ribotype 027 isolate

resistant to rifampicin and erythromycin (Accession: CP073752.1), was used for routine studies

to assess *C. difficile* colonization. To test the role of reductive fermentation of proline in

colonization, a previously described ClosTron mutant in proline reductase (*prdB*::ClosTron;

(43)) along with its congenic wild type strain were tested. Although originally reported as

mutants in the R20291 background, these strains are in the CD196 background (E. Skaar,

personal communication). To test the role of reductive fermentation of glycine and leucine,

deletions of the coding sequences of glycine reductase (*grdA*) and 2-hydroxyisocaproate

dehydrogenase (*ldhA*) were generated by riboswitch-mediated allelic exchange in strain R20291

(57). Successful deletion of genes and absence of second site mutations was verified by Illumina

sequencing. Specifically, short read, whole-genome Illumina sequencing was performed by SeqCoast

Genomics (Portsmouth, NH USA). Sequencing reads for mutant starins were both compared to the

parental *C. difficile* R20291 strain by Geneious Prime's Geneious alignment algorithm. Sequence variants

were then analyzed and both strains were found to be free of mutations outside of the expected gene

deletions. Sequences were deposited to NCBI SRA with accession numbers: SRR31632038 and

SRR31632038. These mutants were tested along with their congenic wild-type R20291 strain.

Fecal sample collection and preparation. Fecal samples were collected from 18 healthy individuals who had not been treated with oral antibiotics within the previous six months. Samples were collected from children aged 4-17 (n=3), adults aged 18-65 (n=12), and older adults aged >65 (n=2). Similar numbers of male and female participants agreed to provide samples. However, only donor age categorization (child, adult, older adult) remained linked to specific fecal samples following de-identification. Studies were designed to collect samples across the age span rather than specifically powered to evaluate differences in community composition by age. All adult participants provided consent to participate in the study. Children provided assent to participate along with parental consent to participate. Protocols for collection and use of fecal samples were reviewed and approved by Institutional Review Boards at Baylor College of Medicine (protocol number H-38014) and University of Nebraska-Lincoln (protocol numbers 18585 and 20186).

Samples were self-collected by participants in commode specimen collection containers, sealed in a plastic bag containing a gas pack (BBL GasPak Anaerobe sachet), packed in ice packs, placed in a sealed container, and returned to the lab within 24 hours as previously described (39). Fecal samples were manually homogenized and subdivided under anaerobic conditions (Anaerobic Chamber with 5% H_2, 5%CO_2, 90% N_2 atmosphere) and frozen at -80°C until later resuspended at 25% w/v in reduced phosphate buffered saline (PBS). Fecal suspensions were vortex-mixed for 5 min at $\geq$ 2500 rpm, centrifuged at 200 X g for 5 min to settle large particulates, and the supernatants were either used immediately or amended with 7.5% dimethylsulfoxide and preserved at -80°C until use. Previous work had shown that freezing samples at -80°C did not significantly change the composition of communities cultured in bioreactors (44).

Bioreactor experiments. MBRAs were assembled and operated under anoxic conditions (5% H_2, 5% CO_2, 90% N_2) at 37□C as described previously (39). MBRAs are strips of six independent continuous flow bioreactors that operate at a 15 mL volume and are continuously stirred with small magnetic stir bars positioned over a stir plate. Each independent reactor in the array contains three ports, one for delivery of fresh medium at a slow continuous rate, one for removal of waste, and a sample port for sampling of medium and bacteria in suspension. Bioreactor medium version 3 (BRM3;(58); Table S1), which simulates nutritional conditions of the distal colon was used for all studies, with the exception of studies testing the effect of bovine bile, when this component was excluded from BRM3. Fecal suspensions from individual donors were inoculated into sterile, anoxic BRM3 medium at 1% w/v final concentration and allowed to grow for 16-24 hr in batch, prior to initiation of continuous flow of medium at 1.875 mL/hr (16 hr retention time for 15 mL bioreactors). Communities were allowed to equilibrate under continuous flow for 1 or 6 days as indicated in figures. Antibiotics were either administered twice daily to individual bioreactors or added directly to source medium as indicated in figures. Research grade antibiotics were obtained from the following sources: Augmentin (5:1 amoxicillin sodium salt to potassium clavulanate; Research Products International); azithromycin dihydrate (Thermo Scientific Chemicals); cefaclor (Thermo Scientific Chemicals); ceftriaxone sodium salt hemiheptahydrate (Thermo Scientific Chemicals); clindamycin phosphate (TCI America); and fidaxomicin (Apexbio Technology, LLC). Antibiotic concentrations used for twice daily dosing were estimated based upon previously published measurements in human feces and/or bile (59–65), although fidaxomicin dosing was limited by its maximal solubility. Clindamycin concentrations used in media were previously described (42). One or two days

following cessation of antibiotics as indicated in figure legends, *C. difficile* spores or vegetative

cells were administered to reactors at ~ 1 X 10^5 CFU/mL (vegetative cells) or 1 X 10^6 CFU/mL

(spores) and levels were enumerated over time. Samples were collected from bioreactors at the

time points indicated using a sterile needle and syringe to collect bacteria in suspension. When

appropriate, an aliquot of this sample was used to enumerate *C. difficile* by serial dilution and

plating on taurocholate cefoxitin cycloserine fructose agar (TCCFA (39), for CD196 and CD196

prdB::CT), TCCFA supplemented with 50 µg/mL rifampicin and 20 µg/mL erythromycin (for

CD2015), or TCCFA supplemented with 25 µg/mL kanamycin (for R20291, R20291 Δ*grdA*, and

R20291 Δ*ldhA*). Sample were centrifuged at ~3000 X *g* for 5 min to pellet cells. Supernatants

were removed, and pellets and supernatants were stored at $\leq$ -20°C.

Microbial community analysis by 16S rRNA gene sequencing. DNA was extracted from cell

pellets using the BioSprint 96 One-For-All Vet processing kit (Qiagen) according to instructions

with the following modifications. Prior to extraction, cells were resuspended in Buffer ASL

(Qiagen) and added to sterile deep well plates (Axygen) containing 0.1 mm Zirconia beads

(BioSpec Products). Cells were disrupted by bead beating for 2 minutes at 1800 rpm on a

FastPrep 96 homogenizer (MP Biomedicals). DNA was amplified in duplicate with Phusion

polymerase using barcoded primers 515F and 806R that target the 16S rRNA gene as previously

described (41, 44), then sequenced on an Illumina MiSeq using 2 x 250 kits according to

manufacturer's protocol. All sample processing and sequencing was performed by the

investigators at the University of Nebraska-Lincoln using equipment shared by members of the

Nebraska Food for Health Center. Fastqs were processed by mothur 1.41.3, removing chimeras

identified by uchime, mapping sequences against Silva release 132, and clustering OTUs at 99%

identity using the OptiClust algorithm (66–68). Mothur 1.48.1 was used to rarefy samples to

6944 sequences, to calculate alpha (observed OTUs, Inverse Simpson) and beta diversity

(Jaccard and Bray-Curtis dissimilarity) metrics on rarefied data. Code for sequence processing

can be found in Supplementary File 3; Excel worksheets for processing of beta diversity data can

be found in Supplementary File 4. An OTU table of rarefied data can be found in Table S2 and

compiled data from sequencing analysis can be found in Table S3.

Bile salt measurements by liquid chromatography-tandem mass spectrometry (LC-

MS/MS). Quantitative analysis of the bile salt levels contained in conditioned-BRM3 culture

media sample supernatants was performed by the Texas Children's Microbiome Center

Metabolomics and Proteomics Mass Spectrometry Laboratory (TCMC-MPMSL) using

previously published methods(69, 70). Briefly, conditioned-BRM3 growth media samples were

removed from the fecal communities cultured in the individual bioreactors using sterile needles

and syringes. Samples were added into sterile tubes, and the bacterial cells were pelleted by

centrifugation at 3,000 X g for 5 min. Clarified supernatants were then sterile filtered using 96-

well 0.2 μM polyvinyldifluoride (PVDF) filter plates by centrifugation at 200 X g for 5 min. The

96-well, deep well plates used to capture sample filtrates were covered with silicone cap mats

and the capped samples were wrapped in parafilm to ensure the fixture of the capmats during

shipment. The cell-free conditioned media samples were stored frozen at -80°C pending

shipment to the TCMC-MPMSL. The cell-free conditioned media samples were shipped frozen

on dry ice, and upon receipt by the TCMC-MPMSL, were stored at -80°C until analysis.

All cell-free conditioned media samples were thawed at ambient temperature on a

laboratory benchtop. Once thawed, the entire sample volume was transferred into individual 0.6

or 1.5 mL Eppendorf tubes, and all samples were vortex-mixed using a multi-tube vortexer.

Preliminary work indicated that 10-fold and 1,000-fold dilutions were suitable to measure the

taurocholic acid (TCA) content, and the cholic acid (CA) and deoxycholic acid (DCA) content,

respectively, in each of the cell-free conditioned media samples using a common linear dynamic

range of 0.977-1,000 ng/mL for CA, DCA, and TCA – the two different dilution procedures for

each metabolite were described previously (70). These sample dilution steps were performed

using a working internal standard (WIS) solution prepared in 1:1 methanol: water that contained

250 ng/mL each of D4-CA and D4-DCA as described (70). A 5 μL volume of sample was

injected onto the SCIEX QTRAP 6500-based LC-MS/MS system, and bile acid concentrations in

the conditioned media samples were back calculated using the regression parameters of the

standard curve as described previously (70). Sample data was filtered to remove any with total

concentrations of cholate, deoxycholate and taurcholate that were less than 1 microgram/mL.

The median level of total bile salts measured was 246 ug/mL; filtering removed 18 of 336

samples. All bile salt data, including the 18 samples that were filtered, can be found in Table S4.

While the levels of TCA, CA and DCA in fresh bioreactor medium was measured at the time

cultured samples were tested (69, 70), concentrations were not reported and were lost to follow up.

Therefore, we performed a broad screen of the bile acids/salt content of two independently

prepared fresh preparations of BRM3-based bioreactor medium (with bovine bile) as well as a

batch of BRM3 prepared without bile using another previously reported targeted metabolomics

method that is capable of quantitatively measuring 16 bile acids/salts (71). This expanded

method can quantify the bile acid/salt content of microbial media samples: CA, glycocholic acid

(GCA), TCA, beta-muricholic acid (β-MCA), DCA, glycodeoxycholic acid (GDCA),

taurodeoxycholic acid (TDCA), chenodeoxycholic acid (CDCA), glycochenodeoxycholic acid

(GCDCA), taurochenodeoxycholic acid (TCDCA), ursodeoxycholic acid (UDCA),

glycoursodeoxycholic acid (GUDCA), tauroursodeoxycholic acid (TUDCA), lithocholic acid

(LCA), glycolithocholic acid (GLCA), and taurolithocholic acid (TLCA). Briefly, samples were

prepared by diluting a 10 µL volume of a thawed cell-free, conditioned BRM3 media sample in a

90 µL volume of a WIS solution that contains deuterated analogs of each of the analytes listed

above at concentrations of 250 nM for each prepared in 1:1 methanol: water (71). The samples

were prepared directly in glass autosampler vials and the samples were vortex-mixed briefly

prior to injecting a 10 µL volume onto the SCIEX QTRAP 6500-based LC-MS/MS system. Bile

acid/salt concentrations in the fresh BRM3 medium samples were back calculated using the

linear regression parameters of the standard curve as described previously (71). All samples were

tested in triplicate. Bile salt data for media can be found in Table S5. These studies confirmed

the absence of major forms of bile acids/salts in BRM3 medium made without bile (Table S5).

Data visualization and statistical analysis. The implementation of the LEfSe (47) algorithim

in mothur 1.48.1 was used to identify potential OTUs correlated with *C. difficile* colonization

from rarefied data as described in Supplementary File 3. Abundance data for OTUs correlated

with *C. difficile* colonization was manually curated from OTU abundance tables in Excel and

visualized in GraphPad Prism version 10.2.3. R studio version 2022.07.0+548 running R version

4.2.3 was used to generate the heatmap in Figure S2. Code and dependent software versions used

for data analysis can be found in Supplementary File 3. The remaining statistical analyses and

visualization was performed using GraphPad Prism. Unless otherwise noted (Figure 5; Figure

S8), significance of differences at a single time point between two treatment groups was

determined with a Mann-Whitney test and between more than two groups was determined with

Kruskal-Wallis testing with Dunn's correction for multiple comparisons. In Figure 5 and Figure

S8, parametric statistics were used to determine significance due to the low number of replicates

in one or more samples tested. One-way ANOVA with Brown-Forsythe and Welch correction

for unequal variances and Dunnett T3 correction for multiple comparisons was used in Figure 5,

whereas a two-tailed student's t-test with Welch's correction for unequal variances was used in

Figure S8. For analysis of repeated measures shown in Figures 1C, 1D, Figure S1E and Figure

S1F, a mixed effects model was used to infer statistical significance with time and reactor as the

fixed effects and treatment as the random effect. For CFU/mL data, values below the limit of

detection of the assay (333 CFU/mL), were reported as "300" to facilitate plotting.

ACKNOWLEDGEMENTS

We thank Robert Britton (Baylor College of Medicine) for access to data collected in his

laboratory by J.M.A that contributed to Figure 4B and analysis to bile salt levels in medium. We

thank former University of Nebraska-Lincoln undergraduate students Keegan Schuchart, Yining

He, and Austin Johnson for technical assistance. We thank Eric Skaar for providing CD196 and

CD196 *prdB*::CT mutants.

J.M.A., X.H., and A.E.J. were responsible for conceptualization and design of the studies.

Mutants in *grdA* and *ldhA* were generated and validated by J.N.B. and J.A.S. Data collection

was performed by X.H., A.E.J., T.A.A., A.I.L, T.V.T.D., H.C.M., S.J.H., T.D.H., and J.M.A.

Analysis was performed by X.H., A.E.J., T.A.A, S.J.H., K.M.H., T.D.H., A.M.H. and J.M.A.

J.M.A. and A.M.H acquired funding for the studies. The initial draft of the manuscript was

written by A.E.J. and J.M.A., and all authors revised, edited, and approved the final manuscript .

The Texas Children's Hospital Department of Pathology provides salary support to the Texas

Children's Microbiome Center-Metabolomics and Proteomics Mass Spectrometry Laboratory

staff, and purchased all of the reagents, the consumables and durable supplies, and the LC–

MS/MS equipment described. Sequence analysis was completed utilizing the Holland

Computing Center, which receives support from the UNL Office of Research and Economic

Development and the Nebraska Research Initiative.

DATA AVAILABILITY

16S rRNA gene data has been deposited in NCBI's Sequence Read Archive (SRA) under

BioProject ID PRJNA729569. Sequence data from *grdA* and *ldhA* were deposited to NCBI SRA

with accession numbers: SRR31632038 and SRR31632038.

ETHICS APPROVAL

Protocols for collection and use of fecal samples were reviewed and approved by Institutional

Review Boards at Baylor College of Medicine (protocol number H-38014) and University of

Nebraska-Lincoln (protocol numbers 18585 and 20186).

FUNDING

This work was partially supported by funding from the Centers for Disease Control awards 200-

2017-96080 and 75D301-18C-02909 (J.M.A. and A.M.H.), the National Institute of General

Medical Sciences awards P20GM103420 and P20GM113126 (J.M.A. received seed funding

through the Nebraska Prevention for Obesity Diseases and the Nebraska Center for Integrated

Biomolecular Communication, respectively), the Nebraska Tobacco Settlement Biomedical

Research Development Fund (J.M.A.) and the Agricultural Research Division at the University

of Nebraska-Lincoln (J.M.A.). This project was partially supported by award 5R01AI172043 to

J.A.S. from the National Institute of Allergy and Infectious Diseases. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the NIAID.

The funders had no role in study design, data collection and interpretation, or the decision to

submit the work for publication. The Texas Children's Microbiome Center – Metabolomics &

Proteomics Mass Spectrometry Laboratory (TCMC-MPMSL) is supported by an NIH S10

Shared Instrumentation Grant (S10ODO36416; T.D.H.).

CONFLICTS OF INTEREST

J.M.A. and T.A.A. have a significant financial interest in Synbiotic Health. T.D.H. is a member

of the Editorial Advisory board and is contracted as an Associate Editor for a Cell Press Journal

called STAR Protocols, and is as a member of the SCIEX Global Thought Leader in Mass

Spectrometry Program. There are no other conflicts of interest to declare.

REFERENCES

1. Guh AY, Mu Y, Winston LG, Johnston H, Olson D, Farley MM, Wilson LE, Holzbauer

SM, Phipps EC, Dumyati GK, Beldavs ZG, Kainer MA, Karlsson M, Gerding DN,

McDonald LC. 2020. Trends in U.S. Burden of Clostridioides difficile infection and

outcomes. N Engl J Med 382:1320–1330.

2. Magill SS, O'Leary E, Janelle SJ, Thompson DL, Dumyati G, Nadle J, Wilson LE, Kainer

MA, Lynfield R, Greissman S, Ray SM, Beldavs Z, Gross C, Bamberg W, Sievers M,

Concannon C, Buhr N, Warnke L, Maloney M, Ocampo V, Brooks J, Oyewumi T, Sharmin

S, Richards K, Rainbow J, Samper M, Hancock EB, Leaptrot D, Scalise E, Badrun F,

Phelps R, Edwards JR. 2018. Changes in prevalence of health care–associated infections in

U.S. hospitals. N Engl J Med 379:1732–1744.

3. Patangia DV, Anthony Ryan C, Dempsey E, Paul Ross R, Stanton C. 2022. Impact of

antibiotics on the human microbiome and consequences for host health. Microbiologyopen

11:e1260.

4. Baggs J, Jernigan JA, Halpin AL, Epstein L, Hatfield KM, McDonald LC. 2017. Risk of

subsequent sepsis within 90 days after a hospital stay by type of antibiotic exposure. Clin

Infect Dis 66:1004–1012.

5. Vardakas KZ, Trigkidis KK, Boukouvala E, Falagas ME. 2016. Clostridium difficile

infection following systemic antibiotic administration in randomised controlled trials: a

systematic review and meta-analysis. Int J Antimicrob Agents 48:1–10.

6.   Zhang Y, Limaye PB, Renaud HJ, Klaassen CD. 2014. Effect of various antibiotics on

modulation of intestinal microbiota and bile acid profile in mice. Toxicol Appl Pharmacol

277:138–145.

7.   Theriot CM, Bowman AA, Young VB. 2016. Antibiotic-induced alterations of the gut

microbiota alter secondary bile acid production and allow for Clostridium difficile spore

germination and outgrowth in the large intestine. mSphere 1:e00045-15.

8.   Seekatz AM, Theriot CM, Rao K, Chang Y-M, Freeman AE, Kao JY, Young VB. 2018.

Restoration of short chain fatty acid and bile acid metabolism following fecal microbiota

transplantation in patients with recurrent Clostridium difficile infection. Anaerobe 53:64–

73.

9.   Kang JD, Myers CJ, Harris SC, Kakiyama G, Lee I-K, Yun B-S, Matsuzaki K, Furukawa

M, Min H-K, Bajaj JS, Zhou H, Hylemon PB. 2019. Bile Acid 7α-Dehydroxylating Gut

Bacteria Secrete Antibiotics that Inhibit Clostridium difficile: Role of Secondary Bile

Acids. Cell Chem Biol 26:27-34.e4.

10. Battaglioli EJ, Hale VL, Chen J, Jeraldo P, Ruiz-Mojica C, Schmidt BA, Rekdal VM, Till

LM, Huq L, Smits SA, Moor WJ, Jones-Hall Y, Smyrk T, Khanna S, Pardi DS, Grover M,

Patel R, Chia N, Nelson H, Sonnenburg JL, Farrugia G, Kashyap PC. 2018. Clostridioides

difficile uses amino acids associated with gut microbial dysbiosis in a subset of patients

with diarrhea. Sci Transl Med 10:eaam7019.

11.   Fletcher JR, Erwin S, Lanzas C, Theriot CM. 2018. Shifts in the gut metabolome and

699   Clostridium difficile transcriptome throughout colonization and infection in a mouse model.

700   mSphere 3:e00089-18.

701   12.   Bustin KA, Abbas A, Wang X, Abt MC, Zackular JP, Matthews ML. 2023. Characterizing

702   metabolic drivers of Clostridioides difficile infection with activity-based hydrazine probes.

703   Front Pharmacol 14:1074619.

704   13.   Girinathan BP, DiBenedetto N, Worley JN, Peltier J, Arrieta-Ortiz ML, Immanuel SRC,

705   Lavin R, Delaney ML, Cummins CK, Hoffman M, Luo Y, Gonzalez-Escalona N, Allard M,

706   Onderdonk AB, Gerber GK, Sonenshein AL, Baliga NS, Dupuy B, Bry L. 2021. In vivo

707   commensal control of Clostridioides difficile virulence. 11. Cell Host Microbe 29:1693-

708   1708.e7.

709   14.   Ridlon JM, Kang D-J, Hylemon PB. 2006. Bile salt biotransformations by human intestinal

710   bacteria. J Lipid Res 47:241–259.

711   15.   Sorg JA, Sonenshein AL. 2008. Bile salts and glycine as cogerminants for Clostridium

712   difficile spores. J Bacteriol 190:2505–2512.

713   16.   Sorg JA, Sonenshein AL. 2009. Chenodeoxycholate Is an inhibitor of Clostridium difficile

714   spore germination. J Bacteriol 191:1115–1117.

715   17.   Bhattacharjee D, Francis MB, Ding X, McAllister KN, Shrestha R, Sorg JA. 2016.

716   Reexamining the Germination Phenotypes of Several Clostridium difficile Strains Suggests

717   Another Role for the CspC Germinant Receptor. J Bacteriol 198:777–786.

18. Shrestha R, Sorg JA. 2018. Hierarchical recognition of amino acid co-germinants during

Clostridioides difficile spore germination. Anaerobe 49:41–47.

19. Kochan TJ, Shoshiev MS, Hastie JL, Somers MJ, Plotnick YM, Gutierrez-Munoz DF, Foss

ED, Schubert AM, Smith AD, Zimmerman SK, Carlson PE, Hanna PC. 2018. Germinant

synergy facilitates Clostridium difficile spore germination under physiological conditions.

mSphere 3:e00335-18.

20. Usui Y, Ayibieke A, Kamiichi Y, Okugawa S, Moriya K, Tohda S, Saito R. 2020. Impact

of deoxycholate on Clostridioides difficile growth, toxin production, and sporulation.

Heliyon 6:e03717.

21. Allegretti JR, Kearney S, Li N, Bogart E, Bullock K, Gerber GK, Bry L, Clish CB, Alm E,

Korzenik JR. 2016. Recurrent Clostridium difficile infection associates with distinct bile

acid and microbiome profiles. Aliment Pharmacol Ther 43:1142–1153.

22. Weingarden AR, Chen C, Bobr A, Yao D, Lu Y, Nelson VM, Sadowsky MJ, Khoruts A.

2014. Microbiota transplantation restores normal fecal bile acid composition in recurrent

Clostridium difficile infection. Am J Physiol Gastrointest Liver Physiol 306:G310–G319.

23. Theriot CM, Koenigsknecht MJ, Carlson PE, Hatton GE, Nelson AM, Li B, Huffnagle GB,

Z Li J, Young VB. 2014. Antibiotic-induced shifts in the mouse gut microbiome and

metabolome increase susceptibility to Clostridium difficile infection. Nat Commun 5:3114.

24. Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H,

Kinnebrew M, Viale A, Littmann E, van den Brink MRM, Jenq RR, Taur Y, Sander C,

Cross JR, Toussaint NC, Xavier JB, Pamer EG. 2015. Precision microbiome reconstitution

restores bile acid mediated resistance to Clostridium difficile. Nature 517:205–208.

25. Aguirre AM, Yalcinkaya N, Wu Q, Swennes A, Tessier ME, Roberts P, Miyajima F,

Savidge T, Sorg JA. 2021. Bile acid-independent protection against Clostridioides difficile

infection. 10. PLoS Pathog 17:e1010015.

26. Saenz C, Fang Q, Gnanasekaran T, Trammell SAJ, Buijink JA, Pisano P, Wierer M, Moens

F, Lengger B, Brejnrod A, Arumugam M. 2023. Clostridium scindens secretome suppresses

virulence gene expression of Clostridioides difficile in a bile acid-independent manner.

Microbiol Spectr 11:e03933-22.

27. Sievers S, Metzendorf NG, Dittmann S, Troitzsch D, Gast V, Tröger SM, Wolff C, Zühlke

D, Hirschfeld C, Schlüter R, Riedel K. 2019. Differential view on the bile acid stress

response of Clostridioides difficile. Front Microbiol 10:258.

28. Wilson KH, Freter R. 1986. Interaction of Clostridium difficile and Escherichia coli with

microfloras in continuous-flow cultures and gnotobiotic mice. Infect Immun 54:354–358.

29. Wilson KH, Perini F. 1988. Role of competition for nutrients in suppression of Clostridium

difficile by the colonic microflora. Infect Immun 56:2610–2614.

30. Engevik MA, Engevik AC, Engevik KA, Auchtung JM, Chang-Graham AL, Ruan W, Luna

RA, Hyser JM, Spinler JK, Versalovic J. 2021. Mucin-degrading microbes release

monosaccharides that chemoattract Clostridioides difficile and facilitate colonization of the

human intestinal mucus layer. ACS Infect Dis 7:1126–1142.

31.   Janoir C, Deneve C, Bouttier S, Barbut F, Hoys S, Caleechum L, Chapeton-Montes D,

759   Pereira FC, Henriques AO, Collignon A, Monot M, Dupuy B. 2013. Adaptive strategies

760   and pathogenesis of Clostridium difficile from in vivo transcriptomics. Infect Immun

761   81:3757–3769.

762   32.   Jenior ML, Leslie JL, Young VB, Schloss PD. 2017. Clostridium difficile colonizes

763   alternative nutrient niches during infection across distinct murine gut microbiomes.

764   mSystems 2:e00063-17.

765   33.   Bouillaut L, Self WT, Sonenshein AL. 2013. Proline-dependent regulation of Clostridium

766   difficile Stickland metabolism. 4. J Bacteriol 195:844–854.

767   34.   Pavao A, Graham M, Arrieta-Ortiz ML, Immanuel SRC, Baliga NS, Bry L. 2022.

768   Reconsidering the *in vivo* functions of *Clostridial* Stickland amino acid fermentations.

769   Anaerobe 76:102600.

770   35.   Robinson JI, Weir WH, Crowley JR, Hink T, Reske KA, Kwon JH, Burnham C-AD,

771   Dubberke ER, Mucha PJ, Henderson JP. 2019. Metabolomic networks connect host-

772   microbiome processes to human Clostridioides difficile infections. J Clin Invest 129:3792–

773   3806.

774   36.   Fletcher JR, Pike CM, Parsons RJ, Rivera AJ, Foley MH, McLaren MR, Montgomery SA,

775   Theriot CM. 2021. Clostridioides difficile exploits toxin-mediated inflammation to alter the

776   host nutritional landscape and exclude competitors from the gut microbiota. Nat Commun

777   12:462.

37.   Kim J, Darley D, Selmer T, Buckel W. 2006. Characterization of (R)-2-Hydroxyisocaproate

Dehydrogenase and a Family III Coenzyme A Transferase Involved in Reduction of l-

Leucine to Isocaproate by Clostridium difficile. Applied and Environmental Microbiology

72:6062–6069.

38.   Rizvi A, Vargas-Cuebas G, Edwards AN, DiCandia MA, Carter ZA, Lee CD, Monteiro

MP, McBride SM. 2023. Glycine fermentation by C. difficile promotes virulence and spore

formation, and is induced by host cathelicidin. Infection and Immunity 91:e00319-23.

39.   Robinson CD, Auchtung JM, Collins J, Britton RA. 2014. Epidemic Clostridium difficile

strains demonstrate increased competitive fitness compared to nonepidemic isolates. Infect

Immun 82:2815–2825.

40.   Spinler JK, Auchtung J, Brown A, Boonma P, Oezguen N, Ross CL, Luna RA, Runge J,

Versalovic J, Peniche A, Dann SM, Britton RA, Haag A, Savidge TC. 2017. Next-

generation probiotics targeting Clostridium difficile through precursor-directed

antimicrobial biosynthesis. Infect Immun 85:e00303-17.

41.   Auchtung JM, Preisner EC, Collins J, Lerma AI, Britton RA. 2020. Identification of

simplified microbial communities that inhibit Clostridioides difficile infection through

dilution/extinction. mSphere 5:e00387-20.

42.   Mahnic A, Auchtung JM, Poklar Ulrih N, Britton RA, Rupnik M. 2020. Microbiota in vitro

modulated with polyphenols shows decreased colonization resistance against Clostridioides

difficile but can neutralize cytotoxicity. 1. Sci Rep 10:8358.

43. Lopez CA, Beavers WN, Weiss A, Knippel RJ, Zackular JP, Chazin W, Skaar EP. 2019.

The immune protein Calprotectin impacts Clostridioides difficile metabolism through zinc

limitation. 6. mBio 10:e02289-19.

44. Auchtung JM, Robinson CD, Britton RA. 2015. Cultivation of stable, reproducible

microbial communities from different fecal donors using minibioreactor arrays (MBRAs).

Microbiome 3:42.

45. Aluthge ND, Tom WA, Bartenslager AC, Burkey TE, Miller PS, Heath KD, Kreikemeier-

Bower C, Kittana H, Schmaltz RJ, Ramer-Tait AE, Fernando SC. 2020. Differential

longitudinal establishment of human fecal bacterial communities in germ-free porcine and

murine models. Commun Biol 3:760.

46. Collins J, Auchtung JM, Schaefer L, Eaton KA, Britton RA. 2015. Humanized microbiota

mice as a model of recurrent Clostridium difficile disease. Microbiome 3:35.

47. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011.

Metagenomic biomarker discovery and explanation. Genome Biol 12:R60.

48. Collins J, Robinson C, Danhof H, Knetsch CW, van Leeuwen HC, Lawley TD, Auchtung

JM, Britton RA. 2018. Dietary trehalose enhances virulence of epidemic Clostridium

difficile. Nature 553:291–294.

49. Wilson KH. 1983. Efficiency of various bile salt preparations for stimulation of Clostridium

difficile spore germination. J Clin Microbiol 18:1017–1019.

50. Francis MB, Allen CA, Shrestha R, Sorg JA. 2013. Bile acid recognition by the Clostridium

difficile germinant receptor, CspC, is important for establishing infection. PLoS Pathog

9:e1003356.

51. Stabler RA, He M, Dawson L, Martin M, Valiente E, Corton C, Lawley TD, Sebaihia M,

Quail MA, Rose G, Gerding DN, Gibert M, Popoff MR, Parkhill J, Dougan G, Wren BW.

2009. Comparative genome and phenotypic analysis of Clostridium difficile 027 strains

provides insight into the evolution of a hypervirulent bacterium. Genome Biol 10:R102.

52. Zhang J, Chen L, Gomez-Simmonds A, Yin MT, Freedberg DE. 2022. Antibiotic-specific

risk for community-acquired Clostridioides difficile infection in the United States from

2008 to 2020. Antimicrob Agents Chemother 66:e01129-22.

53. Brown KA, Khanafer N, Daneman N, Fisman DN. 2013. Meta-analysis of antibiotics and

the risk of community-associated Clostridium difficile infection. Antimicrob Agents

Chemother 57:2326–2332.

54. Reeves AE, Koenigsknecht MJ, Bergin IL, Young VB. 2012. Suppression of Clostridium

difficile in the Gastrointestinal Tracts of Germfree Mice Inoculated with a Murine Isolate

from the Family Lachnospiraceae. Infect Immun 80:3786–3794.

55. Song Z, Cai Y, Lao X, Wang X, Lin X, Cui Y, Kalavagunta PK, Liao J, Jin L, Shang J, Li

J. 2019. Taxonomic profiling and populational patterns of bacterial bile salt hydrolase

(BSH) genes based on worldwide human gut microbiome. Microbiome 7:9.

56.   Kim KH, Park D, Jia B, Baek JH, Hahn Y, Jeon CO. 2022. Identification and

837   characterization of major bile acid 7α-dehydroxylating bacteria in the human gut.

838   mSystems 7:e00455-22.

839   57.   Brehm JN, Sorg JA. 2024. Theophylline-based control of repA on a Clostridioides difficile

840   plasmid for use in allelic exchange. Anaerobe 88:102858.

841   58.   Auchtung TA, Fofanova TY, Stewart CJ, Nash AK, Wong MC, Gesell JR, Auchtung JM,

842   Ajami NJ, Petrosino JF. 2018. Investigating Colonization of the Healthy Adult

843   Gastrointestinal Tract by Fungi. mSphere 3:e00092-18.

844   59.   Neu HC. 1974. Antimicrobial activity and human pharmacology of amoxicillin. J Infect Dis

845   129:S123–S131.

846   60.   Owen AW, Faragher EB. 1986. Biliary pharmacokinetics of ticarcillin and clavulanic acid.

847   J Antimicrob Chemother 17:65–70.

848   61.   Brogard J-M, Pinget M, Comte F, Adloff M, Lavillaureix J. 2009. Biliary excretion of

849   cefaclor: experimental and clinical study. Chemotherapy 28:189–199.

850   62.   Pletz MWR, Rau M, Bulitta J, De Roux A, Burkhardt O, Kruse G, Kurowski M, Nord CE,

851   Lode H. 2004. Ertapenem pharmacokinetics and impact on intestinal microflora, in

852   comparison to those of ceftriaxone, after multiple dosing in male and female volunteers.

853   Antimicrob Agents Chemother 48:3765–3772.

63.   Kager L, Liljeqvist L, Malmborg AS, Nord CE. 1981. Effect of clindamycin prophylaxis on

the colonic microflora in patients undergoing colorectal surgery. Antimicrob Agents

Chemother 20:736–740.

64.   Steinbakk M, Lingaas E, Carlstedt-Duke B, Høverstad T, Midtvedt A-C, Norin KE,

Midtvedt T. 1992. Faecal concentration of ten antibiotics and influence on some microflora-

associated characteristics (MACs). Microb Ecol Health Dis 5:269–276.

65.   Sears P, Crook DW, Louie TJ, Miller MA, Weiss K. 2012. Fidaxomicin attains high fecal

concentrations with minimal plasma concentrations following oral administration in

patients with Clostridium difficile infection. Clin Infect Dis 55:S116-120.

66.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,

Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-

supported software for describing and comparing microbial communities. Appl Environ

Microbiol 75:7537–7541.

67.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data

on the MiSeq Illumina sequencing platform. Appl Environ Microbiol 79:5112–5120.

68.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Res 41:D590–D596.

69.   Horvath TD, Haidacher SJ, Oezguen N, Hoch KM, Auchtung JM, Haag AM. 2020.

875   Ruggedness testing of liquid chromatography-tandem mass spectrometry system

876   components using microbiome-relevant methods and matrices. J Microbiol Methods

877   177:106020.

878   70.   Horvath TD, Haidacher SJ, Hoch KM, Auchtung JM, Haag AM. 2020. A high-throughput

879   LC-MS/MS method for the measurement of the bile acid/salt content in microbiome-

880   derived sample sets. MethodsX 7:100951.

881   71.   Engevik MA, Thapa S, Lillie IM, Yacyshyn MB, Yacyshyn B, Percy AJ, Chace D, Horvath

882   TD. 2024. Repurposing dried blood spot device technology to examine bile acid profiles in

883   human dried fecal spot samples. Am J Physiol Gastrointest Liver Physiol 326:G95–G106.

 FIGURE LEGENDS

Figure 1: **Alterations in microbiota diversity varied by type of antibiotic administered.** (**A**)

Antibiotics administered during experiment. (**B**) Experimental timeline indicating points where

samples were collected; time is indicated relative to the point of *C. difficile* challenge (Day 0). In

(**C**)-(**F**), samples were collected from duplicate reactors inoculated with one of twelve healthy

human fecal samples and treated as indicated. (**C**) Microbiota richness (Observed OTUs with ≥

99% identity) and (**D**) microbial diversity (Inverse Simpson measure) were determined from 16S

rRNA gene sequences. Statistical significance of changes between time points (Days -7, -2, and

0) are indicated for each antibiotic. (**E**) Jaccard and (**F**) Bray-Curtis dissimilarities were

determined from samples collected on Day -7 and plotted for each reactor and the fecal inocula.

Statistical significance of differences between each antibiotic-treated sample and untreated

communities at each time point are indicated. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****,

$p<0.0001$. All data is plotted, with boxes representing the interquartile range (IQR) and cross

lines representing the median values.

Figure 2: Clindamycin-treatment increases susceptibility to *C. difficile* colonization. (A)

Levels of *C. difficile* measured in all reactors on Day 7 of infection. Data is pooled by treatment,

with each reactor and replicate indicated by a distinct symbol consistent across treatment groups.

(Effects on individual fecal samples can be better visualized in Supplementary Figure S3.)

Statistical significance of differences between each antibiotic-treated community and untreated

communities are indicated; ***, $p<0.001$. **(B)** – **(E)** Simple linear regression between levels of *C.*

difficile on Day 7 and Jaccard (**B**-**C**) and Bray-Curtis (**D**-**E**) dissimilarity on Day -2 (**B,D**) or Day

0 (**C,E**). (**F**) OTUs that differed significantly in abundance on Day -2 and/or Day 0 in

communities that were colonized or not colonized with *C. difficile* on Day 7 were identified with

LEfSe analysis of rarefied OTU abundance data. Abundance of OTUs across all samples are

plotted for colonized and not colonized communities, with linear discriminant analysis (LDA)

and p-values determined by LEfSe reported below abundance plots.

Figure 3: Cholate, deoxycholate, and taurocholate levels in untreated and antibiotic-treated

bioreactors. Levels of (**A**, **D**) cholate, (**B**, **E**) deoxycholate, and (**C**, **F**) taurocholate were

measured from the communities described in Figure 1 at the end of antibiotic treatment (Day -2;

A-**C**) and just prior to *C. difficile* challenge (Day 0; **D**-**F**). Significance of differences between

antibiotic-treated and untreated samples at each time point are reported. *, $p<0.05$; **, $p<0.01$;

***, $p<0.001$; ****, $p<0.0001$. (**G**) Simple linear correlation between levels of cholate measured

on Day -2 and *C. difficile* levels on Day 7. (**H**) Levels of deoxycholate are plotted for the

communities identified as colonized and not colonized in Figure 2. (**I**) Simple linear correlation

between levels of deoxycholate measured on Day 0 and *C. difficile* levels on Day 7. Colonized

communities with deoxycholate levels greater than 1 mM are indicated in red.

Figure 4. Bile acids were not required for colonization resistance and did not enhance

colonization in communities of human fecal microbes cultured in bioreactors. (**A**)

Experimental timeline indicating points where samples were collected; time is indicated relative

to the point of *C. difficile* challenge. The final collection timepoint was on Day 5 for all samples

with the exception of FS22, which was collected on Day 6 and is represented by open symbols

(**B**) *C. difficile* levels in untreated and clindamycin-treated communities cultured in the presence

and absence of bile. Significance of differences between untreated and clindamycin-treated

samples are shown. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$. (**C**) *C. difficile* levels

in clindamycin-treated communities challenged with *C. difficile* vegetative (V) cells or spores

(S). Significance of differences between vegetative and spores are shown at each time point. (**D**)

C. difficile levels in clindamycin-treated communities cultured in the presence or absence of bile

challenged with *C. difficile* spores. No significant differences were observed between

communities cultured in the presence or absence of bile.

Figure 5. Low levels of spore germination in spent culture medium from communities

cultured in the presence of bile. Germination of CD2015 spores was determined in filter-

sterilized spent culture medium from clindamycin-treated fecal communities cultured in the

presence or absence of bile. Spores were incubated in medium for one hour followed by

enumeration on agar medium with or without added taurocholate. The percent germination on

taurocholate indicates the percent of colonies recovered after overnight incubation in the absence

of taurocholate compared to incubation in the presence of taurocholate for each sample. Fresh

bioreactor medium made with or without bovine bile were included as controls. Significant

difference observed in comparisons between all samples are reported. ns, $p > 0.05$; **, $p < 0.01$;

***, $p < 0.001$.

Figure 6. *C. difficile prdB*::CT mutants fail to persist in most clindamycin-treated

communities. (**A**) Levels of wt (*prdB*$^+$) and *prdB*::CT (*prdB*$^-$) CD196 strains were measured

over time following introduction into fecal communities treated with clindamycin as outlined in

Figure 4A. Data reported are pooled from all six fecal samples tested. Significant differences

between wild type and mutant strain at each time point are reported. ***, $p < 0.001$; ***,

$p < 0.0001$. (**B**) Levels of wt and *prdB*::CT CD196 strains measured on day 5 of colonization from

each of the six fecal samples tested (FS515, FS228, FS235, FS685, FS133, and FS583).

Significant differences between wild type and mutant strain for each fecal sample are reported. *,

p<0.05; **, p<0.01; ****, p<0.001. (C) Levels of wt, Δ*ldhA*, and Δ*grdA* R20291 strains were

measured over time following introduction into fecal communities treated with clindamycin as

outlined in Figure 4A. No significant differences between wild type and mutant strains were

observed at any time point.

SUPPLEMENTAL MATERIAL

Supplementary File 1: Figures S1 to S8

Figure Legends:

Figure S1. Changes in microbial richness and diversity across individual reactors in response

to time and treatment with antibiotics. (**A**,**C**) Microbial richness, (**B**,**D**) diversity, and (**E**,**F**)

community dissimilarity were re-plotted and re-analyzed from Figure 1C-1D to show statistically

significant differences between treatments at each time point (**A**,**B**) and to show values connected

over time across individual reactors (**C**,**D**). Note that no communities had been treated with

antibiotics at Day -7, but reactors would commence treatment after sampling. For comparisons

between fecal inocula and Day -7 samples in A and B, all differences were statistically significant

with $p<0.0001$. In (**E**) and (**F**), treatment groups are shown in the same order with abbreviated

designations for treatment groups. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$.

Figure S2. Distribution of 100 most abundant OTUs across fecal inocula and cultured fecal

communities prior to antibiotic treatment (Day -7). Log_4 OTU abundances are indicated as

shaded in fecal inocula (gray boxes) and cultured fecal communities (white boxes) for each fecal

sample as indicated. Genus-level classifications of each OTU, organized by phyla, are indicated

on the y-axis. Full taxonomy can be found in Table S2.

Figure S3. *C. difficile* levels vary across microbial communities from different fecal samples

treated with antibiotics. *C. difficile* levels measured from duplicate reactors on (**A**) Day 7 and (**B**)

Day 3 were pooled and plotted for each fecal sample (FS1-FS12) and each antibiotic tested as

indicated. Circles indicate first replicate sample, while squares indicate second replicate samples.

Day 3 data is shown for completeness, although it was not used for determining colonization.

Statistical analyses were not performed on each treatment because they were not sufficiently

powered. Data is shown to indicate trends in variation observed. Points at the lowest point visible on

the y-axis values indicate samples with *C. difficile* levels below the level of detection.

Figure S4. Correlations between levels of *C. difficile* colonization on Day 7 and changes in

microbiota composition. Simple linear regression was used to investigate relationships between *C.*

difficile colonization on Day 7 and changes in microbiota composition. (**A**-**B**) Correlation with

richness (Observed OTUs) on (**A**) Day -2 and (**B**) Day 0. (C-D) Correlation with changes in richness

from Day -7 to (**C**) Day -2 and (**D**) Day 0. (**E**-**F**) Correlation with changes in microbial diversity

(Inverse Simpson Measure) on (**E**) Day -2 and (**F**) Day 0. Goodness of fit (R^2)and significance of

deviation from zero (p-value) are reported.

Figure S5. Correlations between levels of *C. difficile* colonization on Day 3 and changes in

microbiota composition. Simple linear regression was used to investigate relationships between

colonization on Day 3 and changes in microbiota composition. (**A**-**B**) Correlation with richness

(Observed OTUs) on (**A**) Day -2 and (**B**) Day 0. (**C**-**D**) Correlation with changes in richness from

(**C**) Day -7 to Day -2 and (**D**) Day -7 to Day 0. (**E**-**F**) Correlation with changes microbial diversity

(Inverse Simpson) on (**E**) Day -2 and (**F**) Day 0. (**G**-**H**) Correlation with changes in Jaccard

dissimilarity before antibiotic treatment (Day -7) and after antibiotic treatment on (**G**) Day -2 or (**H**)

Day 0. (**I**-**J**) Correlation with changes in Bray-Curtis dissimilarity before antibiotic treatment (Day -

7) and after antibiotic treatment on (**I**) Day -2 or (**J**) Day 0. Goodness of fit (R^2)and significance of

deviation from zero (p-value) are reported.

1004

Figure S6. Bile salts in bioreactor medium and cultured bioreactor communities. (**A**) Cholate

and (**B**) chenodeoxycholate/ursodeoxycholate family bile salts were measured in two batches of

bioreactor medium. Independent batches are indicated by shape, with three technical replicates

tested. In (**A** and **B**), the column labeled "Total" indicates the sum of all (**A**) cholate or (**B**)

chenodeoxycholate/ursodeoxycholate family bile salts measured. While the levels of taurocholate,

cholate and deoxycholate in fresh bioreactor medium was measured at the time cultured samples

were tested (described in references 69 and 70), concentrations were not reported and were lost to

follow up. (**C**) Concentrations of taurocholate, cholate and deoxycholate in fresh medium and in

untreated communities on Day -2 are shown for comparison. For fresh medium, total indicates total

cholate family bile salts (including glycocholate) whereas total indicates the sum of taurocholate,

cholate and deoxycholate for spent culture medium. Although glycocholate was not measured in

spent culture medium, levels would likely be low based on levels of taurocholate measured.

1017

Figure S7. Changes in levels of cholate and deoxycholate across fecal communities. (**A-B**)

Levels of cholate and (**C-D**) deoxycholate measured from duplicate reactors on Day -2 (**A,C**) and

Day 0 (**B,D**) were pooled and plotted for each fecal sample (FS1-FS12) and each antibiotic tested as

indicated. Statistical analyses were not performed on each treatment because they were not

sufficiently powered. Data is shown to indicate trends in variation observed. Circles indicate first

replicate sample, while squares indicate second replicate samples. (**E**) Cholate and (**F**) deoxycholate

levels on Day -2 or Day 0 from all antibiotic-treated reactors from a fecal sample were pooled.

Statistical significance of bile salt levels on Day -2 were compared to FS3, which had median levels

of cholate that were >2.9 fold higher than all other samples (2.9-46 fold higher across all samples).

*, p<0.05; **, p<0.01; ***, p<0.001; ****, p<0.0001.

1028

Figure S8: *C. difficile* **spores infect clindamycin-treated microbial communities from different**

fecal samples similarly to vegetative cells. *C. difficile* levels measured from replicate reactors on

Days 0, 1, 3 and 5 or 6 are plotted for each fecal sample (FS11, FS16, FS22, FS27, and FS288)

infected with *C. difficile* vegetative cells (circles, filled bars) or spores (squares, open bars). n=3 for

all conditions except FS288 challenged with spores, which was n=2. All communities were cultured

in media containing bile. *, p<0.05, **, p<0.01, ***, p<0.001.

1035

Supplementary File 2: Tables S1 to S5

Table S1: BRM3 recipe and instructions

Table S2: OTU abundance data for sequenced samples.

Table S3: Compiled data used for Figure 1-3 and Supplementary Figures S1-S7

Table S4: Filtering of raw bile salt data

Table S5: Raw and processed bile salt data obtained from analysis of two independent batches of

BRM3

1043

1044   **Supplementary File 3: Mothur and R code for analysis of microbiota data.**

1045

1046   **Supplementary File 4: Excel worksheets for selection of beta diversity values from mothur**

1047   **sequence analysis output.**

1048

1049 **Supplementary File 5: Tables S6 to S8**

1050 **Table S6**: Huang.final.abund.opti_mcc.0.01.shared

1051 **Table S7**: Data for simple.metadata.csv

1052 **Table S8**: Huang.final.abund.opti_mcc.0.01.cons.subsample.cons.taxonomy

1053

1054

Figure 2

Figure 4

A

B

C