Title: Viral delivery of an RNA-guided genome editor for transgene-free germline editing in *Arabidopsis*

Authors: Trevor Weiss[1], Maris Kamalu[1], Honglue Shi[2,6], Zheng Li[1], Jasmine Amerasekera[1], Zhenhui Zhong[5], Benjamin A. Adler[2,11], Michelle Song[1], Kamakshi Vohra[2,11], Gabriel Wirnowski[1], Sidharth Chitkara[1], Charlie Ambrose[1], Noah Steinmetz[1], Ananya Sridharan[1], Diego Sahagun[1], Jillian F. Banfield[2,8,9,10], Jennifer A. Doudna[2,4,6,11,12,13,14], Steven. E. Jacobsen[1,7,15]

Affiliations:

[1]Department of Molecular, Cell and Developmental Biology, University of California at Los Angeles, Los Angeles, CA 90095, USA

[2]Innovative Genomics Institute, University of California, Berkeley, CA 94720 USA

[3]Department of Bioengineering, University of California, Berkeley, CA 94720 USA

[4]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

[5]Department of Biotechnology, Sichuan University, Chengdu, China

[6]Howard Hughes Medical Institute, University of California, Berkeley, CA 94720, USA.

[7]Howard Hughes Medical Institute (HHMI), University of California at Los Angeles, Los Angeles, CA 90095, USA

[8]Department of Earth and Planetary Science, University of California, Berkeley, CA 94720, USA

[9]Department of Environmental Science, Policy and Management, University of California, Berkeley, CA 94720, USA

[10]University of Melbourne, Melbourne, Australia

[11]California Institute for Quantitative Biosciences (QB3), University of California, Berkeley,

CA 94720, USA

[12]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[13]MBIB Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[14]Gladstone Institutes, University of California, San Francisco, CA 94158, USA

[15]Lead contact

*Corresponding author. Email: jacobsen@ucla.edu

Abstract

Genome editing is transforming plant biology by enabling precise DNA modifications.

However, delivery of editing systems into plants remains challenging, often requiring slow,

genotype-specific methods such as tissue culture or transformation. Plant viruses, which

naturally infect and spread to most tissues, present a promising delivery system for editing

reagents. But most viruses have limited cargo capacities, restricting their ability to carry large

CRISPR-Cas systems. Here, we engineered tobacco rattle virus (TRV) to carry the compact RNA-

guided TnpB enzyme ISYmu1 and its guide RNA. This innovation allowed transgene-free editing

of *Arabidopsis thaliana* in a single step, with edits inherited in the subsequent generation. By

overcoming traditional reagent delivery barriers, this approach offers a novel platform for

genome editing, which can greatly accelerate plant biotechnology and basic research.

Main text

Programmable RNA-guided endonucleases including CRISPR-Cas9 are driving advances

in genome editing for both fundamental research and biotechnology. The ability to genetically

modify plant genomes has allowed for the creation of rationally designed phenotypes.

However, efficient delivery of genome editing reagents to plants remains a major challenge.

The most common strategy is to encode RNA-guided genome editors (e.g. CRISPR-Cas enzymes)

within transgenes and use tissue culture and plant transformation approaches to make

transgenic plants, after which genetic crosses are required to remove the transgenic material

but retain the edits [1–3]. However, current plant transformation methods are limited to specific

plant species and genotypes, often require considerable time, resources and technical

expertise, and can cause unintended changes to the genome and epigenome [1].

One approach to circumvent these limitations is to use plant viral vectors to deliver

genome editing reagents. For example, several viral vectors have been engineered to encode

guide RNAs (gRNAs) for delivery to transgenic plants already expressing Cas9, resulting in

somatic and germline editing and transmission of edits to the next generation [4–7]. Because

plants have evolved mechanisms to restrict viral infection of meristems and germ cells, most

viruses are rarely sexually transmitted [8]. However, transient invasion of meristem cells by viral

RNAs encoding gRNAs can allow these cells to be edited, and for these edits to be seed

transmissible [4–7]. While these approaches represent significant advances, they still require the

use of transgenic plants to express the nuclease protein.

A strategy to avoid the need for transgenic plant materials has been the use of viral

vectors with large cargo capacities, capable of expressing entire editing systems (e.g. Cas9 and

the gRNA). This approach has been met with some success, however it still requires plant

regeneration steps because these viruses do not cause germline editing and heritability of the

edits [9–12]. On the other hand, encoding entire CRISPR systems in viruses that are capable of

germline transmission has not been possible because of their limited cargo capacity [4–7,13].

To overcome this cargo size limit, we explored the potential of TnpB, a class of

ultracompact RNA-guided endonucleases (~400 amino acids) [14–16], to be encoded in a plant RNA

viral vector. As ancestors of Cas enzymes, TnpBs similarly utilize a programmable RNA guide,

called an omega RNA (ωRNA), to be directed to any target site and induce genome edits.

Previously, TnpBs ISDra2, ISYmu1 and ISAam1 were shown to be capable of targeted genome

editing in mammalian cells, and ISDra2 and ISYmu1 in monocot rice plant cells [14,17–19]. Here, we

tested the ISDra2, ISYmu1 and ISAam1 TnpBs for genome editing in the dicot plant, *Arabidopsis*.

Given the single cargo site in the TRV vector that is typically used, we sought to express both

the TnpB protein and its guide RNA within the same mRNA transcript under a single promoter,

similar to their natural expression arrangement [14–16].

To test the activities of TnpB and its gRNA encoded in a single transcript, we first

expressed these three TnpBs and assessed their RNA-guided plasmid interference activities in

bacteria. We co-expressed the TnpB and gRNA from the same promoter as a single transcript,

maintaining their natural sequences without codon optimization. We compared two

configurations of the 3'-guide region: one extended continuously without a terminator to mimic

the natural TnpB condition, and another capped by the hepatitis delta virus (HDV) ribozyme, as

previously used in bacteria [14] (Extended Data Fig. 1). Our results showed that without the HDV

ribozyme, only ISDra2 demonstrated plasmid interference activity whereas with the HDV

ribozyme, all three TnpBs exhibited robust activity at both 26°C and 37°C (Fig. 1a, Extended

Data Fig. 2). These findings revealed that single transcript expression cassettes with an HDV

ribozyme sequence at the 3' end are capable of cleaving plasmid DNA in bacteria.

To test the single expression cassette for targeted genome editing in *Arabidopsis*, we

used the *AtUBQ10* promoter to drive expression of the TnpB-ωRNA and a gRNA targeting the

PHYTOENE DESATURASE3 (*AtPDS3*) gene region, followed by the HDV ribozyme and rbcS-E9

terminator (Fig. 1b). We tested twenty ISDra2 sites, ten ISYmu1 sites, and seven ISAam1 sites

for editing capabilities in *Arabidopsis* protoplast cells (Supplementary Table 1) [20]. ISDra2 and

ISYmu1 demonstrated active editing ranging from 0-4.8% and 0.1-4.2%, respectively, as

measured by next generation amplicon sequencing (amp-seq) (Extended Data Fig. 3a). ISAam1

was much less active, ranging from 0-0.3% editing efficiency (Extended Data Fig. 3a). On

average, we observed editing efficiencies of 1% for ISDra2, 2.1% for ISYmu1 and 0.1% for

ISAam1 (Fig. 1c). In line with previous reports, the DNA repair profiles consisted of deletion-

dominant repair outcomes for all three TnpBs (Extended Data Fig. 3b) [14,17,18]. These data

demonstrate that ISDra2, ISYmu1 and ISAam1 are all capable of targeted genome editing in

Arabidopsis plant cells using the single transcript expression design.

To evaluate TnpB-mediated editing in transgenic plants we selected ISYmu1, as it

demonstrated the highest average editing efficiency in *Arabidopsis* protoplast cells and was

shown to exhibit no off-target editing in rice [18]. Two gRNAs with the most active editing were

selected, each targeting a unique genomic context. gRNA2 targeted the coding region of

AtPDS3 whereas gRNA12 targeted the promoter region directly upstream of the *AtPDS3* gene.

Transgenic plants were created via standard floral dip transformation utilizing the same

plasmids as for the protoplast experiments [21]. To test for sensitivity to temperature, transgenic

plants expressing ISYmu1 were either grown at room temperature or subjected to a heat shock

treatment. We tested editing in wild type (WT) plants, as well as in the *rna dependent rna*

polymerase 6 (*rdr6*) mutant which is known to reduce transgene silencing [22]. Analysis using

amp-seq revealed an average editing efficiency of 1.6% and 2.5% for gRNA2 in WT and *rdr6*,

respectively (Fig. 1d). Analysis of gRNA12 revealed greater editing than gRNA2, averaging 44.9%

editing in WT and 75.5% in *rdr6* (Fig. 1e). Comparison of editing efficiency in the plants grown at

room temperature with those that received the heat shock treatment revealed a preference for

increased temperature for both target sites in the WT background, demonstrating 6.3-fold and

1.4-fold increases in editing for gRNA2 and gRNA12, respectively (Fig. 1d, e). In *rdr6* we

observed a 13-fold increase in editing for gRNA2, but little change in editing for gRNA12 (Fig.

1d, e). These data demonstrate that ISYmu1, encoded as a transgene, is capable of performing

efficient genome editing in *Arabidopsis* plants, and that heat treatment and the *rdr6* silencing

mutant can be used to increase editing efficiency.

Encouraged by the ISYmu1 activity in transgenic *Arabidopsis* plants, we next tested

ISYmu1 for TRV-mediated genome editing. TRV is a bipartite RNA virus composed of TRV1 and

TRV2 (Fig. 2a). Previous work has shown that the TRV2 RNA can be engineered by inserting a

cargo expression cassette downstream of the pea early browning virus promoter (pPEBV) (Fig.

2a) [23,24]. To test ISYmu1 for genome editing capabilities via TRV delivery to *Arabidopsis,* we

engineered two TRV2 Cargo Architectures. In TRV2 Architecture_A, the tRNA[Ileu] was directly

downstream of the TnpB and gRNA sequences (Fig. 2a). In TRV Architecture_B, we included an

HDV ribozyme sequence between the guide and tRNA[Ileu] sequences.(Fig. 2a). We included

tRNA[Ileu] in both designs as it was previously shown to promote systemic TRV movement and

transmission of edited alleles to the next generation [23,24].

First, we evaluated TRV-mediated editing potential with gRNA2 using both TRV2

Architecture_A and Architecture_B. gRNA2 was selected because it targets the *AtPDS3* coding

sequence, enabling easy phenotypic screening for editing due to white photobleaching of cells

containing biallelic mutations [23,24]. We delivered TRV vectors to both WT and the *ku70* genetic

mutant. *Ku70* plays a role in the nonhomologous end joining (NHEJ) double strand break (DSB)

repair pathway [25]. ISYmu1-mediated editing efficiency should be greater in the *ku70* genotype if

double stranded breaks generated by ISYmu1 are repaired through NHEJ. Each TRV2 plasmid

was co-delivered with the TRV1 plasmid to *Arabidopsis* plants using the agroflood method [24].

White speckles were observed on some of the leaves around three weeks post agroflooding,

suggesting that sectors of cells contained biallelic mutations in the target *AtPDS3* gene (Fig. 2b).

Amp-seq analysis revealed an average of 0.1% and 0% editing efficiency in leaf tissue of WT and

ku70 plants agroflooded with TRV2 Architecture_A and grown under room temperature,

respectively (Fig. 2c). For the heat shock treated plants, we observed an average editing

efficiency of 0.4% in WT and 0.7% in *ku70* plants agroflooded with TRV2 Architecture_A (Fig.

2c). Using TRV2 Architecture_B we observed an average 0.6% and 2% editing in WT and *ku70*,

respectively, for the room temperature grown plants. For the plants that received TRV

Architecture_B and a heat shock, we observed an average editing efficiency of 3.3% in WT and

8.9% in *ku70* (Fig. 2c). These results show that Architecture_B, containing the HDV ribozyme,

generated higher editing than Architecture_A, and that the *ku70* mutant can enhance editing

efficiency.

Next, we tested gRNA12 utilizing the TRV2 Architecture_B, since this architecture

demonstrated the highest levels of editing for gRNA2. Using the same agroflood TRV delivery

method, we observed an average 8.51% and 4.27% editing efficiency in room temperature and

heat treatment growth conditions, respectively (Fig. 2d). Further, 6/57 plants displayed editing

greater than 40%, with four greater than 75%, when the room temperature treatment was

used (Fig. 2d). Again, analysis of the repair outcomes showed deletion dominant profiles for

ISYmu1 gRNA2 and gRNA12 (Fig. 2e, f).

To test for transmission of edited alleles to the next generation, we first screened the

progeny of a WT plant showing 54.54% somatic editing using the TRV2 Architecture_B design

with gRNA2 that underwent heat shock treatment. In total, 2318 seeds were sown on ½ MS

plates containing 3% sucrose. After ten days, 68 albino seedlings were observed, suggesting

biallelic mutations in the *PDS3* gene (Fig. 3a). To confirm *AtPDS3* was mutated we performed

Sanger sequencing on the two white seedlings shown in Fig. 3a, which revealed both plants to

be homozygous for a 4bp frame-shift deletion (Fig. 3b). To further characterize transmission of

edited alleles, amp-seq on 209 seedlings (41 albino and 168 green) showed that all of the albino

seedlings contained biallelic mutations, with the majority of mutations being the 4bp deletion

observed in Fig. 3b (Supplementary Table 2). Of the 168 green seedlings, eight were

heterozygous (4bp deletion/WT) (Supplementary Table 2).

Next, we characterized transmission of edited alleles from two individual lines, plant 54

(80.5% somatic editing) and plant 69 (77.1% somatic editing), that underwent agroflood using

gRNA12 TRV2 Architecture_B with the room temperature condition. Using Sanger sequencing,

we analyzed the genotypes of 148 and 75 progeny seedlings from plants 54 and 69,

respectively. Sanger sequencing analysis of the progeny from plant 54 revealed 27 (18%)

biallelic and 25 (17%) monoallelic edited plants (Supplementary Table 2, Extended Data Fig. 4a)

[26]. For plant 69 we observed higher transmission of edited alleles, totaling 32 (43%) biallelic and

15 (20%) monoallelic edited plants (Supplementary Table 2, Extended Data Fig. 4b) [26]. These

data demonstrate the heritability of edits generated via TRV-delivery of ISYmu1 at two distinct

target sites.

It has been demonstrated that TRV is not transmitted to the next generation following

agroflood inoculation of plants [23,24]. To confirm the TRV was not present in the progeny of a

TRV-infected plant, RT-PCR was performed on five albino plants harboring homozygous 4bp

deletions at *AtPDS3*. Consistent with the literature, TRV was not detected in any of the albino

plants (Extended Data Fig. 5) [23,24]. These data indicate that TRV-mediated biallelic edits using

ISYmu1 are heritable and virus-free.

To evaluate off-target editing we surveyed three individual albino plants harboring

biallelic mutations generated by ISYmu1 TRV2 Architecture_B gRNA2. Whole genome

sequencing was performed to generate an average 770x coverage, with greater than 99% of the

genome covered by mapped reads (Supplementary Table 3). In all three samples, we confirmed

the targeted mutations in the *AtPDS3* gene, as previously identified using amp-seq.

Additionally, we found a large number of variant differences compared to the Col-0 reference

genome both in the control and the edited plants (Supplementary Table 4), suggesting that

most of the variants detected are due to spontaneous mutations present in our lab strain of

Arabidopsis. To screen for variants potentially caused by ISYmu1 off-target editing, all variants

in the edited plants were filtered with variants already present in the control background.

Variants with coverage lower than 30-fold were also filtered out. The remaining variants were

checked manually for any false positive variant calling. In the three albino plants we sequenced,

only 5, 5, and 4 variants were detected, and these variants are all outside the predicted

potential off-target sites based on sequence similarity to the PDS3 gRNA2 sequence

(Supplementary Tables 4, 5) [27]. In line with ISYmu1 off-target analysis reported in rice and

human cells [17,18] , these data further demonstrate the high target site specificity of ISYmu1.

A long-term goal of plant scientists has been the development of fast and easy means of

editing plant genomes without the need for tissue culture and transgenesis. Here, we

developed an approach utilizing the ultracompact site-specific TnpB genome editor, ISYmu1,

together with tobacco rattle virus, for heritable plant genome editing. These results should

accelerate high throughput genome editing for both basic and applied research. We anticipate

this approach will be applicable to other novel TnpBs, various viral vectors and a number of

plant species for genome editing. Recent work has uncovered many TnpB systems from diverse

microbial sources, including enzymes with unique PAM sequence specificities [28], which can

increase the range of target DNA sequences that could be edited using this approach. The TRV

virus used in this study has a broad host range of over 400 species, including many solanaceous

plants such as tomato, ornamental plants, and other crops [29]. Additionally, plant viruses with

similar cargo capacities such as Potato Virus X and Barley Stripe Mosaic Virus are likely to be

amenable to this approach since it has been demonstrated they are capable of viral-mediated

heritable gene editing by delivering the gRNA to a Cas9-expressing transgenic plant [30]. In

addition to being an important tool for crop biotechnology, viral delivery of TnpBs could enable

high throughput CRISPR screens in model plant species such as *Arabidopsis*, further unlocking

their potential for genetic discovery.

Materials and Methods:

<u>Plasmids used in this study</u>

Plasmids used for bacterial assay were generated as follows. The single expression

cassette containing TnpB and ωRNA sequences were synthesized as geneblocks from IDT

(Integrated DNA Technologies), and were golden-gate cloned using BsmbI restriction enzyme

(cat E1602S) into a vector (chloramphenicol resistance) under a single tetracycline-inducible

promoter (TetR/pTet) to make the TnpB-ωRNA plasmid. Target sites with various PAM

sequences and target sites were golden-gate cloned with BBsI restriction enzyme(cat R3539S)

into a vector (ampicillin/carbenicillin resistance).

Plasmids were generated for protoplast and floral dip experiments in a two-step cloning

strategy. In step one, the ISDra2, ISYmu1, and ISAam1 protein coding sequences and their

ωRNAs were synthesized as geneblocks by IDT (Integrated DNA Technologies). Then, starting

with the pC1300_pUB10_pcoCASphi_E9t_MCS_version2 vector (Li et al. 2023), we used

NEBuilder HiFi DNA Assembly (catE2621) and PCR to assemble the TnpB-ωRNA geneblocks into

plant expression vectors with a toxic *ccdB* insert flanked by PaqCI sites immediately

downstream of the ωRNA scaffold and preceding an HDV ribozyme sequence. The HiFi

reactions were then transformed into One Shot *ccdB* Survival 2 T1R Competent Cells

(catA10460) to obtain the pMK003 (ISDra2), pMK025 (ISYmu1), and pMK024 (ISAam1)

intermediate vectors for facile guide sequence cloning (Supplementary Table 6). In step two,

guide sequences were synthesized as individual top and bottom strands with 4 base pair

overhangs from IDT, phosphorylated and annealed, and then used for golden-gate assembly

using the NEB PaqCI (catR0745) enzyme (Supplementary Table 7). When transformed into NEB

10-beta Competent E. coli (catC3019) vectors that still contained the *ccdB* gene would kill the

cells, leaving behind only the transformants possessing successfully assembled TnpB plant

expression vectors harboring a guide RNA sequence. A full list of all plasmids can be found in

Supplementary Table 6.

270   TRV vectors were created with the pDK3888 TRV2 plasmid as a base vector [24]. NEBuilder

HiFi DNA Assembly (catE2621) was used to clone the ISYmu1 gRNA2 Architecture_A, ISYmu1

gRNA2 Architecture_B, and ISYmu1 gRNA12 Architecture_B into the TRV2 cargo slot. First,

pDK3888 was digested using NEB ZraI (catR0659), NEB PmlI (catR0532), and NEB Quick CIP

(catM0525) overnight, and purified using Qiagen QiaQuick purification column (cat28104).

Next, three PCR reactions were performed to amplify the fragments needed for NEBuilder HiFi

DNA Assembly, followed by purification using Qiagen QiaQuick purification column (cat28104)

(Supplementary Table 8). Then, the digested and purified pDK3888 plasmid, and purified PCR

fragments were used to assemble the final TRV2 plasmid using NEBuilder HiFi DNA Assembly

according to the manufacturer's protocol. Finally, the NEBuilder HiFi DNA Assembly reaction

was transformed into NEB 10-beta Competent E.coli (catC3019). Correct plasmids were

confirmed using Primordium whole plasmid sequencing. Plasmids and their descriptions can be

found in Supplementary Table 6.

<u>Bacterial interference assay</u>

For the bacterial interference assay, we co-transfected 100 ng of the TnpB-ωRNA

plasmid and 100 ng of the target plasmid to 33 µl of NEB 10-beta electrocompetent E. coli cells

(cat C3020K). Specifically, the target plasmid contains a target site either flanking the canonical

PAM (TTGAT for ISYmu1 and ISDra2 and TTTAA for ISAam1) or flanking a non-canonical PAM

(GGGGG). The cells were recovered in 1 ml of NEB 10-Beta Stable/Outgrowth media(cat

B9035S) for 1 hour. Following recovery, a series of 5-fold dilutions of the recovery culture were

prepared. Each dilution (5 µl) was spot-plated onto LB-Agar plates containing double antibiotics

(34 µg/ml chloramphenicol, 100 µg/ml carbenicillin, and 2 nM anhydrotetracycline) and onto

control plates with a single antibiotic (34 µg/ml chloramphenicol and 2 nM

anhydrotetracycline). If no colonies were visible on the serial dilution plates, 400 µl of the 1 ml

recovery culture was plated entirely on the double antibiotic plate to enhance detection

sensitivity. Plates were left overnight at either 26°C or 37°C, and colony-forming units (CFU)

were counted on all plates the next morning. The normalized CFU were calculated by taking the

ratio of CFU on the double antibiotic plates to the CFU on the single antibiotic plates. The

normalized CFU in the canonical PAM conditions were compared to the one in the non-

canonical PAM conditions. Experiments were performed in triplicate.

<u>Plant materials and growth conditions</u>

For protoplast preparation, *Arabidopsis* Columbia ecotype (Col-0) seeds were suspended

in a 0.1% agarose solution and kept at 4°C in the dark for three days to stratify. Following

stratification, seeds were planted on Jiffy pucks and grown under a 12-h light/12-h dark

photoperiod, with low light condition for 3 to 4 weeks [20].

For the creation of transgenic plants, the *Arabidopsis* Col-0 ecotype was used. The *ku70*

(SALK_123114) genotype was provided from Feng Zhang lab at the University of Minnesota. The

rdr6 genotype was created using CRISPR-Cas9, resulting in a 616 bp deletion in the gene body of

rdr6. Floral dip transformation was performed according to the protocol as previously outlined

using the Agl0 agrobacterium strain [21]. Transgenic T1 plants were screened using ½ MS plates

with 40 µg/mL hygromycin B under a 16-/8-h light/dark cycle at 23°C. After one week,

transgenic seedlings that passed selection were transferred to soil and moved to a greenhouse

(23°C) for the rest of their life cycle.

For agroflood experiments, sterilized seeds were sown on ½ MS agar plates and

stratified for five days. After five days, the seeds were moved to a growth room and grown

under a 16-/8-h light/dark cycle at 23°C for 8-10 days. The seedlings were then used for TRV

delivery.

A subset of transgenic T1 plants, and plants that underwent agroflood, were subjected

to a heat shock treatment modified from LeBlanc et al [31]. Seedlings that passed selection, or

underwent agroflood, were then transplanted to soil and grown in a greenhouse (23°C) for one

week. After one week, plants that did not receive a heat shock treatment continued to grow in

the greenhouse (23°C); however, plants that underwent heat shock treatment were exposed to

8 hours (9am - 5pm) of heat exposure at 37°C every day for 5 days, followed by 2 days of

recovery a greenhouse (23°C). This heat shock regime lasted for two weeks.

Protoplast isolation and transfection

Arabidopsis mesophyll protoplast isolation was performed as previously described [20].

Plasmid transfections into *Arabidopsis* protoplasts were performed using 20µg of plasmid,

similarly as [32]. The concentrations of plasmids were determined by nanodrop. Plasmids were

added to the bottom of each transfection tube, and the volume of the plasmids was

supplemented with water to reach 20 µL. 200 µL of protoplasts were added followed by 220 µL

of fresh and sterile polyethylene glycol (PEG)-CaCl2 solution. The samples were mixed by gently

tapping the tubes, and incubated at room temperature for 10 minutes. After 10 minutes, 880

µL of W5 solution was added and mixed with the protoplasts by inverting the tube two to three

times to stop the transfection. Next, protoplasts were harvested by centrifuging the tubes at

100 relative centrifugal force (RCF) for 2 min and resuspended in 1 mL of WI solution. The

protoplast cells were then plated in 6-well plates pre-coated with 5% calf serum. Protoplast

cells in the 6-well plates were incubated at 26°C for 48 hours. During the 48 hour incubation,

the protoplast cells were subjected to a 37°C heat shock treatment for 2 hours at 16 hours post

transfection. At 48 hours post transfection, protoplasts were harvested for genomic DNA

extraction.

<u>TRV delivery to *Arabidopsis* seedlings</u>

TRV delivery was performed as previously described [24]. TRV1 and TRV2 vectors were

first introduced into the GV3101 agrobacterium strain. The agrobacterium harboring TRV

vectors were then grown in 200 ml of lysogeny broth (LB) with antibiotics for 18 hours at 28°C.

Agrobacterium cultures were centrifuged for twenty minutes at 3500 RCF. The LB was

discarded and the agrobacterium cells were resuspended in 200 ml of sterile water. The

resuspended agrobacterium was centrifuged for 10 minutes at 3500 RPM. The supernatant was

discarded and the pellet was resuspended in sterile agro-infiltration buffer containing 10 mM

MgCl2, 10 mM 2-(N-morpholino) ethanesulfonic acid, and 250 µM acetosyringone to OD600 =

1.5. The agrobacterium cells were then incubated at 23°C for three hours with slow shaking.

After three hours, the agrobacterium harboring TRV1 and TRV2 were mixed in a 1:1 ratio. 15 ml

of the 1:1 ratio of TRV was delivered to seedlings at 8-10 days old. After four days of agroflood

co-culture, seedlings were transplanted to soil.

<u>Screening the progeny of TRV-infected plants for edits</u>

Seeds were harvested from the TRV-infected plants about 12 weeks after TRV delivery.

The seeds were sown on ½ MS plates supplemented with 3% sucrose and stored at 4°C in the

dark for five days to stratify. After five days, the seeds were moved to a growth room and

grown under a 16-/8-h light/dark cycle at 23°C for 10-12 days. Amp-seq or Sanger sequencing

was then performed on a subset of the progeny plants using primers listed in Supplementary

Table 9.

<u>Next generation amplicon sequencing</u>

DNA was extracted from protoplast samples with Qiagen DNeasy plant mini kit (Qiagen

69106). Tissue was collected from transgenic plants by sampling and pooling leaf tissue from

three random leaves on a single plant three weeks after being transplanted to soil. For the

plants that underwent agroflood, leaf tissue was sampled by collecting and pooling tissue from

three random (however, if white sectors visible they were sampled) leaves on a single plant

distal to the TRV delivery site three weeks after being transplanted to soil. Once tissue samples

were collected, they were frozen at -80C overnight. The samples were then ground and DNA

was extracted using the invitrogen Platinum Direct PCR Universal Master Mix (refA44647500)

according to the manufacturer's instructions. For the progenies of plants that underwent

agroflood, a single leaf tissue was sampled and DNA was extracted using invitrogen Platinum

Direct PCR Universal Master Mix (REF A44647500) according to the manufacturer's instructions.

The DNA was then used for Next Generation Amplicon Sequencing.

As similarly performed by Li et al [32], editing efficiency was characterized using single-

end Next Generation Sequencing on the Illumina NovaSeqX platform. Libraries were prepared

via a 2-step PCR amplification method. In the first round of amplification, each target site was

amplified using primers flanking the target site (Supplementary Table 9). After 25 cycles of

amplification, the reactions were cleaned using a 1.0X Ampure XP bead purification (Beckman

Coulter A63881). Next, each sample went through 12 additional cycles of amplification using

Illumina indexing primers. The samples were cleaned using a 0.7X Ampure XP purification.

Samples were checked for purity on a 2% agarose gel, quantified using Nanodrop, normalized,

and pooled.

Next generation amplicon sequencing analysis

Amplicon sequencing analysis was performed similarly as Li et al [32]. Only single-end

reads were used for analysis. Reads were adapter trimmed using Trim Galore default settings.

Remaining reads were mapped to the target genome region using the BWA aligner (v0.7.17,

BWA-MEM algorithm). Sorted and indexed bam files were used as input files for further analysis

by the CrispRvariants R package (v1.14.0). Each mutation pattern with corresponding read

counts was exported by the CrispRvariants R package. After assessing all control samples, a

criterion to classify reads as edited was established: only reads with a >= 3 bp deletion or

insertion (indel) of the same pattern (indels of same size starting at the same location) with

>=10 read counts from a sample were counted as edited reads. SNVs were also filtered out.

Off-target analysis

Off-target analysis was performed as previously described in Li et al[32]. DNA from single

Arabidopsis seedlings was extracted with the Qiagen DNeasy plant mini kit and sheared to 300

bp size with a Covaris. Library preparation was performed with Tecan Ovation Ultralow V2 DNA-

seq kit. For variant calling, WGS reads were aligned to the TARI10 reference genome using BWA

mem (v0.7.17)[33] with default parameters. GATK (4.2.0.0)[34] MarkDuplicatesSpark was used to

remove PCR duplicate reads. Then GATK HaplotypeCaller was used to call raw variants. Raw SNPs

were filtered with QD < 2.0, FS > 60.0, MQ < 40.0, and SOR > 4.0. Raw InDels were filtered with

QD < 2.0, FS > 200.0, and SOR > 10.0 and used for base quality score recalibration. The

recalibrated bam was further applied to GATK and Strelka (v2.9.2) SNPs/InDel calling. Only

SNPs/InDels called by both GATK and Strelka were used for further filtering. The intersection of

SNPs/InDel called by GATK with Strelka (v2.9.2)[35] is obtained by BedTools (v2.26.0) [36]. SNPs/InDel

filtered with wild type background were conducted by BedTools (v2.26.0). Variants with coverage

lower than 30 depth were filtered.

RT-PCR

Total RNA from TRV-infected progeny plants was extracted using Zymo Research Direct-

zol RNA MiniPrep kit (catR2052). Total RNA was converted to cDNA using the invitrogen

SuperScript IV VILO Master Mix (cat11766050). The RT-PCR control was performed using

primers targeting the *AtIPP2* gene (Supplementary Table 10). PCR was performed to check for

the presence/absence of the TRV vector using SP9238 and SP9239 (Supplementary Table 10) [24].

PCR was performed with New England Biolabs Q5 High-Fidelity 2X Master Mix (catM0492L)

according to the manufacturer's instructions, using 2 µl of cDNA in a 25 µl reaction. PCR

conditions included a 98°C initial denaturation step for 30 seconds, 35x(98°C, 10 sec; 55°C, 20

sec; 72°C, 10 sec), 72°C for 2 min. 10 µl of PCR amplicons were analyzed by 2% agarose gel

electrophoresis.

Data availability:

All the amp-seq data generated in this study will be accessible at NCBI Sequence Read Archive

under BioProject PRJNA1124592. Whole genome sequencing data is accessible at BioProject

PRJNA1146711

Acknowledgements:

This work was supported by an NSF Plant Genome Research Program grant to SEJ, JAD and JFB.

SEJ and JAD are Investigators of the Howard Hughes Medical Institute. H.S. is an HHMI Fellow of

The Jane Coffin Childs Fund for Medical Research. B.A.A. is supported by m-CAFEs Microbial

Community Analysis & Functional Evaluation in Soils, a Science Focus Area led by Lawrence

Berkeley National Laboratory based upon work supported by the US Department of Energy,

Office of Science, Office of Biological & Environmental Research (DE-AC02-05CH11231). We

thank Suhua Feng, Mahnaz Akhavan and the Broad Stem Cell Research Center Biosequencing

Core for sequencing support, Colette Picard for assistance with analysis of amplicon sequencing

data, Dinesh Kumar for supplying the TRV2 plasmid pDK3888, Feng Zhang for providing the

ku70 (SALK123114) mutant, and all members of the Jacobsen lab for their helpful insight and

suggestions.

Author Contributions:

TW, MK, HS, ZL, BAA, JB, JAD and SEJ designed the research; TW, HS, ZZ, JAD, and SEJ

interpreted the data; TW and SEJ wrote the manuscript; TW, MK, HS, ZL, JA, MS, KV, GW, SC,

CA, NS, AS, and DS performed experiments.

Competing interests:

T.W. ,M.K, J.A, Z.L, H.S, B.A.A., J.A.D. and S.E.J. have filed a patent covering aspects of this work.

Figures and Tables:

Figure 1: Expression of TnpB and guide RNA in a single transcript for plant genome editing. (a)

Barplots of interference assay testing the single transcript expression TnpB vectors for cleavage

in E. coli. The barplot on the top displays data from experiments performed at 26°C and the

barplot below depicts data from experiments performed at 37°C. Black bars indicate absence of

a PAM on the target plasmid and purple bars indicate the presence of a PAM on the target

plasmid. The Y-axis is a log10 scale of the normalized colony forming units (CFU)/milliliter (mL).

The X-axis displays the three TnpBs tested using the single expression transcript design without

a HDV ribozyme or the single expression design with a HDV ribozyme. The standard error of the

mean (SEM) was calculated for each experiment. **(b)** Schematic of the single expression

transcript TnpB-ωRNA plasmid design used for plant genome editing. The green arrow

symbolizes the At*UBQ10* promoter; the dark gray boxes indicate the 2X-FLAG, SV40 NLS and

HDV ribozyme sequences; the light gray boxes indicate the TnpB-ωRNA and guide sequences;

the red box symbolizes the rbcS-E9 terminator; the black arrow indicate the orientation of the

TnpB-ωRNA expression cassette. **(c)** Barplot displaying the average editing frequency for

protoplast experiments using ISDra2, ISYmu1 and ISAam1. Each dot represents the average

editing efficiency of a gRNA from Extended Data Fig. 3a. The Y-axis indicates the editing

efficiencies (percent indel reads (%)). The X-axis displays the ISDra2, ISYmu1 and ISAam1 TnpBs

tested. The standard error of the mean (SEM) was calculated for each experiment. **(d-e)** ISYmu1

somatic editing in T1 transgenic plants. Panels d and e display a barplot for ISYmu1 gRNA2 and

ISYmu1 gRNA12, respectively. Each dot indicates a single T1 transgenic plant. The room and HS

treatments stand for room temperature and heat shock plant growth conditions, respectively.

The genotypes are plotted along the X-axis and the editing efficiencies (percent indel reads (%))

are plotted on the Y-axis. The standard error of the mean (SEM) was calculated for each

experiment. WT is an abbreviation for wild type, and *rdr6* is an abbreviation for a mutation in

the *RNA-DEPENDENT RNA POLYMERASE 6* gene.

Figure 2: Somatic editing of *Arabidopsis* using TRV to deliver ISYmu1 TnpB and guide RNA. (a)
Schematic of the TRV1 and TRV2 plasmids. Green arrows indicate the RNA dependent RNA
polymerase (RDRP) and pea early browning virus (pPEBV) promoters for TRV1 and TRV2,
respectively; the gray boxes in TRV1 and TRV2 indicate the native TRV components; the red
Cargo box in TRV2 indicates the location of either Architecture_A or Architecture_B; below
TRV2 are schematics of the components, Architecture_A or Architecture_B, cloned into the
TRV2 Cargo slot. **(b)** Representative picture of a plant displaying white sectors about three
weeks after TRV delivery. The yellow arrows indicate leaves containing white sectors. **(c)**
Barplot displaying the somatic editing efficiencies for ISYmu1 gRNA2 in wild type (WT) and *ku70*
genetic backgrounds. WT samples are black and *ku70* samples are blue. The TRV2 Cargo
Architectures are plotted along the X-axis with either room or HS treatment. The room and HS
along the X-axis stand for room temperature and heat shock plant growth conditions,
respectively. The Y-axis indicates the editing efficiencies (percent indel reads (%)). Each dot

represents an individual plant that underwent agroflood TRV delivery. The standard error of the

mean (SEM) was calculated for each experiment. **(d)** Barplot displaying the somatic editing

efficiencies for ISYmu1 gRNA12 in wild type (WT). Room temperature treatment (room)

samples are black and heat shock (HS) samples are red. The TRV2 Cargo Architecture_B is

plotted along the X-axis. The Y-axis indicates the editing efficiencies (percent indel reads (%)).

Each dot represents an individual plant that underwent agroflood TRV delivery. The standard

error of the mean (SEM) was calculated for each experiment. **(e-f)** Panel e displays the DNA

indel repair profile for an individual WT plant that underwent delivery of TRV Cargo

Architecture_B with ISYmu1 gRNA2 under the heat shock treatment. Panel f shows the DNA

indel repair profile for an individual WT plant that underwent delivery of TRV Cargo

Architecture_B with ISYmu1 gRNA12 under the heat shock treatment. The top five most

common indel types are listed on the left. The read counts for each indel are listed on the right.

The PAM is identified by the red box, and the target site is outlined by the black box, in the

Reference sequence. The total read number and editing efficiency are listed below each indel

profile.

a

b

tggttgtgtttgggaatgtttctgcggcgaatttgccttatcaaaacgggtttttgg

Figure 3: Inheritance of edited alleles. (a) Representative image of albino and green progeny

seedlings from a WT plant showing 54.54% somatic editing using the TRV2 Architecture_B

design with gRNA2 that underwent heat shock treatment. **(b)** Sanger sequencing trace file

screenshot from one of the albino plants in Fig 3a. The sequence at the top is the wild type

reverse complement; below that are the ISYmu1 gRNA2 target and PAM (gray box); the ab1

trace file displays a homozygous 4bp deletion.

Extended Data Fig. 1: Schematic of interference assay plasmids containing either the TnpB-

ωRNA Single or TnpB-ωRNA Single-HDV design. The blue arrow indicates the TnpB sequence;

the yellow arrow indicates the ωRNA sequence; the black rectangle indicates the guide

sequence; the green arrow indicates the HDV ribozyme sequence. The plasmids contain the

tetracycline resistance gene (TetR). A tetracycline promoter (pTet) was used to drive expression

of the TnpB-ωRNA single or Single-HDV sequences, and the tetracycline resistance gene.

Extended Data Fig. 2: Plate images of bacterial plasmid interference assay. Five-fold serial

dilutions (5 µL) from the 1 mL recovery culture post transformation were plated on both single

antibiotic LB-Agar plates (Cam, upper row) and double antibiotic LB-Agar plates (Cam+Carb,

lower row). Plates without visible colonies (indicated by an asterisk) had 400 µL of the original 1

mL recovery culture plated on the double antibiotic plates (dash insets). Experiments were

performed in triplicates.

Extended Data Fig. 3: Editing efficiency and repair profiles for ISDra2, ISYmu1 and ISAam1 in protoplast experiment. (a) The name of each TnpB tested is at the upper left of each barplot. The gRNAs are plotted along the X-axis and the editing efficiency (percent indel reads (%)) is plotted on the Y-axis. Each dot indicates a single transfection. The standard error of the mean (SEM) was calculated for each target site. **(b)** DNA repair indel profiles for individual transfection samples. The top five most common indel types are listed on the left. The read counts for each indel are listed on the right. The PAM is identified by the red box, and the target site is outlined by the black box, in the Reference sequence. The total read number and editing efficiency are listed below each indel profile. The name of each TnpB and gRNA is displayed above the reference sequence of each indel repair profile.

Extended Data Fig. 4: Sanger sequencing screenshots of the progeny plant genotypes for

ISYmu1 gRNA12. (a-b) Panels a and b correspond to the progeny from plants 54 and 69,

respectively. The sequence at the top is the wild type genomic sequence; below that are the

ISYmu1 gRNA12 target and PAM (gray box); the ab1 trace file displays the mutation.

Extended Data Fig. 5: RT-PCR gel showing absence of the TRV in the progeny of a TRV-infected

plant. RT-PCR was performed using total RNA extracted from albino plants homozygous for a

4bp deletion at the *AtPDS3* gene. Gel electrophoresis image for RT-PCR performed using

primers targeting TRV (upper panel) and *AtIPP2* gene (bottom panel). The lanes are indicated

(from left to right) as ladder, six individual albino plants, plasmid control, and a water control.

Black arrows indicate the amplicon size in base pairs (bp).

TnpB	Guide	gRNA sequence	Genomic location
ISDra2	gRNA1	ttacgaattgatgaccatat	4:8195186-8195206
ISDra2	gRNA2	aaggcaaattcgccgcagaa	4:8194730-8194749
ISDra2	gRNA3	tcacattaagcctagaaact	4:8193921-8193940
ISDra2	gRNA4	cccaagttctccaaataaat	4:8193476-8193495
ISDra2	gRNA5	tacccatcctaaagtatggg	4:8192944-8192963
ISDra2	gRNA7	aaattcaacatctttctcta	4:8190835-8190854
ISDra2	gRNA8	tggtctcactttccgaatta	4:8194167-8194186
ISDra2	gRNA9	agctttgaaccggtttcttc	4:8192305-8192324
ISDra2	gRNA11	cgaaaactgaagaacacata	4:8191494-8191513
ISDra2	gRNA12	gcgttggagcatataacaga	4:8195556-8195575
ISDra2	gRNA13	taaagagaggaaattgcagg	4:8194513-8194532
ISDra2	gRNA14	ggtagagctgataagatata	4:8193625-8193644
ISDra2	gRNA15	aaaattggattaatgtgcac	4:8193203-8193222
ISDra2	gRNA16	tactattaaatgtcaaaatc	4:8193077-8193096
ISDra2	gRNA17	caatacaaataaatacatgc	4:8193117-8193136
ISDra2	gRNA18	caattcaagctaattataga	4:8193095-8193114
ISDra2	gRNA19	gagcttaacttggtagagta	4:8192577-8192596
ISDra2	gRNA20	ttgtcagctttcttatggat	4:8192528-8192547
ISDra2	gRNA21	gttgattaacttgtactacc	4:8192131-8192150
ISDra2	gRNA22	taacttgtactacctcatcc	4:8192125-8192144
ISYmu1	gRNA1	ttacgaattgatgacc	4:8195187-8195202

ISYmu1	gRNA2	aaggcaaattcgccgc	4:8194730-8194745
ISYmu1	gRNA3	tcacattaagcctaga	4:8193921-8193936
ISYmu1	gRNA4	cccaagttctccaaat	4:8193476-8193491
ISYmu1	gRNA5	tacccatcctaaagta	4:8192944-8192959
ISYmu1	gRNA7	aaattcaacatctttc	4:8190839-8190854
ISYmu1	gRNA8	tggtctcactttccga	4:8194171-8194186
ISYmu1	gRNA9	agctttgaaccggttt	4:8192309-8192324
ISYmu1	gRNA10	taacttgtactacctc	4:8192129-8192144
ISYmu1	gRNA12	gcgttggagcatataa	4:8195560-8195575
ISAam1	gRNA11	caattcatctggtatc	4:8191755-8191770
ISAam1	gRNA12	ccaagaacaagcctta	4:8191347-8191362
ISAam1	gRNA13	gttttgtcctcttctc	4:8190412-8190427
ISAam1	gRNA14	caatttacctatctta	4:8193797-8193812
ISAam1	gRNA15	cacataattgaaaaga	4:8194919-8194934
ISAam1	gRNA16	aatttgttacacaact	4:8195755-8195770
ISAam1	gRNA17	tttgtgtggtatttaa	4:8195158-8195173

559 **Supplementary Table 1: ISDra2, ISYmu1 and ISAam1 target sites.** The table consists of four

560 columns. The TnpB and Guide columns indicate the TnpB gRNA used. The gRNA sequence

561 column lists the gRNA sequence used for targeted genome editing. The Genomic location

562 column provides the location of each gRNA target site in the *Arabidopsis* genome.

Plant ID	Phenotype	Genotype (allele1 / allele2)	Sequencing type
gRNA2_HS_116-1	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-2	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-3	WT	WT / WT	amp-seq
gRNA2_HS_116-4	WT	WT / WT	amp-seq
gRNA2_HS_116-5	WT	4bp del / WT	amp-seq
gRNA2_HS_116-6	WT	WT / WT	amp-seq
gRNA2_HS_116-9	WT	WT / WT	amp-seq
gRNA2_HS_116-10	WT	WT / WT	amp-seq
gRNA2_HS_116-12	WT	WT / WT	amp-seq
gRNA2_HS_116-13	WT	WT / WT	amp-seq
gRNA2_HS_116-14	WT	WT / WT	amp-seq
gRNA2_HS_116-15	WT	WT / WT	amp-seq
gRNA2_HS_116-16	WT	WT / WT	amp-seq
gRNA2_HS_116-17	WT	WT / WT	amp-seq
gRNA2_HS_116-18	WT	WT / WT	amp-seq
gRNA2_HS_116-19	WT	WT / WT	amp-seq
gRNA2_HS_116-20	WT	WT / WT	amp-seq
gRNA2_HS_116-21	WT	WT / WT	amp-seq
gRNA2_HS_116-22	WT	WT / WT	amp-seq
gRNA2_HS_116-23	WT	WT / WT	amp-seq
gRNA2_HS_116-24	WT	WT / WT	amp-seq

gRNA2_HS_116-25	WT	WT / WT	amp-seq
gRNA2_HS_116-26	WT	WT / WT	amp-seq
gRNA2_HS_116-27	WT	WT / WT	amp-seq
gRNA2_HS_116-28	WT	4bp del / WT	amp-seq
gRNA2_HS_116-29	WT	WT / WT	amp-seq
gRNA2_HS_116-30	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-31	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-32	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-33	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-34	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-35	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-36	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-38	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-39	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-40	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-41	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-42	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-43	Albino	11 bp del / 17bp del	amp-seq
gRNA2_HS_116-45	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-46	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-47	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-48	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-49	Albino	10bp del / 12 bp del	amp-seq

gRNA2_HS_116-50	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-51	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-52	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-53	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-54	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-56	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-57	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-58	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-59	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-60	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-61	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-65	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-72	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-79	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-80	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-84	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-85	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-86	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-90	Albino	17bp del / 11bp del	amp-seq
gRNA2_HS_116-91	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-92	Albino	4bp del / 4bp del	amp-seq
gRNA2_HS_116-94	WT	WT / WT	amp-seq
gRNA2_HS_116-95	WT	WT / WT	amp-seq

gRNA2_HS_116-96	WT	WT / WT	amp-seq
gRNA2_HS_116-97	WT	WT / WT	amp-seq
gRNA2_HS_116-98	WT	WT / WT	amp-seq
gRNA2_HS_116-99	WT	WT / WT	amp-seq
gRNA2_HS_116-100	WT	4bp del / WT	amp-seq
gRNA2_HS_116-101	WT	WT / WT	amp-seq
gRNA2_HS_116-102	WT	WT / WT	amp-seq
gRNA2_HS_116-103	WT	WT / WT	amp-seq
gRNA2_HS_116-104	WT	WT / WT	amp-seq
gRNA2_HS_116-105	WT	WT / WT	amp-seq
gRNA2_HS_116-106	WT	WT / WT	amp-seq
gRNA2_HS_116-107	WT	WT / WT	amp-seq
gRNA2_HS_116-108	WT	4bp del / WT	amp-seq
gRNA2_HS_116-109	WT	WT / WT	amp-seq
gRNA2_HS_116-110	WT	WT / WT	amp-seq
gRNA2_HS_116-111	WT	WT / WT	amp-seq
gRNA2_HS_116-112	WT	WT / WT	amp-seq
gRNA2_HS_116-113	WT	WT / WT	amp-seq
gRNA2_HS_116-114	WT	WT / WT	amp-seq
gRNA2_HS_116-115	WT	WT / WT	amp-seq
gRNA2_HS_116-116	WT	WT / WT	amp-seq
gRNA2_HS_116-117	WT	WT / WT	amp-seq
gRNA2_HS_116-118	WT	WT / WT	amp-seq

gRNA2_HS_116-119	WT	WT / WT	amp-seq
gRNA2_HS_116-120	WT	WT / WT	amp-seq
gRNA2_HS_116-122	WT	WT / WT	amp-seq
gRNA2_HS_116-123	WT	WT / WT	amp-seq
gRNA2_HS_116-124	WT	WT / WT	amp-seq
gRNA2_HS_116-125	WT	WT / WT	amp-seq
gRNA2_HS_116-126	WT	WT / WT	amp-seq
gRNA2_HS_116-127	WT	WT / WT	amp-seq
gRNA2_HS_116-129	WT	WT / WT	amp-seq
gRNA2_HS_116-132	WT	WT / WT	amp-seq
gRNA2_HS_116-133	WT	WT / WT	amp-seq
gRNA2_HS_116-134	WT	WT / WT	amp-seq
gRNA2_HS_116-135	WT	WT / WT	amp-seq
gRNA2_HS_116-137	WT	WT / WT	amp-seq
gRNA2_HS_116-138	WT	WT / WT	amp-seq
gRNA2_HS_116-140	WT	WT / WT	amp-seq
gRNA2_HS_116-141	WT	WT / WT	amp-seq
gRNA2_HS_116-142	WT	WT / WT	amp-seq
gRNA2_HS_116-143	WT	WT / WT	amp-seq
gRNA2_HS_116-145	WT	WT / WT	amp-seq
gRNA2_HS_116-147	WT	WT / WT	amp-seq
gRNA2_HS_116-148	WT	WT / WT	amp-seq
gRNA2_HS_116-149	WT	WT / WT	amp-seq

gRNA2_HS_116-150	WT	WT / WT	amp-seq
gRNA2_HS_116-151	WT	WT / WT	amp-seq
gRNA2_HS_116-152	WT	WT / WT	amp-seq
gRNA2_HS_116-153	WT	WT / WT	amp-seq
gRNA2_HS_116-154	WT	WT / WT	amp-seq
gRNA2_HS_116-156	WT	WT / WT	amp-seq
gRNA2_HS_116-157	WT	WT / WT	amp-seq
gRNA2_HS_116-158	WT	WT / WT	amp-seq
gRNA2_HS_116-160	WT	WT / WT	amp-seq
gRNA2_HS_116-161	WT	WT / WT	amp-seq
gRNA2_HS_116-162	WT	WT / WT	amp-seq
gRNA2_HS_116-163	WT	WT / WT	amp-seq
gRNA2_HS_116-164	WT	WT / WT	amp-seq
gRNA2_HS_116-165	WT	WT / WT	amp-seq
gRNA2_HS_116-166	WT	WT / WT	amp-seq
gRNA2_HS_116-167	WT	WT / WT	amp-seq
gRNA2_HS_116-168	WT	WT / WT	amp-seq
gRNA2_HS_116-169	WT	WT / WT	amp-seq
gRNA2_HS_116-170	WT	WT / WT	amp-seq
gRNA2_HS_116-171	WT	WT / WT	amp-seq
gRNA2_HS_116-172	WT	WT / WT	amp-seq
gRNA2_HS_116-173	WT	WT / WT	amp-seq
gRNA2_HS_116-174	WT	WT / WT	amp-seq

gRNA2_HS_116-175	WT	WT / WT	amp-seq
gRNA2_HS_116-176	WT	WT / WT	amp-seq
gRNA2_HS_116-177	WT	WT / WT	amp-seq
gRNA2_HS_116-178	WT	WT / WT	amp-seq
gRNA2_HS_116-179	WT	WT / WT	amp-seq
gRNA2_HS_116-180	WT	4bp del / WT	amp-seq
gRNA2_HS_116-181	WT	WT / WT	amp-seq
gRNA2_HS_116-182	WT	WT / WT	amp-seq
gRNA2_HS_116-183	WT	WT / WT	amp-seq
gRNA2_HS_116-184	WT	WT / WT	amp-seq
gRNA2_HS_116-185	WT	WT / WT	amp-seq
gRNA2_HS_116-186	WT	WT / WT	amp-seq
gRNA2_HS_116-187	WT	WT / WT	amp-seq
gRNA2_HS_116-188	WT	WT / WT	amp-seq
gRNA2_HS_116-189	WT	WT / WT	amp-seq
gRNA2_HS_116-190	WT	WT / WT	amp-seq
gRNA2_HS_116-191	WT	WT / WT	amp-seq
gRNA2_HS_116-192	WT	WT / WT	amp-seq
gRNA2_HS_116-193	WT	WT / WT	amp-seq
gRNA2_HS_116-194	WT	WT / WT	amp-seq
gRNA2_HS_116-195	WT	WT / WT	amp-seq
gRNA2_HS_116-196	WT	WT / WT	amp-seq
gRNA2_HS_116-198	WT	WT / WT	amp-seq

gRNA2_HS_116-199	WT	WT / WT	amp-seq
gRNA2_HS_116-200	WT	WT / WT	amp-seq
gRNA2_HS_116-201	WT	WT / WT	amp-seq
gRNA2_HS_116-202	WT	WT / WT	amp-seq
gRNA2_HS_116-203	WT	WT / WT	amp-seq
gRNA2_HS_116-204	WT	WT / WT	amp-seq
gRNA2_HS_116-205	WT	WT / WT	amp-seq
gRNA2_HS_116-206	WT	WT / WT	amp-seq
gRNA2_HS_116-207	WT	WT / WT	amp-seq
gRNA2_HS_116-208	WT	WT / WT	amp-seq
gRNA2_HS_116-209	WT	WT / WT	amp-seq
gRNA2_HS_116-210	WT	WT / WT	amp-seq
gRNA2_HS_116-211	WT	WT / WT	amp-seq
gRNA2_HS_116-212	WT	WT / WT	amp-seq
gRNA2_HS_116-213	WT	WT / WT	amp-seq
gRNA2_HS_116-214	WT	WT / WT	amp-seq
gRNA2_HS_116-215	WT	4bp del / WT	amp-seq
gRNA2_HS_116-216	WT	WT / WT	amp-seq
gRNA2_HS_116-217	WT	4bp del / WT	amp-seq
gRNA2_HS_116-218	WT	WT / WT	amp-seq
gRNA2_HS_116-219	WT	WT / WT	amp-seq
gRNA2_HS_116-220	WT	WT / WT	amp-seq
gRNA2_HS_116-221	WT	WT / WT	amp-seq

gRNA2_HS_116-222	WT	WT / WT	amp-seq
gRNA2_HS_116-223	WT	WT / WT	amp-seq
gRNA2_HS_116-224	WT	WT / WT	amp-seq
gRNA2_HS_116-225	WT	WT / WT	amp-seq
gRNA2_HS_116-226	WT	WT / WT	amp-seq
gRNA2_HS_116-227	WT	WT / WT	amp-seq
gRNA2_HS_116-228	WT	WT / WT	amp-seq
gRNA2_HS_116-229	WT	WT / WT	amp-seq
gRNA2_HS_116-230	WT	WT / WT	amp-seq
gRNA2_HS_116-231	WT	WT / WT	amp-seq
gRNA2_HS_116-232	WT	WT / WT	amp-seq
gRNA2_HS_116-233	WT	WT / WT	amp-seq
gRNA2_HS_116-234	WT	WT / WT	amp-seq
gRNA2_HS_116-235	WT	WT / WT	amp-seq
gRNA2_HS_116-236	WT	WT / WT	amp-seq
gRNA2_HS_116-237	WT	WT / WT	amp-seq
gRNA2_HS_116-238	WT	WT / WT	amp-seq
gRNA2_HS_116-239	WT	WT / WT	amp-seq
gRNA2_HS_116-240	WT	WT / WT	amp-seq
gRNA2_HS_116-241	WT	WT / WT	amp-seq
gRNA2_HS_116-242	WT	WT / WT	amp-seq
gRNA2_HS_116-243	WT	WT / WT	amp-seq
gRNA2_HS_116-244	WT	4bp del / WT	amp-seq

gRNA2_HS_116-245	WT	WT / WT	amp-seq
gRNA2_HS_116-246	WT	WT / WT	amp-seq
gRNA2_HS_116-247	WT	WT / WT	amp-seq
gRNA2_HS_116-248	WT	WT / WT	amp-seq
gRNA12_room_54-35	WT	4bp del / 4bp del	Sanger
gRNA12_room_54-36	WT	9bp del / 9bp del	Sanger
gRNA12_room_54-37	WT	3bp del / 3bp del	Sanger
gRNA12_room_54-40	WT	9bp del / 9bp del	Sanger
gRNA12_room_54-42	WT	37bp del / WT	Sanger
gRNA12_room_54-43	WT	WT / WT	Sanger
gRNA12_room_54-44	WT	WT / WT	Sanger
gRNA12_room_54-45	WT	WT / WT	Sanger
gRNA12_room_54-46	WT	WT / WT	Sanger
gRNA12_room_54-47	WT	WT / WT	Sanger
gRNA12_room_54-48	WT	WT / WT	Sanger
gRNA12_room_54-49	WT	17bp del / 36bp del	Sanger
gRNA12_room_54-50	WT	WT / WT	Sanger
gRNA12_room_54-52	WT	WT / WT	Sanger
gRNA12_room_54-53	WT	WT / WT	Sanger
gRNA12_room_54-54	WT	3bp del / WT	Sanger
gRNA12_room_54-55	WT	WT / WT	Sanger
gRNA12_room_54-58	WT	17bp del / 36bp del	Sanger
gRNA12_room_54-59	WT	9bp del / 9bp del	Sanger
gRNA12_room_54-60	WT	WT / WT	Sanger

gRNA12_room_54-61	WT	17bp del / 36bp del	Sanger
gRNA12_room_54-62	WT	WT / WT	Sanger
gRNA12_room_54-63	WT	WT / WT	Sanger
gRNA12_room_54-64	WT	WT / WT	Sanger
gRNA12_room_54-65	WT	WT / WT	Sanger
gRNA12_room_54-66	WT	WT / WT	Sanger
gRNA12_room_54-68	WT	12bp del / 12bp del	Sanger
gRNA12_room_54-69	WT	3bp del / 3bp del	Sanger
gRNA12_room_54-70	WT	WT / WT	Sanger
gRNA12_room_54-71	WT	WT / WT	Sanger
gRNA12_room_54-72	WT	10bp del / WT	Sanger
gRNA12_room_54-73	WT	WT / WT	Sanger
gRNA12_room_54-74	WT	WT / WT	Sanger
gRNA12_room_54-75	WT	WT / WT	Sanger
gRNA12_room_54-76	WT	WT / WT	Sanger
gRNA12_room_54-77	WT	WT / WT	Sanger
gRNA12_room_54-79	WT	4bp del / WT	Sanger
gRNA12_room_54-80	WT	WT / WT	Sanger
gRNA12_room_54-84	WT	9bp del / WT	Sanger
gRNA12_room_54-86	WT	WT / WT	Sanger
gRNA12_room_54-90	WT	WT / WT	Sanger
gRNA12_room_54-91	WT	9bp del / 9bp del	Sanger
gRNA12_room_54-94	WT	WT / WT	Sanger
gRNA12_room_54-95	WT	WT / WT	Sanger

gRNA12_room_54-97	WT	3bp del / 3bp del	Sanger
gRNA12_room_54-98	WT	WT / WT	Sanger
gRNA12_room_54-104	WT	WT / WT	Sanger
gRNA12_room_54-105	WT	7bp del / 7bp del	Sanger
gRNA12_room_54-106	WT	WT / WT	Sanger
gRNA12_room_54-108	WT	WT / WT	Sanger
gRNA12_room_54-109	WT	17bp del / 21bp del	Sanger
gRNA12_room_54-110	WT	3bp del / WT	Sanger
gRNA12_room_54-111	WT	WT / WT	Sanger
gRNA12_room_54-112	WT	7bp del / 7bp del	Sanger
gRNA12_room_54-113	WT	26bp del / WT	Sanger
gRNA12_room_54-114	WT	WT / WT	Sanger
gRNA12_room_54-115	WT	WT / WT	Sanger
gRNA12_room_54-116	WT	WT / WT	Sanger
gRNA12_room_54-117	WT	WT / WT	Sanger
gRNA12_room_54-119	WT	WT / WT	Sanger
gRNA12_room_54-120	WT	11bp del / 11bp del	Sanger
gRNA12_room_54-121	WT	10bp del / WT	Sanger
gRNA12_room_54-122	WT	WT / WT	Sanger
gRNA12_room_54-123	WT	WT / WT	Sanger
gRNA12_room_54-124	WT	10bp del / WT	Sanger
gRNA12_room_54-125	WT	WT / WT	Sanger
gRNA12_room_54-126	WT	WT / WT	Sanger
gRNA12_room_54-127	WT	WT / WT	Sanger

gRNA12_room_54-128	WT	4bp del / WT	Sanger
gRNA12_room_54-130	WT	38bp del / WT	Sanger
gRNA12_room_54-132	WT	WT / WT	Sanger
gRNA12_room_54-133	WT	WT / WT	Sanger
gRNA12_room_54-134	WT	WT / WT	Sanger
gRNA12_room_54-135	WT	WT / WT	Sanger
gRNA12_room_54-136	WT	WT / WT	Sanger
gRNA12_room_54-138	WT	9bp del / 9bp del	Sanger
gRNA12_room_54-139	WT	2bp del / 2bp del	Sanger
gRNA12_room_54-140	WT	WT / WT	Sanger
gRNA12_room_54-141	WT	WT / WT	Sanger
gRNA12_room_54-142	WT	37bp del / WT	Sanger
gRNA12_room_54-143	WT	WT / WT	Sanger
gRNA12_room_54-144	WT	WT / WT	Sanger
gRNA12_room_54-145	WT	WT / WT	Sanger
gRNA12_room_54-146	WT	WT / WT	Sanger
gRNA12_room_54-148	WT	17bp del / 36bp del	Sanger
gRNA12_room_54-149	WT	WT / WT	Sanger
gRNA12_room_54-152	WT	WT / WT	Sanger
gRNA12_room_54-153	WT	WT / WT	Sanger
gRNA12_room_54-154	WT	WT / WT	Sanger
gRNA12_room_54-155	WT	WT / WT	Sanger
gRNA12_room_54-156	WT	11bp del / 11bp del	Sanger
gRNA12_room_54-157	WT	5bp del / WT	Sanger

gRNA12_room_54-158	WT	37bp del / WT	Sanger
gRNA12_room_54-159	WT	62bp del / 62bp del	Sanger
gRNA12_room_54-160	WT	WT / WT	Sanger
gRNA12_room_54-161	WT	8bp del / WT	Sanger
gRNA12_room_54-162	WT	8bp del / WT	Sanger
gRNA12_room_54-163	WT	WT / WT	Sanger
gRNA12_room_54-164	WT	4bp del / WT	Sanger
gRNA12_room_54-165	WT	WT / WT	Sanger
gRNA12_room_54-166	WT	2bp del / WT	Sanger
gRNA12_room_54-167	WT	WT / WT	Sanger
gRNA12_room_54-168	WT	WT / WT	Sanger
gRNA12_room_54-169	WT	WT / WT	Sanger
gRNA12_room_54-170	WT	WT / WT	Sanger
gRNA12_room_54-171	WT	WT / WT	Sanger
gRNA12_room_54-172	WT	WT / WT	Sanger
gRNA12_room_54-173	WT	WT / WT	Sanger
gRNA12_room_54-175	WT	WT / WT	Sanger
gRNA12_room_54-176	WT	WT / WT	Sanger
gRNA12_room_54-178	WT	WT / WT	Sanger
gRNA12_room_54-179	WT	WT / WT	Sanger
gRNA12_room_54-180	WT	WT / WT	Sanger
gRNA12_room_54-181	WT	WT / WT	Sanger
gRNA12_room_54-182	WT	WT / WT	Sanger
gRNA12_room_54-183	WT	WT / WT	Sanger

gRNA12_room_54-184	WT	5bp del / WT	Sanger
gRNA12_room_54-185	WT	WT / WT	Sanger
gRNA12_room_54-186	WT	WT / WT	Sanger
gRNA12_room_54-190	WT	9bp del / WT	Sanger
gRNA12_room_54-191	WT	15bp del / WT	Sanger
gRNA12_room_54-192	WT	WT / WT	Sanger
gRNA12_room_54-193	WT	WT / WT	Sanger
gRNA12_room_54-194	WT	7bp del / 7bp del	Sanger
gRNA12_room_54-195	WT	WT / WT	Sanger
gRNA12_room_54-196	WT	WT / WT	Sanger
gRNA12_room_54-197	WT	10bp del / 10bp del	Sanger
gRNA12_room_54-198	WT	10bp del / WT	Sanger
gRNA12_room_54-199	WT	WT / WT	Sanger
gRNA12_room_54-200	WT	2bp del / 2bp del	Sanger
gRNA12_room_54-202	WT	WT / WT	Sanger
gRNA12_room_54-205	WT	WT / WT	Sanger
gRNA12_room_54-208	WT	WT / WT	Sanger
gRNA12_room_54-209	WT	WT / WT	Sanger
gRNA12_room_54-211	WT	WT / WT	Sanger
gRNA12_room_54-212	WT	WT / WT	Sanger
gRNA12_room_54-213	WT	3bp del / WT	Sanger
gRNA12_room_54-214	WT	WT / WT	Sanger
gRNA12_room_54-215	WT	4bp del / 4bp del	Sanger
gRNA12_room_54-216	WT	WT / WT	Sanger

gRNA12_room_54-217	WT	3bp del / WT	Sanger
gRNA12_room_54-218	WT	WT / WT	Sanger
gRNA12_room_54-219	WT	17bp del / 17bp del	Sanger
gRNA12_room_54-220	WT	4bp del / WT	Sanger
gRNA12_room_54-221	WT	7bp del / 10bp del	Sanger
gRNA12_room_54-222	WT	WT / WT	Sanger
gRNA12_room_54-223	WT	WT / WT	Sanger
gRNA12_room_54-224	WT	WT / WT	Sanger
gRNA12_room_69-2	WT	15bp del / 15bp del	Sanger
gRNA12_room_69-4	WT	27bp del / WT	Sanger
gRNA12_room_69-5	WT	5bp del / 24bp del	Sanger
gRNA12_room_69-7	WT	WT / WT	Sanger
gRNA12_room_69-8	WT	7bp del / 15bp del	Sanger
gRNA12_room_69-9	WT	1bp del / 1bp del	Sanger
gRNA12_room_69-10	WT	WT / WT	Sanger
gRNA12_room_69-12	WT	5bp del / 15bp del	Sanger
gRNA12_room_69-14	WT	22bp del / 22bp del	Sanger
gRNA12_room_69-16	WT	10bp del / 10bp del	Sanger
gRNA12_room_69-17	WT	1bp del / 17bp del	Sanger
gRNA12_room_69-18	WT	11bp del / WT	Sanger
gRNA12_room_69-21	WT	5bp del / 24bp del	Sanger
gRNA12_room_69-22	WT	17bp del / WT	Sanger
gRNA12_room_69-23	WT	WT / WT	Sanger
gRNA12_room_69-28	WT	5bp del / 4bp del	Sanger

gRNA12_room_69-30	WT	29bp del / 29bp del	Sanger
gRNA12_room_69-33	WT	WT / WT	Sanger
gRNA12_room_69-34	WT	5bp del / 5bp del	Sanger
gRNA12_room_69-37	WT	WT / WT	Sanger
gRNA12_room_69-41	WT	WT / WT	Sanger
gRNA12_room_69-44	WT	5bp del / 15bp del	Sanger
gRNA12_room_69-51	WT	5bp del / WT	Sanger
gRNA12_room_69-53	WT	15bp del / 15bp del	Sanger
gRNA12_room_69-55	WT	5bp del / 5bp del	Sanger
gRNA12_room_69-59	WT	24bp del / 24bp del	Sanger
gRNA12_room_69-63	WT	5bp del / 24bp del	Sanger
gRNA12_room_69-66	WT	WT / WT	Sanger
gRNA12_room_69-67	WT	WT / WT	Sanger
gRNA12_room_69-70	WT	7bp del / WT	Sanger
gRNA12_room_69-73	WT	17bp del / WT	Sanger
gRNA12_room_69-74	WT	5bp del / 10bp del	Sanger
gRNA12_room_69-78	WT	WT / WT	Sanger
gRNA12_room_69-80	WT	WT / WT	Sanger
gRNA12_room_69-87	WT	13bp del / 14bp del	Sanger
gRNA12_room_69-88	WT	11bp del / WT	Sanger
gRNA12_room_69-91	WT	WT / WT	Sanger
gRNA12_room_69-92	WT	5bp del / 10bp del	Sanger
gRNA12_room_69-93	WT	WT / WT	Sanger
gRNA12_room_69-94	WT	WT / WT	Sanger

gRNA12_room_69-95	WT	3bp del / 23bp del	Sanger
gRNA12_room_69-101	WT	7bp del / WT	Sanger
gRNA12_room_69-108	WT	WT / WT	Sanger
gRNA12_room_69-111	WT	WT / WT	Sanger
gRNA12_room_69-115	WT	WT / WT	Sanger
gRNA12_room_69-117	WT	10bp del / WT	Sanger
gRNA12_room_69-120	WT	17bp del / WT	Sanger
gRNA12_room_69-123	WT	WT / WT	Sanger
gRNA12_room_69-124	WT	15bp del / 15bp del	Sanger
gRNA12_room_69-125	WT	9bp del / 22bp del	Sanger
gRNA12_room_69-126	WT	17bp del / 17bp del	Sanger
gRNA12_room_69-127	WT	WT / WT	Sanger
gRNA12_room_69-129	WT	WT / WT	Sanger
gRNA12_room_69-130	WT	WT / WT	Sanger
gRNA12_room_69-131	WT	4bp del / 5bp del	Sanger
gRNA12_room_69-132	WT	WT / WT	Sanger
gRNA12_room_69-134	WT	WT / WT	Sanger
gRNA12_room_69-137	WT	4bp del / WT	Sanger
gRNA12_room_69-142	WT	15bp del / 5bp del	Sanger
gRNA12_room_69-148	WT	WT / WT	Sanger
gRNA12_room_69-149	WT	22bp del / 22bp del	Sanger
gRNA12_room_69-150	WT	WT / WT	Sanger
gRNA12_room_69-151	WT	WT / WT	Sanger
gRNA12_room_69-154	WT	11bp del / 11bp del	Sanger

gRNA12_room_69-157	WT	WT / WT	Sanger
gRNA12_room_69-158	WT	11bp del / WT	Sanger
gRNA12_room_69-159	WT	18bp del / WT	Sanger
gRNA12_room_69-163	WT	WT / WT	Sanger
gRNA12_room_69-170	WT	7bp del / 26bp del	Sanger
gRNA12_room_69-171	WT	3bp del / 5bp del	Sanger
gRNA12_room_69-172	WT	10bp del / 10bp del	Sanger
gRNA12_room_69-175	WT	5bp del / WT	Sanger
gRNA12_room_69-183	WT	WT / WT	Sanger
gRNA12_room_69-185	WT	5bp del / 5bp del	Sanger
gRNA12_room_69-188	WT	1bp del / WT	Sanger

Supplementary Table 2: Phenotype and genotype of the progenies from three individual TRV-

infected plants. The plant ID column indicates an internal identifier for each plant sampled. The

Phenotype column indicates whether the plant was green (WT) or albino. The Genotype (allele1

/ allele2) column lists the alleles of the *AtPDS3* gene determined by amp-seq or Sanger

Sequencing. WT is an abbreviation for wild type, bp is an abbreviation for base pair, and del is

an abbreviation for deletion.

Sample	Reads count	Unmapped reads count	Mapping rate (%)	Coverage
WT-rep1	331996121	2122433	99.36	817.69
WT-rep2	314298698	1969352	99.37	774.20
gRNA2_HS_116-2	315840117	1628611	99.48	778.87
gRNA2_HS_116-32	357227432	2397598	99.33	879.55
gRNA2_HS_116-43	265745567	1688428	99.36	654.55

571 **Supplementary Table 3: Whole genome sequencing read count and mapping rate of WT**

572 **control and albino mutant plants.**

Sample	GATK	Strealk2	GATK+Strealk2	Filter with WT	Filter by depth > 30	Manually check
WT	40852	24406	NA	NA	NA	NA
albino-plant2	36346	20659	16296	28	25	5
albino-plant32	36342	20991	16629	35	30	5
albino-plant43	36989	21928	17111	80	71	4

574 **Supplementary Table 4: WGS variant detection of WT control and albino mutant plants using**

575 **GATK and Strelka2.**

DNA	Chromosome	Location	Direction	Mismatches	Variants
TTcATtAGGCAAATTCaCtGC	chr1	1799061	+	4	No SNP/Indel
TTGATAAGGaAgATTCGtCtC	chr1	21688156	+	4	No SNP/Indel
TTGATAAGcCAtATTCtCCcC	chr1	28196564	+	4	No SNP/Indel
TTGATcAGGaAAATTCGCgGa	chr1	29172329	-	4	No SNP/Indel
TTGATcAGGCAAATTCGgaGC	chr2	2183162	+	3	No SNP/Indel
aaGcTAAGGCAAATcCGCCGC	chr2	10574134	-	4	No SNP/Indel
TTGATAAGaacAtTTCGCCGC	chr2	11679602	-	4	No SNP/Indel
gTcATAAGtCAcATTCGCCGC	chr2	12173059	+	4	No SNP/Indel
TTGATAAGtCAAAgTCtaCGC	chr2	13554186	-	4	No SNP/Indel
TTGATActGCAAAgTCGCCtC	chr2	16402169	+	4	No SNP/Indel
TacAgAAGGCAAATTCcCCGC	chr3	381121	-	4	No SNP/Indel
TTGATAAtGCAAcTTCtCCGt	chr3	527026	-	4	No SNP/Indel
TTcATcAGGCAtATTCGCCGt	chr3	13200296	-	4	No SNP/Indel
TTGgTAAGGCAAATTCttaGC	chr4	8560337	+	4	No SNP/Indel
gTGAaAAcGCAAATTCGCgGC	chr5	3918356	+	4	No SNP/Indel
TTGATAAGtCAAAaTCcCCaC	chr5	10937249	-	4	No SNP/Indel
TTGATAAtGCAAATTacCCGt	chr5	18593314	-	4	No SNP/Indel
TTGATAAGGCAAtcTCtaCGC	chr5	26138716	+	4	No SNP/Indel

Supplementary Table 5: Off-target editing analysis using Cas-OFFinder. The DNA column

contains the sequence of the potential off-target site, with lower case letters indicating a

mismatch compared to the actual target site. The Chromosome, Location, and Direction

columns indicate where in the *Arabidopsis* genome that potential off-target site is located. The

mismatch column indicates the number of mismatches in the potential off-target site relative to

the actual target site sequence. The Variants column lists the off-target editing result for that

site, with every off-target site analyzed as wild type (No SNP/Indel).

Plasmid	Description
pMK003	Cloning vector for PaqCI Golden Gate assembly of ISDra2 guides into ISDra2 plant expression vector
pMK025	Cloning vector for PaqCI Golden Gate assembly of ISYmu1 guides into ISYmu1 plant expression vector
pMK024	Cloning vector for PaqCI Golden Gate assembly of ISAam1 guides into ISAam1 plant expression vector
pMK026	ISDra2 g12 cloned into pMK003
pMK027	ISDra2 g13 cloned into pMK003
pMK028	ISDra2 g8 cloned into pMK003
pMK029	ISDra2 g14 cloned into pMK003
pMK030	ISDra2 g4 cloned into pMK003
pMK031	ISDra2 g15 cloned into pMK003
pMK032	ISDra2 g17 cloned into pMK003
pMK033	ISDra2 g18 cloned into pMK003
pMK034	ISDra2 g5 cloned into pMK003
pMK035	ISDra2 g20 cloned into pMK003
pMK036	ISDra2 g22 cloned into pMK003
pMK037	ISDra2 g1 cloned into pMK003
pMK038	ISDra2 g2 cloned into pMK003
pMK039	ISDra2 g3 cloned into pMK003
pMK040	ISDra2 g16 cloned into pMK003

pMK041	ISDra2 g19 cloned into pMK003
pMK042	ISDra2 g9 cloned into pMK003
pMK043	ISDra2 g21 cloned into pMK003
pMK044	ISDra2 g11 cloned into pMK003
pMK045	ISDra2 g7 cloned into pMK003
pMK060	ISYmu1 g1 cloned into pMK025
pMK061	ISYmu1 g2 cloned into pMK025
pMK062	ISYmu1 g3 cloned into pMK025
pMK063	ISYmu1 g4 cloned into pMK025
pMK064	ISYmu1 g5 cloned into pMK025
pMK065	ISYmu1 g7 cloned into pMK025
pMK066	ISYmu1 g8 cloned into pMK025
pMK067	ISYmu1 g9 cloned into pMK025
pMK068	ISYmu1 g10 cloned into pMK025
pMK070	ISYmu1 g12 cloned into pMK025
pMK050	ISAam1 g11 cloned into pMK024
pMK051	ISAam1 g12 cloned into pMK024
pMK052	ISAam1 g13 cloned into pMK024
pMK053	ISAam1 g14 cloned into pMK024
pMK054	ISAam1 g15 cloned into pMK024
pMK055	ISAam1 g16 cloned into pMK024
pMK056	ISAam1 g17 cloned into pMK024

pTW2065	ISYmu1 gRNA2 TRV2 Architecture_A
pTW2066	ISYmu1 gRNA2 TRV2 Architecture_B
pTW2082	ISYmu1 gRNA12 TRV2 Architecture_B

585 **Supplementary Table 6: Plasmids used in this study.** The Plasmid column lists the name of

586 each plasmid used. The Description column provides a TnpB and gRNA description of each

587 plasmid used, along with the intermediate plant expression vector that was used to create it.

Oligo name	Oligo sequence	Oligo description
MK_29476	TCAAgcgttggagcatataacaga	cloning ISDra2 gRNA_20_bp_gRNA_12_Top
Trevor_28675	GGCCtctgttatatgctccaacgc	cloning ISDra2 gRNA_20_bp_gRNA_12_Bottom
MK_29477	TCAAttacgaattgatgaccatat	cloning ISDra2 gRNA_20_bp_gRNA_1_Top
Trevor_28677	GGCCatatggtcatcaattcgtaa	cloning ISDra2 gRNA_20_bp_gRNA_1_Bottom
MK_29478	TCAAaaggcaaattcgccgcagaa	cloning ISDra2 gRNA_20_bp_gRNA_2_Top
Trevor_28679	GGCCttctgcggcgaatttgcctt	cloning ISDra2 gRNA_20_bp_gRNA_2_Bottom
MK_29479	TCAAtaaagagaggaaattgcagg	cloning ISDra2 gRNA_20_bp_gRNA_13_Top
Trevor_28681	GGCCcctgcaatttcctctcttta	cloning ISDra2 gRNA_20_bp_gRNA_13_Bottom
MK_29480	TCAAtggtctcactttccgaatta	cloning ISDra2 gRNA_20_bp_gRNA_8_Top
Trevor_28683	GGCCtaattcggaaagtgagacca	cloning ISDra2 gRNA_20_bp_gRNA_8_Bottom
MK_29481	TCAAtcacattaagcctagaaact	cloning ISDra2 gRNA_20_bp_gRNA_3_Top
Trevor_28685	GGCCagtttctaggcttaatgtga	cloning ISDra2 gRNA_20_bp_gRNA_3_Bottom
MK_29482	TCAAggtagagctgataagatata	cloning ISDra2 gRNA_20_bp_gRNA_14_Top
Trevor_28687	GGCCtatatcttatcagctctacc	cloning ISDra2 gRNA_20_bp_gRNA_14_Bottom
MK_29483	TCAAcccaagttctccaaataaat	cloning ISDra2 gRNA_20_bp_gRNA_4_Top
Trevor_28689	GGCCatttatttggagaacttggg	cloning ISDra2 gRNA_20_bp_gRNA_4_Bottom
MK_29484	TCAAaaaattggattaatgtgcac	cloning ISDra2 gRNA_20_bp_gRNA_15_Top
Trevor_28691	GGCCgtgcacattaatccaatttt	cloning ISDra2 gRNA_20_bp_gRNA_15_Bottom
MK_29485	TCAAcaatacaaataaatacatgc	cloning ISDra2 gRNA_20_bp_gRNA_17_Top
Trevor_28693	GGCCgcatgtatttatttgtattg	cloning ISDra2 gRNA_20_bp_gRNA_17_Bottom
MK_29486	TCAAcaattcaagctaattataga	cloning ISDra2 gRNA_20_bp_gRNA_18_Top
Trevor_28695	GGCCtctataattagcttgaattg	cloning ISDra2 gRNA_20_bp_gRNA_18_Bottom
MK_29487	TCAAtactattaaatgtcaaaatc	cloning ISDra2 gRNA_20_bp_gRNA_16_Top

Trevor_28697	GGCCgattttgacatttaatagta	cloning ISDra2 gRNA_20_bp_gRNA_16_Bottom
MK_29488	TCAAtacccatcctaaagtatggg	cloning ISDra2 gRNA_20_bp_gRNA_5_Top
Trevor_28699	GGCCcccatactttaggatgggta	cloning ISDra2 gRNA_20_bp_gRNA_5_Bottom
MK_29489	TCAAgagcttaacttggtagagta	cloning ISDra2 gRNA_20_bp_gRNA_19_Top
Trevor_28701	GGCCtactctaccaagttaagctc	cloning ISDra2 gRNA_20_bp_gRNA_19_Bottom
MK_29490	TCAAttgtcagctttcttatggat	cloning ISDra2 gRNA_20_bp_gRNA_20_Top
Trevor_28703	GGCCatccataagaaagctgacaa	cloning ISDra2 gRNA_20_bp_gRNA_20_Bottom
MK_29491	TCAAagctttgaaccggtttcttc	cloning ISDra2 gRNA_20_bp_gRNA_9_Top
Trevor_28705	GGCCgaagaaaccggttcaaagct	cloning ISDra2 gRNA_20_bp_gRNA_9_Bottom
MK_29492	TCAAgttgattaacttgtactacc	cloning ISDra2 gRNA_20_bp_gRNA_21_Top
Trevor_28707	GGCCggtagtacaagttaatcaac	cloning ISDra2 gRNA_20_bp_gRNA_21_Bottom
MK_29493	TCAAtaacttgtactacctcatcc	cloning ISDra2 gRNA_20_bp_gRNA_22_Top
Trevor_28709	GGCCggatgaggtagtacaagtta	cloning ISDra2 gRNA_20_bp_gRNA_22_Bottom
MK_29494	TCAAcgaaaactgaagaacacata	cloning ISDra2 gRNA_20_bp_gRNA_11_Top
Trevor_28711	GGCCtatgtgttcttcagttttcg	cloning ISDra2 gRNA_20_bp_gRNA_11_Bottom
MK_29495	TCAAaaattcaacatctttctcta	cloning ISDra2 gRNA_20_bp_gRNA_7_Top
Trevor_28713	GGCCtagagaaagatgttgaattt	cloning ISDra2 gRNA_20_bp_gRNA_7_Bottom
MK_29758	TCAAttacgaattgatgacc	cloning ISYmu1 gRNA_16_bp_gRNA_1_Top
MK_29759	GGCCggtcatcaattcgtaa	cloning ISYmu1 gRNA_16_bp_gRNA_1_Bottom
MK_29760	TCAAaaggcaaattcgccgc	cloning ISYmu1 gRNA_16_bp_gRNA_2_Top
MK_29761	GGCCgcggcgaatttgcctt	cloning ISYmu1 gRNA_16_bp_gRNA_2_Bottom
MK_29762	TCAAtcacattaagcctaga	cloning ISYmu1 gRNA_16_bp_gRNA_3_Top
MK_29763	GGCCtctaggcttaatgtga	cloning ISYmu1 gRNA_16_bp_gRNA_3_Bottom
MK_29764	TCAAcccaagttctccaaat	cloning ISYmu1 gRNA_16_bp_gRNA_4_Top

MK_29765	GGCCatttggagaacttggg	cloning ISYmu1 gRNA_16_bp_gRNA_4_Bottom
MK_29766	TCAAtacccatcctaaagta	cloning ISYmu1 gRNA_16_bp_gRNA_5_Top
MK_29767	GGCCtactttaggatgggta	cloning ISYmu1 gRNA_16_bp_gRNA_5_Bottom
MK_29770	TCAAaaattcaacatctttc	cloning ISYmu1 gRNA_16_bp_gRNA_7_Top
MK_29771	GGCCgaaagatgttgaattt	cloning ISYmu1 gRNA_16_bp_gRNA_7_Bottom
MK_29772	TCAAtggtctcactttccga	cloning ISYmu1 gRNA_16_bp_gRNA_8_Top
MK_29773	GGCCtcggaaagtgagacca	cloning ISYmu1 gRNA_16_bp_gRNA_8_Bottom
MK_29774	TCAAagctttgaaccggttt	cloning ISYmu1 gRNA_16_bp_gRNA_9_Top
MK_29775	GGCCaaaccggttcaaagct	cloning ISYmu1 gRNA_16_bp_gRNA_9_Bottom
MK_29776	TCAAtaacttgtactacctc	cloning ISYmu1 gRNA_16_bp_gRNA_10_Top
MK_29777	GGCCgaggtagtacaagtta	cloning ISYmu1 gRNA_16_bp_gRNA_10_Bottom
MK_29780	TCAAgcgttggagcatataa	cloning ISYmu1 gRNA_16_bp_gRNA_12_Top
MK_29781	GGCCttatatgctccaacgc	cloning ISYmu1 gRNA_16_bp_gRNA_12_Bottom
MK_29818	TCACcaattcatctggtatc	cloning ISAam1 gRNA_16_bp_gRNA_11_Top
MK_29819	GGCCgataccagatgaattg	cloning ISAam1 gRNA_16_bp_gRNA_11_Bottom
MK_29820	TCACccaagaacaagcctta	cloning ISAam1 gRNA_16_bp_gRNA_12_Top
MK_29821	GGCCtaaggcttgttcttgg	cloning ISAam1 gRNA_16_bp_gRNA_12_Bottom
MK_29822	TCACgttttgtcctcttctc	cloning ISAam1 gRNA_16_bp_gRNA_13_Top
MK_29823	GGCCgagaagaggacaaaac	cloning ISAam1 gRNA_16_bp_gRNA_13_Bottom
MK_29824	TCACcaatttacctatctta	cloning ISAam1 gRNA_16_bp_gRNA_14_Top
MK_29825	GGCCtaagataggtaaattg	cloning ISAam1 gRNA_16_bp_gRNA_14_Bottom
MK_29826	TCACcacataattgaaaaga	cloning ISAam1 gRNA_16_bp_gRNA_15_Top
MK_29827	GGCCtcttttcaattatgtg	cloning ISAam1 gRNA_16_bp_gRNA_15_Bottom
MK_29828	TCACaatttgttacacaact	cloning ISAam1 gRNA_16_bp_gRNA_16_Top

MK_29829	GGCCagttgtgtaacaaatt	cloning ISAam1 gRNA_16_bp_gRNA_16_Bottom
MK_29830	TCACtttgtgtggtatttaa	cloning ISAam1 gRNA_16_bp_gRNA_17_Top
MK_29831	GGCCttaaataccacacaaa	cloning ISAam1 gRNA_16_bp_gRNA_17_Bottom

589 **Supplementary Table 7: Oligos used for cloning gRNA sequences into intermediate TnpB plant**

590 **expression vectors.** The Oligo name column is an internal name for each oligo used. The Oligo

591 sequence column lists the sequence of each oligo used for cloning the gRNA. The Oligo

592 description column provides details for cloning each TnpB gRNA target site.

Plasmid	PCR template	Primer 1	Primer 2	Reaction #
pTW2065	pDK3888	gaatttcctttaccattgacgtcagtgtcgttggtagcat	tgtaatccatCTCGTTAACTCGGGTAAGTGATACA	1
pTW2065	pMK061	AGTTAACGAGatggattacaaggatgatgatgataagga	CCAACTGAGCTACGGgcggcgaatttgccttTTGA	2
pTW2065	pDK3888	aggcaaattcgccgcCCGTAGCTCAGTTGGTTAGAGC	TGGTCACCTGTAATTCACACGTGGTGGTGGTGGTGGT	3
pTW2066	pDK3888	gaatttcctttaccattgacgtcagtgtcgttggtagcat	tgtaatccatCTCGTTAACTCGGGTAAGTGATACA	1
pTW2066	pMK061	AGTTAACGAGatggattacaaggatgatgatgataagga	TCTAACCAACTGAGCTACGGgtcccattcgccatgcc	2
pTW2066	pDK3888	catggcgaatgggacCCGTAGCTCAGTTGGTTAGAGC	TGGTCACCTGTAATTCACACGTGGTGGTGGTGGTGGT	3
pTW2082	pDK3888	gaatttcctttaccattgacgtcagtgtcgttggtagcat	tgtaatccatCTCGTTAACTCGGGTAAGTGATACA	1
pTW2082	pMK070	AGTTAACGAGatggattacaaggatgatgatgataagga	TCTAACCAACTGAGCTACGGgtcccattcgccatgcc	2
pTW2082	pDK3888	catggcgaatgggacCCGTAGCTCAGTTGGTTAGAGC	TGGTCACCTGTAATTCACACGTGGTGGTGGTGGTGGT	3

Supplementary Table 8: Cloning TRV Cargo Architecture_A and Architecture_B plasmids. The

Plasmid column lists the name of each plasmid created. The PCR template column lists the

name of each plasmid used for PCR DNA template. The Primer 1 and Primer 2 columns provide

the sequence used to amplify fragments using the corresponding PCR template. The reaction

column indicates the individual reaction for each PCR. The three PCR reactions, along with the

restriction enzyme digested pDK3888, were used for NEB Hifi assembly to create the TRV2

vector listed in the Plasmid column.

TnpB	Guide	F primer	R primer	Sequencing type
ISDra2	gRNA1	ACACTCTTTCCCTACACGACGCTCTTCCGATCTaccaaattacgttgagatgcatggtc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTgctctttttgttgcttacgaaacatgt	amp-seq
ISDra2	gRNA2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgaagcagttgtgagttaagttggaga	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtttgtcttaagcgcttgagaagtgg	amp-seq
ISDra2	gRNA3	ACACTCTTTCCCTACACGACGCTCTTCCGATCTatgtcttcagtcttgtgctacacacc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaaacggaacaaacaagaacctaacct	amp-seq
ISDra2	gRNA4	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtctcttcatgttcttgttggaagtcgg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTgtgctggtaggacatctgggaagt	amp-seq
ISDra2	gRNA5	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgacgtcaggaagaacatggtcatttg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTagaacatttcagcgctaatgctacaa	amp-seq
ISDra2	gRNA7	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtagcttcaaatgagggaccaggctt	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtgtagaggacgacatggttcacagtt	amp-seq
ISDra2	gRNA8	ACACTCTTTCCCTACACGACGCTCTTCCGATCTcacaaacctctgttgcttgaagc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcattgggtacctaggagctgtaac	amp-seq
ISDra2	gRNA9	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgcgctaaactttataaaccctgatg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtatgattgccagtgagataaagattc	amp-seq
ISDra2	gRNA11	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgaatcgagattacatgcaggtctta	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTccaacactgtatccaactaagtattc	amp-seq
ISDra2	gRNA12	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgaaccgacccgagaagagatttg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtggaatacacacatttgtacaacca	amp-seq
ISDra2	gRNA13	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtggagttctcgaaatttcatctagtg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcggaaggatgcagataaactagca	amp-seq
ISDra2	gRNA14	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtctgtaagtccaaaactcataccct	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcttccaacaagaacatgaagagagt	amp-seq
ISDra2	gRNA15	ACACTCTTTCCCTACACGACGCTCTTCCGATCTacagtggttgaactattagttagtgg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcaaaatctataattagcttgaattgatcaatac	amp-seq
ISDra2	gRNA16	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgtgaggttagtaactatgatgcatgt	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaagacaaatgaccatgttcttcctg	amp-seq
ISDra2	gRNA17	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTacagtggttgaactattagttagtgg	ACACTCTTTCCCTACACGACGCTCTTCCGATCTcctcaaacattcatcagttctttgat	amp-seq
ISDra2	gRNA18	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTacagtggttgaactattagttagtgg	ACACTCTTTCCCTACACGACGCTCTTCCGATCTcctcaaacattcatcagttctttgat	amp-seq

ISDra2	gRNA19	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTaagtaggctattgagttcacagttt	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTatgaaaccacaac aaccacatg	amp-seq
ISDra2	gRNA20	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTcttggtagagtagttagcatgctga	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTccctggacaagta tacaattaatcag	amp-seq
ISDra2	gRNA21	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgttgcagcttatgacacacaataag	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaatctttatctcact ggcaatcatag	amp-seq
ISDra2	gRNA22	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgttgcagcttatgacacacaataag	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaatctttatctcact ggcaatcatag	amp-seq
ISYmu1	gRNA1	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTaccaaattacgttgagatgcatggtc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTgctcttttgttgctt acgaaacatgt	amp-seq
ISYmu1	gRNA2	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgaagcagttgtgagttaagttggaga	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTttgtcttaagcgct tgagaagtgg	amp-seq
ISYmu1	gRNA3	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTatgtcttcagtcttgtgctacacacc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaaacggaacaaa caagaacctaacct	amp-seq
ISYmu1	gRNA4	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTtctcttcatgttcttgttggaagtcgg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTgtgctggtaggac atctgggaagt	amp-seq
ISYmu1	gRNA5	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgacgtcaggaagaacatggtcatttg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTagaacatttcagc gctaatgctacaa	amp-seq
ISYmu1	gRNA7	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTtagcttcaaatgagggaccaggctt	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTaagtacttctgtttt gtgtaatctcc	amp-seq
ISYmu1	gRNA8	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTcacaaacctctgttgcttgaagc	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcattgggtacctag gagctgtaac	amp-seq
ISYmu1	gRNA9	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgcgctaaactttataaaccctgatg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtatgattgccagtg agataaagattc	amp-seq
ISYmu1	gRNA10	GTGACTGGAGTTCAGACGTGTGCTCTTCCG ATCTgaatctttatctcactggcaatcata	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtgttgcagcttatgaca cacaataag	amp-seq
ISYmu1	gRNA12	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgaaccgacccgagaagagatttg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtggaatacacaca tttgtacaacca	amp-seq
ISAam1	gRNA11	GTGACTGGAGTTCAGACGTGTGCTCTTCCG ATCTagtttcttactcactaatggaagcac	ACACTCTTTCCCTACACGACGCTCTTCCGATCTaaggagcttcaggata tcgactacaa	amp-seq
ISAam1	gRNA12	GTGACTGGAGTTCAGACGTGTGCTCTTCCG ATCTgcaggtgaatttcttatttatgcagt	ACACTCTTTCCCTACACGACGCTCTTCCGATCTgaaggttacttctgca aaaaacaaca	amp-seq
ISAam1	gRNA13	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTcgtctggaccaagaaagttgt	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTcagaaacgtgaga tgtcaaatctgt	amp-seq
ISAam1	gRNA14	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTgttaagttgttattgtgttctctgg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTgaaaatatgtaaa ccagtctcatacc	amp-seq

ISAam1	gRNA15	GTGACTGGAGTTCAGACGTGTGCTCTTCCG ATCTcgctgcgtctcctgtttctctac	ACACTCTTTCCCTACACGACGCTCTTCCGATCTccggtggaagcaaat cagac	amp-seq
ISAam1	gRNA16	ACACTCTTTCCCTACACGACGCTCTTCCGAT CTtgcaaggaggaggagtatttgg	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtatccgattccgct tagtttgc	amp-seq
ISAam1	gRNA17	GTGACTGGAGTTCAGACGTGTGCTCTTCCG ATCTaccaaattacgttgagatgcatggtc	ACACTCTTTCCCTACACGACGCTCTTCCGATCTtggtcttgtttggttttc agacc	amp-seq
ISYmu1	gRNA12	tgttacacaacttattatgattgggct	ggaagtagccgataacaaaatggag	Sanger

603 **Supplementary Table 9: Primers used for amp-seq or Sanger sequencing.** The TnpB and Guide

604 columns indicate the site being targeted for each TnpB. The F primer and R primer columns list

605 the oligo sequences used to amplify genomic DNA for amp-seq or Sanger Sequencing.

oligo name	Oligo sequence	Description
SP9238	TGTATAGACTGTTTGAGATCGGCG	RT-PCR primer to check for the presence/absence of TRV in mutant albino plants
SP9239	GTAATAACGCTTACGTAGGCGAGG	RT-PCR primer to check for the presence/absence of TRV in mutant albino plants
IPP2 RT-PCR Fw	GTATGAGTTGCTTCTCCAGCAAAG	RNA control for RT-PCR
IPP2 RT-PCR Rev	GAGGATGGCTGCAACAAGTGT	RNA control for RT-PCR

607 **Supplementary Table 10: Oligos for RT-PCR.** The name of each oligo is listed in the Oligo name

608 column. Each oligo sequence is provided in the Oligo sequence column, with a description of

609 what the oligos were used for in the Description column.

References:

1.   Altpeter, F. *et al.* Advancing Crop Transformation in the Era of Genome Editing. *Plant Cell*

28, 1510–1520 (2016).

2.   Chen, K., Wang, Y., Zhang, R., Zhang, H. & Gao, C. CRISPR/Cas Genome Editing and

Precision Plant Breeding in Agriculture. *Annu. Rev. Plant Biol.* **70**, 667–697 (2019).

3.   Nasti, R. A. & Voytas, D. F. Attaining the promise of plant gene editing at scale. *Proceedings*

of the National Academy of Sciences **118**, e2004846117 (2021).

4.   Ali, Z. *et al.* Efficient Virus-Mediated Genome Editing in Plants Using the CRISPR/Cas9

System. *Mol. Plant* **8**, 1288–1291 (2015).

5.   Ellison, E. E., Chamness, J. C. & Voytas, D. F. Viruses as vectors for the delivery of gene-

editing reagents. (2021).

6.   Beernink, B. M., Lappe, R. R., Bredow, M. & Whitham, S. A. Impacts of RNA Mobility Signals

on Virus Induced Somatic and Germline Gene Editing. *Front Genome Ed* **4**, 925088 (2022).

7.   Uranga, M. *et al.* Efficient Cas9 multiplex editing using unspaced sgRNA arrays engineering

in a Potato virus X vector. *Plant J.* **106**, 555–565 (2021).

8.   Bradamante, G., Mittelsten Scheid, O. & Incarbone, M. Under siege: virus control in plant

meristems and progeny. *Plant Cell* **33**, 2523–2537 (2021).

9.   Ma, X., Zhang, X., Liu, H. & Li, Z. Highly efficient DNA-free plant genome editing using

virally delivered CRISPR–Cas9. *Nature Plants* **6**, 773–779 (2020).

10. Liu, Q., Zhao, C., Sun, K., Deng, Y. & Li, Z. Engineered biocontainable RNA virus vectors for

non-transgenic genome editing across crop species and genotypes. *Mol. Plant* **16**, 616–631

(2023).

11. Lee, S.-Y. *et al.* Development of virus-induced genome editing methods in Solanaceous

crops. *Hortic Res* **11**, uhad233 (2023).

12. Ariga, H., Toki, S. & Ishibashi, K. Potato Virus X Vector-Mediated DNA-Free Genome Editing

in Plants. *Plant Cell Physiol.* **61**, 1946–1953 (2020).

13. Ghoshal, B. *et al.* A viral guide RNA delivery system for CRISPR-based transcriptional

activation and heritable targeted DNA demethylation in Arabidopsis thaliana. *PLoS Genet.*

16, e1008983 (2020).

14. Karvelis, T. *et al.* Transposon-associated TnpB is a programmable RNA-guided DNA

endonuclease. *Nature* **599**, 692–696 (2021).

15. Altae-Tran, H. *et al.* The widespread IS200/IS605 transposon family encodes diverse

programmable RNA-guided endonucleases. *Science* **374**, 57–65 (2021).

16. Meers, C. *et al.* Transposon-encoded nucleases use guide RNAs to promote their selfish

spread. *Nature* **622**, 863–871 (2023).

17. Xiang, G. *et al.* Evolutionary mining and functional characterization of TnpB nucleases

identify efficient miniature genome editors. *Nat. Biotechnol.* 1–13 (2023)

doi:10.1038/s41587-023-01857-x.

18. Li, Q. *et al.* Genome editing in plants using the TnpB transposase system. *aBIOTECH* (2024)

doi:10.1007/s42994-024-00172-6.

19. Li, Z. *et al.* Engineering a transposon-associated TnpB-ωRNA system for efficient gene

editing and phenotypic correction of a tyrosinaemia mouse model. *Nat. Commun.* **15**, 1–11

(2024).

20. Yoo, S.-D., Cho, Y.-H. & Sheen, J. Arabidopsis mesophyll protoplasts: a versatile cell system

for transient gene expression analysis. *Nat. Protoc.* **2**, 1565–1572 (2007).

21. Zhang, X., Henriques, R., Lin, S.-S., Niu, Q.-W. & Chua, N.-H. Agrobacterium-mediated

transformation of Arabidopsis thaliana using the floral dip method. *Nat. Protoc.* **1**, 641–

646 (2006).

22. Wang, M. *et al.* A gene silencing screen uncovers diverse tools for targeted gene

repression in Arabidopsis. *Nature Plants* **9**, 460–472 (2023).

23. Ellison, E. E. *et al.* Multiplexed heritable gene editing using RNA viruses and mobile single

guide RNAs. *Nat Plants* **6**, 620–624 (2020).

24. Nagalakshmi, U., Meier, N., Liu, J.-Y., Voytas, D. F. & Dinesh-Kumar, S. P. High-efficiency

multiplex biallelic heritable editing in Arabidopsis using an RNA virus. *Plant Physiol.* **189**,

1241–1245 (2022).

25. Qi, Y. *et al.* Increasing frequencies of site-specific mutagenesis and gene targeting in

Arabidopsis by manipulating DNA repair pathways. *Genome Res.* **23**, 547–554 (2013).

26. Brinkman, E. K. & van Steensel, B. Rapid Quantitative Evaluation of CRISPR Genome Editing

by TIDE and TIDER. *Methods Mol. Biol.* **1961**, 29–44 (2019).

27. Bae, S., Park, J. & Kim, J.-S. Cas-OFFinder: a fast and versatile algorithm that searches for

potential off-target sites of Cas9 RNA-guided endonucleases. *Bioinformatics* **30**, 1473–

1475 (2014).

28. Badon, I. W., Oh, Y., Kim, H.-J. & Lee, S. H. Recent application of CRISPR-Cas12 and OMEGA

system for genome editing. *Mol. Ther.* **32**, 32–43 (2024).

29. Macfarlane, S. A. Tobraviruses--plant pathogens and tools for biotechnology. *Mol. Plant*

Pathol. **11**, 577–583 (2010).

30. Zhang, C., Liu, S., Li, X., Zhang, R. & Li, J. Virus-Induced Gene Editing and Its Applications in

692   Plants. *Int. J. Mol. Sci.* **23**, (2022).

693 31. LeBlanc, C. *et al.* Increased efficiency of targeted mutagenesis by CRISPR/Cas9 in plants

694   using heat stress. *Plant J.* **93**, 377–386 (2018).

695 32. Li, Z. *et al.* Genome editing in plants using the compact editor CasΦ. *Proc. Natl. Acad. Sci.*

696   *U. S. A.* **120**, e2216822120 (2023).

697 33. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

698   transform. *Bioinformatics* **25**, 1754–1760 (2009).

699 34. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing

700   next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

701 35. Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nat.*

702   *Methods* **15**, 591–594 (2018).

703 36. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

704   features. *Bioinformatics* **26**, 841–842 (2010).