Temporal dynamics analysis reveals that concurrent working

memory load eliminates the Stroop effect through disrupting

stimulus-response mapping

Yafen Li[a], Yixuan Lin[b], Qing Li[c], Yongqiang Chen[c], Zhifang Li[b], Antao Chen[a,*]

[a] School of Psychology, Shanghai University of Sport, Shanghai 200438, China

[b] School of Psychology and Cognitive Science, East China Normal University,

Shanghai 200062, China

[c] Faculty of Psychology, Southwest University, Chongqing 400715, China

*Corresponding author:

School of Psychology, Shanghai University of Sport, Yangpu District, Shanghai 200438,

China.

Email: chenantao@sus.edu.cn

Abstract

Concurrent verbal working memory task can eliminate the color-word Stroop effect. Previous research, based on specific and limited resources, suggested that the disappearance of the conflict effect was due to the memory information preempting the resources for distractors. However, it remains unclear which particular stage of Stroop conflict processing is influenced by working memory loads. In this study, electroencephalography (EEG) recordings with event-related potential (ERP) analyses, time-frequency analyses, multivariate pattern analyses (MVPA), and representational similarity analyses (RSA) were applied to provide an in-depth investigation of the aforementioned issue. Subjects were required to complete the single task (the classical manual color-word Stroop task) and the dual task (the Sternberg working memory task combined with the Stroop task), respectively. Behaviorally, the results indicated that the Stroop effect was eliminated in the dual-task condition. The EEG results showed that the concurrent working memory task did not modulate the P1 and alpha bands. However, it modulated the sustained potential (SP), late theta (740-820 ms), and beta (920-1040 ms) power, showing no difference between congruent and incongruent trials in the dual-task condition but significant difference in the single-task condition. Importantly, the RSA results revealed that the neural activation pattern of the late theta was similar to the response interaction pattern. Together, these findings implied that concurrent working memory task eliminated the Stroop effect through disrupting stimulus-response mapping.

Keywords: Stroop effect; working memory; electroencephalography; multivariate pattern analysis

1. Introduction

It is critical to effectively deal with the conflict between task-relevant and task-irrelevant information to successfully complete goal-directed behaviors. One of the most commonly employed experimental paradigms for evaluating conflict processing is the color-word Stroop task, which is characterized by lower accuracy and/or longer reaction times (RT) in incongruent trials compared to congruent trials (Stroop, 1935). Of note, an interesting phenomenon is that concurrent verbal working memory task can reduce or even eliminate the Stroop conflict (Kim et al., 2005; Luna et al., 2020; Park et al., 2007; Zhao et al., 2014). In terms of the load theory based on multiple resources, cognitive resources are limited while specific to different cognitive processes. When substantial resources are allocated to maintaining memory information, the resources required to process same-dimension distractors are occupied, reducing the interference caused by the distractors (Brockhoff et al., 2022; Luna et al., 2020; Park et al., 2007). Although this theory attempts to explain the disappearance of the Stroop conflict, its description remains relatively vague, and it does not further reveal which specific processes are influenced during Stroop processing.

Stroop conflict processing should be understood not as a single process but as multiple successive subprocesses, including stimulus processing, stimulus-response mapping, and response output stage (Banich, 2019; Parris et al., 2022; Viviani et al., 2023). Related research suggests that there might be two possibilities. The first one is that concurrent working memory task primarily decreases the neural encoding of

perceptual visual features, known as the early-stage modulation hypothesis. The

sensory recruitment hypothesis posits that the sensory cortex is crucial for

maintaining visual content by modulating sensory gain, thereby biasing processing

towards relevant stimuli (Hallenbeck et al., 2021; Wolff et al., 2020). Moreover,

several researchers have suggested that the Stroop effect relies more on the input-

driven allocation of attention to perceptual features (Algom et al., 2022; Bugg &

Crump, 2012; Haciahmet et al., 2023; Schmidt, 2019). Accordingly, these views

suggest that working memory maintenance competes for neural resources with the

lower-level visual encoding of distractors, hindering conflict formation at an early

stage. Besides, the second one is that concurrent working memory task primarily

weaken higher-level processing, termed the late-stage modulation hypothesis. The

theoretical underpinning is that the working memory system comprises not only

sensory buffers but also a central system (Baddeley, 2012) that mainly recruits the

prefrontal cortex, partially overlapping with the brain networks involved in Stroop

conflict processing (Friehs et al., 2020; Okayasu et al., 2023). Furthermore, some

research indicated that filtering out highly familiar and easily processed semantic

information during the early sensory processing stage is difficult for individuals

(Augustinova et al., 2019; Cavanagh & Frank, 2014; Itthipuripat et al., 2019).

Consistent with this hypothesis, both relevant and irrelevant sensory inputs receive

comprehensive semantic processing, while working memory load constrains the

process of binding higher-level semantic codes with responses, potentially influencing

the subsequent response output stage. Overall, the two hypotheses, although distinct,

are not mutually exclusive and can be inspected from a time-dynamic perspective.

In the present study, a single task (the classical manual Stroop task) and a dual task (the Sternberg working memory task combined with the Stroop task) were employed to explore the aforementioned hypotheses with electroencephalography recording of high temporal resolution. Several event-related potentials (ERP) highly relevant to Stroop conflict processing will be focused on in this study. Previous studies have proposed that within cognitive control mechanisms, P1 signifies the earliest stage of top-down processes in early sensory processing, gating the direction of information processing in the brain (Klimesch et al., 2007). Moreover, the incongruent trials in the Stroop task evoke smaller positive P1 than congruent trials, supporting that P1 is a valid index in conflict processing (Yu et al., 2015). Additionally, numerous studies have identified two ERP components exclusively elicited by the Stroop task (i.e., the N450 and the sustained potential (SP)). N450 is a frontal-central negative deflection that exhibits greater negativity in incongruent trials compared to congruent trials within 350-500 ms after stimulus onsets, which is typically interpreted as conflict monitoring, the process of evaluating conflict levels and communicating the conflict information to control systems (Coderre et al., 2011; Heidlmayr et al., 2020; Xu et al., 2024). As for the SP component, it shows a centro-parietal positive deflection from approximately 600 ms to 1000 ms in incongruent conditions, which is regarded as an index of conflict resolution (Di Russo & Bianco, 2023; Donohue et al., 2016; Heidlmayr et al., 2020; Vo et al., 2021). In all, these ERP components, which are distinctly time-sequential, can help explore the dynamic

effects of working memory on Stroop conflict processing.

In addition to the ERP components, conflict processing also involves various

neural oscillatory activities. First, extensive research has found that mid-frontal theta

(4-7 Hz) power increases in conflict situations with the time windows of early and late

theta band effects in stimulus-locked analyses corresponding to those of the N450 and

SP effects, which suggests that early and later theta power reflects conflict detection

and resolution processes, respectively (Brilliant et al., 2024; Haciahmet et al., 2023;

Senoussi et al., 2022; Xu et al., 2024; Zhao et al., 2015). Second, alpha oscillations

(8-13 Hz) can serve as a neural inverse index of mental activity or alertness, while a

decrease in alpha power reflects increased alertness or enhanced attentional inhibition

of distractors (Arakaki et al., 2022; Tafuro et al., 2019; Zhou et al., 2023; Zhu et al.,

2023). Third, previous study has shown that the response incongruent trial evokes a

small beta power than the stimulus incongruent trial in the Stroop task (Zhao et al.,

2015). The suppressed effect of the beta power most likely reflects the inhibition of

proponent responses, associated with motion preparation (Böttcher et al., 2023;

Chung et al., 2024; Wendiggensen et al., 2022; Zhao et al., 2015). Accordingly, these

frequency bands can provide a comprehensive characterization of the Stroop conflict

processing.

It is worth realizing that neural oscillations are not characterized by a pre-

determined time window like ERPs. However, multivariate pattern analysis (MVPA)

can help to solve the difficulty of hypothesizing which frequency band or the time

window is changed in the Stroop task. As a data-driven technique, MVPA provides

novel insight into the differences between two conditions by identifying the features

that contribute most to classification (Li et al., 2023; Lin et al., 2023; Meier et al.,

2012). Therefore, the present study will combine time-frequency analysis and MVPA

to inspect the dynamic neural characteristics of the modulation of working memory on

Stroop conflict processing at the neural oscillatory level.

Notably, even if differences are found in ERPs or neural oscillations, the extent

to which they are linked to behavior is an interesting question. Representational

similarity analysis (RSA) is applied to build a connection between behavior and

neural activities in the present study (Kriegeskorte & Kievit, 2013). Recent studies

have suggested that RSA is a reliable way to uncover the neural mechanisms behind

conflict processing (Freund et al., 2021; Yang et al., 2023), which helps identify

which neural activities are central to the influence of working memory on Stroop

conflict processing. Overall, the present study employs EEG technology, through

MVPA and RSA, in an attempt to comprehensively uncover the dynamic neural

underlying the elimination of the Stroop effect influenced by working memory when

the dimensions of memory information and distractors overlap.

2. Results

2.1. Behavioral results

For behavioral analysis, a two-way repeated measures ANOVA (task type: single

task vs. dual task × congruency: congruent vs. incongruent) was performed on RTs.

The results showed that the RTs of the single-task condition were faster than dual-task

condition (main effect of task type: $F_{(1, 31)} = 113.28$, $p < 0.001$, $\eta_p^2 = 0.79$) and a

reliable Stroop interference effect (main effect of congruency: $F_{(1, 31)} = 32.53$, $p <$

0.001, $\eta_p^2 = 0.51$). Notably, the interaction between task type and congruency was

significant ($F_{(1, 31)} = 8.62$, $p = 0.006$, $\eta_p^2 = 0.22$). Simple effect analysis revealed that,

in the single-task condition, the RTs of incongruent trials were significantly slower (M

$= 851.15$, $SE = 32.16$) than congruent trials ($M = 736.10$, $SE = 26.29$) ($p < 0.001$).

However, in the dual-task condition, the RTs did not differ between incongruent trials

($M = 1073.04$, $SE = 39.73$) and congruent trials ($M = 1033.68$, $SE = 49.43$) ($p =$

0.089), indicating that the Stroop effect of RTs only occurred in single-task condition

(Fig. 1).

156 2.2. ERP results

157 2.2.1. P1

Two-way repeated-measures ANOVA was conducted on the P1 component,

revealing a significant main effect of task type ($F_{(1, 31)} = 29.92$, $p < 0.001$, $\eta_p^2 = 0.49$).

Specifically, the P1 amplitude of the dual task ($M = 1.63$, $SE = 0.30$) was significantly

greater than the single task ($M = 0.39$, $SE = 0.28$). However, neither the main effect of

congruency ($F_{(1, 31)} = 0.00$, $p = 0.99$, $\eta_p^2 = 0.00$) nor the two-way interaction ($F_{(1, 31)} =$

0.23, $p = 0.63$, $\eta_p^2 = 0.007$) was significant (Fig. 2).

164 2.2.2. N450

Analysis of the N450 component showed the main effect of task type was

significant ($F_{(1, 31)} = 23.79$, $p < 0.001$, $\eta_p^2 = 0.43$), resulting in greater N450 amplitude

in the single task ($M = 1.90$, $SE = 0.33$) compared to the dual task ($M = 0.84$, $SE =$

0.34). However, the main effect of current congruency ($F_{(1, 31)} = 1.60$, $p = 0.22$, $\eta_p^2 =$

0.05) and the interaction ($F_{(1, 31)} = 0.02$, $p = 0.90$, $\eta_p^2 = 0.001$) were not significant

(Fig. 3A, B, C).

2.2.3. SP

For the SP component, two-way repeated-measures ANOVA showed the main

effect of task type was significant ($F_{(1, 31)} = 14.54$, $p = 0.001$, $\eta_p^2 = 0.32$) but the main

effect of congruency was not ($F_{(1, 31)} = 2.89$, $p = 0.10$, $\eta_p^2 = 0.09$). Importantly, the

interaction between congruency and task type was significant ($F_{(1, 31)} = 5.68$, $p < 0.05$,

$\eta_p^2 = 0.16$). Simple effect analysis showed that, in the single task, the SP amplitude in

incongruent trials ($M = 1.29$, $SE = 0.39$) was significantly greater than in congruent

trials ($M = 0.70$, $SE = 0.30$) ($p = 0.009$). On the contrary, in the dual task, the SP

amplitude did not differ between congruent trials ($M = 0.27$, $SE = 0.34$) and

incongruent trials ($M = 0.22$, $SE = 0.37$) ($p = 0.98$) (Fig. 3A, D, E).

2.3. MVPA results

Cluster-level permutation tests of the theta band indicated that only one cluster in

740-820 ms could significantly classify the Stroop effect of the single-task and dual-

task (Fig. 4A). Regarding the beta band, tests showed a significant above-chance

difference between the two task types from 920 ms to 1040 ms (Fig. 4C). However,

no reliable decoding was observed between the two task types in the alpha band (Fig.

4B).

2.4. Time-Frequency results

2.4.1. Theta band

The two-way repeated-measures ANOVA on theta power revealed that the main effect of congruency was significant ($F_{(1, 31)}$ = 14.68, p = 0.001, η_p^2 = 0.32), but the main effect of task type was not ($F_{(1, 31)}$ = 0.30, p = 0.59, η_p^2 = 0.01). Notably, the interaction was significant ($F_{(1, 31)}$ = 7.71, p = 0.009, η_p^2 = 0.20). Simple effect analysis showed that, in the dual task, the theta power did not differ between congruent trials (M = 3.12, SE = 0.24) and incongruent trials (M = 3.27, SE = 0.24) (p = 0.63). However, in the single task, the theta power was significantly higher in incongruent trials (M = 4.00, SE = 0.32) than in congruent trials (M = 2.68, SE = 0.20) (p < 0.001) (Fig. 5).

2.4.2. Beta band

Analysis of the beta power showed the main effect of task type ($F_{(1, 31)}$ = 34.55, p < 0.001, η_p^2 = 0.53) and the main effect of congruency ($F_{(1, 31)}$ = 11.61, p = 0.002, η_p^2 = 0.27) were both significant. Importantly, the interaction between task type and congruency reached a significant level ($F_{(1, 31)}$ = 15.64, p < 0.001, η_p^2 = 0.34). Simple effect analysis revealed that, in the single-task condition, the beta power was significantly lower in the incongruent trials (M = 0.53, SE = 0.06) than in the congruent trials (M = 0.80, SE = 0.08) (p < 0.001). While no significant difference was found between them in the dual-task condition (p = 0.40) (Fig. 6).

2.5. RSA results

The RSA results showed that the interaction pattern in theta band significantly

contributes to variation in the interaction pattern of response time (estimate = 0.17, *SE*

= 0.05, *t* = 3.22, $p_{corrected}$ = 0.004). However, the correlation between the dissimilarity

matrix of the SP (estimate = 0.02, *SE* = 0.05, *t* = 0.48, $p_{corrected}$ = 0.63) and theta

(estimate = 0.10, *SE* = 0.05, *t* = 2.03, $p_{corrected}$ = 0.06) representation and the

interaction pattern of response time was not significant (Fig. 7).

3. Discussion

The present study systematically investigated the dynamic neural mechanisms

underlying the modulation of concurrent working memory task on Stroop conflict

processing when the dimensions of memory information and distractors overlap. At

first, we verified behaviorally that working memory processing eliminated the conflict

effect. Specifically, in the single task, the response times of incongruent trials were

significantly slower than congruent trials, but there was no significant difference

between the two in the dual task. For the ERP results, no significant interaction was

found in P1 or N450, but in SP. Namely, it showed that the incongruent trials evoked

more positive SP than the congruent trials in the single task. While there was no

difference between the two in the dual task. Moreover, MVPA results showed that

only the theta power (740-820 ms) and the beta power (920-1040 ms) of conflict

effect could successfully classify different task types. Further, in the corresponding

time windows, we found that only in the single task, the incongruent trial induced

greater theta power and smaller beta power than the congruent trial. Finally, RSA

showed that only the interaction pattern in theta power correlated with response time.

All these results collectively suggested that, when the memorized information

overlaps with the dimension of distractors, working memory processing modulates the

late stage rather than early stage of Stroop conflict processing.

Behaviorally, we found that the Stroop interference effect was reduced under the

dual-task condition, replicating previous findings (Kim et al., 2005; Luna et al., 2020;

Park et al., 2007; Zhao et al., 2014). Consistent with the latest load theory (Murphy et

al., 2016; Park et al., 2007), there are multiple resources, each with its own

processing-specific, limited-capacity pool. Working memory load will tie up the

limited resources necessary for distractor processing, diminishing the intensity of

distractor processing, and thereby reducing their interference with the task. In our

study, concurrent visual short-term working memory maintenance occupied the

semantic resources required for distractor processing, resulting in the elimination of

the Stroop effect. Interestingly, a previous study (Zhao et al., 2014) that reported a

diminished Stroop interference effect employed the classic Chinese character Stroop

task as well, but the working memory task involved memorizing meaningless visual

shapes made from hand-drawn lines. This seems inconsistent with the load theory

hypothesis, as the images appear to occupy spatial-related resources rather than

overlapping with the distractor stimulus processing dimension. However, since

Chinese characters are pictographs that require visual shape encoding and memory

(Chaoran, 2019), they inherently occupy the same resources for processing Chinese

characters as in our study. Additionally, this study suggested that the modulation

effect of the working memory task on the typical Chinese character Stroop task was

relatively minor. We speculate that this is because graphic processing occupies lower-

level resources related to the overall shape and spatial structure of Chinese characters

(Zhang et al., 2020). In contrast, our study, which employed Chinese characters as the

working memory load, could directly influence higher-level resources involved in

semantic processing. Therefore, we believe that verbal working memory load can

significantly modulate the Stroop effect. Overall, the improvement in conflict

processing observed in our study is consistent with the core hypotheses of load theory

based on multiple resources, paving the way for further exploration into the neural

mechanisms behind this improvement.

The ERP findings indicated that the working memory task influenced only the SP

component, which reflects the process of conflict resolution. Consistent with most

research, our study showed that the SP amplitude is more positive in conflict

situations (Heidlmayr et al., 2020; Zhao et al., 2015). Crucially, our study found that

verbal working memory load eliminated the difference in SP between congruent and

incongruent trials, indicating that conflict resolution did not occur in the dual-task

condition. Furthermore, Some studies have suggested that N450 and SP are related to

stimulus conflict (or semantic conflict) and response conflict, respectively (Coderre et

al., 2011; West et al., 2005; Zhao et al., 2015). Therefore, this result may imply that

participants neither encountered stimulus conflict nor response conflict in the dual-

task condition. Additionally, our study did not find any differences in P1 and N450

between the different trial types. For P1, most studies indicate that Stroop interference

occurs only in the later stages of response selection, rather than during the sensory

input processing stage (David et al., 2011; Di Russo & Bianco, 2023). For N450, our

previous research shows that semantic conflict in the Stroop task only appears in the

early practice stages, and the manual Stroop task should include only "pure" response

conflict (Chen et al., 2013). Therefore, due to the practice effect, the semantic conflict

represented by N450 may have disappeared in this study. Generally, the overall ERP

results suggest that working memory influences the later stage of response conflict

resolution, while not influencing the early stage of perceptual processing and semantic

encoding.

In addition to the ERP results, the MVPA results from time-frequency also

provide supportive evidence that working memory modulates Stroop conflict

processing in the later stage. First, in line with the previous study (Zhao et al., 2014),

there are no continuous time periods in which alpha activities (incongruency minus

congruency) can successfully discriminate the single-task and dual-task. Alpha is

regarded as an indicator of top-down attentional modulation, with an increase in alpha

reflecting the inhibition of distractor input (Van Diepen et al., 2019; Zhou et al.,

2023), implying that the patterns of attentional adjustment are similar across different

task types in our study. This finding may be related to the attentional rejection

template (Wen et al., 2022). In the Sternberg task, participants are required to

remember only seven Chinese characters and reject the others, which is similar to the

attention template used for rejecting characters in the Stroop task. Second, the

different task types can be successfully discriminated by late theta activity after the

stimulus presentation (740-820 ms). Similarly, previous research has found that

current working memory leads to smaller differences in theta activity (400-1000 ms)

between incongruent and neutral trials (Zhao et al., 2014). According to the

framework of event coding theory (TEC), late theta may represent the process of

stimulus-response binding or mapping (Beste et al., 2023; Wendiggensen et al., 2023).

And this has been demonstrated in several empirical experiments. For example, theta

is thought to be involved in adjusting the processing of stimulus information to

correspond with appropriate responses (Cavanagh et al., 2009). In line with this

opinion, about -200 to 0 ms before a response, higher theta activity is observed in

high-conflict conditions compared to low-conflict conditions (Cohen & Donner,

2013). Accordingly, working memory may eliminate the Stroop effect by influencing

the stimulus-response mapping process. Lastly, the present study also shows that beta

activity between 920-1040 ms after stimulus presentation can successfully

discriminate between the two task types. Beta activity is often thought to be

associated with motor priming. Specifically, it decreases at the onset of movement

(Jenkinson & Brown, 2011), but increases during holding a static status (Baker et al.,

2001). Since response button selection definitely occurs after conflict resolution and is

a late motor adjustment according to the stimulus, this result suggests that working

memory may also influence the response output stage of Stroop conflict processing.

The RSA results further reveal that the neural activity pattern of theta is most

similar to the behavioral Stroop effect modulation pattern. This result can be

interpreted from the standpoint of neural efficiency, suggesting that a reduction in

activation patterns reflects greater neural efficiency (Rypma et al., 2006). Therefore,

we speculate that the core reason for the disappearance of the behavioral Stroop effect

under the dual-task condition is that, compared to congruent trials, neural activation in

theta decreases and response speed increases in incongruent trials (Fig. 1 and Fig.

5C). Moreover, this result aligns with many studies that have found theta can predict

reaction times (Kaiser & Schütz-Bosbach, 2021; Senoussi et al., 2022). However,

although SP and beta exhibited interaction effects in the data patterns, they may not be

the core indicators directly determining the behavioral modulation effect. SP, as an

ERP component, contains multiple oscillatory bands, indicating several parallel brain

processes (Brilliant et al., 2024). Beta, as a general response-related indicator, may

not be directly linked to conflict processing. Combined with the MVPA decoding

results indicating that the beta band (920-1040 ms) follows the theta band (740-820

ms), we speculate that changes in beta bands may be influenced by theta bands,

thereby indirectly influencing the behavioral Stroop effect. Overall, when the

dimensions of working memory load overlap with those of distractor stimuli,

individuals do not need to engage in conflict resolution. Specifically, the memory

information occupies the resources for the late-stage stimulus-response mapping of

the distractors, weakening the mapping of distractors to incorrect responses, and

subsequently inhibiting the response output of the distractors, resulting in the

disappearance of response conflict.

In summary, this study demonstrated that the concurrent working memory task

can eliminate Stroop conflict through disrupting stimulus-response mapping. The

results indicated that concurrent working memory load influenced the late stage

related to stimulus-response mapping and response selection processes, indexed by

SP, late theta band, and beta band; while it did not influence the early stage associated

with perceptual processing and attention adjustment, indexed by P1 and alpha band.

RSA further identified theta as the neural indicator directly associated with the

behavioral modulation pattern. These findings suggested that when the working

memory load overlaps with the dimension of distractor stimuli, it occupies the

resources for the stimulus-response mapping of the distractor, weakening the stronger

response tendency of the distractor, and ultimately leading to the disappearance of

response conflict. Accordingly, the present study comprehensively characterized the

dynamic neural mechanisms by which working memory modulates Stroop conflict

processing, further deepening our understanding of the mechanisms underlying

conflict generation.

4. Materials and methods

4.1. Participants

Using G*Power software (Faul et al., 2007), we calculated the sample size for

the experiment. The power analysis (power ≥ 0.9) on within-factors designs, assuming

a moderate effect size of 0.25, indicated a sample size of 30. We recruited 34 healthy

subjects (17 females; age: 21.38 ± 1.92 years) through advertisements. Two subjects

were excluded from further analyses due to excessive EEG artifacts. Finally, a total of

32 subjects (16 females; age: 21.44 ± 1.97 years) remained for behavioral and EEG

analyses. All participants were right-handed, had normal or corrected-to-normal

vision and normal color perception assessed by the Ishihara Color Test, and reported

no history of neurologic or psychiatric disease. They all signed written informed

consent before the experiment. All procedures were conducted in accordance with the

principles of the Declaration of Helsinki and its later amendments and were approved

by the Human Research Human Ethics Committee of the Shanghai University of

Sport.

4.2. Apparatus and Stimuli

Participants were instructed to perform experiments in a soundproof room. The

experimental stimuli were presented on a 19-inch color monitor (resolution 1920 ×

1080, refresh rate 60 Hz) with a viewing distance of approximately 65 cm. Stimulus

presentation and response recording were controlled by a Microsoft PC running

Matlab (Mathworks, Inc.) with Psychophysics Toolbox extensions (Brainard, 1997;

Pelli, 1997).

The experiment employed the single task (the classical manual Stroop task) and

the dual task (the Sternberg working memory task combined with the Stroop task).

The Stroop task stimuli consisted of four Chinese words: "红" (red, RGB: 255, 0, 0),

"绿" (green, RGB: 0, 255, 0), "蓝" (blue, RGB: 0, 0, 255), and "黄" (yellow, RGB:

255, 255, 0). They were presented in the semantically corresponding font color

(congruent trials) or a different font color (incongruent trials) in the center of a black

background (visual angle, ~1.15×1.15°). The Sternberg working memory task

involved seven Chinese characters chosen randomly among 14 familiar Chinese

characters: "人" (human), "刀" (knife), "大" (big), "水" (water), "天" (heaven),

"口" (mouth), "月" (moon), "立" (stand), "山" (mountain), "上" (up), "手" (hand),

"石" (stone), "女" (woman), and "火" (fire) (Zhao et al., 2010). They were equally

spaced on a horizontal row at the center of the display (visual angle for each

character, ~1.15×1.15°).

4.3. Procedure

For the dual task (Fig.8A), the subjects were required to memorize seven

Chinese characters in anticipation of a recognition test at the end of the trial. Each

trial began with a white fixation point for a random duration from 400 to 600 ms,

followed by seven white Chinese characters that required memorization for 2000 ms.

After a jittered 800-1200 ms fixation point, participants performed a Stroop task in

which the color of the words were identified as quickly as possible by pressing the

corresponding button ("F" with the left index finger, "D" with the left middle finger,

"J" with the right index finger, and "K" with the right middle finger), while

attempting to ignore the task-irrelevant semantic meaning. The display for the words

remained on the screen until a response was made. After the Stroop task, a fixation

point was randomly presented for 400-600 ms, and terminated by the appearance of a

probe character, to which the subject had to determine if the character was part of

their memorized list of items. During 50% of the trials, the probe letter was included

as part of the memorized set. Participants were asked to answer by pressing one of

two response buttons ("S" with the left ring finger and "L" with the left ring finger).

All response buttons in this experiment were counterbalanced across subjects. After a

response was given, the intertrial interval was randomized between 600 and 1000 ms

until the onset of the next trial. For the single task (Fig.8B), subjects were only

required to complete the individual Stroop task. Each block alternated between single

and dual tasks in an ABAB fashion, and the task type in the first block was

counterbalanced across participants. Participants were permitted to participate in the

formal experiment when their average error rate was below 15%. The formal

experiment included six blocks, totaling 144 trials, with an equal distribution of 50%

congruent and 50% incongruent trials presented randomly in each block. The entire

experiment lasted for around 30 minutes.

414 4.4. EEG recording and preprocessing

EEG data were recorded using standard 64 in-cap Ag/AgCl electrodes, following

the extended international 10-20 system (Brain Products GmbH, Germany; pass band,

0.01-100 Hz; sampling rate, 1000 Hz). Vertical and horizontal electrooculograms

(EOGs) were recorded from below the left eye and outer canthus of the right eye,

respectively. The FCz served as an online reference. The impedance of all electrodes

was maintained below 5 kΩ during the entire recording process.

The offline EEG data were preprocessed by EEGLAB V2023.0 (Delorme &

Makeig, 2004) and custom scripts in MATLAB 2021a. First, the continuous data was

down-sampled from the original sampling rate of 1000 Hz to 500 Hz. Then, the data

were filtered with a 30 Hz low-pass filter and a 0.1 Hz high-pass filter. After that, we

re-referenced the data to the average activity of all scalp channels and manually

inspected the data to remove obvious artifacts. Subsequently, we carried out an

independent component analysis (ICA, infomax algorithm) and thereby identified and

manually removed artifacts such as eyeblinks, horizontal eye movements, heart,

muscle, and line noise. As a next step, the data was segmented based on the trial type

(single-task congruent vs. single-task incongruent vs. dual-task congruent vs. dual-

task incongruent) and was time-locked to the Stroop stimulus onset. Continuous EEG

data were epoched from −200 to 1000 ms relative to word onset, with a baseline

correction applied using the 200 ms pre-stimulus interval. Finally, trials with

amplitude values exceeding ±100 μV at any electrode were rejected.

4.5. Data analysis

4.5.1. Behavioral analysis

Only trials that were correct in both the memory task and the Stroop task were

included in all subsequent analyses. In addition, trials in which response times (RTs)

deviated by more than three standard deviations from the condition mean were

excluded from behavioral analyses. We performed repeated-measures analysis of

variance (ANOVA) with Greenhouse-Geisser correction if needed, in which the

within-subject factors were congruency (congruent, incongruent) and task type (single

task, dual task). Adjustments for multiple comparisons were realized using Bonferroni

correction.

4.5.2. ERP analysis

Considering previous studies (Coderre et al., 2011; Xu et al., 2024; Zivony &

Lamy, 2022), P1 was calculated using a time window of 40 ms (110-150 ms) at

bilateral occipito-parietal electrodes (PO3, PO4, PO7, PO8, P7, P8). N450 was

computed using a time window of 150 ms (350-500 ms) at centro-parietal electrodes

(Cz, CPz, Pz). SP was scored as the mean amplitude at centro-parietal electrodes (Cz,

CPz, Pz) during a time window of 300 ms (650-950 ms).

452   4.5.3. Multivariate pattern analysis

Our multivariate pattern analysis was based on support vector machine (SVM), a

supervised machine learning method that operated on labeled samples. A binary label

with 1 for the single task and -1 for the dual task was used in this study. The

classification process at a given time point included a training phase and a testing

phase. During the training step, the SVM would find a decision boundary that

separated the samples in the input features using class labels. Next, the trained

classifier is used to predict the class labels of new examples. For the spectral MVPA,

we defined three frequency regions of interest (F-ROIs): theta (4-7 Hz), alpha (8-13

Hz), and beta (14-25 Hz). Then, three spatial regions of interest (S-ROIs) were

chosen: fronto-central (Fz, FCz, Cz), occipital-parietal (POz, Oz, PO3, PO4, PO7,

PO8, O1, O2) and central (C3, CP3, Cz, CP4, C4) to measure theta, alpha and beta

power, respectively (Zhao et al., 2015; Li et al., 2021). For each frequency band, the

selected features were the power difference between congruent and incongruent trials

in the single task and dual task in every 20 ms of each ROI (See "Time-frequency

analysis" section for the power transformation method). All data were normalized and

mapped to the range [-1, 1] within each class to mitigate scale effects.

A linear kernel SVM was chosen and trained through the LIBSVM toolbox

(Chang & Lin, 2011). The linear SVM has only one parameter C that determines the

trade-off between allowing misclassifications and training error minimization. The

functions were as follows:

$$K\left(X_i,X_j\right)=X_i^T X_j$$

$$\min \frac{1}{2}\|\omega\|^2 + C \sum_{i=1}^{m} \xi(i)$$

A classifier was trained and tested at each time point by using a 5-fold cross-

validation procedure. Trials were randomly divided into 5 equal-sized folds. Then, a

leave-one-out procedure was run in which the classifier was trained on 4-folds and

tested on the remaining fold. In this case, different C values were tried within a

certain range to find the best cross-validation accuracy (or the smallest value if there

were more than one). We set the range of C values to base 2, and the exponent range

to [10, 10] with a step size of 0.2. Here, accuracy was used to quantify the

performance of the classifier based on the results of cross-validation.

For statistical analyses, a nonparametric statistical test based on the cluster-level

permutation was implemented to deal with multiple comparisons while maintaining

strict control of the false-alarm rate (Maris & Oostenveld, 2007). In detail, t-tests

against chance level (acc = 0.5) were carried out at each data point to obtain the

statistic values. Statistical values with p-values < 0.01 were clustered together by

summing their statistic values. The cluster alpha was set to 0.01 to reduce the

likelihood of large clusters spanning the entire dataset (Mensen & Khatami, 2013).

Additionally, to avoid using extremely transient intervals that may not capture

meaningful psychological phenomena, we specified that at least two significant time

points formed a cluster. Then, we permuted the data for 1000 times by randomly

swapping condition labels to obtain a permutation distribution of the maximum

cluster-level statistic under the null hypothesis of no condition difference. If there was

any cluster with its real statistic larger than the threshold, we rejected the null

hypothesis and concluded that there was a significant difference.

4.5.4. Time-frequency analysis

To avoid edge artifacts, EEG data were re-epoched from -1000 ms to 2000 ms

relative to stimulus onset. EEG data were converted into time-frequency domain data

using continuous wavelet transform in Letswave Software

(https://www.letswave .org/; Mouraux & Iannetti, 2008). The cmor1-1.5 was chosen

as the mother wavelet function for the time-frequency representations ranging from 1

to 30 Hz with a step size of 0.29 Hz. The single-trial time-frequency representation

was obtained after wavelet transform, and they were then averaged throughout all

trials per condition. The baseline normalizations were performed at each time point

for each frequency using an event-related spectral perturbation (ERSP) transformation

$[ER_{t,f}\% = [A_{t,f} - R_f]/R_f]$ (Pfurtscheller & Lopes da Silva, 1999). A 400-200 ms pre-

stimulus time window was selected as the baseline time window. For the statistical

test, we only analyzed the frequency bands and corresponding time windows that

could successfully classify different tasks in MVPA, namely theta (740-820 ms) and

beta (920-1040 ms).

4.5.5. Representational similarity analysis

Our RSA analysis aimed to explore the correlation between neural interaction patterns and behavioral interaction patterns. First, we extracted data that could reflect interaction patterns, namely Stroop effect differences between single-task and dual-task conditions based on the subject-level in RT, SP, theta, and beta. Prior to analysis, data were normalized by z-scoring the values across all subjects. We then constructed Representational Dissimilarity Matrices (RDMs) measuring the geometrical distances between all subjects, yielding 32×32 symmetrical matrices. Here, Euclidean distance was employed to measure the distance between subjects (Edelman, 1998). Finally, we fitted a linear regression with the neural RDMs as regressors and the behavioral RDM as the dependent variable to test which neural RDMs could predict behavioral RDMs. *P*-values were corrected by the false discovery rate.

Data availability

Data and codes presented in this work are available at: https://osf.io/89bpz/.

Author Contributions

Yafen Li: Conceptualization, Data curation, Methodology, Investigation, Visualization, Writing - original draft. Yixuan Lin: Methodology, Visualization, Writing - review & editing. Qing Li: Conceptualization, Methodology, Validation. Yongqiang Chen: Methodology, Validation. Zhifang Li: Methodology, Validation. Antao Chen: Conceptualization, Supervision, Writing - review & editing.

Funding information

This work was supported by grants from the National Natural Science Foundation of China (32371105).

Competing interests

The authors declare that they have no competing interests.

References

Algom, D., Fitousi, D., & Chajut, E. (2022). Can the Stroop effect serve as the gold standard of conflict monitoring and control? A conceptual critique. *Memory & Cognition*, *50*(5), 883–897. https://doi.org/10.3758/s13421-021-01251-5

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., van Langen, J., & Kievit, R. A. (2021). Raincloud plots: A multi-platform tool for robust data visualization. *Wellcome Open Research*, *4*, 63. https://doi.org/10.12688/wellcomeopenres.15191.2

Arakaki, X., Hung, S.-M., Rochart, R., Fonteh, A. N., & Harrington, M. G. (2022). Alpha desynchronization during Stroop test unmasks cognitively healthy individuals with abnormal CSF Amyloid/Tau. *Neurobiology of Aging*, *112*, 87–101. https://doi.org/10.1016/j.neurobiolaging.2021.11.009

Augustinova, M., Parris, B. A., & Ferrand, L. (2019). The Loci of Stroop Interference and Facilitation Effects With Manual and Vocal Responses. *Frontiers in Psychology*, *10*. https://www.frontiersin.org/articles/10.3389/fpsyg.2019.01786

Baddeley, A. (2012). Working Memory: Theories, Models, and Controversies. *Annual Review of Psychology*, *63*(Volume 63, 2012), 1–29. https://doi.org/10.1146/annurev-psych-120710-100422

Baker, S. N., Spinks, R., Jackson, A., & Lemon, R. N. (2001). Synchronization in Monkey Motor Cortex During a Precision Grip Task. I. Task-Dependent Modulation in Single-Unit Synchrony. *Journal of Neurophysiology*, *85*(2), 869–885. https://doi.org/10.1152/jn.2001.85.2.869

Banich, M. T. (2019). The Stroop Effect Occurs at Multiple Points Along a Cascade of

Control: Evidence From Cognitive Neuroscience Approaches. *Frontiers in*

Psychology, *10*, 2164. https://doi.org/10.3389/fpsyg.2019.02164

Beste, C., Münchau, A., & Frings, C. (2023). Towards a systematization of brain

oscillatory activity in actions. *Communications Biology*, *6*(1), Article 1.

https://doi.org/10.1038/s42003-023-04531-9

Böttcher, A., Wilken, S., Adelhöfer, N., Raab, M., Hoffmann, S., & Beste, C. (2023).

A dissociable functional relevance of theta- and beta-band activities during

complex sensorimotor integration. *Cerebral Cortex*, *33*(14), 9154–9164.

https://doi.org/10.1093/cercor/bhad191

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Brilliant, Yaar-Soffer, Y., Herrmann, C. S., Henkin, Y., & Kral, A. (2024). Theta and

alpha oscillatory signatures of auditory sensory and cognitive loads during

complex listening. *NeuroImage*, *289*, 120546.

https://doi.org/10.1016/j.neuroimage.2024.120546

Brockhoff, L., Schindler, S., Bruchmann, M., & Straube, T. (2022). Effects of

perceptual and working memory load on brain responses to task-irrelevant

stimuli: Review and implications for future research. *Neuroscience &*

Biobehavioral Reviews, *135*, 104580.

https://doi.org/10.1016/j.neubiorev.2022.104580

Bugg, J. M., & Crump, M. J. C. (2012). In Support of a Distinction between Voluntary

and Stimulus-Driven Control: A Review of the Literature on Proportion

Congruent Effects. *Frontiers in Psychology*, *3*.

https://doi.org/10.3389/fpsyg.2012.00367

Cavanagh, J. F., Cohen, M. X., & Allen, J. J. B. (2009). Prelude to and Resolution of

an Error: EEG Phase Synchrony Reveals Cognitive Control Dynamics during

Action Monitoring. *Journal of Neuroscience*, *29*(1), 98–105.

https://doi.org/10.1523/JNEUROSCI.4137-08.2009

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive

control. *Trends in Cognitive Sciences*, *18*(8), 414–421.

https://doi.org/10.1016/j.tics.2014.04.012

Chang, C.-C., & Lin, C.-J. (2011). LIBSVM: A library for support vector machines.

ACM Transactions on Intelligent Systems and Technology, *2*(3), 27:1-27:27.

https://doi.org/10.1145/1961189.1961199

Chaoran, Y. (2019). Pictographs of gastroparesis in Chinese characters.

Neurogastroenterology and Motility, *31*(8). https://doi.org/10.1111/NMO.13660

Chen, Z., Lei, X., Ding, C., Li, H., & Chen, A. (2013). The neural mechanisms of

semantic and response conflicts: An fMRI study of practice-related effects in the

Stroop task. *NeuroImage*, *66*, 577–584.

https://doi.org/10.1016/j.neuroimage.2012.10.028

Chung, R. S., Cavaleri, J., Sundaram, S., Gilbert, Z. D., Del Campo-Vera, R. M.,

Leonor, A., Tang, A. M., Chen, K.-H., Sebastian, R., Shao, A., Kammen, A.,

Tabarsi, E., Gogia, A. S., Mason, X., Heck, C., Liu, C. Y., Kellis, S. S., & Lee, B.

601 (2024). Understanding the human conflict processing network: A review of the

602 literature on direct neural recordings during performance of a modified stroop

603 task. *Neuroscience Research*. https://doi.org/10.1016/j.neures.2024.03.006

604 Coderre, E., Conklin, K., & van Heuven, W. J. B. (2011). Electrophysiological

605 measures of conflict detection and resolution in the Stroop task. *Brain Research*,

606 *1413*, 51–59. https://doi.org/10.1016/j.brainres.2011.07.017

607 Cohen, M. X., & Donner, T. H. (2013). Midfrontal conflict-related theta-band power

608 reflects neural oscillations that predict behavior. *Journal of Neurophysiology*,

609 *110*(12), 2752–2763. https://doi.org/10.1152/jn.00479.2013

610 David, I. A., Volchan, E., Vila, J., Keil, A., De Oliveira, L., Faria-Júnior, A. J. P.,

611 Perakakis, P., Dias, E. C., Mocaiber, I., Pereira, M. G., & Machado-Pinheiro, W.

612 (2011). Stroop matching task: Role of feature selection and temporal modulation.

613 *Experimental Brain Research*, *208*(4), 595–605. https://doi.org/10.1007/s00221-

614 010-2507-9

615 Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of

616 single-trial EEG dynamics including independent component analysis. *Journal of*

617 *Neuroscience Methods*, *134*(1), 9–21.

618 https://doi.org/10.1016/j.jneumeth.2003.10.009

619 Di Russo, F., & Bianco, V. (2023). Time Course of Reactive Brain Activities during a

620 Stroop Color-Word Task: Evidence of Specific Facilitation and Interference

621 Effects. *Brain Sciences*, *13*(7), Article 7.

622 https://doi.org/10.3390/brainsci13070982

623 Donohue, S. E., Appelbaum, L. G., McKay, C. C., & Woldorff, M. G. (2016). The

624  neural dynamics of stimulus and response conflict processing as a function of

625  response complexity and task demands. *Neuropsychologia*, *84*, 14–28.

626  https://doi.org/10.1016/j.neuropsychologia.2016.01.035

627 Edelman, S. (1998). Representation is representation of similarities. *The Behavioral*

628  *and Brain Sciences*, *21*(4), 449–467; discussion 467-498.

629  https://doi.org/10.1017/s0140525x98001253

630 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible

631  statistical power analysis program for the social, behavioral, and biomedical

632  sciences. *Behavior Research Methods*, *39*(2), 175–191.

633  https://doi.org/10.3758/BF03193146

634 Freund, M. C., Etzel, J. A., & Braver, T. S. (2021). Neural Coding of Cognitive

635  Control: The Representational Similarity Analysis Approach. *Trends in Cognitive*

636  *Sciences*, *25*(7), 622–638. https://doi.org/10.1016/j.tics.2021.03.011

637 Friehs, M. A., Klaus, J., Singh, T., Frings, C., & Hartwigsen, G. (2020). Perturbation

638  of the right prefrontal cortex disrupts interference control. *NeuroImage*, *222*,

639  117279. https://doi.org/10.1016/j.neuroimage.2020.117279

640 Haciahmet, C. C., Frings, C., Beste, C., Münchau, A., & Pastötter, B. (2023).

641  Posterior delta/theta EEG activity as an early signal of Stroop conflict detection.

642  *Psychophysiology*, *60*(3), e14195. https://doi.org/10.1111/psyp.14195

643 Hallenbeck, G. E., Sprague, T. C., Rahmati, M., Sreenivasan, K. K., & Curtis, C. E.

644  (2021). Working memory representations in visual cortex mediate distraction

effects. *Nature Communications*, *12*(1), 4714. https://doi.org/10.1038/s41467-

021-24973-1

Heidlmayr, K., Kihlstedt, M., & Isel, F. (2020). A review on the

electroencephalography markers of Stroop executive control processes. *Brain*

and Cognition, *146*, 105637. https://doi.org/10.1016/j.bandc.2020.105637

Itthipuripat, S., Deering, S., & Serences, J. (2019). When Conflict Cannot be Avoided:

Relative Contributions of Early Selection and Frontal Executive Control in

Mitigating Stroop Conflict. *Cerebral Cortex*, *29*, 1–12.

https://doi.org/10.1093/cercor/bhz042

Jenkinson, N., & Brown, P. (2011). New insights into the relationship between

dopamine, beta oscillations and motor function. *Trends in Neurosciences*, *34*(12),

611–618. https://doi.org/10.1016/j.tins.2011.09.003

Kaiser, J., & Schütz-Bosbach, S. (2021). Motor Interference, But Not Sensory

Interference, Increases Midfrontal Theta Activity and Brain Synchronization

during Reactive Control. *The Journal of Neuroscience*, *41*(8), 1788–1801.

https://doi.org/10.1523/JNEUROSCI.1682-20.2020

Kim, S.-Y., Kim, M.-S., & Chun, M. M. (2005). Concurrent working memory load

can reduce distraction. *Proceedings of the National Academy of Sciences of the*

United States of America, *102*(45), 16524–16529.

https://doi.org/10.1073/pnas.0505454102

Klimesch, W., Hanslmayr, S., Sauseng, P., Gruber, W. R., & Doppelmayr, M. (2007).

P1 and Traveling Alpha Waves: Evidence for Evoked Oscillations. *Journal of*

Neurophysiology, *97*(2), 1311–1318. https://doi.org/10.1152/jn.00876.2006

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating

cognition, computation, and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–

412. https://doi.org/10.1016/j.tics.2013.06.007

Li, Q., Hu, N., Li, Y., Long, Q., Gu, Y., Tang, Y., & Chen, A. (2021). Error-induced

adaptability: Behavioral and neural dynamics of response-stimulus interval

modulations on posterror slowing. *Journal of Experimental Psychology:*

General, *150*(5), 851–863. https://doi.org/10.1037/xge0000978

Li, Q., Yin, S., Wang, J., Zhang, M., Li, Z., Chen, X., & Chen, A. (2023). Not all

errors are created equal: Decoding the error-processing mechanisms using alpha

oscillations. *Cerebral Cortex*, bhad102. https://doi.org/10.1093/cercor/bhad102

Lin, Y., Li, Q., & Chen, A. (2023). The causal mechanisms underlying analogical

reasoning performance improvement by executive attention intervention. *Human*

Brain Mapping, *44*(8), 3241–3253. https://doi.org/10.1002/hbm.26278

Luna, F. G., Telga, M., Vadillo, M. A., & Lupiáñez, J. (2020). Concurrent working

memory load may increase or reduce cognitive interference depending on the

attentional set. *Journal of Experimental Psychology: Human Perception and*

Performance, *46*(7), 667–680. https://doi.org/10.1037/xhp0000740

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and

MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Meier, T. B., Desphande, A. S., Vergun, S., Nair, V. A., Song, J., Biswal, B. B.,

Meyerand, M. E., Birn, R. M., & Prabhakaran, V. (2012). Support vector

machine classification and characterization of age-related reorganization of

functional brain networks. *NeuroImage*, *60*(1), 601–613.

https://doi.org/10.1016/j.neuroimage.2011.12.052

Mensen, A., & Khatami, R. (2013). Advanced EEG analysis using threshold-free

cluster-enhancement and non-parametric statistics. *NeuroImage*, *67*, 111–118.

https://doi.org/10.1016/j.neuroimage.2012.10.027

Mouraux, A., & Iannetti, G. D. (2008). Across-trial averaging of event-related EEG

responses and beyond. *Magnetic Resonance Imaging*, *26*(7), 1041–1054.

https://doi.org/10.1016/j.mri.2008.01.011

Murphy, G., Groeger, J. A., & Greene, C. M. (2016). Twenty years of load theory—

Where are we now, and where should we go next? *Psychonomic Bulletin &*

Review, *23*(5), 1316–1340. https://doi.org/10.3758/s13423-015-0982-5

Okayasu, M., Inukai, T., Tanaka, D., Tsumura, K., Shintaki, R., Takeda, M., Nakahara,

K., & Jimura, K. (2023). The Stroop effect involves an excitatory–inhibitory

fronto-cerebellar loop. *Nature Communications*, *14*(1), Article 1.

https://doi.org/10.1038/s41467-022-35397-w

Park, S., Kim, M.-S., & Chun, M. (2007). Concurrent Working Memory Load Can

Facilitate Selective Attention: Evidence for Specialized Load. *Journal of*

Experimental Psychology. Human Perception and Performance, *33*, 1062–1075.

https://doi.org/10.1037/0096-1523.33.5.1062

Parris, B., Hasshim, N., Wadsley, M., Augustinova, M., & Ferrand, L. (2022). The loci

of Stroop effects: A critical review of methods and evidence for levels of

processing contributing to color-word Stroop effects and the implications for the

loci of attentional selection. *Psychological Research*, *86*, 1029–1053.

https://doi.org/10.1007/s00426-021-01554-x

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics:

Transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442.

https://doi.org/10.1163/156856897X00366

Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG

synchronization and desynchronization: Basic principles. *Clinical*

Neurophysiology: Official Journal of the International Federation of Clinical

Neurophysiology, *110*(11), 1842–1857. https://doi.org/10.1016/s1388-

2457(99)00141-8

Rypma, B., Berger, J. S., Prabhakaran, V., Martin Bly, B., Kimberg, D. Y., Biswal, B.

B., & D'Esposito, M. (2006). Neural correlates of cognitive efficiency.

NeuroImage, *33*(3), 969–979. https://doi.org/10.1016/j.neuroimage.2006.05.065

Schmidt, J. R. (2019). Evidence against conflict monitoring and adaptation: An

updated review. *Psychonomic Bulletin & Review*, *26*(3), 753–771.

https://doi.org/10.3758/s13423-018-1520-z

Senoussi, M., Verbeke, P., Desender, K., De Loof, E., Talsma, D., & Verguts, T.

(2022). Theta oscillations shift towards optimal frequency for cognitive control.

Nature Human Behaviour, *6*(7), Article 7. https://doi.org/10.1038/s41562-022-

01335-5

Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of*

Experimental Psychology, *18*(6), 643–662. https://doi.org/10.1037/h0054651

Tafuro, A., Ambrosini, E., Puccioni, O., & Vallesi, A. (2019). Brain oscillations in

cognitive control: A cross-sectional study with a spatial stroop task.

Neuropsychologia, *133*, 107190.

https://doi.org/10.1016/j.neuropsychologia.2019.107190

Van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-

band activity in attentional processing: The current zeitgeist and future outlook.

Current Opinion in Psychology, *29*, 229–238.

https://doi.org/10.1016/j.copsyc.2019.03.015

Viviani, G., Visalli, A., Montefinese, M., Vallesi, A., & Ambrosini, E. (2023). The

Stroop legacy: A cautionary tale on methodological issues and a proposed spatial

solution. *Behavior Research Methods*. https://doi.org/10.3758/s13428-023-

02215-0

Vo, K. D., Siqi-Liu, A., Chaire, A., Li, S., Demeter, E., Egner, T., & Woldorff, M. G.

(2021). Neural Dynamics of Conflict Control in Working Memory. *Journal of*

Cognitive Neuroscience, *33*(10), 2079–2092.

https://doi.org/10.1162/jocn_a_01744

Wen, W., Huang, Z., Hou, Y., & Li, S. (2022). Tracking Neural Markers of Template

Formation and Implementation in Attentional Inhibition under Different

Distractor Consistency. *Journal of Neuroscience*, *42*(24), 4927–4936.

https://doi.org/10.1523/JNEUROSCI.1705-21.2022

Wendiggensen, P., Adelhöfer, N., Jamous, R., Mückschel, M., Takacs, A., Frings, C.,

Münchau, A., & Beste, C. (2022). Processing of embedded response plans is

modulated by an interplay of frontoparietal theta and beta activity. *Journal of*

Neurophysiology, *128*(3), 543–555. https://doi.org/10.1152/jn.00537.2021

Wendiggensen, P., Prochnow, A., Pscherer, C., Münchau, A., Frings, C., & Beste, C.

(2023). Interplay between alpha and theta band activity enables management of

perception-action representations for goal-directed behavior. *Communications*

Biology, *6*(1), Article 1. https://doi.org/10.1038/s42003-023-04878-z

West, R., Jakubek, K., Wymbs, N., Perry, M., & Moore, K. (2005). Neural correlates

of conflict processing. *Experimental Brain Research*, *167*(1), 38–48.

https://doi.org/10.1007/s00221-005-2366-y

Wolff, M. J., Kandemir, G., Stokes, M. G., & Akyürek, E. G. (2020). Unimodal and

Bimodal Access to Sensory Working Memories by Auditory and Visual

Impulses. *Journal of Neuroscience*, *40*(3), 671–681.

https://doi.org/10.1523/JNEUROSCI.1194-19.2019

Xu, H., Yang, G., Wu, H., Xiao, J., Li, Q., & Liu, X. (2024). Distinct mechanisms

underlying cross-modal semantic conflict and response conflict processing.

Cerebral Cortex, *34*(2), bhad539. https://doi.org/10.1093/cercor/bhad539

Yang, G., Wu, H., Li, Q., Liu, X., Fu, Z., & Jiang, J. (2023). Conflicts are represented

in a cognitive space to reconcile domain-general and domain-specific cognitive

control. *eLife*, *12*. https://doi.org/10.7554/eLife.87126.1

Yu, B., Wang, X., Ma, L., Li, L., & Li, H. (2015). The Complex Pre-Execution Stage

777   of Auditory Cognitive Control: ERPs Evidence from Stroop Tasks. *PLOS ONE*,

778   *10*(9), e0137649. https://doi.org/10.1371/journal.pone.0137649

779 Zhang, X., Yang, S., & Jiang, M. (2020). Rapid implicit extraction of abstract

780   orthographic patterns of Chinese characters during reading. *PLOS ONE*, *15*(2),

781   e0229590. https://doi.org/10.1371/journal.pone.0229590

782 Zhao, J., Liang, W.-K., Juan, C.-H., Wang, L., Wang, S., & Zhu, Z. (2015).

783   Dissociated stimulus and response conflict effect in the Stroop task: Evidence

784   from evoked brain potentials and brain oscillations. *Biological Psychology*, *104*,

785   130–138. https://doi.org/10.1016/j.biopsycho.2014.12.001

786 Zhao, X., Chen, A., & West, R. (2010). The influence of working memory load on the

787   Simon effect. *Psychonomic Bulletin & Review*, *17*(5), 687–692.

788   https://doi.org/10.3758/PBR.17.5.687

789 Zhao, Y., Tang, D., Hu, L., Zhang, L., Hitchman, G., Wang, L., & Chen, A. (2014).

790   Concurrent working memory task decreases the Stroop interference effect as

791   indexed by the decreased theta oscillations. *Neuroscience*, *262*, 92–106.

792   https://doi.org/10.1016/j.neuroscience.2013.12.052

793 Zhou, Y. J., Ramchandran, A., & Haegens, S. (2023). Alpha oscillations protect

794   working memory against distracters in a modality-specific way. *NeuroImage*,

795   *278*, 120290. https://doi.org/10.1016/j.neuroimage.2023.120290

796 Zhu, Y., Wu, D., Sun, K., Chen, X., Wang, Y., He, Y., & Xiao, W. (2023). Alpha and

797   Theta Oscillations Are Causally Linked to Interference Inhibition: Evidence from

798   High-Definition Transcranial Alternating Current Stimulation. *Brain Sciences*,

799   *13*(7), Article 7. https://doi.org/10.3390/brainsci13071026

800   Zivony, A., & Lamy, D. (2022). What processes are disrupted during the attentional

801   blink? An integrative review of event-related potential research. *Psychonomic*

802   *Bulletin & Review*, *29*(2), 394–414. https://doi.org/10.3758/s13423-021-01973-2

804
805 **Fig. 1.** Raincloud plots (Allen et al., 2021) of behavioral Data. The plot consists of a probability
806 density plot, a line graph, and raw data points. ** $p < 0.01$, *** $p < 0.001$.

Fig. 2. The P1 results in the single task and the dual task. (A) The event-related potential (ERP) waveforms for P1 component. The shaded rectangle represents the defined time windows and the shading above and below the ERP waveform represents the 95% confidence intervals. (B) Topographic distributions of average amplitude for each condition within the time window (110-150 ms) and the amplitude differences between congruent and incongruent trials in single and dual tasks. (C) Raincloud plots of amplitude for P1 across conditions. n.s., not significant.

Fig. 3. The N450 and SP results in the single task and the dual task. (A) The event-related potential (ERP) waveforms for P1 and SP components. The shaded rectangle represents the defined time windows and the shading above and below the ERP waveform represents the 95% confidence intervals. (B) Topographic distributions of average amplitude for each condition within the time window (350-500 ms) and the amplitude differences between congruent and incongruent trials in single and dual tasks. (C) Raincloud plots of amplitude for N450 across conditions. (D) Topographic distributions of average amplitude for each condition within the time window (650-950 ms) and the amplitude differences between congruent and incongruent trials in single and dual tasks. (E) Raincloud plots of amplitude for SP across conditions. n.s., not significant, * $p < 0.05$, ** $p < 0.01$.

Fig. 4. MVPA results for (A) the theta band, (B) the alpha band, and (C) the beta band. Dots denote significant differences from chance (P < 0.01). Shading represents clusters of significant time points (cluster-corrected).

Fig. 5. Time-frequency results for the theta band. (A) Grand-average time-frequency representations for the theta band over fronto-central area (Fz, FCz, Cz) for each condition, and the difference activity between congruent and incongruent trials in single and dual tasks. (B) Topographic distributions of average amplitude for each condition within the defined time-frequency region of interest (4-7 Hz, 740-820 ms) and the magnitude differences between congruent and incongruent trials in single and dual tasks. (C) Raincloud plots of ERSP magnitudes for theta band across conditions. n.s., not significant. ** $p < 0.01$, *** $p < 0.001$.

Fig. 6. Time-frequency results for the beta band. (A) Grand-average time-frequency representations for the beta band over the central area (C3, CP3, Cz, CP4, C4) for each condition, and the difference activity between congruent and incongruent trials in single and dual tasks. (B) Topographic distributions of average amplitude for each condition within the defined time-frequency region of interest (14-25 Hz, 920-1040 ms) and the magnitude differences between congruent and incongruent trials in single and dual tasks. (C) Raincloud plots of ERSP magnitudes for the beta band across conditions. n.s., not significant. *** $p < 0.001$.

Fig. 7. Representational dissimilarity matrices for the interaction pattern in RT, SP, theta and beta. Each RDM is calculated as the Euclidean distances of data reflecting interaction pattern from 32 subjects.

Fig. 8. Schematic illustration of experimental trial structures and stimuli types. (A) Dual-task structure, (B) Single-task structure, (C) Representative stimuli for congruent and incongruent conditions.