Sex-dependent gastrointestinal colonization resistance to MRSA is microbiota and Th17

dependent

Alannah Lejeune [1,2], Chunyi Zhou [1,2], Defne Ercelen [3], Gregory Putzel [1,4], Xiaomin Yao [2], Alyson R Guy [5],

Miranda Pawline [3], Magdalena Podkowik [2,4], Alejandro Pironti[1,4], Victor J. Torres [1,6,*], Bo Shopsin [1,2,4*], Ken

Cadwell[7,8,*]

[1]Department of Microbiology, New York University School of Medicine, New York, NY, 10016, USA.

[2]Department of Medicine, Division of Infectious Diseases, New York University School of Medicine, New York, NY 10016, USA.

[3]Department of Medicine, Division of Gastroenterology and Hepatology, New York University Langone Health, New York, NY,

10016, USA.

[4]Antimicrobial-Resistant Pathogens Program, New York University School of Medicine, United States

[5]NYU-Regeneron Veterinary Postdoctoral Training Program in Laboratory Animal Medicine, Division of Comparative Medicine, New

York University School of Medicine, New York, NY 10016, USA.

[6]Department of Host-Microbe Interactions, St. Jude Children's Research Hospital, Memphis, TN, 38105, USA.

[7]Department of Medicine, Division of Gastroenterology and Hepatology, University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA, 19104, USA.

[8]Department of Pathobiology, University of Pennsylvania Perelman School of Veterinary Medicine, Philadelphia, PA, 19104, USA.

*Co-corresponding

Contact: ken.cadwell@pennmedicine.upenn.edu, bo.shopsin@nyulangone.org, victor.torres@stjude.org

ABSTRACT

Gastrointestinal (GI) colonization by methicillin-resistant *Staphylococcus aureus* (MRSA) is

associated with a high risk of transmission and invasive disease in vulnerable populations. The

immune and microbial factors that permit GI colonization remain unknown. Male sex is

correlated with enhanced *Staphylococcus aureus* nasal carriage, skin and soft tissue infections,

and bacterial sepsis. Here, we established a mouse model of sexual dimorphism during GI

colonization by MRSA. Our results show that in contrast to male mice that were susceptible to

persistent colonization, female mice rapidly cleared MRSA from the GI tract following oral

inoculation in a manner dependent on the gut microbiota. This colonization resistance displayed

by female mice was mediated by an increase in IL-17A+ CD4+ T cells (Th17) and dependent on

neutrophils. Ovariectomy of female mice increased MRSA burden, but gonadal female mice that

have the Y chromosome retained enhanced Th17 responses and colonization resistance. Our

study reveals a novel intersection between sex and gut microbiota underlying colonization

resistance against a major widespread pathogen.

INTRODUCTION

Methicillin resistant *Staphylococcus aureus* (MRSA) is a major global health concern due to its multidrug resistance, wide range of infections, and high morbidity and mortality rates[1–4]. Gastrointestinal *S. aureus* colonization is present in an estimated 20% of the healthy population[5,6], and an estimated 1-6% are persistently colonized with MRSA[5,6]. MRSA carriage increases the likelihood of invasive infections, especially while in the hospital or post discharge[7,8]. Intestinal carriage is associated with higher rates of infections and bacteremia than nasal carriage alone[4,9]. The risk of transmission to other individuals in the hospital and community settings through fomite spread is increased by intestinal carriage[10,11].

A better understanding of why a subset of individuals are susceptible to colonization may inform decolonization strategies[5–7]. Although little is known mechanistically, population level studies have identified correlates of colonization. For example, male sex is correlated with *S. aureus* nasal carriage, skin and soft tissue infections and bacterial sepsis in adulthood[12–15]. High free testosterone levels are a risk factor for *S. aureus* throat carriage in women[12]. However, experimental systems are necessary to clarify to what extent male sex as a risk factor for carriage is biological or behaviorally based.

Sex steroid hormones and sex chromosomes can modulate the scale and type of immune response[16–18]. Sex hormones such as estrogen and testosterone can directly influence immune cell activation, proliferation and cytokine response through receptor signaling[19–23]. In general, males are more susceptible to gastrointestinal and respiratory infections, whereas females are more affected by autoimmune diseases in part due to the pro-inflammatory effect of estradiol, but these responses are tissue and cell specific[17,19]. In the few studies examining GI MRSA colonization in mice, the focus has been on the adaptation of *S. aureus* to the host[24–27]. These studies generally rely on depletion of the mouse microbiota with antibiotics to establish long-term colonization[24,25,28]. To investigate mechanisms of colonization resistance in a setting with an intact microbiota, we established a GI colonization model that does not rely on antibiotic treatment, better recapitulating MRSA colonization in a healthy host. We report that following MRSA oral inoculation, female mice with a non-permissive microbiota were resistant to sustained colonization compared with male mice that remain persistently colonized. Colonization resistance displayed by female mice was mediated by an increase in T helper 17 (Th17) cells and associated with sex hormones rather than sex chromosomes. Thus, our study demonstrates a role for the Th17 response and microbiota in GI colonization resistance against MRSA, while also highlighting the sex-dependent susceptibility to this major pathogen.

RESULTS

Female mice are protected from MRSA gastrointestinal colonization in a microbiota dependent manner.

We and others have shown that inbred laboratory mice from different sources, or even the same institution, can display substantial differences in mucosal immune responses and susceptibility to GI colonization by microorganisms[29–34]. Therefore, when we set out to establish a model of GI MRSA colonization, we examined two sources of C57BL/6J (B6) mice - those bred in Jackson Laboratory (JAX) and genetically identical mice bred within our institutional animal facility (referred to as NYU mice). Adult B6 mice received a single oral gavage with 10^8 colony forming units (CFU) of MRSA USA300 LAC, the predominant MRSA clone responsible for community associated MRSA infections and a growing number of hospital-acquired infections in the United States[2]. Prior to inoculation, we confirmed the absence of pre-existing *S. aureus* colonization by plating stool from individual mice on ChromAgar plates which select for *S. aureus* growth. We confirmed that mice inoculated with MRSA do not display signs of disease such as weight loss **(Supplemental Fig1 A)**. Consistent with previous findings[35], we did not observe extraintestinal dissemination of bacteria or histological signs of intestinal inflammation **(Supplemental Fig1 B-D).**

In JAX mice, we detected 10^4 to 10^5 CFU per gram of stool of MRSA on day 2 that remained stable for at least 35 days post-inoculation (dpi) **(Fig1 A)**. However, we noticed bimodal distribution of the data for NYU mice where some mice remained stably colonized while detectable MRSA burden diminished rapidly in others. Segregation of the groups by sex revealed that NYU mice that displayed resistance to MRSA colonization were females. Despite originating from the same litters, male NYU mice displayed sustained colonization at levels similar to male and female JAX mice, while MRSA levels were already lower by day 2 and undetectable by 2-3 weeks post inoculation in female NYU mice **(Fig1 B-C)**. Commensal microbes that are constituents of the microbiota can inhibit *S. aureus* nasal, skin, and intestinal colonization[36–38]. In support of a role for commensal microbes in promoting colonization resistance in NYU female mice, we observed stable MRSA GI colonization in germ-free (GF) mice of both sexes **(Fig1 D).** We next examined germ-free mice that were colonized with a defined consortium of 15 bacterial strains representative of a mouse gut microbiota (Oligo-MM$_{12}$ + FA3), previously shown to be sufficient to confer colonization resistance against the enteric pathogen *Salmonella enterica* serovar Typhimurium[39]. We observed sustained MRSA GI colonization and no sex difference in these minimal flora mice, indicating that additional intestinal commensals contribute to sex-dependent colonization resistance **(Fig1 E).**

Given these results, we examined whether we could transfer the MRSA resistance displayed by
NYU female mice to permissive JAX female mice through co-housing mice in the same cage,
which allows for the transfer of microbiota between mice[40]. JAX and NYU female mice were co-
housed together for a week prior to MRSA inoculation and kept together for the duration of the
experiment. When housed together, JAX female mice cleared MRSA colonization similarly to
their NYU cage mates **(Fig1 F,G)** indicating that the microbiota of NYU female mice promotes
resistance to colonization and that this protective property can be transferred.

Fig.1 Female mice are protected from MRSA gastrointestinal colonization in a microbiota
dependent manner.
(A) MRSA colony forming units (CFU) per gram of stool following oral inoculation of B6 mice
purchased from Jackson Laboratory (JAX). Males n=10, females n=13. **(B)** MRSA CFU in stool
following oral inoculation of B6 mice generated from breeders in the NYU animal facility. Males
n=13, females n=13. **(C)** Proportion of JAX and NYU female mice with detectable MRSA GI
colonization over time. **(D)** MRSA CFU stool burden following oral inoculation of germ-free mice.
Males n=8, females n=7. **(E)** MRSA CFU in stool following oral inoculation of mice with a
defined minimal microbiota. Males n=12, females n=11. **(F)** MRSA CFU stool burden of female
NYU or JAX mice housed separately or **(G)** co-housed with each other. NYU n=9, JAX n=11,
cohoused NYU n=14, cohoused JAX n=14. Data points represent mean ± SEM from at least

two independent experiments. Statistical analysis: area under the curve followed by a two-tailed

t-test for (A), (B), (D)-(G) and Log-rank Mantel Cox test for (C). ns: not significant.

Microbiota is not sufficient to explain sex bias in MRSA GI colonization.

126   To compare the microbiome composition of male and female NYU and JAX mice, we

performed 16S ribosomal DNA sequencing of stool collected prior to inoculation with MRSA.

Principal Coordinates Analysis (PCoA) of operational taxonomic units (OTUs) showed that

samples were distinguished based on the source of mice (NYU versus JAX) rather than sex

(Fig2 A). Overall alpha diversity as measured by Shannon index was increased in JAX mice

compared to NYU mice prior to MRSA inoculation, as well as in JAX females compared to NYU

females **(Fig2 B)**. Alpha diversity by Shannon index between NYU males and females shows a

similar spread when comparing sexes **(Fig2 B)**. JAX females had higher abundance of

Clostridiaceae and *Lachnospiraceae* family members, while NYU female microbiomes were

dominated by the *Muribaculaceae* family **(Fig2 C)**. The NYU male and female microbiomes

clustered together **(Fig2 A)** and had similar relative taxonomic abundances **(Fig2 D).** These

findings suggest that the observed sex bias in colonization resistance may not be due to

microbiome composition alone.

139   As male and female mice are housed separately post weaning, and weaning is

associated with microbiome changes[41], we tested if the colonization resistance displayed by

female NYU mice could be transferred to male NYU mice. Instead of co-housing male and

female mice, which could introduce variables through social stress or changes in hormone

levels due to mating,[42] we performed fecal microbiota transplantations (FMT) by repetitively

gavaging recipient mice with donor stool. JAX female mice receiving a FMT from NYU female,

but not male donor mice, displayed colonization resistance to MRSA. However, an FMT using

NYU female donor stool was unable to confer colonization resistance to JAX male recipients

(Fig2 E). Therefore, the microbiota contributes to the difference between female JAX and NYU

mice, but other factors mediate the divergence of female and male NYU mice. From here on, we

use NYU bred mice to investigate host factors that mediate this sex specific colonization

resistance.

Fig. 2 Microbiome is not sufficient to explain sex bias in MRSA GI colonization.

(A) Principal Coordinates Analysis (PCoA) based on Bray-Curtis distances of 16S sequences

obtained from stool of JAX and NYU male and female mice prior to MRSA inoculation.

Proportion of variance explained by each axes shown in parentheses. JAX M n=5, JAX F n=5,

NYU M n=11, NYU F n=13. **(B)** Alpha diversity of the microbiomes of male and female JAX and

NYU mice. Wilcoxon rank sum test used for pairwise statistical comparisons to NYU females;

ns: not significant. **(C)** Relative abundance of bacterial families in female NYU and JAX mice

prior to MRSA inoculation. **(D)** Relative abundance of bacterial families in male and female NYU

mice prior to MRSA inoculation. **(E)** MRSA colony forming units (CFU) in stool following oral

inoculation of JAX female (JAX F) and male (JAX M) recipients of fecal microbiome transplants

(FMT) from female (NYU F) and male (NYU M) donor mice compared with JAX male controls

(control JAX M) that did not receive an FMT. Mean MRSA burden ± SEM. Area under the curve

analysis + 1 way ANOVA Sidak's multiple comparisons test for (E). Control M n=6, M +NYU F

stool n=7, M +NYU M stool n=6, F +NYU F stool n=6. ns: not significant.

MRSA GI colonization elicits sex-dependent gene expression changes in the gut

To gain insight into the host response, we performed bulk RNA-seq analysis of cecal-

colonic tissue obtained two days post inoculation (2 dpi) of NYU mice (both sexes) with either

MRSA or PBS (mock), as 2 dpi is the timepoint at which MRSA burden begins to diverge

between males and females. The lamina propria compartment, enriched with immune cells, and

the epithelial cell fraction were isolated and sequenced separately. In the lamina propria, 795

genes were upregulated in both male and female mice following MRSA inoculation, with an

additional 352 genes upregulated in females only and 648 genes upregulated in males only, as

determined by a log2 fold change cutoff of 1.2 and a false discovery rate (FDR) of 0.05

(**Supplemental Fig2 A**). Following MRSA inoculation in both sexes, genes with the largest

downregulation included those involved with immunoglobulins (*Ighv1-19*, *Igkv4-91, Bcam*),

cytochrome p450 genes (*Cyp1a1, Cyp1b1*), and von Willebrand factor (*Vwf*), a secreted protein

that MRSA binds to disrupt platelet recruitment and coagulation[43] **(Fig3 A)**. Interestingly,

constitutive cytochrome P4501A1 (*Cyp1a1*) expression impairs the intestinal immune response

against enteric pathogens[44], and its downregulation in mice inoculated with MRSA may be

supportive of a mucosal immune response. Genes that were upregulated in the lamina propria

following MRSA inoculation included those involved in T cell and neutrophil activity such as

Ccl28[45], *Sectm1b*[46], *S100g*[47], *Lrrc19*[48], and *H2-q1*[49] **(Fig3 A)**.

Pathway analysis of the lamina propria fraction indicated that both males and females

have downregulation of immune related Th1, Th2, and granzyme A signaling pathways, and

upregulation of oxidative phosphorylation, neutrophil extracellular trap formation, glucose

metabolism, and xenobiotic metabolism pathways **(Fig3 B)**. Downregulation of interleukin (IL)-4

and IL-13 pathway and upregulation of retinoid X receptor (RXR) and the pregnane X receptor

(PXR) pathways were specific to female mice. Unique to male mice was the downregulation of

IL-5, IL-3, and GM-CSF signaling, and upregulation of the pyroptosis and ion channel transport

pathways.

Known X and Y chromosome linked genes *Xist*, *Ddx3y*, *Eif2s3y*, *Kdm5d*, and *Uty* were
differentially present in females and males as expected but were removed from male vs female
analysis plots to highlight other differentially expressed genes. Other than known sex
chromosome linked genes, when comparing male and female mice prior to MRSA inoculation,
we observed several genes without a descriptive name or suggested pseudogenes such as
Gm12070, Gm13772, and *Gm4968* **(Fig3 C)**. In the MRSA condition, there were 13 genes
differentially expressed in the intestinal lamina propria between male and female mice.
Glycosylation-dependent cell adhesion molecule-1 (*Glycam1*), which mediates lymphocyte
trafficking to lymphoid tissues[50], was upregulated in female mice **(Fig3 D)**. Genes selectively
upregulated in male mice 2 dpi included *Complement protein C7* and several immunoglobulin
heavy chain variants (*Ighv5-4, Ighv1-49, Ighv8-5*).

In the intestinal epithelial fraction, we observed only 15 genes upregulated in males 2 dpi
with MRSA, while 2,037 were upregulated in females with no overlap between sexes
(**Supplemental Fig2 B**). Among the most differentially downregulated genes following MRSA
inoculation when combining both sexes were *Irf5*, a pivotal transcription factor in the type I IFN
pathway, *Csf2rb*, part of the receptor for IL-3, IL-5, and CSF signaling, and *Ciita*, a mediator of
major histocompatibility complex activation. Among those upregulated were genes involved in
limiting inflammation like *Cav1*, involved in epithelial barrier maintenance, *Cfh*, a regulator of the
complement pathway, and *Slpi* which functions to protect tissue from the detrimental
consequences of inflammation **(Supplemental Fig2 C)**. Pathway analysis indicated that both
sexes had upregulation of cell cycle and rRNA processing pathways and downregulation of
neutrophil degranulation. Unique to females was upregulation of PTEN signaling and
SUMOlyation of DNA damage response pathways **(Supplemental Fig2 D)**. There were few
differentially expressed genes between males and females in the mock epithelial condition **(Fig
3E)** but following MRSA inoculation males had increased expression of the antimicrobial peptide
Defa30 and *Tnip3*, a negative regulator of NFκβ and Toll-like receptors **(Fig 3F)**. Thus, MRSA
inoculation elicits distinct transcriptional responses in males and females in both the lamina
propria and epithelial fractions, and many of the differentially regulated genes have known
functions in neutrophils and lymphocyte migration and function.
223
224

Fig 3. MRSA GI colonization elicits sex-dependent gene expression changes in the gut.
(A) Volcano plot of differentially expressed genes identified by RNA-seq analyses of the
intestinal lamina propria of NYU mice 2 dpi with MRSA compared with PBS mock inoculated
NYU mice. **(B)** Ingenuity Pathway Analysis (IPA) of downregulated and upregulated genes in
the intestinal lamina propria upon MRSA inoculation of male and female mice. **(C)** Volcano plot
of differentially expressed genes in the intestinal lamina propria comparing mock treated males
and females. **(D)** Volcano plot of differentially expressed genes in the intestinal lamina propria
comparing males and females 2 dpi with MRSA. **(E)** Volcano plot of differentially expressed
genes in the intestinal epithelial fraction of males and females prior to MRSA inoculation. **(F)**

235 Volcano plot of differentially expressed genes in the intestinal epithelial fraction of males and
236 females 2 dpi with MRSA. Four mice were used for each sequencing experimental condition.
237 Genes shown in red have a false discovery rate (FDR) of <0.05 and an absolute log2 fold
238 change (abslog2FC) of >1.2. Genes shown in blue have a FDR of <0.05. Genes linked to X and
239 Y chromosomes were removed from volcano plots.
240

CD4+ T cells mediate colonization resistance in female mice.

Female mice have been shown to be protected from dermonecrosis in a model of
invasive MRSA skin infection due to reduced expression of genes associated with the NLRP3
inflammasome (*Nlrp3* and *Il-1β*) compared to males[14]. However, we did not observe sex-specific
differences in NLRP3-associated transcripts with RNA-seq **(Supplemental Fig3 A,B)** and
female *Nlrp3*$^{-/-}$ mice displayed similar resistance to MRSA GI colonization as female *Nlrp3*$^{+/-}$
controls **(Supplemental Fig3 C)**. Thus, the mechanism of sex specific resistance to MRSA
colonization of the GI tract may be different from that observed during a skin infection where
barrier breach triggers an exaggerated innate immune response that may cause more harm
than benefit.

The T cell signature in the RNA-seq analyses of the intestinal lamina propria of MRSA
colonized mice was unexpected given that the day 2 time point is earlier than what is typically
required for an adaptive immune response. To test the role of lymphocyte responses in GI
colonization resistance, we measured bacteria in stool following oral inoculation with MRSA of
recombination activating gene 2 (*Rag2*) deficient mice, which lack mature B and T cells. We
observed no difference in burden between male *Rag2*$^{-/-}$ and *Rag2*$^{+/-}$ controls **(Fig4 A).**
However, female *Rag2*$^{-/-}$ mice had significantly increased MRSA burden compared to *Rag2*$^{+/-}$
littermates **(Fig4 A).** These data suggest that a B or T cell mediated response is required for the
sex bias we observe in colonization resistance.

B cell associated *Ighv* genes were differentially regulated between males and females.
However, *Igh-μ*$^{-/-}$ (μMT) mice, which lack mature B lymphocytes, were similar to *Igh-μ*$^{+/-}$
controls and retained sex differences; males displayed prolonged colonization and females were
resistant **(Fig4 B)**. In contrast, antibody-mediated depletion of CD4+ T cells **(Supplemental
Fig3 D)** substantially increased MRSA GI colonization of female mice, similar to *Rag2*$^{-/-}$ mice
(Fig4 C). Therefore, CD4+ T cells are required for MRSA colonization resistance associated
with female sex.
267

Fig 4. CD4+ T cells mediate colonization resistance in female mice.

(A) MRSA CFU in stool following oral inoculation of $Rag2^{-/-}$ and $Rag2^{+/-}$ mice bred at NYU. Male $Rag2^{-/-}$ n=12, male $Rag2^{+/-}$ n=12, female $Rag2^{-/-}$ n=12, female $Rag2^{+/-}$ n=11. **(B)** MRSA CFU in stool following oral inoculation of $Igh\text{-}\mu^{-/-}$ and $Igh\text{-}\mu^{+/-}$ mice bred at NYU. Male $Igh\text{-}\mu^{-/-}$ n=6, male $Igh\text{-}\mu^{+/-}$ n=6, female $Igh\text{-}\mu^{-/-}$ n=8, female $Igh\text{-}\mu^{+/-}$ n=5. **(C)** MRSA CFU in stool following oral inoculation of female NYU B6 mice injected I.P. with 250 µg of anti-CD4 depleting antibody or anti-IgG control. Anti-IgG n=8, anti-CD4 n=10. Data points represent mean ± SEM from at least two independent experiments. Statistical analysis: area under the curve analyzed by a 1 way ANOVA with Sidak's multiple comparison test for (A) and (B) and a two-tailed t test for (C). ns: not significant.

MRSA colonization resistance in female mice is dependent on Th17 cells and neutrophils.

In addition to T cells, our transcriptome analysis showed upregulation of genes associated with neutrophil function **(Fig3 B)**. In mouse models of nasal colonization, clearance is mediated by neutrophil influx downstream of the type 17 response, which includes interleukin-17A (IL-17A) producing lymphoid cells such as Th17 cells[51,52]. As an extracellular Gram-positive bacterium at the mucosal surface in the gut, MRSA may be subject to regulation by pre-existing Th17 cells[53–55]. Although the total numbers of CD4+ T cells were similar across conditions, we

observed an increase in IL-17A+ CD4+ T cells in the small intestine and colon of female mice

2dpi with MRSA that was not observed in males **(Fig5 A,B, Supplemental Fig4 A,B and 5**

A,B). There were no differences in γδ T cells and innate lymphoid cells (ILCs) or the proportion

of these lymphoid subsets that were IL-17A+ **(Fig5 C,D** and **Supplemental Fig5 C,D)**. To

determine the importance of the type 3 immune response that encompasses Th17 cells, we

inoculated *RAR-related orphan receptor gamma* (*Rorγt*) deficient mice that lack the

transcriptional regulator required for the differentiation of these cell types. Although there was

no difference in MRSA burden between male *Rorγt$^{+/-}$* and *Rorγt$^{-/-}$* mice that remained

colonized, female *Rorγt$^{-/-}$* mice had higher MRSA burden compared to *Rorγt$^{+/-}$* controls **(Fig5**

E). Responses to IL-17 itself was required, because female mice deficient in the IL-17 receptor

(*IL-17ra*) had increased MRSA burden compared to heterozygous littermates **(Fig5 F)**. There

was no difference in MRSA burden between *Tcrd$^{-/-}$* and *Tcrd$^{+/-}$* male or female mice **(Fig5 G)**,

confirming that γδ T cells were not required for colonization resistance. Taken together, these

data implicate Th17 cells and IL-17 in colonization resistance against MRSA observed in female

mice.

303   Th17 cells and IL-17 can recruit and promote antimicrobial functions of neutrophils[56].

Neutrophils from female mice and humans display signs of increased maturity and activation[57-

60]. Although we did not observe significant differences in the number of neutrophils between

conditions **(Fig5 H)**, neutrophils from female mice 2 dpi with MRSA had increased levels of

CD11b, a sign of activation[61], compared to males **(Fig5 I)**. Antibody- mediated depletion of

neutrophils **(Supplemental Fig5 E)** led to increased MRSA burden compared with isotype

control treated mice **(Fig5 J)**. Although colonization was increased in males, neutrophil

depletion in female mice led to an earlier and more pronounced increase in MRSA colonization

that was prolonged. These results are consistent with the RNA-seq analysis suggesting that a

neutrophil response occurs in both sexes. It is likely that neutrophils control bacterial burden to

some degree but fail to promote clearance in males. In contrast, neutrophils are required for

clearance of MRSA in the gut of female mice.

Fig 5. Female mice have increased IL-17A+ CD4+ T cells and require neutrophils for MRSA clearance.

(A) Flow cytometry of cecal-colonic lamina propria CD4+ T cells as a percentage of CD45+ cells in male and female NYU mice treated with PBS or MRSA 2 dpi. **(B)** Flow cytometry of cecal-colonic lamina propria IL17A+ CD4+ T cells as a percentageof total CD4+ T cells in male and female NYU mice treated with PBS or MRSA 2 dpi. **(C)** Flow cytometry of cecal-colonic lamina propria γδ T cells as a percentage of CD45+ cells in male and female NYU mice treated with PBS or MRSA 2 dpi. **(D)** Flow cytometry of cecal-colonic lamina propria IL17A+ γδ T cells as a percentage of total CD4+ T cells in male and female NYU mice treated with PBS or MRSA 2 dpi. **(E)** MRSA CFU in stool following oral inoculation of $Roryt^{-/-}$ and $Roryt^{+/-}$ mice bred at NYU. Male $Roryt^{+/-}$ n=5, male $Roryt^{-/-}$ n=7, female $Roryt^{+/-}$ n=9, female $Roryt^{-/-}$ n=9. **(F)** MRSA CFU in stool following oral inoculation of female $IL-17ra^{+/-}$ and $IL-17ra^{-/-}$ mice bred at NYU. $IL-17ra^{+/-}$ n=6, $IL-17ra^{-/-}$ n=6. **(G)** MRSA CFU in stool following oral inoculation of $Tcrd^{+/-}$ and $Tcrd^{-/-}$ mice bred at NYU. Male $Tcrd^{+/-}$ n=6, male $Tcrd^{-/-}$ n=6, female $Tcrd^{+/-}$ n=8, and female $Tcrd^{-/-}$ n=12. **(H)** Flow cytometry of cecal-colonic lamina propria Ly6G+CD11b+ neutrophils as a percentage

of CD45+ cells in male and female NYU mice treated with PBS or MRSA 2 dpi. **(I)** Quantification

of mean fluorescence intensity (MFI) of surface CD11b on neutrophils by flow cytometry

normalized to mock treated controls. **(J)** MRSA CFU in stool following oral inoculation of NYU

B6 mice treated with anti-Ly6G neutrophil depleting antibody or anti-IgG control. Male anti-IgG

n=6, male anti-Ly6G n=8, female anti-IgG n=12, and female anti-CD4+ n=15. Data points

represent mean ± SEM from at least two independent experiments. Statistical analysis: 2 way

ANOVA + Sidak's multiple comparisons test for (A)-(D) and (H), area under the curve followed

by a 1 way ANOVA + Sidak's multiple comparisons test for (E), (G), (J) or a two-tailed t test for

(F) and a two-tailed t test for (I). ns: not significant.

Sex hormones, not sex chromosomes, mediate MRSA colonization resistance in female

mice

Sex bias in immunity could be mediated by genes that are encoded on sex

chromosomes. Many X linked genes are involved in immunity, such as pattern recognition

receptors, and incomplete X inactivation can lead to higher expression of such genes[16]. To test

whether colonization resistance was due to sex chromosomes within a hematopoietic cell type,

we generated chimeras in which wildtype male mice were reconstituted with T-lymphocyte

depleted bone marrow (BM) from female donors and compared with male and female mice that

received BM from same sex donors. The reciprocal chimera in which female mice are

reconstituted with male BM was not feasible due to transplant rejection. Although female mice

that received female donor BM were resistant to MRSA, male mice that received male or female

donor BM remained colonized **(Fig6 A)**. Thus, it is unlikely that a gene expressed on a sex

chromosome in immune cells such as T cells mediates colonization resistance in females,

raising the possibility of a role for sex hormones in our model. The hormones estrogen,

progesterone, and testosterone are known to modulate trafficking and function of immune cells.

For example, estrogen can enhance responses to extracellular pathogens[21,62,63]. To test

whether sex hormones are involved in GI colonization resistance to MRSA, female mice at six

weeks of age were ovariectomized (OVX) or given a sham operation (SO) and allowed to

recover for two weeks prior to oral inoculation with MRSA. OVX mice had increased MRSA

burden and duration of carriage compared to SO controls **(Fig6 B)**. Estrogen receptor-α (ERα,

encoded by *Esr1*) signaling increases Th17 cell proliferation and production of IL-17[21,64]. We

confirmed *Esr1* was expressed in cecal-colonic immune cells, and that it displayed a non-

significant trend suggesting an increase upon MRSA inoculation **(Supplemental Fig6 A)**.

Female *Esr1*[-/-] mice had increased MRSA burden compared to heterozygous littermates **(Fig6**

C) indicating that estrogen mediates the enhanced colonization resistance in female mice.

To formally decouple the role of sex hormones and sex chromosome encoded gene

products, we used the mouse four core genotype (FCG) model in which the gene sex

determining region Y (*Sry*) that defines male sex has been moved from the Y chromosome to an

autosome[65]. Thus, the sex chromosome complement (XX or XY) does not relate to gonadal sex

in the FCG model. We found that XY*(-Sry)* gonadal female mice lose MRSA carriage analogous

to XX females, consistent with prior studies showing that they display similar estradiol levels[66,67],

and both XY and XX*(+Sry)* gonadal male mice remain persistently colonized at similar levels

(Fig6 D). Like their wildtype counterparts, we do not observe a change in the proportions of

CD4+ T cells in the lamina propria between XX males and XX females with and without MRSA

(Fig6 E). Instead, XX*(-Sry)* females displayed an increase in IL17A+ CD4+ T cells in the lamina

propria compared with XX(+*Sry*) males 2 dpi MRSA inoculation **(Fig6 F).** These findings further

support a hormone mediated effect on the Th17 cell response in female mice rather than a

chromosomal one.

Fig 6. Sex hormones, not sex chromosomes, mediate MRSA colonization resistance in female mice.

(A) MRSA CFU in stool following oral inoculation of male or female mice that were irradiated and reconstituted with bone marrow (BM) from donor male or female mice. Female BM into female recipients (F --> F) n=8, male BM into male recipients (M--> M) n=7, female BM into male recipients (F--> M) n=8. **(B)** MRSA CFU in stool following oral inoculation of ovariectomized female mice or sham operated littermate controls. OVX n=10, sham n=10. **(C)** MRSA CFU in stool following oral inoculation of *Esr1*$^{+/-}$ and *Esr1*$^{-/-}$ female mice bred at NYU. *Esr1*$^{+/-}$ n=6, *Esr1*$^{-/-}$ n=6. **(D)** MRSA CFU in stool following oral inoculation of four core genotype mice. XX n=5, XY(-*Sry*) n=5. XY n=5, XX(+*Sry*) n=4. **(E)** CD4+ T cells from cecal-colonic lamina propria of XX females and XX(+*Sry*) males 2 dpi inoculation with MRSA or mock control. **(F)** Percentage of IL-17A+ CD4+ T cells in cecal-colonic lamina propria of XX females and XX(+*Sry*) males 2dpi inoculation with MRSA or mock control. Data points represent mean ± SEM from at least two independent experiments. Statistical analysis: area under the curve followed by a 1 way ANOVA with Sidak's multiple comparisons for (A),(C) and two-tailed t-test for (B),(F) and 2 way ANOVA + Sidak's multiple comparisons test for (D)-(E). ns: not significant.

DISCUSSION

S. aureus intestinal carriage is common and associated with infection, but how variables
such as sex contribute to colonization susceptibility have been obscure. We established a
mouse model to investigate MRSA intestinal colonization, which revealed a sex-specific effect in
which female mice rapidly cleared MRSA, while their male counterparts remained colonized.
This protection was microbiota dependent because mice lacking a microbiota or those with a
less diverse microbiota were susceptible to persistent colonization. Microbiota composition
alone, however, was insufficient to explain the sex dependent colonization resistance observed.
Females displayed an enhanced immune response to MRSA colonization characterized by
increases in Th17 cells and neutrophil activation. Increase in MRSA burden in ovariectomized or
ERα-deficient females and XX (+*Sry*) gonadal male mice indicate that this effect of the female
sex is hormonally mediated rather than dependent on genes present on sex chromosomes.
Collectively, our results support a model in which GI colonization resistance against MRSA in
female mice is dependent on the microbiota and an enhanced Th17 response downstream of
sex hormones.

Sex steroid hormones have well documented effects on the immune response, including
CD4+ T cells[21–23,68–70]. Our results are consistent with studies showing that ERα signaling
increases differentiation and cytokine production of Th1 and Th17 cells[21,71,72], although higher
levels of estrogen characteristic of pregnancy can stimulate immunosuppressive regulatory
CD4+ T cell conversion[73]. Recent work identified a parallel role for the male sex hormone
androgen in suppressing lymphoid and neutrophil responses during intradermal infection of
mice with *S. aureus*[22,23]. In this context, introduction of a complex microbiota into germ-free mice
amplified the skin type 17 sex bias in females[22]. Sex biases in neutrophil function, including
increased phagocytosis and extracellular trap formation in female mice have been
observed[14,62,74,75]. In line with our findings of increased CD11b on intestinal neutrophils isolated
from female mice exposed to MRSA, studies examining MRSA skin and soft tissue infections
found that bone marrow neutrophils from female mice have an enhanced ability to kill MRSA *ex*
vivo compared to those from male mice, which was linked to increases in surface CD11b and
antimicrobial production[14,75].

Although sex difference was not reported, colonization resistance against *S. aureus* in
the nares is also associated with Th17 cells and neutrophils[51,52], supporting the general
importance of the CD4+ T cell response over antibodies and B cells in determining susceptibility
to colonization. The absence of a sterilizing B cell response may reflect immune evasion

strategies such as the bacterially produced Protein A that binds to the Fcγ portion of

immunoglobulins, protecting *S. aureus* from opsonophagocytic killing[76,77]. Our finding that

female mice lacking mature B cells were still able to resist MRSA GI colonization is mirrored by

a human volunteer study demonstrating that antibodies prior to intranasal inoculation did not

block persistent colonization by their cognate *S. aureus* strains and that antibody levels did not

distinguish intermittent and non-carriers[78]. In contrast, low CD4+ T cell counts in HIV+

individuals are a risk factor for *S. aureus* nasal colonization[79], and rates of colonization were

higher in HIV+ males compared to HIV+ females[80]. Understanding the cellular response to

MRSA colonization and the impact of host sex may inform vaccination strategies[81].

We observed an increase in IL-17A+ CD4+ T cells in colonization resistant females at

two days post inoculation with MRSA, a duration that is typically insufficient for a *de novo*

antigen-specific adaptive immune response. It is likely that the gut harbors pre-existing T cells

that can rapidly respond. The microbiota-dependence of colonization resistance provides a

potential explanation for this observation. Superantigens from *S. aureus* bind directly to Vβ

regions of T cell receptors (TCRs) and major histocompatibility complex (MHC) class II on

antigen-presenting cells, resulting in hyperactivation of T lymphocytes and

monocytes/macrophages. *S. aureus* superantigens induce a robust IL-17 response from

memory Th17 cells from adult humans but not from naive T cells[82], raising the possibility that

such a mechanism can act on Th17 cells selected by the microbiota. Also, CD4+ T cells in the

gut, including Th17 cells, can recognize antigens that are shared by taxonomically diverse

bacteria in the gut[53]. In the colon of healthy humans and mice, MHC-II restricted CD4+ T cells

with Th17 functionality have been identified that are responsive to commensal microbial

antigens in an innate-like manner[83]. Additionally, intestinal colonization by the fungus *Candida*

albicans favors Th17 polarization associated with increased IL-17 and neutrophils that protect

mice from intravenous infection with MRSA[84,85]. Thus, it is possible that we are observing either

a cross reactive Th17 response or a bystander effect initiated by a commensal.

Investigating the antigen specific response to MRSA is challenging due to bacterially-

encoded superantigens and other secreted virulence factors, such as α-hemolysin and LukED,

that induce cell death of lymphocytes[86]. Therefore, the relationship between the microbiota and

sex-dependent colonization resistance remains unanswered in our model. Other possibilities

include microbiota-mediated regulation of estrogen and its receptor or signaling. However, given

the strong connection between microbiota and Th17[33,87], we propose a speculative model in

which initial microbiota differences prime the gut for a Th17 and neutrophil response to MRSA

that is enhanced by estrogen. Identifying the microbe(s) that are necessary for colonization

resistance in females and careful examination of the dynamic regulation of sex hormones in this

model will be insightful.

Animal models that incorporate sex as a variable are critical to conduct rigorous,

translational science and build towards personalized medicine. Our study reveals an interplay

between host microbiota, immune response, and sex steroid hormones in response to intestinal

exposure to MRSA, a common commensal and major source of life-threatening invasive

infections. Given the importance of the gut as a site of colonization and transmission for many

medically important infectious agents, we suggest careful documentation of sex specific effects

in future studies examining this host-microbe interface.

MATERIALS AND METHODS

Mice

Mice designated as JAX refer to male and female 6- to 8-week-old C57BL/6J mice

purchased from Jackson Laboratory and used directly for experiments. NYU C57BL/6J breeders

were originally purchased from Jackson Laboratory and bred onsite at New York University

Grossman School of Medicine to generate littermate male and female mice for comparison.

Every six months breeders were replaced by using a new male from Jackson Laboratory to pair

with a NYU female to reduce genetic drift. *Rag2$^{-/-}$, Tcrd$^{-/-}$, Igh-μ$^{-/-}$, Ptprc^a (B6 CD45.1),* and

Esr1$^{tm1Ksk+/-}$ mice were purchased from Jackson Laboratory. *Rorc(γt)–enhanced GFP* (*Roryt$^{-/-}$*)

mice were previously described[85]. *Il17ra$^{-/-}$* mice were generated by Amgen Inc. (Seattle, WA)

and provided by Dr. Jeffrey Weiser. Littermate heterozygous controls for *Rag2$^{-/-}$*, *Igh-μ$^{-/-}$*, and

Roryt$^{-/-}$ mice were generated by initially crossing homozygous knockout mice with wild-type

C57BL/6J mice to establish heterozygotes that were then used to generate homozygous and

heterozygous breeder pairs.

Germfree (GF) C57BL/6J were bred and maintained in flexible-film isolators at the New

York University Grossman School of Medicine Gnotobiotics Animal Facility[88–90]. Absence of

fecal bacteria was confirmed monthly by evaluating the presence of 16S rDNA in stool samples

by qPCR. Minimal flora mice harboring the consortium of 15 bacteria (Oligo-MM$_{12}$ + FA3)[39] were

maintained in a separate isolator as previously described[88,89]. For inoculation with bacteria, GF

mice and minimal flora mice were housed in Bioexclusion cages (Tecniplast) with access to
sterile food and water.

The sample size for animal experiments was chosen based on previous data generated
in the laboratory. All animal studies were performed according to protocols approved by the
NYU Grossman School of Medicine Institutional Animal Care and Use Committee

MRSA intestinal colonization in mice

Prior to inoculation, stool from C57BL/6J mice was homogenized and plated on
CHROMagar *Staph aureus* (CHROMagar, Paris, France) selective plates to ensure mice were
free of *S. aureus* carriage. C57BL/6J mice were orally gavaged with ~1 × 10^8 colony-forming
units (CFU) of CA-MRSA strain USA300 (LAC). Stool samples were collected from each mouse
on indicated days post inoculation. Screw-cap tubes (2 mL) filled with 1.0 mm beads were
weighed before and after the addition of stool to determine weight. Sterile phosphate-buffered
saline (PBS) (1 mL) was added to each tube, which were vigorously shaken in a bead-beater
(MP Biomedicals, Santa Ana, CA) for 60 s. Stool aliquots were diluted and plated to enumerate
viable bacteria. Samples were plated on CHROMagar MRSA II (BBL) /Mannitol salt agar plates,
incubated at 37ºC for 24 hours, and colonies were counted to determine MRSA burden per
gram of stool.

Intestinal lamina propria and epithelial cell isolation

Colonic and cecal tissues were flushed with HBSS (Gibco), fat and Peyer's patches
were removed, and the tissue was cut into 6-8 pieces. Tissue bits were incubated first with 20
mL of HBSS with 2% HEPES (Corning), 1% sodium pyruvate (Corning), 5mM EDTA, and 1 mM
dithiothreitol (DTT) (Sigma-Aldrich) for 15 min at 37°C with shaking, and then with new 10 mL of
HBSS with 2% HEPES, 1% sodium pyruvate, 5mM EDTA for 10 min at 37°C with shaking. The
samples were filtered by 40 µm cell strainer (BD) and the supernatant containing intestinal
epithelial cells was collected and subjected to gradient centrifugation using 40% and 80%
Percoll (Sigma-Aldrich). The upper layer containing epithelial cells between the 40% and 80%
gradients was collected. Tissue bits were washed in HBSS + 5% FCS, minced, and then
enzymatically digested with collagenase D (0.5 mg/mL, Roche) and DNase I (0.01 mg/mL,
Sigma-Aldrich) for 30 min at 37°C with shaking. Digested solutions were passed through a 40
mm cell strainer and cells were subjected to gradient centrifugation using 40% and 80% Percoll
(Sigma-Aldrich). The lower layer containing immune cells between the 40% and 80% gradients
was collected.

Flow cytometry

Lamina propria cells from either cecal and colonic or small intestinal tissue were

harvested as described. For intracellular cytokine staining, cells were stimulated using the

eBioscience cell stimulation cocktail for 4 h at 37 °C. The cells were fixed and permeabilized

using the Biolegend fixation and permeabilization buffer. The following antibodies (clones) were

used for staining: CD45 (30-F11), Ly6G (1A8), TCR-β (H57-597), CD4 (GK1.5), CD8 (53–6.7),

IL-17A (TC11-18H10.1), TCR γ/δ (GL3), CD127(SB/199), and CD11b (M1/70). All samples were

blocked with Fc Block (TruStain FcX). Zombie UV Fixable Viability Kit (Biolegend) was used to

exclude dead cells prior to gating for other markers. Samples were run on BD FACS Symphony

A5 and FACsFlowJo v.10 was used to analyze the flow cytometry data.

Antibody mediated depletion experiments

C57BL/6J mice bred at NYU were injected intraperitoneally (I.P.) with either 200 μg rat

anti-mouse Ly6G or rat IgG2a isotype control antibody to deplete neutrophils and 250 μg rat

anti-mouse CD4 or rat IgG2a isotype control antibody to deplete CD4+ T cells (Bio-X-Cell, West

Lebanon, NH). Anti-Ly6G injections occurred 1 day prior to MRSA inoculation and then every 3

days until mice were sacrificed 14 days post inoculation. Anti-CD4 injections occured 3 days

prior to inoculation and every 7 days after initial injection until mice were sacrificed 14 days post

inoculation.

Bone marrow chimera experiments

Eight-week-old recipient CD45.1 congenic C57BL/6J mice bred at NYU received 550

rads in 2 doses over 2 sequential days and were injected retro-orbitally with 2×10^6 T

lymphocyte-depleted bone marrow cells from either female or male CD45.2 congenic C57BL/6J

donors bred at NYU. Mature T lymphocytes were depleted from bone marrow cell suspension

using the CD3ε MicroBead Kit (Miltenyi Biotech). Mice were allowed 8 weeks for reconstitution

before oral inoculation with 1×10^8 CFU MRSA. Bone marrow reconstitution of the CD45+

compartment in CD45.1 mice was confirmed by flow cytometry analysis of CD45.2+ cells in the

bone marrow at the time of sacrifice.

Ovariectomy surgery

6-week-old female mice bred at NYU were anesthetized with isoflurane and placed in

ventral recumbency with tail towards surgeon. Ophthalmic ointment was applied bilaterally and

heat support was provided throughout the procedure. The dorsal mid-lumbar area was shaved

in an area 150% greater than anticipated incision length and swabbed three times with

alternating scrubs of betadine and alcohol. A 1-1.5 cm dorsal midline skin incision was made

halfway between the caudal edge of the ribcage and the base of the tail. A single incision of 5-

7mm long was made into the muscle wall on both the right and left sides approximately 1/3 of

the distance between the spinal cord and the ventral midline. The ovary and the oviduct were

exteriorized through the muscle wall. A hemostat was clamped around the uterine vasculature

between the oviduct and uterus. Each ovary and part of the oviduct was removed with single

cuts through the oviducts near the ovary. The hemostat was removed and the remaining oviduct

was assessed for hemorrhage. Hemostasis was confirmed prior to placing the remaining tissue

into the peritoneal cavity. The ovary on the other side was removed in a similar manner. The

muscle incision was closed with monofilament absorbable suture in a cruciate pattern.

Bupivacaine was applied on the closed muscle layer. The skin incision was closed in a cruciate

pattern with sterile skin sutures (monofilament nonabsorbable) and a small amount of tissue

glue. The skin sutures were removed 10 days after surgery. Sham surgery mice underwent the

same surgery but did not have their ovaries removed. Mice were administered Carprofen SQ

every 24 hours for 3 days following the procedure. Mice were allowed two weeks to fully recover

before oral inoculation with $1x10^8$ CFU of MRSA.

RNA-sequencing

RNA was extracted from lamina propria cells and epithelial fraction isolated from the

cecum-colon using RNeasy Plus Mini Kit (Qiagen). Four male and four female B6 mice bred at

NYU were used for each experimental condition. An RNA library was prepared using the

Illumina TruSeq RNA sample preparation kit and sequenced with the Illumina HiSeq 2000 using

the TruSeq RNA v.2 protocol. Illumina CASAVA v.1.8.2 was used to generate FASTQ files

containing 29.5–53.5 million qualified reads per sample. Alignment and gene expression count

were computed using default settings, which aligns reads to the union of all RefSeq-annotated

exons for each gene.

RNA-seq results were processed using the v.4 R package DESeq2 v.3 to obtain

variance stabilized count reads, fold changes relative to specific condition, and statistical P

value. Analysis of the transcriptome focused on differentially expressed genes (DEGs), defined

as the genes with an absolute log2 fold change relative to specific condition >1.2 and an

adjusted P value < 0.05. Enriched pathways for the DEGs were analyzed by Ingenuity Pathway

Analysis (Qiagen). The analyses were visualized using R package ggplot2[91].

DNA extraction from stool

DNA were extracted from stool using the MagMAX™ Microbiome Ultra Nucleic Acid Isolation Kit (Thermo Fischer) following the manufacturer's instructions with modifications. An initial bead-beating step using differentially sized beads (glass beads, 0.5–0.75 mm; zirconia beads, <100 µm) was included and lysozyme (20 mg mL–1) was added to the lysis buffer.

16S library preparation and sequencing analysis

Bacterial 16S rRNA genes were amplified at the V4 region using 16S universal primer pairs and amplicon sequencing was performed on the Illumina MiSeq system, yielding 150-bp paired reads.

16S Amplicon PCR Forward Primer = 5'

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG,

16S Amplicon PCR Reverse Primer = 5'

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC

The sequencing reads were processed using the DADA2 v.1 pipeline in the QIIME2 v.2022.2 software package. The cloud based platform Nephele v.2 was used to run QIIME2 analysis[92]. The minimum Phred quality score of 20 was applied to ensure high-quality sequence data. An open-reference clustering algorithm was used to identify operational taxonomic units (OTUs) based on a 97% similarity threshold to a reference database. Chimera removal was conducted to eliminate artifacts in the data. Taxonomic classification was performed using a search-based method to assign taxonomy to the OTUs. Barplots were generated with a minimum frequency filter of 1000 to visualize the abundance of microbial taxa. A percentage identity threshold of 97% was used to assign taxonomy at the genus level using the SILVA rRNA database. Principal coordinates analysis (PCoA) and calculation of Shannon index were performed using the phyloseq R package[93] version 1.46 after rarefaction to a depth of 1000.

Statistical analysis

The number of animals per group is annotated in corresponding figure legends. GraphPad Prism v.10 was used to generate graphs and assess significance for bacterial burden, weight loss, and flow cytometry data were analyzed using FlowJo v.10. The MRSA burden curves were analyzed using area under the curve followed by a 1 way ANOVA with Sidak's multiple comparisons test when comparing more than 2 experimental conditions and a two-tailed t-test when comparing 2 experimental conditions. An unpaired two-tailed Student's t-test was used to evaluate the differences between two groups. Welch's correction was used

when variances were significantly different between groups. A 2-way ANOVA with Sidak's
multiple comparisons test was used to evaluate experiments involving multiple groups. All P
values are shown in the figures.

DATA AVAILABILITY

The sequencing accession number for the bulk RNA sequencing is PRJNA1134782 and
the accession number for the 16S rRNA sequencing is PRJNA1135964.

ACKNOWLEDGEMENTS

We would like to thank Margie Alva, Juan Carrasquillo and David Basnight for their help
in the NYU Gnotobiotic Facility, the NYU Flow Cytometry Core for training and access to
equipment, the NYU Genome Technology Center for processing and sequencing of 16S and
RNA samples, the NYU Experimental Pathology Research Laboratory for processing of H&E
tissue samples, and the NYU Reagent Preparation service for providing bacterial media and
plates. Core facilities were supported by NIH grant P31CA016087. We would like to thank Dr.
Mariya London for her help with flow cytometry technique and analysis. We would also like to
thank members of the Cadwell, Shopsin, and Torres Labs for their constructive comments. This
work was supported in part by NIH grants DK093668 (KC), AI121244 (KC, VJT), HL123340
(KC), AI130945 (KC), AI140754 (BS, VJT, KC), AI179896 (KC), DK 050306 (KC), and
DK124336 (KC) and NIH grant 2T32AI100853-11 (AL). We would like to acknowledge the
Vilcek Institute of Graduate Biomedical Sciences for their support.

COMPETING INTERESTS

KC has received research support from Pfizer, Takeda, Pacific Biosciences, Genentech, and
Abbvie. KC has consulted for or received an honoraria from Puretech Health, Genentech, and
Abbvie. KC is an inventor on U.S. patent 10,722,600 and provisional patent 62/935,035 and
63/157,225. V.J.T. has received honoraria from Pfizer and MedImmune and is an inventor on
patents and patent applications (US8431,687B2; US2019135900-A1; EP4313303A1) filed by
New York University, which are currently under commercial license to Janssen Biotech Inc.
Janssen Biotech Inc. provides research funding and other payments associated with a licensing
agreement.

REFERENCES

1. David, M. Z. & Daum, R. S. Community-associated methicillin-resistant Staphylococcus aureus: Epidemiology and clinical consequences of an emerging epidemic. *Clinical Microbiology Reviews* vol. 23 616–687 (2010).

2. Seybold, U. *et al.* Emergence of community-associated methicillin-resistant Staphylococcus aureus USA300 genotype as a major cause of health care-associated blood stream infections. *Clin. Infect. Dis.* **42**, 647–656 (2006).

3. Boucher, H. W. & Corey, G. R. Epidemiology of methicillin-resistant Staphylococcus aureus. *Clin. Infect. Dis.* **46**, (2008).

4. de Kraker, M. E. A., Davey, P. G. & Grundmann, H. Mortality and hospital stay associated with resistant Staphylococcus aureus and Escherichia coli bacteremia: Estimating the burden of antibiotic resistance in Europe. *PLoS Med.* **8**, (2011).

5. Acton, D. S., Tempelmans Plat-Sinnige, M. J., Van Wamel, W., De Groot, N. & Van Belkum, A. Intestinal carriage of Staphylococcus aureus: How does its frequency compare with that of nasal carriage and what is its clinical impact? *European Journal of Clinical Microbiology and Infectious Diseases* vol. 28 115–127 (2009).

6. Gagnaire, J. *et al.* Epidemiology and clinical relevance of Staphylococcus aureus intestinal carriage: a systematic review and meta-analysis. *Expert Rev. Anti. Infect. Ther.* **15**, 767–785 (2017).

7. Huang, S. S. *et al.* Decolonization to Reduce Postdischarge Infection Risk among MRSA Carriers. *N. Engl. J. Med.* **380**, 638–650 (2019).

8. von Eiff, C., Becker, K., Machka, K., Stammer, H. & Peters, G. Nasal Carriage as a Source of Staphylococcus aureus Bacteremia . *N. Engl. J. Med.* **344**, 11–16 (2001).

9. Squier, C. *et al.* Staphylococcus aureus rectal carriage and its association with infections in patients in a surgical intensive care unit and a liver transplant unit. *Infect. Control Hosp. Epidemiol.* **23**, 495–501 (2002).

10. Boyce, J. M., Havill, N. L., Otter, J. A. & Adams, N. M. T. Widespread Environmental Contamination Associated With Patients With Diarrhea and Methicillin-Resistant

Staphylococcus aureus Colonization of the Gastrointestinal Tract . *Infect. Control Hosp.*
Epidemiol. **28**, 1142–1147 (2007).

11. Bhalla, A., Aron, D. C. & Donskey, C. J. Staphylococcus aureus intestinal colonization is
associated with increased frequency of S. aureus on skin of hospitalized patients. *BMC*
Infect. Dis. **7**, 105 (2007).

12. Nowak, J. E., Borkowska, B. A. & Pawlowski, B. Z. Sex differences in the risk factors for
Staphylococcus aureus throat carriage. *Am. J. Infect. Control* **45**, 29–33 (2017).

13. Humphreys, H., Fitzpatick, F. & Harvey, B. J. Gender Differences in Rates of Carriage
and Bloodstream Infection Caused by Methicillin-Resistant *Staphylococcus aureus* : Are
They Real, Do They Matter and Why? *Clin. Infect. Dis.* **61**, civ576 (2015).

14. Castleman, M. J. *et al.* Innate Sex Bias of Staphylococcus aureus Skin Infection Is
Driven by α-Hemolysin . *J. Immunol.* **200**, 657–668 (2018).

15. Tacconelli, E. & Foschi, F. Does gender affect the outcome of community-acquired
Staphylococcus aureus bacteraemia? *Clin. Microbiol. Infect.* **23**, 23–25 (2017).

16. Schurz, H. *et al.* The X chromosome and sex-specific effects in infectious disease
susceptibility. *Hum. Genomics* **13**, 2 (2019).

17. Klein, S. L. & Flanagan, K. L. Sex differences in immune responses. *Nat. Rev. Immunol.*
2016 1610 **16**, 626–638 (2016).

18. Jaillon, S., Berthenet, K. & Garlanda, C. Sexual Dimorphism in Innate Immunity. *Clinical*
Reviews in Allergy and Immunology vol. 56 308–321 (2019).

19. Vázquez-Martínez, E. R., García-Gómez, E., Camacho-Arroyo, I. & González-Pedrajo, B.
Sexual dimorphism in bacterial infections. *Biology of Sex Differences* vol. 9 (2018).

20. Dias, S. P., Brouwer, M. C., Van De Beek, D. & Richardson, A. R. Sex and Gender
Differences in Bacterial Infections. *Infect. Immun.* (2022) doi:10.1128/IAI.00283-22.

21. Fuseini, H. *et al.* ERα Signaling Increased IL-17A Production in Th17 Cells by
Upregulating IL-23R Expression, Mitochondrial Respiration, and Proliferation. *Front.*
Immunol. **10**, 2740 (2019).

22. Chi, L. *et al.* Sexual dimorphism in skin immunity is mediated by an androgen-ILC2-
dendritic cell axis. *Science* eadk6200 (2024)

doi:10.1126/SCIENCE.ADK6200/SUPPL_FILE/SCIENCE.ADK6200_MDAR_REPRODU
CIBILITY_CHECKLIST.PDF.

23. Li, F. *et al.* Sex differences orchestrated by androgens at single-cell resolution. *Nat. 2024*
1–8 (2024) doi:10.1038/s41586-024-07291-6.

24. Misawa, Y. *et al.* Staphylococcus aureus Colonization of the Mouse Gastrointestinal Tract
Is Modulated by Wall Teichoic Acid, Capsule, and Surface Proteins. *PLOS Pathog.* **11**,
e1005061 (2015).

25. Kernbauer, E., Maurer, K., Torres, V. J., Shopsin, B. & Cadwell, K. Gastrointestinal
dissemination and transmission of Staphylococcus aureus following bacteremia. *Infect.*
Immun. **83**, 372–378 (2015).

26. Piewngam, P. *et al.* Pathogen elimination by probiotic Bacillus via signaling interference.
Nature **562**, 532 (2018).

27. Flaxman, A. *et al.* Development of persistent gastrointestinal S. aureus carriage in mice.
Sci. Rep. **7**, 1–13 (2017).

28. Gries, D. M., Pultz, N. J. & Donskey, C. J. Growth in cecal mucus facilitates colonization
of the mouse intestinal tract by methicillin-resistant Staphylococcus aureus. *J. Infect. Dis.*
192, 1621–1627 (2005).

29. Jang, K. K. *et al.* Antimicrobial overproduction sustains intestinal inflammation by
inhibiting Enterococcus colonization. *Cell Host Microbe* **31**, 1450-1468.e8 (2023).

30. Ramanan, D., Tang, M. S., Bowcutt, R., Loke, P. & Cadwell, K. Bacterial sensor Nod2
prevents small intestinal inflammation by restricting the expansion of the commensal
Bacteroides vulgatus. *Immunity* **41**, 311 (2014).

31. Cadwell, K. *et al.* Virus-Plus-Susceptibility Gene Interaction Determines Crohn's Disease
Gene Atg16L1 Phenotypes in Intestine. *Cell* **141**, 1135 (2010).

32. Moon, C. *et al.* Vertically transmitted fecal IgA levels distinguish extra-chromosomal
phenotypic variation. *Nature* **521**, 90 (2015).

33. Ivanov, I. I. *et al.* Induction of intestinal Th17 cells by segmented filamentous bacteria.
Cell **139**, 485 (2009).

34. Mamantopoulos, M. *et al.* Nlrp6- and ASC-Dependent Inflammasomes Do Not Shape the

Commensal Gut Microbiota Composition. *Immunity* **47**, 339-348.e4 (2017).

35. Zhou, C. *et al.* Microbiota and metabolic adaptation shape Staphylococcus aureus virulence and antimicrobial resistance during intestinal colonization. *bioRxiv : the preprint server for biology* (2024) doi:10.1101/2024.05.11.593044.

36. Piewngam, P. *et al.* Pathogen elimination by probiotic Bacillus via signalling interference. *Nature* **562**, 532–537 (2018).

37. Krismer, B., Weidenmaier, C., Zipperer, A. & Peschel, A. The commensal lifestyle of Staphylococcus aureus and its interactions with the nasal microbiota. *Nat. Rev. Microbiol.* **15**, 675–687 (2017).

38. T, N. *et al.* Antimicrobials from human skin commensal bacteria protect against Staphylococcus aureus and are deficient in atopic dermatitis. *Sci. Transl. Med.* **9**, (2017).

39. Brugiroux, S. *et al.* Genome-guided design of a defined mouse microbiota that confers colonization resistance against Salmonella enterica serovar Typhimurium. *Nat. Microbiol.* **2**, 16215 (2016).

40. Robertson, S. J. *et al.* Comparison of Co-housing and Littermate Methods for Microbiota Standardization in Mouse Models. *Cell Rep.* **27**, 1910-1919.e2 (2019).

41. Schloss, P. D. *et al.* Stabilization of the murine gut microbiome following weaning. *Gut Microbes* **3**, 383 (2012).

42. Palanza, P., Gioiosa, L. & Parmigiani, S. Social stress in mice: gender differences and effects of estrous cycle and social dominance. *Physiol. Behav.* **73**, 411–420 (2001).

43. Steinert, M., Ramming, I. & Bergmann, S. Impact of Von Willebrand Factor on Bacterial Pathogenesis. *Front. Med.* **7**, 565911 (2020).

44. Schiering, C. *et al.* Feedback control of AHR signalling regulates intestinal immunity. *Nat. 2017 5427640* **542**, 242–245 (2017).

45. Mohan, T., Deng, L. & Wang, B. Z. CCL28 chemokine: An anchoring point bridging innate and adaptive immunity. *Int. Immunopharmacol.* **51**, 165–170 (2017).

46. Huyton, T., Göttmann, W., Bade-Döding, C., Paine, A. & Blaszczyk, R. The T/NK cell co-stimulatory molecule SECTM1 is an IFN 'early response gene' that is negatively regulated by LPS in human monocytic cells. *Biochim. Biophys. Acta* **1810**, 1294–1301

(2011).

47. Donato, R. RAGE: a single receptor for several ligands and different cellular responses:
the case of certain S100 proteins. *Curr. Mol. Med.* **7**, 711–724 (2007).

48. Cao, S. *et al.* The Gut Epithelial Receptor LRRC19 Promotes the Recruitment of Immune
Cells and Gut Inflammation. *Cell Rep.* **14**, 695 (2016).

49. Anderson, C. K. & Brossay, L. The role of MHC class Ib-restricted T cells during infection.
Immunogenetics **68**, 677 (2016).

50. Brustein, M., Kraal, G., Mebius, R. E. & Watson, S. R. Identification of a soluble form of a
ligand for the lymphocyte homing receptor. *J. Exp. Med.* **176**, 1415–1419 (1992).

51. Archer, N. K., Harro, J. M. & Shirtliff, M. E. Clearance of Staphylococcus aureus Nasal
Carriage Is T Cell Dependent and Mediated through Interleukin-17A Expression and
Neutrophil Influx. *Infect. Immun.* **81**, 2070 (2013).

52. Archer, N. K. *et al.* Interleukin-17A (IL-17A) and IL-17F Are Critical for Antimicrobial
Peptide Production and Clearance of Staphylococcus aureus Nasal Colonization. *Infect.*
Immun. **84**, 3575 (2016).

53. Nagashima, K. *et al.* Mapping the T cell repertoire to a complex gut bacterial community.
Nat. 2023 6217977 **621**, 162–170 (2023).

54. Bacher, P. *et al.* Human Anti-fungal Th17 Immunity and Pathology Rely on Cross-
Reactivity against Candida albicans. *Cell* **176**, 1340-1355.e15 (2019).

55. Cassotta, A. *et al.* Broadly reactive human CD4+ T cells against Enterobacteriaceae are
found in the naïve repertoire and are clonally expanded in the memory repertoire. *Eur. J.*
Immunol. **51**, 648–661 (2021).

56. McGeachy, M. J., Cua, D. J. & Gaffen, S. L. The IL-17 family of cytokines in health and
disease. *Immunity* **50**, 892 (2019).

57. Blazkova, J. *et al.* Multicenter Systems Analysis of Human Blood Reveals Immature
Neutrophils in Males and During Pregnancy. *J. Immunol.* **198**, 2479–2488 (2017).

58. Er-Lukowiak, M. *et al.* Testosterone affects type I/type II interferon response of
neutrophils during hepatic amebiasis. *Front. Immunol.* **14**, 1279245 (2023).

59. Gupta, S. *et al.* Sex differences in neutrophil biology modulate response to type I

interferons and immunometabolism. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 16481–16491 (2020).

60. Lu, R. J. *et al.* Multi-omic profiling of primary mouse neutrophils predicts a pattern of sex- and age-related functional regulation. *Nat. Aging 2021 18* **1**, 715–733 (2021).

61. Parkos, C. A., Colgan, S. P. & Madara, J. L. Interactions of neutrophils with epithelial cells. *J. Am. Soc. Nephrol.* **5**, 138–152 (1994).

62. Yasuda, H. *et al.* 17-β-estradiol enhances neutrophil extracellular trap formation by interaction with estrogen membrane receptor. *Arch. Biochem. Biophys.* **663**, 64–70 (2019).

63. Békési, G. *et al.* Induced myeloperoxidase activity and related superoxide inhibition during hormone replacement therapy. *BJOG* **108**, 474–481 (2001).

64. Newcomb, D. C. *et al.* Estrogen and progesterone decrease let-7f microRNA expression and increase IL-23/IL-23 receptor signaling and IL-17A production in patients with severe asthma. *J. Allergy Clin. Immunol.* **136**, 1025-1034.e11 (2015).

65. Arnold, A. P. & Chen, X. What does the "four core genotypes" mouse model tell us about sex differences in the brain and other tissues? *Front. Neuroendocrinol.* **30**, 1 (2009).

66. Palaszynski, K. M. *et al.* A yin-yang effect between sex chromosome complement and sex hormones on the immune response. *Endocrinology* **146**, 3280–3285 (2005).

67. Sasidhar, M. V., Itoh, N., Gold, S. M., Lawson, G. W. & Voskuhl, R. R. The XX Sex Chromosome Complement in Mice is Associated with Increased Spontaneous Lupus as compared to XY. *Ann. Rheum. Dis.* **71**, 1418 (2012).

68. Taneja, V. Sex Hormones Determine Immune Response. *Front. Immunol.* **9**, 1931 (2018).

69. Karpuzoglu-Sahin, E., Hissong, B. D. & Ansar Ahmed, S. Interferon-γ levels are upregulated by 17-β-estradiol and diethylstilbestrol. *J. Reprod. Immunol.* **52**, 113–127 (2001).

70. Karpuzoglu-Sahin, E., Zhi-Jun, Y., Lengi, A., Sriranganathan, N. & Ansar Ahmed, S. Effects of long-term estrogen treatment on IFN-γ, IL-2 and IL-4 gene expression and protein synthesis in spleen and thymus of normal C57BL/6 mice. *Cytokine* **14**, 208–217 (2001).

71.　Mohammad, I. *et al.* Estrogen receptor α contributes to T cell-mediated autoimmune
inflammation by promoting T cell activation and proliferation. *Sci. Signal.* **11**, (2018).

72.　Maret, A. *et al.* Estradiol enhances primary antigen-specific CD4 T cell responses and
Th1 development in vivo. Essential role of estrogen receptor alpha expression in
hematopoietic cells. *Eur. J. Immunol.* **33**, 512–521 (2003).

73.　Tai, P. *et al.* Induction of regulatory T cells by physiological level estrogen. *J. Cell.*
Physiol. **214**, 456–464 (2008).

74.　Spitzer, J. A. & Zhang, P. Gender differences in neutrophil function and cytokine-induced
neutrophil chemoattractant generation in endotoxic rats. *Inflammation* **20**, 485–498
(1996).

75.　S, P. *et al.* Complement Receptor 3 Contributes to the Sexual Dimorphism in Neutrophil
Killing of Staphylococcus aureus. *J. Immunol.* **205**, 1593–1600 (2020).

76.　Falugi, F., Kim, H. K., Missiakas, D. M. & Schneewind, O. Role of Protein A in the
Evasion of Host Adaptive Immune Responses by Staphylococcus aureus. *MBio* **4**,
(2013).

77.　Pauli, N. T. *et al.* Staphylococcus aureus infection induces protein A–mediated immune
evasion in humans. *J. Exp. Med.* **211**, 2331 (2014).

78.　Van Belkum, A. *et al.* Reclassification of Staphylococcus aureus Nasal Carriage Types. *J.*
Infect. Dis. **199**, 1820–1826 (2009).

79.　De Carvalho Ferreira, D. *et al.* Methicillin-resistant Staphylococcus aureus in HIV
patients: Risk factors associated with colonization and/or infection and methods for
characterization of isolates – a systematic review. *Clinics* **69**, 770 (2014).

80.　Neupane, K. *et al.* Comparison of Nasal Colonization of Methicillin-Resistant
Staphylococcus aureus in HIV-Infected and Non-HIV Patients Attending the National
Public Health Laboratory of Central Nepal. *Can. J. Infect. Dis. Med. Microbiol.* **2018**,
(2018).

81.　Brown, A. F., Leech, J. M., Rogers, T. R. & McLoughlin, R. M. Staphylococcus aureus
Colonization: Modulation of Host Immune Response and Impact on Human Vaccine
Design. *Front. Immunol.* **4**, (2013).

82.   Islander, U. *et al.* Superantigenic Staphylococcus aureus stimulates production of
interleukin-17 from memory but not naive T cells. *Infect. Immun.* **78**, 381–386 (2010).

83.   Hackstein, C. P. *et al.* A conserved population of MHC II-restricted, innate-like,
commensal-reactive T cells in the gut of humans and mice. *Nat. Commun.* **13**, (2022).

84.   Shao, T. Y. *et al.* Commensal Candida albicans positively calibrates systemic Th17
immunological responses. *Cell Host Microbe* **25**, 404 (2019).

85.   Chen, Y. H. *et al.* Rewilding of laboratory mice enhances granulopoiesis and immunity
through intestinal fungal colonization. *Sci. Immunol.* **8**, (2023).

86.   Tam, K. & Torres, V. J. Staphylococcus aureus Secreted Toxins and Extracellular
Enzymes. *Microbiol. Spectr.* **7**, (2019).

87.   Atarashi, K. *et al.* Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial
Cells. *Cell* **163**, 367–380 (2015).

88.   Sargsian, S. *et al.* Clostridia isolated from helminth-colonized humans promote the life
cycle of Trichuris species. *Cell Rep.* **41**, 111725 (2022).

89.   Dallari, S. *et al.* Enteric viruses evoke broad host immune responses resembling those
elicited by the bacterial microbiome. *Cell Host Microbe* **29**, 1014 (2021).

90.   Kernbauer, E., Ding, Y. & Cadwell, K. An enteric virus can replace the beneficial function
of commensal bacteria. *Nat. 2014 5167529* **516**, 94–98 (2014).

91.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag, 2016).

92.   Weber, N. *et al.* Nephele: a cloud platform for simplified, standardized and reproducible
microbiome data analysis. *Bioinformatics* **34**, 1411–1413 (2018).

93.   McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, (2013).

882