1

Prebiotic Supplementation Modulates the Gut Microbiome for Improving Oral Antipsychotic Bioavailability

5   *Kate Collins, [a] Srinivas Kamath,[a] Tahlia R. Meola,[a] Anthony Wignall,[a] and Paul Joyce[a] **

6   [a] UniSA Clinical & Health Sciences, University of South Australia, Adelaide, South Australia 5000, Australia

*Corresponding Author. Phone: +61 8 8302 1429; Email: paul.joyce@unisa.edu.au'; ORCID:

0000-0003-3619-7901

Abstract:

This study investigates the impact of gut microbiome composition on the bioavailability and

biodistribution of lurasidone hydrochloride, an atypical antipsychotic used in treating mental

health conditions. The research aims to uncover the relationship between gut metabolome,

specifically short-chain fatty acids (SCFAs), and lurasidone absorption through modulation of

intestinal pH. Male Sprague-Dawley rats underwent a 14-day microbiome depletion using

antibiotics or prebiotics administered via drinking water, resulting in depleted, enriched, or

normal (control) microbiomes. Following this treatment, lurasidone was orally administered to

fasted rats, and blood samples were collected at specific intervals to correlate pharmacokinetic

analysis with changes in microbiota and metabolome composition. Results revealed a

significant 4.3-fold increase in lurasidone bioavailability following prebiotic administration,

while antibiotic treatment decreased bioavailability. This effect is attributed to prebiotics

increasing microbial diversity, which strongly correlates with SCFA production ($R^2 = 0.93$).

The subsequent lowering of intestinal pH created a more suitable environment for lurasidone

solubility and absorption. This novel study establishes a clear relationship between gut

microbiome composition and lurasidone bioavailability, with prebiotic administration

positively impacting both gut microbial diversity and drug absorption. Thus, these findings

suggest that gut microbiome manipulation may present an innovative approach to improving

therapeutic outcomes for antipsychotic medications, potentially addressing poor response rates

in some patients. Furthermore, this research highlights the importance of considering gut

microbiome composition, in particular the use of SCFAs as a biomarker to produce predictive

models of drug pharmacokinetics, especially for pH-dependent soluble drugs. In the context of

drug efficacy these gut microbiome-based predictive pharmacokinetic models will pave the

path for personalised medicine approaches for mental health treatment.

Keywords:

Gut microbiome; Lurasidone; Bioavailability; Solubility; Antipsychotics; Short-chain fatty

acids

Graphical Abstract:

1. Introduction:

Atypical antipsychotics pharmacotherapeutics are utilised for a broad spectrum of mental health conditions (such as major depressive disorder, anxiety, schizophrenia, and bipolar disorder) [1–3]. However, these medications are often prescribed without considering the potential negative effects, such as metabolic dysfunction and worsening mental health. Antipsychotics interact with the gut microbiome in a bidirectional way, which is frequently overlooked [1,4–6]. This interaction can lead to adverse effects, including metabolic disruptions like weight gain [4]. The bidirectional relationship involves two aspects: drugs and their excipients can disrupt the gut microbiome leading to adverse metabolic effects (*e.g.*, weight gain in the case of antipsychotics), and alternatively the gut microbiome can alter the pharmacokinetics of orally administered drugs (*e.g.*, absorption, biodistribution, metabolism, and excretion) [1,3,7,8].

A noteworthy example of a drug-gut interaction occurs with lurasidone, a benzothiazole derivative atypical antipsychotic that possesses pH-dependent solubility. This impact was previously explored by Cussotto et al. on similar atypical antipsychotics, olanzapine and risperidone [6]. Post microbiome depletion on adult male Sprague-Dawley rats (n: 6-7/group, weight: 0.200-0.250 kg, dosed: ampicillin 1g/L, vancomycin 500mg/L, and imipenem 250mg/L for 14 days in drinking water), olanzapine bioavailability increased 82%, whilst risperidone did not show any significant changes [6]. Cussotto et al. suggest pH changes and enzymatic degradation to be causative culprits however, they admit the discrepancy demonstrates that the impacts of the gut microbiome are not uniform [6]. Such mechanisms rationalise the real-world interindividual pharmacokinetic variation observed when taking antipsychotics and are corroborated by Minichino et al. [3].

Furthermore, short-chain fatty acids (SCFAs; acetate, propionate, and butyrate) are produced by microbial metabolism of indigestible dietary fibres which, as explored by Walker et al.,

influence intestinal pH [9]. This phenomenon in turn drives intestinal environments toward or

away from the pKa of administered drugs [10]. Enrichment of the gut microbiome enhances

SCFAS production, as several studies by Venkataraman et al., Deehan et al., McOrist et al. and

Gibson et al. [11–14]. However, the implications of this for individual drugs are poorly understood.

This novel study aims to address this critical gap in our understanding of atypical antipsychotic

pharmacokinetics by investigating the bidirectional relationship between microbiota and

lurasidone. Unravelling these complex interactions and mechanisms underlying the poor

solubilisation and bioavailability of lurasidone, may pave the way for more personalised and

effective treatment strategies, potentially mitigating adverse effects and improving therapeutic

outcomes for patients with mental health disorders. Furthermore, insights gained from this

research could have far-reaching implications for drug development and administration across

various classes of psychotropic medications, ultimately contributing to more informed and

nuanced approaches in psychopharmacology.

2. Materials and methods

2.1 Animal Ethics & Husbandry

The *in vivo* study received approval from the Animal Ethics Committee at the University of

South Australia (approval #U24-21), adhering to the NIH's Principles of Laboratory Animal

Care (NIH publication #85-23), revised in 1985) and the Australian code for the care and use

of animals for scientific purposes (8th edition 2013, revised 2021), and was conducted and

reported following ARRIVE guidelines. The study was conducted on 8-week-old male

Sprague-Dawley rats sourced from Ozgene (Canning Vale, Australia) with a mean starting

weight of 0.270-0.330 kg. Sprague-Dawley rats were utilised in this study because 1) their gut

microbiome closely mimics the human gut microbiome [15,16], 2) they are widely utilised in

pharmacomicrobiomics studies [17,18] and have been used previously to investigate the impact of

olanzapine on the gut microbiome [4] and 3) their docile nature allows for ease-of-handling for

daily dosing via oral gavage [19]. Rats were housed in groups of 2 in a temperature-, humidity-,

pressure-controlled animal facility with a 12h/12h light/dark cycle with access to standard

rodent chow and water. Only healthy rats were included in the study, and those showing signs

of illness or abnormal behaviour during the acclimation period were excluded. Randomisation

and blinding were implemented during the allocation of animals to treatment groups and

throughout the data collection process. The researchers administering treatments, as well as

those assessing outcomes, were blinded to the group allocations to prevent bias.

2.2 In vivo study design

The oral bioavailability of lurasidone (Hangzhou Dayangchem Co. Ltd., Hangzhou City,

China) was investigated in three randomly assigned treatment groups ($n = 5$). The sample size

for each treatment group ($n = 5$) was based on Power calculations for anticipated changes in

the lurasidone area under the curve, using a power level of ≥ 0.8 and a significance level of

0.05. A single-gender was used for the pharmacokinetics study to minimise gender-based

variation commonly observed in oral pharmacokinetics investigations. Treatment groups were

as follows: 1) control group where rats were provided *ad libitum* access to standard drinking

water; 2) enriched microbiome group where rats were provided *ad libitum* access to drinking

water with inulin (100 mg/L) dissolved; and, 3) depleted microbiome group where rats were

provided *ad libitum* access to drinking water with vancomycin (500 mg/L), ampicillin (1 g/L),

and imipenem (250 mg/L) dissolved. Following 14 days of pre-treatment with prebiotics (*i.e.,*

inulin, inulin from dahlia tubers (with a degree of polymerisation of 35) were purchased from

Merck [Bayswater, Australia]) or antibiotics (ampicillin and vancomycin were purchased from

Merck [Bayswater, Australia]) to manipulate the gut microbiome, each treatment group was

administered a single dose of lurasidone (7.5 mg/mL in PBS) via oral gavage. Blood samples

were collected over various time points, (0, 0.33, 0.66, 1, 1.5, 2, 4, 6, 8, 24h) for

pharmacokinetic analysis. Caecal samples were collected for gut microbiome analysis. After

the final samples were collected, rats were anesthetised using 5% isoflurane, followed by

cardiac puncture and cervical dislocation.

2.3 Gut microbiota analysis

Following treatment and a single dose of lurasidone animal caecal contents were collected and

stored in a -80°C freezer for future analysis. Samples were sent to the Australian Genomics

Research Facility (Brisbane, Australia) for DNA extraction, PCR amplification, and 16S rRNA

sequencing. Using the Quantitative Insights to Microbiology Ecology (QIIME 1.8and the

Silvia reference database the 16S rRNA sequences were processed for the V3-V4 hypervariable

regions, with raw reads clustered into operational taxonomic units (OTUs). OTUs were

assigned taxonomy using the QIAGEN Microbial Insights – Prokaryotic Taxonomy Database

(QMI-PTDB) within the QIAGEN CLC Genomics Workbench version 23.0.4 (Hilden,

Germany). Alpha diversity was measured via Shannon's Index at the genus level, and beta

diversity analyses were conducted using Principal Coordinate Analyses (PCoAs) based on

Bray-Curtis' dissimilarity metrics. Statistical significance was evaluated through Permutational

Multivariate ANOVA (PERMANOVA) on the beta diversity plots.

2.4 Short-chain fatty acid analysis

SCFA concentration in caecal contents was analysed using a modified gas chromatography-

mass spectrometry (GCMS) protocol (Moreau et al, 2003), modified for the GC–MS (7890A

GC System and 5975C inert XI EI/CI MSD with an EI inert 350 source) and MassHunter

Workstation Software (MassHunter, Agilent Technologies). The protocol details have been

previously published [9].

2.5 Intestinal pH analysis

Rat intestinal media was analysed to investigate the effect of the gut microbiome on intestinal

pH. A Hanna instruments Bluetooth pH electrode (HI10832) was used to measure intestinal

pH. The electrode was placed in the intestinal media and the measurement was recorded.

2.6 Lurasidone drug solubility

Lurasidone drug solubilisation was analysed by adjusting the pH of the simulated intestinal

buffer solution and adding lurasidone in excess. pH levels 6.25, 6.75, and 7.5 were of interest

due to being common within rat intestinal media. The different pH levels were made using

intestinal buffer and hydrochloric acid to achieve the pH level desired. The Hanna instruments

Bluetooth pH electrode (HI10832) was used to verify the pH level. Lurasidone was added to

the pH buffers in excess, above its solubility, with 1 mg of lurasidone added to 1 mL of each

different pH solution. Investigations were performed in triplicate. Lurasidone and pH buffer

solution were left on a moving plate at 37 degrees for 2 days covered in aluminium. Samples

were centrifuged at max speed for 20 minutes to allow lurasidone precipitate to be separated.

The supernatant was collected and analysed using liquid chromatography-mass spectrometry

(LCMS). The protocol details have been previously published [10].

2.7 Pharmacokinetic analysis

Blood samples were collected from the saphenous vein and centrifuged to collect the plasma.

For LCMS analysis the plasma was diluted with acetonitrile (5:9 ratio) and spiked using an

internal standard (ziprasidone; 10μl). Pharmacokinetic analyses were performed using

Phoenix® WinNonlin® Version 8.3 (Pharsight®, a Certara™ company), using non-

compartmental methods (Model 200–202, Extravascular Input) and a linear-trapezoidal

approach.

 2.8 Statistical Analysis

GraphPad Prism Version 8.0 (GraphPad Software Inc., California) was used for all statistical

analyses (excluding 16S sequencing analyses) of experimental data. An unpaired t-test and one-

way ANOVA were used to determine statistically significant differences, followed by Tukey's

post-test for multiple comparisons. Values are expressed as the mean ± standard deviation (SD),

with statistical significance set at $p < 0.05$. In the figures, statistical significance is denoted by

* $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$, and **** $p < 0.00005$.

3. Results

3.1 Gut microbial enrichment via prebiotic supplementation correlates with metabolome

enrichment through increased SCFA concentrations in the caecum.

An overview of the dosing regimen and study design to investigate the impact of the gut

microbiome on lurasidone bioavailability is presented in **Fig. 1A.** The 14-day pre-treatment

approach, whereby treatment groups were administered 1) no pre-treatment (*i.e.*, control), 2)

antibiotics, or 3) prebiotics *ad libitum* in drinking water. Analysis of microbial abundance at

the family taxonomic level from the caecal contents from each animal, illustrated in **Fig. 1B**,

with grouped data highlighted in **Fig. 1C**, indicates that the pre-treatment led to treatment

groups with a 1) control microbiome, 2) depleted microbiome, and 3) enriched microbiome.

Microbial abundance increased following prebiotic pre-treatment, in particular members of the

Lachnospiraceae and *Bacteroidacae* families. In contrast, Enterobacteriaceae show

predominance in the antibiotic-treated group. Observed changes in total operation taxonomic

units (OTUs) and alpha diversity (measured by Shannon's index) are shown in Fig. 1D and Fig.

1E for the three groups. The prebiotic treatment group exhibits an enriched microbiome via an

increase in OTUs (+50 OTUs) and Shannon's index (+1 increase in Shannon's index) compared

to the control group. Contrary to this, the antibiotic treatment group displays a significant

decrease in OTUs (-50 OTUs) and Shannon's index (-1 decrease), when compared to the

control group. PERMANOVA of principal component analysis (PCoA) reveals statistically

significant differences in beta diversity between the three treatment groups (Fig 1F), further

highlighting that three distinct microbial populations were present for the respective treatment

groups.

Figure 1. *Microbial pretreatment resulted in three distinct microbiome compositions between*

the treatment groups. [A] A schematic representation of the treatment protocols for each

group. Relative abundance of gut microbiota families post-treatment for individual animals

(Panel [B]) and grouped averages (Panel [C]). Changes in OTUs (Panel [D]) and Shannon's

Index (Panel [E]) reveal statistically significant changes in microbial diversity across the three

treatment groups. Principal component analysis demonstrates statistically significant shifts in

microbiota composition across the three treatment groups (Panel [F]). Statistical significance

 was assessed using unpaired t-test and one-way ANOVA for α-diversity and PERMANOVA for

β-diversity. The asterisk (****) denotes statistically significant results (p < 0.0001).

The concentration of key SCFAs was quantified within caecal contents. A significant difference

($p = 0.0004$) in acetic acid concentration was observed between the enriched microbiome and

depleted microbiome groups (**Fig. 2A**). Similarly, propionic acid concentrations are reduced (p

$= 0.0296$) in microbiome-depleted groups compared to the control group (**Fig. 2B**). No

significant difference in butyric acid concentration is noted between control and enriched

microbiome groups, with a similar concentration range (10-20 μmol g^{-1}), however, depleted

microbiome group showcased a significant decrease (5 μmol g^{-1}), as depicted in **Fig. 2C**. Total

SCFA concentrations vary between all groups as depicted in **Fig. 2D**. The enriched microbiome

demonstrates the highest concentration (200-300μmol g^{-1}), while the depleted microbiome has

the lowest total SCFA concentration (< 50 μmol g^{-1}). A strong correlation between total SCFA

(μmol g^{-1}) and Shannon's index ($R^2 = 0.93$) is displayed in **Fig. 2E**, suggesting microbial

diversity increases total SCFA concentration. Similarly, a correlation ($R^2 = 0.83$) can be noted

between total SCFA (μmol g^{-1}) and OTUs (**Fig. 2F**).

Figure 2. *Enriched microbiota diversity correlates directly with an increase in SCFA*

concentrations in rats. [A] Acetic acid concentration across control, depleted, and enriched

microbiome groups. [B] Propionic acid concentration across the three groups. [C] Butyric

acid concentration across the three groups. [D] Total SCFA concentration across the three

groups, demonstrating statistical significance across all three groups investigated. Total SCFA

concentration directly correlates with alpha diversity, as demonstrated through strong

correlations with Shannon's index, showing strong correlation ($R^2 = 0.93$) (Panel [E]) and

OTUs ($R^2 = 0.83$) (Panel [F]). Values are presented as the mean ± standard deviation (SD).

Statistical significance was assessed using an unpaired t-test and one-way ANOVA. The

asterisk (*), (**), (***) or (****) denote statistically significant results ($p < 0.05$), ($p < 0.01$),

($p < 0.001$) or ($p < 0.0001$) respectively.

 3.2 Increased lurasidone plasma concentration in enriched microbiome

The pharmacokinetic profiles for each group after a single oral dose of lurasidone (20 mg/kg)

to male Sprague Dawley rats are displayed in **Fig. 3A** with interindividual pharmacokinetic

variability highlighted in **Fig. 3B-D.** Noteworthy is the outlier present in the depleted

microbiome group (**Fig. 3C**) which showcases uncharacteristically prolonged lurasidone

absorption/ delayed clearance. When excluded from consideration, the mean lurasidone

bioavailability is diminished. The enriched microbiome group demonstrates a 4.3-fold increase

in lurasidone bioavailability over the 24 hours in comparison to the control, based on mean

area-under-the-curve (AUC_{0-24}) values, presented in **Fig. 3E**. Further, C_{max} values increased by

3.3-fold and 3.9-fold for the enriched microbiome group compared to the control and depleted

microbiome group, respectively (**Fig. 3F**). The degree of drug absorption between the control

and depleted microbiome groups were statistically equivalent, but significant differences were

observed in T_{max} values. The pharmacokinetic results are summarised in Table 1.

Figure 3. *Microbiome enrichment via prebiotic supplementation increases lurasidone*

bioavailability. [A] Pharmacokinetic profiles for the oral administration of lurasidone (20

mg/kg) across the three groups (n = 5) revealed an increase in oral bioavailability for the

*enriched microbiome group. Panels [**B-D**] present the pharmacokinetic analysis of each rat*

*within the three groups. [**E**] Showcasing area-under-the-curve across the groups with*

statistical significance observed between enriched microbiome and depleted microbiome

*groups, and enriched microbiome and control groups respectively. [**F**] Maximum lurasidone*

plasma concentration observed across the three groups trialled. Values are presented as the

mean ± standard deviation (SD). Statistical significance was assessed using an unpaired t-test

and one-way ANOVA. The asterisk (**) or (****) denote statistically significant results (p<

0.01) or (p < 0.0001) respectively.

Table 1: Pharmacokinetic parameters following oral administration of lurasidone (20 mg/kg)

to male Sprague Dawley rats in the fasted state following a 14-day pre-treatment with

antibiotics and prebiotics to deplete and enrich the gut microbiome, respectively. Values

represent the mean ± SD, $n = 5$.

Pharmacokinetic parameter	Control	Depleted microbiome	Enriched microbiome
T_{max} (h)	1.40 ± 0.548	0.39 ± 0.15	0.736 ± 0.148
C_{max} (ng/mL)	36.2 ± 18.6	30.3 ± 6.30	119 ± 24.6
AUC_{0-24} (ng.h/mL)	154 ± 77.3	144 ± 241	669 ± 207
$T_{1/2}$ (h)	3.07 ± 2.13	0.863 ± 0.20	Not determined

        3.3 SCFAs lower intestinal pH and promote lurasidone solubility

The impact of microbiome compositional changes on luminal pH was investigated, with **Fig.**

4A. demonstrating significant alteration to intestinal pH accompanied by gut microbiome

changes. An enriched microbiome results in a lower pH (6.0-6.5), and a depleted microbiome

shows a higher pH (7.0-8.0), compared to the control group exhibiting a pH range of 6.5-7.0.

Strong negative correlations existed between intestinal pH and Total SCFA (**Fig. 4B**) and

Shannon's Index (**Fig. 4C**), demonstrating that the intestinal pH decreases as a function of

increasing SCFA concentrations. Lower pH levels in the enriched microbiota groups also

demonstrated 2-fold increases in lurasidone solubility compared to control groups (**Fig. 4D**).

Fig. 4E and **4F** demonstrate correlations between area-under-the-curve and total SCFA (R^2 of

0.72) and Shannon's index (R^2 of 0.71) respectively.

Figure 4. *An enriched microbiome demonstrates increased lurasidone solubility due to a*

decreased intestinal pH. *[A] Intestinal pH levels across control, depleted microbiome, and*

enriched microbiome. The enriched microbiome displays the lowest pH level and is statistically

significant. [B] Illustrating the relationship between total SCFAs and intestinal pH there is a

strong correlation (R^2 = 0.82). [C] Exhibiting Shannon's index against intestinal pH, with a

strong correlation (R^2 = 0.87). [D] Lurasidone solubility across the three groups. An enriched

microbiome demonstrates the highest lurasidone solubility and statistical significance when

compared with the depleted microbiome. [E] and [F] Highlighting area-under-the-curve

against total SCFA and Shannon's index. Both show strong correlation ($R^2 = 0.71$ and 0.72)

Values are presented as the mean ± standard deviation (SD). Statistical significance was

assessed using an unpaired t-test and one-way ANOVA. The asterisk (**) or (****) denote

statistically significant results (p< 0.01) or (p < 0.0001) respectively.

4. Discussion

Pharmacomicrobiomics explores the bidirectional relationships between drugs and the gut

microbiome and is of particular importance when prescribing antipsychotic medication [1,20–22].

Whilst the impact of psychotropics on the gut microbiome is gaining traction with

comprehensive reviews from Kamath et al. and Minichino et al., the reciprocal relationships

are less explored [1,3].

To investigate the pharmacomicrobiomics of lurasidone, an *in vivo* animal model of Sprague-

Dawley rats was utilised with an antibiotic, control and prebiotic group (as per **Fig. 1A**) with

doses interpreted from prior work by Huang et al. and Davey et al. [4,23]. Antibiotic

supplementation decreased several commensal bacterial families such as *Lachnospiraceae* and

Bacteroidacae both of which are also vital for SCFA production as per prior work by Fusco et

al. [24]. *Lachnospiraceae, in* particular, is suggested by Vacca et al. to possess several members

(such as *Roseburia* and *Eubacterium rectale*) which ferment starch and simple sugars into

butyrate (via the butyryl-CoA pathway), formate and lactate [25]. Similarly, *Bacteroidacae*

harbour polysaccharide utilisation loci responsible for saccharification of complex

carbohydrates, which result in SCFA production as suggested by Zhang et al. [26]. Enrichment of

the gut microbiome increased both *Lachnospiraceae* and *Bacteroidacae* relative abundance

alongside holistic improvements in OTUs (**Fig. 1D**) and increases in β-diversity (**Fig. 1F**) both

of which align with previous studies by Carlson et al. and Tian et al. [27,28]. Such increases in the

relative abundance of the aforementioned SCFA-producing bacteria and overall OTUs led to

increases in SCFA concentrations (**Fig. 2D**) with strong correlations being noted between SCFA

concentration and Shannon's index and OTUs (**Fig. 2E**: R^2 0.93, **Fig. 2F**: R^2 0.83). Such results

are echoed by Gurry et al. with similar correlations between functional heterogeneity in gut

microbiome and SCFA production [29].

The increases in SCFA bear importance in exploring pharmacomicrobiomics through their

influence on luminal pH. SCFAs are acidic (propionate: pH 2.4, butyrate: pH 5.5, acetate: pH

4.6), which Liu et al. suggest can decrease luminal pH through the release of H^+ ions [30].

Specifically, the acidity of SCFAs arises from the presence of a carboxyl group (-COOH),

which can donate a proton (H^+) to the solution, forming a carboxylate anion ($R\text{-}COO^-$)

stabilised by resonance as suggested by Parada Venegas et al. [31]. Acetic acid (CH_3COOH)

releases a proton to form acetate (CH_3COO^-), propionic acid (CH_3CH_2COOH) forms

propionate ($CH_3CH_2COO^-$), and butyric acid ($CH_3CH_2CH_2COOH$) forms butyrate

($CH_3CH_2CH_2COO^-$). The inductive effect of the alkyl chains (methyl, ethyl and propyl) also

slightly influences their acidity, with longer chains being less acidic due to electron-donating

effects. Sun et al. explored the physiological role of SCFAs, suggesting this luminal pH

reduction to be integral in inhibiting pathogenic invasion and colonisation [32]. However, in the

context of pharmacomicrobiomics, the acidic environment can drive drugs toward or further

from their pKa, influencing drug absorption, metabolism, and efficacy as noted by Panebianco

et al. [20].

Lurasidone, (chemical formula: $C_{28}H_{36}N_4O_2S$) contains a benzothiazole derivative with a

piperazine ring (providing basicity and contributing to pKa), a sulphonamide group (neutral

but capable of influencing hydrogen bonding capacity), and multiple aromatic rings (impacting

lipophilicity and drug-receptor interactions) which result in a pKa of 7.6 [33]. Below the pKa, the

piperazine nitrogen is protonated, resulting in a positively charged species (Lurasidone-NH_3^+)

which possesses greater aqueous solubility and hydrophilicity, through the formation of ion-

dipole interactions between the positively charged amine and water molecules [34,35]. This results

in the maximum aqueous solubility of 0.349 mg/mL in pH 3.5 buffer [33].

Whilst the specifics of lurasidone absorption and active transporter receptors is poorly

understood (possibly due to its recency in development), increased solubility is generally

accepted as being favourable towards increasing absorption and bioavailability as covered in a

comprehensive review by Kumari et al. [36]. Alternative atypical antipsychotics also possess

similar pH-dependent solubility characteristics, such as olanzapine (protonation of amine

group) and risperidone (protonation of the piperidine nitrogen) [37]. However, unlike alternatives,

lurasidone is hampered by poor bioavailability (9-19%) with a 2-fold positive food effect owing

to its poor aqueous solubility [38]. Such findings may explain why lurasidone prescription rates

are considerably lower than alternatives such as olanzapine (*e.g.,* 111,911 prescriptions for

lurasidone in Australia compared to 719,915 for olanzapine) [39].

In the context of SCFA luminal accumulation, the lowered pH ensures lurasidone remains in

its ionised form, thereby enhancing aqueous solubility [40]. Enriched microbiota groups

correlated with increased OTUs (**Fig. 2D)** and subsequently increased SCFA production (**Fig.**

2F). Ultimately, these findings correlated strongly (**Fig. 4B:** $R^2 = 0.82$ and **Fig. 4C** $R^2 = 0.87$)

with intestinal pH, suggesting enriched microbiomes presented decreased luminal pH as

corroborated by Walker et al. [9]. The aforementioned pH-dependent solubility of lurasidone

benefits from this SCFA-induced decrease in pH and thus the gut microbiome emerges as a key

mediator between SCFAs, luminal pH and subsequent drug solubility and bioavailability. Such

assertions are corroborated by **Fig. 4E** and **F**, with both exhibiting strong correlations between

the gut microbiome, SCFAs and subsequent impacts on lurasidone AUC_{0-24} (R^2 0.72 and 0.71

respectively). Whilst not previously explored for lurasidone, similar relationships are

summarised in a comprehensive review by Kamath et al. [1].

By extension, the correlations between SCFA, pH and subsequent solubilisation, absorption

and bioavailability of lurasidone illustrate a future wherein predictive models may use SCFAs

as biomarkers for managing personalised therapy. Currently, SCFAs are tested as putative

biomarkers for colorectal and non-small cell lung cancers [41]. However, these scenarios are

limited in scope, restricted to merely suggesting gut inflammation characteristics and immune

activation [42]. Emerging work by Zhao et al. and Tsunoda et al. suggests the gut microbiome

composition as a whole to be capable of predicting drug pharmacokinetics [43,44]. A case example

is demonstrated by Duan et al. which showcased discernible differences in the gut microbiome

of escitalopram responders and non-responders (C57BL/6 male mice model, n: 7/group, mean

age: 4-6 weeks) [45]. Furthermore, the gut microbiome has shown capacity to modulate hepatic

cytochrome P450 (CYP450) characteristics as explored by Wang et al. and Bai et al. [46,47]. In

the context of lurasidone, it presents another avenue of pharmacokinetic modulation as

lurasidone is extensively metabolised (thought oxidative N-dealkylation between the

piperazine and cyclohexane rings) by CYP450, forming lurasidone metabolites 14326, 14283

(both active) and 20219, 20220 (inactive) [48]. By harnessing the gut-liver axis (facilitated by

intestinal permeability), the gut microbiome is capable of altering CYP450 levels and thereby

lurasidone pharmacokinetics [49,50]. As such, identification of specific gut biomarkers (such as

SCFAs metabolites) together with machine-learning based predictive models may provide

personalised pharmacotherapy to mental health patients as explored in reviews by Keutzer et

al. [51].

The insights derived from this study are valuable in broadening our understanding of

pharmacomicrobiomics, in particular of poorly understood lurasidone and are summarised in

Fig 5.

Figure 5. *Enriched microbiome increases bioavailability and solubility, in contrast the*

depleted microbiome decreases bioavailability and solubility. *Three treatment groups and*

their subsequent effects on the gut microbiome and drug bioavailability and solubility.

Given the role of pH and the gut microbiome in modulating lurasidone solubility and

absorption, it may necessitate re-evaluation of treatment regimens for schizophrenia to

accommodate gut microbiome interventions to maximise pharmacotherapy efficacy. Recent

evidence that lurasidone presents 'gut neutral' characteristics and imparts generally favourable

impacts on the gut microbiome, thus it may serve as a better-suited therapeutic option for

metabolic sensitive patients [5]. Furthermore, it is hoped that the exploration of lurasidone's pH-

dependent solubility spur strategies to maximise solubility, absorption and bioavailability, for

example as achieved by Meola et al. through the formulation of a lipid-based nanostructure and

porous silica nanostructure which enhanced bioavailability of lurasidone by 8.8-fold. [52].

Similarly, Gamboa-Arancibia et al. formed a lurasidone-hydroxypropyl-β-cyclodextrin

(obtained through the hydroxylation of β-cyclodextrin) and observed increases in solubility (up

to 20 mg/mL) attributable to the presence of the hydroxyl substituent in this cyclodextrin [53].

Thus, it is expected that the current findings will guide future research in optimising therapy

and potentially lead to increased prescription and clinical use of lurasidone in standard therapy

for treating schizophrenia and bipolar disorder.

5. Conclusion and Prospects:

This study on lurasidone's pharmacomicrobiomics reveals the intricate relationships between

the gut microbiome, SCFA production, luminal pH, and drug bioavailability. The critical role

of commensal bacteria, particularly *Lachnospiraceae* and *Bacteroidaceae*, in modulating the

gut environment highlights the importance of considering microbiome dynamics in

antipsychotic treatment strategies. Lurasidone's pH-dependent solubility and 'gut neutral'

characteristics position it as a promising option for metabolically sensitive patients, despite its

poor bioavailability. Future research should focus on validating these findings in human

studies, interpreting specific mechanisms of lurasidone absorption, and developing novel drug

delivery systems to enhance bioavailability. The success of lipid-based nanostructures and

cyclodextrin complexes in improving solubility suggests promising avenues for formulation

development. These findings underscore the need to re-evaluate treatment regimens for

schizophrenia and bipolar disorder, potentially incorporating additional gut microbiome

interventions to maximise pharmacotherapy efficacy. Additionally, exploring personalised

medicine approaches based on individual gut microbiome profiles and investigating the broader

implications for other pH-sensitive drugs could lead to significant advancements in drug
development and administration.

Acknowledgements: The Hospital Research Foundation (THRF) Group is gratefully acknowledged for their
EMCR Fellowship funding and generous support provided to Dr Paul Joyce (2022-CF-EMCR-004-25314). The
authors acknowledge the University of Adelaide, enabled by NCRIS, university, and state government support.

References:

1. Kamath, S., Stringer, A. M., Prestidge, C. A. & Joyce, P. Targeting the gut microbiome to
control drug pharmacomicrobiomics: the next frontier in oral drug delivery. *Expert Opinion*
on Drug Delivery **20**, 1315–1331 (2023).

2. Leucht, S. *et al.* Dose-Response Meta-Analysis of Antipsychotic Drugs for Acute
Schizophrenia. *AJP* **177**, 342–353 (2020).

3. Minichino, A. *et al.* Psycho-Pharmacomicrobiomics: A Systematic Review and Meta-
Analysis. *Biological Psychiatry* **95**, 611–628 (2024).

4. Davey, K. J. *et al.* Antipsychotics and the gut microbiome: olanzapine-induced metabolic
dysfunction is attenuated by antibiotic administration in the rat. *Transl Psychiatry* **3**, e309–
e309 (2013).

5. Kamath, S., Hunter, A., Collins, K., Wignall, A. & Joyce, P. The atypical antipsychotic
lurasidone positively modulates the gut microbiota in rats: A comparative study to
olanzapine. Preprint at https://doi.org/10.1101/2024.02.28.582623 (2024).

6. Cussotto, S. *et al.* The gut microbiome influences the bioavailability of olanzapine in rats.
eBioMedicine **66**, 103307 (2021).

7. Subramaniam, S. *et al.* Self-emulsifying drug delivery systems (SEDDS) disrupt the gut
microbiota and trigger an intestinal inflammatory response in rats. *International Journal*
of Pharmaceutics **648**, 123614 (2023).

8. Cussotto, S., Clarke, G., Dinan, T. G. & Cryan, J. F. Psychotropics and the Microbiome: a

Chamber of Secrets…. *Psychopharmacology (Berl)* **236**, 1411–1432 (2019).

9. Walker, A. W., Duncan, S. H., McWilliam Leitch, E. C., Child, M. W. & Flint, H. J. pH and

Peptide Supply Can Radically Alter Bacterial Populations and Short-Chain Fatty Acid

Ratios within Microbial Communities from the Human Colon. *Appl Environ Microbiol* **71**,

3692–3700 (2005).

10. Firrman, J. *et al.* The impact of environmental pH on the gut microbiota community

structure and short chain fatty acid production. *FEMS Microbiol Ecol* **98**, fiac038 (2022).

11. Venkataraman, A. *et al.* Variable responses of human microbiomes to dietary

supplementation with resistant starch. *Microbiome* **4**, 33 (2016).

12. Deehan, E. C. *et al.* Precision Microbiome Modulation with Discrete Dietary Fiber

Structures Directs Short-Chain Fatty Acid Production. *Cell Host & Microbe* **27**, 389-404.e6

(2020).

13. McOrist, A. L. *et al.* Fecal Butyrate Levels Vary Widely among Individuals but Are Usually

Increased by a Diet High in Resistant Starch1,2. *The Journal of Nutrition* **141**, 883–889

(2011).

14. Gibson, G. R. & Roberfroid, M. B. Dietary Modulation of the Human Colonic Microbiota:

Introducing the Concept of Prebiotics. *The Journal of Nutrition* **125**, 1401–1412 (1995).

15. Nguyen, T. T. *et al.* Differences in gut microbiome composition between persons with

chronic schizophrenia and healthy comparison subjects. *Schizophrenia Research* **204**, 23–

29 (2019).

16. Wos-Oxley, M. L. *et al.* Comparative evaluation of establishing a human gut microbial

community within rodent models. *Gut Microbes* **3**, 234–249 (2012).

451 17. Forsgård, R. A. *et al.* Chemotherapy-induced gastrointestinal toxicity is associated with

452   changes in serum and urine metabolome and fecal microbiota in male Sprague–Dawley

453   rats. *Cancer Chemother Pharmacol* **80**, 317–332 (2017).

454 18. Letertre, M. P. M. *et al.* A Two-Way Interaction between Methotrexate and the Gut

455   Microbiota of Male Sprague–Dawley Rats. *J. Proteome Res.* **19**, 3326–3339 (2020).

456 19. Hickman, D. L., Johnson, J., Vemulapalli, T. H., Crisler, J. R. & Shepherd, R. Commonly

457   Used Animal Models. *Principles of Animal Research for Graduate and Undergraduate*

458   *Students* 117–175 (2017) doi:10.1016/B978-0-12-802151-4.00007-4.

459 20. Panebianco, C., Andriulli, A. & Pazienza, V. Pharmacomicrobiomics: exploiting the drug-

460   microbiota interactions in anticancer therapies. *Microbiome* **6**, 92 (2018).

461 21. Doestzada, M. *et al.* Pharmacomicrobiomics: a novel route towards personalized medicine?

462   *Protein Cell* **9**, 432–445 (2018).

463 22. Dikeocha, I. J., Al-Kabsi, A. M., Miftahussurur, M. & Alshawsh, M. A.

464   Pharmacomicrobiomics: Influence of gut microbiota on drug and xenobiotic metabolism.

465   *The FASEB Journal* **36**, e22350 (2022).

466 23. Huang, J. *et al.* Probiotics Plus Dietary Fiber Supplements Attenuate Olanzapine-Induced

467   Weight Gain in Drug-Naïve First-Episode Schizophrenia Patients: Two Randomized

468   Clinical Trials. *Schizophrenia Bulletin* **48**, 850–859 (2022).

469 24. Fusco, W. *et al.* Short-Chain Fatty-Acid-Producing Bacteria: Key Components of the

470   Human Gut Microbiota. *Nutrients* **15**, 2211 (2023).

471 25. Vacca, M. *et al.* The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms*

472   **8**, 573 (2020).

473 26. Zhang, Z. J. *et al.* Comprehensive analyses of a large human gut Bacteroidales culture

474   collection reveal species and strain level diversity and evolution. Preprint at

475   https://doi.org/10.1101/2024.03.08.584156 (2024).

27. Carlson, J., Erickson, J., Hess, J., Gould, T. & Slavin, J. Prebiotic Dietary Fiber and Gut

Health: Comparing the in Vitro Fermentations of Beta-Glucan, Inulin and

Xylooligosaccharide. *Nutrients* **9**, 1361 (2017).

28. Tian, R. *et al.* Effect of inulin, galacto-oligosaccharides, and polyphenols on the gut

microbiota, with a focus on *Akkermansia muciniphila*. *Food Funct.* **15**, 4763–4772 (2024).

29. Gurry, T., Nguyen, L. T. T., Yu, X. & Alm, E. J. Functional heterogeneity in the

fermentation capabilities of the healthy human gut microbiota. *PLoS ONE* **16**, e0254004

(2021).

30. Liu, L., Li, Q., Yang, Y. & Guo, A. Biological Function of Short-Chain Fatty Acids and Its

Regulation on Intestinal Health of Poultry. *Front. Vet. Sci.* **8**, 736739 (2021).

31. Parada Venegas, D. *et al.* Short Chain Fatty Acids (SCFAs)-Mediated Gut Epithelial and

Immune Regulation and Its Relevance for Inflammatory Bowel Diseases. *Front. Immunol.*

10, 277 (2019).

32. Sun, Y. & O'Riordan, M. X. D. Regulation of Bacterial Pathogenesis by Intestinal Short-

Chain Fatty Acids. in *Advances in Applied Microbiology* vol. 85 93–118 (Elsevier, 2013).

33. Ishibashi, T. *et al.* Pharmacological Profile of Lurasidone, a Novel Antipsychotic Agent

with Potent 5-Hydroxytryptamine 7 (5-HT $_7$) and 5-HT $_{1A}$ Receptor Activity. *J Pharmacol*

Exp Ther **334**, 171–181 (2010).

34. Lacivita, E., Leopoldo, M., Giorgio, P. D., Berardi, F. & Perrone, R. Determination of 1-

aryl-4-propylpiperazine pKa values: The substituent on aryl modulates basicity. *Bioorganic*

& Medicinal Chemistry **17**, 1339–1344 (2009).

35. Bergström, C. A. S. & Avdeef, A. Perspectives in solubility measurement and

interpretation. *ADMET DMPK* **7**, 88–105 (2019).

36. Kumari, L. *et al.* Advancement in Solubilization Approaches: A Step towards

Bioavailability Enhancement of Poorly Soluble Drugs. *Life* **13**, 1099 (2023).

501 37. Selmin, F. *et al.* Relevance of production method on the physical stability and in vitro

502   biopharmaceutical performances of olanzapine orodispersible film. *International Journal*

503   *of Pharmaceutics* **603**, 120697 (2021).

504 38. Meola, T. R., Joyce, P., Wignall, A., Bremmell, K. E. & Prestidge, C. A. Harnessing the

505   potential of nanostructured formulations to mimic the food effect of lurasidone.

506   *International Journal of Pharmaceutics* **608**, 121098 (2021).

507 39. Medicare Benefits Scheme funded services: monthly data, About. *Australian Institute of*

508   *Health and Welfare* https://www.aihw.gov.au/reports/medicare/mbs-funded-services-

509   data/contents/about (2024).

510 40. Meola, T. R., Paxton, K., Joyce, P., Schultz, H. B. & Prestidge, C. A. The effect of drug

511   ionization on lipid-based formulations for the oral delivery of anti-psychotics. *ADMET &*

512   *DMPK* **8**, 437 (2020).

513 41. Chalova, P. *et al.* Determination of short-chain fatty acids as putative biomarkers of cancer

514   diseases by modern analytical strategies and tools: a review. *Front Oncol* **13**, 1110235

515   (2023).

516 42. Carretta, M. D., Quiroga, J., López, R., Hidalgo, M. A. & Burgos, R. A. Participation of

517   Short-Chain Fatty Acids and Their Receptors in Gut Inflammation and Colon Cancer. *Front*

518   *Physiol* **12**, 662739 (2021).

519 43. Zhao, Q., Chen, Y., Huang, W., Zhou, H. & Zhang, W. Drug-microbiota interactions: an

520   emerging priority for precision medicine. *Sig Transduct Target Ther* **8**, 1–27 (2023).

521 44. Tsunoda, S. M., Gonzales, C., Jarmusch, A. K., Momper, J. D. & Ma, J. D. Contribution of

522   the Gut Microbiome to Drug Disposition, Pharmacokinetic and Pharmacodynamic

523   Variability. *Clin Pharmacokinet* **60**, 971–984 (2021).

524 45. Duan, J. *et al.* Characterization of gut microbiome in mice model of depression with

525   divergent response to escitalopram treatment. *Transl Psychiatry* **11**, 1–12 (2021).

46. Wang, S. *et al.* Gut microbiota and host cytochrome P450 characteristics in the pseudo

germ-free model: co-contributors to a diverse metabolic landscape. *Gut Pathogens* **15**, 15

(2023).

47. Bai, X. *et al.* Regulation of CYP450 and drug transporter mediated by gut microbiota under

high-altitude hypoxia. *Front Pharmacol* **13**, 977370 (2022).

48. Kot, M., Haduch, A., Papp, M. & Daniel, W. A. The Effect of Chronic Treatment with

Lurasidone on Rat Liver Cytochrome P450 Expression and Activity in the Chronic Mild

Stress Model of Depression. *Drug Metab Dispos* **45**, 1336–1344 (2017).

49. Yan, M. *et al.* Gut liver brain axis in diseases: the implications for therapeutic interventions.

Sig Transduct Target Ther **8**, 1–26 (2023).

50. Albillos, A., de Gottardi, A. & Rescigno, M. The gut-liver axis in liver disease:

Pathophysiological basis for therapy. *J Hepatol* **72**, 558–577 (2020).

51. Keutzer, L. *et al.* Machine Learning and Pharmacometrics for Prediction of

Pharmacokinetic Data: Differences, Similarities and Challenges Illustrated with

Rifampicin. *Pharmaceutics* **14**, 1530 (2022).

52. Meola, T. R. *et al.* Inulin-Lipid Core–Shell Microcapsules Target the Gut Microbiota and

Mimic the Pharmaceutical Food Effect for Improved Oral Antipsychotic Delivery. *Adv*

Funct Materials 2403914 (2024) doi:10.1002/adfm.202403914.

53. Gamboa-Arancibia, M. E. *et al.* Improving Lurasidone Hydrochloride's Solubility and

Stability by Higher-Order Complex Formation with Hydroxypropyl-β-cyclodextrin.

Pharmaceutics **15**, 232 (2023).

547