Navigating the Cold: Integrative Transcriptome Sequencing Approach Reveals Ionoregulatory and Whole-Body Responses to Cold Acclimation in *Drosophila ananassae*

Vera Miyase <u>Yılmaz</u>[1], Zhihui <u>Bao</u>[1], Sonja <u>Grath</u>[1*]

[1]Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany

[*]Corresponding author: grath@bio.lmu.de

Running title: Adaptation to cold in *Drosophila*

Abstract

Understanding how species adapt to changing environments is a major goal in evolutionary biology and can elucidate the impact of climate change. Climate imposes inevitable effects on the geographical distribution of insects as their body temperature primarily depends on the environment. The vinegar fly *Drosophila ananassae* expanded from its tropical ancestral range to more temperate regions, which requires adaptation to colder temperatures. Transcriptome and genome-wide association studies focusing on the ancestral-range population identified the targets of selection related to ionoregulatory tissues. However, how cosmopolitan *D. ananassae* adapted to colder environments, where low temperatures last longer, is still unknown. Here, we present a study on the effect of long-term cold exposure on *D. ananassae*, examining the gene expression variation in the whole body and the ionoregulatory tissues, namely the hindgut and the Malpighian tubule. To elucidate molecular mechanisms of cold adaptation during species expansion, we included cold-tolerant and cold-sensitive strains from the ancestral species range and cold-tolerant strains from the derived species range. We show that cold acclimation improves cold tolerance and results in differential expression of more than half of the transcriptome in the ionoregulatory tissues and the whole body. Notably, we provide complementary insight into molecular processes at four levels: strains, populations, phenotypes, and tissues. By determining the biochemical pathways of phenotypic plasticity underlying cold tolerance, our results enhance our understanding of how environmental changes affect thermal adaptation in natural populations.

36   **Keywords:** thermoregulation, ectothermy, metabolic cold adaptation, gene expression,

37   phenotypic plasticity, ionoregulatory evolution, acclimation

Significance

The physiological mechanisms underlying cold tolerance of insects are complex. Our study integrates tissue-specific responses with whole-body response to cold and compares these responses in different populations, phenotypes, and strains of the vinegar fly *Drosophila ananassae*. The study reveals that the response to extended cold in the renal and intestinal tissues are different, that the tissue-specific responses can be masked in the whole body, and that the transcriptomic response to extended cold can depend on population of origin, cold tolerance phenotype, or strain. Overall, this study sheds light on insect cold adaptation in terms of transcriptomic response to cold acclimation in the ionoregulatory tissues and whole bodies.

Introduction

Temperature is a major factor influencing how species adapt to changing environments and colonize new ecosystems. How animals generally cope with thermal stressors and which genes are specifically responsible for cold adaptation has been a topic of intense research for decades in molecular and evolutionary biology. Recent developments in high-throughput sequencing methods allow insights into molecular mechanisms at the whole transcriptome and genome level, but integrating the complementary information of various experimental designs remains challenging. For example, studies on small insects often rely on either systemic or tissue-specific methodologies or single phenotypes, resulting in incomplete conclusions (Andersen *et al.*, 2015; Hoedjes *et al.*, 2024). Therefore, integrating different experimental designs and phenotypes to study transcriptional responses to cold is critical to uncovering mechanisms underlying physiological adaptation. Elucidating the mechanisms that underlie adaptation to changing environments in natural populations and integrating different approaches can help us understand

which species prove to be successful in the future and have potential applications for

conservation, agriculture, and pest management.

Many insects evolved and diversified in warm climates (Bale & Hayward, 2010). Nevertheless,

insects have developed various strategies to cope with thermal stress and to successfully expand

their thermal and habitat ranges (Overgaard & MacMillan, 2017). Some insects even tolerate

freezing and remain active, such as the Antarctic midge *Belgica antarctica* (Teets *et al.*, 2012) or

the alpine beetle *Pytho deplanatus* (Ring, 1982). However, a wide range of insect species suffer

from chilling injuries and eventually die when they are exposed to cold. This chill-susceptibility

has been reported for several orders of insects (Overgaard & MacMillan, 2017). Chill-susceptible

species include some of great economic and ecological importance, such as the honeybee *Apis*

mellifera (Free & Spencer-Booth, 1960), but also agricultural pests, for instance, the migratory

locust *Locusta migratoria* (Bayley *et al.*, 2018) and disease vectors such as mosquitoes

(Reinhold *et al.*, 2018).

The physiology of cold stress resistance in insects is highly complex. When insects are exposed

to cold, their ion and water homeostasis is disrupted (MacMillan *et al.*, 2015). Additionally,

transcriptomic and metabolic profiles are changed (MacMillan *et al.*, 2016). Along with these

systemic changes, chilling injuries affect the body at the cell level as they cause phase changes in

the cell membrane, resulting in a stiff membrane with decreased permeability (Steponkus, 1984;

Koštál, 2010; Drobnis *et al.*, 1993). Further, cold exposure results in an imbalance of passive

diffusion and active ion transport (MacMillan & Sinclair, 2011; Koštál *et al.*, 2004; Zachariassen

et al., 2004). An excess of potassium ions in the extracellular space leads to depolarization of the

membrane (Findsen *et al.*, 2014; Hoyle, 1953), eventually activating the voltage-gated calcium

channels (Salkoff & Wyman, 1983; Findsen *et al.*, 2016). The cellular influx of calcium ions

through voltage-gated channels activates apoptotic and necrotic pathways, leading to chill injury

(Koštál *et al.*, 2004, 2007, 2016; Zachariassen *et al.*, 2004; MacMillan *et al.*, 2015, 2016).

Consequently, the regulation of ion and water homeostasis has been identified as a key

physiological feature for cold tolerance in insects with specifically the ionoregulatory tissues

being essential for maintaining the water-ion balance in an insect body (Andersen *et al.*, 2017;

Andersen & Overgaard, 2020; Des Marteaux *et al.*, 2018; Gerber & Overgaard, 2018; MacMillan

et al., 2015, 2016, 2017; Yerushalmi *et al.*, 2018). Despite several in-depth studies aiming to

resolve this complexity, it remains challenging to identify and functionally characterize specific

genes and genetic elements that contribute to the diversity of cold stress-resistant phenotypes

(Udaka *et al.*, 2010; von Heckel *et al.*, 2016; MacMillan *et al.*, 2016). With their worldwide

distribution, Drosophilids are excellent models for studying these questions.

Flies from the genus *Drosophila* suffer from impaired synaptic transmission and muscular failure

when they are exposed to cold; in other words, they lose their neuromuscular coordination and

enter a comatose condition referred to as chill coma in which they are not able to move, feed,

mate or escape predators (Kelty *et al.*, 1996; MacMillan & Sinclair, 2011; Findsen *et al.*, 2014).

However, flies can recover from such a coma given that time and temperature of exposure to

cold do not exceed certain thresholds. The time the fly needs for recovery from a cold-induced

coma (chill coma recovery time, CCRT) is widely used to assess the cold hardiness of insects in

the laboratory and serves as a predictor for the climatic condition at the geographical sampling

site of a particular species (Gibert *et al.*, 2001). Further, CCRT follows latitudinal clines within

Drosophila species in Australia (Hoffmann *et al.*, 2002; Hoffmann & Weeks, 2007; Hallas *et al.*,

2002), Asia (Sisodia & Singh, 2010), Africa and Europe (David *et al.*, 2003). Cold tolerance in

Drosophilids is polygenic. Despite this complexity, variation in gene expression of only a few

loci is associated with cold tolerance in subtropical populations of *Drosophila ananassae*

(Königer & Grath, 2018), and variation of just three genomic regions explained more than 60%

of the phenotypic variation between cold-tolerant and cold-sensitive strains (Königer *et al.*,

2019). Recently, we provided Cas9-integrated fly strains that facilitate the use of the CRISPR-

Cas9 system in this species and allow functional characterization of individual candidate loci

(Yılmaz *et al.*, 2023). In sum, *D. ananassae* has been established as a promising model for

studying the genetic basis of cold tolerance.

Previous studies mostly neglected the potential of integrating two specific aspects to better

inform about cold tolerance in insects. First, how insects respond to low temperatures can be

plastic: the ability to tolerate cold and the genes responsible for the survival can change

depending on prior exposure to cold. Such acclimation has been shown to alleviate the adverse

effects of cold and can increase cold tolerance (Bubliy *et al.*, 2002; Enriquez & Colinet, 2019;

Hoffmann & Watson, 1993; MacMillan *et al.*, 2016, 2017; Watson & Hoffmann, 1996; Yılmaz *et*

al., 2023). Cold acclimation triggers substantial changes in the transcriptional response to cold in

Drosophila suzukii (Enriquez & Colinet 2019). It reorganizes the transcriptome and metabolome

of whole animals (MacMillan *et al.*, 2016) and the gut transcriptome in *Drosophila melanogaster*

(MacMillan *et al.*, 2017). Second, conclusions on potentially ecologically relevant adaptations

should not rely on a single phenotype (Andersen *et al.*, 2015). Consequently, incorporating

distinct cold tolerance phenotypes and systemic and ionoregulatory tissue-specific transcriptome

data will allow pinpointing cold-related physiological adaptations.

In this study, we use isofemale lines collected from a population of ancestral origin in Bangkok,

Thailand (from here on mentioned as ancestral population) and a derived population originating

from Kathmandu, Nepal (Das *et al.*, 2004). We examined the transcriptional response to cold

acclimation in both the ancestral and the derived populations to shed light on the genetic

mechanisms underlying cold tolerance. In addition, we integrated the original results of this

study with previously gained insight into the distinct phenotype of cold shock. To this end, we

contrasted the transcriptional response to cold at four levels: (1) population (ancestral vs.

derived), (2) strain (cold-sensitive vs. cold-tolerant), (3) tissue (whole-body vs. ionoregulatory

tissues, Malpighian tubules and hindgut), and (4) phenotype (cold acclimation vs. cold shock).

We find empirical evidence for the influence of population and tissue on gene expression

variation upon cold stress. Using an RNA sequencing approach that combined systemic with

tissue-specific transcriptomes, we show that cold acclimation improves cold tolerance and results

in more than 50% of the transcriptome being differentially expressed in the whole body and the

ionoregulatory tissues.

Our results demonstrate that an RNA sequencing strategy combining a systemic overview with

tissue-specific data can be a cost-effective and informative way to elucidate genes that have

major effects on a crucial adaptation-related trait, namely, thermoregulation.

143 Results

Complementary elucidation of transcriptional response to cold acclimation with systemic

whole body and tissue-specific data

Flies from ancestral and derived populations were previously described as having different

sensitivity to cold shock as determined by chill coma recovery time, CCRT (Königer & Grath,

2018). Additionally, preliminary tests revealed that acclimation to cold improves the tolerance

and decreases the chill coma recovery time (see Supplementary Text). To receive an informative

overview of the transcriptional response to cold acclimation in *D. ananassae*, we combined high-

throughput RNA sequencing of the whole body (wb), as well as of two ionoregulatory tissues,

Malpighian tubules (Mt) and hindguts (hg). We obtained RNA sequencing data from male flies

upon 4-day acclimation at 10°C. In the whole body and hindgut samples, 54% of all expressed

genes were differentially expressed in response to cold acclimation, and the percentage of DE

genes in the Malpighian tubules was 50% (Table 1). The principal component analysis of all

samples revealed that they clustered according to the tissue RNA was extracted from, whereas no

separation according to the treatment was observed (Figure 1A). The first principal component

explained 68% of the variance, separating the ionoregulatory tissues from the whole-body

samples. The second principal component explained 19% of the variance, separating the two

ionoregulatory tissues. Though pairwise comparisons revealed that 2952 differentially expressed

genes were shared between sample types, a considerable number of genes (ranging from 505

genes that were explicitly up-regulated in Malpighian tubules to 1436 genes that were down-

regulated explicitly in the whole body) were specifically up- or down-regulated only in the

individual tissues (Figure 1B-C). Notably, these genes were enriched in distinct GO categories

(Supplementary Table 3-4). These results suggest that the combination of systemic and informed

165 tissue-specific RNA sequencing data allows for complementary conclusions on cold tolerance in

166 *Drosophila*.

167 **Table 1:** Number of differentially expressed genes in different sample types regarding the general,
168 population-specific, phenotype-specific, and line-specific response to cold acclimation

	total expressed genes	up regulated	down regulated	total DE
General response to acclimation				
Whole body	14304	3825 (27%)	3930 (27%)	7755 (54%)
Malpighian tubule	12189	3280 (27%)	2874 (24%)	6154 (50%)
Hindgut	13272	3715 (28%)	3480 (26%)	7195 (54%)
Population-specific response				
Whole body				
BKK	14300	3618 (25%)	3668 (26%)	7286 (51%)
KATH	14300	3414 (24%)	3497 (24%)	6911 (48%)
KATH vs BKK*	13191	1181 (9%)	827 (6.3%)	2008 (15%)
Malpighian tubule				
BKK	11721	3077 (26%)	2606 (22%)	5683 (48%)
KATH	11191	2288 (20%)	2174 (19%)	4462 (40%)
KATH vs BKK*	9125	124 (1.4%)	92 (1%)	216 (2.4%)
Phenotype-specific response				
Hindgut				
Fast	12821	3203 (25%)	3570 (28%)	6773 (53%)
Slow	12547	2907 (23%)	2784 (22%)	5691 (45%)
Fast vs Slow*	12547	182 (1.5%)	246 (2%)	428 (3.4 %)
Line-specific response**				
Whole body				
FastBKK	14300	3333 (23%)	3434 (24%)	6767 (47%)
SlowBKK	14300	3091 (22%)	3300 (23%)	6391 (45%)
FastBKK vs SlowBKK*	13468	508 (3.8%)	472 (3.5%)	980 (7.3%)

*   For these comparisons, the columns show the number of genes that are significantly differentially
expressed between the compared groups. Up and down-regulated columns indicate direction regarding the
effect of acclimation on the former group compared to the latter group. BKK = Bangkok, KATH =
Kathmandu.

**   Only isofemale lines of the ancestral population (BKK) are taken into consideration. The terms
'fast' and 'slow' indicate faster and slower recovery, respectively.

Figure 1: Transcriptomic response to cold acclimation in ionoregulatory tissues and the whole body. A) Principal component analysis. The samples are shown according to the tissue (color) and treatment (shape). Whole body (Wb), Malpighian tubule (Mt), and hindgut (Hg) tissues are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Venn diagram showing the number of DE genes specific to or shared among sample types: Whole body, Malpighian tubule and Hindgut samples exhibited both shared, as well as exclusive differentially expressed genes. C) Upset plot showing differentially expressed genes and their overlap between the three different sample types. Black, horizontal bars indicate the total number of differentially expressed genes that were significantly up- or down-regulated in a sample in response to cold acclimation. A black circle indicates that a sample and direction (up/down) combination is included in an intersection class. Intersection classes with a single circle are comprised of genes significantly differentially expressed in a single tissue and direction. Circles connected by a line indicate intersection classes comprised of multiple tissue and direction combinations. Vertical bars correspond to the number of genes in each intersection class, colors represent the respective sections of the Venn diagram.

Strong systemic transcriptional response to cold acclimation

We performed RNA sequencing on whole-body samples to obtain a systemic overview of the

transcriptional response to cold acclimation in *Drosophila ananassae*. This approach has the

unique advantage of being both cost-efficient and unbiased in terms of preconditions on

transcription. Samples of acclimation and control treatments were clustered according to the

populations of origin (Figure 2A). The first principal component explained 58% of the variance.

It separated cold-acclimated samples from control samples, while the second principal

component explained 14% of the variance and separated the derived population from the

ancestral population. Within the ancestral population, the samples also clustered according to

their CCRT phenotype (Figure 2B). We addressed three questions with this data: (1) What is the

general, systemic transcriptional response to cold acclimation? (2) How does transcriptional

response differ between ancestral and derived populations? (3) How does transcriptional

response differ between cold-tolerant and cold-sensitive strains in the ancestral population?

More than half of the transcriptome (54%) was differentially expressed between control and

treatment, with 3825 genes significantly up-regulated and 3930 genes significantly down-

regulated (Table 1). Gene ontology enrichment analysis revealed that up-regulated genes were

enriched in biological processes related to translation and cell junction organization (Figure 2C,

Supplementary Table 5). On the contrary, down-regulated genes were enriched in metabolic

processes (Figure 2D, Supplementary Table 5). These results suggest a solid response to cold

acclimation in the whole body, and the transcriptional response of populations differs not only in

the presence of cold treatment but also at the basal level.

Figure 2: Transcriptomic response to cold acclimation in the whole body. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Heatmap for the sample pools. Sample clusters are shown on the left and top of the heatmap. The letters indicate the sample pools FastBKK (F), SlowBKK (S), Kath (K). Cold acclimated samples are

219 indicated by delta (Δ) sign. The numbers correspond to the replicates. The similarity matrix indicates high
220 (red) and low (blue) similarity of samples. Gene ontology enrichment analysis results for genes C) up-
221 regulated and D) down-regulated in the whole body in response to cold acclimation. Biological processes
222 (red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate
223 panels.

224 Systemic differences between populations are mediated by genes involved in detoxification

225 and maintaining ion balance

To characterize the differences between populations, we contrasted the transcriptional response

of ancestral and derived populations. A total of 14300 genes were expressed in each population

(Table 1). In the ancestral population, 51% of these genes were differentially expressed upon

cold acclimation, while 3618 genes were significantly up- and 3668 down-regulated,

respectively. In comparison, 48% of these genes were differentially expressed in the derived

population, while 3414 genes were significantly up-regulated, and 3497 genes were significantly

down-regulated. Although the populations shared over half of the differentially expressed genes

(Supplementary Figure 2A), 2008 genes showed significant differential expression between

populations in response to cold acclimation. The GO analysis suggested that the genes with

differential expression between the populations were mainly located in muscle tissues and

enriched in cilium movement and various metabolic processes, including fatty acid metabolism

(Supplementary Figure 2B). Notably, these genes were also enriched in molecular functions

related to ion binding, ion channel, and oxidoreductase activity (Supplementary Table 6).

Overall, these results suggest the difference between the acclimation response of populations

ground on ion transport and detoxification.

241 The actin cytoskeleton differentiates the acclimation response of tolerant and sensitive

242 strains in the ancestral population

To address the question of how the transcriptional response differs within the ancestral

population between cold-tolerant and cold-sensitive strains, we compared gene expression upon

cold acclimation in both types of strains. For the cold-tolerant strains, 3333 genes were

significantly up-regulated, and 3434 genes were significantly down-regulated (Table 1). For the

sensitive strains, 3091 genes were significantly up-regulated, and 3300 genes were significantly

down-regulated. However, 980 genes were differentially expressed between cold-sensitive and

cold-tolerant strains. GO enrichment analysis revealed significant enrichment of these genes in

various biological processes related to the actin cytoskeleton. (Supplementary Table 7)

Adaptive transcriptional response is evident in the renal tissues

Malpighian tubules (Mt) are the renal equivalent of mammals in insects and contribute to

maintaining homeostasis, which can be disrupted by cold exposure. Therefore, we investigated

the transcriptional response of Mt to characterize the impact of cold acclimation on renal tissues

in *D. ananassae*. Like the whole-body samples, RNA samples extracted from Mt of acclimated

and control flies were clustered according to the populations of origin, suggesting that the

populations are distinct in their response to acclimation and at the basal level of gene expression

(Figure 3A). Principal component analysis indicated that 37% of the variation can be explained

by the first principal component, which separated cold-acclimated and control samples, and 14%

of the variation can be explained by the second principal component, which separated the

ancestral and derived populations. Therefore, we investigated the general transcriptional

response of Mt to cold acclimation and the population-specific response.

In the Mt samples, 3280 genes were up-regulated, and 2874 were down-regulated (Table 1).

Similar to the whole-body samples, up-regulated genes of Mt were enriched in GO terms related

to translation and transcription (Supplementary Figure 3A, Supplementary Table 8), and the

down-regulated genes were enriched in processes related to fatty acid degradation

(Supplementary Figure 3B, Supplementary Table 8). However, contrary to the whole-body

samples, the down-regulated genes of the Mt were also enriched in chromatin remodeling.

Additionally, a major part of the down-regulated genes was located in the mitochondrion and,

more specifically, the respiratory chain complex.

Figure 3: Transcriptomic response to cold acclimation in the Malpighian tubules. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Venn diagram showing the number of DE genes specific to or shared between populations: Bangkok and Kathmandu populations exhibited both shared, as well as exclusive differentially expressed genes in response to acclimation. Gene ontology (GO) enrichment analysis results for genes C) up-regulated and

D) down-regulated in the Malpighian tubules of Kathmandu population compared to Bangkok population
in response to cold acclimation. Biological processes (red bars), cellular components (blue bars), and
molecular functions (yellow bars) are shown in separate panels.

Differences in renal tissues of populations are mediated by genes involved in ion transport

and lipid metabolism

In order to understand the differences in metabolic processes involved in cold adaptation

between the ancestral and derived populations, we compared the transcriptional response to cold

acclimation in flies from the Bangkok and Kathmandu populations. In the ancestral population,

11721 genes were expressed, with 3077 up-regulated and 2606 down-regulated (Figure 3B, Table

1). In the derived population, 11191 genes were expressed, with 2288 up-regulated and 2174

down-regulated. Among these DE genes, 216 genes showed differential expression between the

populations. We conducted a GO analysis to explore the functions of these genes and found GO

terms related to transposition. To enhance the analysis, we performed separate analyses for gene

sets: genes up regulated in the derived population compared to the ancestral one, and *vice versa*.

The analysis showed that genes up regulated in the Kathmandu population compared to Bangkok

were enriched in molecular functions related to ion transport (Figure 3C, Supplementary Table

9). Conversely, genes up regulated in the ancestral population compared to the derived one were

enriched in processes and functions related to transposition (Figure 3D, Supplementary Table 9).

Additionally, this analysis revealed an enrichment in molecular functions related to fatty acid

ligase activity.

Hindgut responds to cold acclimation with respect to phenotype, but not to population

Along with the Malpighian tubules, hindguts (hg) are essential in regulating ion and water

balance. Therefore, we investigated the tissue-specific transcriptional response to cold

acclimation in the hindgut of *D. ananassae* flies to elucidate molecular mechanisms of cold

adaptation in insects. Initial PCA revealed two clusters and an outlier sample (Figure 4A). When

the outlier was removed to increase the resolution, PC1 explained 48% of the variance, and PC2

explained 12%, separating the treatments and phenotypes, respectively (Supplementary Figure

4A). The acclimated samples clustered according to their cold tolerance phenotype, in other

words, according to their recovery time from chill coma. However, as opposed to the whole body

and MT, the control samples showed no apparent separation, which suggests that the strains do

not differ at their basal gene expression level according to their cold tolerance phenotype or

population of origin. Therefore, we addressed two questions with this data: (1) What is the

general response to cold acclimation in the hindguts? (2) How does transcriptional response

differ between cold-tolerant and cold-sensitive strains?

Figure 4: Transcriptomic response to cold acclimation in the hindguts. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. Gene ontology (GO) enrichment analysis results for genes B) up-regulated and C) down-regulated in the hindguts of tolerant flies (fast phenotype) compared to sensitive flies (slow phenotype) in response to cold

acclimation. Biological processes (red bars), cellular components (blue bars), and molecular functions
(yellow bars) are shown in separate panels.

In the hindgut samples, we observed a significant up-regulation of 3835 genes and a down-

regulation of 3483 genes in response to cold acclimation (Table 1). Like the whole body and

Malpighian tubules, the up-regulated genes in the hindguts were enriched in GO terms related to

translation and transcription (Supplementary Figure 4B, Supplementary Table 10). In contrast,

the down-regulated genes showed enrichment in biological processes related to cellular

respiration and lipid metabolism, with many of these genes located in the mitochondrial

membrane (Supplementary Figure 4C, Supplementary Table 10).

Cellular respiration and chitin metabolism play roles in the acclimation response of
tolerant and sensitive strains in the hindgut

We characterized the transcriptional response to cold acclimation in cold-tolerant and cold-

sensitive strains with differential gene expression analysis. In the cold-sensitive strains, 3046

genes were significantly up-regulated, and 2886 genes were significantly down-regulated in

response to acclimation (Table 1). On the other hand, in the cold-tolerant strains, 3401 genes

were significantly up-regulated, and 3224 genes were significantly down-regulated in response to

cold acclimation. Between the tolerant and sensitive strains, 330 genes were differentially

expressed. GO enrichment analysis revealed that these genes were enriched in biological

processes related to translation and ribosome biogenesis and in molecular functions related to

chitin binding. We separately investigated the genes up-regulated in the cold-tolerant strains

compared to the sensitive strains and *vice versa*. Genes significantly up-regulated in the cold-

tolerant strains compared to the cold-sensitive strains were enriched in GO terms related to

cuticle development, respiration, and metabolism (Figure 4B, Supplementary Table 11).

Additionally, the products of these genes were localized mainly to the membrane and

extracellular matrix. Genes significantly up-regulated in the cold-sensitive strains compared to

the cold-tolerant strains were enriched in GO terms related to translation (Figure 4C,

Supplementary Table 11).

Distinct transcriptional response to cold acclimation and cold shock

To receive a comprehensive overview of cold tolerance measured by distinct phenotypes, we

contrasted our previous results of the transcriptional response to cold shock as measured by chill

coma recovery time upon cold shock (Königer & Grath, 2018) with the transcriptome data from

this study. To this end, we considered the genes differentially expressed in the cold-tolerant and

cold-sensitive strains of the ancestral population from Bangkok, Thailand, in two conditions: (1)

cold shock and (2) cold acclimation.

Our findings reveal the intricate nature of gene expression in response to cold shock and cold

acclimation. We observed that more genes are differentially expressed in response to cold

acclimation than cold shock (Supplementary Figure 5), indicating a more complex regulatory

network in the former. To understand these responses, we investigated the up- and down-

regulated genes in both treatments and performed a GO enrichment analysis.

The genes up-regulated in response to both treatments were enriched in actin cytoskeleton-

related GO terms (Supplementary Table 12). In contrast, the down-regulated ones showed

enrichment in various metabolic processes such as organic acid, lipid, and carbohydrate

metabolic processes (Supplementary Table 12). We also investigated the genes with opposite

expression (i.e., up-regulated in one treatment but down-regulated in the other) in response to

treatments. Among the 1932 genes that showed differential expression in both studies, 390 genes

had opposite directions. The genes up-regulated upon cold shock but down-regulated upon

acclimation were enriched in the immune response (Supplementary Table 13). On the other hand,

the genes that were up-regulated in response to acclimation but down-regulated after cold shock

showed enrichment in biological processes related to transcription and translation

(Supplementary Table 13).

Finally, we investigated the genes that show differential expression in response to only one of the

treatments. Over 5000 genes were up- or down-regulated upon acclimation, but only 1070 genes

were uniquely expressed after cold shock. The genes differentially expressed only upon the cold

shock treatment were enriched in protein folding, actin cytoskeleton organization, replication,

and glycosylation (Supplementary Table 14). The genes that were differentially expressed upon

acclimation showed enrichment in translation and transcription as well as amino acid metabolism

and copulation (Supplementary Table 14).

Discussion

Understanding the differences between the ancestral and derived populations and the variance

within the ancestral Bangkok population sheds light on the adaptive processes leading to

increased cold tolerance of the species. The two populations were tested for their acclimation

response. Both populations showed improved cold tolerance upon acclimating to mild

temperatures as expected, but the underlying mechanisms were unknown. Comparative

transcriptomics using the whole-body samples provided the systemic overview to understand the

differences in the acclimation response between the two populations and within the ancestral

population. Additionally, the tissue-specific response to acclimation provided further insight into

population and phenotypic differences.

In a complex phenotype, such as cold response, transcriptomic analysis of a whole body might

lead to missing information (reviewed in Hoedjes *et al.*, 2024). However, tissue-specific RNA-

Seq can unmask details otherwise lost in the systemic approach. Both whole-body and tissue-

specific approaches were used in this study to increase the strength of the comparison of

populations, phenotypes, and strains. The DE genes in the Malpighian tubule samples enriched in

hypoxia response and HSP binding activity are not differentially expressed in the whole-body

samples. Similarly, the DE genes in the hindgut samples enriched in sodium channel activity and

cuticle structure are not differentially expressed in the whole body. On the other hand, even

though the genes enriched in cellular respiration were up-regulated in the whole body, they were

down-regulated in both ionoregulatory tissues. Taken together, combining whole-body and

tissue-specific RNA-Seq allowed combining advantages while eliminating the disadvantages of

each approach.

Previous studies on the cold acclimation response of other *Drosophila* species suggest that the

percentage of differentially expressed genes in response to acclimation decreases as the species

becomes more tolerant to cold (Parker *et al.*, 2015, 2021; MacMillan *et al.*, 2016; Enriquez &

Colinet, 2019). Even though these differences can be due to technological advancements,

acclimation conditions, or analysis methods, the current study's large-scale differential

expression aligns with this claim. In addition, the general response to cold acclimation consisted

of differential expression of genes related to translation, transcription, and metabolism. Over-

expression of translation-related genes parallels the high number of DE genes responding to

acclimation. The populations also showed a slight difference in DE gene percentage, supporting

the previous suggestion.

Exposure to low temperatures decreases membrane fluidity (Steponkus, 1984; Drobnis et al.,

1993; Koštál, 2010). Homeoviscious adaptation suggests that the organism can adjust its

membrane fluidity by increasing the proportion of unsaturated fatty acids or the ratio of short-

chain fatty acids to long-chain fatty acids (Sinensky, 1974; Koštál, 2010). Insects can adjust their

membrane composition through fatty acid metabolism to keep the membrane fluidity consistent,

which may increase cold tolerance during acclimation (reviewed in Teets & Denlinger, 2013).

Aquaporins, essential membrane proteins supporting cold tolerance in many insects (Zhao *et al.*,

2023), were suggested to regulate cold shock response by modulating water balance and fatty

acid metabolism in ticks (Wang *et al.*, 2025). In our study, the fatty acid elongation was down-

regulated in response to acclimation in both populations, preventing a decrease in the short-

chain-to-long-chain ratio. However, the differential expression analysis between the populations

suggested that the down-regulation of genes related to fatty acid elongase activity was

significantly higher in the Bangkok population than in the Kathmandu population. These results

suggest that the ancestral population prevents the elongation of short-chain fatty acids into long-

chain ones to maintain the membrane fluidity. In contrast, the acclimation response of the

derived population does not solely depend on membrane structure.

As discussed previously, cold exposure increases potassium ion concentration in the extracellular

space, resulting in membrane depolarization (Hoyle, 1953; Findsen *et al.*, 2014; reviewed in

Overgaard & MacMillan, 2017 and in Overgaard *et al.*, 2021). As acclimation increases

tolerance, adjustments are expected to be made to improve the maintenance of the water-ion

balance. Hence, the up-regulation of genes regulating ion transport mechanisms can improve the

cold tolerance phenotype. The genes significantly up-regulated in the derived population

compared to the ancestral one showed enrichment in ion transport and channel activity.

Specifically, the up-regulation of genes regulating the potassium channel activity in the

Kathmandu population indicates that the derived population takes advantage of preventing

depolarization by maintaining the potassium concentration. On the other hand, in the Malpighian

tubules, the ion transport genes were responsible for the active transmembrane transport rather

than ion-specific channels. As active transport is affected more than passive diffusion in the case

of cold exposure (Koštál *et al.*, 2004; Zachariassen *et al.*, 2004; MacMillan & Sinclair, 2011),

maintaining the ion balance in response to acclimation depends on regulating the active transport

mechanism.

Taken together, these results suggest that the ancestral population maintains the membrane

fluidity to prevent the loss of water-ion balance, whereas the derived population can maintain the

active transport in the Malpighian tubules and regulate the potassium ion concentration in the

extracellular space in response to acclimation, improving the cold tolerance. Therefore,

maintaining ion transport plays a crucial role in the adaptation of the derived Kathmandu

population to cold temperatures.

Considering the phenotypic variance within and between populations, comparative

transcriptomic analysis of the cold acclimation response also sheds light on the differences

between the phenotypes. The analysis focused on the hindgut tissue, where cellular respiration

and lipid metabolism-related genes were down-regulated in response to acclimation, whereas

cuticle development was up-regulated.

A previous study showed that prolonged exposure to cold reduces insect respiration rate (Colinet

et al., 2017). However, the same study also suggested that high tolerance can be achieved by

maintaining the energy requirement under cold stress. In line with these suggestions, even though

the respiration-related genes were down-regulated in response to acclimation in both phenotypes,

the difference between the tolerant and sensitive phenotypes was significant: sensitive lines

showed significant down-regulation of these genes compared to the tolerant lines. These

indications suggest that maintaining the respiration-related processes can be associated with fast

recovery from chill coma.

Together with respiration, lipid metabolism-related genes were significantly down-regulated in

hindguts in response to cold acclimation. As discussed above, low temperatures affect the lipid

bilayer structure of the membrane, and these impacts can be overcome by homeoviscious

adaptation. Even though the genes associated with fatty acid elongase activity were down-

regulated, no phenotype-specific difference was observed.

Previous studies showed that reabsorption in the hindgut plays a crucial role in cold tolerance of

Drosophila species (Andersen *et al.*, 2017; Andersen & Overgaard, 2020). Therefore, any

adaptation preventing or minimizing the leak in the hindgut is expected to increase tolerance to

cold. As the cuticle structure of the hindgut prevents water loss (Phillips, 1970), activation of

related pathways is expected to be observed in cold-tolerant strains. Interestingly, genes related

to cuticle development and chitin metabolism were up-regulated in the hindguts of all strains in

response to cold acclimation. However, the up-regulation significantly differed between the

phenotypes. As expected, tolerant strains showed more up-regulation of related genes than the

sensitive strains, suggesting that cuticle formation plays a vital role in the cold tolerance level of

the species.

The comparative transcriptomic analysis of the whole-body cold acclimation response also

explains the variation within the ancestral population. Differences were observed in GO terms

related to actin cytoskeleton and calcium channels between the fast and slow-recovering

ancestral strains. Genes related to both were up-regulated in response to acclimation, but the

upregulation was significantly more in the tolerant fast-recovering strains.

Actin cytoskeleton plays roles in various biological processes, including cell structure, shape,

migration, and signaling. Previous studies suggested that the actin cytoskeleton is essential in

cold tolerance (Cottam *et al*., 2006; Kim *et al*., 2006; von Heckel *et al*., 2016; Chen *et al*., 2017;

Des Marteaux *et al*., 2018; Bowman *et al*., 2018). One of these studies also showed that the

tolerant and sensitive Bangkok strains differed in actin polymerization: six genes had differential

expression in response to cold shock (Königer & Grath, 2018). On the other hand, five out of six

genes were up-regulated in both phenotypes in response to acclimation. However, they did not

show phenotype-specific differential expression even though the initial expression levels were

significantly lower in the sensitive phenotype.

In addition to the roles above, the actin cytoskeleton also acts as an anchor to ion channels and

regulates ion transport (reviewed in Denker & Barber, 2002). Previous studies showed that

together with microtubules, actin cytoskeleton dynamics affect the calcium ion influx. Calcium

ion concentration is shown to be affected by cold temperatures (Salkoff & Wyman, 1983;

Findsen *et al.*, 2016) and plays an essential role in the activation of apoptotic and necrotic

pathways (Koštál et al., 2004, 2007, 2016; Zachariassen et al., 2004; MacMillan et al., 2015,

2016). Genes related to calcium channels were also up-regulated in response to acclimation in

the ancestral population. However, the differential expression analysis between the two

phenotypes of the ancestral population showed that the calcium transport-related genes were

significantly more up-regulated in the fast-recovering phenotype than in the slow-recovering one.

These results suggest that the acclimation response of the two phenotypically distinct ancestral

pools differ in actin cytoskeleton and ion transport, focusing on calcium ions.

In this study, we showed that acclimation to mild temperatures profoundly affected the cold

tolerance of *D. ananassae* and investigated the mechanisms leading to increased tolerance. We

showed that different populations had distinct transcriptomic responses to acclimation in the

whole body and the Malpighian tubules. The differences between populations manifested

through ion transport and lipid metabolism. In addition, the cold tolerance phenotype affected the

acclimation response in the hindgut: tolerant strains could better maintain the cellular respiration

that may generate the energy required by the active transport mechanism and took advantage of

the cuticle structure that prevents water loss. Finally, we showed that the phenotypic variation

within the ancestral population resulted in differences in the acclimation response, including

actin cytoskeleton and calcium channel-related processes.

Further studies should be performed to validate these differences between the two phenotypes,

focusing on the function of the hindgut and cellular respiration. Mitochondria activity has been

described to undergo various changes under cold stess in insects, both to minimize cell damage

as well as due to a decline in mitochondrial homeostasis caused by cell injuries and also

depending on experimental conditions, what should be investigated further (Lubawy *et al.*,

2022). Hindgut reabsorption functional assays were previously used in larger insects such as

grasshoppers (Gerber *et al.*, 2021) but also successfully performed in other *Drosophila* species

(Andersen & Overgaard, 2020; Andersen *et al.*, 2017). Similarly, cellular respiration

measurements were successful in other species (Colinet *et al.*, 2017) and can be used in *D.*

ananassae. In addition to functional assays, candidate genes arising from further analyses can be

edited through CRISPR-*Cas9* technology previously established in the species (Yılmaz *et al.*,

2023) for functional validation.

Materials and Methods

Fly strains

This study used eight *D. ananassae* isofemale lines from Thailand (Bangkok, BKK) and three

isofemale lines from Nepal (Kathmandu, KATH). The lines from Bangkok were the same as used

in previous studies (Königer and Grath, 2018; Königer *et al.*, 2019). All lines were collected in

2002 (Das *et al.*, 2004). The fly strains were maintained under standard laboratory conditions

(Rearing temperature – RT: 22°C ± 1°C 14:10 hours light:dark cycle) at low density on corn-

molasses medium. Experiments were performed on flies that were expanded in large vials for

two generations under the same conditions.

Cold Acclimation Assay

Acclimation conditions were determined via preliminary tests (see Supplementary Text,

Supplementary Methods). Acclimation vials contained standard food and were kept at 10°C for

24 hours prior to the experiment to allow the food to cool down to acclimation temperature. Male

flies were pooled per population and phenotype on the day of emergence. After 24 hours, flies

were transferred to vials containing standard medium as described above. The vials were kept at

10°C for 4 days, followed by 90 minutes recovery time at rearing temperature.

RNA Extraction and Sequencing

Ten individual males per sample were snap-frozen in liquid nitrogen for whole-body samples,

and RNA was extracted using MasterPure™ RNA Purification Kit (Epicentre, Madison, WI,

USA) following the manufacturer's protocol. For hindgut and Malpighian tubule samples, tissues

from 60 males per sample were dissected, and the tissues were collected into 1.5 mL Eppendorf

tubes containing 10-20 μl 1X PBS. RNA extraction from tissues was performed using RNeasy

Mini Kit (Qiagen). RNA quantity and quality for all samples were confirmed with NanoDrop©

(ND 1000, VWR International, Radnor, PA, USA) and a bioanalyzer (Bioanalyzer 2001, Agilent

Technologies, Santa Clara, CA, USA, provided by the LMU genomics service unit). All samples

were sent for sequencing to an external facility (Novogene, Cambridge, UK), which performed

poly-A mRNA purification, reverse transcription, and high-throughput using a Novaseq 6000

sequencer (Illumina, San Diego, CA, USA) to generate 150-bp paired-end reads. On average, 88

million reads were generated per sample. An average of 73 million reads (83%) were

successfully mapped to the *D. ananassae* transcriptome (dana_r1.05_FB2016_01) using

NextGenMap (version 0.5.2) (Sedlazeck *et al.*, 2013) and the annotation of FlyBase release 1.05.

This release was used to compare the results of the present study with previously described data

on transcriptome response following cold shock (Königer and Grath, 2018). All subsequent

analyses were performed in R (version 4.2.2) (R Core Team, 2022).

Differential Gene Expression Analysis

Differentially expressed genes were called using DESeq2 (version 1.38.3) (Love *et al.*, 2014).

The general response to treatment was assessed using ~Treatment as the design formula. The

differences between other variables (populations, phenotypes, or strains) were assessed using the

design formula ~Variable + Treatment + Variable:Treatment. Significant log2 fold changes

between control and cold-acclimated samples were determined using a Wald test and a

subsequent p-value multiple testing correction according to Benjamini-Hochberg. Genes with a

corrected p-value less than 0.05 were considered as significantly differentially expressed.

Gene Ontology (GO) Enrichment Analysis

GO analysis was performed using clusterProfiler (version 4.6.2) (Wu *et al.*, 2021) implemented

in R to determine the GO terms associated with *D. melanogaster* orthologs of the differentially

expressed genes. The p-values were corrected using FDR method with a p-value cut-off of 0.01.

The significant GO terms were visualized using the genekitr package (version 1.2.5) (Liu, 2023).

Comparison with Cold Shock

Previously, Königer and Grath (2018) analyzed differential gene expression before, at 15

minutes, and 90 minutes after a cold shock in cold-sensitive and cold-tolerant strains of the

Bangkok population. The study used the same Bangkok strains used in the current study, but the

samples were not pooled for RNA extraction. Additionally, the data contained an extra time point

(15 minutes after cold shock), which we did not test in the current study. Therefore, we used

mapped cold shock reads that were kindly provided by Königer and Grath, analyzed them with

the same DESeq2 model as the cold acclimation reads, and compared the DE genes between two

phenotypes of the BKK population upon cold acclimation to the DE genes between the same

groups at 90 minutes after cold shock.

Data Availability

Gene expression data are available at GEO under the accession number GSE270239. All other data and scripts are available in the article and in its Supplementary Material.

Author Contributions

Conceptualization S.G.; investigation V.M.Y., Z.B.; writing-original draft V.M.Y., S.G.; visualization V.M.Y., S.G.; funding acquisition S.G.

Acknowledgements

Our work was supported by the Deutsche Forschungsgemeinschaft (project 271330745 to S. G.). We thank Hilde Lainer and Bianca Hölldobler for technical assistance in the laboratory, Amanda Glaser-Schmitt and John Parsch for their valuable input on the analysis of raw reads, and Annabella Königer for cold shock read counts. We extend our appreciation to all current and previous members of the Division of Evolutionary Biology, as well as the Graduate School Life Science Munich (LSM) at Ludwig-Maximilians-University Munich (LMU).

Conflicts of Interest

The authors declare no conflicting interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Andersen JL, Manenti T, Sørensen JG, MacMillan HA, Loeschcke V, Overgaard J. 2015. How to assess *Drosophila* cold tolerance: Chill coma temperature and lower lethal temperature are the best predictors of cold distribution limits. *Functional Ecology*, 29(1), 55–65. https://doi.org/10.1111/1365-2435.12310

Andersen MK, & Overgaard J. 2020. Maintenance of hindgut reabsorption during cold exposure is a key adaptation for *Drosophila* cold tolerance. *Journal of Experimental Biology*, jeb.213934. https://doi.org/10.1242/jeb.213934

Andersen MK, MacMillan HA, Donini A, Overgaard J. 2017. Cold tolerance of *Drosophila* species is tightly linked to epithelial K+ transport capacity of the Malpighian tubules and rectal pads. *Journal of Experimental Biology*, jeb.168518. https://doi.org/10.1242/jeb.168518

Bale JS, Hayward SAL. 2010. Insect overwintering in a changing climate. *Journal of Experimental Biology*, *213*(6), 980-994.

Bayley JS, Winther CB, Andersen MK, Grønkjær C, Nielsen OB, Pedersen TH, Overgaard J. 2018. Cold exposure causes cell death by depolarization-mediated Ca2+ overload in a chill-susceptible insect. *Proceedings of the National Academy of Sciences*, *115*(41), E9737-E9744.

Bowman LL, Kondrateva ES, Timofeyev MA, Yampolsky LY. 2018. Temperature gradient affects differentiation of gene expression and SNP allele frequencies in the dominant Lake Baikal zooplankton species. *Molecular Ecology*, *27*(11), 2544–2559. https://doi.org/10.1111/mec.14704

Bubliy OA, Riihimaa A, Norry FM, Loeschcke V. 2002. Variation in resistance and acclimation to low-temperature stress among three geographical strains of *Drosophila melanogaster*. *Journal of Thermal Biology*, 27(5), 337–344. https://doi.org/10.1016/S0306-4565(01)00098-5

Chen Y, Jiang T, Zhu J, Xie Y, Tan Z, Chen Y, Tang S, Hao B, Wang S, Huang J, Shen X. 2017. Transcriptome sequencing reveals potential mechanisms of diapause preparation in bivoltine silkworm *Bombyx mori* (Lepidoptera: *Bombycidae*). *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, *24*, 68–78. https://doi.org/10.1016/j.cbd.2017.07.003

Colinet H, Renault D, Roussel D. 2017. Cold acclimation allows *Drosophila* flies to maintain

mitochondrial functioning under cold stress. *Insect Biochemistry and Molecular Biology*, 80,

52–60. https://doi.org/10.1016/j.ibmb.2016.11.007

Cottam DM, Tucker JB, Rogers-Bald MM, Mackie JB, Macintyre J, Scarborough JA, Ohkura H,

Milner MJ. 2006. Non-centrosomal microtubule-organising centres in cold-treated cultured

Drosophila cells. *Cell Motility*, *63*(2), 88–100. https://doi.org/10.1002/cm.20103

Das A, Mohanty S, Stephan W. 2004. Inferring the Population Structure and Demography of

Drosophila   ananassae   From   Multilocus   Data.   *Genetics*,   168(4),   1975–1985.

https://doi.org/10.1534/genetics.104.031567

David JR, Gibert P, Moreteau B, Gilchrist GW, Huey RB. 2003. The fly that came in from the

cold: Geographic variation of recovery time from low-temperature exposure in *Drosophila*

subobscura.   *Functional   Ecology*,   17(4),   425–430.   https://doi.org/10.1046/j.1365-

2435.2003.00750.x

Denker SP, Barber DL. 2002. Ion transport proteins anchor and regulate the cytoskeleton.

Current   Opinion   in   Cell   Biology,   *14*(2),   214–220.   https://doi.org/10.1016/S0955-

0674(02)00304-6

Des Marteaux LE, Khazraeenia S, Yerushalmi GY, Donini A, Li NG, Sinclair BJ. 2018. The

effect of cold acclimation on active ion transport in cricket ionoregulatory tissues.

Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, *216*,

28–33. https://doi.org/10.1016/j.cbpa.2017.11.005

Drobnis EZ, Crowe LM, Berger T, Anchordoguy TJ, Overstreet JW, Crowe JH. 1993. Cold

shock damage is due to lipid phase transitions in cell membranes: a demonstration using

sperm as a model. *Journal of Experimental Zoology*, *265*(4), 432-437.

Enriquez T, Colinet H. 2019. Cold acclimation triggers major transcriptional changes in

Drosophila suzukii. *BMC Genomics*, 20(1), 413. https://doi.org/10.1186/s12864-019-5745-7

Findsen A, Overgaard J, Pedersen TH. 2016. Reduced L-type Ca2+ current and compromised

excitability induce loss of skeletal muscle function during acute cooling in locust. *Journal of*

Experimental Biology, *219*(15), 2340-2348.

Findsen A, Pedersen TH, Petersen AG, Nielsen OB, Overgaard J. 2014. Why do insects enter

and recover from chill coma? Low temperature and high extracellular potassium compromise

muscle function in *Locusta migratoria*. *Journal of Experimental Biology*, *217*(8), 1297-1306.

Free JB, Spencer-Booth Y. 1960. Chill-coma and cold death temperatures of *Apis mellifera*.
Entomologia Experimentalis et Applicata, 3(3), 222–230. https://doi.org/10.1111/j.1570-
7458.1960.tb00451.x

Gerber L, Overgaard J. 2018. Cold tolerance is linked to osmoregulatory function of the hindgut
in *Locusta migratoria*. *Journal of Experimental Biology*, jeb.173930.
https://doi.org/10.1242/jeb.173930

Gerber L, Kresse JC, Šimek P, Berková P, Overgaard J. 2021. Cold acclimation preserves
hindgut reabsorption capacity at low temperature in a chill-susceptible insect, *Locusta
migratoria*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative
Physiology*, *252*, 110850. https://doi.org/10.1016/j.cbpa.2020.110850

Gibert P, Moreteau B, Pétavy G, Karan D, David JR. 2001. Chill-Coma Tolerance, a Major
Climatic Adaptation Among *Drosophila* Species. *Evolution*, *55*(5), 1063–1068.
https://doi.org/10.1111/j.0014-3820.2001.tb00623.x

Hallas R, Schiffer M, Hoffmann AA. 2002. Clinal variation in *Drosophila serrata* for stress
resistance and body size. *Genetical Research*, 79(2), 141–148.
https://doi.org/10.1017/S0016672301005523

Hoedjes KM, Grath S, Posnien N, Ritchie MG, Schlötterer C, Abbott JK, Almudi I, Coronado-
Zamora M, Durmaz Mitchell E, Flatt T, Fricke C, Glaser-Schmitt A, González J, Holman L,
Kankare M, Lenhart B, Orengo DJ, Snook RR, Yılmaz VM, Yusuf L. 2024. From whole
bodies to single cells: A guide to transcriptomic approaches for ecology and evolutionary
biology. *Molecular Ecology*, e17382. https://doi.org/10.1111/mec.17382

Hoffmann AA, Watson M. 1993. Geographical Variation in the Acclimation Responses of
Drosophila to Temperature Extremes. *The American Naturalist*, 142, S93–S113.
https://doi.org/10.1086/285525

Hoffmann AA, Weeks AR. 2007. Climatic selection on genes and traits after a 100 year-old
invasion: A critical look at the temperate-tropical clines in *Drosophila melanogaster* from
eastern Australia. *Genetica*, 129(2), 133. https://doi.org/10.1007/s10709-006-9010-z

Hoffmann AA, Anderson A, Hallas R. 2002. Opposing clines for high and low temperature
resistance in *Drosophila melanogaster*. *Ecology Letters*, 5(5), 614–618.
https://doi.org/10.1046/j.1461-0248.2002.00367.x

Hoyle G. 1953. Potassium ions and insect nerve muscle. *Journal of Experimental Biology*, *30*(1),
121-135.

Kelty JD, Killian KA, Lee RE. 1996. Cold shock and rapid cold-hardening of pharate adult flesh
flies (*Sarcophaga crassipalpis*): Effects on behaviour and neuromuscular function following
eclosion. *Physiological Entomology*, 21(4), 283–288. https://doi.org/10.1111/j.1365-
3032.1996.tb00866.x

Kim M, Robich RM, Rinehart JP, Denlinger DL. 2006. Upregulation of two actin genes and
redistribution of actin during diapause and cold stress in the northern house mosquito, *Culex*
pipiens.   *Journal   of   Insect   Physiology*,   *52*(11–12),   1226–1233.
https://doi.org/10.1016/j.jinsphys.2006.09.007

Königer A, Grath S. 2018. Transcriptome Analysis Reveals Candidate Genes for Cold Tolerance
in *Drosophila ananassae*. *Genes*, 9(12), 624. https://doi.org/10.3390/genes9120624

Königer A, Arif S, Grath S. 2019. Three Quantitative Trait Loci Explain More than 60% of
Variation for Chill Coma Recovery Time in a Natural Population of *Drosophila ananassae*.
G3 Genes|Genomes|Genetics, 9(11), 3715–3725. https://doi.org/10.1534/g3.119.400453

Koštál V. 2010. Cell structural modifications in insects at low temperature. *Low temperature*
biology of insects, 116-140.

Koštál V, Korbelová J, Štětina T, Poupardin R, Colinet H, Zahradníčková H, Šimek P. 2016.
Physiological basis for low-temperature survival and storage of quiescent larvae of the fruit
fly *Drosophila melanogaster*. *Scientific Reports*, *6*(1), 32346.

Koštál V, Renault D, Mehrabianová A, Bastl J. 2007. Insect cold tolerance and repair of chill-
injury at fluctuating thermal regimes: Role of ion homeostasis. *Comparative Biochemistry*
and Physiology Part A: Molecular & Integrative Physiology, 147(1), 231–238.
https://doi.org/10.1016/j.cbpa.2006.12.033

Koštál V, Vambera J, Bastl J. 2004. On the nature of pre-freeze mortality in insects: water
balance, ion homeostasis and energy charge in the adults of *Pyrrhocoris apterus*. *Journal of*
experimental biology, *207*(9), 1509-1521.

Liu Y. 2023. genekitr: Gene Analysis Toolkit (R package version 1.2.5). https://CRAN.R-
project.org/package=genekitr

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology, 15*(12), 550.

Lubawy J, Chowański S, Adamski Z, Słocińska M. 2022. Mitochondria as a target and central
hub of energy division during cold stress in insects. *Frontiers in Zoology, 19*(1).
https://doi.org/10.1186/s12983-021-00448-3

MacMillan HA, Sinclair BJ. 2011. Mechanisms underlying insect chill-coma. *Journal of Insect
Physiology*, *57*(1), 12–20. https://doi.org/10.1016/j.jinsphys.2010.10.004

MacMillan HA, Andersen JL, Davies SA, Overgaard J. 2015. The capacity to maintain ion and
water homeostasis underlies interspecific variation in *Drosophila* cold tolerance. *Scientific
Reports*, *5*(1), 18607. https://doi.org/10.1038/srep18607

MacMillan HA, Knee JM, Dennis AB, Udaka H, Marshall KE, Merritt TJS, Sinclair BJ. 2016.
Cold acclimation wholly reorganizes the *Drosophila melanogaster* transcriptome and
metabolome. *Scientific Reports*, *6*(1), 28999. https://doi.org/10.1038/srep28999

MacMillan HA, Yerushalmi GY, Jonusaite S, Kelly SP, Donini A. 2017. Thermal acclimation
mitigates cold-induced paracellular leak from the *Drosophila* gut. *Scientific Reports*, *7*(1),
Article 1. https://doi.org/10.1038/s41598-017-08926-7

Overgaard J, MacMillan HA. 2017. The Integrative Physiology of Insect Chill Tolerance. *Annual
Review of Physiology*, *79*(1), 187–208. https://doi.org/10.1146/annurev-physiol-022516-
034142

Overgaard J, Gerber L, Andersen MK. 2021. Osmoregulatory capacity at low temperature is
critical for insect cold tolerance. *Current Opinion in Insect Science*, 47, 38–45.
https://doi.org/10.1016/j.cois.2021.02.015

Parker DJ, Envall T, Ritchie MG, Kankare M. 2021. Sex-specific responses to cold in a very
cold-tolerant, northern *Drosophila* species. *Heredity*, 126(4), 695–705.
https://doi.org/10.1038/s41437-020-00398-2

Parker DJ, Vesala L, Ritchie MG, Laiho A, Hoikkala A, Kankare M. 2015. How consistent are
the transcriptome changes associated with cold acclimation in two species of the *Drosophila
virilis* group? *Heredity*, 115(1), 13–21. https://doi.org/10.1038/hdy.2015.6

Phillips JE. 1970. Apparent Transport of Water by Insect Excretory Systems. *American
Zoologist*, 10(3), 413–436. https://doi.org/10.1093/icb/10.3.413

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for
Statistical Computing. https://www.R-project.org/

Reinhold JM, Lazzari CR, Lahondère C. 2018. Effects of the environmental temperature on

Aedes aegypti and *Aedes albopictus* mosquitoes: a review. *Insects*, *9*(4), 158.

Ring RA. 1982. Freezing-tolerant insects with low supercooling points. *Comparative*

Biochemistry and Physiology Part A: Physiology, *73*(4), 605-612.

Salkoff LB, Wyman RJ. 1983. Ion currents in *Drosophila* flight muscles. *The Journal of*

physiology, *337*(1), 687-709.

Sedlazeck FJ, Rescheneder P, Von Haeseler A. 2013. NextGenMap: Fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics*, 29(21), 2790–2791.

https://doi.org/10.1093/bioinformatics/btt468

Sinensky M. 1974. Homeoviscous Adaptation—A Homeostatic Process that Regulates the

Viscosity of Membrane Lipids in *Escherichia coli*. *Proceedings of the National Academy of*

Sciences, 71(2), 522–525. https://doi.org/10.1073/pnas.71.2.522

Sisodia S, Singh BN. 2010. Resistance to environmental stress in *Drosophila ananassae*:

Latitudinal variation and adaptation among populations. *Journal of Evolutionary Biology*,

23(9), 1979–1988. https://doi.org/10.1111/j.1420-9101.2010.02061.x

Steponkus PL. 1984. Role of the plasma membrane in freezing injury and cold

acclimation. *Annual Review of Plant Physiology*, *35*(1), 543-584.

Teets NM, Denlinger DL. 2013. Physiological mechanisms of seasonal and rapid cold-hardening

in insects. *Physiological Entomology*, 38(2), 105–116. https://doi.org/10.1111/phen.12019

Teets NM, Kawarasaki Y, Lee RE, Denlinger DL. 2012. Energetic consequences of repeated and

prolonged dehydration in the Antarctic midge, *Belgica antarctica*. *Journal of Insect*

Physiology, 58(4), 498–505. https://doi.org/10.1016/j.jinsphys.2011.11.011

Udaka H, Ueda C, Goto SG. 2010. Survival rate and expression of *Heat-shock protein 70* and

Frost genes after temperature stress in *Drosophila melanogaster* lines that are selected for

recovery time from temperature coma. *Journal of Insect Physiology*, *56*(12), 1889-1894.

von Heckel K, Stephan W, Hutter S. 2016. Canalization of gene expression is a major signature

of regulatory cold adaptation in temperate *Drosophila melanogaster*. *BMC Genomics*, 17(1),

574. https://doi.org/10.1186/s12864-016-2866-0

Wang H, Bai R, Pei T, et al. Aquaporins modulate the cold response of Haemaphysalis

longicornis via changes in gene and protein expression of fatty acids. *Parasites & Vectors*.

2025;18(1):70. https://doi.org/10.1186/s13071-025-06718-x

Watson MJO, Hoffmann AA. 1996. Acclimation, cross-generation effects, and the response to
selection for increased cold resistance in *Drosophila*. *Evolution*, 50(3), 1182–1192.
https://doi.org/10.1111/j.1558-5646.1996.tb02359.x

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Liu S, Bo X,
Yu G. 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The
Innovation, 2*(3), 100141.

Yerushalmi GY, Misyura L, MacMillan HA, Donini A. 2018. Functional plasticity of the gut and
the Malpighian tubules underlies cold acclimation and mitigates cold-induced hyperkalemia
in *Drosophila melanogaster*. *Journal of Experimental Biology*, jeb.174904.
https://doi.org/10.1242/jeb.174904

Yılmaz VM, Ramnarine TJS, Königer A, Mussgnug S, Grath S. 2023. Tropical super flies:
Integrating *Cas9* into *Drosophila ananassae* and its phenotypic effects. *Journal of Insect
Physiology*, 147, 104516. https://doi.org/10.1016/j.jinsphys.2023.104516

Zachariassen KE, Kristiansen E, Pedersen SA. 2004. Inorganic ions in cold-hardiness.
Cryobiology, *48*(2), 126-133

Zhao C, Liu Z, Liu Y, Zhan Y. 2023. Identification and characterization of cold-responsive
aquaporins from the larvae of a crambid pest *Agriphila aeneociliella* (Eversmann)
(Lepidoptera: Crambidae). *PeerJ*. 11:e16403. Published 2023 Nov 13.
doi:10.7717/peerj.16403