Article Type – Discovery

Navigating the Cold: Integrative Transcriptome Sequencing Approach Reveals Ionoregulatory and Whole-Body Responses to Cold Acclimation in *Drosophila ananassae*

Vera Miyase Yılmaz[1], Zhihui Bao[1], Sonja Grath[1*]

[1]Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany

[*]Corresponding author: grath@bio.lmu.de

Running title: Adaptation to cold in *Drosophila*

Abstract

Understanding how species adapt to changing environments is a major goal in evolutionary biology and can elucidate the impact of climate change. Climate imposes inevitable effects on the geographical distribution of insects as their body temperature primarily depends on the environment. The vinegar fly *Drosophila ananassae* expanded from its tropical ancestral range to more temperate regions, which requires adaptation to colder temperatures. Transcriptome and genome-wide association studies focusing on the ancestral-range population identified the targets of selection related to ionoregulatory tissues. However, how cosmopolitan *D. ananassae* adapted to colder environments, where low temperatures last longer, is still unknown. Here, we present a study on the effect of long-term cold exposure on *D. ananassae*, examining the gene expression variation in the whole body and the ionoregulatory tissues, namely the hindgut and the Malpighian tubule. To elucidate molecular mechanisms of cold adaptation during species expansion, we included cold-tolerant and cold-sensitive strains from the ancestral species range and cold-tolerant strains from the derived species range. We show that cold acclimation improves cold tolerance and results in differential expression of more than half of the transcriptome in the ionoregulatory tissues and the whole body. Notably, we provide complementary insight into molecular processes at four levels: strains, populations, phenotypes, and tissues. By determining the biochemical pathways of phenotypic plasticity underlying cold tolerance, our results enhance our understanding of how environmental changes affect thermal adaptation in natural populations.

Keywords: thermoregulation, ectothermy, metabolic cold adaptation, gene expression, phenotypic plasticity, ionoregulatory evolution, acclimation

Introduction

Temperature is a major factor influencing how species adapt to changing environments and

colonize new ecosystems. How animals generally cope with thermal stressors and which genes are

specifically responsible for cold adaptation has been a topic of intense research for decades in

molecular and evolutionary biology. Recent developments in high-throughput sequencing methods

allow insights into molecular mechanisms at the whole transcriptome and genome level, but

integrating the complementary information of various experimental designs remains challenging.

For example, studies on small insects often rely on either systemic or tissue-specific methodologies

or single phenotypes, resulting in incomplete conclusions (Andersen *et al.*, 2015; Hoedjes *et al.*,

2024). Therefore, integrating different experimental designs and phenotypes to study

transcriptional responses to cold is critical to uncovering mechanisms underlying physiological

adaptation. Elucidating the mechanisms that underlie adaptation to changing environments in

natural populations and integrating different approaches can help us understand which species

prove to be successful in the future and have potential applications for conservation, agriculture,

and pest management.

Many insects evolved and diversified in warm climates (Bale & Hayward, 2010). Nevertheless,

insects have developed various strategies to cope with thermal stress and to successfully expand

their thermal and habitat ranges (Overgaard & MacMillan, 2017). Some insects even tolerate

freezing and remain active, such as the Antarctic midge *Belgica antarctica* (Teets *et al.*, 2012) or

the alpine beetle *Pytho deplanatus* (Ring, 1982). However, a wide range of insect species suffer

from chilling injuries and eventually die when they are exposed to cold. This chill-susceptibility

has been reported for several orders of insects (Overgaard & MacMillan, 2017). Chill-susceptible

species include some of great economic and ecological importance, such as the honeybee *Apis*

mellifera (Free & Spencer-Booth, 1960), but also agricultural pests, for instance, the migratory

locust *Locusta migratoria* (Bayley *et al.*, 2018) and disease vectors such as mosquitoes (Reinhold

et al., 2018).

The physiology of cold stress resistance in insects is highly complex. When insects are exposed to

cold, their ion and water homeostasis is disrupted (MacMillan *et al.*, 2015). Additionally,

transcriptomic and metabolic profiles are changed (MacMillan *et al.*, 2016). Along with these

systemic changes, chilling injuries affect the body at the cell level as they cause phase changes in

the cell membrane, resulting in a stiff membrane with decreased permeability (Steponkus, 1984;

Koštál, 2010; Drobnis *et al.*, 1993). Further, cold exposure results in an imbalance of passive

diffusion and active ion transport (MacMillan & Sinclair, 2011; Koštál *et al.*, 2004; Zachariassen

et al., 2004). An excess of potassium ions in the extracellular space leads to depolarization of the

membrane (Findsen *et al.*, 2014; Hoyle, 1953), eventually activating the voltage-gated calcium

channels (Salkoff & Wyman, 1983; Findsen *et al.*, 2016). The cellular influx of calcium ions

through voltage-gated channels activates apoptotic and necrotic pathways, leading to chill injury

(Koštál *et al.*, 2004, 2007, 2016; Zachariassen *et al.*, 2004; MacMillan *et al.*, 2015, 2016).

Consequently, the regulation of ion and water homeostasis has been identified as a key

physiological feature for cold tolerance in insects with specifically the ionoregulatory tissues being

essential for maintaining the water-ion balance in an insect body (Andersen *et al.*, 2017; Andersen

& Overgaard, 2020; Des Marteaux *et al.*, 2018; Gerber & Overgaard, 2018; MacMillan *et al.*,

2015, 2016, 2017; Yerushalmi *et al.*, 2018). Despite several in-depth studies aiming to resolve this

complexity, it remains challenging to identify and functionally characterize specific genes and

genetic elements that contribute to the diversity of cold stress-resistant phenotypes (Udaka *et al.*,

2010; von Heckel *et al.*, 2016; MacMillan *et al.*, 2016). With their worldwide distribution,

Drosophilids are excellent models for studying these questions.

Flies from the genus *Drosophila* suffer from impaired synaptic transmission and muscular failure

when they are exposed to cold; in other words, they lose their neuromuscular coordination and

enter a comatose condition referred to as chill coma in which they are not able to move, feed, mate

or escape predators (Kelty *et al.*, 1996; MacMillan & Sinclair, 2011; Findsen *et al.*, 2014).

However, flies can recover from such a coma given that time and temperature of exposure to cold

do not exceed certain thresholds. The time the fly needs for recovery from a cold-induced coma

(chill coma recovery time, CCRT) is widely used to assess the cold hardiness of insects in the

laboratory and serves as a predictor for the climatic condition at the geographical sampling site of

a particular species (Gibert *et al.*, 2001). Further, CCRT follows latitudinal clines within

Drosophila species in Australia (Hoffmann *et al.*, 2002; Hoffmann & Weeks, 2007; Hallas *et al.*,

2002), Asia (Sisodia & Singh, 2010), Africa and Europe (David *et al.*, 2003). Cold tolerance in

Drosophilids is polygenic. Despite this complexity, variation in gene expression of only a few loci

is associated with cold tolerance in subtropical populations of *Drosophila ananassae* (Königer &

Grath, 2018), and variation of just three genomic regions explained more than 60% of the

phenotypic variation between cold-tolerant and cold-sensitive strains (Königer *et al.*, 2019).

Recently, we provided Cas9-integrated fly strains that facilitate the use of the CRISPR-Cas9

system in this species and allow functional characterization of individual candidate loci (Yılmaz

et al., 2023). In sum, *D. ananassae* has been established as a promising model for studying the

genetic basis of cold tolerance.

Previous studies mostly neglected the potential of integrating two specific aspects to better inform

about cold tolerance in insects. First, how insects respond to low temperatures can be plastic: the

ability to tolerate cold and the genes responsible for the survival can change depending on prior

exposure to cold. Such acclimation has been shown to alleviate the adverse effects of cold and can

increase cold tolerance (Bubliy *et al.*, 2002; Enriquez & Colinet, 2019; Hoffmann & Watson, 1993;

MacMillan *et al.*, 2016, 2017; Watson & Hoffmann, 1996; Yılmaz *et al.*, 2023). Cold acclimation

triggers substantial changes in the transcriptional response to cold in *Drosophila suzukii* (Enriquez

& Colinet 2019). It reorganizes the transcriptome and metabolome of whole animals (MacMillan

et al., 2016) and the gut transcriptome in *Drosophila melanogaster* (MacMillan *et al.*, 2017).

Second, conclusions on potentially ecologically relevant adaptations should not rely on a single

phenotype (Andersen *et al.*, 2015). Consequently, incorporating distinct cold tolerance phenotypes

and systemic and ionoregulatory tissue-specific transcriptome data will allow pinpointing cold-

related physiological adaptations.

In this study, we use iso-female lines collected from a population of ancestral origin in Bangkok,

Thailand (from here on mentioned as ancestral population) and a derived population originating

from Kathmandu, Nepal (Das *et al.*, 2004). We examined the transcriptional response to cold

acclimation in both the ancestral and the derived populations to shed light on the genetic

mechanisms underlying cold tolerance. In addition, we integrated the original results of this study

with previously gained insight into the distinct phenotype of cold shock. To this end, we contrasted

the transcriptional response to cold at four levels: (1) population (ancestral vs. derived), (2) strain

(cold-sensitive vs. cold-tolerant), (3) tissue (whole-body vs. ionoregulatory tissues, Malpighian

tubules and hindgut), and (4) phenotype (cold acclimation vs. cold shock). We find empirical

evidence for the influence of population and tissue on gene expression variation upon cold stress.

Using an RNA sequencing approach that combined systemic with tissue-specific transcriptomes,

we show that cold acclimation improves cold tolerance and results in more than 50% of the

transcriptome being differentially expressed in the whole body the ionoregulatory tissues.

Our results demonstrate that an RNA sequencing strategy combining a systemic overview with

tissue-specific data can be a cost-effective and informative way to elucidate genes that have major

effects on a crucial adaptation-related trait, namely, thermoregulation.

Results

Complementary elucidation of transcriptional response to cold acclimation with systemic whole body and tissue-specific data

Flies from ancestral and derived populations were previously described as having different sensitivity to cold shock as determined by CCRT (Königer & Grath, 2018). Additionally, preliminary tests revealed that acclimation to cold improves the tolerance and decreases the chill coma recovery time (see Supplementary Text). To receive an informative overview of the transcriptional response to cold acclimation in *D. ananassae*, we combined high-throughput RNA sequencing of the whole body (wb), as well as of two ionoregulatory tissues, Malpighian tubules (Mt) and hindguts (hg). We obtained RNA sequencing data from male flies upon 4-day acclimation at 10°C. In the whole body and hindgut samples, 54% of all expressed genes were differentially expressed in response to cold acclimation, and the percentage of DE genes in the Malpighian tubules was 50% (Table 1). The principal component analysis revealed that the samples clustered according to the tissue RNA was extracted from, whereas no separation according to the treatment was observed (Figure 1A). The first principal component explained 68% of the variance, separating the ionoregulatory tissues from the whole-body samples. The second principal component explained 19% of the variance, separating the two ionoregulatory tissues. Though pairwise comparisons revealed that 2952 differentially expressed genes were shared between sample types, a considerable number of genes (ranging from 505 genes that were explicitly up-regulated in Malpighian tubules to 1436 genes that were down-regulated explicitly in the whole body) were specifically up- or down-regulated only in the individual tissues (Figure 1B-C). Notably, these genes were enriched in distinct GO categories (Supplementary Table 3-4). These

154 results suggest that the combination of systemic and informed tissue-specific RNA sequencing

155 data allows for complementary conclusions on cold tolerance in *Drosophila*.

156 **Table 1:** Number of differentially expressed genes in different sample types regarding the general,
157 population-specific, phenotype-specific, and line-specific response to cold acclimation

	total expressed genes	up regulated	down regulated	total DE
General response to acclimation				
Whole body	14304	3825 (27%)	3930 (27%)	7755 (54%)
Malpighian tubule	12189	3280 (27%)	2874 (24%)	6154 (50%)
Hindgut	13272	3715 (28%)	3480 (26%)	7195 (54%)
Population-specific response				
<u>Whole body</u>				
BKK	14300	3618 (25%)	3668 (26%)	7286 (51%)
KATH	14300	3414 (24%)	3497 (24%)	6911 (48%)
KATH vs BKK*	13191	1181 (9%)	827 (6.3%)	2008 (15%)
<u>Malpighian tubule</u>				
BKK	11721	3077 (26%)	2606 (22%)	5683 (48%)
KATH	11191	2288 (20%)	2174 (19%)	4462 (40%)
KATH vs BKK*	9125	124 (1.4%)	92 (1%)	216 (2.4%)
Phenotype-specific response				
<u>Hindgut</u>				
Fast	12821	3203 (25%)	3570 (28%)	6773 (53%)
Slow	12547	2907 (23%)	2784 (22%)	5691 (45%)
Fast vs Slow*	12547	182 (1.5%)	246 (2%)	428 (3.4)
Line-specific response**				
<u>Whole body</u>				
FastBKK	14300	3333 (23%)	3434 (24%)	6767 (47%)
SlowBKK	14300	3091 (22%)	3300 (23%)	6391 (45%)
FastBKK vs SlowBKK*	13468	508 (3.8%)	472 (3.5%)	980.3%)

158 * For these comparisons, the columns show the number of genes that are significantly differentially
159 expressed between the compared groups. Up and down-regulated columns indicate direction regarding the
160 effect of acclimation on the former group compared to the latter group.

161 ** Only iso-female lines of the ancestral population are taken into consideration.

Figure 1: Transcriptomic response to cold acclimation in ionoregulatory tissues and the whole body. A) Principal component analysis. The samples are shown according to the tissue (color) and treatment (shape). Whole body (Wb), Malpighian tubule (Mt), and hindgut (Hg) tissues are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Venn diagram showing the number of DE genes specific to or shared among sample types: Whole body, Malpighian tubule and Hindgut samples exhibited both shared, as well as exclusive differentially expressed genes. C) Upset plot showing differentially expressed genes and their overlap between the three different sample types. Black, horizontal bars indicate the total number of differentially expressed genes that were significantly up- or down-regulated in a sample in response to cold acclimation. A black circle indicates that a sample and direction (up/down) combination is included in an intersection class. Intersection classes with a single circle are comprised of genes significantly differentially expressed in a single tissue and direction. Circles connected by a line indicate intersection classes comprised of multiple tissue and direction combinations. Vertical bars correspond to the number of genes in each intersection class, colors represent the respective sections of the Venn diagram.

Strong systemic transcriptional response to cold acclimation

We performed RNA sequencing on whole-body samples to obtain a systemic overview of the transcriptional response to cold acclimation in *Drosophila ananassae*. This approach has the unique advantage of being both cost-efficient and unbiased in terms of preconditions on

transcription. Samples of acclimation and control treatments were clustered according to the

populations of origin (Figure 2A). The first principal component explained 58% of the variance.

It separated cold-acclimated samples from control samples, while the second principal component

explained 14% of the variance and separated the derived population from the ancestral population.

Within the ancestral population, the samples also clustered according to their CCRT phenotype

(Figure 2B). We addressed three questions with this data: (1) What is the general, systemic

transcriptional response to cold acclimation? (2) How does transcriptional response differ between

ancestral and derived populations? (3) How does transcriptional response differ between cold-

tolerant and cold-sensitive strains in the ancestral population?

More than half of the transcriptome (54%) was differentially expressed between control and

treatment, with 3825 genes significantly up-regulated and 3930 genes significantly down-

regulated (Table 1). Gene ontology enrichment analysis revealed that up-regulated genes were

enriched in biological processes related to translation and cell junction organization (Figure 2C,

Supplementary Table 5). On the contrary, down-regulated genes were enriched in metabolic

processes (Figure 2D, Supplementary Table 5). These results suggest a solid response to cold

acclimation in the whole body, and the transcriptional response of populations differs not only in

the presence of cold treatment but also at the basal level.

Figure 2: Transcriptomic response to cold acclimation in the whole body. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Heatmap for the sample pools. Sample clusters are shown on the left and top of the heatmap. The similarity matrix indicates high (red) and low (blue) similarity of samples. Gene ontology enrichment analysis results for genes C) up-

regulated and D) down-regulated in the whole body in response to cold acclimation. Biological processes
(red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate
panels.

Systemic differences between populations are mediated by genes involved in detoxification
and maintaining ion balance

To characterize the differences between populations, we contrasted the transcriptional response of

ancestral and derived populations. A total of 14300 genes were expressed in each population (Table

1). In the ancestral population, 51% of these genes were differentially expressed upon cold

acclimation, while 3618 genes were significantly up- and 3668 down-regulated, respectively. In

comparison, 48% of these genes were differentially expressed in the derived population, while

3414 genes were significantly up-regulated, and 3497 genes were significantly down-regulated.

Although the populations shared over half of the differentially expressed genes (Supplementary

Figure 2A), 2008 genes showed significant differential expression between populations in

response to cold acclimation. The GO analysis suggested that the genes with differential expression

between the populations were mainly located in muscle tissues and enriched in cilium movement

and various metabolic processes, including fatty acid metabolism (Supplementary Figure 2B).

Notably, these genes were also enriched in molecular functions related to ion binding, ion channel,

and oxidoreductase activity (Supplementary Table 6). Overall, these results suggest the difference

between the acclimation response of populations ground on ion transport and detoxification.

The actin cytoskeleton differentiates the acclimation response of tolerant and sensitive
strains in the ancestral population

To address the question of how the transcriptional response differs within the ancestral population

between cold-tolerant and cold-sensitive strains, we compared gene expression upon cold

acclimation in both types of strains. For the cold-tolerant strains, 3333 genes were significantly

up-regulated, and 3434 genes were significantly down-regulated (Table 1). For the sensitive

strains, 3091 genes were significantly up-regulated, and 3300 genes were significantly down-

regulated. However, 980 genes were differentially expressed between cold-sensitive and cold-

tolerant strains. GO enrichment analysis revealed significant enrichment of these genes in various

biological processes related to the actin cytoskeleton. (Supplementary Table 7)

Adaptive transcriptional response is evident in the renal tissues

Malpighian tubules (Mt) are the renal equivalent of mammals in insects and contribute to

maintaining homeostasis, which can be disrupted by cold exposure. Therefore, we investigated the

transcriptional response of Mt to characterize the impact of cold acclimation on renal tissues in *D.*

ananassae. Like the whole-body samples, RNA samples extracted from Mt of acclimated and

control flies were clustered according to the populations of origin, suggesting that the populations

are distinct in their response to acclimation and at the basal level of gene expression (Figure 3A).

Principal component analysis indicated that 37% of the variation can be explained by the first

principal component, which separated cold-acclimated and control samples, and 14% of the

variation can be explained by the second principal component, which separated the ancestral and

derived populations. Therefore, we investigated the general transcriptional response of Mt to cold

acclimation and the population-specific response.

In the Mt samples, 3280 genes were up-regulated, and 2874 were down-regulated (Table 1).

Similar to the whole-body samples, up-regulated genes of Mt were enriched in GO terms related

to translation and transcription (Supplementary Figure 3A, Supplementary Table 8), and the down-

regulated genes were enriched in processes related to fatty acid degradation (Supplementary Figure

3B, Supplementary Table 8). However, contrary to the whole-body samples, the down-regulated

genes of the Mt were also enriched in chromatin remodeling. Additionally, a major part of the

down-regulated genes was located in the mitochondrion and, more specifically, the respiratory

chain complex.

Figure 3: Transcriptomic response to cold acclimation in the Malpighian tubules. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. B) Venn diagram showing the number of DE genes specific to or shared between populations: Bangkok and Kathmandu populations exhibited both shared, as well as exclusive differentially expressed genes in response to acclimation. Gene ontology (GO) enrichment analysis results for genes C) up-regulated and D) down-regulated in the Malpighian tubules of Kathmandu population compared to Bangkok population in response to cold acclimation. Biological processes (red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate panels.

Differences in renal tissues of populations are mediated by genes involved in ion transport and lipid metabolism

In order to understand the differences in metabolic processes involved in cold adaptation between the ancestral and derived populations, we compared the transcriptional response to cold acclimation in flies from the Bangkok and Kathmandu populations. In the ancestral population, 11721 genes were expressed, with 3077 up-regulated and 2606 down-regulated (Figure 3B, Table 1). In the derived population, 11191 genes were expressed, with 2288 up-regulated and 2174 down-regulated. Among these DE genes, 216 genes showed differential expression between the populations. We conducted a GO analysis to explore the functions of these genes and found GO terms related to transposition. To enhance the analysis, we performed separate analyses for gene sets: genes up regulated in the derived population compared to the ancestral one, and *vice versa*. The analysis showed that genes up regulated in the Kathmandu population compared to Bangkok were enriched in molecular functions related to ion transport (Figure 3C, Supplementary Table 9). Conversely, genes up regulated in the ancestral population compared to the derived one were enriched in processes and functions related to transposition (Figure 3D, Supplementary Table 9). Additionally, this analysis revealed an enrichment in molecular functions related to fatty acid ligase activity.

Hindgut responds to cold acclimation with respect to phenotype, but not to population

Along with the Malpighian tubules, hindguts (hg) are essential in regulating ion and water balance.

Therefore, we investigated the tissue-specific transcriptional response to cold acclimation in the

hindgut of *D. ananassae* flies to elucidate molecular mechanisms of cold adaptation in insects.

Initial PCA revealed two clusters and an outlier sample (Figure 4A). When the outlier was removed

to increase the resolution, PC1 explained 48% of the variance, and PC2 explained 12%, separating

the treatments and phenotypes, respectively (Supplementary Figure 4A). The acclimated samples

clustered according to their cold tolerance phenotype, in other words, according to their recovery

time from chill coma. However, as opposed to the whole body and MT, the control samples showed

no apparent separation, which suggests that the strains do not differ at their basal gene expression

level according to their cold tolerance phenotype or population of origin. Therefore, we addressed

two questions with this data: (1) What is the general response to cold acclimation in the hindguts?

(2) How does transcriptional response differ between cold-tolerant and cold-sensitive strains?

Figure 4: Transcriptomic response to cold acclimation in the hindguts. A) Principal component analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. Gene ontology (GO) enrichment analysis results for genes B) up-regulated and C) down-regulated in the hindguts of tolerant flies (fast phenotype) compared to sensitive flies (slow phenotype) in response to cold acclimation.

Biological processes (red bars), cellular components (blue bars), and molecular functions (yellow bars) are
shown in separate panels.

In the hindgut samples, we observed a significant up-regulation of 3835 genes and a down-

regulation of 3483 genes in response to cold acclimation (Table 1). Like the whole body and

Malpighian tubules, the up-regulated genes in the hindguts were enriched in GO terms related to

translation and transcription (Supplementary Figure 4B, Supplementary Table 10). In contrast, the

down-regulated genes showed enrichment in biological processes related to cellular respiration

and lipid metabolism, with many of these genes located in the mitochondrial membrane

(Supplementary Figure 4C, Supplementary Table 10).

Cellular respiration and chitin metabolism play roles in the acclimation response of tolerant
and sensitive strains in the hindgut

We characterized the transcriptional response to cold acclimation in cold-tolerant and cold-

sensitive strains with differential gene expression analysis. In the cold-sensitive strains, 3046 genes

were significantly up-regulated, and 2886 genes were significantly down-regulated in response to

acclimation (Table 1). On the other hand, in the cold-tolerant strains, 3401 genes were significantly

up-regulated, and 3224 genes were significantly down-regulated in response to cold acclimation.

Between the tolerant and sensitive strains, 330 genes were differentially expressed. GO enrichment

analysis revealed that these genes were enriched in biological processes related to translation and

ribosome biogenesis and in molecular functions related to chitin binding. We separately

investigated the genes up-regulated in the cold-tolerant strains compared to the sensitive strains

and *vice versa*. Genes significantly up-regulated in the cold-tolerant strains compared to the cold-

sensitive strains were enriched in GO terms related to cuticle development, respiration, and

metabolism (Figure 4B, Supplementary Table 11). Additionally, the products of these genes were

localized mainly to the membrane and extracellular matrix. Genes significantly up-regulated in the

cold-sensitive strains compared to the cold-tolerant strains were enriched in GO terms related to

translation (Figure 4C, Supplementary Table 11).

The network analysis provides an overview of the cold acclimation response

In addition to differential expression and gene ontology analyses, we conducted a weighted gene

co-expression network analysis (WGCNA) to uncover the pathway networks crucial to the cold

acclimation response in the whole body, Malpighian tubules, and hindguts of *D. ananassae*. The

general response to acclimation in the whole body was found to be centered around signaling

pathways and nervous system development (Supplementary Figure 5A), as per WGCNA, while

the populations showed variations in lipid and amino acid metabolism (Supplementary Figure 5B-

C). In the Malpighian tubules, the WGCNA revealed pathways involved in RNA metabolism and

WNT signaling (Supplementary Figure 5D-E). Conversely, the analysis of hindgut samples

unveiled carbohydrate metabolism-related pathways (Supplementary Figure 5F). Notably, no

population- or phenotype-specific differences were observed in the ionoregulatory tissues.

Distinct transcriptional response to cold acclimation and cold shock

To receive a comprehensive overview of cold tolerance measured by distinct phenotypes, we

contrasted our previous results of the transcriptional response to cold shock as measured by chill

coma recovery time upon cold shock (Königer & Grath, 2018) with the transcriptome data from

this study. To this end, we considered the genes differentially expressed in the cold-tolerant and

cold-sensitive strains of the ancestral population from Bangkok, Thailand, in two conditions: (1)

cold shock and (2) cold acclimation.

Our findings reveal the intricate nature of gene expression in response to cold shock and cold

acclimation. We observed that more genes are differentially expressed in response to cold

acclimation than cold shock (Supplementary Figure 6), indicating a more complex regulatory

network in the former. To understand these responses, we investigated the up- and down-regulated

genes in both treatments and performed a GO enrichment analysis.

The genes up-regulated in response to both treatments were enriched in actin cytoskeleton-related

GO terms (Supplementary Table 12). In contrast, the down-regulated ones showed enrichment in

various metabolic processes such as organic acid, lipid, and carbohydrate metabolic processes

(Supplementary Table 12). We also investigated the genes with opposite expression (i.e., up-

regulated in one treatment but down-regulated in the other) in response to treatments. Among the

1932 genes that showed differential expression in both studies, 390 genes had opposite directions.

The genes up-regulated upon cold shock but down-regulated upon acclimation were enriched in

the immune response (Supplementary Table 13). On the other hand, the genes that were up-

regulated in response to acclimation but down-regulated after cold shock showed enrichment in

biological processes related to transcription and translation (Supplementary Table 13).

Finally, we investigated the genes that show differential expression in response to only one of the

treatments. Over 5000 genes were up- or down-regulated upon acclimation, but only 1070 genes

were uniquely expressed after cold shock. The genes differentially expressed only upon the cold

shock treatment were enriched in protein folding, actin cytoskeleton organization, replication, and

glycosylation (Supplementary Table 14). The genes that were differentially expressed upon

acclimation showed enrichment in translation and transcription as well as amino acid metabolism

and copulation (Supplementary Table 14).

Discussion

Understanding the differences between the ancestral and derived populations and the variance within the ancestral Bangkok population sheds light on the adaptive processes leading to increased cold tolerance of the species. The two populations were tested for their acclimation response. Both populations showed improved cold tolerance upon acclimating to mild temperatures as expected, but the underlying mechanisms were unknown. Comparative transcriptomics using the whole-body samples provided the systemic overview to understand the differences in the acclimation response between the two populations and within the ancestral population. Additionally, the tissue-specific response to acclimation provided further insight into population and phenotypic differences.

In a complex phenotype, such as cold response, transcriptomic analysis of a whole body might lead to missing information (reviewed in Hoedjes *et al.*, 2024). However, tissue-specific RNA-Seq can unmask details otherwise lost in the systemic approach. Both whole-body and tissue-specific approaches were used in this study to increase the strength of the comparison of populations, phenotypes, and strains. The DE genes in the Malpighian tubule samples enriched in hypoxia response and HSP binding activity are not differentially expressed in the whole-body samples. Similarly, the DE genes in the hindgut samples enriched in sodium channel activity and cuticle structure are not differentially expressed in the whole body. On the other hand, even though the genes enriched in cellular respiration were up-regulated in the whole body, they were down-regulated in both ionoregulatory tissues. Taken together, combining whole-body and tissue-specific RNA-Seq allowed combining advantages while eliminating the disadvantages of each approach.

Previous studies on the cold acclimation response of other *Drosophila* species suggest that the

percentage of differentially expressed genes in response to acclimation decreases as the species

becomes more tolerant to cold (Parker *et al.*, 2015, 2021; MacMillan *et al.*, 2016; Enriquez &

Colinet, 2019). Even though these differences can be due to technological advancements,

acclimation conditions, or analysis methods, the current study's large-scale differential expression

aligns with this claim. In addition, the general response to cold acclimation consisted of differential

expression of genes related to translation, transcription, and metabolism. Over-expression of

translation-related genes parallels the high number of DE genes responding to acclimation. The

populations also showed a slight difference in DE gene percentage, supporting the previous

suggestion.

Exposure to low temperatures decreases membrane fluidity (Steponkus, 1984; Drobnis et al., 1993;

Koštál, 2010). Homeoviscious adaptation suggests that the organism can adjust its membrane

fluidity by increasing the proportion of unsaturated fatty acids or the ratio of short-chain fatty acids

to long-chain fatty acids (Sinensky, 1974; Koštál, 2010). Insects can adjust their membrane

composition through fatty acid metabolism to keep the membrane fluidity consistent, which may

increase cold tolerance during acclimation (reviewed in Teets & Denlinger, 2013). The fatty acid

elongation was down-regulated in response to acclimation in both populations, preventing a

decrease in the short-chain-to-long-chain ratio. However, the differential expression analysis

between the populations suggested that the down-regulation of genes related to fatty acid elongase

activity was significantly higher in the Bangkok population than in the Kathmandu population.

These results suggest that the ancestral population prevents the elongation of short-chain fatty

acids into long-chain ones to maintain the membrane fluidity. In contrast, the acclimation response

of the derived population does not solely depend on membrane structure.

As discussed previously, cold exposure increases potassium ion concentration in the extracellular

space, resulting in membrane depolarization (Hoyle, 1953; Findsen *et al.*, 2014; reviewed in

Overgaard & MacMillan, 2017 and in Overgaard *et. al.*, 2021). As acclimation increases tolerance,

adjustments are expected to be made to improve the maintenance of the water-ion balance. Hence,

the up-regulation of genes regulating ion transport mechanisms can improve the cold tolerance

phenotype. The genes significantly up-regulated in the derived population compared to the

ancestral one showed enrichment in ion transport and channel activity. Specifically, the up-

regulation of genes regulating the potassium channel activity in the Kathmandu population

indicates that the derived population takes advantage of preventing depolarization by maintaining

the potassium concentration. On the other hand, in the Malpighian tubules, the ion transport genes

were responsible for the active transmembrane transport rather than ion-specific channels. As

active transport is affected more than passive diffusion in the case of cold exposure (Koštál *et al.*,

2004; Zachariassen *et al.*, 2004; MacMillan & Sinclair, 2011), maintaining the ion balance in

response to acclimation depends on regulating the active transport mechanism.

Taken together, these results suggest that the ancestral population maintains the membrane fluidity

to prevent the loss of water-ion balance, whereas the derived population can maintain the active

transport in the Malpighian tubules and regulate the potassium ion concentration in the

extracellular space in response to acclimation, improving the cold tolerance. Therefore,

maintaining ion transport plays a crucial role in the adaptation of the derived Kathmandu

population to cold temperatures.

Considering the phenotypic variance within and between populations, comparative transcriptomic

analysis of the cold acclimation response also sheds light on the differences between the

phenotypes. The analysis focused on the hindgut tissue, where cellular respiration and lipid

metabolism-related genes were down-regulated in response to acclimation, whereas cuticle

development was up-regulated.

A previous study showed that prolonged exposure to cold reduces insect respiration rate (Colinet

et al., 2017). However, the same study also suggested that high tolerance can be achieved by

maintaining the energy requirement under cold stress. In line with these suggestions, even though

the respiration-related genes were down-regulated in response to acclimation in both phenotypes,

the difference between the tolerant and sensitive phenotypes was significant: sensitive lines

showed significant down-regulation of these genes compared to the tolerant lines. These

indications suggest that maintaining the respiration-related processes can be associated with fast

recovery from chill coma.

Together with respiration, lipid metabolism-related genes were significantly down-regulated in

hindguts in response to cold acclimation. As discussed above, low temperatures affect the lipid

bilayer structure of the membrane, and these impacts can be overcome by homeoviscious

adaptation. Even though the genes associated with fatty acid elongase activity were down-

regulated, no phenotype-specific difference was observed.

Previous studies showed that reabsorption in the hindgut plays a crucial role in cold tolerance of

Drosophila species (Andersen *et al.*, 2017; Andersen & Overgaard, 2020). Therefore, any

adaptation preventing or minimizing the leak in the hindgut is expected to increase tolerance to

cold. As the cuticle structure of the hindgut prevents water loss (Phillips, 1970), activation of

related pathways is expected to be observed in cold-tolerant strains. Interestingly, genes related to

cuticle development and chitin metabolism were up-regulated in the hindguts of all strains in

response to cold acclimation. However, the up-regulation significantly differed between the

phenotypes. As expected, tolerant strains showed more up-regulation of related genes than the

sensitive strains, suggesting that cuticle formation plays a vital role in the cold tolerance level of

the species.

The comparative transcriptomic analysis of the whole-body cold acclimation response also

explains the variation within the ancestral population. Differences were observed in GO terms

related to actin cytoskeleton and calcium channels between the fast and slow-recovering ancestral

strains. Genes related to both were up-regulated in response to acclimation, but the upregulation

was significantly more in the tolerant fast-recovering strains.

Actin cytoskeleton plays roles in various biological processes, including cell structure, shape,

migration, and signaling. Previous studies suggested that the actin cytoskeleton is essential in cold

tolerance (Cottam *et al*., 2006; Kim *et al*., 2006; von Heckel *et al*., 2016; Chen *et al*., 2017; Des

Marteaux *et al*., 2018; Bowman *et al*., 2018). One of these studies also showed that the tolerant

and sensitive Bangkok strains differed in actin polymerization: six genes had differential

expression in response to cold shock (Königer & Grath, 2018). On the other hand, five out of six

genes were up-regulated in both phenotypes in response to acclimation. However, they did not

show phenotype-specific differential expression even though the initial expression levels were

significantly lower in the sensitive phenotype.

In addition to the roles above, the actin cytoskeleton also acts as an anchor to ion channels and

regulates ion transport (reviewed in Denker & Barber, 2002). Previous studies showed that together

with microtubules, actin cytoskeleton dynamics affect the calcium ion influx. Calcium ion

concentration is shown to be affected by cold temperatures (Salkoff & Wyman, 1983; Findsen *et*

al., 2016) and plays an essential role in the activation of apoptotic and necrotic pathways (Koštál

et al., 2004, 2007, 2016; Zachariassen et al., 2004; MacMillan et al., 2015, 2016). Genes related

to calcium channels were also up-regulated in response to acclimation in the ancestral population.

However, the differential expression analysis between the two phenotypes of the ancestral

population showed that the calcium transport-related genes were significantly more up-regulated

in the fast-recovering phenotype than in the slow-recovering one. These results suggest that the

acclimation response of the two phenotypically distinct ancestral pools differ in actin cytoskeleton

and ion transport, focusing on calcium ions.

In this study, we showed that acclimation to mild temperatures profoundly affected the cold

tolerance of *D. ananassae* and investigated the mechanisms leading to increased tolerance. We

showed that different populations had distinct transcriptomic responses to acclimation in the whole

body and the Malpighian tubules. The differences between populations manifested through ion

transport and lipid metabolism. In addition, the cold tolerance phenotype affected the acclimation

response in the hindgut: tolerant strains could better maintain the cellular respiration that may

generate the energy required by the active transport mechanism and took advantage of the cuticle

structure that prevents water loss. Finally, we showed that the phenotypic variation within the

ancestral population resulted in differences in the acclimation response, including actin

cytoskeleton and calcium channel-related processes.

Further studies should be performed to validate these differences between the two phenotypes,

focusing on the function of the hindgut and cellular respiration. Hindgut reabsorption functional

assays were previously used in larger insects such as grasshoppers (Gerber *et al.*, 2021) but also

successfully performed in other *Drosophila* species (Andersen & Overgaard, 2020; Andersen *et*

al., 2017). Similarly, cellular respiration measurements were successful in other species (Colinet

et al., 2017) and can be used in *D. ananassae*. In addition to functional assays, candidate genes

arising from further analyses can be edited through CRISPR-*Cas9* technology previously

established in the species (Yılmaz *et al*., 2023) for functional validation.

Materials and Methods

Fly strains

This study used eight *D. ananassae* isofemale lines from Thailand (Bangkok, BKK) and three isofemale lines from Nepal (Kathmandu, KATH). The lines from Bangkok were the same as used in previous studies (Königer and Grath, 2018; Königer *et al.*, 2019). All lines were collected in 2002 (Das *et al.*, 2004). The fly strains were maintained under standard laboratory conditions (Rearing temperature – RT: 22˚C ± 1˚C 14:10 hours light:dark cycle) at low density on corn-molasses medium. Experiments were performed on flies that were expanded in large vials for two generations under the same conditions.

Cold Acclimation Assay

Acclimation conditions were determined via preliminary tests (see Supplementary Text, Supplementary Methods). Acclimation vials contained standard food and were kept at 10°C for 24 hours prior to the experiment to allow the food to cool down to acclimation temperature. Male flies were pooled per population and phenotype on the day of emergence. After 24 hours, flies were transferred to vials containing standard medium as described above. The vials were kept at 10°C for 4 days, followed by 90 minutes recovery time at rearing temperature.

RNA Extraction and Sequencing

Ten individual males per sample were snap-frozen in liquid nitrogen for whole-body samples, and RNA was extracted using MasterPure™ RNA Purification Kit (Epicentre, Madison, WI, USA) following the manufacturer's protocol. For hindgut and Malpighian tubule samples, tissues from 60 males per sample were dissected, and the tissues were collected into 1.5 mL Eppendorf tubes containing 10-20 µl 1X PBS. RNA extraction from tissues was performed using RNeasy Mini Kit

(Qiagen). RNA quantity and quality for all samples were confirmed with NanoDrop© (ND 1000,

VWR International, Radnor, PA, USA) and a bioanalyzer (Bioanalyzer 2001, Agilent

Technologies, Santa Clara, CA, USA, provided by the LMU genomics service unit). All samples

were sent for sequencing to an external facility (Novogene, Cambridge, UK), which performed

poly-A mRNA purification, reverse transcription, and high-throughput using a Novaseq 6000

sequencer (Illumina, San Diego, CA, USA) to generate 150-bp paired-end reads. On average, 88

million reads were generated per sample. An average of 73 million reads (83%) were successfully

mapped to the *D. ananassae* transcriptome (dana_r1.05_FB2016_01) using NextGenMap (version

0.5.2) (Sedlazeck *et al.*, 2013) and the annotation of FlyBase release 1.05. This release was used

to compare the results of the present study with previously described data on transcriptome

response following cold shock (Königer and Grath, 2018). All subsequent analyses were

performed in R (version 4.2.2) (R Core Team, 2022).

Differential Gene Expression Analysis

Differentially expressed genes were called using DESeq2 (version 1.38.3) (Love *et al.*, 2014). The

general response to treatment was assessed using ~Treatment as the design formula. The

differences between other variables (populations, phenotypes, or strains) were assessed using the

design formula ~Variable + Treatment + Variable:Treatment. Significant log2 fold changes

between control and cold-acclimated samples were determined using a Wald test and a subsequent

p-value multiple testing correction according to Benjamini-Hochberg. Genes with a corrected p-

value less than 0.05 were considered as significantly differentially expressed.

Gene Ontology (GO) Enrichment Analysis

GO analysis was performed using clusterProfiler (version 4.6.2) (Wu *et al.*, 2021) implemented in

R to determine the GO terms associated with *D. melanogaster* orthologs of the differentially

expressed genes. The p-values were corrected using FDR method with a p-value cut-off of 0.01.

The significant GO terms were visualized using the genekitr package (version 1.2.5) (Liu, 2023).

Weighted Gene Co-expression Network Analysis (WGCNA)

The network analysis was performed using the WGCNA package (version 1.72-1) (Langfelder &

Horvath, 2008) to determine co-expressed gene modules and their association with cold

acclimation response in specific tissues. The functional annotation of the genes in given modules

was performed using packages ReactomePA (version 1.42.0) (Yu & He, 2016) and clusterProfiler

to determine the enrichment of Reactome and KEGG pathways, respectively. The visualization

enrichment results were visualized using the enrichplot package (version 1.18.4) (Yu, 2023).

Comparison with Cold Shock

Previously, Königer and Grath (2018) analyzed differential gene expression before, at 15 minutes,

and 90 minutes after a cold shock in cold-sensitive and cold-tolerant strains of the Bangkok

population. The study used the same Bangkok strains used in the current study, but the samples

were not pooled for RNA extraction. Additionally, the data contained an extra time point (15

minutes after cold shock), which we did not test in the current study. Therefore, we used mapped

cold shock reads that were kindly provided by Königer and Grath, analyzed them with the same

DESeq2 model as the cold acclimation reads, and compared the DE genes between two phenotypes

of the BKK population upon cold acclimation to the DE genes between the same groups at 90

minutes after cold shock.

Data Availability

Gene expression data are available at GEO under the accession number GSE270239. All other data and scripts are available in the article and in its Supplementary Material.

Author Contributions

Conceptualization S.G.; investigation V.M.Y., Z.B.; writing-original draft V.M.Y., S.G.; visualization V.M.Y., S.G.; funding acquisition S.G.

Acknowledgements

Our work was supported by the Deutsche Forschungsgemeinschaft (project 271330745 to S. G.). We thank Hilde Lainer and Bianca Hölldobler for technical assistance in the laboratory, Amanda Glaser-Schmitt and John Parsch for their valuable input on the analysis of raw reads, and Annabella Königer for cold shock read counts. We extend our appreciation to all current and previous members of the Division of Evolutionary Biology, as well as the Graduate School Life Science Munich (LSM) at Ludwig-Maximilians-University Munich (LMU).

Conflicts of Interest

The authors declare no conflicting interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Andersen, J. L., Manenti, T., Sørensen, J. G., MacMillan, H. A., Loeschcke, V., & Overgaard, J. (2015). How to assess *Drosophila* cold tolerance: Chill coma temperature and lower lethal temperature are the best predictors of cold distribution limits. *Functional Ecology*, 29(1), 55–65. https://doi.org/10.1111/1365-2435.12310

Andersen, M. K., & Overgaard, J. (2020). Maintenance of hindgut reabsorption during cold exposure is a key adaptation for *Drosophila* cold tolerance. *Journal of Experimental Biology*, jeb.213934. https://doi.org/10.1242/jeb.213934

Andersen, M. K., MacMillan, H. A., Donini, A., & Overgaard, J. (2017). Cold tolerance of *Drosophila* species is tightly linked to epithelial K+ transport capacity of the Malpighian tubules and rectal pads. *Journal of Experimental Biology*, jeb.168518. https://doi.org/10.1242/jeb.168518

Bale, J. S., & Hayward, S. A. L. (2010). Insect overwintering in a changing climate. *Journal of Experimental Biology*, *213*(6), 980-994.

Bayley, J. S., Winther, C. B., Andersen, M. K., Grønkjær, C., Nielsen, O. B., Pedersen, T. H., & Overgaard, J. (2018). Cold exposure causes cell death by depolarization-mediated Ca2+ overload in a chill-susceptible insect. *Proceedings of the National Academy of Sciences*, *115*(41), E9737-E9744.

Bowman, L. L., Kondrateva, E. S., Timofeyev, M. A., & Yampolsky, L. Y. (2018). Temperature gradient affects differentiation of gene expression and SNP allele frequencies in the dominant Lake Baikal zooplankton species. *Molecular Ecology*, *27*(11), 2544–2559. https://doi.org/10.1111/mec.14704

Bubliy, O. A., Riihimaa, A., Norry, F. M., & Loeschcke, V. (2002). Variation in resistance and acclimation to low-temperature stress among three geographical strains of *Drosophila melanogaster*. *Journal of Thermal Biology*, 27(5), 337–344. https://doi.org/10.1016/S0306-4565(01)00098-5

Chen, Y., Jiang, T., Zhu, J., Xie, Y., Tan, Z., Chen, Y., Tang, S., Hao, B., Wang, S., Huang, J., & Shen, X. (2017). Transcriptome sequencing reveals potential mechanisms of diapause preparation in bivoltine silkworm *Bombyx mori* (Lepidoptera: *Bombycidae*). *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, *24*, 68–78. https://doi.org/10.1016/j.cbd.2017.07.003

Colinet, H., Renault, D., & Roussel, D. (2017). Cold acclimation allows *Drosophila* flies to
maintain mitochondrial functioning under cold stress. *Insect Biochemistry and Molecular*
Biology, 80, 52–60. https://doi.org/10.1016/j.ibmb.2016.11.007

Cottam, D. M., Tucker, J. B., Rogers-Bald, M. M., Mackie, J. B., Macintyre, J., Scarborough, J.
A., Ohkura, H., & Milner, M. J. (2006). Non-centrosomal microtubule-organising centres in
cold-treated cultured *Drosophila* cells. *Cell Motility*, *63*(2), 88–100.
https://doi.org/10.1002/cm.20103

Das, A., Mohanty, S., & Stephan, W. (2004). Inferring the Population Structure and Demography
of *Drosophila ananassae* From Multilocus Data. *Genetics*, 168(4), 1975–1985.
https://doi.org/10.1534/genetics.104.031567

David, J. R., Gibert, P., Moreteau, B., Gilchrist, G. W., & Huey, R. B. (2003). The fly that came
in from the cold: Geographic variation of recovery time from low-temperature exposure in
Drosophila subobscura. *Functional Ecology*, 17(4), 425–430. https://doi.org/10.1046/j.1365-
2435.2003.00750.x

Denker, S. P., & Barber, D. L. (2002). Ion transport proteins anchor and regulate the cytoskeleton.
Current Opinion in Cell Biology, *14*(2), 214–220. https://doi.org/10.1016/S0955-
0674(02)00304-6

Des Marteaux, L. E., Khazraeenia, S., Yerushalmi, G. Y., Donini, A., Li, N. G., & Sinclair, B. J.
(2018). The effect of cold acclimation on active ion transport in cricket ionoregulatory tissues.
Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, *216*,
28–33. https://doi.org/10.1016/j.cbpa.2017.11.005

Drobnis, E. Z., Crowe, L. M., Berger, T., Anchordoguy, T. J., Overstreet, J. W., & Crowe, J. H.
(1993). Cold shock damage is due to lipid phase transitions in cell membranes: a demonstration
using sperm as a model. *Journal of Experimental Zoology*, *265*(4), 432-437.

Enriquez, T., & Colinet, H. (2019). Cold acclimation triggers major transcriptional changes in
Drosophila suzukii. *BMC Genomics*, 20(1), 413. https://doi.org/10.1186/s12864-019-5745-7

Findsen, A., Overgaard, J., & Pedersen, T. H. (2016). Reduced L-type Ca2+ current and
compromised excitability induce loss of skeletal muscle function during acute cooling in
locust. *Journal of Experimental Biology*, *219*(15), 2340-2348.

Findsen, A., Pedersen, T. H., Petersen, A. G., Nielsen, O. B., & Overgaard, J. (2014). Why do
insects enter and recover from chill coma? Low temperature and high extracellular potassium

compromise muscle function in *Locusta migratoria*. *Journal of Experimental Biology*, *217*(8), 1297-1306.

Free, J. B., & Spencer-Booth, Y. (1960). Chill-coma and cold death temperatures of *Apis mellifera*. *Entomologia Experimentalis et Applicata*, 3(3), 222–230. https://doi.org/10.1111/j.1570-7458.1960.tb00451.x

Gerber, L., & Overgaard, J. (2018). Cold tolerance is linked to osmoregulatory function of the hindgut in *Locusta migratoria*. *Journal of Experimental Biology*, jeb.173930. https://doi.org/10.1242/jeb.173930

Gerber, L., Kresse, J.-C., Šimek, P., Berková, P., & Overgaard, J. (2021). Cold acclimation preserves hindgut reabsorption capacity at low temperature in a chill-susceptible insect, *Locusta migratoria*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *252*, 110850. https://doi.org/10.1016/j.cbpa.2020.110850

Gibert, P., Moreteau, B., Pétavy, G., Karan, D., & David, J. R. (2001). Chill-Coma Tolerance, a Major Climatic Adaptation Among *Drosophila* Species. *Evolution*, *55*(5), 1063–1068. https://doi.org/10.1111/j.0014-3820.2001.tb00623.x

Hallas, R., Schiffer, M., & Hoffmann, A. A. (2002). Clinal variation in *Drosophila serrata* for stress resistance and body size. *Genetical Research*, 79(2), 141–148. https://doi.org/10.1017/S0016672301005523

Hoedjes, K. M., Grath, S., Posnien, N., Ritchie, M. G., Schlötterer, C., Abbott, J. K., Almudi, I., Coronado-Zamora, M., Durmaz Mitchell, E., Flatt, T., Fricke, C., Glaser-Schmitt, A., González, J., Holman, L., Kankare, M., Lenhart, B., Orengo, D. J., Snook, R. R., Yılmaz, V. M., & Yusuf, L. (2024). From whole bodies to single cells: A guide to transcriptomic approaches for ecology and evolutionary biology. *Molecular Ecology*, e17382. https://doi.org/10.1111/mec.17382

Hoffmann, A. A., & Watson, M. (1993). Geographical Variation in the Acclimation Responses of *Drosophila* to Temperature Extremes. *The American Naturalist*, 142, S93–S113. https://doi.org/10.1086/285525

Hoffmann, A. A., & Weeks, A. R. (2007). Climatic selection on genes and traits after a 100 year-old invasion: A critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern Australia. *Genetica*, 129(2), 133. https://doi.org/10.1007/s10709-006-9010-z

Hoffmann, A. A., Anderson, A., Hallas, R. (2002). Opposing clines for high and low temperature
resistance in *Drosophila melanogaster*. *Ecology Letters*, 5(5), 614–618.
https://doi.org/10.1046/j.1461-0248.2002.00367.x

Hoyle, G. (1953). Potassium ions and insect nerve muscle. *Journal of Experimental Biology*, *30*(1),
121-135.

Kelty, J. D., Killian, K. A., & Lee, R. E. (1996). Cold shock and rapid cold-hardening of pharate
adult flesh flies (*Sarcophaga crassipalpis*): Effects on behaviour and neuromuscular function
following eclosion. *Physiological Entomology*, 21(4), 283–288.
https://doi.org/10.1111/j.1365-3032.1996.tb00866.x

Kim, M., Robich, R. M., Rinehart, J. P., & Denlinger, D. L. (2006). Upregulation of two actin
genes and redistribution of actin during diapause and cold stress in the northern house
mosquito, *Culex pipiens*. *Journal of Insect Physiology*, *52*(11–12), 1226–1233.
https://doi.org/10.1016/j.jinsphys.2006.09.007

Königer, A., & Grath, S. (2018). Transcriptome Analysis Reveals Candidate Genes for Cold
Tolerance in *Drosophila ananassae*. *Genes*, 9(12), 624. https://doi.org/10.3390/genes9120624

Königer, A., Arif, S., & Grath, S. (2019). Three Quantitative Trait Loci Explain More than 60%
of Variation for Chill Coma Recovery Time in a Natural Population of *Drosophila ananassae*.
G3 Genes|Genomes|Genetics, 9(11), 3715–3725. https://doi.org/10.1534/g3.119.400453

Koštál, V. (2010). Cell structural modifications in insects at low temperature. *Low temperature
biology of insects*, 116-140.

Koštál, V., Korbelová, J., Štětina, T., Poupardin, R., Colinet, H., Zahradníčková, H., ... & Šimek,
P. (2016). Physiological basis for low-temperature survival and storage of quiescent larvae of
the fruit fly *Drosophila melanogaster*. *Scientific Reports*, *6*(1), 32346.

Koštál, V., Renault, D., Mehrabianová, A., & Bastl, J. (2007). Insect cold tolerance and repair of
chill-injury at fluctuating thermal regimes: Role of ion homeostasis. *Comparative
Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 147(1), 231–238.
https://doi.org/10.1016/j.cbpa.2006.12.033

Koštál, V., Vambera, J., & Bastl, J. (2004). On the nature of pre-freeze mortality in insects: water
balance, ion homeostasis and energy charge in the adults of *Pyrrhocoris apterus*. *Journal of
experimental biology*, *207*(9), 1509-1521.

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics, 9*, 559. https://doi.org/10.1186/1471-2105-9-559

Liu, Y. (2023). genekitr: Gene Analysis Toolkit (R package version 1.2.5). https://CRAN.R-
project.org/package=genekitr

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology, 15*(12), 550.

MacMillan, H. A., & Sinclair, B. J. (2011). Mechanisms underlying insect chill-coma. *Journal of*
Insect Physiology, *57*(1), 12–20. https://doi.org/10.1016/j.jinsphys.2010.10.004

MacMillan, H. A., Andersen, J. L., Davies, S. A., & Overgaard, J. (2015). The capacity to maintain
ion and water homeostasis underlies interspecific variation in *Drosophila* cold tolerance.
Scientific Reports, *5*(1), 18607. https://doi.org/10.1038/srep18607

MacMillan, H. A., Knee, J. M., Dennis, A. B., Udaka, H., Marshall, K. E., Merritt, T. J. S., &
Sinclair, B. J. (2016). Cold acclimation wholly reorganizes the *Drosophila melanogaster*
transcriptome and metabolome. *Scientific Reports*, *6*(1), 28999.
https://doi.org/10.1038/srep28999

MacMillan, H. A., Yerushalmi, G. Y., Jonusaite, S., Kelly, S. P., & Donini, A. (2017). Thermal
acclimation mitigates cold-induced paracellular leak from the *Drosophila* gut. *Scientific*
Reports, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-08926-7

Overgaard, J., & MacMillan, H. A. (2017). The Integrative Physiology of Insect Chill Tolerance.
Annual Review of Physiology, *79*(1), 187–208. https://doi.org/10.1146/annurev-physiol-
022516-034142

Overgaard, J., Gerber, L., & Andersen, M. K. (2021). Osmoregulatory capacity at low temperature
is critical for insect cold tolerance. *Current Opinion in Insect Science*, 47, 38–45.
https://doi.org/10.1016/j.cois.2021.02.015

Parker, D. J., Envall, T., Ritchie, M. G., & Kankare, M. (2021). Sex-specific responses to cold in
a very cold-tolerant, northern *Drosophila* species. *Heredity*, 126(4), 695–705.
https://doi.org/10.1038/s41437-020-00398-2

Parker, D. J., Vesala, L., Ritchie, M. G., Laiho, A., Hoikkala, A., & Kankare, M. (2015). How
consistent are the transcriptome changes associated with cold acclimation in two species of the
Drosophila virilis group? *Heredity*, 115(1), 13–21. https://doi.org/10.1038/hdy.2015.6

Phillips, J. E. (1970). Apparent Transport of Water by Insect Excretory Systems. *American*
Zoologist, 10(3), 413–436. https://doi.org/10.1093/icb/10.3.413

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation for
Statistical Computing. https://www.R-project.org/

Reinhold, J. M., Lazzari, C. R., & Lahondère, C. (2018). Effects of the environmental temperature
on *Aedes aegypti* and *Aedes albopictus* mosquitoes: a review. *Insects*, *9*(4), 158.

Ring, R. A. (1982). Freezing-tolerant insects with low supercooling points. *Comparative*
Biochemistry and Physiology Part A: Physiology, *73*(4), 605-612.

Salkoff, L. B., & Wyman, R. J. (1983). Ion currents in *Drosophila* flight muscles. *The Journal of*
physiology, *337*(1), 687-709.

Sedlazeck, F. J., Rescheneder, P., & Von Haeseler, A. (2013). NextGenMap: Fast and accurate read
mapping in highly polymorphic genomes. *Bioinformatics*, 29(21), 2790–2791.
https://doi.org/10.1093/bioinformatics/btt468

Sinensky, M. (1974). Homeoviscous Adaptation—A Homeostatic Process that Regulates the
Viscosity of Membrane Lipids in *Escherichia coli*. *Proceedings of the National Academy of*
Sciences, 71(2), 522–525. https://doi.org/10.1073/pnas.71.2.522

Sisodia, S., & Singh, B. N. (2010). Resistance to environmental stress in *Drosophila ananassae*:
Latitudinal variation and adaptation among populations. *Journal of Evolutionary Biology*,
23(9), 1979–1988. https://doi.org/10.1111/j.1420-9101.2010.02061.x

Steponkus, P. L. (1984). Role of the plasma membrane in freezing injury and cold
acclimation. *Annual Review of Plant Physiology*, *35*(1), 543-584.

Teets, N. M., & Denlinger, D. L. (2013). Physiological mechanisms of seasonal and rapid cold-
hardening in insects. *Physiological Entomology*, 38(2), 105–116.
https://doi.org/10.1111/phen.12019

Teets, N. M., Kawarasaki, Y., Lee, R. E., & Denlinger, D. L. (2012). Energetic consequences of
repeated and prolonged dehydration in the Antarctic midge, *Belgica antarctica*. *Journal of*
Insect Physiology, 58(4), 498–505. https://doi.org/10.1016/j.jinsphys.2011.11.011

Udaka, H., Ueda, C., & Goto, S. G. (2010). Survival rate and expression of *Heat-shock protein 70*
and *Frost* genes after temperature stress in *Drosophila melanogaster* lines that are selected for
recovery time from temperature coma. *Journal of Insect Physiology*, *56*(12), 1889-1894.

von Heckel, K., Stephan, W., & Hutter, S. (2016). Canalization of gene expression is a major signature of regulatory cold adaptation in temperate *Drosophila melanogaster*. *BMC Genomics*, 17(1), 574. https://doi.org/10.1186/s12864-016-2866-0

Watson, M. J. O., & Hoffmann, A. A. (1996). Acclimation, cross-generation effects, and the response to selection for increased cold resistance in *Drosophila*. *Evolution*, 50(3), 1182–1192. https://doi.org/10.1111/j.1558-5646.1996.tb02359.x

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X., & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation, 2*(3), 100141.

Yerushalmi, G. Y., Misyura, L., MacMillan, H. A., & Donini, A. (2018). Functional plasticity of the gut and the Malpighian tubules underlies cold acclimation and mitigates cold-induced hyperkalemia in *Drosophila melanogaster*. *Journal of Experimental Biology*, jeb.174904. https://doi.org/10.1242/jeb.174904

Yılmaz, V. M., Ramnarine, T. J. S., Königer, A., Mussgnug, S., & Grath, S. (2023). Tropical super flies: Integrating *Cas9* into *Drosophila ananassae* and its phenotypic effects. *Journal of Insect Physiology*, 147, 104516. https://doi.org/10.1016/j.jinsphys.2023.104516

Yu, G. (2023). enrichplot: Visualization of Functional Enrichment Result (R package version 1.18.4). https://yulab-smu.top/biomedical-knowledge-mining-book/

Yu, G., & He, Q. Y. (2016). ReactomePA: an R/Bioconductor package for reactome pathway analysis and visualization. *Molecular BioSystems, 12*(2), 477-479.

Zachariassen, K. E., Kristiansen, E., & Pedersen, S. A. (2004). Inorganic ions in cold-hardiness. *Cryobiology*, *48*(2), 126-133

Supplementary Material

Navigating the Cold: Integrative Transcriptome Sequencing Approach Reveals Ionoregulatory and Whole-Body Responses to Cold Acclimation in *Drosophila ananassae*

Vera Miyase Yılmaz, Zhihui Bao and Sonja Grath

Supplementary Text

Chill Coma Recovery Time Improves with Acclimation to Mild Temperatures

Prior exposure to mild stress conditions can improve the tolerance phenotype. To test the effect of temperature and duration of acclimation on CCRT, we tested four temperatures (6-12˚C) for different periods (1-5 days). We used three strains representing fast-recovering Bangkok (BKK12 for FastBKK), slow-recovering Bangkok (BKK13 for SlowBKK), and Kathmandu (KATH14 for KATH) populations. Flies exposed to 6˚C for more than a day showed high mortality, exceeding 50% on day three and reaching 100% on day 5 (Supplementary Figure 1A, Supplementary Table 1). Additionally, the surviving flies showed CCRT values higher than the control (Supplementary Figure 1B). The remaining temperatures resulted in minimal mortality, reaching an average of 10% at 8˚C and almost no mortality at 10˚C and 12˚C. These temperatures also lead to an improvement in the CCRT phenotype (Supplementary Figure 1B, Supplementary Table 2). Even though the effect of the remaining temperatures did not significantly differ from one another, the difference to control was largest when the acclimation temperature was 10˚C, and the duration was four days, which was chosen as the acclimation set-up for further experiments.

Supplementary Methods

Cold Acclimation Preliminary Assays

To determine the temperature and duration of acclimation, we tested five time periods (1, 2, 3, 4, and 5 days) and four temperatures (6°C, 8°C, 10°C, and 12°C) using BKK12, BKK13, and KATH14, representing FastBKK, SlowBKK, and KATH populations, respectively. The ranges of temperature and duration were determined based on acclimation studies on other *Drosophila* species. The flies were controlled for age and the effect of acclimation was tested using the survival of acclimated flies and chill coma recovery time (CCRT) upon acclimation. Five replicates were used for each acclimation regime. On the day of emergence, 10 male flies were collected and placed into 50 ml vials with 5 ml food at room temperature. On the following day, test flies were flipped into new vials that were kept at the acclimation temperature and placed into cold incubators for 5-, 4-, 3-, 2-, and 1-day acclimation. The control flies were kept at rearing temperature (22 ± 1°C). After acclimation, the flies were placed back to room temperature for 24 hours for recovery, after which the dead flies were counted and removed from the vials, whereas the surviving flies were flipped into empty vials and placed on melting ice for 3 hours for CCRT measurement. At the end of 3 hours, the vials were placed into room temperature and the recovery time of each fly was measured. The temperature and duration that lead to minimal mortality and improved the CCRT were determined.

Supplementary Figures

Supplementary Figure 1: Effect of duration of acclimation on A) mortality and B) chill coma recovery time (CCRT) phenotype. The lines that were used were representatives of the respective population and phenotype. Ten flies per replicate in five replicates were used for mortality measurements and dead flies were discarded. The remaining flies were used for chill coma recovery assays. Colored dots indicate proportional mortality of each replicate and black dots indicate outliers with high or low CCRT.

Supplementary Figure 2: A) Venn diagram with the number of differentially expressed (DE) genes specific to or shared between populations. Bangkok and Kathmandu populations exhibited both shared, as well as exclusive differentially expressed genes in response to acclimation. B) Gene ontology (GO) enrichment results for genes differentially expressed between the populations in response to cold acclimation treatment in the whole body. Biological processes (red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate panels.

Supplementary Figure 3: Transcriptional response to cold acclimation in the Malpighian tubules. Gene enrichment analysis results for genes A) up-regulated and B) down-regulated in response to cold acclimation. Biological processes (red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate panels.

Supplementary Figure 4: Transcriptional response to cold acclimation in the hindguts. A) Principal component analysis (PCA) after the outlier was removed from the analysis. The samples are shown according to the pooled strains (color) and treatment (shape). FastBKK, SlowBKK, and KATH sample pools are represented by purple, blue, and green colors, respectively. The control samples are represented by circles and the cold acclimated samples are represented by triangles. Gene enrichment analysis results for genes B) up-regulated and C) down-regulated in response to cold acclimation. Biological processes (red bars), cellular components (blue bars), and molecular functions (yellow bars) are shown in separate panels. Cellular respiration and chitin metabolism play roles in the acclimation response of tolerant and sensitive strains in the hindgut.

Supplementary Figure 5: Weighted gene co-expression network analysis (WGCNA). Dot-plots for functional annotation of genes in modules that are significantly enriched in differentially expressed genes in response to cold acclimation in the whole body, Malpighian tubule, and hindgut samples. A) General response to acclimation in the whole body, B-C) population specific response to acclimation in the whole body, D-E) general response to acclimation in the Malpighian tubules, and F) general response to acclimation in the hindguts are shown. The size of the dot represents the gene count associated with the term and the color represents the adjusted p-value.

Supplementary Figure 6: Upset plot showing differentially expressed genes and their overlap between the cold shock (CS) and cold acclimation (CA) assays. Black, horizontal bars indicate the total number of differentially expressed genes that were significantly up- or down-regulated in a sample in response to cold acclimation. A black circle indicates that a sample and direction (up/down) combination is included in an intersection class. Intersection classes with a single circle are comprised of genes significantly differentially expressed in a single treatment and direction. Circles connected by a line indicate intersection classes comprised of multiple treatment and direction combinations. Vertical bars correspond to the number of genes in each intersection class.

Supplementary Tables

Supplementary Table 1: Tukey's HSD test for proportional mortality after cold exposure.

	1-day	**2-day**	**3-day**	**4-day**	**5-day**
6˚C	e	d	c	b	a
8˚C	e	e	de	de	de
10˚C	e	e	e	e	e
12˚C	e	e	e	e	e
control	e	e	e	e	e

Note 1: Samples that do not share any letter(s) are significantly different. *E.g.* Mortality upon 2-day exposure to 6˚C (**d**) significantly differs from mortality upon 3-day exposure to 6˚C (**c**), but not from mortality upon 3-day exposure to 8˚C (**de**).

Supplementary Table 2: Tukey's HSD test for CCRT values upon acclimation.

	1-day	**2-day**	**3-day**	**4-day**	**5-day**
8˚C	def	ef	cdef	def	cde
10˚C	def	def	def	f	ef
12˚C	cde	def	def	def	cde
control	cd	c	b	b	a

Note 1: Samples that do not share any letter(s) are significantly different. *E.g.* CCRT of 2-day control samples (**c**) significantly differs from CCRT of 3-day control samples (**b**), but not from CCRT of 1-day control samples (**cd**).
Note 2: Since the flies that survived the acclimation were used in the chill coma recovery assay, 6˚C acclimation was not included in the statistical test for CCRT due to high mortality (50-100%).

Supplementary Table 3: GO enrichment analysis results for differentially expressed genes exclusively in the whole body, Malpighian tubule, or hindgut samples in response to treatment.

Supplementary Table 4: GO enrichment analysis results for differentially expressed genes in the whole body, Malpighian tubule, and hindgut samples in response to treatment in opposite directions.

Supplementary Table 5: GO enrichment analysis results for differentially expressed genes in the whole body in response to treatment.

Supplementary Table 6: GO enrichment analysis results for differentially expressed genes in the whole body in response to treatment. The comparison of Kathmandu and Bangkok populations is shown. The column "Population" indicates which population showed positive log2FC value.

Supplementary Table 7: GO enrichment analysis results for differentially expressed genes in the whole body in response to treatment. The comparison of fast- and slow-recovering Bangkok lines is shown. The column "Line" indicates which line showed positive log2FC value.

Supplementary Table 8: GO enrichment analysis results for differentially expressed genes in the Malpighian tubules in response to treatment.

Supplementary Table 9: GO enrichment analysis results for differentially expressed genes in the Malpighian tubules in response to treatment. The comparison of Kathmandu and Bangkok populations is shown. The column "Population" indicates which population showed positive log2FC value.

Supplementary Table 10: GO enrichment analysis results for differentially expressed genes in the hindguts in response to treatment.

Supplementary Table 11: GO enrichment analysis results for differentially expressed genes in the hindguts in response to treatment. The comparison of fast and slow phenotypes is shown. The column "Phenotype" indicates which phenotype showed positive log2FC value.

Supplementary Table 12: GO enrichment analysis results for differentially expressed genes in the whole body in response to cold acclimation and cold treatment.

Supplementary Table 13: GO enrichment analysis results for differentially expressed genes in the whole body in response to cold acclimation or cold treatment.

Supplementary Table 14: GO enrichment analysis results for differentially expressed genes in the whole body in response to cold acclimation and cold treatment in opposite directions.