TITLE: Three-dimensional culture in a bioengineered matrix and somatic cell

complementation to improve growth and survival of bovine preantral follicles

Juliana I. Candelaria[1#], Ramon C. Botigelli[1], Carly Guiltinan[1], Ariella Shikanov[2,3,4], Anna

C. Denicol[1*]

[1]Department of Animal Science, University of California Davis, Davis, CA, USA.
[2] Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI, USA.
[3]Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI, USA.
[4]Department of Macromolecular Science and Engineering, University of Michigan, Ann
Arbor, MI, USA.
[#]Current address: Postdoctoral Fellow, Department of Biochemistry and Molecular
Biology, Uniformed Services University of the Health Sciences, Bethesda, MD, USA.
*Corresponding author: Anna C. Denicol
acdenicol@ucdavis.edu

STRUCTURED ABSTRACT

Purpose: Here we explored poly(ethylene glycol) (PEG) bioengineered hydrogels for

bovine preantral follicle culture with or without ovarian cell co-culture and examined the

potential for differentiation of bovine embryonic stem cells (bESCs) towards gonadal

somatic cells to develop a system more similar to the ovarian microenvironment.

Methods: Bovine preantral follicles were first cultured in two-dimensional (2D) control or

within PEG hydrogels (3D) and then co-cultured within PEG hydrogels with bovine

ovarian cells (BOCs) to determine growth and viability. Finally, we tested conditions to

drive differentiation of bESCs towards the intermediate mesoderm and bipotential gonad

fate.

Results: Primary follicles grew over the 10-day culture period in PEG hydrogels compared

to 2D control. Early secondary follicles maintained a similar diameter within the PEG while

control follicles decreased in size. Follicles lost viability after co-encapsulation with BOCs;

BOCs lost stromal cell signature over the culture period within hydrogels. Induction of

bESCs towards gonadal somatic fate under WNT signaling was sufficient to upregulate

intermediate mesoderm (*LHX1*) and early coelomic epithelium/bipotential gonad markers

(*OSR1*, *GATA4*, *WT1*). Higher BMP4 concentrations upregulated the lateral plate

mesoderm marker *FOXF1*. *PAX3* expression was not induced, indicating absence of the

paraxial mesoderm lineage.

Conclusions: Culture of primary stage preantral follicles in PEG hydrogels promoted

growth compared to controls; BOCs did not maintain identity in the PEG hydrogels.

Collectively, we demonstrate that PEG hydrogels can be a potential culture system for

early preantral follicles pending refinements, which could include addition of ESC-derived

ovarian somatic cells using the protocol described here.

39
40    KEYWORDS
41
42    Preantral follicle, three-dimensional culture, embryonic stem cells, ovary, bovine,

43    biomimetic

44
45    CAPSULE SUMMARY
46
47    We demonstrate that three-dimensional bioengineered hydrogels could aid in the survival

48    and growth of small bovine preantral follicles. Moreover, bovine embryonic stem cells

49    have the potential to differentiate towards precursors of somatic gonadal cell types,

50    presenting an alternative cell source for preantral follicle co-culture.

## INTRODUCTION

Folliculogenesis is a complex and selective process resulting in production of a limited quantity of meiotically and developmentally competent oocytes over a female's lifespan. The molecular processes that coordinate the activation, growth and maturation of ovarian follicles have yet to be fully elucidated, especially during the early preantral stages of development. Progress in understanding preantral folliculogenesis in non-rodent species has been hindered by the lack of appropriate in vitro culture systems and an insufficient knowledge base of ovarian development and function. This is particularly true in large mammal (i.e. cows and humans) preantral folliculogenesis which have prolonged duration and sizeable volumetric changes between the primordial and pre-ovulatory follicle stages. A great advantage of performing in vitro folliculogenesis beginning at the preantral stage includes establishing an alternative fertility preservation and restoration method by utilizing the vast preantral follicle pool for oocyte production [1]. More specifically, this approach would be beneficial for prepubertal girls suffering from blood-borne malignancies as isolated preantral follicles could be matured in vitro without the risk of reintroducing malignant cells or surgical intervention, as could be the case with the culture and reintroduction of ovarian tissue fragments.

Recapitulation of bovine preantral folliculogenesis has been attempted by several groups, but there has been little success in efficiently growing follicles to the preovulatory stage and yielding metaphase II oocytes. Although 2-dimensional culture systems have shown some promise in supporting bovine preantral follicle growth [2–4], mounting evidence suggests that 3-dimensional (3D) culture systems better promote development due to their ability to maintain a follicle's spherical configuration and preserve cell contact

both within the follicular unit (i.e. granulosa cell and oocyte interface) and outside the

basement membrane where theca cells are located [5–8]. Matrigel and alginate are

commonly used biomaterials to create hydrogels for in vitro follicle culture, however

pitfalls include uncontrollable degradation and biological inertness, respectively [9].

Therefore, a highlighted alternative is the use of bioengineered 3D hydrogel culture

systems that permit timely degradation by the follicle yet retaining the mechanical support

necessary to keep its spherical structure [10,11]. Poly(ethylene glycol) (PEG) hydrogels

constructed with proteolytically-degradable peptide crosslinkers have been employed for

mouse in vitro folliculogenesis and demonstrated success in growing preantral follicles to

the antral stage and producing metaphase II oocytes [12]. Likewise, the addition of

adipose-derived stem cells or conditioned media to PEG hydrogels containing mouse

preantral follicles further aided development and survival [13]. Despite the promise

demonstrated in mice, the use of PEG hydrogels for large mammal in vitro preantral

follicle culture has not been reported. Moreover, investigating cell supplementation in the

hydrogel system to determine their propensity to become theca cells and/or

advantageous effect on follicle growth has not been directly assessed. Indeed, early

preantral follicles do not contain theca cells and must recruit precursor cells from the

stromal environment to build the essential theca layer, hence addition of cells to the

system would be essential when starting with primordial- or primary-stage follicles. More

specifically, the hedgehog signaling pathway is key to driving theca cell differentiation and

recruitment [14–16], therefore cells added to the culture system should harbor the cellular

machinery to respond to hedgehog ligands secreted by the follicular granulosa cells.

Although native ovarian stromal cells have been used as feeder cells for 2D and

3D in vitro follicle culture [17–19], surgical intervention is required to obtain ovarian tissue

and harvest cells. Moreover, their abilities to phenotypically-mirror theca cells upon co-

culture with follicles has not been shown to date. Alternative cell sources, particularly

those of renewable capacity (i.e. stem cells) present a unique advantage to generating

cells that could give rise to a theca cell-like phenotype. To that end, stepwise in vitro

differentiation of pluripotent stem cells into gonadal-like cells has been demonstrated in

humans and mice [20–22] thus highlighting the potential to create a substitute cell source

with prospective theca cell characteristics. Cell fate decisions that result in formation of

the bipotential gonad in the embryo are the culmination of precisely timed cell signaling

via gradients of morphogens such as FGF, BMP, and WNT proteins. Hence, protocols

have been devised employing a stepwise induction of pluripotent stem cells using FGF,

BMP, and WNT signaling such that cells progress through a mesoderm-like state, and

finally a bipotential gonad-like state [23–25].

In this study, we describe the use of a PEG hydrogel culture system with or without

native ovarian cells for the in vitro development of bovine preantral follicles and explore

the potential of bovine embryonic stem cells to be induced towards gonadal-like cells in

vitro. We first hypothesized that culture in PEG hydrogels would improve the growth and

viability of bovine preantral follicles. By examining the gene expression of ECM-degrading

enzymes of primary- and early secondary-stage follicles, we tailored the hydrogel design

to respond to bovine follicle enzyme secretion. Based on the evidence that cell

supplementation improves mouse follicle development, we also examined the effect of

dissociated bovine ovarian cells (BOCs) on bovine preantral follicle growth and if BOCs

exhibit a pre-theca cell phenotype when cultured in vitro using PEG hydrogels. Finally,

we hypothesized that bovine embryonic stem cells (bESCs) can be differentiated into the

intermediate mesoderm and early coelomic epithelium/bipotential gonad.

MATERIAL AND METHODS
124
All materials were purchased from Thermo Fisher Scientific unless otherwise specified.

_Follicle Isolation and bovine ovary single-cell dissociation_

Fresh bovine preantral follicles were isolated as previously described [26]. Follicles

were maintained in warmed follicle wash media until subsequent experiments. For bovine

ovary single-cell dissociation (n = 3 replicates), cortical tissue (250-300 mg) was

fragmented into 500 $\mu m^2$ pieces incubated with Hank's balanced salt solution ($Ca^{2+}/Mg^{2+}$)

containing 1 mg/mL collagenase IV, 1 U/mL DNase I, and 50 µg/mL Liberase

(5401119001, Sigma Aldrich). The solution with tissue was shaken at 38.5 °C for 20 min,

pipetted 15 times using a serological pipette, and shaken again for 20 min to mechanically

aid in the dissociation of cells. Enzymatic activity was stopped by adding 20% v/v fetal

bovine serum to the solution. The tissue-dissociated solution was filtered through a 100

µm then through a 35 µm cell strainer. Cells collected after final filtration were centrifuged

(300 x$g$ for 5 min) and the resulting pellet was resuspended in 1 mL Hank's balanced salt

solution without $Ca^{2+}/Mg^+$ until encapsulation.

_PEG hydrogel materials, preparation, and follicle encapsulation_

PEG hydrogel experiments were conducted using 8-arm PEG vinyl sulfone (PEG-

VS) (40 kDa, >99% purity, JenKem Technology) and crosslinked with MMP- and Plasmin-

sensitive peptide sequences (Ac-GCRD**VPMS**MRGGDRCG**YKNS**CG, i.e. YKNS/VPMS)

(2391.8 g/mol, >90%, Celtek Peptides). PEG and crosslinker peptide were dissolved in

isotonic 50 mM HEPES buffer and mixed to a final composition containing 5% PEG-VS

with 1:1 stoichiometric ratio of VS to thiol (side chain present on cysteines in peptide

sequence). Sixteen-µl hydrogels containing preantral follicles were formed between

parafilm-lined glass slides and allowed to gelatinize for up to 10 min. Then, hydrogels

were transferred to warmed follicle maintenance media to swell and placed in a humidified

incubator (37°C). Freshly isolated follicles were maintained in warmed follicle wash media

until encapsulation. In experiment 1, follicles (n = 1-4) were first washed in 10% PEG-VS

and either encapsulated in PEG hydrogels or placed into 2D culture. Hydrogel-

encapsulated and 2D culture follicles were placed into individual wells of a 96-well plate

containing follicle culture media (1:1 ratio of α-MEM and F12 with glutamine

supplemented with 3 mg/mL bovine serum albumin, 100 ng/mL penicillin/streptomycin,

1 mM sodium pyruvate, 1 % (v/v) non-essential amino acids, 10 mg/l insulin, 5.5 mg/l

transferrin and 6.7 µg/L selenium (ITS), 100 ng/mL human recombinant follicle stimulating

hormone (228-12609-2, RayBiotech), and 50 µg/mL ascorbic acid. Half of the spent media

was changed every 2 days and follicles were cultured for 10 days. To assess follicle

growth over the culture period, the average of 2 perpendicular diameter measurements

were taken at day 0, day 5, and day 10 using an inverted microscope (Revolve, Echo

Labs). The percentage of growing and non-growing follicles (growth defined as having at

least 5% increase in diameter by day 10 compared to day 0) and percent increase in size

in growing-only follicles was calculated. In total, 15 replicates were conducted with a total

of 216 follicles (n = 153 in PEG and n = 63 in 2D control)

*Cell viability staining and quantification*

Bovine ovarian cells and MeLCs were resuspended in PEG-VS + crosslinker

peptide VPMS/YKNS at a concentration of 1, 2, 10, and 20 x $10^6$ cells/mL to test cell

viability immediately after encapsulation, 5 and 10 days after in vitro culture. Hydrogels

were cultured individually in 96-well plates with 150 µl of follicle culture media with half

media changes every 2 days. After culture, hydrogels were incubated for 10 min with pre-

equilibrated alpha-MEM containing 2 µM propidium iodide (PI) and 1:1000 Hoechst (1

µg/mL final concentration). Negative control hydrogels were void of propidium iodide and

used for setting exposure limit during imaging. Hydrogels were imaged using an

epifluorescent microscope (Revolve) and the total number of cells and number of viable

cells were quantified using ImageJ after converting to binary images where PI/Hoechst-

positive cells were white, and background was black. Total number of cells positive for PI

were divided by total cells quantified by Hoechst staining to determine percentage of

unviable cells. Percentage of viable cells was determined by subtracting percentage of

unviable cells from 100. The experiment was repeated 4 times using 3 hydrogels per cell

density for each replicate.

*Bovine ovarian cells with follicle encapsulation and culture in PEG*

To test the impact of BOC and co-encapsulation on follicle viability and growth and

theca cell differentiation during in vitro culture, hydrogels were formed with either 1) only

follicles, 2) 5 x $10^6$ BOCs/mL only, or 3) follicles and 5 x $10^6$ BOCs/mL BOCs. Each

hydrogel contained 1-5 follicles (n = 65 follicles); 5 replicates). Hydrogel formation and

follicle encapsulation was performed as previously described with the addition of staining

follicles with the cell membrane marker PKH26 (2 μM final concentration; Sigma-Aldrich)

prior to encapsulation to improve visualization. Hydrogels were cultured for 10 days with

half of the medium changed every other day. Follicles were considered viable if the

basement membrane was attached, the follicles were void of vacuoles, and granulosa

cells were not darkened or mishappened. To recover cells for RT-qPCR, the hydrogels

were incubated in phenol red-free alpha-MEM with Liberase (1.3 wünsch units/mL) and

incubated at 37 °C and 5% $CO_2$ for 30 min with repeated pipetting to degrade the hydrogel

and extract the cells. After centrifugation, the cell pellet was used for RNA extraction,

cDNA synthesis and RT-qPCR as described below.

_Culture of bovine embryonic stem cells (bESCs)_

Bovine embryonic stem cells (n = 2 female lines) were routinely cultured feeder-

free in NBFR medium (N2B27 medium [1:1 DMEM/F12 and Neurobasal media, 0.5% v/v

N-2 supplement, 1% v/v B-27 supplement, 2 mM MEM Non-Essential Amino Acid

solution, 1% v/v GlutaMAX supplement, 0.1 mM 2-mercaptoethanol, 100 U/mL Penicillin,

and 100 μg/mL Streptomycin] supplemented with 1% bovine serum albumin (BSA), 20

ng/mL bFGF (100-18B, PeproTech), 2.5 μM IWR-1 (I0161, Sigma), and 20 ng/mL Activin

A (338-AC, R&D Systems)]. Cells were passaged onto vitronectin-coated plates every 2-

4 days using TrypleExpress at 1:4-1:6 split ratio and cultured in humidified incubators at

37 ºC and 5% $CO_2$. When cells were passaged, media was supplemented with 10 μm

ROCK inhibitor Y-27632. Media was changed daily and bESCs used for mesoderm

induction were between passage 18-26.

For mesoderm induction, bESCs were passaged onto 12-well vitronectin-coated

plates at 40,000 cells per well in mesoderm-induction media [GMEM medium, 15%

knockout serum replacement, 0.1 mM NEAA, 2mM GlutaMAX, 1mM sodium pyruvate,

0.1mM β-mercaptoethanol, 100 U/mL Penicillin, and 100 μg/mL streptomycin and

supplemented with 3 μm CHIR99021 and 70 ng/mL Activin A].  Cells were cultured in

mesoderm induction medium for 48 hours with daily medium change and thereafter the

cells were considered "mesoderm-like cells" (MeLCs) and used for subsequent

experiments.

Mesoderm-like cells were cultured in mesoderm induction medium without activin

A for an additional 72 hours and supplemented with 10 ng/mL bFGF beginning

immediately after MeLC induction, 24 hours after MeLC induction, 48 hours after MeLC

induction, or no supplementation (mesoderm medium only). Based on the finding of which

bFGF regimen best upregulated gene expression, in a second experiment, MeLCs were

cultured in mesoderm induction media without activin A, with 10 ng/mL bFGF and with

either 0, 1, 10, or 20 ng/mL of BMP4 for an additional 72 hours. All experiments contained

duplicate wells for each condition tested and had 3 biological replicates/independent

differentiations. Cells were imaged every day during each differentiation on the EVOS

(Thermo Fisher) inverted microscope.

*End-point (RT-PCR) and quantitative reverse transcription PCR (qRT-PCR)*

Freshly isolated follicles were pooled by stage and snap frozen in minimal PBS. To

examine the expression of transcripts for extracellular-matrix degrading enzymes,

primary follicles (n = 3 pools of 46-56 follicles per pool) and early secondary follicles (n =

3 pools of 34-40 follicles per pool) were subjected to RNA isolation using a protocol with

a combination of Trizol, Purelink DNase treatment, and Qiagen RNeasy Micro Kit. Final

total RNA was eluted into 14 µl of RNase-free water. Bovine ovarian cells (BOCs), bESCs,

MeLCs and differentiated cells were pelleted, snap frozen, and subjected to RNA isolation

following manual instructions using the Qiagen RNeasy Micro Kit. Maximum available

RNA mass was used for cDNA synthesis of primary and early secondary follicles and 250

ng - 1 µg of total RNA was used for cDNA synthesis of bESCs, BOCs, MeLCs, and

differentiated cells (RevertAid cDNA Reverse Transcription Kit). For RT-qPCR, 1-2 µl of

cDNA (50 ng total), 1x SsoAdvanced Universal SYBR Green Supermix (Bio-Rad,

Hercules, CA), and 250 nM of forward and reverse primers were mixed for 10 µl total

volume and pipetted into 96-well plates. The reaction was carried out in the CFX96 Touch

Deep Well Real-Time PCR Detection System (Bio-Rad). Cycling conditions were an initial

denaturation step at 95 ºC for 30 sec followed by 40 cycles of denaturation at 95 ºC for

10 sec, annealing at 60 ºC for 30 sec, and extension at 60 ºC for 5 sec. To confirm

specificity of PCR products for each gene, each assay included a melt curve analysis and

non-template control. Data was normalized to *ACTB* by the delta-Ct method and relative

expression data are represented as 1/ΔCT. For conventional PCR, PCR products from

quantitative PCR reactions were ran on a 2% agarose gel to visualize the PCR products.

For RT-qPCR experiments using *BAX* and *BCL2* primers, a standard curve of fresh

bovine ovarian cells was used to estimate absolute quantity of mRNA present in unknown

samples of BOCs. The ratio of *BAX/BCL2* was calculated from mRNA quantity of

respective genes and used to determine if cells were in a pro- or anti-apoptotic state. All

primers used in these experiments are listed in Table 1.

*Statistical analysis*

All data was first assessed for normality using a Shapiro-Wilk test. PEG growth

data was subjected to a two-way ANOVA using RStudio with effect of replicate included

in the model. Cell viability and RT-qPCR data were subjected to a one-way ANOVA or

Kruskal-Wallis test if data was not normally distributed using Graph Prism. Post-hoc tests

(Tukey's and Dunn's for ANOVA and Kruskal-Wallis, respectively) were conducted when

appropriate. RT-qPCR data were analyzed based on delta-Ct values. Comparisons were

considered significantly different when associated with $P < 0.05$.

RESULTS

To design a proteolytically-degradable PEG hydrogel that would most likely be

compatible with bovine preantral follicles, we first examined the gene expression of ECM-

degrading enzymes in primary and early secondary follicles that express the key follicle

marker *FSHR*. Matrix metalloproteinases (MMPs) and plasminogen activators are ECM-

degrading enzymes and play a pivotal role in tissue remodeling. We found the bovine

primary and early secondary follicles express *MMP2*, although in one pool we only found

*MMP2* in early secondary follicles. However, neither follicle stage expressed *MMP9*

(Figure 1A)*.* Plasminogen-activator urokinase (*PLAU*) was expressed in primary and early

secondary follicles from one pool, but not in the second or third pool. Similarly,

Plasminogen-activator tissue (*PLAT*) was expressed in primary follicles of one pool and

in early secondary follicles from two other pools (Figure 1A). Since we collectively found

mRNA expression of *MMP2* and plasminogen enzymes, we proceeded with using a PEG

hydrogel that is functionalized with VPMS (MMP-sensitive) and YKNS (plasminogen-
sensitive) peptide sequences (Figure 1B).

Next, we tested the hypothesis that bovine preantral follicles are more likely to
grow and demonstrate greater growth rate when encapsulated in PEG hydrogels
compared to 2D traditional methods after 10 days of culture. We found no difference in
the percentage of growing follicles between PEG (35.8% ± 6.23%) and 2D control (30.3%
± 7.88%) ($P$ = 0.56) (Figure 2A). We found that replicate (i.e. individual ovary) affected
the percentage of growing follicles ($P$ < 0.05) regardless of culture condition. It is worth
noting that in the control group, no follicle growth was observed in four out of the 15
replicates, whereas in the PEG group there was growth, although variable, in every
replicate (Figure 1A). Of the replicates where at least one follicle demonstrated growth in
either condition, we found no difference ($P$ = 0.50) in the growth rate from initial size (PEG:
18.8% ± 3.32%, control: 22.7% ± 3.49%) (Figure 2B). There was no effect of replicate on
growth rate ($P$ = 0.78).

We further examined the diameter changes between day 0, 5, and 10 in follicles
within a narrow range of starting diameters corresponding to their stage of development
between the primary to early secondary. When we examined follicles with starting
diameters corresponding to the early primary stage (40-60 μm), we found PEG hydrogels
outperformed the control conditions as day 5 follicles were larger in PEG compared to 2D
control ($P$ < 0.05). Moreover, by day 10, PEG hydrogel follicles had a significant increase
in diameter compared to day 0 ($P$ < 0.05), whereas control follicles showed no difference
($P$ = 0.72). Follicles starting at 61-70 μm showed a decrease in diameter by day 10 of
culture in control conditions ($P$ < 0.05), whereas follicles cultured in PEG maintained a

similar diameter over time ($P$ = 0.68; Figure 2C). Therefore, day 5 and 10 follicles were

significantly smaller ($P$ < 0.05) in control follicles when compared to similar timepoints of

PEG follicles. A representative image of a follicle grown in a PEG hydrogel over the 10-

day period is depicted in Fig. 2D.

We evaluated cell viability of BOCs when encapsulated in PEG hydrogels and

cultured for 10 days (Figure 3A-C). When encapsulated at 1 or 2 x $10^6$ cells/mL, BOCs

showed low cell viability (data not shown), whereas concentrations of 10 or 20 x $10^6$

cells/mL showed high viability (70-90%) and cell aggregation (Figure 3B and C).  In this

experiment, we found that neither control (no BOCs encapsulated) or hydrogels with

BOCs supported follicle growth over the 10 day-culture period (Figure 3D). Moreover,

gene expression analysis of pre-theca cell markers in the BOCs after 10 days of culture

revealed a decrease in expression of *PTCH1* ($P$ < 0.01) and *GLI1* ($P$ < 0.01) when

compared to day 0 fresh cells in both BOCs cultured alone and co-encapsulated with

follicles (Figure 3E). We found a decrease in both *BAX* and *BCL2* in BOCs that were

cultured with follicles and a decrease in *BCL2* in BOCs cultured in PEG hydrogels alone

($P$ < 0.05; Figure 3F). However, there was no difference in *BAX/BCL2* ratio, thus indicating

no difference in the pro- versus anti-apoptotic balance between cultured and fresh BOCs.

Given that native BOCs did not show maintenance of pre-theca identity or

contribute to follicle growth, we then explored the differentiation of bESCs into progenitor

cells of the bipotential gonad by testing effect of WNT, bFGF and BMP4 signaling (Figure

4A). We first tested the hypothesis that MeLCs exposed to a pulse of bFGF during

intermediate mesoderm/bipotential gonad differentiation would upregulate gene

expression of key markers at the end of induction (Figure 4B and C). Compared to bESCs

and 0, 24, 48, and 72 hours of 10 ng/mL bFGF exposure, MeLCs had higher expression

of the early lateral plate/intermediate mesoderm marker *LHX1* (Figure 4B; $P < 0.001$). We

also found the intermediate mesoderm marker *OSR1* to be upregulated in all bFGF

regimens compared to bESCs ($P < 0.01$). When analyzing expression of bipotential gonad

markers, there was no difference in *GATA4* ($P = 0.15$) across all experimental groups. We

also found higher expression of *WT1* after 0, 24, 48, and 96 hours of bFGF (i.e., after 4

additional days of culture) compared to bESCs and MeLCs ($P < 0.001$). There was no

difference in expression of *WT1* between bESCs and MeLCs ($P = 0.15$). There was also

no difference in expression of the paraxial mesoderm marker *PAX3* across all

experimental groups ($P = 0.06$), indicating induction conditions did not show strong

propensity to drive cells to the paraxial mesoderm fate. However, the lateral plate

mesoderm marker *FOXF1* was significantly upregulated compared to bESCs in 0, 24, and

48 hours of bFGF exposure ($P < 0.05$).

Because all bFGF regimens tested caused elevation in *WT1* and *OSR1*

expression, we chose to use a 24-hour pulse of bFGF in the next experiments when

testing the synergistic effects of including various concentrations of BMP4 over the same

culture period time and in two bESC lines (Figure 5A and B). Similar to the bFGF

experiment, we found the highest level of *LHX1* expression in MeLCs (Figure 5A; $P <$

0.0001). However, 10 and 20 ng/mL of BMP4 led to lower expression when compared to

bESCs ($P < 0.05$). Also confirming the previous results, *OSR1* was elevated in all

experimental groups, including MeLCs, when compared to bESCs ($P < 0.01$). Unlike in

the bFGF experiment where there was no difference amongst exposure times to bFGF,

we found an increase in *GATA4* in MeLCs and when 1, 10 or 20 ng/mL of BMP4 were

present ($P < 0.05$). *WT1* was upregulated in 0, 1, 10, or 20 ng/mL BMP4, but not in MeLCs

($P < 0.05$). Again, we found no difference in *PAX3* expression among all concentrations

of BMP4 when compared to bESCs. As expected, increasing the concentration of BMP4

led to a significant increase in *FOXF1* ($P < 0.05$), indicating a likely shift towards the lateral

plate mesoderm lineage with rising levels of BMP4 which is similar to in vivo development.

We further assessed WT1 and OCT4 expression in bESCs, MeLCs, and cells

cultured with 0 and 1 ng/mL BMP4 as these concentrations led to elevated *WT1* without

greatly upregulating *FOXF1* (Figure 5B). We found positive nuclear staining for OCT4 in

bESCs and MeLCs and no signal in cells exposed to 0 or 1 ng/mL BMP4, indicating likely

loss of pluripotency after mesoderm induction. We found nuclear localization of WT1 in

bESCs and nuclear and cytoplasmic localization in MeLCs (48 h culture) and 0 ng/mL

BMP4 (120 h culture). Interestingly, we did not observe WT1 expression at the protein

level when cells were exposed to 1 ng/mL BMP4.

DISCUSSION

To our knowledge, this study is the first to utilize a novel bioengineered PEG

hydrogel system for in vitro culture of bovine preantral follicles and assess whether

bESCs could be differentiated towards progenitor cells of the bipotential gonad using

stepwise in vitro culture protocols. The ultimate goal of this work is to develop novel

assisted reproductive technologies that enable the use of small preantral follicles in

conjunction with renewable theca-like cells for oocyte production. Since maintaining

structural integrity of the follicle is a main concern with 2D culture, hydrogels represent a

viable option as they provide framework to maintain the 3D configuration of follicles while

permitting growth. Indeed, the theory to build a culture system complete with extracellular

matrix components like the ovary has been suggested when early studies first explored

bovine preantral in vitro development [27]. Further adaptations to include sophisticated

ECM components in the hydrogel design have permitted mouse follicle growth in vitro

starting as early as the primordial stage [28]. Our finding that ECM-degrading enzymes

such as MMPs, PLAT and PLAU are expressed by bovine preantral follicles agrees with

previous investigations in follicles and spent medium [37,38]. However, the fact that these

transcripts were not uniformly expressed in all pools examined could indicate an intrinsic

variability in the ability of the follicles to remodel the ECM. This could explain at least in

part the variability observed in growth rate in the present study. Although we found that

PEG hydrogels outperformed 2D control over 10 days in primary follicles, there was a

notable difference in the percentage of follicles that grew between replicates (each

represented by one ovary), further underlining the biological variability between samples

and possibly an inherent propensity of some follicles for growth versus atresia. A limitation

of this study using slaughterhouse-derived oocytes is the inability to track factors known

to affect follicle viability such as animal age, nutritional and health status. Of note, early

secondary follicles showed a stagnant rate of growth compared to primary follicles

cultured in PEG hydrogels; this stagnation suggests that either 1) the 10 day-culture

conditions remain substandard for bovine preantral follicles or 2) follicles canonically slow

their growth at this stage of development which is a phenomenon seen in mice [29]. It is

worth noting that in vivo, 10 days represents a very narrow window within the estimated

180 days of folliculogenesis in the cow, and therefore relatively small changes in growth

could be expected in normally-developing follicles.

We tested the hypothesis of enhancing the bovine follicle culture system with

cellular supplementation as others have shown that the addition of ovarian stromal cells,

fibroblasts, and adipose-derived stem cells promote survival and growth of preantral

follicles in several species [13,17–19,30]. When BOCs were encapsulated in PEG

hydrogels and cultured, we found cell aggregates which may be due to absence of

basement membrane binding proteins in the PEG composition, rendering the cells unable

to adhere to the matrix and resulting in self-aggregation. Indeed, cell-extracellular matrix

interactions are important for mediating complex cellular responses such as migration,

proliferation, and differentiation [31], thus BOCs were likely incapable of integrating well

throughout the hydrogel. Furthermore, the finding of decreased expression of genes

known to be important for the pre-theca cell phenotype after culture shows that the BOCs

did not maintain their identity, which could have contributed to the lack of success of this

3D co-culture system. Others have reported that mouse ovarian stromal cells transition

from a predominately theca-like identity to almost entirely macrophage identity after 12

days of in vitro culture with follicles, partially corroborating our findings of theca cell loss

[32]. Since we found similarly high signs of atresia between follicles co-encapsulated and

follicles in PEG-only hydrogels, we pose the possibility that the prolonged protocol

required to dissociate ovarian tissue, isolate and label follicles, and process for

encapsulation may have contribute to decrease follicle viability beginning at day 0. In

future experiments, pre-theca cells could be sorted (pending availability of reliable

antibodies against specific theca cell surface markers) before incorporation into the PEG

hydrogel. Furthermore, culture conditions that specifically support theca cell phenotype

would likely need to be investigated.

In an attempt to produce gonadal-like cells for supplementation as pre-theca-like

cells in follicle culture, we employed bovine embryonic stem cells to recapitulate

mesoderm formation and early differentiation of the bipotential gonad with defined media

conditions. We show that inclusion of exogenous signaling factors such as the GSK3β

inhibitor CHIR 99021 for activation of WNT. WNT signaling can drive bESCs into

intermediate mesoderm-like cells that show early signs of coelomic epithelial-like

phenotype as indicated by gene and protein expression. During embryonic development,

the sexually dimorphic gonad (ovary or testis) must develop beginning from the

mesoderm germ layer, and then further differentiate into the intermediate mesoderm, then

coelomic epithelium which ingresses and gives rise to the sexually amorphic "bipotential

gonad" [33,34]. More specifically, upon Nodal, WNT and FGF signaling to induce

mesoderm differentiation [35], a subset of cells in the caudal aspect of the early

gastrulated embryo are specified to become precursors of the bipotential gonad.

Signaling by WNT, Nodal, and FGF during anterior-posterior (A-P) body axis extension,

which occurs at peri-gastrulation and is regulated by TBXT and other morphogens in the

mesoderm, drives cell fate decisions that result in caudal body axis formation [36,37].

Concomitantly, mediolateral segmentation occurs when a low, medium, and high gradient

of BMP signaling drives mesoderm cells into the paraxial, intermediate, and lateral plate

mesoderm domains, respectively [38,39]. Following this knowledge of developmental

biology, we chose to modulate WNT, Nodal, FGF and BMP signaling to mimic the

molecular events that drive cell differentiation for pre-gonad development during early

embryogenesis as shown in several mammalian species [34]. However, through these

experiments, we found that under WNT activation, bFGF and BMP4 are not essential to

induce the intermediate mesoderm or early coelomic epithelium. We observed peak

expression of the intermediate mesoderm marker *LHX1* after 48 h of mesoderm induction

under WNT and activin A, the latter a known inducer of *Lim1* (i.e. *Lhx1*) that has an

important role in IM specification/kidney development [40]. We also found *OSR1*

upregulated in all conditions after MeLC induction. *OSR1* is an early marker of IM and

others have shown its necessity for the urogenital phenotype [41–43] and generating

somatic cells of the gonad from PSCs [44]. Thus, we propose these cells may be reaching

the IM state with possible capability of becoming progenitor bipotential gonad-like.

Nevertheless, *OSR1* is not exclusive of the IM or gonads; kidney progenitor and lateral

plate mesoderm cells express *OSR1* [43,45] and this marker is used to demarcate kidney

progenitor-like cells during in vitro differentiation of PSCs [46–48]. Therefore, our

intermediate mesoderm/early coelomic epithelium-like cells should be further evaluated

for markers of early nephrogenesis.

We additionally found a rise in *WT1,* which is also an early, and more restrictive,

marker of gonadal progenitor cells [49], in induced cells across all bFGF and BMP4

regimens compared to bESCs. This indicates that continuous WNT activation for 3

additional days after initial mesoderm induction can provoke *WT1* expression. WT1 is

found along the A-P axis of mice as early as E9.0, with overlapping expression of GATA4

and NR5A1 in the coelomic epithelium by E10.0 and onward [49]. Moreover, WT1

activates the NR5A1 promoter [50], thus driving early gonad development. Interestingly,

*WT1* upregulation was accompanied by *GATA4* upregulation when 10 or 20 ng/mL of

BMP4 was included in culture which aligns with other studies [51]. However, we found

variable expression of the protein WT1 in cells cultured beyond MeLC induction and after

3 additional days under 0 or 1 ng/mL of BMP4. Although typically found in the nucleus,

WT1 isoforms with capability of shuttling between the nucleus and cytoplasm, therefore

we postulate various WT1 isoforms with nuclear and/or cytoplasmic activity may be

expressed in our cells [52]. We additionally highlight that there was no difference in *PAX3*

in all experiments, demonstrating that our culture conditions did not favor differentiation

of the paraxial mesoderm lineage. Interestingly, increasing BMP4 concentrations led to

increased expression of *FOXF1* (lateral plate mesoderm marker), which aligns with in

vivo studies showing a higher gradient of BMPs specifies the lateral plate of developing

embryos [39].

Our findings provide seminal context to previous work looking at differentiation of

ESCs from humans and mouse to gonad-like cells by using combinations of BMP4, FGF,

CHIR, and activin A [53–55]. Alternative approaches to inducing gonad-like cell formation

include overexpression of key transcription factors such as *SF1*, *WT1*, *RUNX1*, and

*GATA4* [56,57]. The efficiency of induction can be variable when using small molecules

or genetic engineering for differentiating stem cells, therefore heterogeneity in cell identity

would be expected in the present study. Although not done in this study, cell-sorting based

on expression of key markers could aid in purifying cell populations of interest and used

for downstream in vitro culture to achieve bipotential and even ovary or testis- like cells.

Yet, in vivo early somatic progenitor cells of the bipotential gonad are not specified

homogeneously, therefore some heterogeneity could be warranted, especially when

forming organoids as self-organizing structures. Finally, future studies should examine

early bovine embryonic development in vivo as findings from these studies could shed

light on the species-specific molecular mechanisms that drive gonad formation and inform

future in vitro culture conditions to recapitulate the process.

In conclusion, these findings provide a key step in expanding current

understanding of in vitro preantral folliculogenesis using a non-rodent model as well as

sheds light on methodologies in differentiating bESCs into precursor-like cells of the

ovary. The convergence of these two technologies will be pivotal to broadening the

knowledge base of follicle development and early gonadogenesis such that novel

assisted reproductive technologies can be expanded.

ACKNOWLEDGEMENTS

The authors acknowledge Stephanie McDonnell and Amanda Morton for aiding in follicle

collection and isolation. Funding: NIH 1F31HD111173, UC Davis Jastro Shields Graduate

Research Award, and Gary B. Anderson Food Animal Fellowship to JC; USDA-NIFA

W4112, 4171 multi-state groups. Some figures created with BioRender.com.

AUTHOR CONTRIBUTION

Conceptualization: Juliana Candelaria, Anna Denicol; Methodology: Juliana Candelaria,

Carly Guiltinan, Ramon Botigelli, Ariella Shikanov, Anna Denicol; Formal analysis and

investigation: Juliana Candelaria, Anna Denicol; Writing - original draft preparation:

Juliana Candelaria; Writing - review and editing: Juliana Candelaria, Anna Denicol, Ariella

Shikanov, Carly Guiltinan, Ramon Botigelli,; Funding acquisition: Juliana Candelaria,

Anna Denicol; Resources: Anna Denicol, Ariella Shikanov; Supervision: Anna Denicol.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Bus A, Langbeen A, Martin B, Leroy JLMR, Bols PEJ. Is the pre-antral ovarian follicle
the 'holy grail'for female fertility preservation? Anim Reprod Sci [Internet]. 2019;207:119–
30. Available from: http://www.sciencedirect.com/science/article/pii/S0378432019302064

2. Gutierrez CG, Ralph JH, Telfer EE, Wilmut I, Webb R. Growth and Antrum Formation
of Bovine Preantral Follicles in Long-Term Culture In Vitro1. Biol Reprod [Internet].
2000;62:1322–8. Available from: https://doi.org/10.1095/biolreprod62.5.1322

3. Araújo VR, Gastal MO, Wischral A, Figueiredo JR, Gastal EL. Long-term in vitro culture
of bovine preantral follicles: Effect of base medium and medium replacement methods.
Anim Reprod Sci [Internet]. 2015;161:23–31. Available from:
http://www.sciencedirect.com/science/article/pii/S0378432015001736

4. Rossetto R, Saraiva MVA, Bernuci MP, Silva GM, Brito IR, Alves A, et al. Impact of
insulin concentration and mode of FSH addition on the in vitro survival and development
of isolated bovine preantral follicles. Theriogenology [Internet]. 2016/05/22.
2016;86:1137–45. Available from: https://www.ncbi.nlm.nih.gov/pubmed/27207475

5. Gomes JE, Correia SC, Gouveia-Oliveira A, Cidadão AJ, Plancha CE. Three-
dimensional environments preserve extracellular matrix compartments of ovarian follicles
and increase FSH-dependent growth. Mol Reprod Dev. 1999;54:163–72.

6. Belli M, Vigone G, Merico V, Redi CA, Zuccotti M, Garagna S. Towards a 3D culture of
mouse ovarian follicles. International Journal of Developmental Biology. 2013;56:931–7.

7. Xu M, West-Farrell ER, Stouffer RL, Shea LD, Woodruff TK, Zelinski MB. Encapsulated
Three-Dimensional Culture Supports Development of Nonhuman Primate Secondary
Follicles1. Biol Reprod. 2009;81:587–94.

8. Xu J, Lawson MS, Yeoman RR, Molskness TA, Ting AY, Stouffer RL, et al. Fibrin
promotes development and function of macaque primary follicles during encapsulated
three-dimensional culture. Hum Reprod [Internet]. 2013/04/21. 2013;28:2187–200.
Available from: https://pubmed.ncbi.nlm.nih.gov/23608357

9. Vanacker J, Luyckx V, Dolmans MM, Des Rieux A, Jaeger J, Van Langendonckt A, et
al. Transplantation of an alginate–matrigel matrix containing isolated ovarian cells: First
step in developing a biodegradable scaffold to transplant isolated preantral follicles and
ovarian cells. Biomaterials. 2012;33:6079–85.

10. Ihm JE, Lee ST, Han DK, Lim JM, Hubbell JA. Murine ovarian follicle culture in PEG-
hydrogel: Effects of mechanical properties and the hormones FSH and LH on
development. Macromol Res [Internet]. 2015;23:377–86. Available from:
https://doi.org/10.1007/s13233-015-3045-x

11. Mendez U, Zhou H, Shikanov A. Synthetic PEG Hydrogel for Engineering the
Environment of Ovarian Follicles. In: Chawla K, editor. Methods in Molecular Biology
[Internet]. New York, NY: Springer New York; 2018. p. 115–28. Available from:
https://doi.org/10.1007/978-1-4939-7741-3_9

12. Shikanov A, Smith RM, Xu M, Woodruff TK, Shea LD. Hydrogel network design using
multifunctional macromers to coordinate tissue maturation in ovarian follicle culture.
Biomaterials [Internet]. 2011;32:2524–31. Available from:
http://dx.doi.org/10.1016/j.biomaterials.2010.12.027

13. Tomaszewski CE, Constance E, Lemke MM, Zhou H, Padmanabhan V, Arnold KB, et
al. Adipose-derived stem cell-secreted factors promote early stage follicle development
in a biomimetic matrix. Biomater Sci. 2019;7:571–80.

14. Wijgerde M, Ooms M, Hoogerbrugge JW, Grootegoed JA. Hedgehog signaling in
mouse ovary: Indian hedgehog and desert hedgehog from granulosa cells induce target
gene expression in developing theca cells. Endocrinology. 2005;146:3558–66.

15. Cowan RG, Quirk SM. Cells responding to hedgehog signaling contribute to the theca
of ovarian follicles. Reproduction. 2021;161:437–48.

16. Russell MC, Cowan RG, Harman RM, Walker AL, Quirk SM. The hedgehog signaling
pathway in the mouse ovary. Biol Reprod. 2007;77:226–36.

17. Grubliauskaitė M, Vlieghe H, Moghassemi S, Dadashzadeh A, Camboni A,
Gudlevičienė Ž, et al. Influence of ovarian stromal cells on human ovarian follicle growth
in a 3D environment. Hum Reprod Open. 2023;2023.

18. Tingen CM, Kiesewetter SE, Jozefik J, Thomas C, Tagler D, Shea L, et al. A
macrophage and theca cell-enriched stromal cell population influences growth and
survival of immature murine follicles in vitro. Reproduction. 2011;141:809–20.

19. Tagler D, Tu T, Smith RM, Anderson NR, Tingen CM, Woodruff TK, et al. Embryonic
Fibroblasts Enable the Culture of Primary Ovarian Follicles Within Alginate Hydrogels.
Tissue Eng Part A [Internet]. 2012;18:1229–38. Available from:
https://doi.org/10.1089/ten.tea.2011.0418

20. Jadhav U, Jameson JL. Steroidogenic factor-1 (SF-1)-driven differentiation of murine
embryonic stem (ES) cells into a gonadal lineage. Endocrinology. 2011;152:2870–82.

21. Lan C-W, Chen M-J, Jan P-S, Chen H-F, Ho H-N. Differentiation of Human Embryonic
Stem Cells Into Functional Ovarian Granulosa-like Cells. J Clin Endocrinol Metab
[Internet]. 2013;98:3713–23. Available from: https://doi.org/10.1210/jc.2012-4302

22. Rodríguez Gutiérrez D, Eid W, Biason-Lauber A. A Human Gonadal Cell Model From
Induced Pluripotent Stem Cells. Front Genet [Internet]. 2018;9. Available from:
https://www.frontiersin.org/articles/10.3389/fgene.2018.00498

23. Knarston IM, Pachernegg S, Robevska G, Ghobrial I, Er PX, Georges E, et al. An In
Vitro Differentiation Protocol for Human Embryonic Bipotential Gonad and Testis Cell
Development. Stem Cell Reports [Internet]. 2020;15:1377–91. Available from:
https://doi.org/10.1016/j.stemcr.2020.10.009

24. Sepponen K, Lundin K, Knuus K, Väyrynen P, Raivio T, Tapanainen JS, et al. The role
of sequential BMP signaling in directing human embryonic stem cells to bipotential
gonadal cells. Journal of Clinical Endocrinology and Metabolism. 2017;102:4303–14.

25. Danti L, Lundin K, Sepponen K, Yohannes DA, Kere J, Tuuri T, et al. CRISPR/Cas9-
mediated activation of NR5A1 steers female human embryonic stem cell-derived

bipotential gonadal-like cells towards a steroidogenic cell fate. J Ovarian Res [Internet].
2023;16:194. Available from: https://doi.org/10.1186/s13048-023-01264-5

26. McDonnell SP, Candelaria JI, Morton AJ, Denicol AC. Isolation of Small Preantral
Follicles from the Bovine Ovary Using a Combination of Fragmentation, Homogenization,
and Serial Filtration. JoVE [Internet]. 2022; Available from: https://www.jove.com/64423

27. Figueiredo JR de, Hulshof SCJ, Thiry M, Van Den Hurk R, Bevers MM, Nusgens B,
et al. Extracellular matrix proteins and basement membrane: Their identification in bovine
ovaries and significance for the attachment of cultured preantral follicles. Theriogenology.
1995;43:845–58.

28. Nason-Tomaszewski CE, Thomas EE, Matera DL, Baker BM, Shikanov A.
Extracellular matrix-templating fibrous hydrogels promote ovarian tissue remodeling and
oocyte growth. Bioact Mater [Internet]. 2024;32:292–303. Available from:
https://doi.org/10.1016/j.bioactmat.2023.10.001

29. Richard S, Anderson NJ, Zhou Y, Pankhurst MW. Mouse primary follicles experience
slow growth rates after activation and progressive increases that influence the duration
of the primary follicle phase. Biol Reprod. 2023;109:684–92.

30. Green LJ, Zhou H, Padmanabhan V, Shikanov A. Adipose-derived stem cells promote
survival, growth, and maturation of early-stage murine follicles. Stem Cell Res Ther.
2019;10:1–13.

31. Hynes RO. Stretching the boundaries of extracellular matrix research. Nat Rev Mol
Cell Biol [Internet]. 2014;15:761–3. Available from: https://doi.org/10.1038/nrm3908

32. Tingen CM, Kiesewetter SE, Jozefik J, Thomas C, Tagler D, Shea L, et al. A
macrophage and theca cell-enriched stromal cell population influences growth and
survival of immature murine follicles in vitro. Reproduction [Internet]. 2011/03/09.
2011;141:809–20. Available from: https://pubmed.ncbi.nlm.nih.gov/21389078

33. Karl J, Capel B. Sertoli cells of the mouse testis originate from the coelomic
epithelium. Dev Biol. 1998;203:323–33.

34. Rotgers E, Jørgensen A, Yao HHC. At the crossroads of fate-Somatic cell lineage
specification in the fetal gonad. Endocr Rev. 2018;39:739–59.

35. Ciruna B, Rossant J. FGF signaling regulates mesoderm cell fate specification and
morphogenetic movement at the primitive streak. Dev Cell. 2001;1:37–49.

36. Amin S, Neijts R, Simmini S, van Rooijen C, Tan SC, Kester L, et al. Cdx and T
Brachyury Co-activate Growth Signaling in the Embryonic Axial Progenitor Niche. Cell
Rep [Internet]. 2016;17:3165–77. Available from:
https://doi.org/10.1016/j.celrep.2016.11.069

37. Brennan J, Lu CC, Norris DP, Rodriguez TA, Beddington RSP, Robertson EJ. Nodal
signalling in the epiblast patterns the early mouse embryo. Nature. 2001;411:965–9.

38. James RG, Schultheiss TM. Bmp signaling promotes intermediate mesoderm gene
expression in a dose-dependent, cell-autonomous and translation-dependent manner.
Dev Biol. 2005;288:113–25.

39. Niehrs C, Dosch R, Onichtchouk D. Embryonic patterning of Xenopus mesoderm by
Bmp-4. Of Fish, Fly, Worm, and Man: Lessons from Developmental Biology for Human
Gene Function and Disease. 2000;165–90.

40. Preger-Ben Noon E, Barak H, Guttmann-Raviv N, Reshef R. Interplay between activin
and Hox genes determines the formation of the kidney morphogenetic field. Development
[Internet]. 2009;136:1995–2004. Available from: https://doi.org/10.1242/dev.035592

41. Wang Q, Lan Y, Cho ES, Maltby KM, Jiang R. Odd-skipped related 1 (Odd1) is an
essential regulator of heart and urogenital development. Dev Biol. 2005;288:582–94.

42. Mae S-I, Shono A, Shiota F, Yasuno T, Kajiwara M, Gotoda-Nishimura N, et al.
Monitoring and robust induction of nephrogenic intermediate mesoderm from human
pluripotent stem cells. Nat Commun [Internet]. 2013;4:1367. Available from:
https://doi.org/10.1038/ncomms2378

43. James RG, Kamei CN, Wang Q, Jiang R, Schultheiss TM. Odd-skipped related 1 is
required for development of the metanephric kidney and regulates formation and
differentiation of kidney precursor cells. Development. 2006;133:2995–3004.

44. Yoshino T, Suzuki T, Nagamatsu G, Yabukami H, Ikegaya M, Kishima M, et al.
Generation of ovarian follicles from mouse pluripotent stem cells. Science (1979).
2021;373.

45. Xu J, Liu H, Park J-S, Lan Y, Jiang R. Osr1 acts downstream of and interacts
synergistically with Six2 to maintain nephron progenitor cells during kidney
organogenesis. Development [Internet]. 2014;141:1442–52. Available from:
https://doi.org/10.1242/dev.103283

46. Araoka T, Mae S, Kurose Y, Uesugi M, Ohta A, Yamanaka S, et al. Efficient and Rapid
Induction of Human iPSCs/ESCs into Nephrogenic Intermediate Mesoderm Using Small
Molecule-Based Differentiation Methods. PLoS One [Internet]. 2014;9:e84881. Available
from: https://doi.org/10.1371/journal.pone.0084881

47. Morizane R, Lam AQ, Freedman BS, Kishi S, Valerius MT, Bonventre J V. Nephron
organoids derived from human pluripotent stem cells model kidney development and
injury. Nat Biotechnol [Internet]. 2015;33:1193–200. Available from:
https://doi.org/10.1038/nbt.3392

48. Takasato M, Er PX, Becroft M, Vanslambrouck JM, Stanley EG, Elefanty AG, et al.
Directing human embryonic stem cell differentiation towards a renal lineage generates a
self-organizing kidney. Nat Cell Biol [Internet]. 2014;16:118–26. Available from:
https://doi.org/10.1038/ncb2894

49. Sasaki K, Oguchi A, Cheng K, Murakawa Y, Okamoto I, Ohta H, et al. The embryonic
ontogeny of the gonadal somatic cells in mice and monkeys. Cell Rep [Internet].
2021;35:109075. Available from: https://doi.org/10.1016/j.celrep.2021.109075

50. Wilhelm D, Englert C. The Wilms tumor suppressor WT1 regulates early gonad
development by activation of Sf1. Genes Dev. 2002;16:1839–51.

51. Rojas A, De Val S, Heidt AB, Xu S-M, Bristow J, Black BL. Gata4 expression in lateral
mesoderm is downstream of BMP4 and is activated directly by Forkhead and GATA

transcription factors through a distal enhancer element. Development [Internet].
2005;132:3405–17. Available from: https://doi.org/10.1242/dev.01913

52. Niksic M, Slight J, Sanford JR, Caceres JF, Hastie ND. The Wilms' tumour protein
(WT1) shuttles between nucleus and cytoplasm and is present in functional polysomes.
Hum Mol Genet [Internet]. 2004;13:463–71. Available from:
https://doi.org/10.1093/hmg/ddh040

53. Gonen N, Eozenou C, Mitter R, Elzaiat M, Stévant I, Aviram R, et al. In vitro cellular
reprogramming to model gonad development and its disorders. Sci Adv. 2023;9.

54. Rore H, Owen N, Piña-Aguilar RE, Docherty K, Sekido R. Testicular somatic cell-like
cells derived from embryonic stem cells induce differentiation of epiblasts into germ cells.
Commun Biol [Internet]. 2021;4. Available from: http://dx.doi.org/10.1038/s42003-021-
02322-8

55. Shin E-Y, Park S, Choi WY, Lee DR. Rapid Differentiation of Human Embryonic Stem
Cells into Testosterone-Producing Leydig Cell-Like Cells In vitro. Tissue Eng Regen Med
[Internet]. 2021;18:651–62. Available from: https://doi.org/10.1007/s13770-021-00359-8

56. Xu C, Mohsin A, Luo Y, Xie L, Peng Y, Wang Q, et al. Differentiation roadmap of
embryonic Sertoli cells derived from mouse embryonic stem cells. Stem Cell Res Ther.
2019;10:1–12.

57. Smela MDP, Kramme CC, Fortuna PRJ, Adams JL, Su R, Dong E, et al. Directed
differentiation of human iPSCs to functional ovarian granulosa-like cells via transcription
factor overexpression. Elife. 2023;12.

**Table 1** Primer list.

| Gene | Forward Sequence (5'-3') | Reverse Sequence (5'-3') | Product size (base pairs) |
|---|---|---|---|
| *BAX* | CATGAGGTCAACCCTCGGGA | TAG GGA AGC CCA TGCTGAGT | 117 |
| *BCL2* | GAGTTCGGAGGGGTCATGTG | GGG CCA TAC AGC TCCACAAA | 158 |
| *CSPG4* | CTGGAGGAACAA AGGTCTCTGG | ACTGCGTGATCTGGAAAAGCA | 140 |
| *DHH* | GCATGCATTGCAGGTCGTC | ACGTGGGCAGAATGCTAGTC | 145 |
| *FOXF1* | GCCTCCTACATCAAGCAGCA | GTTCTGGTGCAGATACGGCT | 108 |
| *GATA4* | TGCGCCCCATCAAAACAGAG | GGGCCATACAGCTCCACAAA | 77 |
| *GLI1* | CCTTCAAAGCCCAGTACATGC | GTGCGTCTTCAGATTTTCCAG | 122 |
| *IHH* | CGGCTTCGACTGGGTGTATT | GGCTGAGTGCTCGGACTTG | 70 |
| *LHX1* | TGGACCGCTTCCTCTTGAAC | CTTGGTACCGAAACACCGGA | 152 |
| *MMP2* | ATCGTCTTCGACGGC ATCTC | TTCGCCAGATGAATCGGTCC | 70 |
| *MMP9* | CTTCACCTTTGAGGGTCGCT | TGGGTGTAGAGTCTCTCGCT | 138 |
| *OSR1* | CCTGTGGAAATCCAGCGCC | CAGTCTGTTCTGACTTCCTGCTTA | 111 |
| *PAX3* | CCTTCAAAGCCCAGTACATGC | GTGCGTCTTCAGATTTTCCAG | 122 |
| *PLAT* | TCGGCAGGCCCTCTATTCTT | ACGTGGCGGTAGCATCTATTT | 89 |
| *PLAU* | TGGGTCAGTCACGGCTTAAC | ACGGATCTTCAGCAAGGCAAT | 131 |
| *WT1* | CCTTCACCGTCCACTTCTCT | CCGTGCTGTATCCCTGGTTG | 183 |
| *H2A.2* | GAGGAGCTGAACAAGCTGTTG | TTGTGGTGGCTCTCAGTCTTC | 104 |
| *ACTB* | CTCTTCCAGCCTTCCTTCCT | GGGCAGTGATCTCTTTCTGC | 178 |

696
**Fig 1** Bovine preantral follicles express extracellular matrix-degrading enzymes, allowing
the design of a PEG hydrogel containing target peptides for controlled degradation. (**A**)
Agarose gel with products of RT-PCR of primary and early secondary follicles expressing
mRNA for *MMP2, MMP9, PLAU, PLAT, FSHR,* and *H2A*. Each pool of primary and early
secondary follicles (n = 3 pools per stage) contained 46-56 and 34-40 follicles,

respectively. (**B**) Schematic of PEG hydrogel for follicle encapsulation. PEG =
poly(ethylene glycol). MMP = matrix metalloproteinase
704

**Fig 2** Bovine preantral follicle in vitro growth using PEG or 2D control culture systems.
Percentage of preantral follicles that grew (**A**) and percent change in diameter from
follicles that grew (**B**). (**C**) Bovine preantral follicle diameter growth during in vitro culture
at indicated starting diameters/stages. Lowercases letters that differ = significant
difference ($P < 0.05$) between treatment groups in the same day. Uppercase letters that
differ = significant difference ($P < 0.05$) within the same treatment group but compared to
day 0. Data are presented as mean $\pm$ SEM. (**D**) Representative images of follicle grown
in PEG over 10-day culture period. Scale bar = 100 µm.

715
**Fig 3** Co-culture of bovine preantral follicles with bovine ovarian cells (BOCs) in PEG
hydrogels. (**A**) Representative image of BOCs in PEG hydrogel (scale bar = 500 µm). (**B**)
Viability of BOCs encapsulated in PEG hydrogels during in vitro culture. (**C**) Propidium

iodine (PI) staining in BOCs encapsulated in PEG hydrogels at day 0 and day 10 of
culture. Counterstained with DAPI (TL = transluminescent; scale bar = 100 μm). (**D**)
Unviable preantral follicle encapsulated in PEG hydrogel with or without BOCs using
PKH26 to label follicles at day 10 of culture (scale bar = 100 μm). (**E**) Gene expression of
pre-theca cell markers in BOCs (relative expression = $1/\Delta CT$). (**F**) Anti- and pro-apoptotic
gene marker expression in BOCs (level of expression = absolute quantification from
standard curve). Data are presented as mean $\pm$ SEM.
726

728
729
730

**Fig 4** In vitro differentiation of bovine embryonic stem cells (bESCs) towards progenitors
of somatic bipotential gonad-like cells. (**A**) Schematic diagram of experimental design for
bFGF and BMP4 experiments. (**B**) Gene expression of intermediate mesoderm,
bipotential gonad, paraxial and lateral plate mesoderm markers before and after bESC
differentiation using bFGF at various days of exposure. (**C**) Representative images of
cells from each treatment group. Data are presented as mean $\pm$ SEM. Significant
differences are noted by * = $P$ <0.05, ** = $P$ <0.01, *** = $P$ <0.001, **** = $P$ <0.0001.
738

740
741
**Fig 5** In vitro differentiation of bovine embryonic stem cells (bESCs) towards progenitors
of somatic bipotential gonad-like cells testing BMP4 concentrations. (**A**) Gene expression
of intermediate mesoderm, bipotential gonad, paraxial and lateral plate mesoderm
markers before and after bESC differentiation (**B**) Representative images of cells from
each treatment group. Data are presented as mean ± SEM. Significant differences are
noted by * = $P$ <0.05, ** = $P$ <0.01, *** = $P$ <0.001, **** = $P$ <0.0001.