# Spatio-temporal dynamics of attacks around deaths of wolves: A statistical assessment of lethal control efficiency in France

Oksana Grente*[1,3], Thomas Opitz[2], Christophe Duchamp[3], Nolwenn Drouet-Hoguet[4], Simon Chamaillé-Jammes[1], Olivier Gimenez[1]

[1] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
[2] Biostatistics and Spatial Processes, INRAE, 228 Route de l'Aérodrome, Avignon, 84914, France
[3] Office Français de la Biodiversité, Direction de la Recherche et Appui Scientifique, Micropolis La Bérardie 05000 Gap, France
[4] Office Français de la Biodiversité, Direction de la Recherche et Appui Scientifique, ZI Mayencin 38610 Gières, France

*Corresponding author
Correspondence: oksana.grente@gmail.com

## Abstract

The lethal control of large carnivores is criticized regarding its efficiency to prevent hotspots of attacks on livestock. Previous studies, mainly focused on North America, provided mixed results. We evaluated the effects of wolf lethal removals on the distribution of attack intensities in the French Alps between 2011 and 2020, using a Before After Control-Impact approach with retrospective data. We built an original framework combining both continuous spatial and temporal scales and a 3D kernel estimation. We compared the attack intensities observed before and after the legal killings of wolves on a period of 90 days and on a range of 10 km, and with control situations where no removal occurred. The analysis was corrected for the presence of livestock. A moderate decrease of attack intensity was the most common outcome after the lethal removal of a single wolf. This reduction was greatly amplified when removing two or three wolves. The scale of analysis also modulated this general pattern, with decreases being generally amplified at a small spatio-temporal range. Contextual factors (*e.g.*, geographical or seasonal variations) could also lead to deviations from this general pattern. Overall, between 2011 and 2020, lethal control of wolves in France generally contributed to reduce livestock attacks, but mainly locally and to a minor extent. Our results highlight the importance of accounting for scale in such assessments and suggest that the evaluation of the effectiveness of lethal removals in reducing livestock predation might be more relevant in a local context. As a next step, we recommend to move forward from patterns to mechanisms by linking the effects of lethal control on wolves to their effects on attacks through analysis of fine-scaled data on wolves and livestock.

*Keywords:* Before-After Control Impact, *Canis lupus*, culling, depredation, French Alps, kernel density estimation, removal, sheep

## 1. Introduction

Conservation of large carnivores while limiting their damages on livestock is a difficult task worldwide (van Eeden et al., 2018). Predation on livestock by large carnivores has been the main fuel of hostility towards them (Lute et al., 2018; van Eeden et al., 2018). Strategies of eradication led to their local extirpations in some cases (Treves and Karanth, 2003), like in many areas across Europe (Chapron et al., 2014). The growing promotion of biodiversity conservation these last decades has reversed the trend, with some populations of large carnivores now expanding in Europe, though at unequal paces (Kaczensky et al., 2024). Diverse measures have been promoted and implemented with the objective of mitigating or limiting predations, whether by increasing acceptance through compensations (Gervasi et al., 2021), by decreasing livestock vulnerability through changes in husbandry practices, or by directly affecting carnivores (van Eeden et al., 2018).

Lethal control, one of the most controversial measures targeting carnivores, aims to decrease carnivore density and so the encounter risk with livestock, or to remove specific individuals inclined to predation on livestock, while maintaining their conservation status as favorable (Lennox et al., 2018). Lethal control has been widely applied worldwide, and can be exclusively directed towards carnivores attacking livestock or be part of a more general harvest (Lorand et al., 2022; Treves and Karanth, 2003).

The side effects of lethal control are often claimed to undermine its efficiency or even to create more damages (Elbroch and Treves, 2023). In the specific case of social canids like wolves, studies have mostly pointed out additive human-caused breeder loss as a risk for increasing damages on livestock  (Jędrzejewska et al., 1996; Žunna et al., 2023). Social instability can indeed cause multiple reproduction (Ausband et al., 2017b) or pack disruption (Cassidy et al., 2023), although their effects on damages have not been assessed.

Implementing randomized controlled experiments to evaluate the efficiency of lethal control of large carnivores is extremely difficult, considering the involved large spatial and temporal scales. To date, no experimental study has been conducted for this matter (Treves et al., 2016, 2019). Only Allen (2014) conducted a controlled experiment to evaluate dingo's baiting efficiency but using the number of lactation failures as a proxy for calves lost to predation instead of direct cases of predation.

Also, retrospective studies have been conducted to answer the question of lethal control efficiency. The data they used were not collected within a scientific frame but generally through management (*e.g.* damages from compensation schemes). Most of retrospective studies on social canids (gray wolves *Canis lupus*, dingoes *Dingo dingo*, coyotes *Canis latrans*) were correlative and tested for a relationship between the number of livestock damages and the number of killed predators (Šuba et al., 2023; Fernández-Gil et al., 2016; DeCesare et al., 2018; Conner et al., 1998; Allen, 2015; Kutal et al., 2023; Merz et al., 2025). However, the ability of this approach to infer causality is being debated (Arif and MacNeil, 2022; Nichols and Cooch, 2025). The approach is also subject to analytical variability (Gould et al., 2025). For example, three correlative studies which used the same dataset concluded differently on the effect of lethal removals of wolves on livestock damages (Kompaniyets and Evans, 2017; Poudyal et al., 2016; Wielgus and Peebles, 2014).

A smaller number of retrospective studies used a comparative approach. Harper et al. (2008), Bradley et al. (2015), Santiago-Avila et al. (2018) and Wagner and Conover (1999) compared situations where lethal control was used and situations where it was not (*i.e.* inter-site comparison). Bjorge and Gunson (1985) and Blejwas et al. (2002) compared the situation before and after the

implementation of lethal control within the same site (*i.e.* intra-site comparison). While the inter-site
comparison is more susceptible to possible spatial confounding factors but minimizes temporal
confounding factors, the opposite is true for the intra-site comparison. Kutal et al. (2023) was the
only study to mix inter-site and intra-site comparisons, by combining comparisons before and after
hunting season and between sites with and without hunted wolves (*i.e.* Before-After Control-Impact,
BACI). The BACI study design showed a better performance to estimate the effect of an impact in a
simulated ecological dataset than intra-site or inter-site comparisons (Christie et al., 2019), and
performed even better than the randomized controlled experiment still considered as gold-standard
of scientific inference (Treves et al., 2019).

Five of these seven comparative studies on social canids showed that lethal control significantly
decreased damage recurrence, whereas Santiago-Avila et al. (2018) and Kutal et al. (2023) found no
significant effect of lethal control on damages. Yet, results suggested variability in the efficiency of
lethal control. For example, Harper et al. (2008) detected that only the death of an adult wolf male
was significantly efficient, whereas Bradley et al. (2015) did not detect sex or age conditions on
efficiency. Variability could also result from the study designs themselves, as these studies used very
different spatial and temporal scales to evaluate lethal control efficiency. Thus, small scale (*e.g.* 4 km
in Harper et al. (2008)), medium scale (*e.g.* pack territory in Bradley et al. (2015)) or large scale (*e.g.*
711 km² in Kutal et al. (2023)) could be used for space, sometimes within the same study (*e.g.*
Santiago-Avila et al., 2018). Time scale could also be disparate, going from several days or months
(*e.g.* Blejwas et al. (2002)) to one year (*e.g.* Kutal et al., (2023)) or more (*e.g.* Bjorge and Gunson
(1985)). Most importantly, none of the mentioned retrospective studies have both used continuous
spatial and temporal scales, making it difficult to identify potential gradients of lethal control effects,
as those suggested in Santiago-Avila et al. (2018). Finally, only two of the seven comparative studies
have controlled for livestock presence or abundance (Blejwas et al., 2002; Wagner and Conover,
1999), yet it is the primary factor structuring damage risk and the highest risk of confounding factor.

The aim of our study was to evaluate the effects of the lethal control applied for gray wolves in
France on the intensity of attacks on sheep. We used the data collected by the French
administrations for the period 2011-2020. Given the retrospective nature of our study, we
attempted to fill the gaps identified in the previously mentioned retrospective studies on the
subject. For that purpose, we used the study design of BACI, used continuous spatial and temporal
study scales and corrected for livestock presence through an original framework of kernel
estimation.

**2. Methods**

**2.1 Management context and study area**

After historical campaigns of destruction, gray wolves disappeared from France at the beginning of
the twentieth century, despite being present on the whole metropolitan territory at the end of the
eighteenth century (de Beaufort, 1987). From the 1970s, the combination of protective legislation
and the increase of forested habitat allowed the recovery of the species in Europe (Chapron et al.,
2014). In the 1990s, wolves from the Apennines population in Central Italy dispersed and settled into
the south-western Alps, straddling Italy and France (Linnell and Cretois, 2018; Fabbri et al., 2007;
Poulle et al., 1997). From there, a wolf population progressively established in the area. On the
French side, the population has expanded but has until now remained mainly restricted to the south-
eastern corner of France, except for a few reproductive packs outside of this area (Louvrier et al.,
2018). The spatial growth reflected a demographic growth that has been exponential. In the first
twenty years of the recovery, the population size progressively increased and reached approximately

200 estimated individuals in 2013 (Réseau Loup-Lynx, 2024). Then, the growth rate accelerated, and
the population more than doubled in five years, with more than 500 estimated individuals in 2018
(Office National de la Chasse et de la Faune Sauvage, 2019). This sustained growth has continued at
least up to 2022, with a population size approaching 800 estimated individuals (Milleret et al.,
2025a).

South-eastern France is characterized by a variety of landscapes, including high altitude mountains,
Piedmont plains, hills and garrigue lands, where mountain and Mediterranean climates converge.
Livestock farming is generally extensive. Herds of cow are principally present in the north of the
region, while those of sheep are present in the whole area (Dobremez et al., 2016). The pastoral
practice of transhumance, consisting in moving all or part of herds from low to high altitude pastures
during summer and vice versa during winter, is still important. Duration of grazing season increases
following a north-south gradient, with some herds from low altitudes in the south grazing outside all
year round (Dobremez et al., 2016). Livestock owners can share pastures, and even gather their
herds to limit the cost of transport for transhumance and of herding. In the Alps, herds of cows are
generally not mixed with flocks of sheep, but the latter can be occasionally mixed with goats. Mean
size of transhumant flocks of sheep varies greatly between alpine departments and ranges between
less than 250 sheep up to 2000 sheep (Agreste, 2021).

To prevent attacks, the authorities subsidize non-lethal measures for protecting flocks of sheep or
goat through livestock guarding dogs, fences and additional shepherds. In situations where attacks
persist despite non-lethal measures, the authorities have authorized lethal removals of wolves by
ground shooting since 2004, based on the article 12 of the Directive Habitats. Other forms of
removals (*e.g.* trapping, aerial shooting) are prohibited. Conditions of application of ground shooting
are defined by ministerial decrees (*e.g.* Ministère de la Transition Ecologique et Solidaire, 2018) and
depend on the implementation of non-lethal protective measures and on the recurrence of attacks.
If the conditions are met, the administration assigns the derogation to the livestock owner for a
duration between two months and five years. The owner itself or a mandated hunter can shoot
wolves during an attack on its livestock (hereafter 'simple defense'). When attacks still remain,
governmental agents or special volunteers who are trained may support owners in protecting their
livestock (hereafter 'reinforced defense'). Eventually, if the level of attacks escalates in an area, the
administration can decide to shoot a specific number of wolves under game hunting conditions
(hereafter 'hunting'). Between 2004 and 2018, the number of wolves that could be annually killed if
required was prevented from exceeding between 10 and 12% of the annual estimated winter wolf
population size, in order to avoid threatening the recovery of the species. Throughout this period,
the number of killed wolves followed the increases of both the wolf population size and the attack
occurrences, and from 2011, wolves have been legally killed every year (Grente, 2021). Since 2019,
considering that the wolf population has reached a satisfying demographic conservation status,
France has increased the possible number of lethal removals to 21% of the annual winter population
size estimate (Ministère de la Transition Ecologique et Solidaire, 2020).

Because wolves were removed sporadically between 2004 and 2010 (range: 0-2 wolves per year), we
started our study period in 2011, until 2020 included. Within the study period, only three lethal
removals were outside south-eastern France and concerned lone wolves, which dispersed far and
whose territories were very isolated. Their deaths could only lead to an absence of attacks. This
context was totally different from the one in the Alpine range where the wolf population was
settled. Therefore, we restricted our study area to the two south-eastern regions, Provence-Alpes-
Côte-d'Azur in its entirety and Auvergne-Rhône-Alpes restricted to its five most south-eastern
departments, Ardèche, Drôme, Isère, Savoie and Haute-Savoie (Figure 1).

 **2.2 Datasets**

 *2.2.1    Lethal removals*

We used the 362 administrative records of wolves killed by legal shootings from 2011-2020 that
were geolocated within the study area. Dates of lethal removals were recorded, but not the hour.
The number of lethal removals increased over time and could be divided into three periods, 2011-
2014, 2015-2018 and 2019-2020, that represented respectively 8%, 44% and 48% of the dataset, for
annual means of 7 (standard deviation, hereafter '±', 6), 40 (± 6) and 87 (± 3) lethal removals
respectively.

Almost half of the records were cases of reinforced defense (48%), while simple defense and hunting
cases represented respectively 27% and 25% of the dataset. There were 58 records (16%) for which
no wolf body was found after the shooting, but they were considered as killed because there was
enough evidence for lethal injuries. For the remaining 304 records, information from the necropsy
reports was available on the sex and the age class (pup, subadult or adult) of the killed wolves and
on the annual reproductive status for females.

*2.2.2    Attacks on sheep*

We used the 21 936 claims of attacks on sheep from 2010-2020 and geolocated within the study
area, that were compiled by the French administration and for which wolf responsibility could not be
discarded.  Each claim corresponded to one or several dead or wounded sheep.

*Figure 1. Study area and location within France. Purple points represent locations of the 188 single lethal removals, colored according to the three administrative types of shooting implementation (simple defense, reinforced defense, hunting) and of the centroids of the 35 multiple lethal removals (darkest color). Pastoral units with or without attacks are shown in green and grey, respectively.*

Apart very few specific cases, all claims were checked in the field by trained public agents. They
followed a standardized procedure to estimate the date of the attack and if wolf responsibility could
be discarded or not. Compensation was attributed regardless of non-lethal measures up to 2019.
From 2020, compensations after two compensated attacks depended on the implementation of
protective measures, with conditions of implementation varying locally according to the level of
attacks of municipalities. Protective measures are widely implemented in south-eastern France
(Meuret et al., 2020). We have therefore confidence in the level of reports of attacks noticed by
breeders.

### 2.2.3   Spatio-temporal sheep distribution

We used the data from the pastoral census of the two regions Provence-Alpes-Côte-d'Azur and
Auvergne-Rhône-Alpes conducted by the National Research Institute for Agriculture, Food and the
Environment (Dobremez et al., 2016). The census indexed and delimited for the period 2012-2014
the pastoral units at medium or high altitude used by flocks, as well as the pastoral zones at lower
altitudes. We restricted the dataset to units and zones that were used for grazing sheep, regardless
of the other livestock. The annual number of grazing sheep was on average 955 (± 764) for units
(range 2-6000) and 412 (± 411) for zones (range 1-3600). The months for which the area was grazed
were known for each pastoral unit. Only the seasons in use were available for pastoral zones.
Therefore, we attributed to the pastoral zones the months corresponding to the seasons for which
they were grazed (spring: March to May, summer: June to August, fall: September to November,
winter: December to February). The annual number of grazed months was on average 6.9 (range 1-
12). The total size of our final dataset of pastoral units and zones (hereafter 'pastoral units') was
4987 (Figure 1).

A substantial part of the attacks (41%) was not occurring within pastoral units during their set of
grazed months. This part was only reduced at 35% when considering the attacks from the period of
the pastoral census (2012-2014). Therefore, most mismatches were not due to the temporal
desynchronization between the two datasets. These mismatches certainly resulted from changes in
the use of pastures that shepherds or breeders need to apply to adapt to real-time conditions. We
corrected minor mismatches in the attack and pastoral datasets by following the procedure
described in Supplementary Information S1. We excluded 13% of the attack dataset from the
analysis, because no pastoral unit could be found within 500 meters from these attacks, or because
they occurred in units that were supposedly not used, with a temporal mismatch of more than 3
months. The final dataset of attacks had 19 302 events. We discuss the limits of this correction in the
Discussion section.

## 2.3  Analysis

233 Apart from being precisely geolocated, our data about attacks and lethal removals were very
234 unbalanced in favor of attacks (19 302 attacks against 362 lethal removals). Consequently, we
235 adopted an original approach consisting in choosing removals as focal points, and in computing
236 attack intensities over continuous spatial and temporal scales.

### 2.3.1   Spatio-temporal distributions of attacks around removals

We grouped the lethal removals separated by less than 5 km and less than 25 days (hereafter,
'multiple removals'), and considered the other removals as independent (hereafter, 'single
removals').

For each single removal, we only selected the attacks in a radius of 10 km from its location (Figure 2)
and across 90 days before and 90 days after its date. The day of the removal was not part of the
buffer, because we could not know if an attack recorded at the date of the removal occurred after or

before the removal, having no information about the hour of both attacks and removals. We
attributed two coordinates *(i, j)* to each attack (Figure 2). Coordinate *i* corresponded to the spatial
distance *i* (km) between the attack and the removal. Coordinate *j* corresponded to the temporal
distance (days) between the date of the attack and the date of the removal. The point of coordinates
*i* = 0 and *j* = 0 represented the spatial location of the removal, and the day before the removal (*i.e.*
the day of the removals was not part of the analysis).

In the case of multiple removals, we selected the attacks in a radius of 10 km from the centroid of
the removals of the group, and across 90 days before the date of the first removal of the group and
90 days after the date of the last removal of the group. As before, we attributed the same two
coordinates *(i, j)* to each attack. Coordinate *i* corresponded to the spatial distance *i* (km) between
the attack and the centroid of the group. Coordinate *j* corresponded to the temporal distance (days)
between the date of the attack and the date of the first (or last) removal of the group if the attack
occurred before (or after) the period covered by the group.

Attributing one spatial coordinate *i* to the attack instead of two traditional coordinates induces a
spatial bias that needed to be corrected.

We ensured that the spatial and temporal selections of the attacks for each removal did not overlap
areas or periods without data about attacks. We excluded from the analysis the 84 lethal removals
occurring at less than 10 km from the limits of the study area or occurring after the 1$^{st}$ October 2020.

Eventually, we randomly simulated attacks around each removal, with a distribution defined
according to the locations of pastoral units within 10 km and their grazed months. The simulated
attack dataset corresponded to the expected distribution of the attacks if they only depended on the
livestock presence. We repeated the simulation 1000 times (Figure 2).

More information about the selection of attacks, the correction of the spatial bias and the simulation
of attacks is available in Supplementary Information S2.

*2.3.2    Intra-site comparison*

To generalize the spatio-temporal distributions of the attacks around removals, we aggregated the
observed attacks of all removals into a unique set (Figure 2). We repeated the same process for each
of the 1000 replications of simulated attacks. We treated separately the dataset of single removals
and the dataset of multiple removals. We also explored other sets of aggregation of single removals
(hereafter, 'subsets') that we described in the next section.

We then estimated the observed and simulated attack intensities $\lambda_{obs}$ and $\lambda_{sim}$ by applying an
anisotropic Gaussian kernel density estimation (Figure 2). Given similar spatial corrections, the
higher the number of attacks within the vicinity of the same pair of coordinates (*i,j*), the higher the
attack intensity estimate of the related pixel *λ(i,j)*. Eventually, we computed the mean simulated
attack intensities $\bar{\lambda}_{sim}$ by averaging $\lambda_{sim}(i,j)$ across the 1000 sets of simulations.

*Figure 2. Example of the spatio-temporal distributions of attacks around three lethal removals. The circles in the center show the spatial distribution of observed attacks (red points) within the pastoral units (green polygons) 10 km from the removal location (purple points). The intermediate graphs show the distributions of the observed attacks (left) and of two sets of simulated attacks (right) around removals along a unique spatial axis (i) and temporal axis (j). The size of points represents the spatial correction (see Supplementary Information S2). The observed and simulated attack intensities $\lambda_{obs}$ and $\lambda_{sim}$ on the sides are calculated by aggregating the observed attacks and the simulated sets of attacks, respectively. The mean simulated attack intensities $\bar{\lambda}_{sim}$ are eventually calculated by averaging $\lambda_{sim}(i,j)$ of all simulated sets (here only two for the example). Rectangles delimit the five nested scales for trend computations: 15, 30, 45, 67 and 90 days and 0, 2, 4, 6, 8 and 10 km.*

289 To correct the observed intensities $\lambda_{obs}$ for livestock presence, we calculated the corrected
290 intensities $\lambda_{corr}(i, j)$ as:

$$\lambda_{corr}(i,j) = \lambda_{obs}(i,j) - \bar{\lambda}_{sim}(i,j)$$

Thus, a value of $\lambda_{corr}(i, j)$ lower, equal or higher than zero indicated that the observed attack intensity
at coordinates *(i, j)* was lower, equal or higher, respectively, than the attack intensity expected
according to pastoral use.

Eventually, we calculated the trends of observed or corrected attack intensities after removals,
taking the intensities before removals as the basis. Both observed and corrected trends could exceed
100% if the total of intensities after removals more than doubled the total of intensities before
removals. Corrected trends could also be lower than -100% because $\lambda_{corr}$ could take negative values.
The standard deviation of each trend was estimated by using the jackknife method (Maesono, 2011).

First, we calculated trends at five nested scales, by progressively increasing both spatial and
temporal scales (Figure 2). We also calculated these trends by increasing the temporal scale only, for
three fixed spatial scales. Second, we investigated the potential spatial shift of attack intensities by
calculating trends at each spatial unit for three fixed temporal scales. Contrary to the nested trends,
we calculated the trend of the attack intensity at *j* km, not from 0 to *j* km. In all cases, trends were
adjusted to allow comparison across scales or sets of removals.

More information about the attack intensity estimation and the calculations of trends is available in
Supplementary Information S3.

### 2.3.3 Inter-site comparison

Lastly, we combined the intra-site comparison performed in the previous section with an inter-site
comparison. In other words, we applied a complete BACI study design (Christie et al., 2019). This
study design allowed us to address confounding factors but also temporal autocorrelation, *i.e.* the

risks that attacks naturally persisted, increased or decreased after lethal removals, without being
related to the application of lethal removals. The correction of attack intensities for livestock
presence allowed us to partly address the risk of confounding factors, because livestock presence is
the primary factor of attack distribution, but there were many other possible factors that could
modify attack distribution beside lethal removals (*e.g.* livestock protection, wild prey distribution,
wolf poaching…).

We compared the trends of the spatio-temporal distributions of attacks around lethal removals and
those around control events where no lethal removal was applied. If the trends of lethal removals
and of control events were similar, we could assume that the trends after lethal removals were not
the result of the removals themselves, but the result of confounding factors or of temporal
autocorrelation. Otherwise, we could assume that the lethal removals mainly caused the observed
changes on attack distribution after their application.

Each removal received one control event. In the case of a single removal, we attributed the date of
the removal to its control event. In the case of a multiple removal, we attributed two dates to its
control event: the date of the first removal of the group and the date of the last removal of the
group.

Then, we delimited the sampling area where the spatial location of the control event could be
randomly sampled. We considered that only the geographic zone the removal was within
represented a similar environment. We delimited these zones by using 25 Alpine mountain ranges
defined by hydrography and orography (Knopf, 2019) and the limits from OpenStreetMap of the
protected area *Parc Naturel Régional des Préalpes d'Azur* (PNRPA) and of the military zone *Canjuers*.
We buffered the zones by 15 km, in order to include removals in their vicinity, provided that the
buffer did not overlap another buffered zone, or the area closer than 10 km from limits of the study
area (Figure 3). We excluded from the analysis the 9 removals that could not be attributed to a zone.
However, the attack distribution close to a control event could be distorted by the lethal removals
occurring shortly before or after the control event. Therefore, the sampling area did not include the
area within a 10-km radius around any known removal occurring less than 90 days from the control
event.

In total, we generated 100 control events per lethal removal. In the same way as for the sets of
lethal removals, we applied the intra-site comparison process previously described to each
generation of control events (hereafter, 'control set').

Eventually, we assessed if the removals significantly modified the intensities of attack intensities (*i.e.*
alternative hypothesis), or if there was no sufficient evidence to reflect an effect of removals on
attacks (*i.e.* null hypothesis). We considered that the effect was significant if the confidence intervals
of the trends following removals were not overlapping the confidence intervals of the trends
following control sets. Following Knol et al. (2011), we adapted the levels of the confidence intervals
(range 83.4–95%) in order to set the significance level at 0.05 (*i.e.* 5% of risks of rejecting the null
hypothesis when it is true).

In the end, we analyzed 188 single removals (1 killed wolf) and 35 multiple removals, with on
average 2.29 killed wolves (± 0.46, range 2—3) per multiple removal. On average, 27.4 attacks were
selected per removal (± 20.7, 1—100), against 16.7 attacks per control event (± 0.9). A total of 6024
attacks was selected across all removals, against 3387 attacks (± 173) on average per control set.

*Figure 3. Sampling areas for the control events of the 223 lethal removals (colored squares). The inset plot shows the sampling area of one focal removal (thick square) in the Vercors zone, and its 100 sampled control events (grey round points).*

354 *2.3.4 Hypotheses*

355 We explored three possible non-exclusive alternative hypotheses:

1. **Hypothesis 1**: The lethal removals decreased attack intensities by removing problem individuals (Linnell et al., 1999).
2. **Hypothesis 2**: The lethal removals decreased attack intensities by decreasing local wolf density (Bradley et al., 2015). Relationship between wolf density and attack intensity is also modulated by local richness of wild prey (Petridou et al., 2019).
3. **Hypothesis 3**: The lethal removals decreased attack intensities at small analysis scales and increased attack intensities at intermediate and large analysis scales, by altering the territorial limits of remaining wolves (Brainerd et al., 2008).

We tested the three hypotheses by applying separately the BACI method on different sets of lethal
removals defined according to different criteria.

Our first test was made by comparing results from the single removals (n = 188) and from the
multiple removals (n = 35). We predicted that a similar decrease of attack intensity observed for
both datasets could be in favor of Hypothesis 1, *i.e.* the removal of only one attacking wolf was
sufficient to reduce attack intensities. We predicted that a stronger decrease of attack intensity for
the dataset of multiple removals could be in favor of Hypothesis 2, *i.e.* attack intensity was

correlated to wolf density. Following Hypothesis 3, we predicted that the dataset of multiple
removals could more likely support this hypothesis than the dataset of single removals, because risks
of territorial alterations increase with the number of killed wolves within a pack (Cassidy et al.,
2023).

Our second test was made by dividing the dataset of single removals according to the 3 removal
administrative types, as explained in section 2.1 (Figure 1). In practice, the higher the attack intensity
in an area, the higher the authorized effort to remove a wolf, from *Simple defense* (n = 65), through
*Reinforced defense* (n = 71), and finally *Hunting* (n = 52). Note, however, that the three subsets still
concerned removals of only one wolf.

Following Hypothesis 1, we predicted that the *Hunting* subset could present a weaker decrease of
attack intensity than the two other subsets, or even no effect, because *Hunting* removals could kill
wolves outside of a context of livestock defense, *i.e.* wolves that could not have been responsible for
attacks. Following Hypothesis 2, we predicted that the efficiency of removals of *Simple defense*,
*Reinforced defense* and *Hunting* to decrease attack intensity would be decreasing following this
order, *i.e.* efficiency of removing one wolf decreases with attack intensity.

Our third test was made by dividing the dataset of single removals according to removal location
(Figure 4). We selected the 8 zones with at least 10 removals (Figure 4): *PNRPA* (n = 33), *Mercantour*
(23), *Vercors* (18), *Trois-Evêchés* (18), *Vanoise* (15), *Préalpes de Digne* (12), *Ecrins* (12) and *Canjuers*
(11).

*Figure 4. Topography of the study area. Points represent single removals according to their zone and the wolf season*
*during which they occurred. The 8 zones selected for the analyses of the geographic subsets are highlighted.*

Following Hypothesis 2, we predicted the weaker decrease of attack intensity for the 3 subsets with
the earliest wolf recolonization (*i.e.* starting before 2000), which could result in high local wolf
density: *Mercantour*, *Préalpes de Digne* and *Vercors*. Conversely, we predicted the strongest
decrease of attack intensity for the 2 subsets with the latest wolf recolonization (*i.e.* after 2015):
*Vanoise* and *Ecrins*. Eventually, we predicted intermediate results for the 3 subsets with wolf
recolonization starting between 2000 and 2010: *Canjuers*, *Trois Evêchés* and *PNRPA*. Still in line with
Hypothesis 2, we also predicted that removals applied in the least mountainous areas, *Canjuers* and
*PNRPA*, could be among the least efficient subsets to decrease attack intensity, because wolves
could rely more heavily on livestock in these two areas due to a weaker wild ungulate species
richness (Petridou et al., 2019). Following Hypothesis 3, we predicted that the subsets with the
earliest wolf recolonization would less likely support this hypothesis because their high density of
pack territories could decrease the possibilities of territorial changes.

Our fourth test, focused on Hypothesis 2, was made by dividing the dataset of single removals
according to the wolf seasons during which removals occurred (Figure 4): *Winter* (November —
January, n = 29), *Reproduction* (February — March, 8), *Denning* (April — June, 33), *Rendez-vous* (July
— August, 56) and *Fall* (September — October, 62). We predicted that the subsets of *Rendez-vous*
and *Fall* could be the least efficient seasonal subsets to decrease attack intensity because of the
higher local wolf density due to the presence of growing pups. Similarly, the subsets of *Reproduction*
and *Denning* could be the most efficient because of the higher risks of reproduction failure due to
removals, leading to reduced pack sizes, but also because of the presence of young wild ungulates
during the *Denning* period that were alternative easy prey. We predicted that the subset of *Winter*
could present intermediate efficiency, because there are less wolves in winter than in summer, while
there are less young wild ungulates, making livestock particularly attractive if available (Mattioli et
al., 2011).

Our fifth and final test, focused on Hypothesis 2, was made by dividing the dataset of single
removals according to the sex of the killed wolf when known, *Male* (n = 83) and *Female* (n = 78). The
removals of the breeding females during spring may greatly impact pup survival, and thus pack size
and local wolf density. However, during fall and winter, breeder's presence also increases pup
survival regardless of breeder's sex (Ausband et al., 2017a). In our dataset, most of female removals
were done during summer and fall, and only 7 females over 78 were declared as being reproductive
the year they were killed (annual reproductive status of males was unknown). Therefore, we did not
predict different results between the two subsets. Breeder removal has greater impacts on the pack
stability than other removals (Cassidy et al., 2023), however we could not test Hypothesis 3 because
the distribution of removals was too unbalanced between female reproductive status or between
age class (177 adults, 24 subadults and 16 pups).

**2.4 Implementation**

The analysis was conducted in R 4.4.2 (R Core Team, 2023). We used the R packages *tidyverse 2.0.0*
(Wickham et al., 2019) and *sf 1.0-12* (Pebesma and Bivand, 2023) to handle the datasets. We used
the R package *spatstat 3.2-1* (Baddeley et al., 2015) to simulate point patterns. We adapted the
function *kde2d.weighted* from *ggtern 3.4.2* (Hamilton and Ferry, 2018) to separate the kernel
estimates before and after the removal date. We used the package *osmdata 0.2.5* (Padgham et al.,
2017) to extract limits of PNRPA and of Canjuers military zone. We used the packages *geodata 0.5-8*
(Hijmans et al., 2023), *ggfx 1.0.1* (Pedersen, 2022a), *ggnewscale 0.5.2* (Campitelli, 2023), *ggpubr*
*0.6.0* (Kassambara, 2023), *metR 0.14.1* (Campitelli, 2021), *patchwork 1.3.1* (Pedersen, 2022b),
*RColorBrewer 1.1-3* (Neuwirth, 2022), *scales 1.3.0* (Wickham and Seidel, 2022), *tidyterra 0.7.2*

(Hernangómez, 2023) and *tidytext 0.4.3* (Silge and Robinson, 2016) for data visualization. The
datasets (with anonymized coordinates), codes and results are available online (Grente et al., 2025).

## 3. Results

### 3.1 General spatio-temporal distribution of attacks around lethal removals

Spatio-temporal distribution of attacks around lethal removals took various forms according to the
datasets or subsets (Figures 5, S1). However, we could extract two features common. First, removals
were generally preceded by high attack intensities at their locations, meaning they were applied in
response to attacks, in accordance with French legislation. Second, attacks tended to mostly
aggregate below 2 km from removal locations, regardless of time. Therefore, the attack process was
generally a persistent phenomenon at these locations, either before or after removals, while
generally presenting several interruptions in time.

### 3.2 General effect of removals

The trends calculated for the dataset of single removals (*i.e.* one killed wolf) and for the dataset of
multiple removals (*i.e.* on average killed 2.29 wolves) were significantly negative across the five
nested analysis scales (Figure 6).

The trend of attack intensity after single lethal removals ranged from -22.5% (-24.2% – -20.8%) to -
6.3% (-7.15% – -5.38%) according to the analysis scale (Figure 6), suggesting the lethal removals of
one wolf moderately decreased the attack intensities.

By comparison, the trend of attack intensity after multiple removals ranged from -85.1% (-93.4 – -
76.9%) to -12.8% (-14.7 – -10.9%) according to the analysis scale (Figure 6), suggesting the lethal
removals of 2 or 3 wolves very strongly to moderately decreased the attack intensities.

### 3.3 Effect of the analysis scale

The analysis scale had a limited effect on the trend for the dataset of single removals (Figure 6). At
fixed spatial scale, the decrease of attack intensity remained moderate over 90 days after the
removal (Figure 7A). However, the short periods after single removals, below 15 days, could show
non-significant trends.

On the contrary, the analysis scale had an important effect on the trend for the dataset of multiple
removals (Figure 6). Over 0–2 km, the trend distinctly increased with time (Figure 7A), going from
the strongest decrease at day 1 (-124% – -140%) to the smallest decrease at day 90 (-30.4% – -
19.5%). The very strong effect of reduction of multiple removals disappeared and became moderate
when focusing on larger spatial scales, joining the trend of single removals over the entire time scale
(Figure 7A). This suggests that the very strong reduction induced by multiple removals was mostly
localized to the targeted pasture and to its closest neighbors over a period of more than one month.

Spatial shifts of attacks after removals were not obvious (Figure 7B). For most distances, trends were
negative, regardless of the fixed temporal scale. Nevertheless, some significant positive trends could
appear at certain distances (e.g. around the 3[rd] and 7[th] km, 15 days after multiple removals) but they
had not the magnitude of negative peaks. This might suggest that a minor part of the attack
intensities shifted towards other pastures after removals.

### 3.4 Effect of the correction for livestock presence

The correction for livestock presence had a limited effect on the trends for the dataset of single
removals, but reinforced the magnitude of negative trends for the dataset of multiple removals
(Figures 6, S2, 7, S3). In other words, raw attack intensities decreased after multiple removals, but
this decrease took on even more significance in light of the fact that most pastures were still in use
after these multiple removals, and could therefore have been attacked but were mainly not.

This difference between single and multiple removals could be explained by the fact that single
removals were more evenly distributed between summer and fall, while multiple removals were
mainly applied in August (Figure 5). The dataset of single removals (n = 188) was also larger than the
dataset of multiple removals (n = 35). Therefore, the needs for a correction would be less important
for a large dataset with various underlying livestock distributions.

### 3.5 Effect of subsets

The main effect of moderate decrease of single removals concealed disparities between single
removals. When dividing this large dataset according to removal's location, removal's season,
removal's administrative type or the sex of the killed wolf, we could observe very different trend
profiles across the subsets (Figures 8, 9).

At the smallest nested scale (*i.e.* over 15 days and 0–2 km), we observed a large gradient of
significant negative trends (Figure 8). Besides 6 subsets that presented moderate decreases as the
general dataset of single removals (from -5% to -24%), 3 subsets presented very strong decreases
(below -50%) and 3 others strong decreases (from -49% to -25%). When increasing the analysis scale,
most subsets showed similar results to the general dataset of single removals (Figures 8, 9).

In addition, there were three subsets that showed significant positive trends (*Vercors*, *Winter* and
*Rendez-vous*), while their trends at the first nested scale were significantly negative (Figure 8).
Besides the increase of attack intensity, spatial shifts could be suspected for these subsets (Figure
S4B). The correction for livestock presence for these subsets played an important role to reveal most
of these positive trends, that could be not significant or even negative without correction (Figures
S5, S6). In these cases, this meant that the raw decrease of attack intensities was in fact expected
given the fact that most pastures were not in use anymore after the removals, and that the attack
intensity actually increased when taking this bias into account. This was especially true for the
Winter subset, because its removals occurred at the limit between an intense and calm grazing
period (Figure S1).

These deviations from the general dataset of single removals, whether through strong decreases or
increases after single removals, revealed particularities that could happen in some situations, but did
not represent the main effect that we could expect after single removals. Except the *Rendez-vous*
subset, the sizes of the subsets with these deviations were indeed below the average subset size
($\bar{n} = 37.3$).

### 3.6 Exploring the variability of the effect

510    This variability among subsets could be explored through the results to the tests of hypotheses we
511    described in section 2.3.4.

512    Hypothesis 1 ("*lethal removals decreased attack intensities by removing problem individuals*") was
513    not supported. Single removals did not show similar trends as multiple removals, and removals from
514    the Hunting subset were not the least efficient among its category (Figures 6, 9).

Hypothesis 2 ("*lethal removals decreased attack intensities by decreasing local wolf density*")
received partial support (Figures 6, 9). As expected, multiple removals presented a better efficiency
to decrease attack intensity than single removals. However, the prediction that removal efficiency
decreased with the intensities of attacks preceding removals, tested with the subsets of the
administrative types, was not verified by the results. Results were mixed regarding the geographic
and season subsets. The subsets of *Canjuers* and *PNRPA* were the only subsets that did not show
strong negative trends after removals, in line with the possible higher reliance of wolves on livestock
there, due to a weaker wild ungulate species richness. However, the order of efficiency of
geographic subsets expected according to the historical wolf recolonization did not clearly support
the hypothesis. The subsets of *Denning* and *Rendez-vous* verified the predictions of Hypothesis 2,
but not the subsets of *Reproduction* and *Fall*. Contrary to the initial prediction, the subset of *Winter*
did not present intermediate efficiency, but instead showed positive trends at the largest scale. This
could match with part of the prediction, where we suggested that wolves may heavily depend on
livestock at this season. Eventually, the absence of large trend differences between the subsets of
female and male removals was consistent with Hypothesis 2.

The results for Hypothesis 3 ("*lethal removals altered wolf territorial limits*") also received partial
support. As predicted, the dataset of multiple removals more supported this hypothesis than the
dataset of single removals (Figure 7B). However, two of the three subsets with the earliest wolf
recolonization (*Vercors* and *Préalpes de Digne*) presented signs of spatial shift (Figure S4B), contrary
to the prediction.

*Figure 5. Corrected attack intensities around lethal removals (middle) and around the first set of control events (right), for*
*the dataset of single removals (A) and of multiple removals (B). Each panel includes the monthly distribution of the dataset.*
*Grey rectangles delimit the five nested scales used to calculate trends in Figures 6, 8 and 9.*

General datasets

*Figure 6. Trends of attack intensities after the applications of single removals and multiple removals, at five nested scales. Points are mean estimates, alongside the confidence intervals (range 83.4%–89.8%). All trends are significant. Trends were corrected for livestock presence.*

*Figure 7. Trends of attack intensities after the applications of single removals (orange) and of multiple removals (purple), alongside continuous temporal (A) and spatial (B) scales, for three fixed distances and periods, respectively. Lines are mean estimates, ribbons are the confidence intervals (range 83.4–90.4%). Dotted lines are not significant trends. Trends were corrected for livestock presence. Note that temporal trends are calculated as nested periods (e.g. from 0 to x days, over 0–2 km for first panel), while spatial shifts are trends calculated for specific distances (e.g. at the $x^{th}$ km, over 15 days for first panel).*

*Figure 8. Distributions of the trends of attack intensities for the 18 subsets of single removals at three nested scales. Points*
*are mean estimates, alongside the confidence intervals (range 83.4%–89.8%). Trends in grey are not significant. Trends*
*were corrected for livestock presence. Confidence intervals of the entire dataset of single removals are in background in*
*dark blue.*

*Figure 9. Trends of attack intensities after the applications of single removals, divided into 18 subsets, at five nested scales.*
*Points are mean estimates, alongside the confidence intervals (range 88.5%–89.8%). Trends in grey are not significant.*
*Trends were corrected for livestock presence. 'R. Def' and 'S. Def' for Reinforced and Simple defense subsets; 'T. Evêc.', 'P.*
*Digne' and 'Mercant.' for Trois Evêchés, Préalpes de Digne and Mercantour subsets; 'Rdv' and 'Repro.' for Rendez-vous and*
*Reproduction subsets.*

## 4. Discussion

We evaluated the effects of lethal control of wolves on attack intensity distribution through kernel density estimation. Our method provided an efficient visualization of attack intensity distributions in the form of heatmaps over continuous spatial and temporal scales, corrected for sheep presence. The resulting attack intensity trends showed the differences in the attack intensities before and after lethal removals (intra-site comparison) and with comparable areas without removals (BACI design).

The aim of our study was to generalize the spatio-temporal patterns of attack intensity surrounding lethal removal applications, without concealing the potential diversity of these patterns across the French Alps that could result from different situations. This is why we compared trends of single lethal removals and trends of multiple lethal removals, but we also divided single lethal removals into 18 subsets according to the application conditions (geographical, seasonal, administrative) or to the removal results (sex).

### 4.1 Assessment of lethal removal efficiency

The distribution of the attacks was usually structured by space, with strong spatial aggregations of attacks, especially close to the locations of the lethal removals. This suggested that some locations may have been more subject to attacks than others, including at a local scale (< 10 km).

Removals of several wolves (2.29 wolves on average) were more efficient to reduce attacks than lethal removals of one wolf, especially at low analysis scales. Multiple removals reduced the attack intensity by 85% over 15 days and 2 km, against 11% for single removals. Results of both datasets were getting closer with increasing analysis scales. If we focus on the pastures targeted by multiple removals and their closest neighbors (< 2 km), the reduction of attack intensity remained very strong or strong during 89 days. These results are consistent with the results from Bradley et al. (2015), where removal efficiency increased with the number of killed wolves in the northern Rocky Mountains of the USA.

The effect of analysis scale was also visible for the 18 subsets of single lethal removals. The moderate decrease of single lethal removals transformed into a large gradient of trends at the subset level. Generally, the smaller the area and the time period surrounding the lethal removals, the stronger the decrease of attacks after removals. Three subsets also showed positive trends while their trends at the lowest analysis scale were negative, and showed signs of spatial shifts of attacks.

Our results highlight the importance of accounting for scale in such assessments. Therefore, the observed disparity between the results of studies evaluating wolf lethal control could partly result from the disparity in the analysis scales. If the effect of removals generally weakens with analysis scale, like in our results, the likelier it is to find non-significant results when assessing removal effects over large spatio-temporal scales. For example, Kutal et al. (2023) found no significant effect of wolf removals on attacks in Slovakia, but analyzed the datasets over 711 km² (314 km² here) and over one year (3 months here) without intermediate scales.

Differences in analysis scales cannot be the only explanation for differences between studies, as Santiago-Avila et al., (2018) found non-significant effects while using a progressive spatial scale comparable to ours. Besides analysis scales, other methodological choices, linked to the analyzed datasets, may explain these differences. Here, our dataset was very unbalanced in favor of attacks, with a ratio of 1 removal for 27 attacks, against 1 for 7 in Santiago-Avila et al., 2018. Thus, the risk of recurrence of a single attack was the chosen metric in several studies with datasets from the USA

605 (Bradley et al., 2015; Harper et al., 2008; Santiago-Avila et al., 2018), whereas we chose to focus on
606 the number of attacks (*i.e.* intensities).

### 4.2 Limits of the analysis

We expect these results to reasonably reflect the effects of lethal removals on wolves. Thus, the
lethal removals should have generally decreased attack intensities by temporarily decreasing local
wolf density or by altering the distribution or the predation behavior of the remaining wolves.
However, as in every study, confounding factors may interfere. Here, the main risk was to confound
the effects of lethal removals on wolves with the effects of livestock distribution dynamics in space
and time. We corrected our analysis for livestock distribution based on a pastoral census in the Alps
that gave the precise limits of pastures and their months or seasons in use (Dobremez et al., 2016).
This type of regional census requires a substantial amount of work. Fine-scale livestock data as
provided by this census is therefore generally missing from analyses on predation, including for
other parts of France (*e.g.* Gastineau et al., 2019). When integrated, livestock data is generally at
large, administrative scales (*e.g.* DeCesare et al., 2018; Merz et al., 2025; Šuba et al., 2023). Thus,
only two comparative studies (Allen, 2015; Conner et al., 1998) and two correlative studies (Blejwas
et al., 2002; Wagner and Conover, 1999) that evaluated lethal control effects on canids corrected
their analysis for livestock presence at the pasture level.

However, despite the quality of the livestock census we used, this may not have been enough to
prevent the main confounding risk for our study. Annual changes in the use of pastures since the
census may have occurred. In addition, temporary changes in livestock presence are common during
a grazing season, as livestock owners and shepherds modulate the use of pastures by their flocks to
optimize grazing according to environmental conditions. Among these conditions, pressure due to
wolf predation or attempts of predation may encourage livestock owners to remove flocks from
some pastures or to reinforce preventive measures. Data on precise husbandry practices, such as
preventive measures, were out of the scope of this census. Hence, resulting potential variations in
livestock distribution and protection, even small and temporary, may drive changes in attack
distribution that would not have been considered in the census and therefore in the livestock
correction of our analysis. To a lesser extent, variations in the spatio-temporal distributions of wild
ungulates may also impact the distribution of attacks on livestock.

In spite of the limits of the livestock dataset, the intra-site comparison design should have helped to
reduce the confounding risks, especially at the smallest scale. Because wolves move over large
territories, we acknowledge that focusing on changes happening only in very close proximity to
removals could not be enough to evaluate removal effects (Treves et al., 2016). To improve the
statistical power of our analysis, especially at the larger analysis scales, we doubled the intra-site
comparison with an inter-site comparison, therefore applying a BACI design, evaluated as robust by
Christie et al. (2019). The main limit of this study design is to use appropriate control sites. In our
case, this was a challenging task because lethal removals in France were generally authorized and
applied in attack hotspots, creating unbalanced initial attack intensities between sites with and
without removals. Nevertheless, the buffers of control events had on average 16.7 attacks (± 0.9)
against 26.3 (± 20.7) for the removal buffers, suggesting that control sites were also subject to
attacks.

### 4.3 Underlying mechanisms

If the confounding risks were as low as possible, we can conclude that lethal removals generally
decreased attack intensities, with only a few situations leading to other results (non-significant or

increase). We explored several hypotheses to understand the underlying mechanisms leading to
these results. Our results did not support the hypothesis of problem individuals (Linnell et al., 1999).
In addition, for most analyzed subsets, attacks persisted after removals at the location of the
removals, including at high intensity, even though most removals targeted wolves approaching
livestock or attempting attacks. This result suggested that attacks on livestock were more a
structural phenomenon than the outcome of some problem individuals.

The better efficiency of multiple removals compared to single removals clearly supported the
hypothesis of a reduction of attacks through the temporary decrease of local wolf density. The initial
local wolf density might also explain the gradient of attack responses between the subsets of single
removals. However, the support to this hypothesis was less clear-cut, probably because our
characterization of the subsets was too rough and simplistic to describe local contexts surrounding
lethal removals.

Therefore, the gradient of attack responses after removals observed between subsets of single lethal
removals remains largely unexplained. Besides effects on wolf density, removals may have altered
the distribution or the predation behavior of the remaining wolves, as spatial shifts could be
suspected for some subsets, like in Santiago-Avila et al. (2018). However, the persistence of attacks
at the removal's locations in most situations, even at lower intensities, challenges the hypothesis of
pack dissolution (Elbroch and Treves, 2023). Indeed, settlement of new wolves at the same location
was unlikely to be as fast, even in the dense population of the Western Alps (Marucco et al., 2023).

### 4.4 Implications for management and research

Concluding that the analyzed removals were an efficient management tool requires to define
efficiency itself. As in most studies on the subject, we considered that the significant decreases of
attack intensities after removals met expectations of efficiency, at least for the analyzed scales.
However, other definitions are possible, depending on the magnitude of the reduction, the spatio-
temporal scales at which the effects apply, and whether or not removals threaten wolf population
viability (*e.g.* Lennox et al., 2018). Expectations regarding removal efficiency also depend on the
benefits and costs of wolf's conservation, that can be very different across people (Linnell et al.,
2010). Thus, research results about management actions can be differently interpreted by people
(Hodgson et al., 2019). This is why clear management objectives of lethal control must be defined
through open dialogue, in view of ecological responses like here, but also by considering
socioeconomic dimensions and wolf population viability (Lennox et al., 2018; Linnell et al., 2010).

We showed that the analyzed lethal removals of wolves generally helped to reduce attack intensities
within the targeted pastures and their neighbors. By contrast, some particular situations did not lead
to this result (*e.g.* removals within *Vercors*, *Canjuers,* or during July-August). Considering our results
are still valid in today's situation, this variability may be a thorn in manager's side, as wolf removals
will probably lead to attack reduction, but not with certainty. Given the strong impact lethal control
can have on wolf population (Grente et al., 2024), the next step for research would be to move
forward from patterns to mechanisms. This understanding is necessary to provide the keys for
managers for the most efficient application of lethal control in terms of attack reduction, and
therefore for a better public acceptance of this tool.

We could only briefly explore the mechanisms of the effects of lethal removals on wolves and
therefore on their attacks. An advanced analysis would have required to have access to pack
compositions and space use by wolves at fine spatio-temporal scale, which were unavailable.
Collection of these datasets on wolves are particularly difficult on large areas, considering the

secrecy of wolves. A trade-off would be to deploy the substantial means required to collect such
datasets in a small study area (*e.g.* one wolf territory and its neighboring packs) for a few years,
ideally replicated at different locations. Effects of lethal control on wolves and their attacks may
then be evaluated through promising methods such as animal social network analysis, which was
applied to study the reaction of other social animals to human-induced removals (*e.g.* Downing et
al., 2023; Goldenberg et al., 2016). Working at small scales would also facilitate a more precise
collection of livestock presence and husbandry practices.

Variability is inherent to any ecological process (*e.g.* human avoidance by wolves, Ferreiro-Arias et
al., (2024); wolf mortality, Milleret et al., (2025)). Predation behavior is no exception. A few fine-
scale datasets on wolves and livestock would not necessarily explain all the observed removal effects
on attacks. A solution would be to upgrade the analysis of observed data by simulations (Railsback et
al., 2025). In this sense, we simulated removal effects on wolf population dynamics and on attacks at
the scale of a mountain range in the Alps (Grente et al., 2024), but our theoretical model would
greatly benefit from a combination with social network analysis based on observed data at small
scale.

**5. Conclusion**

The lethal management strategy implemented in France from 2011 to 2020—primarily involving the
removal of single wolves in defense situations with protected livestock—can be expected to have
produced a moderate reduction in attack intensity following removals. This reduction was greatly
amplified when removing multiple wolves. Our results partly supported the hypothesis of removal
effects through reduction of local wolf density but also through changes in the distribution of the
remaining wolves. However, contextual factors (e.g., geographical or seasonal variations) could lead
to deviations from this general pattern. Future research should focus on understanding wolves'
responses to removals—such as changes in pack composition, space use, and predation behavior—
to better interpret the observed variability in attack trends post-removal.

**Supplementary material**

Three supplementary materials are available online, on Zenodo
(https://doi.org/10.5281/zenodo.17231870).

File 1. Supplementary information (S1–S3)

File 2. Supplementary figures (S1–S9)

File 3. Supplementary tables (S1–S2)

**Acknowledgements**

We are indebted to the governmental agents in charge of recording attack claims and lethal
removals. We are grateful to the regional administration of Auvergne-Rhône-Alpes for the access to
the databases of claims and of wolf lethal removals. We are also grateful to INRAE (previously
IRSTEA) for the access to the pastoral census. We greatly thank Marion Valeix, John Linnell and
Michel Meuret for their valuable comments regarding the first draft of this study as part of the PhD
dissertation of Oksana Grente. We greatly thank Aaron Bott, Vincenzo Gervasi, Peter Sunde and an
anonymous reviewer for their feedbacks, that greatly helped us to improve the manuscript. We also
thank members of the scientific council of the National Plan of Action for wolves and livestock
breeding activities for their feedback and advices.

**Data, scripts and code**

Data, scripts and code are available online, on Zenodo (https://doi.org/10.5281/zenodo.17231870)
and on gitlab (https://gitlab.com/oksanagrente/Kernel_wolf_culling_attacks_p).

**Conflict of interest disclosure**

738 The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
739 relation to the content of the article.

740 Olivier Gimenez and Simon Chamaillé-Jammes are recommenders of the managing board of PCI
741 Ecology.

**Funding**

743 This study was funded by a grant from the French Ministry of Ecological and Solidary Transition and
744 the French Ministry of Agriculture and Food as part of the National Plan of Action for wolves and
745 livestock breeding activities.

**References**

Agreste, 2021. La transhumance collective. Données issues du recensement agricole 2020. [WWW
Document]. VizAgreste. URL https://vizagreste.agriculture.gouv.fr/la-transhumance-collective.html
(accessed 11.24.23).

Allen, B.L., 2015. More buck for less bang: Reconciling competing wildlife management interests in
agricultural food webs. Food Webs 2, 1–9. https://doi.org/10.1016/j.fooweb.2014.12.001

Allen, L.R., 2014. Wild dog control impacts on calf wastage in extensive beef cattle enterprises. Anim.
Prod. Sci. 54, 214. https://doi.org/10.1071/AN12356

Arif, S., MacNeil, M.A., 2022. Predictive models aren't for causal inference. Ecology Letters 25, 1741–
1745. https://doi.org/10.1111/ele.14033

Ausband, D.E., Mitchell, M.S., Stansbury, C.R., Stenglein, J.L., Waits, L.P., 2017a. Harvest and group
effects on pup survival in a cooperative breeder. Proceedings of the Royal Society B: Biological
Sciences 284, 20170580. https://doi.org/10.1098/rspb.2017.0580

Ausband, D.E., Mitchell, M.S., Waits, L.P., 2017b. Effects of breeder turnover and harvest on group
composition and recruitment in a social carnivore. Journal of Animal Ecology 86, 1094–1101.
https://doi.org/10.1111/1365-2656.12707

Baddeley, A., Rubak, E., Turner, R., 2015. Spatial Point Patterns: Methodology and Applications with
R, CRC Press. ed.

Bjorge, R.R., Gunson, J.R., 1985. Evaluation of Wolf Control to Reduce Cattle Predation in Alberta.
Journal of Range Management 38, 483. https://doi.org/10.2307/3899736

Blejwas, K.M., Sacks, B.N., Jaeger, Michael M., McCullough, D.R., 2002. The Effectiveness of Selective
Removal of Breeding Coyotes in Reducing Sheep Predation. The Journal of Wildlife Management 66,
451–462. https://doi.org/10.2307/3803178

Bradley, E.H., Robinson, H.S., Bangs, E.E., Kunkel, K., Jimenez, M.D., Gude, J.A., Grimm, T., 2015.
Effects of wolf removal on livestock depredation recurrence and wolf recovery in Montana, Idaho,
and Wyoming: Effects of Wolf Removal. Jour. Wild. Mgmt. 79, 1337–1346.
https://doi.org/10.1002/jwmg.948

773 Brainerd, S.M., Andrén, H., Bangs, E.E., Bradley, E.H., Fontaine, J.A., Hall, W., Iliopoulos, Y., Jimenez,
774 M.D., Jozwiak, E.A., Liberg, O., Mack, C.M., Meier, T.J., Niemeyer, C.C., Pedersen, H.C., Sand, H.,
775 Schultz, R.N., Smith, D.W., Wabakken, P., Wydeven, A.P., 2008. The Effects of Breeder Loss on
776 Wolves. Journal of Wildlife Management 72, 89–98. https://doi.org/10.2193/2006-305

777 Campitelli, E., 2023. ggnewscale: Multiple fill and colour scales in "ggplot2" (manual).

778 Campitelli, E., 2021. metR: Tools for easier analysis of meteorological fields (manual).
779 https://doi.org/10.5281/zenodo.2593516

780 Cassidy, K.A., Borg, B.L., Klauder, K.J., Sorum, M.S., Thomas-Kuzilik, R., Dewey, S.R., Stephenson, J.A.,
781 Stahler, D.R., Gable, T.D., Bump, J.K., Homkes, A.T., Windels, S.K., Smith, D.W., 2023. Human-caused
782 mortality triggers pack instability in gray wolves. Frontiers in Ecology and the Environment n/a.
783 https://doi.org/10.1002/fee.2597

784 Chapron, G., Kaczensky, P., Linnell, J.D.C., von Arx, M., Huber, D., Andrén, H., López-Bao, J.V.,
785 Adamec, M., Álvares, F., Anders, O., Balčiauskas, L., Balys, V., Bedő, P., Bego, F., Blanco, J.C.,
786 Breitenmoser, U., Brøseth, H., Bufka, L., Bunikyte, R., Ciucci, P., Dutsov, A., Engleder, T., Fuxjäger, C.,
787 Groff, C., Holmala, K., Hoxha, B., Iliopoulos, Y., Ionescu, O., Jeremić, J., Jerina, K., Kluth, G., Knauer, F.,
788 Kojola, I., Kos, I., Krofel, M., Kubala, J., Kunovac, S., Kusak, J., Kutal, M., Liberg, O., Majić, A., Männil,
789 P., Manz, R., Marboutin, E., Marucco, F., Melovski, D., Mersini, K., Mertzanis, Y., Mysłajek, R.W.,
790 Nowak, S., Odden, J., Ozolins, J., Palomero, G., Paunović, M., Persson, J., Potočnik, H., Quenette, P.-
791 Y., Rauer, G., Reinhardt, I., Rigg, R., Ryser, A., Salvatori, V., Skrbinšek, T., Stojanov, A., Swenson, J.E.,
792 Szemethy, L., Trajçe, A., Tsingarska-Sedefcheva, E., Váňa, M., Veeroja, R., Wabakken, P., Wölfl, M.,
793 Wölfl, S., Zimmermann, F., Zlatanova, D., Boitani, L., 2014. Recovery of large carnivores in Europe's
794 modern human-dominated landscapes. Science 346, 1517–1519.
795 https://doi.org/10.1126/science.1257553

796 Christie, A.P., Amano, T., Martin, P.A., Shackelford, G.E., Simmons, B.I., Sutherland, W.J., 2019.
797 Simple study designs in ecology produce inaccurate estimates of biodiversity responses. Journal of
798 Applied Ecology 56, 2742–2754. https://doi.org/10.1111/1365-2664.13499

799 Conner, M.M., Jaeger, Michael M., Weller, T.J., McCullough, D.R., 1998. Effect of Coyote Removal on
800 Sheep Depredation in Northern California. The Journal of Wildlife Management 62, 690–699.
801 https://doi.org/10.2307/3802345

802 de Beaufort, F., 1987. Le loup en France: éléments d'écologie historique. Encyclopédie des
803 carnivores de France.

804 DeCesare, Nicholas.J., Wilson, S.M., Bradley, E.H., Gude, J.A., Inman, R.M., Lance, N.J., Laudon, K.,
805 Nelson, A.A., Ross, M.S., Smucker, T.D., 2018. Wolf-livestock conflict and the effects of wolf
806 management: Wolf-Livestock Conflict. Jour. Wild. Mgmt. 82, 711–722.
807 https://doi.org/10.1002/jwmg.21419

808 Dobremez, L., Bray, F., Borg, D., 2016. Enquête pastorale 2012-2014. IRSTEA.

809 Downing, B.C., Silk, M.J., Delahay, R.J., Bearhop, S., Royle, N.J., 2023. Culling-induced perturbation of
810 social networks of wild geese reinforces rather than disrupts associations among survivors. Journal
811 of Applied Ecology n/a. https://doi.org/10.1111/1365-2664.14522

812 Elbroch, L.M., Treves, A., 2023. Perspective: Why might removing carnivores maintain or increase
813 risks for domestic animals? Biological Conservation 283, 110106.
814 https://doi.org/10.1016/j.biocon.2023.110106

Fabbri, E., Miquel, C., Lucchini, V., Santini, A., Caniglia, R., Duchamp, C., Weber, J.-M., Lequette, B.,
Marucco, F., Boitani, L., Fumagalli, L., Taberlet, P., Randi, E., 2007. From the Apennines to the Alps:
colonization genetics of the naturally expanding Italian wolf (Canis lupus) population. Molecular
Ecology 16, 1661–1671. https://doi.org/10.1111/j.1365-294X.2007.03262.x

Fernández-Gil, A., Naves, J., Ordiz, A., Quevedo, M., Revilla, E., Delibes, M., 2016. Conflict Misleads
Large Carnivore Management and Conservation: Brown Bears and Wolves in Spain. PLoS ONE 11,
e0151541. https://doi.org/10.1371/journal.pone.0151541

Ferreiro-Arias, I., García, E.J., Palacios, V., Sazatornil, V., Rodríguez, A., López-Bao, J.V., Llaneza, L.,
2024. Drivers of Wolf Activity in a Human-Dominated Landscape and Its Individual Variability Toward
Anthropogenic Disturbance. Ecology and Evolution 14, e70397. https://doi.org/10.1002/ece3.70397

Gastineau, A., Robert, A., Sarrazin, F., Mihoub, J.-B., Quenette, P.-Y., 2019. Spatiotemporal
depredation hotspots of brown bears, Ursus arctos, on livestock in the Pyrenees, France. Biological
Conservation 238, 108210. https://doi.org/10.1016/j.biocon.2019.108210

Gervasi, V., Linnell, J.D.C., Berce, T., Boitani, L., Cerne, R., Ciucci, P., Cretois, B., Derron-Hilfiker, D.,
Duchamp, C., Gastineau, A., Grente, O., Huber, D., Iliopoulos, Y., Karamanlidis, A.A., Kojola, I.,
Marucco, F., Mertzanis, Y., Männil, P., Norberg, H., Pagon, N., Pedrotti, L., Quenette, P.-Y., Reljic, S.,
Salvatori, V., Talvi, T., von Arx, M., Gimenez, O., 2021. Ecological correlates of large carnivore
depredation on sheep in Europe. Global Ecology and Conservation 30, e01798.
https://doi.org/10.1016/j.gecco.2021.e01798

Goldenberg, S.Z., Douglas-Hamilton, I., Wittemyer, G., 2016. Vertical Transmission of Social Roles
Drives Resilience to Poaching in Elephant Networks. Current Biology 26, 75–79.
https://doi.org/10.1016/j.cub.2015.11.005

Gould, E., Fraser, H.S., Parker, T.H., Nakagawa, S., Griffith, S.C., Vesk, P.A., Fidler, F., Hamilton, D.G.,
Abbey-Lee, R.N., Abbott, J.K., Aguirre, L.A., Alcaraz, C., Aloni, I., Altschul, D., Arekar, K., Atkins, J.W.,
Atkinson, J., Baker, C.M., Barrett, M., Bell, K., Bello, S.K., Beltrán, I., Berauer, B.J., Bertram, M.G.,
Billman, P.D., Blake, C.K., Blake, S., Bliard, L., Bonisoli-Alquati, A., Bonnet, T., Bordes, C.N.M., Bose,
A.P.H., Botterill-James, T., Boyd, M.A., Boyle, S.A., Bradfer-Lawrence, T., Bradham, J., Brand, J.A.,
Brengdahl, M.I., Bulla, M., Bussière, L., Camerlenghi, E., Campbell, S.E., Campos, L.L.F., Caravaggi, A.,
Cardoso, P., Carroll, C.J.W., Catanach, T.A., Chen, X., Chik, H.Y.J., Choy, E.S., Christie, A.P., Chuang, A.,
Chunco, A.J., Clark, B.L., Contina, A., Covernton, G.A., Cox, M.P., Cressman, K.A., Crotti, M., Crouch,
C.D., D'Amelio, P.B., De Sousa, A.A., Döbert, T.F., Dobler, R., Dobson, A.J., Doherty, T.S., Drobniak,
S.M., Duffy, A.G., Duncan, A.B., Dunn, R.P., Dunning, J., Dutta, T., Eberhart-Hertel, L., Elmore, J.A.,
Elsherif, M.M., English, H.M., Ensminger, D.C., Ernst, U.R., Ferguson, S.M., Fernandez-Juricic, E.,
Ferreira-Arruda, T., Fieberg, J., Finch, E.A., Fiorenza, E.A., Fisher, D.N., Fontaine, A., Forstmeier, W.,
Fourcade, Y., Frank, G.S., Freund, C.A., Fuentes-Lillo, E., Gandy, S.L., Gannon, D.G., García-Cervigón,
A.I., Garretson, A.C., Ge, X., Geary, W.L., Géron, C., Gilles, M., Girndt, A., Gliksman, D., Goldspiel,
H.B., Gomes, D.G.E., Good, M.K., Goslee, S.C., Gosnell, J.S., Grames, E.M., Gratton, P., Grebe, N.M.,
Greenler, S.M., Griffioen, M., Griffith, D.M., Griffith, F.J., Grossman, J.J., Güncan, A., Haesen, S.,
Hagan, J.G., Hager, H.A., Harris, J.P., Harrison, N.D., Hasnain, S.S., Havird, J.C., Heaton, A.J., Herrera-
Chaustre, M.L., Howard, T.J., Hsu, B.-Y., Iannarilli, F., Iranzo, E.C., Iverson, E.N.K., Jimoh, S.O.,
Johnson, D.H., Johnsson, M., Jorna, J., Jucker, T., Jung, M., Kačergytė, I., Kaltz, O., Ke, A., Kelly, C.D.,
Keogan, K., Keppeler, F.W., Killion, A.K., Kim, D., Kochan, D.P., Korsten, P., Kothari, S., Kuppler, J.,
Kusch, J.M., Lagisz, M., Lalla, K.M., Larkin, D.J., Larson, C.L., Lauck, K.S., Lauterbur, M.E., Law, A.,
Léandri-Breton, D.-J., Lembrechts, J.J., L'Herpiniere, K., Lievens, E.J.P., De Lima, D.O., Lindsay, S.,

Luquet, M., MacLeod, R., Macphie, K.H., Magellan, K., Mair, M.M., Malm, L.E., Mammola, S.,
Mandeville, C.P., Manhart, M., Manrique-Garzon, L.M., Mäntylä, E., Marchand, P., Marshall, B.M.,
Martin, C.A., Martin, D.A., Martin, J.M., Martinig, A.R., McCallum, E.S., McCauley, M., McNew, S.M.,
Meiners, S.J., Merkling, T., Michelangeli, M., Moiron, M., Moreira, B., Mortensen, J., Mos, B.,
Muraina, T.O., Murphy, P.W., Nelli, L., Niemelä, P., Nightingale, J., Nilsonne, G., Nolazco, S., Nooten,
S.S., Novotny, J.L., Olin, A.B., Organ, C.L., Ostevik, K.L., Palacio, F.X., Paquet, M., Parker, D.J., Pascall,
D.J., Pasquarella, V.J., Paterson, J.H., Payo-Payo, A., Pedersen, K.M., Perez, G., Perry, K.I., Pottier, P.,
Proulx, M.J., Proulx, R., Pruett, J.L., Ramananjato, V., Randimbiarison, F.T., Razafindratsima, O.H.,
Rennison, D.J., Riva, F., Riyahi, S., Roast, M.J., Rocha, F.P., Roche, D.G., Román-Palacios, C.,
Rosenberg, M.S., Ross, J., Rowland, F.E., Rugemalila, D., Russell, A.L., Ruuskanen, S., Saccone, P.,
Sadeh, A., Salazar, S.M., Sales, K., Salmón, P., Sánchez-Tójar, A., Santos, L.P., Santostefano, F.,
Schilling, H.T., Schmidt, M., Schmoll, T., Schneider, A.C., Schrock, A.E., Schroeder, J., Schtickzelle, N.,
Schultz, N.L., Scott, D.A., Scroggie, M.P., Shapiro, J.T., Sharma, N., Shearer, C.L., Simón, D., Sitvarin,
M.I., Skupien, F.L., Slinn, H.L., Smith, G.P., Smith, J.A., Sollmann, R., Whitney, K.S., Still, S.M., Stuber,
E.F., Sutton, G.F., Swallow, B., Taff, C.C., Takola, E., Tanentzap, A.J., Tarjuelo, R., Telford, R.J.,
Thawley, C.J., Thierry, H., Thomson, J., Tidau, S., Tompkins, E.M., Tortorelli, C.M., Trlica, A., Turnell,
B.R., Urban, L., Van De Vondel, S., Van Der Wal, J.E.M., Van Eeckhoven, J., Van Oordt, F., Vanderwel,
K.M., Vanderwel, M.C., Vanderwolf, K.J., Vélez, J., Vergara-Florez, D.C., Verrelli, B.C., Vieira, M.V.,
Villamil, N., Vitali, V., Vollering, J., Walker, J., Walker, X.J., Walter, J.A., Waryszak, P., Weaver, R.J.,
Wedegärtner, R.E.M., Weller, D.L., Whelan, S., White, R.L., Wolfson, D.W., Wood, A., Yanco, S.W.,
Yen, J.D.L., Youngflesh, C., Zilio, G., Zimmer, C., Zimmerman, G.M., Zitomer, R.A., 2025. Same data,
different analysts: variation in effect sizes due to analytical decisions in ecology and evolutionary
biology. BMC Biol 23, 35. https://doi.org/10.1186/s12915-024-02101-x

Grente, O., 2021. Understanding the depredation process in grey wolf (Canis lupus) and its
interactions with lethal measures: focus on the French Alpine Arc (Biologie des populations et
Écologie). Thèse, Université de Montpellier, Montpellier, France.

Grente, O., Bauduin, S., Santostasi, N.L., Chamaillé-Jammes, S., Duchamp, C., Drouet-Hoguet, N.,
Gimenez, O., 2024. Evaluating the effects of wolf culling on livestock predation when considering
wolf population dynamics in an individual-based model. Wildlife Biology e01227.
https://doi.org/10.1002/wlb3.01227

Grente, O., Opitz, T., Duchamp, C., Drouet-Hoguet, N., Chamaillé-Jammes, S., Gimenez, O., 2025.
Spatio-temporal dynamics of attacks around deaths of wolves: A statistical assessment of lethal
control efficiency in France. https://doi.org/10.5281/zenodo.14893823

Hamilton, N.E., Ferry, M., 2018. ggtern: Ternary Diagrams Using ggplot2. Journal of Statistical
Software, Code Snippets 87, 1–17. https://doi.org/10.18637/jss.v087.c03

Harper, E.K., Paul, W.J., Mech, L.D., Weisberg, S., 2008. Effectiveness of Lethal, Directed Wolf-
Depredation Control in Minnesota. Journal of Wildlife Management 72, 778–784.
https://doi.org/10.2193/2007-273

Hernangómez, D., 2023. tidyterra: tidyverse Methods and ggplot2 Helpers for terra Objects
(manual). https://doi.org/10.5281/zenodo.6572471

Hijmans, R.J., Barbosa, M., Ghosh, A., Mandel, A., 2023. geodata: Download geographic data
(manual).

Hodgson, I.D., Redpath, S.M., Fischer, A., Young, J., 2019. Who knows best? Understanding the use
of research-based knowledge in conservation conflicts. Journal of Environmental Management 231,
1065–1075. https://doi.org/10.1016/j.jenvman.2018.09.023

Jędrzejewska, B., Jędrzejewski, W., Bunevich, A.N., Minkowski, L., Okarma, H., 1996. Population
dynamics of Wolves Canis lupus in Białowieża Primeval Forest (Poland and Belarus) in relation to
hunting by humans, 1847–1993. Mammal Review 26, 103–126. https://doi.org/10.1111/j.1365-
2907.1996.tb00149.x

Kaczensky, P., Ranc, N., Hatlauf, J., Payne, J.C., Acosta-Pankov, l., Álvares, F., Andrén, H., Andri, P.,
Aragno, P., Avanzinelli, E., Bagrade, G., Balys, V., Barroso, I., Bartol, M., Bassano, B., Bauduin, S.,
Bautista, C., Bedő, P., Belotti, E., Berezowska-Cnota, T., Bernicchi, L., Bijl, H., Bionda, R., Bišćan, A.,
Blanco, J.C., Bliem, K., Böcker, F., Bogdanović, N., Boiani, V., Bojda, M., Boljte, B., Bragalanti, N.,
Breitenmoser, U., Brøseth, H., Bučko, J., Budinski, I., Bufka, L., Černe, R., Cherepanyn, R., Chiriac, S.,
Ćirović, D., Csányi, S., DeAngelis, D., de Gabriel Hernando, M., Diószegi-Jelinek, L., Done, G., Drouet-
Hoguet, N., Duľa, M., Dutsov, A., Engleder, T., Fenchuk, V., Ferloni, M., Ferri, M., Filacorda, S., Finďo,
S., Fležar, U., Frangini, L., Frick, C., Fuxjäger, C., Galanaki, A., Genovesi, P., Gentile, D., Gervasi, V., Gil,
P., Giorgos, G., Gomerčić, T., Gonev, A., Gouwy, J., Gregorová, E., Groff, C., Gužvica, G.,
Hadžihajdarević, H., Heikkinen, S., Heltai, M.G., Henttonen, H., Herrero, A., Hoxha, B., Huber, D.,
Iliopoulos, Y., Imeri, M., Ioannis, G., Ivanov, G., Jan, M., Jansman, H., Jeremić, J., Jerina, K., Kapo, N.,
Karaiskou, N., Karamanlidis, A., Kindberg, J., Kluth, G., Knauer, F., Kojola, I., Kominos, T., Konec, M.,
Koubek, P., Krausová, J., Krofel, M., Krojerová, J., Kubala, J., Kübarsepp, M., Kunz, F., Kusak, J., Kutal,
M., Kyriakidis, S., La Morgia, V., Lajçi, F., Lammertsma, D., Lapini, L., Latini, R., Lemaitre, P.-L.,
Licoppe, A., Linnell, J.D.C., López-Bao, J.V., Majic Skrbinsek, A., Männil, P., Marucco, F., Melovski, D.,
Mengüllüoğlu, D., Mergeay, J., Mertzanis, Y., Meytre, S., Mináriková, T., Mokrý, J., Molinari, P.,
Molinari-Jobin, A., Moreno, I., Mysłajek, R.W., Nägele, O., Napotnik, I., Nezaj, M., Nowak, S., Olsen,
K., Omeragić, J., Oreiller, P., Ornicāns, A., Ozoliņš, J., Palomero, G., Pavlov, A., Perovic, A., Pesaro, S.,
Pilāte, D., Pimenta, V., Poledník, L., Pop, M.I., Prakapchuk, V., Pylidis, C., Quenette, P.-Y., Rauer, G.,
Reinhardt, I., Reljić, S., Rigg, R., Riva, V., Rodekirchen, A.M., Ruņģis, D.E., Šálek, M., Salvatori, V.,
Satra, M., Schally, G.T., Schley, L., Selanec, I., Selimovic, A., Selva, N., Sentilles, J., Shyti, I., Signer, S.,
Simčič, G., Sindičić, M., Škapur, V., Skrbinšek, T., Smith, A.F., Smitskamp, L., Solovej, I., Špinkytė-
Bačkaitienė, R., Stepanova, A., Stergar, M., Sterrer, U., Stojanov, A., Šuleková, D., Sunde, P., Šver, L.,
Szewczyk, M., Topličanec, I., Tosoni, E., Trajče, A., Trbojević, I., Trbojević, T., Tsalazidou, T.-M.,
Tsingarska-Sedefcheva, E., Ursitti, J., Valtonen, M., Vandel, J.-M., Vanpé, C., Veeroja, R., von Arx, M.,
Vorel, A., Vykhor, B., Weber, H., Woelfl, S., Yamelynets, T., Zimmermann, F., Zlatanova, D., Žuglić, T.,
Zukal, J., Žunna, A.B., Luigi, 2024. Large carnivore distribution maps and population updates 2017 –
2022/23 (B.4 Update of the distribution maps No. 09.0201/2023/907799/SER/ENV.D.3), Support for
Coexistence with Large Carnivores. IUCN/SSC Large Carnivore Initiative for Europe (LCIE).

Kassambara, A., 2023. ggpubr: "ggplot2" based publication ready plots (manual).

Knol, M.J., Pestman, W.R., Grobbee, D.E., 2011. The (mis)use of overlap of confidence intervals to
assess effect modification. Eur J Epidemiol 26, 253–254. https://doi.org/10.1007/s10654-011-9563-8

Knopf, G., 2019. Délimitation des massifs alpins. Nivologie des Alpes du Sud. URL
https://nivo06.knobuntu.fr/index.php?option=com_content&view=article&id=400:delimitation-des-
massifs-alpins&catid=63&Itemid=408 (accessed 11.27.23).

Kompaniyets, L., Evans, M.A., 2017. Modeling the relationship between wolf control and cattle
depredation. PLoS ONE 12, e0187264. https://doi.org/10.1371/journal.pone.0187264

Kutal, M., Duľa, M., Selivanova, A.R., López-Bao, J.V., 2023. Testing a conservation compromise: No
evidence that public wolf hunting in Slovakia reduced livestock losses. Conservation Letters n/a,
e12994. https://doi.org/10.1111/conl.12994

Lennox, R.J., Gallagher, A.J., Ritchie, E.G., Cooke, S.J., 2018. Evaluating the efficacy of predator
removal in a conflict-prone world. Biological Conservation 224, 277–289.
https://doi.org/10.1016/j.biocon.2018.05.003

Linnell, J.D.C., Cretois, B., 2018. Research for AGRI Committee – The revival of wolves and other large
predators and its impact on farmers and their livelihood in rural regions of Europe. European
Parliament, Policy Department for Structural and Cohesion Policies, Brussels.

Linnell, J.D.C., Odden, J., Smith, M., E., Aanes, R., Swenson, J.E., 1999. Large carnivores that kill
livestock: do "problem individuals" really exist? Wildlife Society Bulletin 27, 698–705.

Linnell, J.D.C., Rondeau, D., Reed, D.H., Williams, R., Altwegg, R., Raxworthy, C.J., Austin, J.D., Hanley,
N., Fritz, H., Evans, D.M., Gordon, I.J., Reyers, B., Redpath, S., Pettorelli, N., 2010. Confronting the
costs and conflicts associated with biodiversity. Animal Conservation 13, 429–431.
https://doi.org/10.1111/j.1469-1795.2010.00393.x

Lorand, C., Robert, A., Gastineau, A., Mihoub, J.-B., Bessa-Gomes, C., 2022. Effectiveness of
interventions for managing human-large carnivore conflicts worldwide: Scare them off, don't
remove them. Science of The Total Environment 838, 156195.
https://doi.org/10.1016/j.scitotenv.2022.156195

Louvrier, J., Duchamp, C., Lauret, V., Marboutin, E., Cubaynes, S., Choquet, R., Miquel, C., Gimenez,
O., 2018. Mapping and explaining wolf recolonization in France using dynamic occupancy models
and opportunistic data. Ecography 41, 647–660. https://doi.org/10.1111/ecog.02874

Lute, M.L., Carter, N.H., López-Bao, J.V., Linnell, J.D.C., 2018. Conservation professionals agree on
challenges to coexisting with large carnivores but not on solutions. Biological Conservation 218, 223–
232. https://doi.org/10.1016/j.biocon.2017.12.035

Maesono, Y., 2011. Jackknife, in: Lovric, M. (Ed.), International Encyclopedia of Statistical Science.
Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 697–699. https://doi.org/10.1007/978-3-642-
04898-2_317

Marucco, F., Reinhardt, I., Avanzinelli, E., Zimmermann, F., Manz, R., Potočnik, H., Černe, R., Rauer,
G., Walter, T., Knauer, F., Chapron, G., Duchamp, C., 2023. Transboundary Monitoring of the Wolf
Alpine Population over 21 Years and Seven Countries. Animals (Basel) 13.
https://doi.org/10.3390/ani13223551

Mattioli, L., Capitani, C., Gazzola, A., Scandura, M., Apollonio, M., 2011. Prey selection and dietary
response by wolves in a high-density multi-species ungulate community. Eur J Wildl Res 57, 909–922.
https://doi.org/10.1007/s10344-011-0503-4

Merz, L.M., Bruskotter, J.T., Carter, N.H., 2025. Elusive effects of legalized wolf hunting on human-
wolf interactions. Sci Adv 11. https://doi.org/10.1126/sciadv.adu8945

Meuret, M., Moulin, C.-H., Bonnet, O., Garde, L., Nozières-Petit, M.-O., Lescureux, N., 2020. Missing
shots: has the possibility of shooting wolves been lacking for 20 years in France's livestock protection
measures? Rangel. J. 42, 401–413.

Milleret, C., Duchamp, C., Bauduin, S., Kaerle, C., Pirog, A., Queney, G., Gimenez, O., 2025a.
Estimating wolf population size in France using non-invasive genetic sampling and spatial capture
recapture models. https://doi.org/10.48550/arXiv.2508.12434

Milleret, C., Dupont, P., Dey, S., Brøseth, H., Kindberg, J., Turek, D., de Valpine, P., Åkesson, M.,
Wabakken, P., Zimmermann, B., Bischof, R., 2025b. Map of death: spatially explicit mortality of the
grey wolf. Proceedings of the Royal Society B: Biological Sciences 292, 20250948.
https://doi.org/10.1098/rspb.2025.0948

Ministère de la Transition Ecologique et Solidaire, 2020. Arrêté du 30 décembre 2019 portant
expérimentation de diverses dispositions en matière de dérogations aux interdictions de destruction
pouvant être accordées par les préfets concernant le loup (Canis lupus), TREL1936320A.

Ministère de la Transition Ecologique et Solidaire, 2018. Arrêté du 19 février 2018  fixant les
conditions et limites dans lesquelles des dérogations aux interdictions de destruction peuvent être
accordées par les préfets concernant le loup (Canis lupus).

Neuwirth, E., 2022. RColorBrewer: ColorBrewer palettes (manual).

Nichols, J.D., Cooch, E.G., 2025. Predictive models are indeed useful for causal inference. Ecology
106, e4517. https://doi.org/10.1002/ecy.4517

Office National de la Chasse et de la Faune Sauvage, 2019. Bilan hivernal de la population de loup : le
seuil des 500 loups est atteint. Office National de la Chasse et de la Faune Sauvage.

Padgham, M., Rudis, B., Lovelace, R., Salmon, M., 2017. osmdata. Journal of Open Source Software 2,
305. https://doi.org/10.21105/joss.00305

Pebesma, E., Bivand, R., 2023. Spatial data science: With applications in R. Chapman and Hall/CRC.

Pedersen, T.L., 2022a. ggfx: Pixel Filters for "ggplot2" and "grid" (manual).

Pedersen, T.L., 2022b. patchwork: The composer of plots (manual).

Petridou, M., Youlatos, D., Lazarou, Y., Selinides, K., Pylidis, C., Giannakopoulos, A., Kati, V.,
Iliopoulos, Y., 2019. Wolf diet and livestock selection in central Greece. Mammalia 83, 530–538.
https://doi.org/10.1515/mammalia-2018-0021

Poudyal, N., Baral, N., Asah, S.T., 2016. Wolf Lethal Control and Livestock Depredations: Counter-
Evidence from Respecified Models. PLoS ONE 11, e0148743.
https://doi.org/10.1371/journal.pone.0148743

Poulle, M.-L., Carles, L., Lequette, B., 1997. Significance of ungulates in the diet of recently settled
wolves in the Mercantour mountains (southeastern France). Revue d'Ecologie, Terre et Vie 52, 657–
368.

R Core Team, 2023. R: A language and environment for statistical computing (manual). R Foundation
for Statistical Computing, Vienna, Austria.

Railsback, S.F., Gallagher, C.A., Grimm, V., McCary, M.A., Harvey, B.C., 2025. Empirical ecology to
support mechanistic modelling: Different objectives, better approaches and unique benefits.
Methods Ecol Evol 16, 1564–1573. https://doi.org/10.1111/2041-210X.70083

Réseau Loup-Lynx, 2024. Bilan annuel du suivi de la population de loups 2022 (No. 17), Lettre
d'information du réseau Loup-lynx. Office Français de la Biodiversité.

Santiago-Avila, F.J., Cornman, A.M., Treves, A., 2018. Killing wolves to prevent predation on livestock
may protect one farm but harm neighbors. PLoS ONE 13, e0189729.
https://doi.org/10.1371/journal.pone.0189729

Silge, J., Robinson, D., 2016. tidytext: Text mining and analysis using tidy data principles in R. JOSS 1.
https://doi.org/10.21105/joss.00037

Šuba, J., Žunna, A., Bagrade, G., Done, G., Ornicāns, A., Pilāte, D., Stepanova, A., Ozoliņš, J., 2023.
Does Wolf Management in Latvia Decrease Livestock Depredation? An Analysis of Available Data.
Sustainability 15. https://doi.org/10.3390/su15118509

Treves, A., Karanth, K.U., 2003. Human-Carnivore Conflict and Perspectives on Carnivore
Management Worldwide. Conservation Biology 17, 1491–1499. https://doi.org/10.1111/j.1523-
1739.2003.00059.x

Treves, A., Krofel, M., McManus, J., 2016. Predator control should not be a shot in the dark. Front
Ecol Environ 14, 380–388. https://doi.org/10.1002/fee.1312

Treves, A., Krofel, M., Ohrens, O., van Eeden, L.M., 2019. Predator Control Needs a Standard of
Unbiased Randomized Experiments With Cross-Over Design. Front. Ecol. Evol. 7, 462.
https://doi.org/10.3389/fevo.2019.00462

van Eeden, L.M., Crowther, M.S., Dickman, C.R., Macdonald, D.W., Ripple, W.J., Ritchie, E.G.,
Newsome, T.M., 2018. Managing conflict between large carnivores and livestock: Livestock-
Carnivore Conflict. Conservation Biology 32, 26–34. https://doi.org/10.1111/cobi.12959

Wagner, K., K., Conover, M., R., 1999. Effect of Preventive Coyote Hunting on Sheep Losses to Coyote
Predation. The Journal of Wildlife Management 63, 606–612. https://doi.org/10.2307/3802649

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G., Hayes,
A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J.,
Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019.
Welcome to the tidyverse. Journal of Open Source Software 4, 1686.
https://doi.org/10.21105/joss.01686

Wickham, H., Seidel, D., 2022. scales: Scale functions for visualization (manual).

Wielgus, R.B., Peebles, K.A., 2014. Effects of Wolf Mortality on Livestock Depredations. PLoS ONE 9,
e113505. https://doi.org/10.1371/journal.pone.0113505

Žunna, A., Ruņģis, D.E., Ozoliņš, J., Stepanova, A., Done, G., 2023. Genetic Monitoring of Grey Wolves
in Latvia Shows Adverse Reproductive and Social Consequences of Hunting. Biology 12, 1255.
https://doi.org/10.3390/biology12091255

1056