Integration of multi-omics data and deep phenotyping of potato enables novel insights into single- and combined abiotic stress responses

Maja Zagorščak[1]*, Lamis Abdelhakim[2]*, Natalia Yaneth Rodriguez-Granados[3]*, Jitka Široká[4]*, Arindam Ghatak[5]*, Carissa Bleker[1], Andrej Blejec[1], Jan Zrimec[1], Ondřej Novák[4], Aleš Pěnčík[4], Špela Baebler[1], Lucia Perez Borroto[6], Christian Schuy[7], Anže Županič[1], Leila Afjehi-Sadat[8], Bernhard Wurzinger[5], Wolfram Weckwerth[5,9], Maruša Pompe Novak[1,10], Marc R. Knight[11], Miroslav Strnad[4], Christian Bachem[6], Palak Chaturvedi[5], Sophia Sonnewald[7+], Rashmi Sasidharan[3+], Klára Panzarová[2+], Kristina Gruden[1+#], Markus Teige[5+#]

Affiliations:
[1] Department of Biotechnology and Systems Biology, National Institute of Biology, Večna pot 121, 1000 Ljubljana, Slovenia
[2] PSI (Photon Systems Instruments), spol. s r.o., Prumyslova 470, CZ-664 24 Drásov, Czech Republic
[3] Plant Stress Resilience, Institute of Environmental Biology, Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, The Netherlands
[4] Laboratory of Growth Regulators, Palacký University in Olomouc & Institute of Experimental Botany AS CR, Šlechtitelů 27, Olomouc, 779 00, Czech Republic
[5] Molecular Systems Biology (MOSYS), Department of Functional and Evolutionary Ecology, University Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[6] Wageningen University and Research, Plant Breeding, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands
[7] Department Biologie, Lehrstuhl für Biochemie, Friedrich-Alexander-Universität Erlangen-Nürnberg, Staudstr. 5, 91058 Erlangen, Germany
[8] Mass Spectrometry unit, Research Support Facilities, Faculty of Life Sciences, University Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[9] Vienna Metabolomics Center (VIME), University Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[10] School for Viticulture and Enology, University of Nova Gorica, Gladni trg 8, 5271 Vipava, Slovenia
[11] Department of Biosciences, Durham University, South Road, Durham DH1 3LE, United Kingdom

* shared first authorship

+ shared last authorship

corresponding authors

Keywords: potato, *Solanum tuberosum*, abiotic stress responses, heat, drought, waterlogging, multi-omics, integrative omics, adaptomics, panomics

1 Abstract

Potato is highly water and space efficient but susceptible to abiotic stresses such as heat,
drought, or flooding, which are severely exacerbated by climate change. Understanding of crop
acclimation to abiotic stress, however, remains limited. Here, we present a comprehensive
molecular and physiological high-throughput profiling of potato (*Solanum tuberosum*, cv.
Desirée) under heat, drought and waterlogging applied as single stresses or in combinations
designed to mimic realistic future scenarios. Stress-responses were monitored via daily
phenotyping and multi-omics analyses of leaf samples comprising transcriptomics, proteomics,
metabolomics and hormonomics at several timepoints during and after stress treatments.
Additionally, critical metabolites of tuber samples were analysed at the end of the stress period.
Integrative analysis of multi-omics data was performed using a bioinformatic pipeline, which was
established here, based on machine learning and knowledge networks. Overall, waterlogging
had the most immediate and dramatic effects on potato plants, interestingly activating ABA-
responses similar to drought stress. In addition, we observed distinct stress signatures at
multiple molecular levels in response to heat or drought and to a combination of both. In
response to all treatments, we found a downregulation of photosynthesis at different molecular
levels, an accumulation of minor amino acids and diverse stress induced hormones. Our
integrative multi-omics analysis provides global insights into plant stress responses, facilitating
improved breeding strategies towards climate-adapted potato varieties.
20
21 (219 words)

22
23
24 **One Sentence Summary**
25
26 Integrated multi-omics analysis of high-throughput phenotyping in potato reveals distinct
27 molecular signatures of acclimation to single and combined abiotic stresses related to climate
28 change.
29
30

1   INTRODUCTION

Improving crop resilience to climate change is a major challenge of modern agriculture (Bailey-
Serres et al., 2019; Rivero et al., 2022). High-yielding crop varieties including potato (*Solanum*
tuberosum), are vulnerable to heat, drought, and flooding (Benitez-Alfonso et al., 2023;
Zandalinas et al., 2023; Renziehausen et al., 2024; Sato et al., 2024). These environmental
stresses affect plant growth, source-sink relationships, sugar and hormone metabolism, among
other processes, which in turn, negatively impact product yield and nutritional status (Lal et al.,
2022). Potato is particularly sensitive to waterlogging (Jovović et al., 2021), and flooding of the
fields can ruin the entire harvest within a few days. Since global warming is increasing the
occurrence of such extreme weather events, crop productivity worldwide is under considerable
threat (FAO, 2023). To ensure future food security, there is an urgent need for sustainable
farming practices including the development of stress tolerant varieties with consistent yields
(Dahal et al., 2019; Lal et al., 2022).

There is already a good understanding of how plants react to single abiotic stresses, which have
profound effects on plant metabolism and development. The primary effects of abiotic stress are
generation of reactive oxygen species (ROS), destabilisation of proteins and changes in
enzyme efficiencies and membrane fluidity and integrity (Zhang et al., 2022). Together, these
impacts reduce plant productivity through changes in photosynthetic capacity, hormone
balance, transport of assimilates from source to sink as well as transport of soil nutrients and
water by the roots. In addition, species-specific vulnerabilities impact agronomic productivity,
such as for instance tuber initiation and tuber growth dynamics with potato. Potato tuber
formation and growth is dependent on tuberisation signals produced in source leaves, such as
the mobile tuberigen StSP6A (Navarro et al., 2011) that also regulates directional transport of
sucrose to the developing tuber (Abelenda et al., 2019). Heat, drought and flooding trigger
strong changes in gene expression and thereby strongly interfere with the regulation of
flowering and tuberisation by the photoperiodic pathway. This leads to a delay in tuberisation
and anomalies in subsequent tuber development such as second-growth and/or internal defects
which together severely impacts marketable yields of the tuber crop (Dahal et al., 2019; Lal et
al., 2022).

On the other hand, the response of plants to combined stresses is unique and cannot be
extrapolated from the response to the corresponding individual stresses (Mittler, 2006).
Considering the increasing occurrences of simultaneous or sequential abiotic stresses in the

field, the relative lack of knowledge on multi-stress resilience is a major shortfall that hinders the
ability to develop effective strategies for crop improvement. Accordingly, the question of how
combinations of different stresses impact plants has recently gained a lot of interest (Zandalinas
et al., 2021). Several studies on combinatorial stress effects have been performed, especially
studying the physiological and molecular responses to combined heat- and drought stress in
potato (Demirel et al., 2020), wheat (Manjunath et al., 2023) and tomato (Zeng et al., 2024). In
nature, heat and drought often occur together, resulting in different physiological responses as
compared to individual stresses. For example, under heat stomatal conductance and
transpiration are increased to reduce leaf temperature, whilst under drought, stomata are closed
to avoid water loss, which leads to a strongly reduced CO_2-assimilation (Zhang and Sonnewald,
2017). The final phenotypic output in a combined stress scenario greatly depends on synergistic
and antagonistic interactions between stress-specific signalling and response pathways. These
interactions can be regulated at various levels (gene expression to metabolism), and on
different scales (cell to system), thus resulting in complex regulatory network perturbations.
Therefore, as information gained by extrapolating from studies on individual stressors is limited,
it is crucial to increase our understanding of crop responses in multi-stress situations.

To this end, high-throughput phenotyping platforms and integrative omics technologies can
measure molecular mechanisms at multiple levels and in multiple processes simultaneously.
This can help us obtain a comprehensive understanding of the intricate dynamics of plant-
environment interactions (Yang et al., 2020; Hall et al., 2022; Zhang et al., 2022). Here,
advanced data integration pipelines can aid with unbiased integration and systematic extraction
of biological knowledge from large multi-omics data sets (Cembrowska-Lech et al., 2023).
However, despite the increasing application of high-throughput approaches in agricultural and
plant research, only a handful of studies have addressed the problem of data integration from
comprehensive multi-omics data (Jamil et al., 2020) . Therefore, to enable molecular insights
across various system levels and disentangle the intricate physiological and molecular crosstalk
in the context of non-additive effects of different stress combinations, it is imperative to develop
and apply multi-omics integrative approaches that leverage statistics, machine learning, and
graph theory.

In this study, we aimed to increase knowledge on multiple abiotic stress responses of potato
plants and to integrate this into a complex knowledge network. Therefore, a comprehensive
assessment of potato responses to single and combined heat, drought, and waterlogging stress

was performed. Using the cv. Désirée, a widely used moderately stress-resistant potato cultivar,

we monitored dynamic changes in morphological, physiological as well as biochemical and

molecular responses under stress conditions. With the application of high throughput

phenotyping, multi-omics technologies, prior-knowledge and multi-level integration approaches,

we identified important molecular signatures, unique to single and different stress combinations.

These results can guide the development of diagnostic markers for rapid detection of stress,

allowing for earlier agricultural interventions to enhance plant resilience towards abiotic stress

and development of novel marker-assisted breeding programs for climate-resilient crops

(Weckwerth et al., 2020; Mishra et al., 2024).

RESULTS

High throughput physiological phenotyping of potato in multi-stress conditions

This study aims to increase understanding of acclimation of potato plants to abiotic stresses by

focusing on most likely weather (stress) scenarios in future agricultural farming: individual heat,

drought and waterlogging stress, and realistic combinations thereof. We studied the cv. Desirée,

a widely used moderately stress-resistant potato cultivar, as a model. Daily phenotyping using

image-based sensors and regular sampling of leaf material for downstream molecular and

biochemical analysis allowed better insights into the mechanisms of the stress response. Single

and combined stresses were applied to plants during the tuber initiation stage at day 32 and 39

respectively, according to the scheme in Figure 1A (for more details see Supp. Table 1). Daily

phenotyping of plants (control and stress conditions) was performed using the PlantScreen[TM]

phenotyping platform to assess quantitative morphological and physiological traits (Supp. Table

2), employing multiple imaging sensors (Figure 1B). Plants were light-adapted before

chlorophyll fluorescence measurements to ensure steady conditions prior to the measurements.

Effects of single and combined stresses on potato growth and morphology

27
By using RGB side and top view imaging we monitored changes in plant growth dynamics

during the stress treatments and recovery phases, including traits such as plant volume, area,

height and compactness. The exponential growth of plants grown under control conditions was

negatively impacted to various extent by different stresses (Figure 1C, Supp. Figure 1). Thus,

under heat stress (H, 30°C during day, 28°C during night) the rate of biomass accumulation

(plant volume) was decreased after three days similarly to drought stress (D, 30% of field

capacity) (Figure 1C, Supp. Figure 1A-C). Additional morpho-physiological alterations became

visible in response to heat and waterlogging (W). Heat clearly reduced plant height and caused

the typical hyponastic movement of leaves (Figure 1C and Supp. Figure 1C). The negative

effects of heat became more severe when combined with drought (HD, water withdrawal

starting after 7 days of H) (Supp. Figure 1C). Under HD, plants phenotypically more resembled

heat-stressed plants, e.g. with respect to top area and compactness, however, with a clearly

more negative effect (Supp. Figure 1B, 1D). Waterlogging led to an epinastic leaf movement

which was accompanied by growth arrest and significant decrease in the top area, compactness

and relative growth rate (RGR) that were observed after one day (Figure 1C, Supp. Figure 1A,

1B, 1D, 1E, Supp. Table 3). In the 3rd week, when the single and HD stress treatments were

finished (at treatment day 15), plants recovered well from H and HD, which was clearly reflected

by resumption of growth, but not from W stress (Figure 1C, Supp. Figure 1B).

Plant performance was the worst in the triple-stress condition (HDW), where 7 days of H were

followed by 7 days of combined HD and 7 days of W. Interestingly, during the first three days of

W that followed the period of heat and drought plants were growing very fast, but with a

prolonged stress exposure, plants collapsed, as indicated from both RGB side and top view

images, from the plant volume and growth dynamics as shown in RGR and other measured

morphological traits (Figure 1C, Supp. Figure 1, Supp. Table 3).

Photosynthesis is most affected by heat- and drought stress

To evaluate photosynthetic performance under single and multiple stresses, a broad range of

physiological traits were extracted from chlorophyll fluorescence images and analysed (Figure

2). Top view images of the operating efficiency of photosystem II in light steady state (QY_Lss)

clearly showed the negative impact of stress on photosynthetic capacity in all stress treatments,

indicated by the reduction of QY_Lss, with D stress causing only a weak negative effect (Figure

2A, 2B).   Moreover, steady-state fluorescence of maximum efficiency of PSII in the light

(F_v/F_m_Lss) showed a significant decrease after 3 days in W alone and when W followed a

period of HD. There was also a progressive decrease in steady-state estimation of the fraction

of open reaction centres in PSII in the light (qL_Lss) under whenever W was applied (Figure 2C,

2D, Supp. Table 3). Both parameters decreased even further after stress recovery following

triple stress, indicating a high stress level.

A decrease in qL_Lss was also observed after one day of H and remained constantly lower than

in other conditions, while D caused no significant effect on these parameters (Figure 2D, Supp.

Table 3). By applying drought in addition to heat stress, an increase in qL_Lss as compared to

H alone was observed. After H and HD stress, qL_Lss values did not reach control levels till the

end of the experiment at day 21 (Figure 2C and 2D), suggesting that photosynthesis is

enduringly affected. In addition, changes in canopy temperature (ΔT) were deduced from the

thermal imaging, while water use efficiency (WUE) was calculated based on plant volume and

water consumption. The rapid increase in ΔT and WUE under W and HDW was most likely

caused by prompt stomata closure (Figure 2E, 2F). The strong response remained over the

entire stress period and plants did not recover from both stress treatments. A steady increase in

ΔT and WUE was observed starting at three days in D, suggesting that the stress was

recognised, and the plants responded by closing stomata. When D stress was released on day

15, the plants recovered immediately (Figure 2E, 2F). An opposing response was observed in

H, where ΔT decreased together with an increase in water consumption, thus indicating

enhanced leaf cooling (Figure 2E, Supp. Table3, 4). During combined HD stress, it appeared

that the drought response became dominant over the heat responses in the physiological

responses as indicated by the increase in ΔT and WUE. At the time when the drought stress

was applied in addition to heat, as both parameters behaved more like drought stress

responses as compared to H alone.

Stress combinations and waterlogging have strong effects on potato yield

At the end of the phenotyping, plants were harvested to assess the total biomass accumulation

and tuber yield (Figure 2G and 2H). Single H stress led to a slightly higher tuber number (Figure

2G). Harvest index was significantly reduced by H as well as by HD, while D alone did not affect

final tuber yield (Figure 2H). W stress strongly inhibited tuber formation and growth and only a

few tubers were formed, leading to a significant reduction in the harvest index compared to the

control treatment (Figure 2H). A combination of all stress factors abolished tuber formation,

reflecting the (near) lethal effect of HDW (Figure 2G, 2H).

Negative effects of the stress treatments on tubers were also observed at the metabolic level

(Supp. Figure 2). Thus, starch content was significantly lower under H, HD and W stress, while

drought stress alone has no negative impact. The accumulation of hexoses under H and HD

may hint to an increased starch degradation and / or to a reduced starch biosynthesis. W

caused a strong accumulation of almost all amino acids, most likely caused by protein

degradation and a hampered metabolism (Supp. Figure 2).

Molecular responses across omics levels reveal mechanistic insights into

multi-stress acclimation

In addition to the morphological and physiological measurements (68 variables, Figure 3D,

Supp. Table 1, 2), leaf samples were taken for parallel multi-omics analysis. Samples from

(mature) leaves two and three were pooled, homogenized and used for further analysis (Figure

3A). For each of the treatments the fast response (one day post treatment) and the status at the

end of a prolonged stress duration (7 or 14 days of stress) was investigated (sampling points

see Figure 1A). While the proteome analysis was untargeted (4258 identified proteins, Supp.

Table5, 6), other omics analyses were targeted comprising 14 pre-selected transcriptional

marker genes involved in stress response and tuberisation, 13 phytohormones encompassing

abscisic acid, ABA; jasmonic acid, JA; salicylic acid, SA; indole-3-acetic acid IAA, and their

derivatives as well as 22 metabolites encompassing amino acids and sugars (Supp. Table4). To

identify processes regulated on proteomics level we performed gene set enrichment analysis

(GSEA, Supp. Table6).

A multi-level data integration protocol was developed to investigate plant signalling/responses

across the different omics levels (Figure 3B). First, we investigated data distribution by

multidimensional scaling. (Figure 3C). This graph shows a clear clustering aside of samples

taken after 7 and 8 days of waterlogging. Therefore, only data from the first week of

waterlogging were included in further analyses, taking also into consideration that after two

weeks of waterlogging all plants were severely damaged. The overview of data distribution also

revealed that the most distant physiological state was that of plants exposed to triple stress

(HDW): first one week of heat, followed by one week of heat combined with drought, and finally

one week of waterlogging (Figure 3C). Because the triple stress treatment turned out to be very

harsh and plants were severely affected in both above ground and below ground growth, we

also excluded data from these samples from all further analyses. Next, we reduced the number

of variables obtained on phenomics and proteomics levels to equalise numbers of variables

across different analysed levels. In order to identify the most informative variables, feature

selection using random forest with recursive feature elimination was conducted on the

phenomics data, keeping 6 variables for downstream analysis (Figure 3D: qL, Fv/Fm, top area,

delta T, compactness and water consumption). The proteomics dataset was reduced to keep

only proteins that were identified as differentially abundant in any comparison of stress vs.

control (135 proteins) and were functionally assigned to pathways that were studied also on

other levels (36 proteins, related to photosynthesis, metabolism of sugars and amino acids,

hormone metabolism and signalling, ROS signalling and stress pathways).

In addition, correlation analysis within each level of omics data was performed, revealing that

these components are only weakly connected in control conditions, while in both heat or

drought, they are highly correlated to each other (see e.g. for hormones and transcripts, Supp.

Figure 3A, B). More severe stresses, such as the combined heat and drought stress and

waterlogging, however, broke this link, suggesting a disorganisation of signalling responses.

The canonical correlation analysis between components of different molecular levels similarly

showed low connection in control samples. In stressed samples, blocks of components

appeared to be strongly regulated, each specific to a particular stress (Supp. Figure 3C).

Variables measured on different omics levels were integrated into a metabolism and signalling

cascade-based knowledge network to capture events at the molecular level (Figure 4A). Finally,

we superimposed the measured data onto this mechanistic knowledge network and visualised

them in parallel for all omics levels per each analysed condition compared to control (Figure 4B,

Supp. Table 3). This provides a comprehensive overview of how these stresses rewire

biochemical pathways and physiological processes. These networks were used for

interpretation of processes in single and combined H and D stress as well as for waterlogging

(W) and are described in the subsequent sections.

Metabolic and molecular responses to individual and combined heat- and drought stress exhibit combinatorial and distinct signatures

While heat stress was effective immediately, drought stress, applied by water withdrawal on day

seven in our set-up, became effective gradually within three days, visible by an increase in the

ΔT values (Figure 2E). Similar to previous reports (Demirel et al., 2020; Zaki and Radwan,

2022), we found that Desirée was moderately drought tolerant and exhibited only minor

morphological and physiological responses at the moderate stress level applied in our study

(30% water capacity of the soil) and the plants fully recovered when the stress treatment was

finished. The potato plants clearly responded to elevated temperatures (heat stress, H) with

morphological adaptation like the upward movement of leaves (Figure 1A), which is part of

thermo-morphogenic responses (Quint et al., 2016). Previous work showed that heat stress
caused an altered biomass allocation between shoots and tubers of potato plants, with less
assimilates allocated to developing tubers (Hancock et al., 2014; Hastilestari et al., 2018),
leading to decreased tuber yield (Figure 2G) and starch accumulation and a decreased harvest
index which was also seen in our study (Figure 2H and Supp. Figure 2).
To investigate the effect of heat and/or drought stress on leaf carbohydrate metabolism,
contents of soluble sugars and starch were measured (Figure 5A). While sucrose levels did not
change (Supp. Table 3, there was about twofold increase in the amounts of fructose and
glucose after 14 days of H and at day 7 of D and HD. The soluble sugars may act as
osmo-protectants under these stress conditions. In addition, we found that protein levels of
enzymes involved in glycolysis or sucrose degradation were upregulated in H (at 7 days) and
combined HD (day 14) as indicated by gene set enrichment analysis which summarizes the
complex proteomics data set (Figure 5D and Supp. Table 6). This may indicate an increased
demand for energy and/ or building blocks for stress defense responses. Under these conditions
also less starch accumulated, visible at day one and eight in H and at day one HD, suggesting
lower carbon availability due to a disturbed photosynthesis and/or an increased demand for
other processes. Expression of the sugar efflux carrier *SWEET11* was upregulated at the end of
both H and D stress presumably to maintain sucrose loading into the phloem and carbon
allocation to sink tissue to counterbalance the decreased carbon assimilation rate of source
leaves. Together with the reduced photosynthetic efficiency, that was measured, these data
imply that a lower amount of carbon is assimilated, especially under H and HD, and available for
allocation to sink organs, such as growing tubers. This most likely contributes to the impaired
tuber growth and lower tuber starch accumulation found under these conditions (Supp. Figure
2).
One key player stimulating tuberisation and tuber growth is the tuberigen *SP6A*. Its expression
was downregulated during the first week of H and in combined HD (Figure 5C). During longer
heat exposure, expression levels of *SP6A* were similar to control levels, but remained low in
HD. Drought alone had little effect on *SP6A*, which is consistent with the low impact on final
tuber yield. The transcriptional regulator protein *Constans-like 1* (CO) was described to act as a
negative regulator of *SP6A* expression (Abelenda et al., 2016). *CO* was upregulated within 7
days of drought stress (D), and it increased with duration of heat, but was unaffected by HD
combination (Figure 5C). Hence, in our experiment, the transcript levels of *CO* did not always
change in the opposite direction as *SP6A*, assuming additional regulatory mechanisms may act
under stress conditions.

Considering the changes in amino acids, the most striking finding was the strongly elevated
levels of histidine (His) in all three stress treatments, with the highest amounts detected in
combined HD stress (Figure 5A). This was accompanied by a significant increase of many
(minor) amino acids, in particular iso-leucine (Ile) and other branched chain amino acids
(BCAs). This observation was in line with previous reports on combined heat- and drought
stress in potato (Demirel et al., 2020), although it's cause and importance needs further
investigations. Accordingly, at the proteome level, proteins involved in BCA synthesis were
significantly enriched among the ones with increased levels in stress (Figure 5D, 5E, Supp.
Table 5, 6).
10
Proline is known as an important regulator of osmotic potential that protects cells by stabilizing
proteins and scavenging of reactive oxygen species. Its amount was elevated at day 7 of D
stress (sampling day 14) (Figure 5A). This was consistent with increased transcript levels of
P5CS (*Pyrroline-5-carboxylate synthase*), the key enzyme for proline synthesis, and of *RD29*
(*Responsive to Desiccation*), both being well-known (drought-) stress marker genes.
Consistently, levels of ABA, the key phytohormone that induced stomal closing, proline
accumulation and other drought stress responses (Cutler et al., 2010; Zhang et al., 2022), were
elevated after 7 days of D but were clearly reduced in H, while no changes were detected in HD
stress. Interestingly, the levels of phaseic acid (PA), and dihydrophaseic acid (DPA), two
breakdown products of ABA, were lower at one day of D but significantly higher after 7 days in
D. The strongest accumulation of DPA levels was detected in the HD treatment, in which DPA
levels were elevated already after one day and further increased until the end of the treatment
(at day 7). The elevated levels after one day of HD can be explained by the experimental setup,
in which the D treatment started after 7 days of H (that also resulted in DPA accumulation).
However, the strong accumulation of ABA breakdown products under D and to even higher level
in HD are in line with their suggested role in long-term stress acclimation. It was shown that PA,
the first degradation product of ABA and precursor of DPA, does also activate a subset of ABA
receptors (Weng et al., 2016). Because ABA has a very short half-live, it was suggested that the
long-lived PA could prime plants for enhanced responses to future drought (Lozano-Juste and
Cutler, 2016).
31
The phytohormone JA is another typical stress hormone known to be involved in many biotic but
also abiotic stress responses (Wasternack and Feussner, 2018). The biologically active form is
jasmonyl-L-isoleucine (JA-Ile) and we found strongly increased levels of JA-Ile under H, D and

HD conditions, (Figure 5B). 12-Hydroxyjasmonic acid (12-OH-JA) is a by-product of switching

off JA signalling with weak signalling activity (Nakamura et al., 2011). It was also described to

function as tuber-inducing factor in potato (Yoshihara et al., 1989). Under H, amounts of 12-OH-

JA as well as free JA switched from higher amounts measured at day one to lower levels at day

8, and decreased further till the end of the experiment. Also, *cis*-12-oxo-phytodienoic acid (*cis*-

OPDA), the biochemical precursor of JA, was detected at much lower levels on day 8 and 14 in

H. Cis-OPDA was also reduced at the start of the combined HD (day one) treatment, most likely

because of prior heat treatment. Altogether, this indicates a strong upregulation in the last step

of conjugation for the synthesis of JA-Ile in H, D and HD.

The accumulation of ROS is a detrimental by-product of photosynthesis and other pathways

under stress conditions. Accordingly, the detoxification of ROS by different enzymes such as

catalase or superoxide dismutase together with induction of Ca^{2+} signals is a typical response

emerging from stressed chloroplasts (Stael et al., 2015). In line with that, we measured

increased transcript levels of catalase (*CAT1*) and a methyl esterase (*MES*), which was

selected as Ca^{2+} signalling marker gene (Marc Knight, unpublished data) at the end of the

drought and heat treatment (sampling day 14). The transcript levels of *pathogenesis-related*

protein 1b1 (*PR1b*), a biotic stress as well as drought and salt stress marker (Akbudak et al.,

2020) were first lower in H but increased from day 8 to 14 in H and at day 7 in D (Figure 5C). A

similar response was seen for the chloroplast-localized lipoxygenase *13-LOX*, which is a well-

known marker gene for different stresses, especially chloroplast generated ROS (Bachmann et

al., 2002) suggesting increased ROS formation and stress level with stress duration. Strikingly,

the expression of these genes was less induced or even inhibited when H and D were

combined, which may indicate the activation of opposing signalling pathways

Heat stress, but also other stresses, induces the production of heat-shock proteins (HSPs),

which is a very conserved process in all organisms. HSPs act as molecular chaperones and

play an important role in maintain the cellular homeostasis and the proteome by supporting

protein folding, preventing misfolding or by assisting in the degradation of irreversibly damaged

polypeptides (Sato et al., 2024). At transcript level, we observed clearly elevated levels of

HSP70 transcripts after single H and D stress. Under H this was accompanied by an

accumulation of numerous HSP proteins as indicated by their significant enrichment among the

identified proteins in the proteome approach (Figure 5D, 5E). This effect was similarly

pronounced in combined HD stress (Figure 5E) with a strong enrichment of HSP70, 90, 101

involved in heat stress as well as in assisting folding (Figure 5D). The category "protein folding"

comprises mainly HSP70 and 60 group members with many of them that are present in the

chloroplast, where they are participate in the repair of photosystem (PS) II components, but also

protect proteins such as RuBisCo. In fact, our physiological data indicate a disturbance in the

electron-transport chain through PSII under heat. For example, we saw a strong decrease of

PSII efficiency under H stress (Fig. 2B). The negative effect of H stress is also reflected in lower

abundance of photosystem II proteins in the gene enrichment analysis (Figure 5D). More

specifically (Supp. Table5), there were reduced amounts of the PsbQ and PsbP subunits of the

oxygen-evolving complex.

Overall, we do see specific stress responses to heat and drought, but also to a combination of

both. This becomes also visible in Figure 5E illustrating the signature which is caused by

combined heat and drought stress in biochemical pathway view (knowledge network overlaid

with multi-omics data). The responses to HD only partly overlap with single D stress, e.g. for the

accumulation of DPA. However, for most measured metabolites the patterns are similar to heat,

but changes to HD were more pronounced (Figure 5E). Here, we cannot exclude that this

domination by heat was linked to rather the mild drought stress level. Interestingly, the

transcriptional changes of selected stress-related enzymes were weakest in HD stress

combination pointing to a redirection and rearrangement of signalling pathways compared to

individual stress factors as suggested by other studies (Zhang and Sonnewald, 2017).

Comprehensive insight into molecular processes mediating the extreme

waterlogging sensitivity of potato

Despite being documented as a highly flood-sensitive species, an in-depth characterization of

flooding-induced stress responses in potato is sparse (Jovović et al., 2021). The waterlogging

sensitivity of potato was evident in the HTP data, with several morpho-physiological traits related

to plant performance being negatively impacted following stress imposition (Figure 1C, Figure

2). This included: leaf epinasty, decreased biomass accumulation and shoot elongation,

impaired photosynthesis and stomatal conductance as well as a dramatic reduction of tuber

yield (see Figure 1,2).

Waterlogging significantly affected primary metabolic pathways as reflected in an increase in

soluble sugars and free amino acids (Figure 6A). We also observed changes in the expression

of stress-associated genes and hormones, thus highlighting potential mechanisms involved in

waterlogging acclimation. This include the increase of ABA (ABA, PA, DPA) metabolism and
response (*RDB29),* as well as the upregulation of the ethylene (ET) biosynthesis gene, *ACO2,*
and the ROS-producing enzyme, *RbohA.* Waterlogging also led to accumulation of different JA
metabolites (9,10-dHJA, cisOPDA, and JA-Ile) together with the upregulation of 13-LOX, an
enzyme involved in JA metabolism. The strong ABA signature observed in waterlogged plants
prompted us to compare waterlogging and drought responses (Figure 6B). This revealed
common stress-associated responses (e.g.: the induction of *ACO2*, *RD29, HSP70*) and a much
stronger ABA response in waterlogging relative to drought. Another notable observation was the
upregulation of the tuberigen signal, *SP6A* after 1d of waterlogging coinciding with the
downregulation of its negative regulator *CO* (Figure 6C).

Proteomics analyses of waterlogged plants revealed mild effects. Six differentially enriched
proteins were identified in response to prolonged (7 days, W07) waterlogging treatment. Among
the strongly upregulated proteins, were a LAP2-like protein (VdnPW4_5460) which encodes an
aminopeptidase, two glucan endo-1,3-beta-glucosidases (VdnPW4_8729, PBdnRY1_427), a
POP2-like protein encoding a GABA transaminase and a SEC31B-like protein, previously
described as a component of coat protein complex II (COPII), involved in vesicular transport
from the endoplasmic reticulum (ER) to (Li et al., 2021). Waterlogging led to the downregulation
of the chloroplast ribosomal protein RPL27-like, demonstrated to be important for protein
synthesis (Figure 6D).

The multi-level integrative analyses enabled visualization of the progression of stress symptoms
in waterlogged plants. In comparison to one day of waterlogging, molecular responses to
prolonged waterlogging stress (7 days, W07) displayed a distinct signature (Figure 6E). While
ABA-, JA- and ROS-biosynthesis and accumulation of free amino acids was further increased,
we observed that prolonged waterlogging led to a general inhibition of the tuberisation process
(i.e. *SP6A* downregulation). In addition, genes related to ethylene biosynthesis and response
were not regulated (*ACO2)* or downregulated (*ERF1),* thus suggesting temporal control of
ethylene signalling. Despite representing opposite ends of the water stress spectrum,
waterlogging and drought, elicited significantly overlapping responses, notably related to ABA
metabolism and proline accumulation (Figure 6A, 6B, 6E).

1   DISCUSSION

Despite its outstanding importance as a major food crop, research into the vulnerability of potato

to abiotic stresses lags that of other staple crops. In recent years, potato yields have been

significantly affected by heat, drought, and flooding, often occurring sequentially or

simultaneously (Dahal et al., 2019; Jovović et al., 2021; von Gehren et al., 2023). Considering

the increasing occurrence of these extreme weather events, this knowledge gap needs to be

urgently addressed. In this study, we leveraged the power of several omics techniques and their

integrated analysis to build a comprehensive global picture of potato responses to single and

combined heat, drought, and waterlogging stress.

Leveraging multi-omics data integration to capture the complexity of biological systems

Several tools have already been developed for integrative analysis of multi-omics data (Joshi et

al., 2024). Most broadly used are the mixOmics package (Rohart et al., 2017; Singh et al.,

2019), integrating datasets based on correlations, and the pathway visualisation tool

PaintOmics (Liu et al., 2022). In this study, however, we integrated five omics-level datasets.

Such complex datasets have rarely been analysed, even in medical research (Lee et al., 2019),

as most studies combine only two to three omics datasets (Ployet et al., 2019; Lozano-Elena et

al., 2022; Núñez-Lillo et al., 2023; Núñez-Lillo et al., 2024; Sinha et al., 2024). Since existing

tools were not directly suitable for our needs, we developed a novel pipeline harnessing the

potential of both integrative and visualization approaches. An additional step, based on machine

learning. was introduced to reduce the number of variables, in particular of phenotypic

physiological data. This reduction of variables was especially important for the correlation

analyses across omics levels, where we kept only the most informative variable. In the first step

we performed statistical modelling and correlation analyses, which provided partial overview of

events. In the second step, mechanistic insights were obtained by generating a customised

biochemical knowledge network. Our network was constructed based on knowledge extracted

from different databases, most notably the Stress Knowledge Map (Bleker et al., 2024) and

KEGG (Kanehisa et al., 2017), as well as from literature, to integrate all components that were

kept after variable selection. The obtained biochemical knowledge network enabled a

comprehensive overview of events at the pathway level and led to the identification of

mechanistic differences occurring in response to different stresses. The developed pipeline is

thus highly useful for integration and interpretation of complex datasets in future studies and can

also be applied to other species.

Integrative omics provides global insights into potato abiotic stress responses

While multi-omics approaches have been successfully applied in numerous crop species to
better understand abiotic stress responses, our study is the first to do so in potato. We
subjected the cultivar Désirée to waterlogging, drought, heat, a combination of heat and
drought, and triple stress combination encompassing all three. Across each stress treatment,
detailed morpho-physiological traits were measured, with a subset of plants sampled for the
probing of a diverse array of molecular stress markers, hormones, metabolites, and proteome
analyses across several time points.

In general, stress combinations appeared to be more detrimental to the plant performance than
individual stress applications. The combination of H, D and W led to a rapid decline in plant
viability and eventually most plants died. However, all individual stress factors caused a
reduced plant growth, had a negative impact on photosynthetic assimilate production, and both
heat and waterlogging stress impaired tuber yield (Figure 7) and tuber starch accumulation
(Supp. Figure 2). Considering all stress responses, it turned out that the cultivar Desirée was
less affected by the applied drought stress indicating that it is quite resilient to drought as
suggested previously (Demirel et al., 2020). A combination of heat and drought caused stronger
growth retardation than both stresses alone with drought responses overwriting heat
adaptations.

Nevertheless, despite the apparent mild drought phenotypes, a clear drought-associated
signature (accumulation of ABA, sugars, proline, histidine and most stress-induced transcripts)
confirmed the activation of stress responsive pathways, particularly at day 7, which contributed
to stress acclimation (Figure 5). Thus, there was a clear activation of the ABA response
pathway, as seen by an increase in the content of the hormone and its degradation products, as
well as proline, and the increased expression of ABA-responsive marker genes *SnRK*, *P5CS*,
RD29, leading to the corresponding physiological responses, e.g. a decreased water-use and
leaf temperature caused by stomata closure. Particularly interesting was the accumulation of the
ABA catabolite DPA after longer drought stress. The precursor of DPA, PA was suggested to
have an important role in priming for increased resilience to future drought stress in Arabidopsis
(Lozano-Juste and Cutler, 2016). It is believed that DPA does not trigger ABA responses, but
that has to our knowledge not been studied in potato. Hence, it might be that DPA acts as
priming signal for stress acclimation and resilience in potato.

Compared to drought, heat stress had a stronger impact on Desirée plants at all levels from

growth to photosynthesis and yield. Thermomorphogenesis is a well-described morphological

response to elevated temperature stress comprising shoot elongation and hyponastic

movement of leaves which together with an increased transpiration are seen as an acclimation

to increase ventilation and to cool the aboveground part (Quint et al. 2016). In our study we

clearly observed the hyponastic movement of leaves and stomatal opening, indicated by a

decrease in ΔT (Figure 7). This physiological response was accompanied by a decreased

amount of ABA. In contrast, the heat-mediated shoot elongation that has been seen in other

potato varieties was not visible in Desirée (Hastilestari et al., 2018; Tang et al., 2018). Instead,

the plant height of Desiree plants was reduced under elevated temperature suggesting cultivar-

specific difference that could be exploited in further studies to entangle different morphological

stress adaption mechanisms in potato. In *A. thaliana*, the thermomorphogenetic hypocotyl

elongation is tightly linked with an increase in auxin levels and is mediated by the transcription

factor Phytochrome interacting factor 4 (PIF4) (Quint et al. 2016). Consistent with the

morphological response, we did not find significantly altered levels of the phytohormone IAA in

leaves of Desiree plants in response to heat (Figure 5B).

High temperature treatment negatively affects photosynthetic capacity, particularly the efficiency

of photosystem II, which is in line with results of other studies (see (Mathur et al., 2014) for

review), and this was confirmed here at both physiological and proteomic level. In a previous

study by (Hancock et al., 2014), which also used cv. Desiree, the net CO_2 assimilation was even

higher under elevated temperatures than in control conditions. This difference might be related

to the different set-up, as in the latter study the night temperature was kept at 20°C, while here it

was adjusted to 28°C, suggesting that a low night temperature is important to maintain

photosynthetic activity. The heat-induced impact on photosynthetic capacity most likely caused

a lower production of assimilates, indicated by the decreased amount of transitory starch in

leaves. Concomitantly, contents of hexoses were found to be increased consistent with earlier

studies (Hastilestari et al. 2018). This increase may contribute to the osmo-protection of cells

and provides energy for the costly heat stress response such as the formation of heat shock

proteins (Guihur et al., 2022). A massive accumulation of heat shock proteins was found after

one week of heat stress, together with elevated levels of *HSP70* transcript levels at day 8

(Figure 5C). Although energy-demanding, the induction of HSP is important for the cellular

homeostasis and maintenance of growth and metabolism at elevated temperature. This is well

demonstrated by transgenic potato plants with increased expression of a beneficial allele of
HSC70, that exhibit improved heat stress tolerance (Trapero-Mozos et al., 2018).
3
A downregulation of photosynthesis is a typical stress response to prevent potential damage, for
example caused by ROS. This has strong implications on plant growth and yield and is
therefore regulated at various levels including light-harvesting and electron transport with high
implications for crop improvement (Kromdijk et al., 2016), particularly under stress conditions
(Grieco et al., 2020). The resulting lower photosynthetic capacity together with an increased
energy demand for stress defence reduces the amounts of assimilate that can be translocated
toward the developing tubers and its availability for storing starch. Beside assimilates, molecular
signal stimulating tuber development and growth. One important regulator is *SP6A* which was
downregulated at transcript level in our studies, similar to previous ones (Hancock et al., 2014;
Lehretz et al., 2020; Park et al., 2022; Koch et al., 2024). Stem-specific overexpression can
overcome heat-mediated yield reduction and enhances delivery of assimilates towards tubers.
15
Looking at plant hormones, we observed changes of stress hormones like SA and JA, that
traditionally have been associated with biotic stress responses. Quite striking in this context was
the increase in amount of JA-Ile under heat, drought and the combination of both (Figure 5B).
This is consistent with previous observations reporting that JA has a positive effect on
thermotolerance in Arabidopsis (Clarke et al., 2009; Balfagon et al., 2019). Heat stress
increased levels of OPDA, JA and JA-Ile and application of 5µM methyl-jasmonate improved
cell viability (Clarke et al., 2009). Using mutants this study also showed, that JA acts in concert
with SA in conferring thermotolerance. Moreover, an accumulation of JA-Ile was also observed
in Arabidopsis under drought (Yoshida and Fernie, 2024), and increased levels of JA-Ile by
overexpression of *JAR1* resulted in improved drought stress tolerance, however resulted in
stunted growth (Mahmud et al., 2022). Detailed analyses how levels of JA and its derivates as
well as biosynthesis and signalling components changes in response to stress are still missing
in potato. Therefore, but a deeper understanding of regulatory factors is required, particularly
the crosstalk with other hormones and the impact on plant growth. However, a tight modulation
of JA metabolism seems as promising target for future engineering of abiotic stress tolerance in
potato (Bittner et al., 2022).
32
Extreme sensitivity to waterlogging in potato – integrative -omics highlights
commonalities with drought

Our study provides the first detailed insight into the molecular responses underlying the high

vulnerability of potato to waterlogging. Water saturation imposes rapid oxygen deficiency in the

soil, thus impairing root respiration and function. Plant survival in flooded soils involves various

morphological and metabolic responses to either escape or cope with hypoxia, which involve

acclimation responses in roots but also in aerial organs (Sauter, 2013; Leeggangers et al.,

2023).

The data showed that plant growth and performance were more drastically affected by

waterlogging as compared to H, D and HD treatments. In addition, HTP data suggests that

waterlogging had a dominant effect even when applied after a previous combined exposure to

heat and drought (HDW) (Figure 1C and 2). When applied as a single stress, detrimental effects

on plant performance increased over time. Waterlogging dramatically impairs root conductance

and water and nutrient uptake, causing tissue dehydration and wilting. This triggers water-saving

responses such as stomatal closure and epinasty, which were reflected in increased leaf

temperatures and reduced plant compactness, respectively (Figure 1, 2E, suppl. Figure 1).

Epinastic leaf movement, a common waterlogging response in Solanaceae, is thought to reduce

photosystem damage by irradiation and transpiration (Jackson and Campbell, 1976; Geldhof et

al., 2023). Both stomatal conductance and epinasty are regulated by the pivotal flooding signal

ethylene (Leeggangers et al., 2023). While ethylene levels were not measured here, the

analyses of synthesis genes (i.e.: *ACO2)* suggested the activation of ethylene production in

waterlogged shoots. Ethylene is also known to trigger *RBOH* expression and can act

synergistically with ABA to reduce stomatal conductance (Zhao et al., 2021). ABA is also

considered to signal root stress during waterlogging (Jackson and Hall, 1987; Zhao et al., 2021).

We observed both the activation of ABA signalling and ABA accumulation and together with

increased levels of proline, this is consistent with a strong drought signature (Figure 6A, 6B, 6E).

While paradoxical, waterlogging is known to elicit shoot drought responses. As root function in

hypoxic soil ceases, it triggers a 'drought like' response in the shoot with the similar goal to

trigger water saving measures. A focus on this ABA and drought-mediated regulatory network

might thus be an attractive target for probing common resilience mechanisms to both drought

and waterlogging.

The energy shortage caused by waterlogging also leads to significant changes in sugar

metabolism. The accumulation of soluble sugars such as glucose and fructose, might be a

consequence of sink-source imbalances during waterlogging and thereby, a decline in shoot-to-

root sugar transport. Strikingly, we observed upregulation of *SP6A,* a positive regulator of

tuberisation, thus suggesting potential roles of this gene in short-term responses to
waterlogging.

Prolonged exposure to waterlogging revealed several aspects of late responses and factors
contributing to potato susceptibility to waterlogging. Leaves of waterlogged plants overcome
energy shortages by recycling carbon from amino acids and GABA. The latter, plays an
important role not only in TCA replenishment but also in ion homeostasis and reduction of
oxidative stress (Lothier et al., 2020; Wu et al., 2021). We observed a strong increase in free
amino acids that, together with the upregulation of POP2 and an aminopeptidase (Figure 6),
suggests increased protein breakdown and utilization of amino acids as alternative energy
sources. Furthermore, the downregulation of RPL27 (and other ribosomal proteins) could
indicate the shutting down of energy-demanding processes, such as protein synthesis, as a
response to this energy shortage.

Potato susceptibility to prolonged waterlogging was evidenced by other multi-level events such
as the upregulation of proteins related to protein and cell wall component turnover, *RboHA*
upregulation and photosynthesis impairment (Figure 2, Supp. Figure 3, Figure 6). It is also
explained by increased ABA signalling and biosynthesis and *RBohA* expression, which
convergently indicate increased tissue dehydration and oxidative stress that is reflected in the
HTP data (Figure1C, Figure 2). This includes decreased tuber number and weight, indicating a
retardation of both tuber initiation and bulking. As tuberisation is a particularly energetically
expensive process, the imposition of root zone hypoxia likely disrupts the underground sink
force essential for stolon development, tuber initiation and bulking.

Altogether, our data suggest that two weeks of waterlogging led to near-lethal effects, and even
if acclimation responses were activated, overall, they could not compensate for maintaining root
function (i.e. unrecovered water consumption, Supp. Table 4) and general plant survival, even
during recovery, thus confirming the high susceptibility of potato to waterlogging.

Conclusion
The present comprehensive approach produced a rich integrated dataset, which enabled
diverse exploration of molecular mechanisms across various levels and processes. Through the
connection of phenotype to molecular responses, we attained deeper insights into the intricate
regulation of metabolic and phenotypic traits. This should now guide the identification of key

regulators that govern the interplay between molecular dynamics and their phenotypic
expressions. The utilization of both knowledge-based approaches and multivariate statistical
methods played a crucial role in deciphering complex molecular regulatory networks and their
association with phenotypic and physiological traits, thereby facilitating the rapid generation of
hypotheses.

In addition to several novel insights into potato stress responses, this study also provides a
blueprint for performing and analysing single and multiple stress and effective integration of
large datasets for potato. Importantly, this setup could be applied also for other plant species.
These advancements hold significant implications for potato breeding strategies, providing a
deeper understanding of plant stress responses and expediting trait selection. As agricultural
landscapes confront challenges like climate change and population growth, embracing multi-
omics integration holds promise for cultivating resilient potato varieties that can thrive in various
conditions.

METHODS

Plant growth conditions and sampling

150 in-vitro potato cuttings (*Solanum tuberosum* cv Desirée) were cultivated and grown as
described in the supplementary methods. After 32 days of cultivation, plants were randomly
distributed into 6 groups (6 plants each) referring to control group and 5 different stress
conditions (heat, drought, combined heat and drought, waterlogging and combination of heat,
drought and waterlogging) (Figure 1). Plants were moved into two growth units of Growth
Capsule (PSI; (Photon Systems Instruments), Czech Republic) where climate conditions for
day/night temperature were set in one unit to 22/19°C, referring to control conditions, and in the
second unit to 30/28 °C, referring to heat conditions. In both units growing light intensity was set
at 330 µmol m^{-2} s^{-1} PPFD and relative humidity was maintained at 55%. All plants were
measured under control in day 0 then the stress treatments depicted schematically in Figure 1.
The treatments were applied as the following: (1) Control conditions – cultivation at 22/19 °C,
watering up to 60% FC; (2) Drought conditions – cultivation at 22/19 °C, watering up to 60% FC
until day 7, then reduce watering to 30% FC for 1 week (until day 14); (3) Heat conditions –
cultivation at 30/28 °C for 2 weeks, watering up to 60% FC until day 14; (4) Heat + Drought
conditions - cultivation at 30/28 °C for 2 weeks, watering up to 60% FC for 1 week (until day 7),
then reduce watering to 30% FC for 1 week (until day 14); (5) Waterlogging conditions –

cultivation at 22/19 °C, watering up to 130% FC for 2 weeks (until day 14); (6) Heat + Drought +

Waterlogging conditions – cultivation at 30/28 °C for 2 weeks with watering up to 60% FC for 1

week (until day 7), then reduce watering to 30% FC until day 14 followed by inducing

waterlogging by cultivation at 22/19 °C for 1 week with watering up to 130% FC until day 21.

Except for Heat + Drought + Waterlogging conditions, all stress treatments were followed by

one week of recovery (from day 15 and until day 21) in control conditions.

Plants were divided into two sets, "phenotyping plants" and "plants for tissue harvest" (see

Supp. Table 1). Phenotyping set consisted of 6 replicates per treatment, in total 36 plants, and

was used for daily image-based phenotyping (for definition of scored traits see Supp. Table 2).

High-throughput phenotyping

Prior to the stress treatment initiation and during the stress treatments, all plants were daily

phenotyped. A comprehensive phenotypical protocol was used for the acquisition of

physiological and morphological traits according to the described method (Abdelhakim et al.,

2024). All imaging sensors for digital analysis are being implemented in the PlantScreen^{TM}

Modular system (PSI, Czech Republic). The photosynthesis☐related traits were determined

using kinetic chlorophyll fluorescence imaging where the selected protocol for measuring plants

was similar to the defined approach (Abdelhakim et al., 2024). The measurement of

temperature profiles of the plants was measured using thermal imaging, where the acquisition

and segmentation of the images were processed as described in (Abdelhakim et al., 2021;

Findurová et al., 2023). The morphological and growth dynamics were determined using both

top and multiple angles (0°, 120°, and 240°) side view RGB imaging, and images were

processed as described by (Awlia et al., 2016). Each pot was loaded onto a transport disk

automatically moving on a conveyor belt between the automatic laser height measuring unit,

acclimation unit, robotic-assisted imaging units, weighing and watering unit, and the cultivation

greenhouse located area. The raw images were automatically processed and parameters were

extracted through PlantScreen^{TM} Analyzer software (PSI, Czech Republic) (Supp. Table 2).

Statistical evaluation was performed to check the differences between the treatments using

Wilcoxon test (Supp. Table 3).

Tissue sampling

Leaf sampling was conducted on days 1, 7, 8, 14, 15, and 21 after stress treatment initiation

(Treatment days). The 2^{nd} and 3^{rd} fully-developed leaves were harvested and flash-frozen in

liquid nitrogen. Subsequently, leaf tissue was homogenised, aliquoted and distributed for

individual follow-up transcriptomics, metabolomics, hormonomics, and proteomics analyses.

Remaining above ground tissue was harvested and total fresh weight (FW) and dry weight (DW)

was defined (Supp. Table 4). In total 112 plants with 4 replicates per sampling time and per

treatment were collected. The 4^{th} leaf was harvested as well to calculate relative water content

(RWC) (Supp. Table4), and three leaf disks were collected and weighed, then soaked in water

to determine the turgor weight (TW) and dried in the oven to calculate RWC (Supp. Table 2). In

addition, at the end of the experiment, the below ground tissue was collected, where number of

tubers per plant and total weight were assessed from four replicates per treatment. Harvest

index was calculated as a ratio between tuber weight and the total biomass.

Multi-omics analysis

Transcriptomic marker analysis

RT-qPCR was performed to assess the expression of 14 marker genes involved in redox

homeostasis) hormonal signalling - ethylene, abscisic acid, cytokinin, salicylic acid, jasmonic

acid, heat stress, tuber development, circadian clock, and calcium signalling using previously

validated reference genes (Supp. Table 7).

RNA was extracted and DNase treated using Direct-zol RNA Miniprep Kit (Zymo Research,

USA) from 80-100 mg of frozen homogenised leaf tissue, followed by reverse transcription using

High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher, USA). The expression of the

target and reference genes was analysed by qPCR, as described previously (Petek et al., 2014;

Abdelhakim et al., 2021). QuantGenius ([http://quantgenius.nib.si)](http://quantgenius.nib.si), was used for quality control,

standard curve-based relative gene expression quantification and imputation of values below

level of detection or quantification (LOD, LOQ) (Baebler et al., 2017).

Hormonomics

Concentration of the endogenous abscisate metabolites, auxin metabolites, jasmonates and

salicylic acid were determined in 10 mg of frozen homogenised leaf tissue according to the

method described (Flokova et al., 2014) and modified by Široká et al. (Siroka et al., 2022) (see

Supplementary methods for details). All experiments were repeated as four biological replicates.

Metabolomics

For determination of soluble sugar, starch and amino acid contents, 30 - 50 mg of freeze-dried

leaf or tuber material were extracted with 1 ml of 80% (v/v) ethanol. Soluble sugar and starch

content was determined as described in Hastilestari et al. (2018), while amino acids sample

preparation and measurements were performed as described elsewhere (Obata et a. 2020).

(Smith and Zeeman, 2006) (Obata et al., 2020).

Proteomics

High-throughput shotgun proteomics was done according to (Hoehenwarter et al., 2008) with

following modifications: 40 mg of leaf tissue from multiple stress conditions were freeze-dried in

liquid N_2 and ground using mortar and pestle. The proteins were extracted, pre-fractionated

(40µg of total protein were loaded onto the gel (1D SDS-PAGE), trypsin digested and desalted

(using a C18 spec plate) according to a previously described method (Chaturvedi et al., 2013;

Ghatak et al., 2016). One µg of purified peptides was loaded onto a C18 reverse-phase

analytical column (Thermo Scientific, EASY-Spray 50 cm, 2 µm particle size). Separation was

achieved using a two-and-a-half-hour gradient method, starting with a 4–35% buffer B (v/v)

gradient [79.9% ACN, 0.1% formic acid (FA), 20% ultra-high purity water (MilliQ) over 90

minutes. Buffer A (v/v) consisted of 0.1% FA in high-purity water (MilliQ). The flow rate was set

to 300 nL/min. Mass spectra were acquired in positive ion mode using a top-20 data-dependent

acquisition (DDA) method. A full MS scan was performed at 70,000 resolution (m/z 200) with a

scan range of 380–1800 m/z, followed by an MS/MS scan at 17,500 resolution (m/z 200). For

MS/MS fragmentation, higher energy collisional dissociation (HCD) was used with a normalized

collision energy (NCE) of 27%. Dynamic exclusion was set to 20 seconds.

Raw data were searched with the SEQUEST algorithm present in Proteome Discoverer version

1.3 (Thermo Scientific, Germany) described previously (Chaturvedi et al., 2015; Ghatak et al.,

2020). Pan-transcriptome (Petek et al., 2020) protein fasta was employed. The identified

proteins were quantitated based on total ion count and normalised using the normalised spectral

abundance factor (NSAF) strategy (Paoletti et al., 2006).

Data analysis

The programming environments R v.4.3 and v4.4 (https://www.r-project.org/) and Python v3.8

(www.python.org) were used. Experimentally acquired data and data required to reproduce the

analysis are available from Supplemental Table 4 and NIB's GitHub repository

(https://github.com/NIB-SI/multiOmics-integration) were used.

Data preprocessing

A master sample description metadata file was constructed (Supp. Table 1). Potential

inconsistencies between replicates were examined using pairwise plots between omics levels,

multidimensional scaling plots and scatterplot matrices within omics' levels using the vegan

v2.6.-4 R package (Oksanen et al., 2022). Missing values were handled as described in the

Supplementary methods. Due to many missing values, the neoPA (hormonomics) variable was

excluded from further analysis.

Variable selection was conducted on the non-invasive phenomics variable sets (Supp. Table 4).

The random forest (RF) algorithm from the R package caret v6.0-94 (Kuhn, 2008) as well as the

python package scikit-learn v1.2.0 were used with default settings, as RF showed the best

performance out of a selection of algorithms. Recursive feature elimination was applied in R and

multiple importance scores, including mutual information, Anova, RF importance and SHAP

values (Lundberg and Lee, 2017) were computed in Python, showing consistencies between the

approaches for the top 5 variables (Top area, compactness, qL_Lss, ΔT, water consumption;

nonredundancy ranking in R). The sixth variable (Fv/Fm_Lss) was selected based on expert

knowledge.

Gene set enrichment was performed on the proteomics dataset using GSEA v4.3.2

(Subramanian et al., 2005) and in-house generated gene sets (Supp. File 3, Supp. Table 6) and

visualised using biokit v0.1.1. Proteomics differential expression was conducted using the DEP

v 1.22.0 package (Zhang et al., 2018) (Supp. Table 5). For downstream proteomics analyses,

differentially abundant and enriched proteins (from pathways important for this experimental

setup) were used. Waterlogging stress was cut-off at one-week duration, while triple stress

(HDW) was not considered in downstream analyses due to poor plant performance.

Analysis of individual omics data layers

Pearson correlation coefficient (PCC) heatmaps (pheatmap v1.0.12, heatmaply v. 1.5.0) were

generated within each treatment and for explicit treatment duration. Permutation-based t-test

(MKinfer v1.2) was used to denote differences between specific treatment and control within the

corresponding time-point (Kohl, 2024). Corresponding log2FC were calculated. For downstream

analyses, 4 out of 6 replicates were chosen from non-invasive phenomics measurements to

allow integration with invasive phenomics and other omics measurements conducted on 4

replicates.

Integration across different omics datasets

Correlations between components measured in various Omics' levels were calculated and

visualised using DIABLO (Singh et al., 2019) as implemented in the mixOmics v6.24.0 package

(Rohart et al., 2017). The correlation matrix was calculated separately for each stress as well as

for control.

Integration of data with prior knowledge

A background knowledge network was manually constructed considering biochemical pathways

between measured variables. Where necessary, pathways were simplified to only include

representative variables, to prevent addition of many unmeasured nodes that would impede the

visualisation. Proteomics differential expression results were merged with t-test and log2FC

results (Supp. Table 3). Final networks were visualised using DiNAR (Zagorscak et al., 2018)

and Cytoscape (Shannon et al., 2003).

For additional reports and some results not used in this manuscript see supplementary methods

and a project's GitHub repository https://github.com/NIB-SI/multiOmics-integration.

24   Acknowledgements

The authors wish to thank Marijke Woudsma and Doretta Boomsma from HZPC Research for

providing the Desiree plantlets for this study and Mirella Sorrentino for help with conducting the

experiments at Photon Systems Instruments (PSI) Research Center (Drásov, Czech Republic).

Moreover, we the thank following colleagues for their assistance and expertise: S. Reid (FAU)

and D. Pscheidt (FAU) for measuring metabolite content, and Katja Stare (NIB) and Nastja
Marondini (NIB) for measuring gene expression.

Author contributions

KP, KG, SS, RS, CBa and MT conceptualized the study and designed the experiments, KP
measured phenotypic traits, ŠB measured gene expression, ON, AP and JŠ measured hormone
content, CS measured metabolite content, PC, AG prepared samples and performed
proteomics analysis, LAS performed LC-MS measurements for proteomics analysis; MZ curated
data; MZ, CB, AB, AZ and KG defined formal analysis methodology; MZ, CB, JZ, and AB
conducted statistical, mathematical and computational analysis; MZ, LA, NR, CB, AB, ŠB and
KG performed data visualisation; MZ, LA, NR, CB, SS, RS, KG, CBa and MT wrote the original
draft. All authors edited the manuscript and approved the final manuscript version. Detailed
author contributions are available from the Supp. Table 8.

Figure legends

Figure 1. Overview of the experimental design for single- and combined stress treatments and
multi-omics sampling. A) Summary of cultivation conditions. Timeline of the experimental set-up and
applied stress treatments, including the recovery phase in potato cv. Desirée. Timing and duration of
stress treatment and days for tissue sampling are shown. (B) Actions comprised cultivation in the growing
chambers and daily phenotyping with a set of sensors using the PlantScreen^{TM} phenotyping platform at
PSI Research Center. (C) Automated image analysis pipeline was used to extract quantitative traits for
morphological, physiological, and biochemical performance characterization of the plants during the
stress treatment and recovery phase. Side view colour segmented RGB images of plants at selected time
points of tissue sampling (left panel) and daily plant volume (m^3) calculated from top and multiple angle
side view RGB images (right panel). N=6.

Figure 2. Physiological profiling using high-throughput phenotyping platforms reveals distinct
responses to single and combined stresses. A) Pixel-by-pixel false colour images of operating
efficiency of photosystem II in light steady state (QY_Lss) captured by kinetic chlorophyll fluorescence
measurement. Images for selected time points of tissue sampling are shown. Colour coding of the
treatments apply for the entire figure. Vertical dashed lines indicate the onset and end of drought. B)
QY_Lss values extracted from images for each individual time point. C) Steady-state fluorescence of
maximum efficiency of PSII photochemistry in the light trait based on chlorophyll fluorescence top view
(Fv/Fm_Lss). D) steady-state estimation of the fraction of open reaction centres in PSII trait in light based
on chlorophyll fluorescence top view (qL_Lss). E) Difference between canopy average temperature
extracted from thermal IR images and air temperature measured in the thermal IR imaging unit (ΔT). F)
Water use efficiency (WUE) based on plant volume and water consumption. A-F) Black dotted lines

reflect the initiation and removal of drought stress, respectively. See Supplementary Table 3 for Statistical

evaluation of differences between groups using Wilcoxon test. G) Tuber numbers counted per plant on

the last day of the experiment (Day 28 = 60 days of cultivation). H) Harvest index calculated from the total

biomass and tuber weight on the last day of the experiment. G-H) Measurements, mean and standard

deviation are shown (n = 4). Statistical evaluation of differences between groups is given by the non-

parametric Kruskal–Wallis test (*one-way ANOVA on ranks*); p-value above x-axis, where asterisk denotes

p-value < 0.05. See Figure 1A for scheme on stress treatments.

Figure 3: Integrated analysis of measured and generated data permits global visualization and

multi-level amalgamation of potato stress responses. A) Schematics of tissue sampling protocol. 2nd

and 3rd leaves were harvested for destructive "omics" analysis, 4th leaf was used for relative water content

calculation. Remaining plant tissue was quantified to obtain total above-ground biomass and tuber yield.

B) Overview of data analysis pipeline. C) Dataset overview: multidimensional scaling shows combined

HDW stressed plants as extremes, the centroid of each plant group is shown. D) Most informative

variables from the phenomics level. Pearson correlation coefficients between them are presented as

hierarchically clustered heat map in waterlogging and heat stress.

Figure 4: Integration of multi-omics data in a knowledge-based metabolic and signalling network.

A) Structure of knowledge network. Individual studied components are coloured according to their

function in different pathways. B) To compare the effects of different stresses on the overall state of the

plant, we overlaid the knowledge networks with measured changes in component concentration. Nodes

are coloured by log2 fold changes (red – increase in stress compared to control, blue – decrease in stress

compared to control, grey – measurement not available) shown for two time points: sampling day 8 and

sampling day 14 for the different stress treatments, days of stress treatment are given with each network

(for more details of the set up see Figure 1A).

Figure 5: Combined heat and drought stress trigger distinct responses compared to each

individual response. Additive effect of combined stress is most pronounced for branched chain amino

acids accumulation and JA signalling response. A-C) Heatmaps showing log2FC (FDR p-value < 0.05) in

individual stress heat (H) or drought (D) stress in comparison to combined one (HD) for targeted

molecular analyses. Label colours indicate pathway associated with each molecule as in the Knowledge

network (see Figure 4 for legend). A) Changes in metabolite levels. B) Changes in hormone levels, and

C) Changes in selected stress-related transcripts. D) Changes observed on proteomics level. Results of

Gene Set Enrichment Analysis (FDR q-value < 0.1) are shown. For more information see Supp. Table 6.

E) Biochemical knowledge network showing changes under combined HD stress at day 14 (treatment day

7). In this version of knowledge network, only nodes that were significantly differentially expressed (vs.

control conditions) are coloured and the connections between two differentially expressed nodes are

coloured black. Node full black border indicates molecules with higher expression levels in HD compared

to H and/or D alone. Dashed black border indicates molecules with lower expression levels in HD
compared to H and/or D alone (difference of log2FC > 0.5)**.**
3
Figure 6: Waterlogging triggers drought-stress like molecular responses in potato. A-D) Heatmaps
showing log2FC (FDR p-value < 0.05) changes in A) metabolite levels, B) phytohormones, C) selected
stress-related transcripts. D) Volcano plot of differential proteomics analysis at day 7. Proteins with FDR
p-value < 0.05 shown as blue (downregulated) and red (upregulated) dots. For more information see
Supp. Table 5. E) Knowledge network of waterlogging stress at day 1 and day 7 (unfiltered, colour range
[-2, 2]). For legend see Figure 4.
10
Figure 7. Schematic summary of multilevel responses to single and combined heat, drought and
waterlogging stresses. Selected variables from each level are shown. Summary of molecular responses
(hormones, metabolites and transcripts) was based on the comparisons illustrated in Figures 5 and 6.
Summary of morphophysiological responses were based on the data from the last day of the experiment
(Day 28), which includes tuber information. Proteomics data set is not included here due to the small
dataset of differentially expressed proteins in the waterlogging treatment. Degree of increase or decrease
is not specified.
18

Supplementary data

Supplementary File 1: Principal coordinates analysis (PCoA) of Bray-Curtis dissimilarity between
samples. Interactive 3D plot shows mapping to three dimensions. Spheres are coloured by condition.
The numerical values used to construct this figure can be found in Supplementary Table 4. Variables
were min-max scaled.
Supplementary File 2: Differential Network Analysis animation. Export from DiNAR app (Zagoršćak
et al., 2018). Node colours correspond to upregulation (red) and downregulation (blue) compared to
control (p-value adjusted < 0.05). The size of nodes corresponds to absolute log2FC values. Order of
visualisation: 0: background network, 1: 1 day of drought, 2: one week of drought 3: one day of heat, 4:
one week of heat, 5: 8 days of heat, 6: two weeks of heat, 7: 8 days of heat + one day of drought, 8: two
weeks of heat + one week of drought, 9: one day of waterlogging, 10: one week of waterlogging.
Supplementary File 3: Pan-transcriptome Gene Matrix Transposed (GMT) file format. Tab delimited
file utilised for gene set enrichment analysis that describes gene sets from pan-transcriptome of three
potato genotypes: Desiree, PW363 and Rywal (Petek et al., 2020).
Supplementary Figure 1: Plant morphological responses of control and stress-treated plants. (A)
Top view RGB images at selected time points of tissue sampling starting from day 1 where i) in the first
week heat stress was induced for H, HD, HDW marked rows, then ii) in the second week drought was
induced for D, HD, HDW up to day 14, and finally iii) in the third week waterlogging was induced for HDW

up to day 21, while the other treatments were recovered. (B-E) Parameters based on analysed RGB

images; Top area, plant height, compactness, and relative growth rate (RGR). The data represent mean

values ± standard error of mean (S.E.M) (n = 6).

Supplementary Figure 2: Log2FC of a tuber metabolite's relative abundance between stress and

control. Numbers in bold correspond to comparisons with p-value < 0.05. Additionally, log2FC for

number of tubers and total tuber weight is shown.

Supplementary Figure 3: Correlation analysis within and between omics levels. For correlation

analysis within omics levels heatmaps display Pearson correlation coefficient (PCC). For analysis

between components of different molecular levels heatmaps display canonical correlation analysis (CCA)

results. Variable prioritisation was conducted using multiblock sPLS-DA (Singh et al., 2019).

Supplementary Table 1: Phenodata. Experimental design and days of tissue sampling.

Supplementary Table 2: Phenomics featuredata. Phenotyping trait description from multiple imaging

sensors and traits from the final destructive harvest.

Supplementary Table 3: Statistical results across different omics levels. Combined table contains

statistical information and various annotations for molecules shown in biochemical knowledge network

(see Figure 2,4,5,6 and Supplementary File 2).

Supplementary Table 4: Omics measurements. Multi-sheet file containing tuber (metabolomics,

harvest information) and leaf data (phenomics, transcriptomics, hormonomics and metabolomics).

Phenomics data is separated to non-invasive and invasive trait measurements. For trait description see

Supplementary Table 2.

Supplementary Table 5: Differential protein expression. Results table of differential protein expression

from R package DEP (Zhang et al., 2018) complemented by MapMan annotations (Ramšak et al., 2014)

and Arabidopsis thaliana orthologue identifiers, short names, and gene descriptions (Bleker at al., 2024,

Zagorščak et al., 2018).

Supplementary Table 6: Gene set enrichment analysis of proteomics data. MapMan v3 pathway

enrichment results containing classes of proteins that are over-represented in stress treatments

compared to control (FDR cut-off 0.05 and 0.1). MapMan pathways are defined in supplementary file 3.

Supplementary Table 7: Genes used for transcriptome analysis using quantitative PCR. Gene

functional group, name (abbreviation) and description, as well as corresponding primer and probe

sequences, source and amplification efficiency are shown. Suitability of reference genes was validated

using RefFinder (Xie et al., 2023).

Supplementary Table 8: CRediT (Contributor Roles Taxonomy) authorship statement.

Competing interests

The authors declare no competing interests.

Funding

This work was funded by the EU H2020-SFS-2019-2 RIA project ADAPT, GA 2020 862-858 and the Slovenian Research Agency (ARIS) under grant agreements P4-0165, J2-3060, and Z4-50146. Moreover, this work was partially supported by the Ministry of Education, Youth and Sports of the Czech Republic with the European Regional Development Fund-Project "SINGING PLANT" (no. CZ.02.1.01/0.0/0.0/16_026/0008446).

Data availability statement

Experimentally acquired data and data required to reproduce the analysis are available from Supp. Table 4 and NIBs' GitHub repository https://github.com/NIB-SI/multiOmics-integration.

The MS/MS spectra of the identified proteins and their meta-information from both databases have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (https://www.ebi.ac.uk/pride) with the dataset identifier PXD052587.

References

Abdelhakim LOA, Palma CFF, Zhou R, Wollenweber B, Ottosen CO, Rosenqvist E (2021) The effect of individual and combined drought and heat stress under elevated CO(2) on physiological responses in spring wheat genotypes. Plant Physiol Biochem **162:** 301-314

Abdelhakim LOA, Pleskacova B, Rodriguez-Granados NY, Sasidharan R, Perez-Borroto LS, Sonnewald S, Gruden K, Vothknecht UC, Teige M, Panzarova K (2024) High Throughput Image-Based Phenotyping for Determining Morphological and Physiological Responses to Single and Combined Stresses in Potato. J Vis Exp

Abdelhakim LOA, Rosenqvist E, Wollenweber B, Spyroglou I, Ottosen C-O, Panzarová K (2021) Investigating Combined Drought- and Heat Stress Effects in Wheat under Controlled Conditions by Dynamic Image-Based Phenotyping. Agronomy **11:** 364

Abelenda JA, Bergonzi S, Oortwijn M, Sonnewald S, Du M, Visser RGF, Sonnewald U, Bachem CWB (2019) Source-Sink Regulation Is Mediated by Interaction of an FT Homolog with a SWEET Protein in Potato. Curr Biol **29:** 1178-1186 e1176

Abelenda JA, Cruz-Oro E, Franco-Zorrilla JM, Prat S (2016) Potato StCONSTANS-like1 Suppresses Storage Organ Formation by Directly Activating the FT-like StSP5G Repressor. Curr Biol **26:** 872-881

Akbudak MA, Yildiz S, Filiz E (2020) Pathogenesis related protein-1 (PR-1) genes in tomato (Solanum lycopersicum L.): Bioinformatics analyses and expression profiles in response to drought stress. Genomics **112:** 4089-4099

Awlia M, Nigro A, Fajkus J, Schmoeckel SM, Negrao S, Santelia D, Trtilek M, Tester M, Julkowska MM, Panzarova K (2016) High-Throughput Non-destructive Phenotyping of Traits that Contribute to Salinity Tolerance in Arabidopsis thaliana. Front Plant Sci **7:** 1414

Bachmann A, Hause B, Maucher H, Garbe E, Voros K, Weichert H, Wasternack C, Feussner I (2002) Jasmonate-induced lipid peroxidation in barley leaves initiated by distinct 13-LOX forms of chloroplasts. Biol Chem **383:** 1645-1657

Baebler S, Svalina M, Petek M, Stare K, Rotter A, Pompe-Novak M, Gruden K (2017) quantGenius: implementation of a decision support system for qPCR-based gene quantification. BMC Bioinformatics **18:** 276

Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GED, Schroeder JI (2019) Genetic strategies for improving crop yields. Nature **575:** 109-118

Balfagon D, Sengupta S, Gomez-Cadenas A, Fritschi FB, Azad RK, Mittler R, Zandalinas SI (2019) Jasmonic Acid Is Required for Plant Acclimation to a Combination of High Light and Heat Stress. Plant Physiol **181:** 1668-1682

Benitez-Alfonso Y, Soanes BK, Zimba S, Sinanaj B, German L, Sharma V, Bohra A, Kolesnikova A, Dunn JA, Martin AC, Khashi URM, Saati-Santamaria Z, Garcia-Fraile P, Ferreira EA, Frazao LA, Cowling WA, Siddique KHM, Pandey MK, Farooq M, Varshney RK, Chapman MA, Boesch C, Daszkowska-Golec A, Foyer CH (2023) Enhancing climate change resilience in agricultural crops. Curr Biol **33:** R1246-R1261

Bittner A, Ciesla A, Gruden K, Lukan T, Mahmud S, Teige M, Vothknecht UC, Wurzinger B (2022) Organelles and phytohormones: a network of interactions in plant stress responses. J Exp Bot **73:** 7165-7181

Bleker C, Ramsak Z, Bittner A, Podpecan V, Zagorscak M, Wurzinger B, Baebler S, Petek M, Kriznik M, van Dieren A, Gruber J, Afjehi-Sadat L, Weckwerth W, Zupanic A, Teige M, Vothknecht UC, Gruden K (2024) Stress Knowledge Map: A knowledge graph resource for systems biology analysis of plant stress responses. Plant Commun**:** 100920

Cembrowska-Lech D, Krzeminska A, Miller T, Nowakowska A, Adamski C, Radaczynska M, Mikiciuk G, Mikiciuk M (2023) An Integrated Multi-Omics and Artificial Intelligence Framework for Advance Plant Phenotyping in Horticulture. Biology (Basel) **12**

Chaturvedi P, Doerfler H, Jegadeesan S, Ghatak A, Pressman E, Castillejo MA, Wienkoop S, Egelhofer V, Firon N, Weckwerth W (2015) Heat-Treatment-Responsive Proteins in Different Developmental Stages of Tomato Pollen Detected by Targeted Mass Accuracy Precursor Alignment (tMAPA). J Proteome Res **14:** 4463-4471

Chaturvedi P, Ischebeck T, Egelhofer V, Lichtscheidl I, Weckwerth W (2013) Cell-specific analysis of the tomato pollen proteome from pollen mother cell to mature pollen provides evidence for developmental priming. J Proteome Res **12:** 4892-4903

Clarke SM, Cristescu SM, Miersch O, Harren FJM, Wasternack C, Mur LAJ (2009) Jasmonates act with salicylic acid to confer basal thermotolerance in Arabidopsis thaliana. New Phytol **182:** 175-187

Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR (2010) Abscisic acid: emergence of a core signaling network. Annu Rev Plant Biol **61:** 651-679

Dahal K, Li XQ, Tai H, Creelman A, Bizimungu B (2019) Improving Potato Stress Tolerance and Tuber Yield Under a Climate Change Scenario - A Current Overview. Front Plant Sci **10:** 563

Demirel U, Morris WL, Ducreux LJM, Yavuz C, Asim A, Tindas I, Campbell R, Morris JA, Verrall SR, Hedley PE, Gokce ZNO, Caliskan S, Aksoy E, Caliskan ME, Taylor MA, Hancock RD (2020) Physiological, Biochemical, and Transcriptional Responses to Single and Combined Abiotic Stress in Stress-Tolerant and Stress-Sensitive Potato Genotypes. Front Plant Sci **11:** 169

FAO (2023) The Impact of Disasters on Agriculture and Food Security 2023 – Avoiding and reducing losses through investment in resilience. . *In.* FAO, Rome, Italy, p #168 p.

Findurová H, Veselá B, Panzarová K, Pytela J, Trtílek M, Klem K (2023) Phenotyping
drought tolerance and yield performance of barley using a combination of imaging
methods. Environmental and Experimental Botany **209:** 105314
Flokova K, Tarkowska D, Miersch O, Strnad M, Wasternack C, Novak O (2014) UHPLC-
MS/MS based target profiling of stress-induced phytohormones. Phytochemistry **105:**
147-157
Geldhof B, Pattyn J, Van de Poel B (2023) From a different angle: genetic diversity underlies
differentiation of waterlogging-induced epinasty in tomato. Front Plant Sci **14:** 1178778
Ghatak A, Chaturvedi P, Bachmann G, Valledor L, Ramsak Z, Bazargani MM, Bajaj P,
Jegadeesan S, Li W, Sun X, Gruden K, Varshney RK, Weckwerth W (2020)
Physiological and Proteomic Signatures Reveal Mechanisms of Superior Drought
Resilience in Pearl Millet Compared to Wheat. Front Plant Sci **11:** 600278
Ghatak A, Chaturvedi P, Nagler M, Roustan V, Lyon D, Bachmann G, Postl W, Schrofl A,
Desai N, Varshney RK, Weckwerth W (2016) Comprehensive tissue-specific proteome
analysis of drought stress responses in Pennisetum glaucum (L.) R. Br. (Pearl millet). J
Proteomics **143:** 122-135
Grieco M, Roustan V, Dermendjiev G, Rantala S, Jain A, Leonardelli M, Neumann K,
Berger V, Engelmeier D, Bachmann G, Ebersberger I, Aro EM, Weckwerth W, Teige
M (2020) Adjustment of photosynthetic activity to drought and fluctuating light in wheat.
Plant Cell Environ **43:** 1484-1500
Guihur A, Rebeaud ME, Goloubinoff P (2022) How do plants feel the heat and survive?
Trends Biochem Sci **47:** 824-838
Hall RD, D'Auria JC, Silva Ferreira AC, Gibon Y, Kruszka D, Mishra P, van de Zedde R
(2022) High-throughput plant phenotyping: a role for metabolomics? Trends Plant Sci
27: 549-563
Hancock RD, Morris WL, Ducreux LJ, Morris JA, Usman M, Verrall SR, Fuller J, Simpson
CG, Zhang R, Hedley PE, Taylor MA (2014) Physiological, biochemical and molecular
responses of the potato (Solanum tuberosum L.) plant to moderately elevated
temperature. Plant Cell Environ **37:** 439-450
Hastilestari BR, Lorenz J, Reid S, Hofmann J, Pscheidt D, Sonnewald U, Sonnewald S
(2018) Deciphering source and sink responses of potato plants (Solanum tuberosum L.)
to elevated temperatures. Plant Cell Environ **41:** 2600-2616
Hoehenwarter W, van Dongen JT, Wienkoop S, Steinfath M, Hummel J, Erban A, Sulpice
R, Regierer B, Kopka J, Geigenberger P, Weckwerth W (2008) A rapid approach for
phenotype-screening and database independent detection of cSNP/protein
polymorphism using mass accuracy precursor alignment. Proteomics **8:** 4214-4225
Jackson MB, Campbell DJ (1976) Waterlogging and petiole epinasty in tomato: The role of
ethylene and low oxygen. New Phytologist **76:** 21-29
Jackson MB, Hall KC (1987) Early stomatal closure in waterlogged pea plants is mediated by
abscisic acid in the absence of foliar water deficits. Plant, Cell & Environment **10:** 121-
130
Jamil IN, Remali J, Azizan KA, Nor Muhammad NA, Arita M, Goh HH, Aizat WM (2020)
Systematic Multi-Omics Integration (MOI) Approach in Plant Systems Biology. Front
Plant Sci **11:** 944
Joshi S, Patil S, Shaikh A, Jamla M, Kumar V (2024) Modern omics toolbox for producing
combined and multifactorial abiotic stress tolerant plants. Plant Stress **11:** 100301
Jovović Z, Broćić A, Velimirović ŽD, A. Komnenić (2021) The influence of flooding on the
main parameters of potato productivity. *In*, Ed 1320. International Society for
Horticultural Science (ISHS), Leuven, Belgium, pp 133-138
Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K (2017) KEGG: new perspectives
on genomes, pathways, diseases and drugs. Nucleic Acids Res **45:** D353-D361

Koch L, Lehretz GG, Sonnewald U, Sonnewald S (2024) Yield reduction caused by elevated temperatures and high nitrogen fertilization is mitigated by SP6A overexpression in potato (Solanum tuberosum L.). Plant J **117:** 1702-1715

Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP (2016) Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. Science **354:** 857-861

Lal MK, Tiwari RK, Kumar A, Dey A, Kumar R, Kumar D, Jaiswal A, Changan SS, Raigond P, Dutt S, Luthra SK, Mandal S, Singh MP, Paul V, Singh B (2022) Mechanistic Concept of Physiological, Biochemical, and Molecular Responses of the Potato Crop to Heat and Drought Stress. Plants (Basel) **11**

Lee AH, Shannon CP, Amenyogbe N, Bennike TB, Diray-Arce J, Idoko OT, Gill EE, Ben-Othman R, Pomat WS, van Haren SD, Cao KL, Cox M, Darboe A, Falsafi R, Ferrari D, Harbeson DJ, He D, Bing C, Hinshaw SJ, Ndure J, Njie-Jobe J, Pettengill MA, Richmond PC, Ford R, Saleu G, Masiria G, Matlam JP, Kirarock W, Roberts E, Malek M, Sanchez-Schmitz G, Singh A, Angelidou A, Smolen KK, Consortium E, Brinkman RR, Ozonoff A, Hancock REW, van den Biggelaar AHJ, Steen H, Tebbutt SJ, Kampmann B, Levy O, Kollmann TR (2019) Dynamic molecular changes during the first week of human life follow a robust developmental trajectory. Nat Commun **10:** 1092

Leeggangers HACF, Rodriguez-Granados NY, Macias-Honti MG, Sasidharan R (2023) A helping hand when drowning: The versatile role of ethylene in root flooding resilience. Environmental and Experimental Botany **213:** 105422

Lehretz GG, Sonnewald S, Lugassi N, Granot D, Sonnewald U (2020) Future-Proofing Potato for Drought and Heat Tolerance by Overexpression of Hexokinase and SP6A. Front Plant Sci **11:** 614534

Li B, Zeng Y, Cao W, Zhang W, Cheng L, Yin H, Wu Q, Wang X, Huang Y, Lau WCY, Yao ZP, Guo Y, Jiang L (2021) A distinct giant coat protein complex II vesicle population in Arabidopsis thaliana. Nat Plants **7:** 1335-1346

Liu T, Salguero P, Petek M, Martinez-Mira C, Balzano-Nogueira L, Ramsak Z, McIntyre L, Gruden K, Tarazona S, Conesa A (2022) PaintOmics 4: new tools for the integrative analysis of multi-omics datasets supported by multiple pathway databases. Nucleic Acids Res **50:** W551-W559

Lothier J, Diab H, Cukier C, Limami AM, Tcherkez G (2020) Metabolic Responses to Waterlogging Differ between Roots and Shoots and Reflect Phloem Transport Alteration in Medicago truncatula. Plants (Basel) **9**

Lozano-Elena F, Fàbregas N, Coleto-Alcudia V, Caño-Delgado AI (2022) Analysis of metabolic dynamics during drought stress in Arabidopsis plants. Scientific Data **9:** 90

Lozano-Juste J, Cutler SR (2016) Hormone signalling: ABA has a breakdown. Nat Plants **2:** 16137

Lundberg SM, Lee S-I (2017) A unified approach to interpreting model predictions. *In* Proceedings of the 31st International Conference on Neural Information Processing Systems. Curran Associates Inc., Long Beach, California, USA, pp 4768–4777

Mahmud S, Ullah C, Kortz A, Bhattacharyya S, Yu P, Gershenzon J, Vothknecht UC (2022) Constitutive expression of JASMONATE RESISTANT 1 induces molecular changes that prime the plants to better withstand drought. Plant Cell Environ **45:** 2906-2922

Manjunath KK, Krishna H, Devate NB, Sunilkumar VP, Patil SP, Chauhan D, Singh S, Kumar S, Jain N, Singh GP, Singh PK (2023) QTL mapping: insights into genomic regions governing component traits of yield under combined heat and drought stress in wheat. Front Genet **14:** 1282240

Mathur S, Agrawal D, Jajoo A (2014) Photosynthesis: Response to high temperature stress. Journal of Photochemistry and Photobiology B: Biology **137:** 116-126

Mishra S, Srivastava AK, Khan AW, Tran LP, Nguyen HT (2024) The era of panomics-driven gene discovery in plants. Trends Plant Sci

Mittler R (2006) Abiotic stress, the field environment and stress combination. Trends Plant Sci **11:** 15-19

Nakamura Y, Mithofer A, Kombrink E, Boland W, Hamamoto S, Uozumi N, Tohma K, Ueda M (2011) 12-hydroxyjasmonic acid glucoside is a COI1-JAZ-independent activator of leaf-closing movement in Samanea saman. Plant Physiol **155:** 1226-1236

Navarro C, Abelenda JA, Cruz-Oro E, Cuellar CA, Tamaki S, Silva J, Shimamoto K, Prat S (2011) Control of flowering and storage organ formation in potato by FLOWERING LOCUS T. Nature **478:** 119-122

Núñez-Lillo G, Ponce E, Arancibia-Guerra C, Carpentier S, Carrasco-Pancorbo A, Olmo-García L, Chirinos R, Campos D, Campos-Vargas R, Meneses C, Pedreschi R (2023) A multiomics integrative analysis of color de-synchronization with softening of 'Hass' avocado fruit: A first insight into a complex physiological disorder. Food Chemistry **408:** 135215

Núñez-Lillo G, Ponce E, Beyer CP, Álvaro JE, Meneses C, Pedreschi R (2024) A First Omics Data Integration Approach in Hass Avocados to Evaluate Rootstock–Scion Interactions: From Aerial and Root Plant Growth to Fruit Development. Plants **13:** 603

Obata T, Klemens PAW, Rosado-Souza L, Schlereth A, Gisel A, Stavolone L, Zierer W, Morales N, Mueller LA, Zeeman SC, Ludewig F, Stitt M, Sonnewald U, Neuhaus HE, Fernie AR (2020) Metabolic profiles of six African cultivars of cassava (Manihot esculenta Crantz) highlight bottlenecks of root yield. Plant J **102:** 1202-1219

Paoletti AC, Parmely TJ, Tomomori-Sato C, Sato S, Zhu D, Conaway RC, Conaway JW, Florens L, Washburn MP (2006) Quantitative proteomic analysis of distinct mammalian Mediator complexes using normalized spectral abundance factors. Proc Natl Acad Sci U S A **103:** 18928-18933

Park JS, Park SJ, Kwon SY, Shin AY, Moon KB, Park JM, Cho HS, Park SU, Jeon JH, Kim HS, Lee HJ (2022) Temporally distinct regulatory pathways coordinate thermo-responsive storage organ formation in potato. Cell Rep **38:** 110579

Petek M, Rotter A, Kogovsek P, Baebler S, Mithofer A, Gruden K (2014) Potato virus Y infection hinders potato defence response and renders plants more vulnerable to Colorado potato beetle attack. Mol Ecol **23:** 5378-5391

Petek M, Zagorscak M, Ramsak Z, Sanders S, Tomaz S, Tseng E, Zouine M, Coll A, Gruden K (2020) Cultivar-specific transcriptome and pan-transcriptome reconstruction of tetraploid potato. Sci Data **7:** 249

Ployet R, Veneziano Labate MT, Regiani Cataldi T, Christina M, Morel M, San Clemente H, Denis M, Favreau B, Tomazello Filho M, Laclau J-P, Labate CA, Chaix G, Grima-Pettenati J, Mounet F (2019) A systems biology view of wood formation in Eucalyptus grandis trees submitted to different potassium and water regimes. New Phytologist **223:** 766-782

Quint M, Delker C, Franklin KA, Wigge PA, Halliday KJ, van Zanten M (2016) Molecular and genetic control of plant thermomorphogenesis. Nat Plants **2:** 15190

Renziehausen T, Frings S, Schmidt-Schippers R (2024) 'Against all floods': plant adaptation to flooding stress and combined abiotic stresses. Plant J

Rivero RM, Mittler R, Blumwald E, Zandalinas SI (2022) Developing climate-resilient crops: improving plant tolerance to stress combination. Plant J **109:** 373-389

Rohart F, Gautier B, Singh A, Le Cao KA (2017) mixOmics: An R package for 'omics feature selection and multiple data integration. PLoS Comput Biol **13:** e1005752

Sato H, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K (2024) Complex plant responses to drought and heat stress under climate change. Plant J

Sauter M (2013) Root responses to flooding. Curr Opin Plant Biol **16:** 282-286

Singh A, Shannon CP, Gautier B, Rohart F, Vacher M, Tebbutt SJ, Lê Cao K-A (2019) DIABLO: an integrative approach for identifying key molecular drivers from multi-omics assays. Bioinformatics **35:** 3055-3062

Sinha R, Pelaez-Vico MA, Shostak B, Nguyen TT, Pascual LS, Ogden AM, Lyu Z, Zandalinas SI, Joshi T, Fritschi FB, Mittler R (2024) The effects of multifactorial stress combination on rice and maize. Plant Physiol **194:** 1358-1369

Siroka J, Brunoni F, Pencik A, Mik V, Zukauskaite A, Strnad M, Novak O, Flokova K (2022) High-throughput interspecies profiling of acidic plant hormones using miniaturised sample processing. Plant Methods **18:** 122

Smith AM, Zeeman SC (2006) Quantification of starch in plant tissues. Nat Protoc **1:** 1342-1345

Stael S, Kmiecik P, Willems P, Van Der Kelen K, Coll NS, Teige M, Van Breusegem F (2015) Plant innate immunity--sunny side up? Trends Plant Sci **20:** 3-11

Tang R, Niu S, Zhang G, Chen G, Haroon M, Yang Q, Rajora OP, Li X-Q (2018) Physiological and growth responses of potato cultivars to heat stress. Botany **96:** 897-912

Trapero-Mozos A, Morris WL, Ducreux LJM, McLean K, Stephens J, Torrance L, Bryan GJ, Hancock RD, Taylor MA (2018) Engineering heat tolerance in potato by temperature-dependent expression of a specific allele of HEAT-SHOCK COGNATE 70. Plant Biotechnol J **16:** 197-207

von Gehren P, Bomers S, Tripolt T, Söllinger J, Prat N, Redondo B, Vorss R, Teige M, Kamptner A, Ribarits A (2023) Farmers Feel the Climate Change: Variety Choice as an Adaptation Strategy of European Potato Farmers. Climate **11:** 189

Wasternack C, Feussner I (2018) The Oxylipin Pathways: Biochemistry and Function. Annu Rev Plant Biol **69:** 363-386

Weckwerth W, Ghatak A, Bellaire A, Chaturvedi P, Varshney RK (2020) PANOMICS meets germplasm. Plant Biotechnol J **18:** 1507-1525

Weng JK, Ye M, Li B, Noel JP (2016) Co-evolution of Hormone Metabolism and Signaling Networks Expands Plant Adaptive Plasticity. Cell **166:** 881-893

Wu Q, Su N, Huang X, Cui J, Shabala L, Zhou M, Yu M, Shabala S (2021) Hypoxia-induced increase in GABA content is essential for restoration of membrane potential and preventing ROS-induced disturbance to ion homeostasis. Plant Commun **2:** 100188

Yang W, Feng H, Zhang X, Zhang J, Doonan JH, Batchelor WD, Xiong L, Yan J (2020) Crop Phenomics and High-Throughput Phenotyping: Past Decades, Current Challenges, and Future Perspectives. Mol Plant **13:** 187-214

Yoshida T, Fernie AR (2024) Hormonal regulation of plant primary metabolism under drought. J Exp Bot **75:** 1714-1725

Yoshihara T, Omir E-SA, Koshino H, Sakamura S, Kkuta Y, Koda Y (1989) Structure of a Tuber-inducing Stimulus from Potato Leaves (Solanum tuberosum L.). Agricultural and Biological Chemistry **53:** 2835-2837

Zaki HEM, Radwan KSA (2022) Response of potato (Solanum tuberosum L.) cultivars to drought stress under in vitro and field conditions. Chemical and Biological Technologies in Agriculture **9:** 1

Zandalinas SI, Fritschi FB, Mittler R (2021) Global Warming, Climate Change, and Environmental Pollution: Recipe for a Multifactorial Stress Combination Disaster. Trends Plant Sci **26:** 588-599

Zandalinas SI, Pelaez-Vico MA, Sinha R, Pascual LS, Mittler R (2023) The impact of multifactorial stress combination on plants, crops, and ecosystems: how should we prepare for what comes next? Plant J

Zeng ZL, Wang XQ, Zhang SB, Huang W (2024) Mesophyll conductance limits photosynthesis in fluctuating light under combined drought and heat stresses. Plant Physiol **194:** 1498-1511

Zhang H, Sonnewald U (2017) Differences and commonalities of plant responses to single and combined stresses. Plant J **90:** 839-855

Zhang H, Zhu J, Gong Z, Zhu JK (2022) Abiotic stress responses in plants. Nat Rev Genet **23:** 104-119

Zhang R, Zhang C, Yu C, Dong J, Hu J (2022) Integration of multi-omics technologies for crop improvement: Status and prospects. Front Bioinform **2:** 1027457

Zhao Y, Zhang W, Abou-Elwafa SF, Shabala S, Xu L (2021) Understanding a Mechanistic Basis of ABA Involvement in Plant Adaptation to Soil Flooding: The Current Standing. Plants (Basel) **10**

Figure 1. Overview of the experimental design for single- and combined stress treatments and multi-omics sampling. A) Summary of cultivation conditions. Timeline of the experimental set-up and applied stress treatments, including the recovery phase in potato cv. Desirée. Timing and duration of stress treatment and days for tissue sampling are shown. (B) Actions comprised cultivation in the growing chambers and daily phenotyping with a set of sensors using the PlantScreen™ phenotyping platform at PSI Research Center. (C) Automated image analysis pipeline was used to extract quantitative traits for morphological, physiological, and biochemical performance characterization of the plants during the stress treatment and recovery phase. Side view colour segmented RGB images of plants at selected time points of tissue sampling (left panel) and daily plant volume (m^3) calculated from top and multiple angle side view RGB images (right panel). N=6.

A

C
D
H
HD
W
HDW

Day 1   Day 7   Day 14   Day 21

Control   Drought   Heat   Heat + Drought   Waterlogging   Heat + Drought + Waterlogging

B
C
H
W
D
HD
HDW
Recovery

QY_Lss

Treatment days

C

Fv/Fm_Lss

Treatment days

D

qL_Lss

Treatment days

E

ΔT (°C)

Treatment days

F

WUE (dm^3 g^{-1})

Treatment days

G

Number of tubers

Kruskal–Wallis, p = 0.015

C   D   H   HD   W
Treatment

H

Harvest Index

Kruskal–Wallis, p = 0.0024

C   D   H   HD   W
Treatment

Figure 2. Physiological profiling using high-throughput phenotyping platforms reveals distinct responses to single and combined stresses. A) Pixel-by-pixel false colour images of operating efficiency of photosystem II in light steady state (QY_Lss) captured by kinetic chlorophyll fluorescence measurement. Images for selected time points of tissue sampling are shown. Colour coding of the treatments apply for the entire figure. Vertical dashed lines indicate the onset and end of drought. B) QY_Lss values extracted from images for each individual time point. C) Steady-state fluorescence of maximum efficiency of PSII photochemistry in the light trait based on chlorophyll fluorescence top view (Fv/Fm_Lss). D) steady-state estimation of the fraction of open reaction centres in PSII trait in light based on chlorophyll fluorescence top view (qL_Lss). E) Difference between canopy average temperature extracted from thermal IR images and air temperature measured in the thermal IR imaging unit (ΔT). F) Water use efficiency (WUE) based on plant volume and water consumption. A-F) Black dotted lines reflect the initiation and removal of drought stress, respectively. See Supplementary Table 3 for Statistical evaluation of differences between groups using Wilcoxon test. G) Tuber numbers counted per plant on the last day of the experiment (Day 28 = 60 days of cultivation). H) Harvest index calculated from the total biomass and tuber weight on the last day of the experiment. G-H) Measurements, mean and standard deviation are shown (n = 4). Statistical evaluation of differences between groups is given by the non-parametric Kruskal–Wallis test (*one-way ANOVA on ranks*); p-value above x-axis, where asterisk denotes p-value < 0.05. See Figure 1A for scheme on stress treatments.

Figure 3: Integrated analysis of measured and generated data permits global visualization and multi-level amalgamation of potato stress responses. A) Schematics of tissue sampling protocol. 2^{nd} and 3^{rd} leaves were harvested for destructive "omics" analysis, 4^{th} leaf was used for relative water content calculation. Remaining plant tissue was quantified to obtain total above-ground biomass and tuber yield. B) Overview of data analysis pipeline. C) Dataset overview: multidimensional scaling shows combined HDW stressed plants as extremes, the centroid of each plant group is shown. D) Most informative variables from the phenomics level. Pearson correlation coefficients between them are presented as hierarchically clustered heat map in waterlogging and heat stress.

Figure 4: Integration of multi-omics data in a knowledge-based metabolic and signalling network. A) Structure of knowledge network. Individual studied components are coloured according to their function in different pathways. B) To compare the effects of different stresses on the overall state of the plant, we overlaid the knowledge networks with measured changes in component concentration. Nodes are coloured by log2 fold changes (red – increase in stress compared to control, blue – decrease in stress compared to control, grey – measurement not available) shown for two time points: sampling day 8 and sampling day 14 for the different stress treatments, days of stress treatment are given with each network (for more details of the set up see Figure 1A).

Figure 5: Combined heat and drought stress trigger distinct responses compared to each individual response. Additive effect of combined stress is most pronounced for branched chain amino acids accumulation and JA signalling response. A-C) Heatmaps showing log2FC (FDR p-value < 0.05) in individual stress heat (H) or drought (D) stress in comparison to combined one (HD) for targeted molecular analyses. Label colours indicate pathway associated with each molecule as in the Knowledge network (see Figure 4 for legend). A) Changes in metabolite levels. B) Changes in hormone levels, and C) Changes in selected stress-related transcripts. D) Changes observed on proteomics level. Results of Gene Set Enrichment Analysis (FDR q-value < 0.1) are shown. For more information see Supp. Table 6. C) Biochemical knowledge network showing changes under combined HD stress at day 14 (treatment day 7). In this version of knowledge network, only nodes that were significantly differentially expressed (vs. control conditions) are coloured and the connections between two differentially expressed nodes are coloured black. Node full black border indicates molecules with higher expression levels in HD compared to H and/or D alone. Dashed black border indicates molecules with lower expression levels in HD compared to H and/or D alone (difference of log2FC > 0.5).

Figure 6: Waterlogging triggers drought-stress like molecular responses in potato. A-D) Heatmaps showing log2FC (FDR p-value < 0.05) changes in A) metabolite levels, B) phytohormones, C) selected stress-related transcripts. D) Volcano plot of differential proteomics analysis at day 7. Proteins with FDR p-value < 0.05 shown as blue (downregulated) and red (upregulated) dots. For more information see Supp. Table 5. E) Knowledge network of waterlogging stress at day 1 and day 7 (unfiltered, colour range [-2, 2]). For legend see Figure 4.

Figure 7. Schematic summary of multilevel responses to single and combined heat, drought and waterlogging stresses. Selected variables from each level are shown. Summary of molecular responses (hormones, metabolites and transcripts) was based on the comparisons illustrated in Figures 5 and 6. Summary of morphophysiological responses were based on the data from the last day of the experiment (Day 28), which includes tuber information. Proteomics data set is not included here due to the small dataset of differentially expressed proteins in the waterlogging treatment. Degree of increase or decrease is not specified.