Interplay of Anchor and Latch Residues at Enzyme Active Site in Preserving Induced Fit Configurations for Efficient Catalysis: A Case Study of CARNMT1 for Histidine Methylation

Hao Deng[#,1,‡], Adua Rahman[#,2], Ping Qian*[,1,4]and Hong Guo*[,2]

[1]*Chemistry and Material Science Faculty, Shandong Agricultural University, Taian 271018, P. R. China*

[2] *Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN 37996, United States*

[3] *Centers for Disease Control and Prevention, Taian 271000, P. R. China*

[4] *Key Laboratory of Agricultural Film Application of Ministry of Agriculture and Rural Affairs, Taian 271018, P. R. China*

[#] *These authors make equal contributions*

[‡] Current Address: *School of Chemistry and Chemical Engineering, Huazhong University of Science and Technology, Wuhan 430074, P. R. China*

E-mail address: qianp@sdau.edu.cn (to P. Qian) and hguo1@utk.edu (to H. Guo)

ABSTRACT

Formation of enzyme-substrate complex is the initial step for enzyme catalysis. Widely used models for explanation of formation of precise orientations of catalytic groups in enzyme-substrate complexes for enzyme's action include lock-and-key, induced-fit and conformational selection models as well as their extensions. Nevertheless, formation of catalytic competent orientations of residues in enzyme-substrate complexes may not be sufficient to lead to efficient catalysis. Such orientation of catalytic groups relative to substrate may need to be preserved during part of enzyme-catalyzed reaction step to allow catalytic residues to exert their effects during catalysis. Pre-mature collapse of such configuration may lead to a significant reduction of catalytic efficiency. Here MD and free energy simulations based on QM/MM potentials are carried out for CARNMT1 to understand preservation of catalytic competent induced-fit configuration during catalysis. We examine interplay of certain anchor residues that occupy similar positions before and after substrate binding (Asp316, Tyr386 and Tyr396) with the latch residue (His347) that undergoes large induced-fit conformational changes. It is demonstrated that the wild-type enzyme can preserve catalytic competent induced-fit configuration well after the reaction passes TS. For Tyr386Ala and Tyr396Asp, the induced-fit configuration can be formed in the mutant-substrate complex but collapses in the early stage of reaction before it reaches TS. His347Phe also fails to preserve the active induced-fit configuration. For Asp316Ala, the active induced-fit configuration cannot be formed in the mutant-substrate complex. Consistent with these observations and experimental results, the free energy barrier for methyl transfer increases significantly because of the mutations. The present study suggests that generating precise orientations of catalytic groups in enzyme-substrate complexes through the well-known models may be a necessary but not sufficient condition for efficient catalysis and that anchor residues may play an important role in generating and preserving catalytic competent induced-fit configurations during catalysis.

KEYWORDS: Molecular Recognition, Lock-and-Key, Induced Fit, Conformational Selection, Enzyme Catalysis, CARNMT1, QM/MM, MD simulations

INTRODUCTION

Molecular recognition is of fundamental importance in biology. Understanding its principles is crucial for designing drugs and studying diverse biological processes, including, but limited to, enzyme catalysis, antibody–antigen recognition, protein-protein interactions, protein synthesis and transcriptional regulation. Therefore, this topic has attracted considerable attention, especially for protein, ligand interactions.[1-11] The early developments of Emil Fisher's lock-and-key hypothesis[12] and Koshland's induced fit model[13] for molecular recognition were driven largely by the efforts to explain enzyme's activity and specificity. These textbook models were later extended to include original and extended conformational selection models.[3, 14, 15] The lock-and-key (LK) model hypothesizes that the way enzymes interact with substrates/non-substrates conceptually resembles that locks match keys (Figure 1a). Complimentary structural fit between substrate and enzyme is important in this model, and conformational changes of either enzyme or substrate are not considered during binding processes. The induced fit (IF) model, on the other hand, recognizes that the binding of a substrate may induce a change in the three-dimensional relationship of the active site residues, and precise orientation of catalytic groups required for enzyme's action can be generated through such induced fit process (Figure 1b, the upper pathway).[6, 13] It should be pointed out that catalytic competent configurations of enzyme-substrate complexes may also be generated through conformational changes of substrates induced by enzymes.[16, 17] Original conformational selection model (CS) recognizes that proteins are under dynamical fluctuations and that ligand may select and bind to the one that is compatible and shift the conformational ensemble towards this state (Figure 1b, the lower pathway).[14, 15] Extended conformational selection model allows conformational selection to be followed by conformational adjustment.[3]

Figure 1. (a) lock-and-key (LK) model. (b) Induced fit model (IF) and original conformational selection model (CS). Blue: enzyme; Yellow: substrate.

As mentioned above, efforts to understand enzyme function and specificity have played a crucial role in shaping our current view for molecular recognition. However, formation of enzyme-substrate complex is only the initial step for enzyme catalysis, and the generation of catalytic competent orientations of residues in enzyme-substrate complexes through the lock-and-key (LK), induced fit (IF) and/or conformational selection (CS) processes may not be sufficient to lead to efficient catalysis. For an enzyme, correct molecular recognition is not only essential for its substrate, but also for the intermediate states for certain part of the reaction that the enzyme catalyzes. The catalytic competent orientation of enzyme's groups relative to substrate may therefore need to be preserved for those intermediate states of the reaction step to allow catalytic residues to exert their effects and make the reaction to pass the transition state (TS) easier. Pre-mature collapse of such configuration (e.g., due to certain mutations) may lead to a significant reduction of catalytic efficiency. Moreover, it is known that, in addition to the residues that undergo induced fit conformational changes upon substrate binding (e.g., latch residues), there may be some other active-site residues that occupy similar positions before and after substrate binding (e.g., anchor residues). Previous studies have shown that protein-protein interactions may go through a local lock-and-key mechanism from anchor residue followed by a induced-fit process involving latch residues.[18] Such step by step process may also exist for protein-ligand interactions.[3] However, it is not clear how anchor and latch residues would work together to preserve the active induced fit configurations during parts of enzyme catalyzed reactions and lead to efficient catalysis.

In this paper, QM/MM MD and PMF simulations were performed for wild-type CARNMT1 and certain mutants (Asp316Ala, His347Phe, Tyr386Ala and Tyr396Asp) to understand the interplay of anchor and latch residues in preserving active induced fit

Figure 2. Superposition of apo enzyme (5YF0, orange) and enzyme-substrate complex (5YF1 with SAM replaced by SFG, blue). The $\chi(C\text{-}C_\alpha\text{-}C_\beta\text{-}C_\gamma)$ dihedral angle changes from 55° in 5YF0 to 157° in 5YF1, and r(Carnosine-Cε1$\cdots$H347-Cε1) decreases from about 7.5Å to 3.5Å accordingly. Part of the SFG/SAM, carnosine and water molecules are shown in ball and stick, and the rest of the active sites are shown in stick. Waters from 5YF1 are colored in red, and those from 5YF0 in orange. Some distances are also given (in angstroms).

configurations for efficient catalysis. CARNMT1 is a histidine $N_{\delta 1}$ position-specific methyltransferase and catalyzes the conversion of carnosine to anserine. Carnosine exerts numerous biological and physiological effects in multiple organs.[19] The crystal structures of the apo enzyme (5YF0) and enzyme-substrate complex (5YF1, with SAM replaced by SFG) have been determined.[20] The superposition of the two structures is shown in Figure 2. It can be seen from Figure 2 that the carnosine histidine is deeply embedded into the active site pocket surrounded by some amino acid residues such as Phe313, Asp316, Leu345, His347, Tyr386, Tyr396, and Tyr398 with the substrate configuration stabilized by the corresponding interactions involving these residues. The structures of the apo enzyme (5YF0) and enzyme-substrate complex (5YF1) are highly similar with RMSD (based on C_α) of only 0.3Å. The RMSD value based on all atoms superposed for the active site residues without His347 is also quite small (0.3Å); the RMSD for His347 is 3.2Å. The site directed mutagenesis study[20] has shown that some of the active site residues are of important for efficient catalysis by the enzyme, including Asp316, Tyr386, Tyr396 as well as His347. CARNMT1 is an ideal system for studying how the active site residues of enzymes can generate and preserve active induced fit configurations during catalysis. This is because the enzyme catalyzes a single chemical step of the S_N2 methyl transfer and the only active site residue that undergoes large induced-fit conformational changes upon substrate binding and product formation is His347. This makes it easier to study the stability of the active induced fit conformation during the enzyme catalyzed reaction with His347 and some other active-site residues being altered. Moreover, the induced fit conformational change of His347 mainly involves the rotation around the C_α-C_β bond of His347 with the corresponding χ(C-C_α-C_β-C_γ) dihedral angle changing from about 55° in the apo enzyme to 157° in the enzyme-substrate complex (Figure 2); the r(Carnosine-$C_{\varepsilon 1}$···H347-$C_{\varepsilon 1}$) distance decreases from about 7.5Å to 3.5Å accordingly. This makes monitoring the formation and collapse of the active induced fit conformation much easier. The enzyme-product complex (5YF2) for CARNMT1 has also been determined,[20] and the structures for the apo enzyme and enzyme-product complex are also highly similar. Interestingly, the positions of His347 in 5YF0 and 5YF2 can be superposed well with similar χ(C-C_α-C_β-C_γ) values (55° in 5YF0 and 51° in 5YF2). Thus, only the substrate can lead to the induced fit conformational change involving His347, while the product (a non-substrate) cannot. It has been proposed that such "hold-release" switching role of His347 can benefit substrate binding and product release.[20]

The results from the QM/MM MD and PMF simulations of this work showed that the wild-type enzyme could preserve the catalytic competent induced-fit configuration even after the reaction well

passes the transition state (TS). For Asp316Ala, the active induced fit configuration cannot be formed in the Asp316Ala-substrate complex due to missing of certain key interaction between Asp316 and the substrate. Although the active induced fit configuration can be formed in Tyr386Ala-substrate, Tyr396Asp-substrate and His347Phe-substrate complexes, the active configuration collapses in each of the cases in the early stage of reaction before it can reach TS, leading to an increase of the free energy barrier at the transition state and a lower catalytic efficiency. Thus, generating precise orientations of catalytic groups in enzyme-substrate complexes through the well-known models may be a necessary but not sufficient condition for efficient catalysis. Moreover, this work shows that anchor residues may play an important role in preserving catalytic competent induced-fit configurations that are required for efficient catalysis during parts of enzyme-catalyzed reactions. The principle discussed here may be used to explain observed losses of activities due to certain mutations for some other enzymes.

METHODS

The CHARMM program[21, 22] was used to perform the QM/MM MD and free energy simulations to study the formation and collapse of the active induced fit configuration in the enzyme-substrate complex and during the catalysis. The effects of the Asp316→Ala, His347→Phe, Tyr386→Ala and Tyr396→Asp mutations on the stability of the induced fit configuration and the activity of the $N_{\delta 1}$ methylation were also determined. The crystal structure of CARNMT1 (PDB ID: 5YF1) containing carnosine and SFG and was used to create the reaction state structure of the enzyme-substrate complex for methylation,[20] and the N and C atoms of SFG was manually changed to C and S atoms, respectively, to generate SAM. The crystal structure of the apo enzyme (PDB ID: 5YF0) containing SAM was also used for the simulations. The simulations were also performed for the case where carnosine was removed from 5YF1 in the initial structure (with SFG changed to SAM). This allows testing whether His347 could move back to its position in the apo enzyme when the substrate was removed from the enzyme-substrate complex. To perform the QM/MM MD simulations on the mutants,

Figure 3. Relative orientation of SAM and carnosine in the reactant complex. Here, θ is defined as the angle between two vectors r_1 and r_2, which represent the direction of the C_M-S_δ bond and the direction of the lone pair of electrons on $N_{\delta 1}$, respectively, which can determine the connection between the distributions and reactant structure properties.

the Asp316→Ala, His347→Phe, Tyr386→Ala, or Tyr396→Asp mutation was performed manually in the crystal structure of the wild type (5YF1).

The −CH$_2$−CH$_2$−S$^+$(Me)−CH$_2$− part of SAM and the whole substrate were treated by QM, and the rest of the system was treated by MM. The QM region and MM region were separated by using the link-atom approach[23] implemented in CHARMM; the QM region contains a total of 44 atoms, with the net charge of +1. The stochastic boundary molecular dynamics method[24] was used for the QM/MM MD and PMF simulations. A modified TIP3 water model[25] was used for the solvent. The DFTB3 method[26-29] was applied to the QM atoms, and the CHARMM potential function (PARAM27)[30] was adopted for the MM atoms. Hydrogen atoms of the system were built with the HBUILD module[31] in CHARMM. The reference center assigned by the system was chosen to be the N$_{\delta 1}$ atom of target substrate carnosine. The system was separated into a reaction zone and a reservoir region, and the reaction zone was further divided into a reaction region (a sphere with radius r of 20 Å) and a buffer region (20 Å ⩽ r ⩽ 22 Å). The whole system contained around 5700 atoms, including about 460 water molecules. The initial structures were optimized for the entire stochastic boundary systems with the steepest descent (SD) and adopted-basis Newton−Raphson (ABNR) methods. The systems were gradually heated from 50.0 to 310.15 K in 50 ps. The time step for the integration of equation of motion was 1 fs, and the coordinates were saved every 50 fs for analyses. For each of the systems, the QM/MM MD simulations of 2 ns were carried out. The distribution maps of r(C$_M$-N$_{\delta 1}$) and θ (Figure 3) were generated in each case and the fluctuations of the χ(C-C$_\alpha$-C$_\beta$-C$_\gamma$) dihedral angle and r(carnosine-C$_{\varepsilon 1}$···H347-C$_{\varepsilon 1}$) distance of His347 were monitored.

The umbrella sampling method[32] was applied along with the weighted histogram analysis method (WHAM)[33] in the determination of the free energy change along the reaction coordinate for the methyl transfer from S-adenosyl-L-methionine (SAM) to the N$_{\delta 1}$ atom of carnosine. The reaction coordinate was defined as a linear combination of r(C$_M$−N$_{\delta 1}$) and r(C$_M$−S$_\delta$) [R = r(C$_M$−S$_\delta$) − r(C$_M$−N$_{\delta 1}$)]. For the methyl transfer process, 28 windows were used, and for each window, the production runs of 100 ps were performed after 50 ps equilibration. The force constants of the harmonic biasing potentials used in the PMF simulations were 50 to 400 kcal·mol^{-1}·Å^{-2}. In each case, four independent PMF simulations were performed. The free energies and statistical errors were taken as the average values and standard deviations from the four runs, respectively. The fluctuations of the χ(C-C$_\alpha$-C$_\beta$-C$_\gamma$) dihedral angle as well as r(carnosine-C$_{\varepsilon 1}$···H347-C$_{\varepsilon 1}$) distance of His347 were monitored in each PMF window along the

reaction coordinate to detect when the induced fit configuration was formed or collapsed during the methyl transfer reaction.

RESULTS AND DISCUSSIONS

Ability of Wild Type to Preserve Active Induced-Fit Configuration well after TS in Enzyme-Catalyzed Reaction

The average active site structure of the CARNMT1-SAM-Carnosine complex obtained from the QM/MM MD simulations is shown in Figure 4a. The interactions between carnosine and the key residues in the active site region (e.g., Asp316, His347, Tyr386, Tyr396, etc.) are likely to play a crucial role in the

Figure 4. (a) The average structure for the active site of the reactant complex containing SAM and carnosine in wild-type CARNMT1 (left) along with the distribution map of $r(C_M-N_{\delta1})$ and θ (right) obtained from the QM/MM MD simulations. The $\chi(C-C_\alpha-C_\beta-C_\gamma)$ dihedral angle of His347 is about 160°, and r(carnosine-$C_{\epsilon1}\cdots$H347-$C_{\epsilon1}$) is 3.7Å. **(b)** The fluctuations of the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ dihedral angle of His347 in the PMF windows along the reaction coordinate. The red vertical bar designates the location for the transition state of the methyl transfer. **(c)** The average structure for the active site near the transition state obtained from the QM/MM PMF simulations for methylation in wild-type CARNMT1. **(d)** Left: The $\chi(C-C_\alpha-C_\beta-C_\gamma)$ values for the apo enzyme during the QM/MM simulations based on the crystal structure of 5YF0. The red circles indicate the system reached the induced fit configuration even without the substrate. Right: The $\chi(C-C_\alpha-C_\beta-C_\gamma)$ values from the simulations after the substrate was first removed from the active site in the enzyme-substrate complex.

substrate binding and catalysis. The interactions obtained from the simulations are quite similar to those observed experimentally (Figure 2).[20] For instance, the two oxygen atoms of the Asp316 side chain interact with the $C_{\varepsilon 1}$ and $N_{\varepsilon 2}$ atoms of the target substrate imidazole ring, respectively, in both crystal structure and reactant state from simulations; the average distance to $C_{\varepsilon 1}/N_{\varepsilon 2}$ is 3.43 Å/2.78 Å obtained from the simulations (Figure 4a), while it is 3.16 Å/2.94 Å in the crystal structure (Figure 2). Other interactions that are similar between the crystal structure and structure from the simulations include the "π-π" stacking interaction between the Phe313 benzene ring and carnosine imidazole ring (e.g., ~3.95 Å and ~3.80 Å in Figure 4a and 2, respectively), the "π-π" stacking interaction between the His347 and carnosine imidazole rings (e.g., 3.67 Å in the simulated structure and 3.50 Å in the crystal structure), the "methyl-π" interaction between carnosine and the Tyr386 phenol ring (e.g., 3.73 Å in the simulated structure and 3.64 Å in the crystal structure), and the hydrogen bond between the phenolic hydroxyl group on Tyr398 and the substrate (3.13 Å in the simulated structure and 3.04 Å in the crystal structure). The positions of the residues mentioned above are fixed by the interactions with the substrate carnosine and their neighboring residues. For instance, the interaction between the backbone carbonyl oxygen atom of Leu345 and $N_{\delta 1}-H$ of His347 stabilizes the His347 position, and His347 then forms the "π-π" interaction with the imidazole ring of the substrate. We have also observed some interactions involving the transferable methyl group in our simulated structure as a result of replacing SFG by SAM. For example, the hydroxyl group of Tyr386 and the backbone carbonyl oxygen of Phe313 form carbon hydrogen bonds with the transferable methyl group of SAM. In general, the average structure obtained from the QM/MM MD simulated and crystal structures are highly similar; the existence of some small differences is expected as SFG in the crystal structure was replaced by SAM for the simulations. The r-θ distribution map of the reactant state of wild type is also given in Figure 4a. There are many structures with relatively short $r(C_M-N_{\delta 1})$ distances and small values of the θ angle. The average distance between the target $N_{\delta 1}$ atom and C_M atom of the transferable methyl group is approximately 3.0 Å, and the θ angle is mainly in the range of 0-30°. Therefore, when carnosine binds to the active site of the enzyme, it can form the near-attack conformations to facilitate methyl transfer.

Figure 4b shows the fluctuations of the $\chi(C$-C_α-C_β-$C_\gamma)$ dihedral angle of His347 collected from each of the PMF windows along the reaction coordinate from the free energy simulations. It can be seen that the wild type enzyme can preserve the catalytic competent induced-fit configuration with His347 in the "hold" position [i.e., $\chi(C$-C_α-C_β-$C_\gamma) \sim 160°$] well after the reaction passes TS [R = $r(C_M-S_\delta) - r(C_M-N_{\delta 1})$]

~ 0.2Å]. The induced-fit configuration collapses at R = $r(C_M–S_\delta) – r(C_M–N_{\delta 1})$ ~ 1.3 Å with χ(C-C$_\alpha$-C$_\beta$-C$_\gamma$) of His347 changing back to the value of the "released" position in the apo enzyme and the product complex (~ 60°), consistent with experimental observations.[20] The average structure of the active site near the transition state of the methylation obtained from the QM/MM PMF simulations is given in Figure 4c. Comparison of the structures in Figure 4a and 4c shows that there are some small changes between the reactant state and transition state for the wild-type enzyme. Nevertheless, the "π-π" stacking interaction between the His347 side chain and the imidazole ring of the substrate remains intact at the transition state that helps to generate the well-positioned interactions at the transition state and protect the imidazole ring of the substrate from being exposed to solvent. To examine whether the apo enzyme can reach the induced fit configuration as proposed in the CS model,[14, 15] we performed the QM/MM MD simulations based on 5YF0. The results in Figure 4d (Left) shows that the induced fit configuration can be formed as monitored by χ(C-C$_\alpha$-C$_\beta$-C$_\gamma$) (red circles), even though the frequency of formation of such structure is rather low. We have also examined what would happen if the substrate in the induced fit configuration of the enzyme-substrate complex was removed from the active site. Figure 4d (Right) shows that if the substrate was removed, His347 would move quickly from the "hold" to the "released" position (as observed in the apo enzyme).

Inability to Form or Early Collapse of Active Induced-Fit Configuration in the Mutants

The structural results given in Figure 4a and 4c and previous experimental data[20] support the suggestion that Asp316 stabilizes the configuration of the $N_{\varepsilon 2}$-H tautomeric form of the substrate and makes it possible for the transferable methyl group of SAM to point towards the electron lone pair of $N_{\delta 1}$ of carnosine prior to the reaction. The Asp316 residue in the CARNMT1 seems to play a similar and presumably more important role as the Asn255 residue in SETD3[34]. If the hydrogen bonding interactions between Asp316 and carnosine are removed (e.g., through mutations), the binding of the correct tautomer and the formation of the near-attack reactive state may be compromised, leading to a reduction or loss of the activity, as observed experimentally.[20] The average active-site structure of the Asp316Ala-substrate complex obtained from the simulations is given in Figure 5a along with the distribution map. Although the other active site residues (e.g., Phe313, His347, Leu345 and Tyr386) may also be important in anchoring the $N_{\varepsilon 2}$-H tautomeric form and the formation of the near-attack conformation, they cannot achieve this by themselves without Asp316. Indeed, the distribution map in Figure 5a shows that the

Figure 5. (a), (c), (e) and **(g)** The average structures for the active sites of the Asp316Ala-substrate, Tyr386Ala-substrate, Tyr396Asp-substrate and His347Phe-substrate complexes, respectively, along with the corresponding distribution maps. **(b), (d), (f)** and **(h)** The fluctuations of the χ(C-C_α-C_β-C_γ) dihedral angle of His347 in the PMF windows along the reaction coordinate for the Asp316Ala, Tyr386Ala, Tyr396Asp and His347Phe complexes, respectively.

population of the structures with relatively short $r(C_M-N_{\delta 1})$ distances (about 3.0 Å) and small values of the θ angle (less than 30°) become much smaller in the Asp316Ala mutant compared to the case of wild-type (Figure 4a). Moreover, comparison of the structures of Figure 4a and Figure 5a shows that the imidazole ring of the substrate underwent a rotation of ~70° in Asp316Ala, and this conformational change of the substrate also affects the correct alignment between the methyl donor and acceptor. Moreover, the imidazole ring of His347 flipped away from its position in Figure 4a and is now far away from the substrate (Figure 5a). Interestingly, it occupies a similar position as in the apo enzyme and product complex of wild-type with a $\chi(C-C_\alpha-C_\beta-C_\gamma)$ value of about 60°. Thus, His347 is already in the "released" position in the Asp316Ala-substrate complex, and the induced fit configuration collapsed. The fluctuations of the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ dihedral angle of His347 along the reaction coordinate obtained from the PMF simulations for are given in Figure 5b. As is evident from Figure 5b, the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ values remained around 60° during the methyl transfer reaction, consistent with the structural data. The loss of such induced fit configuration is expected to affect the efficiency of methylation process,[34-41] contributing to the increase of free energy barrier (see below) and decrease of the activity.[20]

The average active site structure of the Tyr386Ala-substrate complex is given in Figure 5c along with the distribution map. Some interactions involving Tyr386 in wild type are lost in the mutant. For instance, the carbon hydrogen bond between the hydroxyl oxygen of Tyr386 and transferable methyl group and "methyl-π" interaction between the substrate and the ring of Tyr386 are lost. Nevertheless, the Tyr386Ala-substrate complex is in the induced fit configuration with the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ value of His347 about 160° (Figure 5c). Moreover, there is a relatively large population of structures with short $r(C_M-N_{\delta 1})$ distances and small θ values, and the near attack conformations can therefore be formed. Figure 5d shows the fluctuations of the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ dihedral angle of His347 along the reaction coordinate obtained from the PMF simulations for Tyr386Ala. As is evident from Figure 5d, the $\chi(C-C_\alpha-C_\beta-C_\gamma)$ value changed from about 160° to 60° in the early stage of methyl transfer reaction [R = $r(C_M-S_\delta) - r(C_M-N_{\delta 1})$ ~ -1.5 Å], well before the reaction reaches the transition state [R = $r(C_M-S_\delta) - r(C_M-N_{\delta 1})$ ~ 0.2Å]. Thus, the active induced fit orientation of catalytic groups relative to substrate cannot be preserved to allow catalytic residues to exert their catalytic effects. The pre-mature collapse of such configuration is expected to affect the efficiency of methylation process, contributing to an increase of free energy barrier (see below) and decrease of the activity.[20] The similar discussion can be for the Tyr396Asp mutant (Figure 5e and 5f) where the active induced fit configuration collapsed at R = $r(C_M-S_\delta) - r(C_M-N_{\delta 1})$ ~ -0.8 Å.

It would be of interest to examine the effect of replacing the imidazole ring by the phenol ring on the maintenance of the induced fit configuration. The average structure and distribution map for the His347Phe-substrate complex are given in Figure 5g. As in the cases of the Tyr386Ala and Tyr396Asp mutants, the His347Phe-substrate complex also has a large population of structures with relatively short $r(CM-N\delta1)$ distances and small θ values. Nevertheless, the active induced fit configuration also collapsed before the reaction reaches the TS [at $R = r(C_M-S_\delta) - r(C_M-N_{\delta1}) \sim$ -0.2 Å], consistent with the experimental observation[20] that more than 80% of the activity was lost due to the His347Phe mutation.

The Free Energy Profiles for Methylation in the Wild-Type and Mutants

The free-energy (PMF) profiles for the methylation reaction in the wild-type CARNMT1 and Asp316Ala, His347Phe, Tyr386Ala and Tyr396Asp mutants as a function of the reaction are given in Figure 6. As is shown in Figure 6, the free energy barriers increase by 3.72, 3.42, 1.71 and 1.14 kcal·mol^{-1} relative to that of wild type as a result of the Asp316→Ala, Tyr386→Ala, and Tyr396→Asp, His347→Phe mutations, respectively. The increases of the free energy barriers are consistent with the decreases of the activity observed in the previous experimental observations.[20] Interestingly, the increase of the free energy barrier compared to that of wild type seems to be correlated with the timing of the collapse of the induced fit configuration. For instance, the Tyr386Ala mutant has its induced fit configuration collapsed earlier [$R = r(C_M-S_\delta) - r(C_M-N_{\delta1})$ ~ -1.5 Å] than the Tyr396Asp mutant [$R = r(C_M-S_\delta) - r(C_M-N_{\delta1})$ ~ -0.8 Å], while its free energy barrier (~ 20 kcal/mol) for the reaction is also higher than that of Tyr396Asp (18.3 kcal/mol).

Figure 6. Free energy (PMF) profiles for the methylation reaction in wild-type CARNMT1, Asp316Ala, His347Phe, Tyr386Ala and Tyr396Asp, respectively, as a function of the reaction coordinate [R = $r(C_M-S_\delta) - r(C_M-N_{\delta1})$]. Wild type: blue line with the free energy barrier of 16.59 kcal/mol; Asp316Ala: yellow line with the free energy barrier of 20.31 kcal/mol; Tyr386Ala: grey line with the free energy barrier of 20.01 kcal/mol; Tyr396Asp: green line with the free energy barrier of 18.30 kcal/mol; His347Phe: orange line with the free energy barrier of 17.73 kcal/mol.

CONCLUSIONS

In this paper, we applied MD and free energy simulations based on QM/MM potentials to understand the importance of preserving catalytic competent induced-fit configuration during catalysis and examine the interplay of certain anchor residues with the latch residue at the active site in maintaining such configuration during the enzyme-catalyzed reaction. It has been demonstrated that the wild type enzyme can preserve the catalytic competent induced-fit configuration well after the reaction passes TS (Figure 7, the top case). For Asp316Ala, the active induced-fit configuration could not be formed in the mutant-substrate complex or TS (Type 1 case for the mutation of key anchor residue in Figure 7). The induced-fit configuration could be formed in the Tyr386Ala-substrate and Tyr396Asp-substrate complexes, but it collapsed in the early stage of reaction before it could reach TS (Type 2 case for the mutation of key anchor residue in Figure 7). His347Phe, in which the latch residue was mutated, also fails to preserve the active induced fit configuration. Consistent with these

Figure 7. Wild-type enzyme can preserve catalytic competent induced-fit configuration at TS. Type 1 mutation of key anchor residue (e.g, Asp316Ala) destroys the enzyme's ability to form the induced-fit configuration with substrate. Type 2 mutation (e.g., Tyr386Ala) leads to the collapse of the induced-fit configuration before TS, even though such configuration can be formed in enzyme-substrate complex.

observations and experimental results, the free energy barrier for methyl transfer increases significantly because of the mutations. The present study suggests that generating precise orientations of catalytic groups in enzyme-substrate complexes through the well-known models may be a necessary but not sufficient condition for efficient catalysis and that anchor residues may play an important role in generating and preserving catalytic competent induced-fit configurations during catalysis. The principle discussed here may be used to explain observed losses of activities due to certain mutations for some other enzymes.

Acknowledgments

We thank Professor Martin Karplus for a gift of the CHARMM program. This work was partly supported by the Natural Science Foundation of China (No. 22177064 to P.Q.) and the Natural Science Foundation of Shandong Province (No. ZR2021MB050 to P.Q.).

References:

1.   Boehr, D. D.; Nussinov, R.; Wright, P. E., The role of dynamic conformational ensembles in biomolecular recognition. *Nature Chemical Biology* **2009,** *5* (11), 789-796.

2.   Boehr, D. D.; Wright, P. E., How do proteins interact? *Science* **2008,** *320* (5882), 1429-1430.

3.   Csermely, P.; Palotai, R.; Nussinov, R., Induced fit, conformational selection and independent dynamic segments: an extended view of binding events. *Trends in Biochemical Sciences* **2010,** *35* (10), 539-546.

4.   Pang, X. D.; Zhou, H. X., Rate Constants and Mechanisms of Protein-Ligand Binding. In *Annual Review of Biophysics, Vol 46*, Dill, K. A., Ed. 2017; Vol. 46, pp 105-130.

5.   Redhair, M.; Atkins, W. M., Analytical and functional aspects of protein-ligand interactions: Beyond induced fit and conformational selection. *Archives of Biochemistry and Biophysics* **2021,** *714*.

6.   Koshland, D. E., THE KEY-LOCK THEORY AND THE INDUCED FIT THEORY. *Angewandte Chemie-International Edition* **1994,** *33* (23-24), 2375-2378.

7.   Du, X.; Li, Y.; Xia, Y. L.; Ai, S. M.; Liang, J.; Sang, P.; Ji, X. L.; Liu, S. Q., Insights into Protein-Ligand Interactions: Mechanisms, Models, and Methods. *International Journal of Molecular Sciences* **2016,** *17* (2).

8.   Johnson, K. A., Role of induced fit in enzyme specificity: A molecular forward/reverse switch. *Journal of Biological Chemistry* **2008,** *283* (39), 26297-26301.

9.   Houk, K. N.; Leach, A. G.; Kim, S. P.; Zhang, X. Y., Binding affinities of host-guest, protein-ligand, and protein-transition-state complexes. *Angewandte Chemie-International Edition* **2003,** *42* (40), 4872-4897.

10.   Herschlag, D., THE ROLE OF INDUCED FIT AND CONFORMATIONAL-CHANGES OF ENZYMES IN SPECIFICITY AND CATALYSIS. *Bioorganic Chemistry* **1988,** *16* (1), 62-96.

11.   Stank, A.; Kokh, D. B.; Fuller, J. C.; Wade, R. C., Protein Binding Pocket Dynamics. *Accounts of Chemical Research* **2016,** *49* (5), 809-815.

12.   E, F., The influence of configuration on enzyme activity. (Translated from German). *Dtsch Chem Ges* **1894,** *27*, 2984 –2993.

13.   Koshland, D. E., APPLICATION OF A THEORY OF ENZYME SPECIFICITY TO PROTEIN SYNTHESIS. *Proceedings of the National Academy of Sciences of the United States of America* **1958,** *44* (2), 98-104.

14.   Tsai, C. J.; Kumar, S.; Ma, B. Y.; Nussinov, R., Folding funnels, binding funnels, and protein function. *Protein Science* **1999,** *8* (6), 1181-1190.

15.   Tsai, C. J.; Ma, B. Y.; Nussinov, R., Folding and binding cascades: Shifts in energy landscapes. *Proceedings of the National Academy of Sciences of the United States of America* **1999,** *96* (18), 9970-9972.

16.   Guo, H.; Cui, Q.; Lipscomb, W. N.; Karplus, M., Substrate conformational transitions in the active site of chorismate mutase: Their role in the catalytic mechanism. *Proceedings of the National Academy of Sciences of the United States of America* **2001,** *98* (16), 9032-9037.

17.   Guo, H.; Cui, Q.; Lipscomb, W. N.; Karplus, M., Understanding the role of active-site residues in chorismate mutase catalysis from molecular-dynamics simulations. *Angewandte Chemie-International Edition* **2003,** *42* (13), 1508-1511.

18.   Rajamani, D.; Thiel, S.; Vajda, S.; Camacho, C. J., Anchor residues in protein-protein interactions. *Proceedings of the National Academy of Sciences of the United States of America* **2004,** *101* (31), 11287-11292.

19.   Boldyrev, A. A.; Aldini, G.; Derave, W., Physiology and pathophysiology of carnosine. *Physiological reviews* **2013**.

20.   Cao, R.; Zhang, X.; Liu, X.; Li, Y.; Li, H., Molecular basis for histidine N1 position-specific methylation by CARNMT1. *Cell Research* **2018,** *28* (4), 494-496.

21.   Brooks, B. R.; Brooks, C. L.; Mackerell, A. D.; Nilsson, L.; Petrella, R. J.; Roux, B.; Won, Y.; Archontis, G.; Bartels, C.; Boresch, S.; Caflisch, A.; Caves, L.; Cui, Q.; Dinner, A. R.; Feig, M.; Fischer, S.; Gao, J.; Hodoscek, M.; Im, W.; Kuczera, K.;

Lazaridis, T.; Ma, J.; Ovchinnikov, V.; Paci, E.; Pastor, R. W.; Post, C. B.; Pu, J. Z.; Schaefer, M.; Tidor, B.; Venable, R. M.; Woodcock, H. L.; Wu, X.; Yang, W.; York, D. M.; Karplus, M., CHARMM: The Biomolecular Simulation Program. *Journal of Computational Chemistry* **2009,** *30* (10), 1545-1614.

22.   Brooks, B. R.; Bruccoleri, R. E.; Olafson, B. D.; States, D. J.; Swaminathan, S.; Karplus, M., CHARMM - a program for macromolecular energy, minimization, and dynamics calculations. *Journal of Computational Chemistry* **1983,** *4* (2), 187-217.

23.   Field, M. J.; Bash, P. A.; Karplus, M., A COMBINED QUANTUM-MECHANICAL AND MOLECULAR MECHANICAL POTENTIAL FOR MOLECULAR-DYNAMICS SIMULATIONS. *Journal of Computational Chemistry* **1990,** *11* (6), 700-733.

24.   Brooks, C. L.; Brunger, A.; Karplus, M., Active-site dynamics in protein molecules - a stochastic boundary molecular-dynamics approach. *Biopolymers* **1985,** *24* (5), 843-865.

25.   Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L., COMPARISON OF SIMPLE POTENTIAL FUNCTIONS FOR SIMULATING LIQUID WATER. *Journal of Chemical Physics* **1983,** *79* (2), 926-935.

26.   Elstner, M.; Porezag, D.; Jungnickel, G.; Elsner, J.; Haugk, M.; Frauenheim, T.; Suhai, S.; Seifert, G., Self-consistent-charge density-functional tight-binding method for simulations of complex materials properties. *Physical Review B* **1998,** *58* (11), 7260-7268.

27.   Cui, Q.; Elstner, M.; Kaxiras, E.; Frauenheim, T.; Karplus, M., A QM/MM implementation of the self-consistent charge density functional tight binding (SCC-DFTB) method. *Journal of Physical Chemistry B* **2001,** *105* (2), 569-585.

28.   Christensen, A. S.; Kubař, T.; Cui, Q.; Elstner, M., Semiempirical quantum mechanical methods for noncovalent interactions for chemical and biochemical applications. *Chem. Rev.* **2016,** *116* (9), 5301-37.

29.   Lu, X.; Gaus, M.; Elstner, M.; Cui, Q., Parametrization of DFTB3/3ob for magnesium and zinc for chemical and biological applications. *Journal of Physical Chemistry B* **2015,** *119* (3), 1062-1082.

30.   MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.; Field, M. J.; Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir, L.; Kuczera, K.; Lau, F. T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.; Prodhom, B.; Reiher, W. E.; Roux, B.; Schlenkrich, M.; Smith, J. C.; Stote, R.; Straub, J.; Watanabe, M.; Wiorkiewicz-Kuczera, J.; Yin, D.; Karplus, M., All-atom empirical potential for molecular modeling and dynamics studies of proteins. *Journal of Physical Chemistry B* **1998,** *102* (18), 3586-3616.

31.   Brünger, A. T.; Karplus, M., Polar hydrogen positions in proteins: empirical energy placement and neutron diffraction comparison. *Proteins* **1988,** *4* (2), 148-156.

32.   Torrie, G. M.; Valleau, J. P., Monte-Carlo free-energy estimates using Non-Boltzmann sampling - application to subcritical Lennard-Jones fluid. *Chemical Physics Letters* **1974,** *28* (4), 578-581.

33.   Kumar, S.; Bouzida, D.; Swendsen, R. H.; Kollman, P. A.; Rosenberg, J. M., The weighted histogram analysis method for free-energy calculations on biomolecules .1. the method. *Journal of Computational Chemistry* **1992,** *13* (8), 1011-1021.

34.   Deng, H.; Ma, Y.; Ren, W. S.; Vuong, V. Q.; Qian, P.; Guo, H., Structure and Dynamics of the Reactive State for the Histidine Methylation Process and Catalytic Mechanism of SETD3: Insights from Quantum Mechanics/Molecular Mechanics Investigation. *Acs Catalysis* **2020,** *10* (22), 13314-13322.

35.   Chu, Y. Z., Yao, J.Z., Guo, H., QM/MM MD and Free Energy Simulations of G9a-Like Protein (GLP) and Its Mutants: Understanding the Factors that Determine the Product Specificity *PLos One (accepted)* **2012**.

36.   Chu, Y. Z.; Xu, Q.; Guo, H., Understanding Energetic Origins of Product Specificity of SET8 from QM/MM Free Energy Simulations: What Causes the Stop of Methyl Addition during Histone Lysine Methylation? *Journal of Chemical Theory and Computation* **2010,** *6* (4), 1380-1389.

37.   Guo, H. B.; Guo, H., Mechanism of histone methylation catalyzed by protein lysine methyltransferase SET7/9 and origin of product specificity. *Proc Natl Acad Sci U S A* **2007,** *104* (21), 8797-802.

38.   Hu, P.; Wang, S.; Zhang, Y., How do SET-domain protein lysine methyltransferases achieve the methylation state specificity? Revisited by ab initio QM/MM molecular dynamics simulations. *Journal of the American Chemical Society* **2008,** *130* (12), 3806-3813.

39.   Qian, P.; Guo, H. B.; Wang, L. A.; Guo, H., QM/MM Investigation of Substrate and Product Specificities of Suv4-20h2:

How Does This Enzyme Generate Dimethylated H4K20 from Monomethylated Substrate? *Journal of Chemical Theory and Computation* **2017,** *13* (6), 2977-2986.

40.   Xu, Q.; Chu, Y. Z.; Guo, H. B.; Smith, J. C.; Guo, H., Energy Triplets for Writing Epigenetic Marks: Insights from QM/MM Free-Energy Simulations of Protein Lysine Methyltransferases. *Chemistry-a European Journal* **2009,** *15* (46), 12596-12599.

41.   Zhang, X.; Bruice, T. C., Enzymatic mechanism and product specificity of SET-domain protein lysine methyltransferases. *Proceedings of the National Academy of Sciences of the United States of America* **2008,** *105* (15), 5728-5732.