Title

The noncoding circular RNA *circHomer1* regulates synaptic development and

experience-dependent plasticity in mouse visual cortex

Authors

Kyle R. Jenks[1,9], Ying Cai[2,9], Marvin Eduarte Nayan[3,9], Katya Tsimring[1], Keji Li[1], José C.

Zepeda[1], Gregg R. Heller[1], Chloe Delepine[1], Jennifer Shih[1], Shiyang Yuan[2], Yao Zhu[2],

Ye Wang[4], Yangyang Duan[4], Amy K. Y. Fu[4], Taeyun Ku[5], Dae Hee Yun[1], Kwanghun

Chung[1], Nikolaos Mellios[6-7], Mriganka Sur*[1], Jacque Pak Kan Ip*[2,8]

Affiliations

[1]Department of Brain and Cognitive Sciences, The Picower Institute for Learning and

Memory, Massachusetts Institute of Technology, Cambridge, Massachusetts, 02139,

USA

[2]School of Biomedical Sciences, The Chinese University of Hong Kong, Hong Kong,

China.

[3]Department of Biology, Massachusetts Institute of Technology, Cambridge,

Massachusetts, 02139, USA

[4]Division of Life Science, The Hong Kong University of Science and Technology, Clear

Water Bay, Hong Kong, China.

[5]Graduate School of Medical Science and Engineering, Korea Advanced Institute of

Science and Technology, Daejeon, Korea.

[6]Circular Genomics Inc, Albuquerque, New Mexico, 87110, USA

[7]Previously at: University of New Mexico, Department of Neurosciences, Albuquerque,

New Mexico, 87131, USA

[8]Gerald Choa Neuroscience Institute, The Chinese University of Hong Kong, Hong

Kong, China.

[9]These authors contributed equally

*Address correspondence to Jacque Pak Kan Ip (jacqueip@cuhk.edu.hk) and Mriganka

Sur (msur@mit.edu)

Keywords: circular RNA; primary visual cortex; critical period; ocular dominance plasticity; synaptic plasticity; experience-dependent plasticity

Abstract

Circular RNAs (circRNAs) are a class of closed-loop, single stranded RNAs whose expression is particularly enriched in the brain. Despite this enrichment and evidence that the expression of circRNAs are altered by synaptic development and in response to synaptic plasticity *in vitro*, the regulation by and function of the majority of circRNAs in experience-dependent plasticity *in vivo* remain unexplored. Here, we employed transcriptome-wide analysis comparing differential expression of both mRNAs and circRNAs in juvenile mouse primary visual cortex (V1) following monocular deprivation (MD), a model of experience-dependent developmental plasticity. Among the differentially expressed mRNAs and circRNAs following 3-day MD, the circular and the activity-dependent mRNA forms of the *Homer1* gene, *circHomer1* and *Homer1a* respectively, were of interest as their expression changed in opposite directions: *circHomer1* expression increased while the expression of *Homer1a* decreased following 3-day MD. Knockdown of *circHomer1* delayed the depression of closed-eye responses normally observed after 3-day MD. *circHomer1*-knockdown also led to a reduction in average dendritic spine size prior to MD but critically there was no further reduction after 3-day MD, consistent with impaired structural plasticity. *circHomer1*-knockdown also prevented the reduction of surface AMPA receptors after 3-day MD. Synapse-localized puncta of the AMPA receptor endocytic protein Arc increased in volume after MD but were smaller in *circHomer1*-knockdown neurons, suggesting that *circHomer1* knockdown impairs experience-dependent AMPA receptor endocytosis. Thus, the expression of multiple circRNAs are regulated by experience-dependent developmental plasticity, and our findings highlight the essential role of *circHomer1* in V1 synaptic development and experience-dependent plasticity.

Significance Statement

Circular RNAs (circRNAs) are a class of closed-loop RNAs formed through back-

splicing of exon and/or intron junctions. Initially considered as byproducts of aberrant

RNA splicing with limited function, recent studies have implicated circRNAs in various

neurological disorders. Despite their abundant expression in the brain, the role of

circRNAs in experience-dependent plasticity remains poorly understood. We conducted

an *in vivo* transcriptome analysis of circRNAs whose expression was regulated by

experience-dependent plasticity in the mouse visual cortex and highlight *circHomer1*, a

circRNA derived from the *Homer1* gene, as a circRNA critical for synaptic development

and experience-dependent plasticity *in vivo*.

Introduction

Experience-dependent plasticity is a tightly regulated process that requires orchestrated gene transcription to form mRNAs that are subsequently translocated and translated into proteins to induce the functional and structural reorganization of synapses[1]. Though we now know a great deal about the mRNAs recruited by and required for experience-dependent plasticity, it remains unclear if other forms of non-coding/coding RNAs are as tightly regulated as their mRNA counterparts and if they contribute to experience-dependent plasticity[2, 3]. Recent RNA-sequencing screens have shown that a plethora of genes that produce alternatively spliced mRNAs also generate close-looped isoforms composed of backspliced and covalently-joined exons and/or introns known as circular RNAs (circRNAs)[4]. Although much is still unknown about the precise mechanisms that govern the biogenesis, regulation, and function of circRNAs; there is emerging evidence that circRNAs may have important roles in the brain[3, 5]. They are dynamically expressed during brain development, are localized to synapses[6], and a small number of circRNAs have been shown to affect neuronal gene expression, miRNA availability, and regulate behavior[7–9]. However, it remains unclear how many circRNAs might play a role in regulating experience-dependent plasticity, highlighting the need for screens to identify putative candidates.

Ocular dominance plasticity in mouse binocular primary visual cortex (V1) is a well-defined model of cortical plasticity induced by monocular deprivation (MD), typically via suture of the eye-lids, during a developmental critical period[10–15]. During this window, which in mice begins at postnatal day (P)21 and ends around P35[16], MD induces a reduction of V1 responses to the deprived eye after ~3 days of deprivation, followed by an increase of responses to the non-deprived eye[17] after ~7 days of deprivation[11, 13–15, 18–20]. MD also leads to the activity-dependent up- and down-regulation of many mRNAs known to regulate synaptic plasticity[10–15]. We have successfully used MD in the past to screen for and identify miRNAs, another unique class of RNA species, regulated by and critical for experience-dependent plasticity[2]. Thus, we sought to similarly use MD to identify circRNAs in mouse V1 that could be involved in regulating experience-dependent plasticity and synaptic function.

By screening for differentially expressed mRNAs and circRNAs in the developing mouse
V1 following 3 days of MD (3-day MD), we discovered multiple circRNAs whose
expression was regulated by experience-dependent plasticity. We further explored the
role of the circRNA *circHomer1* in experience-dependent plasticity, as the expression of
circHomer1 was opposite that of its linear, activity-dependent form *Homer1a* and
circHomer1 has previously been shown to be localized to synapses and involved in
cognitive flexibility and reversal learning(9, 21, 22)*.* Using *circHomer1*-specific
knockdown in V1, we showed that *circHomer1* delays normal closed-eye response
depression following 3-day MD. While *circHomer1* knockdown alone decreases the
average size of dendritic spines on layer 2/3 neurons, it also prevented the decrease in
spine size following 3-day MD. Knockdown of *circHomer1* also impaired downregulation
of the AMPA receptor subunit GluA1 following 3-day MD, putatively by reducing Arc
protein expression at dendritic spines. Together, these data provide the first evidence of
an experience-dependent circRNA that is required for normal synaptic development and
experience-dependent plasticity induced by short-term MD.
122
Results
circRNA expression is regulated by experience-dependent plasticity
To identify potential experience-regulated circRNAs, we performed MD in mice
beginning at P25, collected tissue from V1 and examined the transcriptomic profile after
3-day MD (Fig. 1A). We normalized RNA expression in the hemisphere contralateral to
the deprived eye, where the majority of change in synaptic drive occurs, to expression
in the ipsilateral hemisphere (within-subject) which receives significantly less input from
the deprived eye and has significantly less change in gene expression(2). Before
exploring the circRNA expression profile, we first examined the mRNA expression
profile to validate our results against previously observed mRNA regulation by MD. We
identified a total of 1173 differentially-expressed mRNAs, with more mRNAs
downregulated than upregulated (472 upregulated and 701 downregulated, Fig. 1B).
mRNAs known to be downregulated during MD, such as *Bdnf* and *Nptx2*(23), were also
downregulated in our data (Supplementary Table 1). Having confirmed that our
preparation replicated the known regulation of mRNA expression by 3-day MD(24), we

next examined the regulation of circRNAs. We used a published circRNA alignment
tool, circtools, to detect circRNA transcripts by virtue of their unique out-of-order
junctional reads(23). A total of 1489 unique circRNAs were detected, with 73
differentially-expressed (Fig. 1C, Supplementary Table 2). Of the differentially-
expressed circRNAs, we observed 27 significantly upregulated circRNAs and 46
downregulated circRNAs. Some of the genes from which these differentially-expressed
circRNAs were derived have known roles in synaptic plasticity and function
(Supplementary Fig. 1A). Indeed, we also identified changes in the corresponding
mRNA isoforms of several differentially expressed circRNAs. Of these, the linear
isoforms of *circHomer1*, *circPrkce*, and *circDcun1d4* were of interest as they exhibited
changes in the opposite direction compared to their circular counterparts, suggesting
their circRNA expression was regulated by experience independently of the overall
gene transcription rate (Fig. 1D). We also performed a transcriptomic analysis of
mRNAs and circRNAs in V1 after 7-day MD (Supplementary Fig. 1B-E and
Supplementary Tables 3 and 4). *circPrkce*, *circDcun1d4*, and their linear isoforms were
not differentially expressed at this timepoint, while linear *Homer1a* remained
downregulated and *circHomer1* was no longer upregulated. Based on these findings,
and the known role of *Homer1a* in synaptic plasticity, we chose to more closely examine
the experience-dependent and developmental regulation and role of *circHomer1* in
mouse V1.
158

circHomer1 **expression is developmentally regulated**
circHomer1 is composed of exons 2 through 5 of the *Homer1* gene, with backsplicing
between exon 2 and 5 (Fig. 2A). Using primers targeting the backsplice junction of
circHomer1, we next sought to validate the observed changes in contralateral V1
circHomer1 expression, normalized to ipsilateral V1, after 3-day and 7d-day MD using
RT-qPCR (Fig. 2B). In addition, we examined several mRNAs known to be regulated by
MD. As expected after 3-day MD, *Bdnf* and *Nptx2* mRNA were downregulated in
contralateral V1(24). *Homer1a* mRNA was also significantly downregulated after 3-day
MD, in contrast to *circHomer1*, validating the dissociation between *Homer1a* and
circHomer1 expression after 3-day MD (Fig. 2B, left). After 7-day MD, *Stat1*(25), a

regulator of homeostatic synaptic plasticity, was upregulated in contralateral V1 (Fig. 2B,

right) as previously shown for prolonged MD(17, 24, 25). We also saw that the

expression of *circHomer1* was significantly decreased after 7-day MD along with

Homer1a (Fig. 2B, right). These results validate the RNAseq data showing that

experience-dependent regulation of *circHomer1* expression is distinct at 3-day MD from

that of its linear experience-dependent counterpart *Homer1a*.

Expression of many mRNAs required for normal, developmental experience-dependent

plasticity peak during the critical period. To determine the developmental expression

profiles of *Homer1a* and *circHomer1* over the mouse's lifespan, we measured their

expression in V1 at various time points using RT-qPCR. *circHomer1* expression

increased significantly starting at ~P24, aligning to the start of the critical period (Fig.

2C, left), while *Homer1a* expression did not increase significantly until P60, past the

close of the critical period (Fig. 2C, middle). *circHomer1* expression appeared to plateau

at P60 and remained at similar levels of expression in mice as old as 1.5 years.

Expression of another circRNA, *circTulp4*, did not change across development or

adulthood indicating that the developmental expression profile of *circHomer1* is not a

shared feature of all circRNAs (Fig. 2C, right).

circHomer1-depletion delays ocular dominance plasticity

The regulation of *circHomer1* expression by MD and the peak in *circHomer1* expression

at the start of the critical period mirrors that of linear mRNAs with known roles in ocular

dominance plasticity. To determine if *circHomer1* is required for ocular dominance

plasticity following MD, we used a short hairpin RNA (shRNA)-mediated depletion

strategy targeting the backsplice junction of *circHomer1*, which has been previously

validated *in vivo* to specifically downregulate *circHomer1*, and not linear *Homer1*

mRNAs (including *Homer1a*, *Homer1b*, and *Homer1c*)(9), in the cortex. We re-

confirmed the specificity of knockdown using adeno-associated viruses (AAV)-mediated

transduction of the shRNA in mouse V1, which resulted in a significant reduction of

circHomer1 RNA with no changes in *Homer1a*, *Homer1b*, or *Homer1c* mRNA

expression (Supplementary Fig. 2A). sh-circHomer1 also had no effect on total Homer1

protein levels (Supplementary Fig. 2B).

To examine the effects of *circHomer1* depletion on ocular dominance plasticity, we
quantified eye-specific responses in the binocular zone of V1 by using optical imaging of
intrinsic hemodynamic signals measured before and after 3-day or 7-day MD (Fig. 3A).
We then compared results between mice injected with a sh-scramble (control) or sh-
circHomer1 virus (Fig. 3B). To compare the relative visual drive elicited by the two eyes
in V1, the normalized difference between contralateral (dominant-eye) and ipsilateral
responses during optical imaging was measured to determine the ocular dominance
index (ODI). Change in ODI following closure of the contralateral eye (MD) was then
used as a measure of experience-dependent plasticity(26).

Prior to MD (preMD), there was no difference in baseline ODI between sh-scramble and
sh-circHomer1 mice, with responses biased towards the dominant, contralateral eye
(Fig. 3C). After 3-day MD, the ODI of sh-scramble mice decreased significantly (shifted
towards the open, ipsilateral eye) as expected(24, 27, 28). In contrast, the ODI of sh-
circHomer1 mice were unchanged after 3-day MD (Fig. 3C). After 7-day MD, however,
the ODI of sh-circHomer1 mice were significantly decreased and no longer significantly
different than that of sh-scramble mice (Fig. 3C).

The ODI shifts at 3-day and 7-day MD are driven by an initial decrease in contralateral
drive and delayed increase in ipsilateral drive, respectively(24, 27, 28). To see if both
these eye-specific changes instead occurred between 3-day and 7-day MD in sh-
circHomer1 mice, we examined the responses driven by the contralateral and ipsilateral
eyes separately. Unlike in the sh-scramble mice, and as expected from the lack of
change in ODI, contralateral closed-eye responses of sh-circHomer1 mice did not
decrease after 3-day MD. Similarly, ipsilateral open-eye responses were already
significantly increased in sh-scramble mice after 3-day MD whereas they were
paradoxically reduced in sh-circHomer1 mice (Fig. 3D). Despite the differences in eye-
specific changes after 3-day MD, however, by 7-day MD both contralateral and
ipsilateral responses in the sh-circHomer1 mice were comparable to the sh-scramble
mice (Fig. 3D). Thus, *circHomer1* depletion does not grossly impair the eye-specific
development of contralateral and ipsilateral visual drive and delays, but does not block,
the expression of binocular experience dependent plasticity following MD.

circHomer1 regulates dendritic spine morphology

MD induces rapid remodeling of the dendritic spines of mouse layer 2/3 V1 neurons,
with 3-day MD significantly decreasing average spine volumes(10). As we had found
that depletion of *circHomer1* blocked the depression of closed eye responses at 3-day
MD (Fig. 3), we hypothesized that depletion of *circHomer1* would likewise impair 3-day
MD-induced spine shrinkage. We injected an AAV expressing RFP and either sh-
circHomer1 or sh-scramble shRNA into the binocular region of V1 at P15, prior to the
start of the critical period. We also injected AAV9-hSyn-DIO-EGFP (structural marker)
and AAV9-CaMKIIa-Cre virus in order to sparsely label neurons and quantified spine
morphology of labeled layer 2/3 V1 neurons from mice that had undergone no MD (No
MD), 3-day MD, or 7-day MD. To achieve a more detailed analysis of the dendritic spine
structure than is possible with traditional confocal imaging, we used epitope-preserving
magnified analysis of the proteome (eMAP) to expand our tissue samples (Fig. 4A). The
apical dendrites of V1 layer 2/3 pyramidal neurons expressing both GFP and RFP were
selected for further analysis (Fig. 4B, left). Based on their morphology, dendritic spines
were classified into four categories: mushroom, stubby, thin, and filopodia using
standard criteria (Fig. 4B-C)(29).

In No MD mice, sh-circHomer1 led to a significant drop in the density and percentage of
mature mushroom spines(30), but not immature spines (stubby, thin, and filopodia)(30)
(Fig. 4D-E). This basal effect was surprising given the normal baseline ODI and eye-
specific responses we had observed in preMD sh-circHomer1 mice. After 3-day MD in
sh-scramble mice, as expected, there was a significant decrease in the density and
percentage of mature mushroom spines when compared to the No MD condition.
Conversely, the density of both mature and immature spines remained unchanged after
3-day MD in the sh-circHomer1 mice. After 7-day MD, in the sh-scramble mice the
density and percentage of mature spines remained lower compared to the No MD
condition, while the density of immature spines showed a significant increase. Mature
and immature spine density in sh-circHomer1 mice after 7-day MD again remained
unchanged compared to the No MD condition, but was no longer significantly different
than the sh-scramble mice (Fig. 4D-E).

As categorical classification could obscure more subtle changes in the dendritic spines
of sh-circHomer1 neurons, we also analyzed the average spine volume by performing
3D reconstruction. *circHomer1* depletion led to a more than 50% decrease in average
spine volume in the No MD condition (Fig. 4F, Supplementary Fig. 3A). After 3-day MD,
the average spine volume decreased by more than 70% in sh-scramble mice compared
to the No MD condition. In contrast, 3-day MD did not significantly alter spine volume in
sh-circHomer1 mice (Fig. 4F-G, Supplementary Fig. 3B). After 7-day MD, spine volume
was still significantly decreased in sh-scramble mice compared to the No MD condition,
and while the sh-circHomer1 spine volume remained unchanged, the sh-scramble and
sh-circHomer1 spine volumes were not significantly different at 7-day MD (Fig. 4F,
Supplementary Fig. 3C). Taken together, our findings suggest that *circHomer1* plays a
crucial role in regulating the structural morphology of dendritic spines both during
normal development and following short-term MD.
Considering the effects *circHomer1* depletion had on the morphology of synapses in the
No MD condition, we sought to determine how *circHomer1* depletion affected the
electrophysiological properties of neurons. We used whole-cell patch clamp recordings
in acute slices from P28-33 mice to measure passive membrane and synaptic
properties of layer 2/3 neurons infected with either sh-circHomer1 or sh-scramble virus.
We did not find any difference in baseline neuronal excitability (Supplementary Fig. 4) or
the amplitude of miniature excitatory postsynaptic currents (mEPSC; Supplementary Fig.
5C), but there was a decrease in mEPSC frequency (Supplementary Fig. 5D),
consistent with a decrease in protrusion density(31). While the basal structural and
electrophysiological deficits following *circHomer1* depletion were surprising given the
apparently normal binocular visual drive observed prior to MD, experience-dependent
plasticity following MD was delayed as opposed to simply occluded by *circHomer1*
depletion. We therefore sought to further investigate the mechanism by which
circHomer1 depletion impaired experience-dependent plasticity.
288
***circHomer1* controls surface AMPA receptor expression**
The major mechanism underlying both reduced closed-eye drive and spine shrinkage
following MD is a reduction in surface AMPA receptors(25–27). Therefore, we

investigated whether *circHomer1* depletion disrupted surface AMPA receptor trafficking

following MD using a biotinylation assay of the AMPA receptor subunit, GluA1 (Fig. 5A).

As expected, 3-day MD led to a significant reduction of surface GluA1 compared to No

MD in sh-scramble mice. There was no reduction of GluA1 levels after 3-day MD in sh-

circHomer1 mice (Fig 5A). By 7-day MD, however, there was no significant difference in

surface GluA1 expression between sh-scramble and sh-circHomer1 mice. Based on

these results, we hypothesized that *circHomer1* was necessary for normal activity-

dependent AMPA receptor endocytosis. To examine this further, we treated primary

neurons *in vitro* with bicuculline, a $GABA_A R$ antagonist, to induce activity-dependent

AMPA receptor internalization(32). 1- or 2-days of treatment with bicuculline induced

upregulation of *circHomer1* in neurons (Supplementary Fig. 6A), and *circHomer1*-

depletion impaired bicuculline induced endocytosis of surface AMPA receptors

visualized via SEP-GluA1 expression (Supplementary Fig. 6B,C). These data

demonstrate that *circHomer1* depletion impairs experience- and activity-dependent

changes in surface GluA1 expression levels, thus indicating a potential functional

mechanism for the disrupted ocular dominance plasticity after 3-day MD.

How might a circRNA contribute to AMPA receptor endocytosis? One possibility, given

the previous observation of dendritically localized *circHomer1*(33) and binding to linear

Homer1b mRNA(21), is through regulating other dendritically localized mRNAs critical to

AMPA receptor endocytosis(22). The activity-regulated cytoskeleton-associated protein

(Arc) is an immediate early gene product that has been shown to play a critical role in

AMPA receptor endocytosis(32, 34), and *Arc* mRNA is trafficked to dendrites and

translated locally in response to synaptic activity. We therefore tested whether Arc

protein expression at dendritic spines is affected by *circHomer1* depletion following MD.

We again injected AAV expressing RFP and either sh-scramble or sh-circHomer1, and

sparsely labeled neurons in V1 with GFP as described in Figure 4. After processing the

tissue with eMAP, postsynaptic Arc protein was labeled by immunostaining, and

pyramidal neurons in layer 2/3 of V1 expressing both RFP and GFP were selected for

analysis. Next, we performed 3D reconstruction of the imaged dendrites (Fig. 5B), and

quantified the volume of Arc puncta expressed at dendritic spines (Fig. 5C). Intriguingly,

the percentage of Arc-containing spines and the volume of Arc puncta increased

significantly after 3- and 7-day MD in both sh-scramble and sh-circHomer1 mice as
compared to the No MD conditions (Fig. 5D,E). However, the mean volume of Arc
puncta and the Arc volume/spine volume ratio in sh-circHomer1 neurons was
significantly lower than that in sh-scramble neurons after 3-day MD (Fig. 5E,F). There
was no difference in the mean volume of Arc puncta and the Arc volume/spine volume
ratio between sh-scramble and sh-circHomer1 neurons at 7-day MD. This is consistent
with the impairment of surface GluA1 removal and spine shrinkage after 3-day MD in
sh-circHomer1 mice as compared to control (Figs. 4, 5A). Together, these results
demonstrate that *circHomer1*-depletion impairs GluA1 internalization following 3-day
MD, which may be explained by *circHomer1*-depletion reducing the quantity of Arc
protein accumulating at dendritic spines in response to MD.

Discussion

circRNAs are abundantly expressed in mammalian brains, are preferentially derived
from genes encoding synaptic proteins(33, 35), and several circRNA have recently
described roles in synaptic plasticity or function(7–9, 21, 22). Based on these findings,
we performed an unbiased screen to identify candidate, novel experience-dependent
plasticity regulated circRNAs using *in vivo* linear- and circular-transcriptome analyses.
We identified several promising circRNAs with differential expression in contralateral
binocular V1 following 3-day MD in critical period age mice. We focused on *circHomer1*,
a well-known circRNA derived from the *Homer1* gene which encodes important synaptic
scaffolding proteins and an activity dependent linear isoform, *Homer1a*, which is a key
regulator of glutamatergic synaptic signaling(36).
circHomer1 and *Homer1a* were differentially regulated by 3-day MD, but in opposing
directions with the expression level of *circHomer1* increasing and that of *Homer1a*
decreasing following short-term MD. Following longer term, 7-day MD, however,
expression levels of both *circHomer1* and *Homer1a* were significantly decreased (as
measured via RT-qPCR). The developmental expression of *circHomer1* increased
rapidly in V1 between P15 and P28, wherein the critical period for ocular dominance
plasticity begins, which in combination with its differential expression following 3-day
MD suggested a role for *circHomer1* in critical period plasticity. Indeed, loss of

circHomer1 disrupted ocular dominance plasticity following 3-day MD without effecting

baseline visual responses. However, depletion of *circHomer1* alone also led to less

mature dendritic spines and decreased mEPSC frequency. The normal, albeit delayed

expression of ocular dominance plasticity following 7-day MD, however, suggests that

the 3-day MD phenotype is not simply a result of occlusion. Indeed, while *circHomer1*

depletion impaired 3-day MD induced spine shrinkage, AMPA receptor endocytosis, and

accumulation of synaptic Arc protein these effects were not apparent when compared to

their respective controls at 7-day MD.

The various proteins derived from the *Homer1* gene play critical roles in regulating

synapse development, synaptic strength, and homeostatic synaptic scaling(37–41). At

the postsynaptic density, the activity-dependent short protein isoform of the *Homer1*

gene, Homer1a, has dominant-negative effects on the constitutive longer forms of

Homer1 (Homer1b/c), and through this interaction regulates the clustering of

mGluRs(38–40) and mGluR-dependent NMDA and AMPA receptor currents(38, 42–45).

circHomer1 appears to have a similar dominant negative effect as its linear, activity-

dependent counterpart by instead binding directly to dendritic *Homer1b* mRNA which is

necessary in the orbitofrontal cortex for normal reversal learning(21). Despite the fact

that Homer1a and *circHomer1* both share activity dependent regulation and antagonize

Homer1b expression at the synapse, their regulation by and role in developmental

experience-dependent V1 plasticity are clearly distinct. In contrast to the upregulation of

circHomer1, *Homer1a* is reduced following 3-day MD (Fig. 2C), and depletion of

Homer1a does not affect the shift in ODI and reduction of contralateral, closed-eye

responses as a result of 3-day MD(46). Homer1a depletion instead alters the

establishment of contralateral bias of V1 responses under basal conditions, which is not

affected by *circHomer1* depletion (Fig. 3C,D). Thus, although both Homer1a and

circHomer1 are regulators of cortical plasticity, they serve different roles in V1

development and experience-dependent plasticity.

It remains mechanistically unclear how neuronal activity regulates *circHomer1*

expression, as the only demonstrated precursor of *circHomer1* is non-activity dependent

Homer1b pre-mRNA(47). Given that activity-dependent *Homer1a* pre-mRNA contains

the relevant regions to form *circHomer1*, including a section of intron 5(47), it is

plausible that *circHomer1* could derive from *Homer1a* pre-mRNA, thereby allowing its

expression to change in tandem with *Homer1a* mRNA even assuming a fixed ratio of

linear/back-splicing, and thus circularization rate. However, during MD and development

in vivo and bicuculline treatment *in vitro*, we observed timepoints where changes in

circHomer1 expression appear decoupled from or even opposing changes in *Homer1a*.

Circular RNA species decay more slowly than their linear counterparts(48) and changes

in transcription thus tend to result in larger, longer relative increases in circRNA

levels(49), but this alone seems insufficient to explain this discrepancy. More plausible,

perhaps, is that the circularization rate is itself regulated by activity(47).

Thus, we asked whether a constant circularization rate of *Homer1b* and *Homer1a* pre-

mRNA, a variable circularization rate of *Homer1b* pre-mRNA*,* or a variable

circularization rate of both *Homer1b* and *Homer1a* pre-mRNA best fit our observed data

(Supplementary Fig. 7). Our models shows that across multiple paradigms, a constant

circularization rate of *Homer1a* and *1b* pre-mRNA or a variable circularization rate of

Homer1b pre-mRNA alone is insufficient to explain the observed expression levels of

circHomer1. Instead, our data are best fit when adding a variable, activity-dependent

circularization rate of both *Homer1b* and *Homer1a* pre-mRNA. In this way, *circHomer1*

levels can increase when *Homer1a* mRNA levels do not change or even decrease.

Importantly, the model accounts for the bidirectional change in *circHomer1* after 3-day

versus 7-day MD. (Fig. 2C-D, Supplementary Fig. 7A). Such regulation could be

facilitated by recently described molecular pathways underlying *circHomer1*

biogenesis(47) or by changes in other *Homer1* mRNA products not measured in our

experiments (for instance, *Homer1b*(21)). Future work is needed to determine if

Homer1a pre-mRNA can indeed give rise to *circHomer1*, whether circularization of pre-

mRNA occurs at a sufficient frequency to meaningfully decreases linear mRNA

availability, and if circular RNA decay rate is static or itself regulated by activity as is the

case for some mRNA species(7, 8, 50, 51).

An important conclusion of our study is that *circHomer1* has a critical role in the

reduction of synaptic drive that accompanies the initiation of MD. Specifically, we found

that *circHomer1* knockdown alters the time course of surface GluA1 trafficking and Arc

protein expression during experience-dependent plasticity. However, the exact

mechanism of *circHomer1* in synaptic development and plasticity warrants further
investigation. Previous studies suggest that *circHomer1* regulates alternative splicing of
mRNA isoforms(9) as well as synaptic expression of *Homer1b* mRNA in the orbitofrontal
cortex via direction competition for HuD binding(21). Overall expression of several
postsynaptic genes was also observed to be downregulated following *circHomer1*
knockdown(22). It remains to be elucidated whether these mechanisms also contribute
to the function of *circHomer1* in experience-dependent plasticity and the normal
synaptic localization of Arc protein following MD.

The mechanism(s) of circRNA function are an area of active investigation. circRNAs are
enriched in neurons and are believed to play critical roles in regulating neuronal
development and synaptic plasticity(7–9, 33, 52). From within the top 100 candidates
identified in our MD RNAseq screen, several circRNAs were already reported in the
literature to have biological functions in other contexts(53–55). These include the
neuronally-enriched circRNA *Cdr1as* (also known as ciRS-7), which has been found to
contain a large number of binding sites for miR-7, a regulator of neurodevelopment(6,
56–59). This enrichment for miR-7 binding sites effectively serves as a "miRNA
sponge", and *Cdr1as* knockout mice exhibit behavioral deficits and altered neuronal
electrophysiological properties(6, 8).

This study identified tens of circRNAs differentially expressed following experience-
dependent plasticity and highlights *circHomer1*, a neuron-enriched circRNA derived
from the Homer1 gene, as a critical regulator of V1 plasticity during the critical period.
Our findings advance the understanding of circRNA regulation during experience-
dependent plasticity and shed light on their functional significance in developmental
processes.

Acknowledgments

We thank Jamie Benoit for his contribution to the initial RNA-seq experiment. We thank Taylor Johns, Austin Sullins and other members of the Sur lab and the Ip lab for their help and support. We thank Hovy Wong for comments on the manuscript. We thank Lorena Rubino for her assistance in analyzing RNA-seq data. We thank Cara Kwong for her technical assistance in neuronal cultures. We thank the MIT BioMicroCenter for running the RNA-seq library preparation and sequencing. We thank scidraw.io for the schematic, licensed under a Creative Commons 4.0 license (https://creativecommons.org /licenses /by/4.0/). This work was supported by NIH grants R01MH085802 (M.S.), R01EY028219 (M.S.), F31EY033649 (K.T.), F32EY032756 (K.J.), the Simons Foundation Autism Research Initiative through the Simons Center for the Social Brain, MIT (M.S.), the National Research Foundation of Korea (RS-2023-00264980; T.K.), the Hong Kong Research Grants Council Early Career Scheme (24117220; J.I), General Research Fund (14117221; J.I.), Area of Excellence Scheme (AoE/M-604/16; J.I., A.F.), Theme-based Research Scheme (T13-605/18-W; J.I., A.F.), Lo Kwee-Seong Biomedical Research Fund (J.I.), Faculty Innovation Awards from the Faculty of Medicine CUHK (FIA2020/A/04; J.I.), and NARSAD Young Investigator Grant from the Brain & Behavior Research Foundation (J.I.).

Author Contributions

J.I., N.M., M.N., M.S., K.J. conceived experiments with input from others. M.N., C.D. and S.Y. performed the cortical dissections and RNA and protein purifications. C.D., M.N and Y.Z. analyzed RNA-seq data. M.N., Y.C. and C.D. performed qPCR experiments. K.J., K.T., J.Z. and J.I. performed surgeries and viral injection, and carried out *in vivo* optical imaging experiments. K.L. analyzed optical imaging data. Y.C., S. Y. and J.Z. performed spine morphology analysis. Y.C., S.Y., T. K., D. Y. and J.I. performed eMAP experiments with inputs from K.C. J.I. performed the surface biotinylation assay and protein analyses. J.S. performed electrophysiological recordings and analyses. Y.C., Y.W., Y.D. performed neuronal culture experiments and relevant analyses. A.F. provided expertise on neuronal culture experiments. N.M. shared the sh-RNA constructs. G.H. performed the modeling of RNA expression. K.J., J.I., and M.S.

contributed to analysis of experiments and interpretation of results. J.I., K.J. and M.S.

wrote the manuscript with input from others. All authors edited the manuscript.

Competing Interest Statement

N.M. is CSO of Circular Genomics Inc, Albuquerque, NM.

Data, Materials, and Software Availability

All RNA sequencing data have been deposited in GEO. The code will be made

available upon reasonable request.

Figure Legends

Figure 1. Experience-dependent circRNAs in V1 identified through MD.

(A) Timeline and schematic of 3-day MD RNAseq screen. (B) Volcano plot of mRNA sequencing of V1 after 3-day MD (n = 3 mice per group). (C) Volcano plot of circRNA sequencing of V1 after 3-day MD (n = 3 mice per group). The ipsilateral hemisphere (ipsi) served as a control for MD induced changes in the contralateral hemisphere (contra) for both mRNA and circRNA expression. (D) Fold change of differentially-expressed circRNAs and their mRNA isoforms after 3-day MD. The mouse and brain silhouette were adapted from https://doi.org/10.5281/zenodo.8044766 and https://doi.org/10.5281/zenodo.3925971.

Figure 2. *circHomer1* expression is regulated by experience-dependent plasticity and upregulated during the ocular dominance critical period.

(A) The mouse Homer1 gene encodes long and short versions of the synaptic protein Homer1 and a noncoding circRNA, *circHomer1*. (B) RT-qPCR quantification of selected mRNAs and *circHomer1* after 3-day or 7-day MD (n = 3-4 mice per group, paired t-test). (C) *circHomer1* (left), *Homer1a* (middle), and *circTulp4* (right) expression measured by RT-qPCR at several timepoints in mouse V1 across development and into adulthood (n = 3-6 mice per group, one-way ANOVA following Tukey multiple comparisons). Data are presented as mean ± SEM. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 3. *circHomer1* depletion delays the expression of ocular dominance plasticity in V1 following MD.

(A) Mice were injected with either sh-circHomer1 or sh-scramble virus at P15, a craniotomy was performed at P22 to implant a cranial window, then a pre-MD optical imaging session was done and the contralateral eye-lid sutured at P25. After either 3-day or 7-day MD, the eyelid was reopened and post-MD imaging done. (B) Example retinotopic maps from a sh-circHomer1 mouse, obtained with optical imaging. The color corresponds with different phases of the visual stimulus (consistent with the visual field map) and brightness shows the amplitude of cortical response. Top, contralateral closed-eye responses, pre-MD (left) and after 7-day MD (right). Bottom, ipsilateral open-

eye responses. Scale bar = 500 µm. (C) Ocular dominance index (ODI) changes after
3-day or 7-day MD for mice injected with sh-scramble or sh-circHomer1 virus. Each dot
shows the average ODI for one animal, and grey lines show the change of ODI for one
animal (n = 5-11 mice per group, mixed-effects model following Holm-Sidak multiple
comparisons test). (D) Normalized response amplitudes driven by the contralateral (left)
and ipsilateral (right) eyes (n = 5-11 mice per group, mixed-effects model following
Holm-Sidak multiple comparisons test). Data are presented as mean ± SEM. *p < 0.05;
**p < 0.01; ***p < 0.001.

Figure 4. Depletion of *circHomer1* affects spine morphology of V1 neurons.
(A) A schematic of tissue preparation and eMAP. (B) Neurons expressing both GFP and
RFP (denoting shRNA) were selected for analysis (left). Dendritic spines were classified
into four categories based on their morphology (right). Scale bars = 100 µm (left,
estimated to be 33.33 µm prior to 3x expansion), and 10 µm (right, estimated to be 3.33
µm prior to 3x expansion). H, head width; N, neck width; L, length; $\overline{\text{H}}$, average head
width in No MD sh-scramble group. (C) Representative images showing apical
dendrites of layer 2/3 neurons in V1 in the No MD, 3-day MD, and 7-day MD condition
from the sh-scramble (top) or sh-circHomer1 (bottom) group. Arrows indicate different
types of dendritic spines. Scale bar = 10 µm (estimated to be 3.33 µm prior to 3x
expansion). (D) Density of mushroom spines, and immature spines (stubby spines, thin
spines and filopodia) on apical dendrites of layer 2/3 neurons in V1 in the No MD, 3-day
MD, and 7-day MD condition from the sh-scramble (grey) or sh-circHomer1 (pink) group
(n = 8-10 dendrites from 3 mice per group, values were normalized to 3x expansion
factor, two-way ANOVA following Tukey multiple comparisons). (E) Spine morphology
types as a percentage of total spines. The percentage of mushroom spines are labeled
on the bar. (F) Volume of dendritic spines on apical dendrites (same data as in D and E,
values were normalized to 3^3 expansion factor, two-way ANOVA following Tukey
multiple comparisons). (G) Distribution of spine volume in sh-scramble and sh-
circHomer1 mice in the 3 conditions (n = 235-334 dendritic spines from 3 mice per
group). Data are presented as mean ± SEM. *p < 0.05; **p < 0.01; ***p < 0.001.

Figure 5. Trafficking of surface AMPA receptors following MD is regulated by
circHomer1.
(A) Left: Western blot of cell surface expression of GluA1 in V1 of mice infected with
lentivirus after MD. Scr, sh-scramble; KD, sh-circHomer1. Right: Quantification of
surface GluA1 level (n = 4 mice per group, two-way ANOVA following Tukey multiple
comparisons). (B) Representative images showing 3D reconstruction of a segment of
dendrites staining with GFP and Arc after eMAP processing. Scale bar = 5 μm
(estimated to be 1.67 μm prior to 3x expansion). (C) Representative images of 3D-
reconstructed Arc puncta at dendritic spines of V1 neurons after MD. Scale bar = 1 μm
(estimated to be 0.33 μm prior to 3x expansion). (D) Percentage of dendritic spines
containing Arc puncta (n = 8-11 dendrites from 3 mice per group, two-way ANOVA
following Tukey multiple comparisons). (E) Arc puncta volume at dendritic spines (n =
29-71 puncta from 8-11 dendrites from 3 mice per group, values were normalized to 3^3
expansion factor, two-way ANOVA following Tukey multiple comparisons). (F) The ratio
of Arc puncta volume and corresponding dendritic spine volume (n = 20-51 dendritic
spines from 8-11 dendrites from 3 mice per group, two-way ANOVA following Tukey
multiple comparisons). Data are presented as mean ± SEM. *$p < 0.05$; **$p < 0.01$; ***$p <$
0.001.

Supplementary Figure 1. Transcriptomic analyses after MD in mouse V1.
(A) GO enrichment of the parental genes of differentially-expressed circRNAs after 3-
day MD. DEG, differentially-expressed gene. (B) Volcano plot of mRNA sequencing of
V1 after 7-day MD (n = 4 mice per group). (C) Volcano plot of circRNA sequencing of
V1 after 7-day MD (n = 4 mice per group). (D) GO enrichment of the parental genes of
differentially-expressed circRNAs after 7-day MD. (E) Fold change of differentially-
expressed circRNAs and their linear isoforms after 7-day MD.

Supplementary Figure 2. sh-circHomer1 specifically targets *circHomer1* without
affecting linear *Homer1* mRNA.
(A) sh-circHomer1 specifically reduces expression of *circHomer1* but not linear *Homer1*
in mouse V1 (n = 4 mice per group, Student's t-test). (B) sh-circHomer1 has no effect on

total Homer1 protein levels as measured by Western blot (n = 3 mice per group). Data

are presented as mean ± SEM. *p < 0.05.

Supplementary Figure 3. Distribution of spine volume on apical dendrites of layer 2/3

pyramidal neurons in V1 during MD.

Distribution of spine volume in (A) No MD, (B) 3-day MD, and (C) 7-day MD conditions

(n = 235-334 dendritic spines from 3 mice per group, values were corrected assuming a

3^3 expansion factor).

Supplementary Figure 4. Characterization of the effects of *circHomer1* depletion on

membrane properties in layer 2/3 neurons of binocular V1.

(A) Representative traces show the response of sh-scramble or sh-circHomer1 infected

neurons to -100, 0, and 300 pA current steps. The duration of each current step was

250 ms. (B) Quantification of the number of action potential spikes per current injection

step shows the input-output relationship in sh-scramble and sh-circHomer1 infected

neurons. (C) Rheobase, defined as the minimum amount of current injected to elicit

action potential firing, was measured in response to the current injection protocol (n=7-

10 neurons per group, Student's t-test). (D-F) Passive membrane properties were

measured in sh-scramble and sh-circHomer1 infected neurons (n=7-10 neurons per

group, Student's t-test). Data are presented as mean ± SEM.

Supplementary Figure 5. Characterization of the effects of *circHomer1* depletion on

synaptic properties in layer 2/3 neurons of binocular V1.

(A) A representative image of the injection site shows an area of lentivirus-infected

neurons in red, and in green three biocytin-filled cells that were recorded. The dotted

line along the pial surface indicates the outline of the imaged slice. Scale bar = 100 µm.

(B) Left, representative traces recorded from neurons in layer 2/3 of V1 expressing sh-

scrambled (top) or sh-circHomer1 (bottom) lentivirus. Right, all events detected were

scaled and averaged to show the average mEPSC kinetics. (C-D) Quantification of

mEPSC amplitude (C) and frequency (D) in sh-scramble or sh-circHomer1 infected

neurons (n=10 neurons per group, two-sample Kolmogorov-Smirnoff test). Data are

presented as mean ± SEM. **$p < 0.01$.

Supplementary Figure 6. Alterations of surface AMPA receptor expression after

circHomer1 depletion.

(A) Expression of *circHomer1* (left), *c-fos* (middle), and *Homer1a* (right) RNAs following

bicuculline treatment in cultured cortical neurons (n=4-5 per group, one-way ANOVA

following Tukey multiple comparisons). (B) Representative images of cultured

hippocampal neurons' dendrites co-transfected with SEP-GluA1 (sGluA1) together with

sh-scramble (top) or sh-circHomer1 (bottom) at DIV 11. Neurons were treated with

bicuculline (20 μM) for 48 hours at DIV 17, and fixed at DIV 19. Con, control treatment;

Bic, bicuculline treatment. Scale bar = 5 μm. (C) Percentage of sGluA1-positive spines

under different conditions. For each neuron, 3 segments of dendrite with length of 30

μm were randomly selected, and the average percentage were calculated (n=12

neurons per group, two-way ANOVA following Tukey multiple comparisons). *:

compared to sh-scramble: Con; †: compared to compared to sh-circHomer1: Con. Data

are presented as mean ± SEM. *$p < 0.05$; **, ††$p < 0.01$; ***$p < 0.001$.

Supplementary Figure 7. Modeling of *circHomer1* expression.

(A) Schematic of the model components for each experiment. Arrows represent rates fit

in each model of either transcription, circularization, or degradation. These values were

fit for each model and experiment combination separately to minimize squared error

from measured *Homer1a* and *circHomer1* levels. (B) Simulated levels of *circHomer1*

RNA for the constant circularization (pink), variable circularization (red), and full models

(dark red) for monocular deprivation (left), development (middle), and bicuculline (right)

experiments. The measured, normalized levels of *circHomer1* in each experiment are

overlaid (black dots). (C) Normalized Akaike information criterion (AIC) scores for each

model tested. (D) Expanded results from the full models, showing the simulated levels

of both *circHomer1* and *Homer1a* RNA (black and purple lines, respectively) overlaid

with the experimentally measured levels of *circHomer1* and *Homer1a* RNA (black and

purple dots, respectively) for each experiment.

Supplementary Table 1: RNAseq results of mRNAs detected from 3-day MD experiment.

Supplementary Table 2: RNAseq results of circRNAs detected from 3-day MD experiment.

Supplementary Table 3: RNAseq results of mRNAs detected from 7-day MD experiment.

Supplementary Table 4: RNAseq results of circRNAs detected from 7-day MD experiment.

Supplementary Table 5: Primers list.

Materials and Methods

Animals

Experiments were carried out in mice under protocols conforming to NIH guidelines and approved by MIT's Animal Care and Use Committee or by CUHK's Animal Experimentation Ethics Committee. Wildtype C57BL/6J mice (JAX000664) were sourced from Jackson Laboratory or CUHK's Laboratory Animal Services Centre. Mice were group-housed whenever possible with up to 5 same-sex mice per cage. The cages were in a standard animal facility room with a 12-hour/12-hour light/dark cycle. Food and water were available ad libitum. Both male and female mice were used for experiments.

Eyelid suture and monocular deprivation

For monocular deprivation experiments, mice were anesthetized using 2-3% isoflurane. The top and bottom eyelid of the right eye was trimmed, and sterile nylon sutures (7-0) were used to suture the eyelid, which were further sealed with Vetbond (3M). The

suture was inspected for the next 3-7 days post-closure to ensure that the eye did not

reopen. Mice that exhibited incomplete eyelid closure were removed from the

experiment.

circHomer1 depletion via shRNA and validation

For *circHomer1* depletion experiments, the following viruses were used: lentivirus

expressing pLV-mU6-scrambled-shRNA::SYN-tdTomato or pLV-mU6-*circHomer1*a-

shRNA::SYN-tdTomato (1.2×10^9 IFU/mL, System Biosciences) , or AAV9 expressing

pAV-U6-sh-scramble-CMV-DsRed or pAV-U6-sh-circHomer1-CMV-DsRed (2×10^{13}

vg/mL, Vigene). For the *in vivo* shRNA experiments, binocular V1 in the left hemisphere

of P15 pups was targeted. Briefly, sh-scrambled or sh-circHomer1 viruses were loaded

into a pulled-glass micropipette with a beveled tip, lowered into L2/3, and virus was

infused at a rate of 100 nL/min. The following coordinates were used (in mm from

lambda at P15): AP: 0.5 to 1.0; ML: -2.8, DV :-0.2 to -0.25. A minimum of 10 days was

allowed for viral transduction and sufficient expression of the constructs in all

experiments. To estimate the efficiency and specificity of sh-circHomer1 after virus

injection, the mouse brain was examined using a dual fluorescent protein flashlight

(Nightsea) after mouse sacrifice and dissection. the tissue expressing RFP was

dissected for RNA purification for RT-qPCR (see below), and protein isolation for

Western blot. For protein isolation, the RFP-expressing tissue were homogenized in

RIPA buffer (1% NP40, 0.1% SDS and 0.5% sodium deoxycholate in PBS). The

homogenate was centrifuged at 14,000 x g for 15 min and the supernatant was

collected. Protein concentration was measured by BCA protein assay, and 20 µg total

protein lysate was boiled for 5 min with 6x loading buffer. Standard Western blot was

performed. Antibodies used include Homer1 (160003; Synaptic Systems), Homer1a

(160013; Synaptic Systems), and α-Tubulin (sc32293; Santa Cruz).

RNA Sample Preparation

RNA was extracted from fresh tissue. Tissue for RNA extraction was isolated from V1

contralateral and ipsilateral to the sutured eye after 3-day MD (at P28), 7-day MD (at

P32) or from mice that had not undergone MD (various ages based on experiment).
Briefly, mice were anesthetized with isoflurane and transcardially perfused with ice-cold
phosphate buffered saline (PBS) for at least 5 min. A sterile scalpel blade and forceps
were used to surgically micro-dissect the appropriate tissue section. The dissected
tissue was immediately placed into a 2-mL lysing tube containing 1.4 mm spherical
ceramic beads (MPBio, Lysing Matrix D, Cat. No. 116913050) and pre-filled with 1 mL
of ice-cold TRIzol (Ambion, Cat. No. 15596-026) and then immediately homogenized
with a bead-based homogenizer (FastPrep-24 5G, MPBio) using the preset settings for
mouse brain tissue (8.0 m/sec, 30 sec). Once foaming has subsided at room
temperature, the homogenate was transferred into 2 mL Phase Lock Tube (Heavy;
5Prime Bio, Cat. No. 2302830) pre-filled with 200 µL chloroform and briefly vortexed.
Samples were centrifuged at 12,000 x g for 10 min at 2-4°C. The colorless upper
aqueous layer was then transferred into a new 2.0 mL LoBind tube that was pre-filled
with 1.5 volumes of 100% EtOH (~600 µL), then mixed thoroughly by pipetting up and
down several times, and then briefly centrifuged. The sample was furthered purified
using Zymo RNA Clean & Concentrator-5 (Zymo Research, R1016) according the
manufacturer's recommended protocol and always performed with the optional in-
column DNase I treatment. RNA was eluted twice with ≥6 µL 1x TE buffer (pH 7.5). The
approximate concentration and relative purity of RNA was determined using a
NanoDrop 2000 spectrophotometer (Thermo Scientific). The purified RNA was either
immediately used for downstream applications (i.e., reverse transcription) or stored at -
80°C.
720

721 **RNA sequencing library preparation**
722 For samples to be sequenced, RNA quality was assessed using AATI Fragment
723 Analyzer. Samples with RNA Quality Number of >8.0 were further processed. Indexed
724 cDNA libraries were generated using the SMARTer Stranded Total RNA-Seq Kit v2 and
725 multiplexed sequencing was performed on Illumina HiSeq 2000 or Novaseq 6000.
726

RNAseq Analysis

For the mRNA sequencing, read alignment was performed using HISAT2(60). Fragment
counts were obtained using featureCounts(61). Differential expression analysis was
performed using DESeq2(62). For the circRNA sequencing, reads were aligned and
fragment counts were obtained using circtools(23). Fragment counts were obtained
using the Cufflinks pipeline(63). Differential expression analysis was performed using
the Bioconductor package Limma(64). Gene Ontology enrichment analysis was
performed using the Bioconductor package clusterProfiler(65). Differential expression
analysis of 3-day MD RNAseq data was performed comparing the contralateral side and
ipsilateral side within animal. For 7-day MD RNAseq, differential expression analysis
was performed comparing control, no MD animals and MD animals (contralateral side).
738

cDNA Preparation for qPCR
Total RNA was reversed transcribed using SuperScript IV VILO with ezDNase treatment
(Thermo Fisher, Cat. No. 11766050) according to manufacturer's recommended
protocol with the following modifications: the maximum input of total RNA was lowered
to 2 µg per 20 µL reaction, ezDNase digestion was extended to 5 min, and reverse
transcription temperature was increased to 55°C. After the RT reaction, cDNA was
diluted with 1x TE buffer (pH 8.0).
746

qPCR Analysis
Applied Biosystems QuantStudio 3 Fast thermocycler was used for all qPCR
amplification and detection in this study. PowerUp SYBR Green Master Mix (Applied
Biosystems, Cat. No. A25742) was used according to the manufacturer's recommended
protocol. Primers used were listed in Supplementary Table 5. Mouse *Gapdh* was used
as the internal control for normalization. For the fold change calculation, the ΔΔCq
method was used.
754

Optical imaging of intrinsic signals
At postnatal day P15, sh-scramble or sh-circHomer lenti-viral vectors were injected into
left binocular V1 as described above, and at P22 a 3 mm craniotomy was performed
over the same area, where a 5 mm diameter glass window stacked on top of a 3 mm

diameter glass window was fitted over the craniotomy with dental cement, together with
a metal headplate. Baseline optical imaging was performed at P25, prior to MD. MD
lasted either 3 days or 7 days, after which sutures were removed and the closed eye
was reopened under isoflurane anesthesia before optical imaging experiments to
assess the effects of 3-day or 7-day MD on V1 responses. Optical imaging was
performed as previously described(26). Briefly, mice were lightly anesthetized with
isoflurane (0.5-1%), the window was cleaned with 70% ethanol and a cotton tipped
applicator, and the headplate was attached to the imaging rig to minimize head
movements. Green light (560 nm) was used to focus 400 µm below the cortical surface.
Red light (630 nm) was used for functional imaging, and the change in reflectance was
captured by an electron multiplying CCD camera (Cascade 512B; Roper Scientific)
during the presentation of visual stimuli.

Visual stimulation and optical imaging analysis
The visual stimulus was a horizontal bar 30° wide, made of flickering checkerboard over
a black background, drifting continuously through the peripheral–central dimension of
the visual field. After moving to the last position, the bar would jump back to the initial
position and start another cycle of movement; therefore, the chosen region of visual
space (72° x 72°) was stimulated in a periodic manner (12 s/cycle, 20 repetitions).
Images were continuously captured at the rate of 30 frames/s during each stimulus
session of 4 min, with a separate stimulus session for each of the 4 cardinal directions.
A temporal component at the stimulus frequency ($12s^{-1}$) was calculated pixel by pixel
from the whole set of images using custom python scripts
(https://github.com/Palpatineli/oi_analyzer). The amplitude of the FFT component was
used to measure the strength of visually driven response for each eye, and the ODI was
derived from the response (R) of each eye at each pixel as ODI = ($R_{contralateral}$ -
$R_{ipsilateral}$)/($R_{contralateral}$ + $R_{ipsilateral}$). The binocular zone was defined as the cortical region
that was driven by both eyes. The response amplitude for each eye was defined as
fractional changes in reflectance over baseline reflectance ($\Delta R/R$ x 10^{-3}), and the top
50% pixels were analyzed to avoid background contamination.

Epitope-preserving magnified analysis of the proteome (eMAP)

Mice were injected at P15as described above with AAV2/9-hSyn-DIO-EGFP (1×10^{13} vg/mL), AAV2/9-CaMKIIa-Cre (5×10^{8} vg/mL), and AAV9 expressing either pAV-U6-sh-scramble-CMV-DsRed or pAV-U6-sh-circHomer1-CMV-DsRed (1×10^{13} vg/mL). Animals were allowed to express virus for a minimum of 10 days before MD. After MD, mice were transcardially perfused under deep anesthesia with ice-cold PBS followed by ice-cold 4% PFA (in PBS), and then brains were post-fixed in 4% PFA at 4°C overnight. eMAP processing was performed following protocols previously described(66, 67). Tissue embedding was performed using eMAP solution (30% acrylamide, 10% sodium acrylate, 0.1% bisacrylamide, 0.03% VA-044 in PBS) under vacuum, following hydration using hydration solution (0.02% sodium azide in PBS) and sectioning into 60um thick using Leica VT1000S vibratome. Tissue clearing was performed by incubation in clearing solution (6% SDS, 0.1 M phosphate buffer, 50 mM sodium sulfite, 0.02% sodium azide in DI water, pH 7.4) at 37°C for 4 hours. If applicable, after washing with PBST (0.1% Triton X-100, 0.02% sodium azide in PBS), samples were stained with primary antibodies against GFP (A10262, Invitrogen) and Arc (156003, Synaptic Systems) diluted in PBST at 37°C for 48 hours. After washing with PBST, samples were stained with secondary antibodies diluted in PBST for 48 hours. Expansion was performed using 0.01× PBS before imaging. Approximately 3× total linear expansion was achieved consistently. Pyramidal neurons in V1 which were double-positive for RFP and GFP were identified and imaged using Leica SP8 confocal microscope with 20× 0.75 NA or 63× 1.2 NA water immersion lens.

Spine morphology and Arc puncta volume analysis

Dendrites and dendritic spines for analysis were selected and analyzed by an experimenter blinded to experiment condition. To analyze dendritic spine morphology, the length (L), head width (H) and neck width (N) were measured by ImageJ. Dendritic spines were classified into 4 types according to the criteria previously described(29). Briefly, the average length of spine head in the No MD sh-scramble group ($\bar{\bar{H}}$) was calculated. Protrusions with H>N and H> $\bar{\bar{H}}$ were defined as mushroom spines; protrusions with H>N and H< $\bar{\bar{H}}$ were defined as thin spines; protrusions with H≤N and

L<$\overline{H}$ were defined as stubby spines, and protrusions with H≤N and L>1.5*$\overline{H}$ were
defined as filopodia. For dendritic spine volume and surface area analysis, 3D
reconstruction and quantification of dendritic spines volume was performed using the
"Filament" function in Imaris software. The analysis of Arc puncta volume was
performed using the "Surface" function in Imaris, and only the Arc puncta located at
dendritic spines were selected for quantification. The raw data was then divided by 3^3
(=27) to convert to values that closely approximates the unexpanded volume.

Preparation of acute brain slices for electrophysiology
Mice injected with either sh-scramble or sh-circHomer1 lentivirus (as described above)
were anesthetized with 2-3% isoflurane and decapitated. Brains were rapidly removed
and placed into ice-cold low-Ca2+, low-Na+ sucrose cutting solution consisting of (in
mM): 234 sucrose, 11 glucose, 24 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 10 MgSO4, and
0.5 CaCl2. Cutting solution was oxygenated with a mixture of 95% O2 and 5% CO2.
300µM slices were cut in a coronal orientation using a vibratome (Leica Biosystems,
Buffalo Grove, IL) and placed into a recovery chamber filled with oxygenated ACSF
consisting of (in mM): 126 NaCl, 2.5 KCl, 1.25 NaH2PO4, 1 MgSO4, 2 CaCl2, 10
glucose, 26 NaHCO3. Slices were allowed to recover at 34°C for 1 hour before being
kept at room temperature (25°C) for the remainder of the experiment.

Current clamp recordings and analysis
TdTomato-positive neurons were identified in Layer 2/3 of the V1 binocular region.
Recordings were made at 34°C in ACSF containing 10 µM CPP, 20 µM DNQX, and 10
µM SR95531 to isolate passive membrane properties. Signals were collected at 10kHz
using a Multiclamp 700B amplifier and digitized using a Digidata 1440A (Molecular
Devices, San Jose, CA). Whole-cell patch clamp recordings were made using
borosilicate glass electrode with an open-tip resistance of 2-5 MΩ filled with an
intracellular solution consisting of (in mM): 130 potassium gluconate, 10 HEPES, 5 KCl,
5 EGTA, 2 NaCl, 1 MgCl2, 10 TEA-Cl, 10 phosphocreatine, 2 Mg-ATP, 0.3 Na-GTP (pH
adjusted to 7.28, osmolarity adjusted to 290 mOsm). Current steps of 250 ms duration
were applied, from -100 pA increasing to 650 pA at 25 pA increments. The holding

current was returned to 0 pA for 750 ms between each current injection step. Ra

(access resistance) and Rm (membrane resistance) were also measured in response to

a 5 mV depolarizing step, and membrane capacitance was calculated as $C = \frac{\tau}{Rm}$, where

т is the decay of the capacitive transient in response to the 5 mV step. Access

resistance was monitored throughout the recording and recordings with >20% change in

access resistance or had access resistance >30 MΩ were excluded.

mEPSC recordings and analysis

Slices were prepared as above. Recordings of tdTomato positive neurons were made at

34°C in ACSF containing 10 µM CPP, 10 µM SR95531, and 1 µM tetrodotoxin to isolate

miniature AMPA receptor-mediated currents. Signals were collected at 10kHz and

filtered at 2kHz using a Multiclamp 700B amplifier and digitized using a Digidata 1440A

(Molecular Devices, San Jose, CA). Whole-cell patch clamp recordings were

established using borosilicate glass electrode with an open-tip resistance of 2-5 MΩ

filled with an intracellular solution consisting of (in mM): 100 potassium gluconate, 10

HEPES, 20 KCl, 0.5 EGTA, 10 NaCl, 8 phosphocreatine, 2 Mg-ATP, 0.3 Na-GTP (pH

adjusted to 7.23, osmolarity adjusted to 290 mOsm). Neurons were voltage-clamped at

-70 mV and stable gap free recordings were made for at least 2 minutes. Access

resistance was monitored throughout the recording and recordings with >20% change in

access resistance or had access resistance >30 MΩ were excluded. mEPSC events

were detected using the MiniAnalysis software (Synaptosoft, Fort Lee, NJ).

Surface Protein Biotinylation Assay

For the measurement of surface proteins, we first prepared 300 µm acute coronal slices

containing V1 in ice-cold ACSF and then washed slices 3 times in ice-cold ACSF in 6-

well plate on shaker. The sections were incubated in 100 µM S-NHS-SS-biotin for 45

min (~1 mL per well). The superficial layers of V1 were dissected into a new 1.5 mL

tube filled with ice-cold ACSF and homogenize in RIPA buffer. The homogenate was

centrifuged at 14,000 x g for 5 min and the supernatant was transferred into a new 1.5

mL tube. ~20% volume was transferred into a new 1.5 mL tube and stored at -80°C to

be used later as a total protein control. ACSF was added to the remaining supernatant

for a final volume of 1 mL per sample. 40 µL of streptavidin beads were added and
incubated on a shaker overnight at 4°C. Samples were then centrifuged at 3,500 x g for
1 min. The supernatant was discarded and then the beads were washed 3x in 1:1 v/v
ice-cold solution of ACSF and RIPA buffer. Then, 40 µL of 2x loading buffer was added,
mixed briefly, and boiled for 5 min. Standard Western blot was performed. Antibodies
used include GluA1 (04-855, clone C3T; Millipore), and pan-Cadherin (ab6529; Abcam).
889

Neuronal cultures, transfection, and pharmacological treatment
Cultured primary rat hippocampal and cortical cells were prepared as previously
described(68). In brief, Sprague–Dawley rat embryos were sacrificed on embryonic day
18. The hippocampus and cortices were dissected, and the cells were dissociated with
trypsin. Hippocampal cells were cultured on 18-mm coverslips coated with 1□mg/mL
poly-D-lysine at 1×10^5 cells per coverslip in Neurobasal Plus medium supplemented
with 2% B27 Plus and 0.5mM L-glutamate. Cortical cells were cultured on 60-mm
cultural dishes coated with 100µg/mL poly-L-lysine at 3×10^6 per dish in Neurobasal
medium supplemented with 2% B27 and 10mM glucose. All cells were maintained at 37
°C in a humidified atmosphere with 5% CO_2. The cortical neurons were treated with
bicuculine (20 µM) at DIV 13. Total RNA of cortical neurons was extracted at 0, 2, 6, 24,
or 48 hours later using NucleoSpin RNA Mini kit (Macherey-Nagel, 740955) according
to the manufacturer's protocol. The hippocampal neurons at DIV 11 were transfected
with a *circHomer1* or scrambled control shRNA construct and SEP-GluA1 using calcium
phosphate precipitation(68). The transfected neurons were then treated with bicuculine
(20 µM, 48 h) for subsequent morphological analysis. At DIV 19 the hippocampal
neurons were fixed with 4% PFA, and blocking was performed for 1 hour at room
temperature with 1% BSA in PBS. Antibody against GFP (A10262, Invitrogen) was
diluted in 1% BSA in PBS and incubated with cells at 4°C overnight. After washing with
PBS, the cells were incubated with secondary antibody for 1 hour at room temperature.
The cells were then washed with PBS and mounting was performed. Imaging was
performed under Leica SP8 confocal microscope using a 63x oil immersion lens with
1.4 numerical aperture.
913

Computational modeling

As it is unclear what stages in the biogenesis of *circHomer1* confer its activity-dependence, we compared three nested dynamical system models(69) of *circHomer1* expression with the same basic architecture to assess which best fit our data. We modeled three time-varying outputs: (1) *Homer1b* mRNA levels (H1b), (2) *Homer1a* mRNA levels (H1a), and (3) *circHomer1* RNA levels (CH). Dynamical changes in gene expression were described by the following differential equations:

$$\Delta H1a = (H1aTR - decL \cdot H1a - H1aC \cdot CR \cdot H1a)dt$$
$$\Delta H1b = (H1bTR - decL \cdot H1b - CR \cdot H1b)dt$$
$$\Delta CH = (CR \cdot H1b + H1aC \cdot CR \cdot H1a - decCH \cdot CH)dt$$

with H1a/bTR denoting transcription rates of pre-mRNA, decL denoting linear mRNA decay rate, decCH denoting circular RNA decay rate, CR denoting circularization rate, and H1aC denoting a logical operator of whether *Homer1a* can be circularized (1 if true, 0 if false). Parameters were fit separately for each model/experiment combination, but the fit parameter values for a particular model were largely consistent across experiments.

We hypothesized(21, 47) that the activity-dependence of *circHomer1* expression was most likely to arise from circularization of an activity-dependent pre-mRNA (*Homer1a*) and/or activity dependent circularization of a constitutively expressed pre-mRNA (*Homer1b*). The three models tested were as follows. (1) The constant circularization model fit a constant circularization rate of pre-*Homer1a* and pre-*Homer1b*, with the only time varying parameter being the transcription rate of *Homer1a* which was fit to measured data. (2) The variable circularization model allowed the circularization rate to vary over time, however only pre-*Homer1b* could be circularized. (3) The third, full model also included time varying circularization but both pre-*Homer1b* and pre-*Homer1a* could be circularized. *Homer1a* and *circHomer1* levels were explicitly fit to minimize squared error from observed datapoints for each experiment. *Homer1b* levels were treated as a 'hidden' variable; as they were not measured experimentally and were not a part of model testing.

Models were initialized 10 days before the first timepoint to allow RNA levels to stabilize
before permitting any changes in transcription or circularization. Because of the varied
measurements and methods across experiments, data in each model were normalized
to the first timepoint, T=0. In models where circularization of pre-*Homer1a* was
permitted, the same circularization rate was applied to both pre- *Homer1a* and *Homer1b*
pools. Fit circularization rates were 1 to 2 orders of magnitude smaller than the linear
splicing rates(70), and varied with activity by at most a factor of 4. In each of the
models, the decay rate of *circHomer1* was constrained to be slower than that of linear
Homer1 mRNA by a factor of ≥2(48). All models were run with 15 minute timesteps.
Homer1b transcription rate, linear decay rate, and circular decay rate were fit as
constant parameters in each model. *Homer1a* transcription rate was fit as a variable
parameter in each model. Circularization rate was fit as a constant parameter in the
constant circularization model, and a variable parameter in the variable circularization
and full model. Fitting was performed and AIC scores for each model were computed
using the python package lmfit.

Statistical analysis

Data are presented as mean ± SEM in quantitative analysis unless other specified.
When two independent experimental groups were analyzed, Student's t-test was
performed, while paired t-test was performed when two paired experimental data were
analyzed. When more than two independent experimental groups were analyzed, one-
way ANOVA followed by Tukey post hoc test was performed. For those experiments
exploring the effect of *circHomer1* during MD by *circHomer1* depletion, two-way ANOVA
followed by Tukey post hoc test was performed. For optical imaging, mixed-effects
model following Holm-Sidak multiple comparisons test was performed. For mEPSC
quantification, two-sample Kolmogorov-Smirnoff test was performed. Statistical
significance was defined as $p < 0.05$.

References

1. E. L. Yap, M. E. Greenberg, Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. *Neuron* **100**, 330–348 (2018).

2. N. Mellios, *et al.*, MiR-132, an experience-dependent microRNA, is essential for visual cortex plasticity. *Nature Neuroscience* **14**, 1240–1242 (2011).

3. W. Chen, E. Schuman, Circular RNAs in Brain and Other Tissues: A Functional Enigma. *Trends Neurosci* **39**, 597–604 (2016).

4. L. Szabo, J. Salzman, Detecting circular RNAs: bioinformatic and experimental challenges. *Nat Rev Genet* **17**, 679–692 (2016).

5. L. L. Chen, The expanding regulatory mechanisms and cellular functions of circular RNAs. *Nat Rev Mol Cell Biol* **21**, 475–490 (2020).

6. B. Kleaveland, C. Y. Shi, J. Stefano, D. P. Bartel, A Network of Noncoding Regulatory RNAs Acts in the Mammalian Brain. *Cell* **174**, 350-362.e17 (2018).

7. S. A. Giusti, *et al.*, A brain-enriched circular RNA controls excitatory neurotransmission and restricts sensitivity to aversive stimuli. *Sci Adv* **10**, eadj8769 (2024).

8. M. Piwecka, *et al.*, Loss of a mammalian circular RNA locus causes miRNA deregulation and affects brain function. *Science* **357**, eaam8526 (2017).

9. A. J. Zimmerman, *et al.*, A psychiatric disease-related circular RNA controls synaptic gene expression and cognition. *Mol Psychiatry* **25**, 2712–2727 (2020).

10. Y. J. Sun, J. Sebastian Espinosa, M. S. Hoseini, M. P. Stryker, Experience-dependent structural plasticity at pre- and postsynaptic sites of layer 2/3 cells in developing visual cortex. *Proceedings of the National Academy of Sciences* **116**, 21812–21820 (2019).

11. B. M. Hooks, C. Chen, Circuitry Underlying Experience-Dependent Plasticity in the Mouse Visual System. *Neuron* **106**, 21–36 (2020).

12. M. Majdan, C. J. Shatz, Effects of visual experience on activity-dependent gene regulation in cortex. *Nature Neuroscience* **9**, 650–659 (2006).

13. T. K. Hensch, E. M. Quinlan, Critical periods in amblyopia. *Vis Neurosci* **35**, E014 (2018).

14.   M. Y. Frenkel, M. F. Bear, How monocular deprivation shifts ocular dominance in

visual cortex of young mice. *Neuron* **44**, 917–923 (2004).

15.   C. N. Levelt, M. Hübener, Critical-period plasticity in the visual cortex. *Annu Rev*

Neurosci **35**, 309–330 (2012).

16.   J. A. Gordon, M. P. Stryker, Experience-Dependent Plasticity of Binocular

Responses in the Primary Visual Cortex of the Mouse. *Journal of Neuroscience* **16**,

3274–3286 (1996).

17.   D. Tropea, A. Van Wart, M. Sur, Molecular mechanisms of experience-

dependent plasticity in visual cortex. *Philos Trans R Soc Lond B Biol Sci* **364**, 341–355

(2009).

18.   J. S. Espinosa, M. P. Stryker, Development and plasticity of the primary visual

cortex. *Neuron* **75**, 230–249 (2012).

19.   K. R. Jenks, K. Tsimring, J. P. K. Ip, J. C. Zepeda, M. Sur, Heterosynaptic

Plasticity and the Experience-Dependent Refinement of Developing Neuronal Circuits.

Front Neural Circuits **15**, 803401 (2021).

20.   M. Sur, I. Nagakura, N. Chen, H. Sugihara, Mechanisms of plasticity in the

developing and adult visual cortex. *Prog Brain Res* **207**, 243–254 (2013).

21.   A. K. Hafez, *et al.*, A bidirectional competitive interaction between circHomer1

and Homer1b within the orbitofrontal cortex regulates reversal learning. *Cell Reports* **38**,

110282 (2022).

22.   A. J. Zimmerman, J. P. Weick, G. Papageorgiou, N. Mellios, J. L. Brigman,

Aberrant encoding of event saliency in the orbitofrontal cortex following loss of the

psychiatric-associated circular RNA, circHomer1. *Transl Psychiatry* **14**, 1–14 (2024).

23.   T. Jakobi, A. Uvarovskii, C. Dieterich, circtools-a one-stop software solution for

circular RNA research. *Bioinformatics* **35**, 2326–2328 (2019).

24.   D. Tropea, *et al.*, Gene expression changes and molecular pathways mediating

activity-dependent plasticity in visual cortex. *Nature Neuroscience* **9**, 660–668 (2006).

25.   I. Nagakura, A. Van Wart, J. Petravicz, D. Tropea, M. Sur, STAT1 regulates the

homeostatic component of visual cortical plasticity via an AMPA receptor-mediated

mechanism. *J Neurosci* **34**, 10256–10263 (2014).

26.   J. P. K. Ip, *et al.*, Major Vault Protein, a Candidate Gene in 16p11.2 Microdeletion

Syndrome, Is Required for the Homeostatic Regulation of Visual Cortical Plasticity. *J*

Neurosci **38**, 3890–3900 (2018).

27.   C. L. McCurry, *et al.*, Loss of Arc renders the visual cortex impervious to the

effects of sensory experience or deprivation. *Nature neuroscience* **13**, 450–7 (2010).

28.   V. A. Kalatsky, M. P. Stryker, New paradigm for optical imaging: temporally

encoded maps of intrinsic signal. *Neuron* **38**, 529–545 (2003).

29.   C. L. Peebles, *et al.*, Arc regulates spine morphology and maintains network

stability in vivo. *Proc Natl Acad Sci U S A* **107**, 18173–18178 (2010).

30.   K. P. Berry, E. Nedivi, Spine Dynamics: Are They All the Same? *Neuron* **96**, 43–

55 (2017).

31.   A. C. McClelland, M. Hruska, A. J. Coenen, M. Henkemeyer, M. B. Dalva, Trans-

synaptic EphB2–ephrin–B3 interaction regulates excitatory synapse density by inhibition

of postsynaptic MAPK signaling. *Proceedings of the National Academy of Sciences* **107**,

8830–8835 (2010).

32.   J. D. Shepherd, *et al.*, Arc/Arg3.1 Mediates Homeostatic Synaptic Scaling of

AMPA Receptors. *Neuron* **52**, 475–484 (2006).

33.   X. You, *et al.*, Neural circular RNAs are derived from synaptic genes and

regulated by development and plasticity. *Nat Neurosci* **18**, 603–610 (2015).

34.   S. Chowdhury, *et al.*, Arc/Arg3.1 interacts with the endocytic machinery to

regulate AMPA receptor trafficking. *Neuron* **52**, 445–459 (2006).

35.   A. Rybak-Wolf, *et al.*, Circular RNAs in the Mammalian Brain Are Highly

Abundant, Conserved, and Dynamically Expressed. *Molecular Cell* **58**, 870–885 (2015).

36.   N. E. Clifton, S. Trent, K. L. Thomas, J. Hall, Regulation and Function of Activity-

Dependent Homer in Synaptic Plasticity. *Mol Neuropsychiatry* **5**, 147–161 (2019).

37.   G. H. Diering, *et al.*, Homer1a drives homeostatic scaling-down of excitatory

synapses during sleep. *Science* **355**, 511–515 (2017).

38.   V. Chokshi, B. Druciak, P. F. Worley, H. K. Lee, Homer1a is required for

establishment of contralateral bias and maintenance of ocular dominance in mouse

visual cortex. *Journal of Neuroscience* **39**, 3897–3905 (2019).

39.   J. H. Hu, *et al.*, Homeostatic Scaling Requires Group I mGluR Activation
Mediated by Homer1a. *Neuron* **68**, 1128–1142 (2010).
40.   S. C. Martin, S. K. Monroe, G. H. Diering, Homer1a and mGluR1/5 Signaling in
Homeostatic Sleep Drive and Output. *Yale J Biol Med* **92**, 93–101 (2019).
41.   K. M. Tyssowski, *et al.*, Firing Rate Homeostasis Can Occur in the Absence of
Neuronal Activity-Regulated Transcription. *J Neurosci* **39**, 9885–9899 (2019).
42.   A. Rozov, A. R. Zivkovic, M. K. Schwarz, Homer1 gene products orchestrate
Ca(2+)-permeable AMPA receptor distribution and LTP expression. *Front Synaptic*
Neurosci **4**, 4 (2012).
43.   D. Shin, M. Nam, Y. Yoon, M. Kim, Membrane-based hybridization capture of
intracellular peptide nucleic acid. *Analytical Biochemistry* **399**, 135–137 (2010).
44.   B. J. Yoon, G. B. Smith, A. J. Heynen, R. L. Neve, M. F. Bear, Essential role for a
long-term depression mechanism in ocular dominance plasticity. *Proc Natl Acad Sci U S*
A **106**, 9860–9865 (2009).
45.   F. Ango, *et al.*, Dendritic and axonal targeting of type 5 metabotropic glutamate
receptor is regulated by homer1 proteins and neuronal excitation. *J Neurosci* **20**, 8710–
8716 (2000).
46.   V. Chokshi, *et al.*, Input-Specific Metaplasticity in the Visual Cortex Requires
Homer1a-Mediated mGluR5 Signaling. *Neuron* **104**, 736-748.e6 (2019).
47.   N. Mellios, *et al.*, Regulation of neuronal circHomer1 biogenesis by
PKA/CREB/ERK-mediated pathways and effects of glutamate and dopamine receptor
blockade. *Res Sq* rs.3.rs-3547375 (2024). https://doi.org/10.21203/rs.3.rs-3547375/v1.
48.   Y. Enuka, *et al.*, Circular RNAs are long-lived and display only minimal early
alterations in response to a growth factor. *Nucleic Acids Research* **44**, 1370–1383
(2016).
49.   Y. Zhang, *et al.*, The Biogenesis of Nascent Circular RNAs. *Cell Reports* **15**,
611–624 (2016).
50.   S. Farris, G. Lewandowski, C. D. Cox, O. Steward, Selective localization of arc
mRNA in dendrites involves activity- and translation-dependent mRNA degradation. *J*
Neurosci **34**, 4481–4493 (2014).

51.   C. Giorgi, *et al.*, The EJC factor eIF4AIII modulates synaptic strength and
neuronal protein expression. *Cell* **130**, 179–191 (2007).

52.   C. Xu, *et al.*, Structure and plasticity of silent synapses in developing
hippocampal neurons visualized by super-resolution imaging. *Cell Discovery* **6**, 1–11
(2020).

53.   W. R. Jeck, *et al.*, Circular RNAs are abundant, conserved, and associated with
ALU repeats. *RNA* **19**, 141–157 (2013).

54.   D. Liang, J. E. Wilusz, Short intronic repeat sequences facilitate circular RNA
production. *Genes Dev.* **28**, 2233–2247 (2014).

55.   Q. Zheng, *et al.*, Circular RNA profiling reveals an abundant circHIPK3 that
regulates cell growth by sponging multiple miRNAs. *Nat Commun* **7**, 11215 (2016).

56.   N. R. Choudhury, *et al.*, Tissue-specific control of brain-enriched miR-7
biogenesis. *Genes and Development* **27**, 24–38 (2013).

57.   T. B. Hansen, *et al.*, Natural RNA circles function as efficient microRNA sponges.
Nature **495**, 384–388 (2013).

58.   S. Memczak, *et al.*, Circular RNAs are a large class of animal RNAs with
regulatory potency. *Nature* **495**, 333–338 (2013).

59.   M. Rajman, G. Schratt, MicroRNAs in neural development: from master
regulators to fine-tuners. *Development* **144**, 2310–2322 (2017).

60.   D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–
915 (2019).

61.   Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930
(2014).

62.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

63.   C. Trapnell, *et al.*, Differential gene and transcript expression analysis of RNA-
seq experiments with TopHat and Cufflinks. *Nature Protocols* **7**, 562–578 (2012).

64.   M. E. Ritchie, *et al.*, limma powers differential expression analyses for RNA-
sequencing and microarray studies. *Nucleic Acids Research* **43**, e47 (2015).

65.   G. Yu, L. G. Wang, Y. Han, Q. Y. He, clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

66.   T. Ku, *et al.*, Multiplexed and scalable super-resolution imaging of three-

dimensional protein localization in size-adjustable tissues. *Nat Biotechnol* **34**, 973–981

(2016).

67.   J. Park, *et al.*, Epitope-preserving magnified analysis of proteome (eMAP). *Sci*

Adv **7**, eabf6589 (2021).

68.   Y. Wang, *et al.*, Astrocyte-secreted IL-33 mediates homeostatic synaptic

plasticity in the adult hippocampus. *Proceedings of the National Academy of Sciences*

118, e2020810118 (2021).

69.   T. Chen, H. L. He, G. M. Church, Modeling gene expression with differential

equations. *Pac Symp Biocomput* 29–40 (1999).

70.   A. Bachmayr-Heyda, *et al.*, Correlation of circular RNA abundance with

proliferation--exemplified with colorectal and ovarian cancer, idiopathic lung fibrosis, and

normal human tissues. *Sci Rep* **5**, 8057 (2015).

1138

Figure 1

Figure 2

Figure 3

A

sh-circHomer1 or sh-scramble

Craniotomy   Lid suture

Virus expression   Homecage   3-day or 7-day MD

P15   P20   P25

postMD acquisition

P28/P32

preMD acquisition

Flicker checkerboard

B

Example sh-circHomer1

preMD   7-day MD

Contralateral

Ipsilateral

Relative phase   -π   0   +π

Response amplitude   0   0.5   1.0

C

Ocular dominance index

sh-scramble   sh-circHomer1

* ** ** ***

D

Contralateral

Response amplitude

sh-scramble   sh-circHomer1

* ** ** * **

Ipsilateral

Response amplitude

sh-scramble   sh-circHomer1

* ** ** * **

preMD   3-day MD   7-day MD

Figure 4

Figure 5

A

	No MD		3-day MD		7-day MD	
	Scr	KD	Scr	KD	Scr	KD

GluA1

pan-Cad

■ sh-scramble
■ sh-circHomer1

Surface GluA1 level

**

B

Fluorescent image   3D reconstruction

GFP
Arc

C

No MD   3-day MD   7-day MD

Scr   KD   Scr   KD   Scr   KD

Arc

GFP

3D reconstruction

D

■ sh-scramble
■ sh-circHomer1

Percentage of Arc containing dendritic spines (%)

***
***
***
***

No MD   3-day MD   7-day MD

E

Arc puncta volume (μm³)

***
*
**
**

No MD   3-day MD   7-day MD

F

● sh-scramble
● sh-circHomer1

Arc volume / dendritic spine volume

***
*
*
***

No MD   3-day MD   7-day MD