An application of the MR-Horse method to reduce selection bias in genome-wide association studies of disease progression

Killian Donovan[1], Jason Torres[1], Doreen Zhu[1], William G Herrington[1], Natalie Staplin[1]

[1]Nuffield Department of Population Health, University of Oxford, Old Road Campus, OX3 7LF, UK

Correspondence to: killian.donovan@ndph.ox.ac.uk

Word count (abstract): 250

Word count (main text): 3909

Figures: 4

Abstract

Genome-wide association studies (GWAS) of disease progression are vulnerable to collider bias caused by selection of participants with disease at study entry. This bias introduces spurious associations between disease progression and genetic variants that are truly only associated with disease incidence. Methods of statistical adjustment to reduce this bias have been published, but rely on assumptions regarding the genetic correlation of disease incidence and disease progression which are likely to be violated in many human diseases.

MR-Horse is a recently published Bayesian method to estimate the parameters of a general model of genetic pleiotropy in the setting of Mendelian Randomisation. We adapted this method to provide bias-reduced GWAS estimates of associations with disease progression,

robust to the genetic correlation of disease incidence and disease progression and robust to the

presence of pleiotropic variants with effects on both incidence and progression.

We applied this adapted method to simulated GWAS of disease incidence and progression with

pleiotropic variants and varying degrees of genetic correlation. When significant genetic

correlation was present, the MR-Horse method produced less biased estimates than

unadjusted analyses or analyses adjusted using other existing methods. Type 1 error rates with

the MR-Horse method were consistently below the nominal 5% level, at the expense of a

modest reduction in power. We then applied this method to summary statistics from the

CKDGen consortium GWAS of kidney function decline. MR-Horse attenuated the effects of

variants with known likely biased effects in the CKDGen GWAS, whilst preserving effects at loci

with likely true effects.

Introduction

Genome-wide association studies (GWAS) of disease progression are increasingly common due

to the increasing availability of sufficiently large cohorts of genotyped participants with specific

diseases and long-term follow-up[1–3]. Selection for disease cases implicitly conditions on the

processes causing incident disease, thus inducing biased associations through any confounders

of the incidence - progression relationship[4,5] (Fig 1). Eliminating this bias by study design is

challenging, as in order to define "disease progression", participants must be stratified in some

way by disease status. Measurement and elimination of this bias is important for understanding

the genomic architecture of disease traits, and also could be applied to reduce the bias of

estimates in non-genetic epidemiological studies.

If individual-level genetic data are available, and the selection-process responsible for the bias

is well understood, then the bias can be addressed by inverse-probability weighting of

participants according to the probability of selection[6]. Unfortunately individual level data are

often not available. Methods have been proposed to adjust GWAS summary statistics in order

to measure and reduce selection bias[6-8]. Given that the bias arises from conditioning on disease

incidence, these methods use knowledge of genetic variant associations with disease incidence

to estimate the bias of each SNP estimate in the stratified progression GWAS.

The method proposed by Dudbridge and colleagues[7] (termed "instrument effect regression" by

some authors[8]) assumes that there is no correlation between direct genetic effects of a variant

on incident disease (β_{GX}) and its effects on disease progression (β_{GY}), and thus any correlation

between estimated effects on incidence and progression must be due to collider bias according

to the equation

$$\beta_{GY}^* = \beta_{GY} + b\beta_{GX}$$

where β_{GY}^* denotes the biased value of β_{GY} observed in a GWAS stratified by incident disease.

The collider bias effect is thus defined by the coefficient b, which can be estimated by linear

regression of estimated progression effects from the biased GWAS on estimated incidence

effects from a separate study (provided incident disease in both studies is defined in an

identical manner). They also propose two possible mechanisms for adjusting results for

measurement error in $\hat{\beta}_{GX}$. The underlying assumption of no correlation is very strong, and in

our view likely to be violated in most disease contexts. In the case of chronic kidney disease for

example, incident disease is generally defined based on the measurement of a threshold value

of a biomarker-based (usually serum creatinine) estimate of glomerular filtration rate (eGFR),

and disease progression is typically defined as rate of change of this value over time (eGFR

slope)[9]. In such an example, given that both phenotypes are defined based on the same

measurements, it is likely that there is significant correlation between genetic effects on the

phenotypes.

The method proposed by Mahmoud and colleagues[10] (Slope-Hunter method) does not assume

that β_{GX} and β_{GY} are uncorrelated for all SNPs. Gaussian mixture models are used to cluster

incidence-associated SNPs into a "hunted" cluster of SNPs with similar values of $\frac{\beta_{GY}^*}{\beta_{GX}}$ and a

second cluster without such a relationship (the "pleiotropic" cluster). Any correlation between

β_{GX} and β_{GY} among the "hunted" SNPs is assumed to be caused by the collider bias effect,

which can thus be estimated by linear regression of estimated progression effects on estimated

incidence effects in this cluster. Accurate identification of the appropriate "hunted" cluster

requires that a plurality of variants with similar values of $\frac{\beta_{GY}^*}{\beta_{GX}}$ only have true effects on disease

incidence and not disease progression. This is termed the **Ze**ro **M**odal **R**esidual **A**ssumption

[ZeMRA], and is less restrictive than that of the Dudbridge method, but still likely to be violated

in certain human diseases. It is therefore desirable to have a method which can reduce the bias

of disease progression associations without assumptions regarding the underlying genetic

correlation of the traits of interest.

The model of genetic causation used by Grant and Burgess in the development of MR-Horse[11]

(Fig 2a) is similar to the proposed model of causation in GWAS of disease progression (Fig 2b),

and its application to GWAS can be summarized as follows, amending some of the notation

used by Grant and Burgess:

5
$$\hat{\beta}^*_{GY_i} \sim N\big(\beta_{GY_i} + b\beta_{GX_i}, \sigma^2_{GY}\big)$$

6
$$\hat{\beta}_{GX_i} \sim N\big(\beta_{GX_i}, \sigma^2_{GX}\big)$$

7
$$\begin{pmatrix} \beta_{GX_i} \\ \beta_{GY_i} \end{pmatrix} \sim N\left(\begin{pmatrix} \mu \\ 0 \end{pmatrix}, \begin{pmatrix} s_x^2 & \rho_i s_x \phi_i \tau \\ \rho_i s_x \phi_i \tau & \phi_i^2 \tau^2 \end{pmatrix} \right)$$

where ρ_i represents the correlation coefficient between β_{GX_i} and β_{GY_i} and ϕ_i and τ are half

Cauchy distributed variables representing the local and global shrinkage parameters

respectively of a horseshoe prior[12] for β_{GY}. There are some differences in the interpretation of

parameters in the setting of collider bias vs in the MR setting. θ in the original MR-Horse setting

(Fig 2) represents the direct causal effect of the exposure on the outcome. In the collider bias

setting this value b incorporates residual direct effects from "Incidence" to "Progression" not

completely accounted for by adjustment for Incidence, in addition to the effects mediated

along the pathway through the confounders U. It therefore defines the magnitude of the

collider bias but is not intuitively interpretable in the same way as θ in the original MR-Horse

formulation. In the MR setting, α_i represents the pleiotropic effects of the genetic variant i and

its value is not of significant interest *per se*, whereas in the application to collider bias, β_{GY_i} is

the direct genetic effect of a variant on disease progression and thus is the estimand of interest.

We aimed to adapt the MR-Horse method to produce bias-reduced estimates of disease-

progression associations in GWAS which adjust for incident disease, robust to correlation

between β_{GX} and β_{GY}. We then tested this method in published GWAS of CKD progression,

taking advantage of the fact that there are known biased associations of genetic variants with

progression[13] which can be used as negative controls.

Methods

Simulations

In order to test this novel application of the method we first simulated GWAS of disease

incidence and progression, using knowledge of chronic kidney disease to approximately inform

modelling choices. Genotypes were simulated for 10,000 individuals at 2,000 independent SNPs

in Hardy-Weinberg equilibrium[14] and in linkage equilibrium. 95% of SNPs had no effect on

disease incidence or progression (G_0), 1% of SNPs had effects on disease incidence only (G_I), 1%

of SNPs had effects on disease progression only (G_P) and 3% of SNPs had effects on both

incidence and progression (G_{IP}), with correlated β_{GX_i} and β_{GY_i} with correlation coefficient ρ

(constant across G_{IP} variants). This set-up produces a large pleiotropic cluster of variants

explaining more of the phenotypic variance of disease progression than the non-pleiotropic

cluster. This situation is likely to violate the assumptions of both the Dudbridge and Slope-

Hunter methods, but is a plausible architecture of chronic kidney disease related traits. In order

to approximate realistic distributions of effect sizes across the range of allele frequencies,

minor allele frequencies were first drawn from a uniform distribution between 0.01 and 0.49,

and standardised effect sizes were defined as $\pm\sqrt{\dfrac{r^2}{2maf(1-maf)}}$ where variant level r^2 values

were drawn from the positive half of a normal distribution with mean 0 and standard deviation

$\dfrac{R^2\sqrt{\pi}}{n\sqrt{2}}$, where R^2 represents the phenotypic variance explained by the cluster, and n represents

the number of SNPs in the cluster. Confounders U were simulated as a random normal variable,

and disease incidence (Inc) was initially simulated as a continuous variable with 40% SNP

heritability, 30% of its variance explained by the confounders U and the remaining variance

explained by a further random normal variable ϵ_I, according to the equation

8
$$Inc_j = \sum_{i=1}^{2000} \beta_{GX_i} G_{ij} + U_j + \epsilon_{I_j}$$

A threshold value of the Inc phenotype was selected to define the binary phenotype of incident

CKD such that the prevalence of CKD in the simulated population was 40%. Kidney function

decline or disease progression was then simulated in all participants as a continuous variable

with genetic effects on progression scaled such that the phenotype has 30-40% SNP heritability

(estimated in GWAS among the entire simulated population) and 30-40% of its variance

explained by the confounders U, according to the equation

15
$$Prog_j = \sum_{i=1}^{2000} \beta_{GY_i} G_{ij} + \lambda Inc_j + \gamma U_j + \epsilon_{P_j}$$

16 The values of λ and γ (Fig 2) were set as 0.1 and 1.5 respectively for all simulations, ensuring

17 that a collider bias effect is induced, and that stratification for binary kidney disease status does

18 not entirely adjust for the causal effect of continuous Inc on disease progression. GWAS of the

19 binary CKD incidence phenotype was then performed in the whole population (logistic

regression), and GWAS of the CKD progression phenotype was performed in only those

simulated participants with incident CKD. Simulations were repeated with differing values of ρ.

We then applied Dudbridge's method, the Slope-Hunter method, and an adapted version of the

MR-Horse method to these GWAS results. For both Dudbridge and Slope-Hunter methods, only

variants associated with disease incidence (we employed a threshold of $p < 0.001$) were used

to estimate b. The Hedges-Olkin approximation was used to adjust Dudbridge estimates for

measurement error, as the alternative simulation extrapolation methods performed

unpredictably at extremes of genetic correlation. Confidence intervals for the Slope Hunter

estimates were bootstrapped over 200 iterations. The prior probabilities for the MR-Horse

model parameters were informed by knowledge of chronic kidney disease. The prior probability

for ρ was $2r-1$ where $r \sim Beta(3,1)$. This prior puts most of the probability weighting around

high positive correlation but allows for some lower and even negative correlation for some

variants. The prior probability distribution for the global shrinkage parameter τ was set as $\tau \sim$

$Cauchy^+(0,1)$, and the prior for the local shrinkage prior ϕ_i was $\phi_i \sim Cauchy^+(0,1)$.

Uninformative priors were used for $b \sim Uniform(-10,10)$, $\mu \sim N(0,1)$ and $s_x^2 \sim N^+(0,1)$.

Posterior parameter estimates were drawn from four parallel Markov Chains simulated in JAGS

using 15,000 iterations with a 20% burn-in. Power, type 1 error and bias were recorded across

1,000 simulation iterations. For better comparison of method performance, power was

measured across the entire range of simulated variant effect sizes. Type 1 error was defined as

the identification of a significant association of a G_I SNP with disease progression. Significant

effects were defined at $p < 0.05$ for the regression based methods, and if the 95% equal-tailed

credible interval for the MR-Horse estimate of β_{GY_i} did not include 0. Bias was defined as

$\left|\hat{\beta}_{GY_i} - \beta_{GY_i}\right|$ for each method.

Sensitivity analyses were performed including simulated populations of different sizes, different

relative cluster sizes (testing relative performances of the adjustment methods when

assumptions of the other methods are met), and sparser or less sparse MR-Horse prior

distributions for β_{GY_i} (Supplemental Data).

All analyses were performed in R 4.3.1 and JAGS 4.3.2 on the Oxford University Medical

Sciences Division Biomedical Research Computing Cluster.

Application to CKDGen data

To illustrate the performance of our method in a real dataset we used data from the largest

extant meta-analyses of GWAS of kidney disease traits performed by the CKDGen consortium.

We used summary statistics from two GWAS: (i) a GWAS of creatinine-based eGFR from Wuttke

et al.[15] and (ii) a GWAS of eGFR decline over time, adjusted for baseline eGFR from Gorski et

al.[13]. As pointed out by the authors, adjustment for eGFR in GWAS (ii) should introduce a

collider bias effect and spurious associations between genetic variants associated with cross-

sectional eGFR alone and eGFR decline. Notably this scenario of adjustment for a baseline

variable is different to our simulated scenario of stratification by incident disease, but the use

of creatinine-based eGFR measurements as the phenotype of interest has the illustrative

advantage that genetic variants with a known functional role in creatinine production (or

associated only with creatinine-based measurements of kidney function and not with cystatin-c

or urea-based measurements) are unlikely to have true causal associations with kidney function

decline over time. Associations with such variants in GWAS (ii) are likely to be due to bias and

should be attenuated by any adjustment method which successfully reduces collider bias, and

they therefore can be used as a form of negative control for any method purporting to reduce

collider bias. GWAS (ii) detected genome-wide significant effects at such loci: *GATM*, *CPS1*, and

SHROOM3. Conversely, associations with genetic variants at loci with probable true effects on

kidney function decline over time (associated with multiple measures of kidney function or

known implication of nearby genes in Mendelian kidney disease) should be relatively preserved.

We identified a set of independent genetic variants in the Gorski et al. summary statistics using

the PLINK 1.9 "clump" algorithm[16], using the p-value of cross-sectional eGFR associations from

GWAS (i) as the clumping parameter with a clumping window of 100 kilobases and an r^2

threshold of 0.001.

The MR-Horse model was specified with the same structure and prior probability distributions

as above (ie prior probability for ρ placing most probability mass around a correlation between

cross-sectional eGFR lowering effects, and faster eGFR decline), and posterior parameter

estimates were drawn from four parallel Markov chains with 15,000 iterations and a burn in of

20%. In order to account for the fact that variants were selected for cross-sectional eGFR

associations, and that the majority of such variants may have true associations with kidney

function decline rendering a sparse horseshoe prior inappropriate, we performed a sensitivity

analysis with less and more sparse prior probability distributions for β_{GY} (Supplemental data).

Given that the true genetic architecture of many disease traits of interest is often unknown,

sensitivity analyses were also performed using larger and smaller clumping windows for variant

selection, and using deliberately mis-specified prior distributions for ρ (Supplemental data).

1 Results

2 Simulations

Power for different adjustment methods by different simulated scenarios are shown in Figure

3a. The adapted MR-Horse model had less power to detect small variant effect sizes than

unadjusted or Dudbridge method across the range of genetic correlations, but had greater

power than the Slope-Hunter method when significant correlation was present. At the sample

size tested, the power to detect variants of explaining ~ 0.5% of the variance of the progression

phenotype was 80-90% for the adapted MR Horse method across the range of genetic

correlations. This range of effect size is relevant to kidney function phenotypes, with variants at

the *UMOD* locus explaining ~ 0.3 - 0.5% of the variance of eGFR decline[13].

A collider bias effect was apparent in unadjusted analyses among variants with effects on

disease incidence (G_I and G_{IP} variants), and was strongest when genetic correlation was

strongly positive (ie in the same direction as confounder effects γ), and weakest when genetic

correlation was negative (Fig 3b; unadjusted analyses). In the setting of no or little genetic

correlation, this bias was reduced by the Dudbridge method, however the bias was markedly

increased by both Dudbridge and Slope-Hunter methods in the setting of strong genetic

correlation (Fig 3b). In our simulation set-up the Slope-Hunter method increased the bias of G_I

and G_{IP} variants across the range of genetic correlations, likely due to ZeMRA violations. The

adapted MR-Horse method reduced the bias of G_I and G_{IP} variants across the range of genetic

correlations (Fig 3b).

The adapted MR-Horse consistently had lower Type 1 Error rates below the nominal (5%) level

and below the other methods across the range of genetic correlations. This observation reflects

the propensity of the method to shrink null effects to zero with a high precision whilst allowing

non-zero effects to escape shrinkage.

Analyses with different sample sizes demonstrated a greater sensitivity of the power of the

adapted MR-Horse method to sample size changes, with a larger proportional increase and

drop-off in power with changes in sample size than the other methods (Supplemental figures 1-

2).

Changes in the relative sizes of G_I, G_{IP} and G_P clusters had significant effects on the

performance of the Slope-Hunter and Dudbridge methods, with both methods acheiving

comparable reductions in bias, and higher power than the adapted MR-Horse when the

pleiotropic cluster was smaller (and explaining less phenotypic variance) than the pleiotropic

cluster. When cluster sizes were equal, the adapted MR-Horse achieved a superior reduction in

bias and type 1 error in the setting of high genetic correlation (Supplemental figures 3-4).

Relative performance of the adjustment methods was not significantly altered by changes in

the parameters of the prior probability distributions for τ (Supplemental figures 5-6).

CKDGen data

Clumping yielded 531 independent genetic variants with available data in both GWAS (i) [GWAS

of cross-sectional eGFR] and (ii) [GWAS of eGFR decline adjusted for baseline eGFR]. This

pruning process removed ten of the lead variants from GWAS (ii) as they were not lead variants

in GWAS (i); these variants were added back to the dataset giving a total set of 541 SNPs for use

in the Dudbridge, Slope-Hunter and MR-Horse methods.

Application of the adapted MR-Horse method to summary statistics from the CKDGen GWAS of

eGFR decline adjusted for baseline eGFR attenuated the magnitude of the estimates at all loci

identified as significantly associated with eGFR decline (Fig 4). Effect estimates for variants that

were identified as likely to have a true effect on kidney function decline (notably at the *UMOD*

locus) remained significantly associated with eGFR decline after MR-Horse adjustment (ie the

95% credible interval for β_{GY} did not include 0). For rs77924615 (considered to be the causal

variant at the *UMOD-PDILT* locus[17]), the effect of one additional A allele on eGFR decline

adjusted for baseline eGFR was 0.092 mls/min/year (95% confidence interval 0.081 - 0.103);

adjusted using the adapted MR-Horse this was attenuated to 0.067 mls/min/year (equal-tailed

95% credible interval: 0.056 - 0.078). Associations at loci likely to have biased associations with

eGFR decline (*GATM*, *CPS1* and *SHROOM3*) were not significantly associated with kidney

function decline after MR-Horse adjustment. In the case of the *GATM* locus, where there is

good evidence that the association is mediated directly by effects on creatine metabolism

rather than kidney function[13], the baseline-eGFR adjusted result was 0.029 mls/min/year per

additional A allele (0.021 - 0.038); adjusted using the adapted MR-Horse model this was

attenuated to 0.002 mls/min/year (-0.001 - 0.010). Sensitivity analyses with less and more

sparse prior probabilities for β_{GY}, alterations in clumping parameters or deliberately mis-

specified prior distributions for ρ did not alter the interpretation of estimates at *GATM*, *CPS1*

and *SHROOM3* loci, although mis-specification of the probability priors did lead to likely true

associations being rejected at the *ACVR2B*, *OVOL1* and *TPPP* loci (Supplemental Figures 7 - 9).

1 Discussion

We set out to assess methods of studying the genetic associations of disease progression,

accounting for effects of selection bias. The presented analyses have shown that an adapted

version of the MR-Horse method appears to reliably reduce selection bias in GWAS of disease

progression in simulated studies of phenotypes with heritability and polygenicity values within

the range of values relevant to human complex traits[18]. When applied to real GWAS of kidney

disease traits, it significantly attenuates associations at loci where currently observed

associations are likely to be biased. Importantly, associations at loci with likely real effects are

largely preserved, although attenuated in magnitude.

The assumptions underlying the structure of the model used here are: (i) confounding is

constant across genetic variants, (ii) genetic effects are independent and additive and (iii) all

effects in Figure 2 are approximately linear. Additionally, it is important that the definitions of

disease incidence used in both GWAS of disease incidence and disease progression are as

similar as possible. These assumptions are less restrictive than those of the Dudbridge or Slope-

Hunter method, and notably there are no assumptions regarding the underlying correlation of

β_{GX} and β_{GY} effects. This is important because the genetic architectures of disease progression

traits are unknown a priori and methods that entail fewer assumptions are thus more useful for

accurate inference.

Violations of assumption (i) could however arise if there was a single locus or small set of loci

responsible for a large proportion of variance of either phenotype. This would mean that the

genetic component of U would vary depending on whether a variant being tested was at or in

LD with such a locus. This could be ameliorated by removing such a locus from the data, and b

could still be estimated from such an analysis. Violation of assumption (ii) is certainly possible,

but non-additive effects are likely a small contributor to heritability of complex traits[19].

Assumption (iii) is likely to hold when the effects concerned are small.

Results of this method will be to some extent influenced by the prior probability distributions

specified by the user. It could be argued that using a horseshoe prior (or alternative shrinkage

prior) for the β_{GY_i} effects in our analysis is inappropriate if we consider pleiotropy to be a major

characteristic of the genetic architecture of disease progression, and some work has shown that

posterior estimates can be very sensitive to hyperprior choice when using horseshoe priors[20].

We have shown in sensitivity analyses that results of our simulations and the CKDGen analyses

were not significantly changed despite large magnitude alterations of (i) the sparseness of the

horseshoe prior distribution and (ii) the prior probability for the correlation of genetic effects.

Using the horseshoe prior therefore retains sufficient power to detect important effects in real

GWAS, whilst biased effects are shrunk to zero. As shown in our data, this will reduce the

power to detect true effects of a small magnitude, but in the modern era of GWAS sample sizes

(where meta-analyses can involve millions of participants), exemplified by the suspected

achievement of "saturation" in GWAS of height[21], this is a limitation of decreasing importance.

A further limitation of the method is the fact that the bias reductions are small relative to

unadjusted analyses in the setting of no or negative genetic correlation (or in the more general

case, where the correlation of genetic effects is in the opposite direction to the correlation of

confounder effects responsible for the collider bias). The power reduction when using MR-

Horse in these analyses is however quite marked. It may be preferable therefore in situations

where correlation appears unlikely to use unadjusted estimates or alternative methods for

adjustment. For example in the case of cancer where incident disease is driven by cell

proliferation, but disease progression driven by invasion and metastasis, it is possible that the

biology and genetic architecture of disease progression and incidence are distinct[22].

Assumptions of no or low correlation may therefore be appropriate. However we emphasise

that in most cases the true underlying correlation structure is unknown, and the relatively

consistent performance of MR-Horse across the range of correlations renders it less vulnerable

to model mis-specification. Unadjusted analyses and Slope-Hunter and Dudbridge methods can

produce extreme biases when significant correlation is present.

There are also potential applications of this method to observational epidemiology more

broadly. The coefficient b in Figure 2, which defines the magnitude of any collider bias effect

and can be estimated from the MR-Horse model, is independent of the specific exposure of

interest and could be applied to reduce bias of estimated effects on disease progression in

other observational studies provided that "incident disease" and "progression" are defined in

the same manner as in the genetic study of interest. For example, if we wished to estimate the

effect of cigarette smoking on CKD progression (β_{SP}) in a cohort of participants with CKD, we

could estimate b from genetic association studies, and then the unbiased point estimate of the

effect of smoking on progression would be given by $\hat{\beta}_{SP}^* - b\hat{\beta}_{SI}$ where $\hat{\beta}_{SP}^*$ denotes the biased

estimate of β_{SP} obtained in an observational study among patients with CKD, and $\hat{\beta}_{SI}$ denotes

the estimated association of cigarette smoking with CKD incidence.

Unbiased estimates of the associations of genetic and environmental factors with disease

prognosis are crucial to understanding disease mechanisms and predicting disease risks. As

drug development pipelines become more reliant on genetic evidence for target selection[23], it

is increasingly important that the genetic evidence base is unbiased. This adaptation of the MR-

Horse method minimises selection bias, which is an under-addressed source of error in studies

of disease progression.

Bibliography

1   Galesloot TE, Grotenhuis AJ, Kolev D, Aben KK, Bryan RT, Catto JWF *et al.* Genome-wide
Meta-analysis Identifies Novel Genes Associated with Recurrence and Progression in Non-
muscle-invasive Bladder Cancer. *European Urology Oncology* 2022; **5**: 70–83.

2   Fogh I, Lin K, Tiloca C, Rooney J, Gellera C, Diekstra FP *et al.* Association of a Locus in the
CAMTA1 Gene With Survival in Patients With Sporadic Amyotrophic Lateral Sclerosis. *JAMA
neurology* 2016; **73**: 812–820.

3   Chang I-S, Jiang SS, Yang JC-H, Su W-C, Chien L-H, Hsiao C-F *et al.* Genetic Modifiers of
Progression-Free Survival in Never-Smoking Lung Adenocarcinoma Patients Treated with First-
Line Tyrosine Kinase Inhibitors. *American Journal of Respiratory and Critical Care Medicine*
2017; **195**: 663–673.

4   Hernán MA, Hernández-Díaz S, Robins JM. A Structural Approach to Selection Bias.
Epidemiology 2004; **15**: 615.

5   Dahabreh IJ, Kent DM. Index event bias: An explanation for the paradoxes of recurrence
risk research. *JAMA : The Journal of the American Medical Association* 2011; **305**: 822–823.

6   Mitchell RE, Hartley AE, Walker VM, Gkatzionis A, Yarmolinsky J, Bell JA *et al.* Strategies
to investigate and mitigate collider bias in genetic and Mendelian randomisation studies of
disease progression. *PLoS genetics* 2023; **19**: e1010596.

7   Dudbridge F, Allen RJ, Sheehan NA, Schmidt AF, Lee JC, Jenkins RG *et al.* Adjustment for
index event bias in genome-wide association studies of subsequent events. *Nature
Communications* 2019; **10**: 1561.

8   Cai S, Hartley A, Mahmoud O, Tilling K, Dudbridge F. Adjusting for collider bias in genetic
association studies using instrumental variable methods. *Genetic Epidemiology* 2022; **46**: 303–
316.

Robinson-Cohen C, Triozzi JL, Rowan B, He J, Chen HC, Zheng NS *et al.* Genome-Wide Association Study of CKD Progression. *Journal of the American Society of Nephrology: JASN* 2023; **34**: 1547–1559.

Mahmoud O, Dudbridge F, Davey Smith G, Munafo M, Tilling K. A robust method for collider bias correction in conditional genome-wide association studies. *Nature Communications* 2022; **13**: 619.

Grant AJ, Burgess S. A Bayesian approach to Mendelian randomization using summary statistics in the univariable and multivariable settings with correlated pleiotropy. *The American Journal of Human Genetics* 2024; **111**: 165–180.

Carvalho CM, Polson NG, Scott JG. The horseshoe estimator for sparse signals. *Biometrika* 2010; **97**: 465–480.

Gorski M, Rasheed H, Teumer A, Thomas LF, Graham SE, Sveinbjornsson G *et al.* Genetic loci and prioritization of genes for kidney function decline derived from a meta-analysis of 62 longitudinal genome-wide association studies. *Kidney International* 2022; **102**: 624–639.

Hardy GH. MENDELIAN PROPORTIONS IN A MIXED POPULATION. *Science (New York, NY)* 1908; **28**: 49–50.

Wuttke M, Li Y, Li M, Sieber KB, Feitosa MF, Gorski M *et al.* A catalog of genetic loci associated with kidney function from analyses of a million individuals. *Nature Genetics* 2019; **51**: 957–972.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics* 2007; **81**: 559–575.

Gorski M, Jung B, Li Y, Matias-Garcia PR, Wuttke M, Coassin S *et al.* Meta-analysis uncovers genome-wide significant variants for rapid kidney function decline. *Kidney international* 2021; **99**: 926–939.

Zeng J, de Vlaming R, Wu Y, Robinson Matthew R *et al.* Signatures of negative selection in the genetic architecture of human complex traits. *Nature Genetics* 2018; **50:** 746-753

Zhu Z, Bakshi A, Vinkhuyzen Anna AE, Hemani G, Lee S, Nolte Ilja M *et al.* Dominance genetic variation contributes little to the missing heritability for human complex traits. *American Journal of Human Genetics* 2015; **96**: 377–385.

Piironen J, Vehtari A. On the hyperprior choice for the global shrinkage parameter in the horseshoe prior. *Artificial Intelligence and Statistics* 2017; : 905–913.

Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S *et al.* A saturated map of common genetic variants associated with human height. *Nature* 2022; **610**: 704–712.

22   Chathrath A, Ratan A, Dutta A Germline variants that affect tumor progression. *Trends in*
Genetics 2021; **37**(5) 433 - 443

23   Minikel EV, Painter JL, Dong CC, Nelson MR. Refining the impact of genetic evidence on
clinical success. *Nature* 2024; **629**: 624–629.

Funding: KD is funded by a Kidney Research UK Clinical Training Fellowship TF_001_20220708.
The computational aspects of this research were supported by the Wellcome Trust Core Award
Grant Number 203141/Z/16/Z and the NIHR Oxford BRC. The views expressed are those of the
author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Figure 1: **Conditioning on incident disease leads to biased associations of genetic variants with disease progression.** 1A: Directed acyclic graph (DAG) representing genetic
effects on disease progression and incidence in the presence of confounders. 1B: Conditioning on incident disease (black rectangle) in GWAS of progression can induce biased
associations through the confounders U. (red dashed line). G: participant genotype, β_{GX}: effect of genetic variant on disease incidence, β_{GY}: direct effect of genetic variant on
disease progression, U: confounders of the incidence/progression relationship, λ: direct effect of disease incidence on disease progression

$$\beta_{GY} = \alpha + \theta\beta_{GX}$$ $$\beta_{GY}^* = \beta_{GY} + b\beta_{GX}$$

Figure 2: **Comparison of MR-Horse model of genetic pleiotropy in Mendelian Randomisation and its adaptation for mitigating collider bias in GWAS of disease progression.**
2A: DAG representing Mendelian Randomisation with pleiotropic variants, incorporating the notation used in original MR-Horse publication: G: participant genotype, β_{GX}: effect of genetic variant on disease exposure, α: direct (pleiotropic) effect of genetic variant on outcome, U: confounders of the incidence/progression relationship, θ: direct causal effect of exposure on outcome 2B: Adaptation of the DAG in 2A to illustrate the applicability of this model to the setting of selection bias in GWAS. λ: direct causal effect of disease incidence on disease progression, γ: effect of confounders on disease progression b: biasing pathway through U and γ, due to conditioning on Incidence (rectangle)

A   B

Figure 3: **The adapted MR-Horse method has lower bias and type 1 error rates across the range of genetic correlations, at the expense of a reduction in statistical power.**
Left: Power of unadjusted and adjusted analyses to detect β_{GY} effects across the range of effect sizes, by different values of ρ. Plotted values are mean power (%) across 1,000
simulation iterations with shaded areas +/- 1.96 * SEM. Phenotypic variance explained per variant defined as: $2\bar{\beta}^2 maf(1-maf)$. Right: Bias (top panel) and type 1 error rates
(bottom panel) of each method (mean bias of G_I and G_{IP} variants [variants with direct effects on incidence] across 1,000 simulation iterations). Unadjusted analyses have
increasing type 1 error rates with increasingly positive correlation. Dudbridge and Slope-Hunter analyses have increasing bias and type 1 error at extremes of correlation. MR-
Horse retains control of type 1 error across range of correlation.

1 **A**   **B**

Locus	Lead Variant	EA/OA	Unadjusted:	Effect on eGFR slope (mls/min/1.73²/year; 95% CI)
Likely true effect in unadjusted GWAS				
UMOD	rs34882080	A/G		0.092 (0.081 to 0.103)*
UMOD	rs77924615	A/G		-0.099 (-0.110 to -0.088)*
PRKAG2	rs10254101	T/C		0.037 (0.027 to 0.046)*
SPATA7	rs1028455	A/T		-0.024 (-0.033 to -0.016)*
FGF5	rs1458038	T/C		-0.028 (-0.037 to -0.019)*
OVOL1	rs4930319	C/G		0.028 (0.019 to 0.036)*
TPPP	rs434215	A/G		0.033 (0.022 to 0.044)*
C15ORF54	rs28857283	A/G		-0.029 (-0.038 to -0.021)*
ACVR2B	rs13095391	A/C		0.025 (0.016 to 0.034)*
Likely biased effect in unadjusted GWAS				
GATM	rs2453533	A/C		0.029 (0.021 to 0.038)*
CPS1	rs1047891	A/G		0.029 (0.020 to 0.039)*
SHROOM3	rs9998485	A/C		0.027 (0.017 to 0.036)*

*Genome-wide significant result in unadjusted GWAS.
^MR-Horse 95% credible interval does not include 0.

-0.1 -0.05   0.05 0.1

MR-Horse Adjusted	Effect on eGFR slope (mls/min/1.73m²/year; 95% CI)
	0.067 (0.056 to 0.078)^
	-0.072 (-0.084 to -0.061)^
	0.015 (0.002 to 0.026)^
	-0.016 (-0.026 to -0.003)^
	-0.017 (-0.027 to -0.005)^
	0.016 (0.004 to 0.026)^
	0.023 (0.002 to 0.036)^
	-0.023 (-0.032 to -0.014)^
	0.013 (0.001 to 0.024)^
	0.002 (-0.001 to 0.010)
	0.005 (0.000 to 0.017)
	0.003 (0.000 to 0.013)

-0.1 -0.05   0.05 0.1

3 *Figure 4:* **Unadjusted and MR-Horse adjusted associations of variants with eGFR decline, adjusted for baseline eGFR (adapted from CKDGen analyses).** 1A: Associations of lead
4 variants in Gorski et al.[13] with eGFR decline, adjusted for baseline eGFR. Variants in top panel have likely true associations with eGFR-decline (associated with decline in analyses
5 not adjusted for baseline eGFR). Variants in bottom panel have observed effects likely due to bias (not associated with eGFR decline in unadjusted analyses, and in the case of
6 *GATM* and *CPS1*, likely related to creatinine metabolism directly rather than kidney function). 1B: Results after adjustment using adapted MR-Horse method. *Genome-wide
7 significant result in GWAS adjusted for baseline eGFR. ^Significant result using MR-Horse method (95% credible interval does not include zero).