Dads on duty: First account of nest sitting in barnacle ganders

Isabella B. R. Scheiber * [1], Annabel J. Slettenhaar [2,3], M. J. J. E. Loonen [3] and Margje E.

de Jong [1,3]

*corresponding author

[1] Dept. of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030

Vienna, Austria

[2] Faculty of Biosciences and Aquaculture, Nord University, Kongens Gate 42, 7713

Steinkjer, Norway

[3] Arctic Centre, Aweg 30, 9718 CW Groningen, The Netherlands

Orcid IDs and E-Mail Addresses:

Isabella B. R. Scheiber: 0000-0003-3438-1836; isabella.scheiber@univie.ac.at

Annabel J. Slettenhaar: 0009-0005-9954-0721; annabel.j.slettenhaar@nord.no

Maarten J. J. E. Loonen: 0000-0002-3426-4595; m.j.j.e.loonen@rug.nl

Margje E. de Jong: 0000-0002-4941-8323; me.d.jong@gmail.com

Conflict of Interest Statement:

The authors declare no conflict of interest.

Data Availability Statement:

The data that support the findings of this study are provided within the article and

corresponding supplemental material.

26
Abstract
In most Anseriformes (ducks, geese and swans) only females are known to incubate. The few
exceptions where males have been described to incubate regularly are several species of
swans (Cygninae) and whistling ducks (Dendrocygninae). Here we describe, never
investigated quantitatively in any goose species, indicidents of male nest sitting in barnacle
geese (*Branta leucopsis*) as a form of paternal care of nest attendance. Based on pictures from
wildlife cameras we identified males, which sat on their nests when their mates took
incubation recesses. Wildlife cameras were placed at nests of which either the male or female
was fitted with a GPS neck collar in the year prior to this study. To attach transmitters, some
geese were caught while defending their nests, thus we may have unintentionally selected
bolder males which defended their nests more aggressively and were easier to catch. Nest
sitting occurred relatively frequent, *i.e.* in 6/15 individuals. Our results show that males with
collars were more likely to nest sit, but this does not deflect from the fact this behaviour is
novel in geese. Nest sitting may even be more common as collared birds are more easily
recognised, but also uncollared males have been observed to nest sit. We discuss several
possible functions of this behaviour, *i.e.* against raiding of nests by aerial predators, thermal
control of nest temperature, and intraspecific brood parasitism. At this time we cannot
demonstrate a possible function, as chances of successful hatching were not increased in nests
of males, which were nest sitting and we lack sample size for more in depth analyses. Lastly,
we discuss whether 'male incubation' is misleading in the waterfowl literature, as it is truly
justified for only two species, the black swan and black-bellied whistling duck.
48
49 **Keywords:**
50 *Branta leucopsis*, geese, waterfowl, paternal care, male incubation, incubation behaviour
51

1 INTRODUCTION

Nest defence by parents is considered a risky behaviour, as they may endure injury or even

death when trying to ward off predators (de Jong *et al.,* 2021), thus the risk of clutch

protection has to be gauged against its benefits (Ringelman & Stupaczuk, 2013). In birds,

there is large variation in how much males and females contribute to parental care and in

more than half of all species, males participate in incubation (Deeming & Reynolds, 2015). In

most waterfowl species, *i.e.* swans, geese and ducks, however, only females incubate. '*Male*

incubation' was only mentioned anecdotally once in geese (Anserinae), *i.e.* in the pink-footed

goose (*Anser brachyrhynchus*) (Inglis, 1977). Therefore, the purpose of this paper is

threefold: First, we revive the term '*nest sitting*' (Henson & Cooper, 1992) and describe it for

the first time in barnacle ganders (*Branta leucopsis*). Nest sitting refers to cases, where the

non-incubating parent is present *directly on the nest* and is different from other forms of nest

attendance, *i.e.* vigilance and/ or active defence, which are performed by ganders in close

proximity to the nest. Furthermore, nest sitting differs from incubation, which entails the

transfer of heat to the developing eggs (Wang & Beissinger, 2011). Besides, nest sitting

should also not be mixed up with the term '*baby-sitting*' in mammals, as here paternal care

may only occur *after* the females give *birth*. Secondly, we will discuss possible functions of

nest sitting in barnacle geese, and, finally, we question whether previously reported studies in

waterfowl, which refer to 'male incubation', are actually misinterpreted.

Barnacle geese are an Arctic breeding species, well-known for nesting high on steep

mountain cliffs and islets as a protection against predators (Mitchell *et al.*, 1998). Our study

population breeds contemporaneously on islands with common eiders (*Somateria mollisma*),

glaucous gulls (*Larus hyperboreus*), Arctic (*Stercorarius parasiticus*) and great skuas (*S.*

skua), the latter three being the main avian predators of barnacle geese (Tombre & Erikstad,

1996; Prop *et al.*, 2015). Just as other geese (*e.g.* lesser snow goose, *A. caerulescens*

caerulescens; pink-footed goose*;* Canada goose, *Branta canadensis),* barnacle geese protect

their nests against avian predators through a high rate of nest attendance (Prop *et al.*, 1984;

Samelius & Alisauskas, 2001; Schreven *et al.,* 2021) and active defence (Dittami *et al.*, 1979;

Clermont *et al.*, 2019; de Jong *et al*., 2021; Speelman *et al.*, 2022).

Nests on islets seem relatively safe from terrestrial predators. Islets in the inner fjord,

however, can stay connected with the mainland via ice-bridges when there is a late break-up

of fjord ice. This allows Arctic foxes (*Vulpes lagopus*) not only to reach the islands, but may

also confine them to stay there, once the ice breaks, at the time when the geese breed (Tombre

et al., 1998). However, as the fjord ice has diminished in recent years (*e.g.* Gilg *et al.*, 2016;

Pavlova *et al.*, 2019), the risks of encountering foxes on the breeding islands has gone down

(Dørum, 2019). Polar bears (*Ursus maritimus*), which are known to prey on eggs and goslings

(Smith *et al.*, 2010), have increased substantially in numbers (Prop *et al.*, 2015; Andersen &

Aars, 2016). They now visit islands more and more during the breeding season to raid nests

(Fuglei *et al.*, 2003; Hanssen *et al.*, 2013; Prop *et al.,* 2015). This results in complete nest

failures in some years (Prop *et al.*, 2015).

Barnacle geese are socially monogamous and biparental with precocial offspring where

parental roles before and after hatching are clearly defined. Before hatching, males normally

do not come straight up to the nest, but stay in close vicinity, maintaining contact with their

incubating mates visually and vocally (Prop *et al.*, 1984; Lorenz, 1991). In case of an intruder

coming close, most males actively protect and defend the nest and incubating female (de Jong

et al., 2021; Speelman *et al.*, 2022).

Just as in all other geese, only females incubate. During nest construction, they develop

thermosensitive brood patches, which guarantee an efficient heat transfer from the body to the

egg (Jones, 1971). Female geese actively pluck feathers, which they add as an insulating inner

layer of nest lining. They then begin to lay one egg per day until a clutch is completed with a

mean clutch size of about five eggs (Pär Forslund & Larsson, 1992). During this time,

barnacle ganders closely guard their receptive mates (Lamprecht, 1989). Incubation in

barnacle geese lasts for 25 days on average (Black *et al.,* 2014).

Incubation starts after the penultimate egg is laid (Hübner *et al.*, 2002). Females only take

brief incubation recesses for feeding and preening, as each time they leave the nest, the eggs

are at risk of dropping below an optimal temperature, possibly overheating, or being

discovered by a predator (Alsos *et al.*, 1998; Ahmad & Li, 2023). In our study population,

geese take five 20-min incubation recesses per day on average, yielding an incubation

constancy of 93% (Tombre *et al.*, 2012). Before leaving the nest, females use the down lining

of the nest feathers to cover the eggs for insulation and/or hide them from predators. Common

comprehension is that males either join their mates during incubation recesses, leaving the

nest unattended, or stay behind close to the nest (Owen & Wells, 1979). Here we report of

several cases, in which some barnacle ganders nest sit, while females are away.

2 METHODS

Our study population resides on the west coast of Spitsbergen, the largest island of the

Svalbard archipelago. In Kongsfjorden an established breeding colony nests on a group of

several islets (Tombre *et al.*, 1998). These islets are characterised by exposed ridges and flat

stretches of tundra (Tombre & Erikstad, 1996). Two are monitored in detail; the larger main

breeding islet, Storholmen (ca. 30 ha, 259 nests in 2021) and Prins Heinrichøya (ca. 3 ha, 29

nests in 2021) located offshore of the village Ny-Ålesund (78°55'N, 11°56'E), Svalbard,

Norway.

Within the scope of a larger study on circadian and circannual rhythmicity (de Jong *et al.*,

2024) we fitted 24 adults (½ males, ½ females of established pairs) in 2020 with solar-

powered GPS-GSM transmitters attached with neckbands (OrniTrack-NL40 3G, Ornitela,

UAB, Lithuania) in order to locate geese during migrations and on the wintering grounds.

Those 24 individuals were chosen following a suite of criteria, including that it would be

desirable if both pair partners were marked with unique colour rings already (for details of

criteria see supplemental material). Sexing of all individuals was performed when birds were

originally ringed by examining the cloaca for the presence of intromittent penises. To attach

transmitters, geese were caught in the vicinity of the nest either by hand, with a small hand-

net, or a fishing rod with an attached nylon snare ('nest catches', Fenstad *et al.,* 2017), or else

during the annually performed mass captures of moulting geese, where considerable numbers

of geese are funnelled into a trap ('moult catches', Loonen *et al.,* 1997). Only males, which

remained close to the nest when a human observer approached, could be caught by hand,

using the fishing rod or hand-net. We are aware that these methods might comprise a bias in

trappability *sensu* STRANGE (Webster & Rutz, 2020), as these males show high nest defence

and risk-taking behaviour (de Jong *et al*., 2024). From here on we consider them 'bold'.

In 2021, we were able to locate 15 nests of the initial 24 pairs, where either the male

(n=7) or female (n=8) was fitted with a transmitter in the previous year. During transmitter

attachment, four males were caught at the nest, while three were caught during the moult

catch. The neck collar allowed a definite recognition of pair partners on and close by the nest,

even when colour rings were not visible. In the following we refer to colour ring codes, when

we mention certain individuals.

Near the 15 nests, we set up wildlife cameras (Usogood TC30 Trail Camera) to get a

more detailed picture of behaviours of males and females during incubation. Cameras were

used in the past and do not seem to disturb the geese (*e.g*. de Jong *et al.*, 2019). At each nest,

we placed the camera, set in time-lapse mode, in one-metre distance of the nest to monitor

primarily the behaviour of the incubating female. Cameras took two pictures every five

minutes to detect any movement of the geese. The time in which cameras provided reliable

pictures varied between nests ($\bar{x}$ days $\pm$ SD: 12 $\pm$ 6.1; for details see supplement, Table S1,

see also de Jong *et al.*, 2024)

When scanning the photos, we noticed that in several instances males, rather than

females, were sitting on the nest (Figure 1). In the goose data base, first sporadic notes of this

behaviour stem from 2006 (M. E. de Jong & M. J. J. E. Loonen, pers. obs.). We thus decided

to investigate this phenomenon further. In particular, we wanted to determine how often nest

sitting of males occurred, whether only certain individuals did so, if male age played a role,

and whether individuals, which performed nest sitting did so multiple times or only once. As

successive photos in a series are contingent, we only counted male nest sitting bouts as

independent if the female returned to the nest and incubated the clutch for a pronounced

period of time. As occurrence of male nest sitting might have been a consequence of our

selection of males (Webster & Rutz, 2020), we also took the method of catching into account.

Finally, we investigated if eggs, were more likely to hatch if males were nest sitting.

2.1 Statistical Analysis

Using the R programming environment (R Core Team, 2021), we (1) applied a Student's *t*-

test to determine whether male age influenced the propensity to nest sit. We applied Fisher's

Exact Tests and give odds ratios and their 95% confidence intervals to determine, (2) if males,

which sat on the nest, were more likely to have been fitted with transmitters, (3) whether the

method of catching, *i.e.* snare/net vs. moult catch, was correlated with male nest sitting, and

(4) if male nest sitting was associated with hatching probability of the clutch, *i.e.* at least one

egg hatched (for details on counts of the three queries see 2 x 2 contingency tables,

supplement Table S2).

3 RESULTS

Observations of the 15 nests where cameras were placed revealed that six males sat on the

nest, whereas nine did not (Table 1). Age of males had no effect on nest sitting [$n_{males\ not\ nest}$

$_{sitting}$ = 9, $\bar{x}$ years $\pm$ SD: 8.33 $\pm$ 3.16; $n_{males\ nest\ sitting}$ = 6, $\bar{x}$ years $\pm$ SD: 11.00 $\pm$ 3.1; $t_{(13)}$ =

1.612, p = 0.131, Figure 2a)]. Of the six nest sitting males, three were observed on the nest

only once (gZUN, oFIJ, oFIA), whereas the others four males were observed multiple times

(oFYJ and yBYI four times, oFZY six times). Males with neck collars were more likely to sit

on nests than males without [Fisher's Exact Test: p = 0.041, $OR_{2\text{-sided}}$ = 13.59 with 95% CI of

[0.86, 934.01], Figure 2b)]. The likelihood of males sitting on the nest was independent of

whether they were caught near the nest during incubation or during the moult catches

[Fisher's Exact Test: p = 0.143, $OR_{2\text{-sided}}$ = 0.00 with 95% CI of [0.00, 3,53], Figure 2c)].

Whether males nest sat had no effect on whether eggs hatched or else were abandoned/

depredated [Fisher's Exact Test: p = 0.329, $OR_{2\text{-sided}}$ = 0.3 with 95% CI of [0.02, 4.30], Figure

2d), see Table S2 for 2 x 2 contingency tables of Fisher's Exact Tests]. Males took on average

12 minutes (SD ± 17 min, range 0 – 42 min) before they started nest sitting and they

averagely stayed for 39 minutes (SD = 21 min, range = 3 – 32 min). This corresponds to

about 60% (SD = ± 27%) of the time the female was gone. After males terminated nest

sitting, it took approximately 17 minutes (SD ± 13 min, range = 3 – 32 min) for the female to

return to the nest (for details see Table S3). Based on examinations of three nests, where we

had more than five days of reliable camera observations, male nest sitting occurred between

~1 – 4 % of the female nest recesses (Table S4). However, throughout this study sample size

is small and these results should be taken with caution.

4 DISCUSSION AND CONCLUSION

4.1 Nest sitting in barnacle ganders – possible functions

To the best of our knowledge, this is the first solid account of males sitting on nests in any

goose species. Although we may have introduced a bias to catch predominantly bold

individuals (Webster & Rutz, 2020), this does not alter the fact that some ganders perform

this behaviour. We oppose to calling this behaviour 'male incubation', as termed repeatedly in

the waterfowl literature (*e.g.* Rollin, 1957; Bruggers, 1979; Hawkins, 1986; Brugger &

Taborsky, 1994), because there is no active transfer of heat from the male to the eggs as

ganders do not form brood patches. Rather, we revive the term 'nest sitting' (Henson &

Cooper, 1992) as the appropriate label. We identified from the photos that male posture on the

nest was somewhat different from the female posture during incubation; *i.e.* males sat higher

than females; not in the nest bowl but rather on top of the nest. Female ducks and geese

sometimes adopt such as posture not only before actual incubation starts, but also during

incubation (Lorenz, 1991; Poussart *et al.*, 2000; Hartman *et al.*, 2023). The posture is

suggested to shield the nest and eggs from *e.g.* predation, inclement weather and/or

intraspecific brood parasitism, and we propose that male nest sitting has equivalent functions

when the female is absent. To date, we cannot provide a definite explanation of why this

behaviour occurs, as neither the age of males seems to play a role, nor could we show an

increase in hatching success of nests, where males nest sat. At this moment, we lack sample

size for other more in depth analyses, such as when nest sitting occurs over the incubation

period or to what extent males may compensate for more or longer incubation recesses by

their mates. In the following we suggest three possible, not mutually exclusive, functions of

male nest sitting, to initiate future studies not only in barnacle geese, but also other species.

The primary cause of hatching failure in ground nesting waterfowl is predation, thus

several different strategies are employed for nest protection (Peterson *et al*., 2022). These

range from high nest attendance (Samelius & Alisauskas, 2001), selecting sites where nests

can either be successfully concealed or are difficult for predators to reach (Schranck, 1972),

breeding in close proximity to more aggressive species to take advantage of their nest defence

(Quinn *et al*., 2003) to actively put predators to rout (Samelius & Alisauskas, 2001). Barnacle

geese historically nested high up on steep rock faces, particularly suitable to avoid terrestrial

predators. There, however, food is scarce for incubating females (Prop *et al*., 1984). Breeding

islands off the coast used to be fairly safe from polar bears and Arctic foxes, but this turns out

to now also be changing (Tombre *et al.*, 1998; Andersen & Aars, 2016). Furthermore,

breeding on islands forces geese to nest in close proximity to their main aerial predators, thus

nest sitting of barnacle ganders might function in shielding the nest from them. This is backed

by our choosing the most aggressive males when attaching transmitters, which might be bold

enough to attack predators actively. What needs to be investigated, is whether males adjust

their behaviour to certain environmental conditions, *i.e.* through behavioural plasticity, or

whether nest sitting confers fitness benefits as an adaptive response to breeding on islands

(Mery & Burns, 2010; Madsen *et al.,* 2019). In the future, attention should be paid to how

often nest sitting of males occurs, how long the incubation recesses of the female last, what

the predation pressure is, and at which stage during incubation the behaviour is performed.

Another possible function of nest sitting by males is to control the temperature in the nest.

To some extent, nest insulation can reduce both the degree of egg cooling when the female is

absent and the time spent to rewarm the eggs after she returns from an incubation recess

(Thompson & Raveling, 1988). Avoidance of egg cooling is particularly important for Arctic

breeding birds, where the difference between optimal temperature for developing embryos (in

geese: 37-38°C) and the average air temperature during incubation (Ny-Ålesund average air

temperature 2021: June 2.6°C, July 5.5°C, Norwegian Meteorological Institute) is striking

(Poussart *et al*., 2001). Thus, eggs should be covered more or less constantly. Indeed, Arctic

geese adjust their incubation behaviour in response to prevailing weather conditions, such as

wind velocity, air temperature and solar radiation (Harvey, 1971; Poussart *et al.,* 2001),

remaining on the nest more continuously during inclement weather (Harvey, 1971; Elkins,

2004). Yet, egg cooling in the absence of the female might be overrated. In greater snow

geese, egg temperature throughout incubation remained high with only slight drops during the

frequent incubation recesses of females (Poussart *et al.,* 2000). This might be, in part,

ascribed to the female's covering the nest with down (Thompson & Raveling, 1988). Even

under unfavourable weather conditions, egg temperature drops during incubation recesses are

probably nowhere near the minimal temperatures, which endanger an embryo's development

or survival. Rather, this will only lengthen the incubation period until the eggs hatch (White

& Kinney, 1974; Tombre & Erikstad, 1996).

Overheating of unprotected eggs in the nest might become an issue in the Arctic in the

future. Also in Ny-Ålesund temperature has risen steadily since the 1970s (*e.g.* Wei *et al.*,

2015). During continuous incubation in climate control chambers, the critical temperature, at

which no embryos will survive, is 40.5°C (White & Kinney, 1974). Even if barnacle nests are

exposed to direct sun, they probably will, at present, not reach this temperature limit long

enough for embryos to die in the Arctic. Yet, common guillemots (*Uria aalge*), a high-

latitude seabird, already showed an increased probability of egg loss at higher temperatures in

a colony in the Baltic Sea (Olin *et al.*, 2024). Whether nest sitting in barnacle ganders occurs

predominantly on extremely warm or cold days could be subject to further investigations. In

sum, whereas nest sitting of barnacle ganders undoubtedly will assist in temperature control

of eggs in the nest, we suggest it to only play a minor role, at least here and now.

If male nest sitting occurs during the period of egg laying, another possible function

might be to prevent intraspecific brood parasitism, which frequently occurs in barnacle geese

(Black *et al*., 2014). When conspecific barnacle geese attempt to dump eggs, females apply

one of three strategies. They either lay eggs in unguarded nests, leave parasitic eggs in close

proximity to the nest for the host female to retrieve, or attempt to lay an egg into the nest

despite vigorous defence by the host female. In contrast, males let those instances happen,

behaving only mildly aggressive, if at all, towards a female intruder (Par Forslund & Larsson,

1995). This tameness of the male towards the parasitic female, when the host female is

present, leads us to suggest that during egg laying his propensity to nest sit in the absence of

his mate is probably low. We cannot answer this question adequately, however, because we

placed cameras not until the host female had started incubation, therefore lack photos from

the laying period.

A question is whether nest sitting of barnacle ganders is a behaviour that has been

overlooked in the past or is a response to a changing environment. Although adult males are

slightly bigger than the females, barnacle geese are monomorphic. It is difficult to determine

the sex of an individual, even if it is marked, as rings are not visible when a bird sits on the

nest. It is comprehensible that a bird on the nest was assumed to be female, just as the

literature suggests. The relatively frequent occurrence of male nest sitting, *i.e.* 6/15 in our

limited number of individuals, was unexpected to us. It is feasible that nest sitting in ganders

occurs frequently also in other goose species if one's attention is turned specifically to

cautiously noting the sex of the bird, which is sitting on the nest.

4.2 'Male incubation' in waterfowl?

Found throughout the literature is the general assumption that in swans, and to some extent

whistling ducks, deviate from the female only incubation pattern in waterfowl, as here 'males

incubate' (Scott, 1977; Hawkins, 1986; Brugger & Taborsky, 1994). But are these males

indeed incubating or is it a term that should be avoided, as is leads to a misconception when

compared with species, where incubation is truly shared (Deeming & Reynolds, 2015)?

In an early study, Hawkins already concluded that the well-developed male incubation

behaviour in tundra swans (*Cygnus columbianus columbianus*.) protected against egg

predation, provided some egg cooling benefits, but was not essential for successful embryo

development (Hawkins, 1986). This led Henson & Cooper to apply the term 'nest sitting' in

the case of trumpeter swans *(C. buccinator)*. They state that the behaviour was not 'as well

developed as in tundra swans', because it was independent of predator presence or adverse

weather conditions, despite males occasionally exhibiting some of the nest-settling motions

characteristic of incubating females (Henson & Cooper, 1992). Our extensive search of the

literature revealed that in most waterfowl, including three single observations in wood duck

(*Aix sponsa*), mandarin duck (*A. galericulata*) and fulvous whistling duck (*Dendrocygna*

bicolor, Rollin, 1957; Flickinger, 1975; Bruggers, 1979), males nest sit rather than incubate.

The two notable exceptions, for which the term incubation is warranted, are the black

swan (*C. atratus*, BLSW) and black-bellied whistling duck (*D. autumnalis*, BBWD*)*. Whereas

BLSW exhibit the Cygnidae typical long-term monogamy and biparental care, this is also true

for BBWD, unusual for Anatidae, although in some species males participate in attending

broods after hatching (McKinney & Brewer, 1989). Besides male incubation, there are some

features, which BLSW and BBWD share. In both species, neither females nor males develop

brood patches (Skutch, 1976; Bolen & Smith, 1979), thus either parent of each species can

incubate in a similar manner. Also, both species are known to re-nest after successful

breeding attempts (Delnicki, 1973; Bolen & Smith, 1979; Brugger & Taborsky, 1994; James

et al., 2012; Coleman, 2014), whereas most other waterfowl may attempt a second clutch only

if the first one failed particularly during early stages of incubation. We speculate that re-

nesting might be the reason of both parents incubating. Near constant incubation minimizes

the time it takes for eggs to develop and hatch, thereby possibly shortening the inter-clutch

interval, but this would need to be tested in mate removal experiments. In waterfowl with

uniparental female care, mate removal did not affect hatching success or body condition of

the female (LeSchack *et al.*, 1998; Manlove & Hepp, 1998).

In conclusion, this is the first report in any goose species, where males expand paternal

duties beyond the established view, which during the breeding period is to protect the nest and

female in close vicinity of the nest. We have pictorial evidence that several individuals

displayed nest sitting behaviour. Although we may have unintentionally selected bolder

individuals, this should not divert from the fact that the behaviour exists. Both predation and

thermal risks are increased for eggs, which are not protected by parents, thus minimising

those risks are the main causes for successful hatching. In barnacle geese, we suggest nest

sitting of males to be a response to avian predator presence. We cannot exclude that

preventing egg cooling (and overheating) in the absence of the female may also play a role,

but consider this unlikely. Our final point is that in most waterfowl species, males do not

participate in incubation. Nest sitting is an unbiased term and should be used for cases, in

which no active warming of eggs has (yet) been demonstrated.

Acknowledgements

We are grateful to Børge Moe (Norwegian Institute for Nature Research, NINA) as the

project administrator of AJS' field grant, the staff of the French - German Arctic Research

Base at Ny-Ålesund, chiefly Bettina Haupt (station leader AWI 2020/21), and KingsBay AS,

Ny-Ålesund for logistic support. We thank Jouke Prop for sharing his unpublished

observations and Kees Schreven for directing out attention to the one dated reference, where

nest sitting in ganders was briefly alluded to. Funding was provided by the Austrian Science

Fund (P32216 to I. B. R. Scheiber) and an Arctic Field Grant (The Research Council of

Norway) in collaboration with NINA (ES676286 - to A. J. Slettenhaar). The study conforms

to Directive 2010/63/EU and was conducted under FOTS ID 23358 from the Norwegian

Animal Research Authority and approved by the Governor of Svalbard (RIS ID 11237).

REFERENCES:

Ahmad, I. M., & Li, D. (2023). More than a simple egg: Underlying mechanisms of cold tolerance in avian embryos. *Avian Res., 14*, 100104. doi: 10.1016/j.avrs.2023.100104

Alsos, I. G., Elvebakk, A., & Gabrielsen, G. W. (1998). Vegetation exploitation by barnacle geese *Branta leucopsis* during incubation on Svalbard. *Polar Res., 17*, 1-14. doi: 10.3402/polar.v17i1.6603

Andersen, M., & Aars, J. (2016). Barents Sea polar bears (*Ursus maritimus*): Population biology and anthropogenic threats. *Polar Res., 35*(1), 26029. doi: 10.3402/polar.v35.26029

Black, J. M., Prop, J., & Larsson, K. (2014). *The Barnacle Goose*. London, UK: Bloomsbury Publishing.

Bolen, E. G., & Smith, E. N. (1979). Notes on the incubation behavior of black-bellied whistling ducks. *Prairie Nat., 11*(4), 119-123.

Brugger, C., & Taborsky, M. (1994). Male incubation and its effect on reproductive success in the black swan, *Cygnus atratus. Ethology, 96*(2), 138-146. doi: 10.1111/j.1439-0310.1994.tb00889.x

Bruggers, R. L. (1979). Nesting patterns of captive mandarin ducks. *Wildfowl, 30*, 45-54.

Clermont, J., Réale, D., & Giroux, J.-F. (2019). Similarity in nest defense intensity in Canada goose pairs. *Behav. Ecol. Sociobiol., 73*(8). doi: 10.1007/s00265-019-2719-3

Coleman, J. T. (2014). Breeding biology of the black swan *Cygnus atratus* in southeast Queensland, Australia. *Wildfowl, 64*, 217-230.

de Jong, M. E., Nicolaus, M., Fokkema, R. W., & Loonen, M. J. J. E. (2021). State dependence explains individual variation in nest defence behaviour in a long-lived bird. *J. Anim. Ecol., 90*(4), 809-819. doi: 10.1111/1365-2656.13411

de Jong, M. E., Slettenhaar, A. J., Fokkema, R. W., Leh, M., Barnreiter, E., Griffin, L., . . . Scheiber, I. B. R. (2024). Diel and seasonal rhythmicity in activity and corticosterone in an Arctic migratory herbivore: A multifaceted approach. *preprint*. doi: 10.1101/2024.08.30.610510

de Jong, M. E., Wetherbee, R., & Loonen, M. J. J. E. (2019). Effects of fleas on nest success of Arctic barnacle geese: Experimentally testing the mechanism. *J. Avian Biol., 50*(5), e01944. doi: 10.1111/jav.01944

Deeming, D. C., & Reynolds, S. J. (2015). *Nests, Eggs, and Incubation*. Oxford, UK: Oxford University Press.

Delnicki, D. E. (1973). *Renesting, incubation behavior, and compound clutches of the black-bellied tree duck in southern Texas*. MSc, Texas Tech University, Lubbock, TX, USA. (31295001778181)

Dittami, J. P., Kennedy, S., & Thomforde, C. (1979). Observations on barnacle goose
breeding, *Branta leucopsis*, in Spitsbergen 1975. *J. Ornithol., 120*, 188-195. doi:
10.1007/BF01642997

Dørum, H. (2019). *Effects of climate change and diminishing sea ice on common eider*
breeding distribution in Kongsfjorden, Svalbard. Master of Science Master, Norwegian
University of Science and Technology, Trondheim, Norway. Retrieved from
http://hdl.handle.net/11250/2617043

Elkins, N. (2004). Breeding: Incubation. In N. Elkins (Ed.), *Weather and Bird Behaviour*
(Vol. 3rd edition, pp. 101-104). London, UK: T.& A.D. Poyser.

Fenstad, A. A., Bustnes, J., Lierhagen, S., Gabrielsen, K. M., Öst, M., Jaatinen, K., . . .
Krøkje, Å. (2017). Blood and feather concentrations of toxic elements in a Baltic and an
Arctic seabird population. *Marine Pollution, 114*(2), 1152-1158. doi: j.marpolbul.2016.10.034

Flickinger, E. L. (1975). Incubation by a male fulvous tree duck. *Wilson Bull., 87*(1), 106-
107.

Forslund, P., & Larsson, K. (1992). Age-related reproductive success in the barnacle goose. *J.*
Anim. Ecol., 61(1), 195. doi: 10.2307/5522

Forslund, P., & Larsson, K. (1995). Intraspecific nest parasitism in the barnacle goose:
Behavioural tactics of parasites and hosts. *Anim. Behav., 50*, 509-517. doi:
10.1006/anbe.1995.0265

Fuglei, E., Øritsland, N. A., & Prestrud, P. (2003). Local variation in Arctic fox abundance on
Svalbard, Norway. *Polar Biol., 26*(2), 93-98. doi: 10.1007/s00300-002-0458-8

Gilg, O., Istomina, L., Heygster, G., Strøm, H., Gavrilo, M. V., Mallory, M. L., . . . Yannic,
G. (2016). Living on the edge of a shrinking habitat: The ivory gull, *Pagophila eburnea*, an
endangered sea-ice specialist. *Biol. Lett., 12*(11), 20160277. doi: 10.1098/rsbl.2016.0277

Hanssen, S. A., Moe, B., Bårdsen, B. J., Hanssen, F., & Gabrielsen, G. W. (2013). A natural
antipredation experiment: Predator control and reduced sea ice increases colony size in a
long-lived duck. *Ecol. Evol., 3*(10), 3554-3564. doi: 10.1002/ece3.735

Hartman, C. A., Ackerman, J. T., Peterson, S. H., Fettig, B., Casazza, M., & Herzog, M. P.
(2023). Nest attendance, incubation constancy, and onset of incubation in dabbling ducks.
PLOS One, 18(5), e0286151. doi: 10.1371/journal.pone.0286151

Harvey, J. M. (1971). Factors affecting blue goose nesting success. *Can. J. Zool., 49*(2), 223-
234. doi: 10.1139/z71-03

Hawkins, L. L. (1986). Nesting behaviour of male and female whistling swans and
implications of male incubation. *Wildfowl, 37*, 5-27.

Henson, P., & Cooper, J. A. (1992). Divison of labour in breeding trumpeter swans *Cygnus*
buccinator. Wildfowl, 43, 40-48.

Hübner, C. E., Tombre, I. M., & Erikstad, K. E. (2002). Adaptive aspects of intraclutch egg-
size variation in the high Arctic barnacle goose (*Branta leucopsis*). *Can. J. Zool., 80*, 1180-
1188. doi: 10.1139/z02-100

Inglis, I. R. (1977). The breeding behaviour of the pink-footed goose: Behavioural correlates
of nesting success. *Anim. Behav., 25*, 747-764. doi: 10.1016/0003-3472(77)90125-7

James, J. D., Thompson, J. E., & Ballard, B. M. (2012). Evidence of double brooding by
black-bellied whistling-ducks. *Wilson J. Ornithol., 124*(1), 183-185. doi: 10.1676/11-082.1

Jones, R. E. (1971). The incubation patch of birds. *Biol. Rev., 46*(3), 315-339. doi:
10.1111/j.1469-185X.1971.tb01048.x

Lamprecht, J. (1989). Mate guarding in geese: Awaiting female receptivity, protection of
paternity or support of female feeding. In A. E. Rasa, C. Vogel & E. Voland (Eds.), *The
Sociobiology of Sexual and Reproductive Strategies*. Dordrecht, NL: Springer Science and
Business Media.

LeSchack, C. R., Afton, A. D., & Alisauskas, R. T. (1998). Effects of male removal on female
reproductive biology in Ross' and lesser snow goose. *Wilson Bull., 110*(1), 56-64. doi:
http://www.jstor.org/stable/4163897

Loonen, M. J. J. E., Oosterbeek, K., & Drent, R. H. (1997). Variation in growth of young and
adult size barnacle geese *Branta leucopsis*: Evidence for density dependence. *Ardea, 85*, 177-
192.

Lorenz, K. (1991). *Here I Am - Where Are You?* New York, NY, USA: Hartcourt Brace
Jovanovich.

Madsen, J., Jaspers, C., Frikke, J., Gundersen, O. M., Nolet, B. A., Nolet, K., . . . de Vries, P.
P. (2019). A gloomy future for light-bellied brent geese in Tusenøyane, Svalbard, under a
changing predator regime. *Polar Res., 38*, 3393. doi: 10.33265/polar.v38.3393

Manlove, C. A., & Hepp, G. R. (1998). Effects of mate removal on incubation behavior and
reproductive success of female wood ducks. *Condor, 100*, 688-693. doi: 10.2307/1369750

McKinney, F., & Brewer, G. L. (1989). Parental attendance and brood care in four species of
Argentine dabbling ducks. *Condor, 91*, 131-138. doi: 10.2307/1368156

Mery, F., & Burns, J. G. (2010). Behavioural plasticity: An interaction between evolution and
experience. *Evol. Ecol., 24*(3), 571-583. doi: 10.1007/s10682-009-9336-y

Mitchell, C., Black, J. M., & Evans, M. (1998). Breeding success of cliff-nesting and island-
nesting barnacle geese in Svalbard. *Norsk Polar. Skrift., 200*, 141-146.

Olin, A. B., Dück, L., Berglund, P. A., Karlsson, E., Bohm, M., Olsson, O., & Hentati-
Sundberg, J. (2024). Breeding failures and reduced nest attendance in response to heat stress
in a high-latitude seabird. *Mar. Ecol. Prog. Ser., 737*, 147-160. doi: 10.3354/meps14244

Owen, M., & Wells, R. L. (1979). Territorial behaviour in breeding geese - A re-examination
of Ryder's hypothesis. *Wildfowl, 30*, 20-26.

Pavlova, O., Gerland, S., & Hop, H. (2019). Changes in sea-ice extent and thickness in
Kongsfjorden, Svalbard (2003-2016). In H. Hop & C. Wiencke (Eds.), *The Ecosystem of
Kongsfjorden, Svalbard. Advances in Polar Ecology* (Vol. 2, pp. 105-136): Springer.

Peterson, S. H., Ackerman, J. T., Keating, M. P., Schacter, C. R., Hartman, C. A., Casazza,
M. L., & Herzog, M. P. (2022). Predator movements in relation to habitat features reveal
vulnerability of duck nests to predation. *Ecol. Evol., 12*(9). doi: 10.1002/ece3.9329

Poussart, C., Gauthier, G., & Jacques, L. (2001). Incubation behaviour of greater snow geese
in relation to weather conditions. *Can. J. Zool., 79*, 671-678. doi: 10.1139/z01-023

Poussart, C., Larochelle, J., & Gauthier, G. (2000). The thermal regime of eggs during laying
and incubation in greater snow geese. *Condor, 102*(2), 292-300. doi:
10.1093/condor/102.2.292

Prop, J., Aars, J., Bårdson, B.-J., Hanssen, S. A., Bech, C., Bourgeon, S., . . . Moe, B. (2015).
Climate change and the increasing impact of polar bears on bird populations. *Frontier in
Ecology and Evolution, 3*, 33. doi: 10.3389/fevo.2015.00033

Prop, J., Eeerden, M. R., & van Drent, R. H. (1984). Reproductive success in the barnacle
goose *Branta leucopsis* in relation to food exploitation on the breeding grounds, western
Spitsbergen. *Norsk Polar. Skrift., 181*, 87-117.

Quinn, J. L., Prop, J., Kokorev, Y., & Black, J. M. (2003). Predator protection or similar
habitat selection in red-breasted goose nesting associations: Extremes along a continuum.
Anim. Behav., 65(2), 297-307. doi: 10.1006/anbe.2003.2063

R Core Team. (2021). R: A language and environment for statistical computing (Version
1.13). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from
https://www.r-project.org/.

Ringelman, K. M., & Stupaczuk, M. J. (2013). Dabbling ducks increase nest defense after
partial clutch loss. *Condor, 115*(2), 290-297. doi: 10.1525/cond.2013.120096

Rollin, N. (1957). Incubation by drake wood duck in eclipse plumage. *Condor, 59*(4), 263-
265. doi: 10.2307/1364656

Samelius, G., & Alisauskas, R. T. (2001). Deterring arctic fox predation: The role of parental
nest attendance by lesser snow geese. *Can. J. Zool., 79*, 861-866. doi: 10.1139/z01-048

Schranck, B. W. (1972). Waterfowl nest cover and some predation relationships. *J. Wildl.
Manag., 36*(1), 182. doi: 10.2307/3799210

Schreven, K. H. T., Stolz, C., Madsen, J., & Nolet, B. A. (2021). Nesting attempts and
success of Arctic-breeding geese can be derived with high precision from accelerometry and
GPS-tracking. *Anim. Biotelem., 9*(1), 25. doi: 10.1186/s40317-021-00249-9

Scott, D. (1977). Breeding behaviour of wild whistling swans. *Wildfowl, 28*, 101-106.

Skutch, A. F. (1976). *Parent Birds And Their Young*: University of Texas Press.

Smith, P. A., Elliott, K. H., Gaston, A. J., & Gilchrist, H. G. (2010). Has early ice clearance
increased predation on breeding birds by polar bears? *Polar Biol., 33*(8), 1149-1153. doi:
10.1007/s00300-010-0791-2

Speelman, F. J. D., Hammers, M., Komdeur, J., & Loonen, M. J. J. E. (2022). Nest defence
behaviour is similar between pair members but only male behaviour predicts nest survival in
barnacle geese. *J. Avian Biol., 2022*(9), e02982. doi: 10.1111/jav.02982

Thompson, S. C., & Raveling, D. G. (1988). Nest insulation and incubation constancy of
Arctic geese. *Wildfowl, 39*, 124-132.

Tombre, I. M., & Erikstad, E. (1996). An experimental study of incubation effort in high-
Arctic barnacle geese. *J. Anim. Ecol., 65*, 325-331. doi: 10.2307/5878

Tombre, I. M., Erikstad, E., & Bunes, V. (2012). State-dependent incubation behaviour in the
high Arctic barnacle geese *Polar Biol., 35*(7), 985-992. doi: 10.1007/s00300-011-1145-4

Tombre, I. M., Mehlum, F., & Loonen, M. J. J. E. (1998). The Kongsfjorden colony of
barnacle geese: Nest distribution and the use of breeding islands 1980-1997. *Norsk Polar.*
Skrift., 200, 57-65.

Wang, J. M., & Beissinger, S. R. (2011). Partial incubation in birds: Its occurrence, function,
and quantification. *The Auk, 128*(3), 454-466. doi: 10.1525/auk.2011.10208

Webster, M. M., & Rutz, C. (2020). How STRANGE are your study animals? *Nature,*
582(7812), 337-340. doi: 10.1038/d41586-020-01751-5

Wei, T., Ding, M., Wu, B., Lu, C., & Wang, S. (2015). Variations in temperature-related
extreme events (1975–2014) in Ny-Ålesund, Svalbard. *Atmos. Sci. Let., 17*(1), 102-108. doi:
10.1002/asl.632

White, F. N., & Kinney, J. L. (1974). Avian incubation: Interactions among behavior,
environment, nest, and eggs result in regulation of egg temperature. *Science, 186*(4159), 197-
115. doi: 10.1126/science.186.4159.1
522

524
525 **Table 1:** Identities of 15 focal pairs, whose nests were monitored with wildlife cameras in 2021. For each
526 individual, we provide codes of unique colour and aluminium rings, and if they were fitted with a transmitter, the
527 neck collar ID. Except two individuals, which were ringed as goslings (*), the minimum age is given ($\bar{\text{x}}$ age $\pm$
528 SD: males 9.4 ± 3.31; females 10.4 ± 4.46). Whether or not males sat on nests is also indicated.
529

Male colour ID	Male ID	Male neck collar ID	Age (years)	Male nest sitting	Female colour ID	Female ID	Female neck collar ID	Age (years)
oFYJ	CA46944	201913	16	yes	oFVD	CA46857		10
gZUN	CA45605	201915	9	yes	oFLC	CA45810		12
alur	CA47072	201917	8	no	yBSV	CA45997		4
oFZY	CA46942	201918	10	yes	gSNF	CA36931		19
oFIJ	CA45800	201924	12	yes	yBHJ	CA45891		5
oFIA	CA45795	201930	12	yes	gLDC	CA41238		14
yBID	CA46840	201932	7	no	yBIB	CA44228		14
oFZX	CA46941		10	no	gTTL	CA41113	201912	16*
yBYI	CA45746		7	yes	yAUF	CA45744	201919	7
yAXY	CA45703		8	no	yBHY	CA46848	201925	5*
gLHX			2	no	yAVV	CA45736	201926	7
yAZT	CA45715		8	no	oFXV	CA45632	201927	10
yAXC	CA45942		8	no	yAPH	CA45929	201931	8
gTCZ	CA46957		10	no	yAXV	CA45701	201934	8
yBYX	CA41397		14	no	yBYV	CA41400	201935	14

FIGURE LEGENDS

Figure 1: An example of one barnacle gander nest sitting. Photos taken on June 18th, 2021

between 15:48 and 16:28. Panel A) Female oFVD without a neck collar is incubating. Panel

B) The female left the nest. Panel C) Male alur with neck collar approaches the nest. Panel D)

He sits on the nest with the neck collar clearly visible. Panel E) The female returns to the nest.

Panel F) She resumes incubation. Temperature and time are shown in the lower right hand

corner of each photo.

Figure 2: Features, which might influence nest sitting in barnacle ganders. a) Age did not

affect the likelihood of male to nest sit (Student's *t*-test: $p > 0.05$). b) Males, which were fitted

with a neck collar, were more likely to nest sit than males without (Fisher's Exact Test: $p =$

0.041). c) The method of catching for transmitter deployment, either near the nest or the

during moult catch, had no effect on nest sitting propensity of males (Fisher's Exact Test: $p >$

0.05). d) Whether eggs hatched or did not hatch was independent of males' nest sitting

(Fisher's Exact Test: $p > 0.05$).

549
550   Figure 1
551

553
554   Figure 2