Dads on duty: First account of nest sitting in barnacle ganders

Isabella B. R. Scheiber * [1], Annabel J. Slettenhaar [2,3], M. J. J. E. Loonen [3] and Margje E.

de Jong [1,3]

*Corresponding author

[1] Dept. of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030

Vienna, Austria

[2] Faculty of Biosciences and Aquaculture, Nord University, Kongens Gate 42, 7713

Steinkjer, Norway

[3] Arctic Centre, Aweg 30, 9718 CW Groningen, The Netherlands

Orcid IDs and E-Mail Addresses:

Isabella B. R. Scheiber: 0000-0003-3438-1836; isabella.scheiber@univie.ac.at

Annabel J. Slettenhaar: 0009-0005-9954-0721; annabel.j.slettenhaar@nord.no

Maarten J. J. E. Loonen: 0000-0002-3426-4595; m.j.j.e.loonen@rug.nl

Margje E. de Jong: 0000-0002-4941-8323; me.d.jong@gmail.com

Conflict of Interest Statement:

The authors declare no conflict of interest.

Data Availability Statement:

The data that support the findings of this study are provided within the article and

corresponding supplemental material.

Abstract

In most Anseriformes (ducks, geese and swans) only females are known to incubate. Here we

scientifically describe, indicidents of male nest sitting in barnacle geese (*Branta leucopsis*) as

a form of paternal care of nest attendance. Based on pictures from wildlife cameras we

identified males, which sat on their nests when their mates took incubation recesses. Wildlife

cameras were placed at nests of which either the male or female was fitted with a GPS neck

collar in the year prior, which aided with identifying individual birds on the nest. To attach

transmitters, some geese were caught while defending their nests, thus we may have

unintentionally selected bolder males which defended their nests more aggressively and were

easier to catch. Nest sitting occurred relatively frequent, *i.e.* in 6/15 individuals. Our results

show that males with collars were more likely to nest sit, but this does not deflect from the

fact this behaviour exists in geese. In the course of this finding we discuss several possible

functions of this behaviour, *i.e.* against raiding of nests by aerial predators, thermal control of

nest temperature, and intraspecific brood parasitism. At this time we cannot demonstrate a

possible function, as chances of successful hatching were not increased in nest-sitting males

and we lack sample size for more in depth analyses. Lastly, we argue that 'male incubation' is

misleading in the waterfowl literature, as it is truly justified for only two species, the black

swan (*Cygnus atratus)* and black-bellied whistling duck (*Dendrocygna autumnalis)*.

Keywords:

1 INTRODUCTION

Nest defence by parents is considered a risky behaviour, as they may endure injury or even

death when trying to ward off predators (de Jong, Nicolaus, Fokkema, & Loonen, 2021;

Samelius & Alisauskas, 2006). Furthermore, nest defence also bears energetic costs (Caro,

2005; Thys, Lambreghts, Pinxten, & Eens, 2019) thus the risk of clutch protection must be

gauged against its benefits (Ringelman & Stupaczuk, 2013). In birds, there is large variation

in how much males and females contribute to parental care and in more than half of all

species, males participate in incubation (Deeming & Reynolds, 2015). In most waterfowl

species, *i.e.* swans, geese and ducks, however, only females incubate. For barnacle geese

(*Branta leucopsis*) breeding on Svalbard anecdotal evidence indicates that males sometimes

sit on the nest (from now on referred to as 'nest sitting'). In this study we dive further into this

behaviour.

The barnacle goose is an Arctic breeding species, renowned for nesting high on steep

mountain cliffs and islets (Mitchell, Black, & Evans, 1998). They often breed on islands with

glaucous gulls (*Larus hyperboreus*), Arctic skuas (*Stercorarius parasiticus*) and great skuas

(*S. skua*), which are the main avian predators of barnacle geese eggs and/ or adults (Prop et

al., 2015; Tombre & Erikstad, 1996). Similar to other geese (*e.g.* lesser snow goose, *A.*

caerulescens caerulescens; pink-footed goose *A. brachyrhynchus;* Canada goose, *B.*

canadensis), barnacle geese protect their eggs against avian predators through a high rate of

nest attendance (Prop, Van Eerden, & van Drent, 1984; Samelius & Alisauskas, 2001;

Schreven, Stolz, Madsen, & Nolet, 2021) and active defence (Clermont, Réale, & Giroux,

2019; de Jong et al., 2021; Dittami, Kennedy, & Thomforde, 1979; Speelman, Hammers,

Komdeur, & Loonen, 2022).

Just as in all other geese, only barnacle females are known to incubate. During nest

construction, geese develop thermosensitive brood patches, which guarantee an efficient heat

transfer from the body to the egg (Jones, 1971). During this time, ganders closely guard their

receptive mates (Lamprecht, 1989). Males stay in close vicinity of the nest, maintaining

contact with their incubating mates visually and vocally (Prop et al., 1984). In case of an

intruder coming close, most males actively protect and defend the nest and incubating female

(Speelman et al., 2022).

Incubation starts after the penultimate egg is laid (Hübner, Tombre, & Erikstad, 2002)

and lasts for 25 days on average (Black, Prop, & Larsson, 2014). Females only take brief

incubation recesses for feeding and preening, as each time they leave the nest, the eggs are at

risk of dropping below an optimal temperature, possibly overheating, or being discovered by a

predator (Ahmad & Li, 2023; Alsos, Elvebakk, & Gabrielsen, 1998). Before leaving the nest,

females often use the down lining of the nest feathers to cover the eggs for insulation and/or

hide them from predators. Common comprehension is that males either join their mates

during incubation recesses, leaving the nest unattended, or stay behind close to the nest

(Owen & Wells, 1979).

Here we report of several cases, in which some barnacle ganders nest sit, while females

are away. First sporadic notes of this behaviour stem from 1979 in a barnacle goose colony on

Nordenskiöldkysten (J. Prop pers. comm.) and 2006 in our Kongsfjorden study colony

database (M. J. J. E. Loonen, pers. obs.). This type of behaviour was also mentioned

anecdotally in a study on pink-footed geese breeding on Iceland, which described a few males

'crouching over the eggs' for up three minutes when their female left (Inglis, 1977). We

decided to investigate this phenomenon further and provide the first scientific account of nest

sitting in any goose species, although we cannot exclude the possibility that reports of this

behaviour may have been disseminated outside peer-reviewed publications. We wanted to

determine how often nest sitting of males occurred, if male age played a role, and whether

individuals, which performed nest sitting did so multiple times or only once. Finally, we

investigated if eggs were more likely to hatch if males were nest sitting.

2 METHODS

Our study population resides on the west coast of Spitsbergen, the largest island of the

Svalbard archipelago. In Kongsfjorden an established breeding colony nests on a group of

several islets (Tombre, Mehlum, & Loonen, 1998). These islets are characterised by exposed

ridges and flat stretches of tundra (Tombre & Erikstad, 1996). Two are monitored in detail;

the larger main breeding islet, Storholmen (ca. 30 ha, 259 nests in 2021) and Prins

Heinrichøya (ca. 3 ha, 29 nests in 2021) located offshore of the village Ny-Ålesund (78°55'N,

11°56'E), Svalbard, Norway.

Within the scope of a larger study on circadian and circannual rhythmicity (de Jong et al.,

2024) we fitted 24 adults (50% males and females of established pairs, respectively) in 2020

with solar-powered GPS-GSM transmitters attached with neckbands (OrniTrack-NL40 3G,

Ornitela, UAB, Lithuania). Except for two geese, which were ringed as goslings, the

individual age was back calculated from the time geese were originally colour marked as

adults; thus, ages given in Table 1 are conservative. Individuals were chosen following a suite

of criteria (for details see supplemental material) including that it would be desirable if both

pair partners were marked with unique colour rings already. The neck collar allowed a

definite identity recognition of pair partners on and close by the nest, even when colour rings

were not visible. Sexing of all individuals was performed when birds were originally ringed

by examining the cloaca for the presence of intromittent penises. To attach transmitters, geese

were caught in the vicinity of the nest either by hand, with a small hand-net, or a fishing rod

with an attached nylon snare ('nest catches', Fenstad et al., 2017), or else during the annually

performed mass captures of moulting geese, where considerable numbers of geese are

funnelled into a trap ('moult catches', Loonen, Oosterbeek, & Drent, 1997). Only males,

which remained close to the nest when a human observer approached, could be caught by

hand, using the fishing rod or hand-net. We are aware that these methods might comprise a

bias in trappability *sensu* STRANGE, an acronym highlighting several possible sources of

sampling biases (Webster & Rutz, 2020), as these males show high nest defence and risk-

taking behaviour (de Jong et al., 2024). From here on we consider them 'bold'.

In 2021, we were able to locate 15 nests of the initial 24 pairs, where either the male

(n=7) or female (n=8) was fitted with a transmitter in the previous year. During transmitter

attachment, four males were caught at the nest, while three were caught during the moult

catch. In the following we refer to unique colour ring codes when mentioning certain

individuals.

Near the 15 nests, we set up wildlife cameras (Usogood TC30 Trail Camera) to get a

more detailed picture of behaviours of males and females during incubation. Cameras were

used in the past and do not seem to disturb the geese (*e.g.* de Jong, Wetherbee, & Loonen,

2019). At each nest, we placed the camera, set in time-lapse mode, at a one-metre distance of

the nest to monitor primarily the behaviour of the incubating female. Cameras took two

pictures every five minutes to detect any movement of the geese. The time in which cameras

provided reliable pictures varied between nests ($\bar{x}$ days $\pm$ SD: 12 ± 6.1; for details see

supplement, Table S1, de Jong et al., 2024), thus we captured part of the incubation period.

When scanning the photos, we noticed that in several instances males, rather than females,

were sitting on the nest (Figure 1). As successive photos in a series are contingent, we only

counted male nest sitting bouts as independent whenever the female returned and sat to the

nest.

2.1 Statistical Analysis

Using the R programming environment (R Core Team, 2021), we (1) applied a Student's *t*-

test to determine whether male age influenced the propensity to nest sit. We applied Fisher's

Exact Tests and give odds ratios and their 95% confidence intervals to determine, (2) if males,

which sat on the nest, were more likely to have been fitted with transmitters, and (3) if male

nest sitting was associated with hatching probability of the clutch, *i.e.* at least one egg hatched

(for details on counts see 2 x 2 contingency tables, supplement Table S2). The sample size

pertaining to the method of catching (nest catch vs. moult catch) was very small and showed

clear separation, because all males that were caught at the nest showed nest sitting (Hauck &

Donner, 1977). Due to the challenges this poses for statistical testing, we have opted to refrain

from a formal statistical analysis in this case.

3 RESULTS

Observations of the 15 nests where cameras were placed revealed that six males sat on the

nest, whereas nine did not (Table 1). Age of males had no effect on nest sitting [$n_{\text{males not nest}}$

$_{\text{sitting}} = 9$, $\bar{x}$ years $\pm$ SD: 8.33 ± 3.16; $n_{\text{males nest sitting}} = 6$, $\bar{x}$ years $\pm$ SD: 11.00 ± 3.1; $t_{(13)} =$

1.612, $p = 0.131$, Figure 2a)]. Of the six nest sitting males, three were observed on the nest

only once (gZUN, oFIJ, oFIA), whereas the others three males were observed multiple times

(oFYJ and yBYI four times, oFZY six times). Males with neck collars were more likely to sit

on nests than males without [Fisher's Exact Test: $p = 0.041$, $OR_{\text{2-sided}} = 13.59$ with 95% CI of

[0.86, 934.01], Figure 2b)]. Among males caught near the nest, 100% were observed sitting

on nests (N = 4), compared to 33.3% of males caught during moult (N on nest = 1, N not on

nest = 2, Figure 2c). Whether males nest sat had no effect on whether eggs hatched or else

were abandoned/ depredated [Fisher's Exact Test: $p = 0.329$, $OR_{\text{2-sided}} = 0.3$ with 95% CI of

[0.02, 4.30], Figure 2d), see Table S2 for 2 x 2 contingency tables]. All but one male (oFIA),

which started to nest sit shortly after the eggs hatched (see supplement Figure S1), sat on the

nest during incubation. Males took on average 12 minutes (SD $\pm$ 17 min, range 0 – 42 min)

after the female left before they started nest sitting and they averagely stayed for 39 minutes

(SD $\pm$ 21 min, range = 12 – 70 min). This corresponds to about 60% (SD $\pm$ 27%, range = 16 –

88%) of the time the female was on incubation recess. After males terminated nest sitting, it

took approximately 17 minutes (SD $\pm$ 13 min, range = 3 – 32 min) for the female to return to

the nest (for details see Table S3). Based on examinations of four nests, where we had at least

three days of camera observations (N = 4), the percentage of time when males nest sat during

the females' incubation recesses varied greatly between individuals (mean 20.0 % of the time,

SD $\pm$ 21, range = 1.02 – 43%, for details see Table S4). However, throughout this study

sample size is small, and these results should be taken with caution.

4 DISCUSSION AND CONCLUSION

4.1 Nest sitting in barnacle ganders – possible functions

To the best of our knowledge, this is the first scientific account of males sitting on nests in

any goose species. Although we may have introduced a bias to catch predominantly bold

individuals (Webster & Rutz, 2020), this does not alter the fact that some ganders perform this

behaviour. We oppose to calling this behaviour 'male incubation', as termed repeatedly in the

waterfowl literature (*e.g.* Brugger & Taborsky, 1994; Bruggers, 1979; Hawkins, 1986; Rollin,

1957), because there cannot be active transfer of heat from the male to the eggs as ganders do

not form brood patches. Thus, we revive the term 'nest sitting' (Henson & Cooper, 1992) as

the appropriate label. We identified from the photos that male posture on the nest was

somewhat different from the female posture during incubation, *i.e.* males sat higher than

females, not in the nest bowl but rather on top of the nest (compare panels A) and D) in

Figure 1). Female ducks and geese sometimes adopt such as posture not only before actual

incubation starts, but also during incubation (Hartman et al., 2023; Lorenz, 1991; Poussart,

Larochelle, & Gauthier, 2000) to shield the nest and eggs from *e.g.* predation, inclement

weather and/or intraspecific brood parasitism. We propose that male nest sitting has

equivalent functions in the absence of the female. To date, we cannot provide a definite

explanation of why this behaviour occurs, as neither the age of males seems to play a role, nor

could we show an increase in hatching success of nests, where males nest sat. At this moment

we lack a large enough sample for more in-depth analyses, such as when nest sitting occurs

across the nesting period or to what extent males may compensate for more or longer

incubation recesses of their mates. Furthermore, it should also be investigated in-depth,

whether males adjust their behaviour in response to certain environmental conditions, such as

nest density and location, and whether nest sitting confers fitness benefits (Madsen et al.,

2019; Mery & Burns, 2010). In the following we suggest three possible, not mutually

exclusive, functions of male nest sitting to initiate future studies not only in barnacle geese,

but also other goose species.

The primary cause of hatching failure in ground nesting waterfowl is predation, thus

various strategies are employed for nest protection (Peterson et al., 2022; Quinn, Prop,

Kokorev, & Black, 2003; Samelius & Alisauskas, 2001; Schranck, 1972). Geese often nest

near their main aerial predators. To not leave the nest unattended when the females are away,

nest sitting of barnacle ganders might function in shielding the nest. This is backed by our

choosing the most aggressive males when attaching transmitters, which might be bold enough

to attack predators actively. In the future, attention should be paid to how often nest sitting of

males occurs, how long the incubation recesses of the female last, what the predation pressure

is, and at which stage during incubation the behaviour is performed to get a deeper

understanding of male nesting as one form of predator protection.

Another possible function of nest sitting by males is to control the temperature in the

nest. To some extent, nest insulation can reduce both the degree of egg cooling when the

female is absent, and the time spent to rewarm the eggs after she returns from an incubation

recess (Thompson & Raveling, 1988). Indeed, Arctic geese adjust their incubation behaviour

in response to prevailing weather conditions (Elkins, 2004; Harvey, 1971; Poussart, Gauthier,

& Jacques, 2001). Yet, egg cooling in the absence of the female might be overrated as even

under unfavourable weather conditions, egg temperature drops during incubation recesses are

nowhere near the minimal temperatures, which endanger an embryo's development or

survival (Poussart et al., 2000). It is worth noting that also overheating of unprotected eggs in

the nest might become an issue in the Arctic in the future. Even if barnacle nests are exposed

to direct sun, they probably will, at present, not reach this temperature limit long enough for

embryos to die. Yet, common guillemots (*Uria aalge*) already showed an increased

probability of egg loss at higher temperatures in a high-latitude colony in the Baltic Sea (Olin

et al., 2024). Taken together, although nest sitting of barnacle ganders might assist in

temperature control of eggs in the nest, we suggest it to only play a minor role, at least here

and now.

If male nest sitting occurs during the period of egg laying, another possible function

might be to prevent intraspecific brood parasitism, which frequently occurs in barnacle geese

(Black et al., 2014). We cannot answer this question adequately, however, because we placed

cameras not until the host female had started incubation, therefore lack photos from the laying

period.

4.2 'Male incubation' in waterfowl?

In the waterfowl literature only swans, and to some extent whistling ducks are described to

deviate from the female only incubation pattern in waterfowl, as here male incubation occurs

(Brugger & Taborsky, 1994; Hawkins, 1986; Scott, 1977). But are these males indeed

incubating or is it a term that should be avoided, as is leads to a misconception when

compared with species, where incubation is truly shared (Deeming & Reynolds, 2015)?

Hawkins already concluded that male incubation in tundra swans (*Cygnus columbianus*

columbianus.) was not essential for successful embryo development (Hawkins, 1986) but

protected against egg predation and provided some egg cooling benefits. This led Henson &

Cooper to apply the term 'nest sitting' in trumpeter swans *(C. buccinator)*, as here males

occasionally exhibiting some of the nest-settling motions characteristic of incubating females,

independent of predator presence or adverse weather conditions (Henson & Cooper, 1992).

Our extensive search of the literature revealed that in most cases, in which male incubation

was described (*e.g.* Bruggers, 1979; Flickinger, 1975; Rollin, 1957), they nest sit rather than

incubate.

The two notable exceptions, for which the term incubation is warranted, are the black

swan (*C. atratus*, BLSW, Cygnidae) and black-bellied whistling duck (*D. autumnalis*,

BBWD, Anatidae*)*. Whereas BLSW exhibit the typical long-term monogamy and biparental

care of swans, this is true also for the BBWD, a feature atypical for ducks. Some other

features, which the two species share are that neither females nor males develop brood

patches (Bolen & Smith, 1979; Skutch, 1976), thus either parent is equally fit to incubate, and

that both species may re-nest after successful breeding attempts (Bolen & Smith, 1979;

Brugger & Taborsky, 1994; Coleman, 2014; Delnicki, 1973; James, Thompson, & Ballard,

2012). Most other waterfowl attempt a second clutch only if the first one failed. BLSWs and

BBWDs might be able to re-nest because both parents incubate. Near constant incubation

minimizes the time of egg development and hatching, thereby shortening the inter-clutch

interval (James et al., 2012).

4.3 Conclusion

To sum up, this is the first scientific report in any goose species, where males expand

paternal duties beyond the established view, namely that during the breeding period they only

protect the nest and female in close vicinity of the nest. We have pictorial evidence that

several barnacle ganders nest sat in the absence of their mates. Both predation and thermal

risks are increased for eggs, which are not protected by parents, thus minimising those risks

are the main causes for successful hatching. In barnacle geese, we suggest nest sitting of

males to be a response to avian predator presence. We cannot exclude that preventing egg

cooling in the absence of the female may also play a role but consider this unlikely. Our final

point is that in most waterfowl species, males do not participate in incubation. Nest sitting is

an unbiased term and should be used for cases, in which no active warming of eggs has (yet)

been demonstrated.

Acknowledgements

We are grateful to Børge Moe (Norwegian Institute for Nature Research, NINA) as the project

administrator of AJS' field grant, the staff of the French - German Arctic Research Base at

Ny-Ålesund, chiefly Bettina Haupt (station leader AWI 2020/21), and KingsBay AS, Ny-

Ålesund for logistic support. We thank Jouke Prop for sharing his unpublished observations

and Kees Schreven for directing out attention to the one dated reference, where nest sitting in

ganders was briefly alluded to. Comments by two anonymous reviewers and Thomas Lameris

from the editorial board of *Ardea* helped to improve an earlier version of this manuscript.

Funding was provided by the Austrian Science Fund (P32216 to I. B. R. Scheiber) and an

Arctic Field Grant (The Research Council of Norway) in collaboration with NINA (ES676286

- to A. J. Slettenhaar). The study conforms to Directive 2010/63/EU and was conducted under

FOTS ID 23358 from the Norwegian Animal Research Authority and approved by the

Governor of Svalbard (RIS ID 11237).

REFERENCES:

Ahmad, I. M., & Li, D. (2023). More than a simple egg: Underlying mechanisms of cold
tolerance in avian embryos. *Avian Research, 14*, 100104. doi: 10.1016/j.avrs.2023.100104

Alsos, I. G., Elvebakk, A., & Gabrielsen, G. W. (1998). Vegetation exploitation by barnacle
geese *Branta leucopsis* during incubation on Svalbard. *Polar Research, 17*, 1-14. doi:
10.3402/polar.v17i1.6603

Black, J. M., Prop, J., & Larsson, K. (2014). *The Barnacle Goose*. London, UK: Bloomsbury
Publishing.

Bolen, E. G., & Smith, E. N. (1979). Notes on the incubation behavior of black-bellied
whistling ducks. *Prairie Naturalist, 11*(4), 119-123.

Brugger, C., & Taborsky, M. (1994). Male incubation and its effect on reproductive success
in the black swan, *Cygnus atratus. Ethology, 96*(2), 138-146. doi: 10.1111/j.1439-
0310.1994.tb00889.x

Bruggers, R. L. (1979). Nesting patterns of captive mandarin ducks. *Wildfowl, 30*, 45-54.

Caro, T. M. (2005). *Antipredator Defenses in Birds and Mammals*. Chicago, USA: University
of Chicago Press.

Clermont, J., Réale, D., & Giroux, J.-F. (2019). Similarity in nest defense intensity in Canada
goose pairs. *Behavioral Ecology and Sociobiology, 73*(8). doi: 10.1007/s00265-019-2719-3

Coleman, J. T. (2014). Breeding biology of the black swan *Cygnus atratus* in southeast
Queensland, Australia. *Wildfowl, 64*, 217-230.

de Jong, M. E., Nicolaus, M., Fokkema, R. W., & Loonen, M. J. J. E. (2021). State
dependence explains individual variation in nest defence behaviour in a long-lived bird.
Journal of Animal Ecology, 90(4), 809-819. doi: 10.1111/1365-2656.13411

de Jong, M. E., Slettenhaar, A. J., Fokkema, R. W., Leh, M., Barnreiter, E., Griffin, L., . . .
Scheiber, I. B. R. (2024). Diel and seasonal rhythmicity in activity and corticosterone in an
Arctic migratory herbivore: A multifaceted approach. *preprint*. doi:
10.1101/2024.08.30.610510

de Jong, M. E., Wetherbee, R., & Loonen, M. J. J. E. (2019). Effects of fleas on nest success
of Arctic barnacle geese: Experimentally testing the mechanism. *Journal of Avian Biology,
50*(5), e01944. doi: 10.1111/jav.01944

Deeming, D. C., & Reynolds, S. J. (2015). *Nests, Eggs, and Incubation*. Oxford, UK: Oxford
University Press.

Delnicki, D. E. (1973). *Renesting, incubation behavior, and compound clutches of the black-
bellied tree duck in southern Texas.* MSc, Texas Tech University, Lubbock, TX, USA.
(31295001778181)

Dittami, J. P., Kennedy, S., & Thomforde, C. (1979). Observations on barnacle goose
breeding, *Branta leucopsis*, in Spitsbergen 1975. *Journal of Ornithology, 120*, 188-195. doi:
10.1007/BF01642997

Elkins, N. (2004). Breeding: Incubation. In N. Elkins (Ed.), *Weather and Bird Behaviour*
(Vol. 3rd edition, pp. 101-104). London, UK: T.& A.D. Poyser.

Fenstad, A. A., Bustnes, J., Lierhagen, S., Gabrielsen, K. M., Öst, M., Jaatinen, K., . . .
Krøkje, Å. (2017). Blood and feather concentrations of toxic elements in a Baltic and an
Arctic seabird population. *Marine Pollution, 114*(2), 1152-1158. doi: j.marpolbul.2016.10.034

Flickinger, E. L. (1975). Incubation by a male fulvous tree duck. *Wilson Bulletin, 87*(1), 106-
107.

Hartman, C. A., Ackerman, J. T., Peterson, S. H., Fettig, B., Casazza, M., & Herzog, M. P.
(2023). Nest attendance, incubation constancy, and onset of incubation in dabbling ducks.
Plos One, 18(5), e0286151. doi: 10.1371/journal.pone.0286151

Harvey, J. M. (1971). Factors affecting blue goose nesting success. *Canadian Journal of
Zoology, 49*(2), 223-234. doi: 10.1139/z71-03

Hauck, W. W., & Donner, A. (1977). Wald's test as applied to hypotheses in logit analysis.
Journal of the American Statistical Association, 72(360a), 851-853. doi:
10.1080/01621459.1977.10479969

Hawkins, L. L. (1986). Nesting behaviour of male and female whistling swans and
implications of male incubation. *Wildfowl, 37*, 5-27.

Henson, P., & Cooper, J. A. (1992). Division of labour in breeding trumpeter swans *Cygnus
buccinator*. *Wildfowl, 43*, 40-48.

Hübner, C. E., Tombre, I. M., & Erikstad, K. E. (2002). Adaptive aspects of intraclutch egg-
size variation in the high Arctic barnacle goose (*Branta leucopsis*). *Canadian Journal of
Zoology, 80*, 1180-1188. doi: 10.1139/z02-100

Inglis, I. R. (1977). The breeding behaviour of the pink-footed goose: Behavioural correlates
of nesting success. *Animal Behaviour, 25*, 747-764. doi: 10.1016/0003-3472(77)90125-7

James, J. D., Thompson, J. E., & Ballard, B. M. (2012). Evidence of double brooding by
black-bellied whistling-ducks. *Wilson Journal of Ornithology, 124*(1), 183-185. doi:
10.1676/11-082.1

Jones, R. E. (1971). The incubation patch of birds. *Biological Reviews, 46*(3), 315-339. doi:
10.1111/j.1469-185X.1971.tb01048.x

Lamprecht, J. (1989). Mate guarding in geese: Awaiting female receptivity, protection of
paternity or support of female feeding. In A. E. Rasa, C. Vogel & E. Voland (Eds.), *The
Sociobiology of Sexual and Reproductive Strategies*. Dordrecht, NL: Springer Science and
Business Media.

Loonen, M. J. J. E., Oosterbeek, K., & Drent, R. H. (1997). Variation in growth of young and
adult size barnacle geese *Branta leucopsis*: Evidence for density dependence. *Ardea, 85*, 177-
192.

Lorenz, K. (1991). *Here I Am - Where Are You?* New York, NY, USA: Hartcourt Brace
Jovanovich.

Madsen, J., Jaspers, C., Frikke, J., Gundersen, O. M., Nolet, B. A., Nolet, K., . . . de Vries, P.
P. (2019). A gloomy future for light-bellied brent geese in Tusenøyane, Svalbard, under a
changing predator regime. *Polar Research, 38*, 3393. doi: 10.33265/polar.v38.3393

Mery, F., & Burns, J. G. (2010). Behavioural plasticity: An interaction between evolution and
experience. *Evolutionary Ecology, 24*(3), 571-583. doi: 10.1007/s10682-009-9336-y

Mitchell, C., Black, J. M., & Evans, M. (1998). Breeding success of cliff-nesting and island-
nesting barnacle geese in Svalbard. *Norsk Polarinstitutt Skrifter, 200*, 141-146.

Olin, A. B., Dück, L., Berglund, P. A., Karlsson, E., Bohm, M., Olsson, O., & Hentati-
Sundberg, J. (2024). Breeding failures and reduced nest attendance in response to heat stress
in a high-latitude seabird. *Marine Ecology Progress Series, 737*, 147-160. doi:
10.3354/meps14244

Owen, M., & Wells, R. L. (1979). Territorial behaviour in breeding geese - A re-examination
of Ryder's hypothesis. *Wildfowl, 30*, 20-26.

Peterson, S. H., Ackerman, J. T., Keating, M. P., Schacter, C. R., Hartman, C. A., Casazza,
M. L., & Herzog, M. P. (2022). Predator movements in relation to habitat features reveal
vulnerability of duck nests to predation. *Ecology and Evolution, 12*(9). doi:
10.1002/ece3.9329

Poussart, C., Gauthier, G., & Jacques, L. (2001). Incubation behaviour of greater snow geese
in relation to weather conditions. *Canadian Journal of Zoology, 79*, 671-678. doi:
10.1139/z01-023

Poussart, C., Larochelle, J., & Gauthier, G. (2000). The thermal regime of eggs during laying
and incubation in greater snow geese. *Condor, 102*(2), 292-300. doi:
10.1093/condor/102.2.292

Prop, J., Aars, J., Bårdson, B.-J., Hanssen, S. A., Bech, C., Bourgeon, S., . . . Moe, B. (2015).
Climate change and the increasing impact of polar bears on bird populations. *Frontier in*
Ecology and Evolution, 3, 33. doi: 10.3389/fevo.2015.00033

Prop, J., Van Eerden, M. R., & van Drent, R. H. (1984). Reproductive success in the barnacle
goose *Branta leucopsis* in relation to food exploitation on the breeding grounds, western
Spitsbergen. *Norsk Polarinstitutt Skrifter, 181*, 87-117.

Quinn, J. L., Prop, J., Kokorev, Y., & Black, J. M. (2003). Predator protection or similar
habitat selection in red-breasted goose nesting associations: Extremes along a continuum.
Animal Behaviour, 65(2), 297-307. doi: 10.1006/anbe.2003.2063

R Core Team. (2021). R: A language and environment for statistical computing (Version
1.13). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from
https://www.R-project.org/.

Ringelman, K. M., & Stupaczuk, M. J. (2013). Dabbling ducks increase nest defense after
partial clutch loss. *Condor, 115*(2), 290-297. doi: 10.1525/cond.2013.120096

Rollin, N. (1957). Incubation by drake wood duck in eclipse plumage. *Condor, 59*(4), 263-
265. doi: 10.2307/1364656

Samelius, G., & Alisauskas, R. T. (2001). Deterring arctic fox predation: The role of parental
nest attendance by lesser snow geese. *Canadian Journal of Zoology, 79*, 861-866. doi:
10.1139/z01-048

Samelius, G., & Alisauskas, R. T. (2006). Sex-biased costs in nest defence behaviours by
lesser snow geese (*Chen caerulescens*): Consequences of parental roles. *Behavioral Ecology*
and Sociobiology, 59, 805-810.

Schranck, B. W. (1972). Waterfowl nest cover and some predation relationships. *The Journal*
of Wildlife Management, 36(1), 182. doi: 10.2307/3799210

Schreven, K. H. T., Stolz, C., Madsen, J., & Nolet, B. A. (2021). Nesting attempts and
success of Arctic-breeding geese can be derived with high precision from accelerometry and
GPS-tracking. *Animal Biotelemetry, 9*(1), 25. doi: 10.1186/s40317-021-00249-9

Scott, D. (1977). Breeding behaviour of wild whistling swans. *Wildfowl, 28*, 101-106.

Skutch, A. F. (1976). *Parent Birds And Their Young*: University of Texas Press.

Speelman, F. J. D., Hammers, M., Komdeur, J., & Loonen, M. J. J. E. (2022). Nest defence
behaviour is similar between pair members but only male behaviour predicts nest survival in
barnacle geese. *Journal of Avian Biology, 2022*(9), e02982. doi: 10.1111/jav.02982

Thompson, S. C., & Raveling, D. G. (1988). Nest insulation and incubation constancy of
Arctic geese. *Wildfowl, 39*, 124-132.

Thys, B., Lambreghts, Y., Pinxten, R., & Eens, M. (2019). Nest defence behavioural reaction
norms: testing life-history and parental investment theory predictions. *R Soc Open Sci, 6*(4),
182180. doi: 10.1098/rsos.182180

Tombre, I. M., & Erikstad, E. (1996). An experimental study of incubation effort in high-
Arctic barnacle geese. *Journal of Animal Ecology, 65*, 325-331. doi: 10.2307/5878

Tombre, I. M., Mehlum, F., & Loonen, M. J. J. E. (1998). The Kongsfjorden colony of
barnacle geese: Nest distribution and the use of breeding islands 1980-1997. *Norsk*
Polarinstitutt Skrifter, 200, 57-65.

Webster, M. M., & Rutz, C. (2020). How STRANGE are your study animals? *Nature,*
582(7812), 337-340. doi: 10.1038/d41586-020-01751-5
440

Table 1: Identities of 15 focal pairs, whose nests were monitored with wildlife cameras in
2021. For each individual, we provide codes of unique colour (o=orange, g=green, y=yellow
plus a three-letter code) and metal (CA***) rings, and if they were fitted with a transmitter,
the neck collar ID. Except two individuals, which were ringed as goslings (*), the minimum
age is given, calculated from the time the geese were originally ringed as adults ($\bar{x}$ age ± SD:
males 9.4 ± 3.31; females 10.4 ± 4.46). Whether or not males sat on nests is also indicated.
448

Male colour ID	Male ID	Male neck collar ID	Age (years)	Male nest sitting	Female colour ID	Female ID	Female neck collar ID	Age (years)
oFYJ	CA46944	201913	16	yes	oFVD	CA46857		10
gZUN	CA45605	201915	9	yes	oFLC	CA45810		12
alur	CA47072	201917	8	no	yBSV	CA45997		4
oFZY	CA46942	201918	10	yes	gSNF	CA36931		19
oFIJ	CA45800	201924	12	yes	yBHJ	CA45891		5
oFIA	CA45795	201930	12	yes	gLDC	CA41238		14
yBID	CA46840	201932	7	no	yBIB	CA44228		14
oFZX	CA46941		10	no	gTTL	CA41113	201912	16*
yBYI	CA45746		7	yes	yAUF	CA45744	201919	7
yAXY	CA45703		8	no	yBHY	CA46848	201925	5*
gLHX			2	no	yAVV	CA45736	201926	7
yAZT	CA45715		8	no	oFXV	CA45632	201927	10
yAXC	CA45942		8	no	yAPH	CA45929	201931	8
gTCZ	CA46957		10	no	yAXV	CA45701	201934	8
yBYX	CA41397		14	no	yBYV	CA41400	201935	14

FIGURE LEGENDS

Figure 1: An example of one barnacle gander nest sitting. Photos taken on June 18th, 2021,

between 15:48 and 16:28. Panel A) The female without a neck collar, fitted with the

individual orange colour ring oFVD, is incubating. Panel B) The female left the nest. Panel C)

Male alur with neck collar approaches the nest. Panel D) He sits on the nest with the neck

collar clearly visible. Panel E) The female (with the colour ring partially visible) returns to the

nest. Panel F) She resumes incubation. Temperature and time are shown in the lower right-

hand corner of each photo.

Figure 2: Features, which might influence nest sitting in barnacle ganders. a) Age did not

affect the likelihood of male to nest sit (Student's *t*-test: $p > 0.05$, whiskers represent the

SEM). b) Males, which were fitted with a neck collar, were more likely to nest sit than males

without (Fisher's Exact Test: $p = 0.041$). c) Among males caught near the nest, 100% were

observed to nest sit, while only 33,3% of males caught during moult did so. d) Whether eggs

hatched or did not hatch was independent of males' nest sitting (Fisher's Exact Test: $p >$

0.05).

469
470   Figure 1
471

472
473   Figure 2
474