Dads on duty: First account of nest sitting in barnacle ganders

Isabella B. R. Scheiber * [1], Annabel J. Slettenhaar [2,3], Maarten J. J. E. Loonen [3] and

Margje E. de Jong [1,3]

*corresponding author

[1] Dept. of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030

Vienna, Austria

[2] Faculty of Biosciences and Aquaculture, Nord University, Kongens Gate 42, 7713

Steinkjer, Norway

[3] Arctic Centre, Aweg 30, 9718 CW Groningen, The Netherlands

Orcid IDs and E-Mail Addresses:

Isabella B. R. Scheiber: 0000-0003-3438-1836; isabella.scheiber@univie.ac.at

Annabel J. Slettenhaar: 0009-0005-9954-0721; annabel.j.slettenhaar@nord.no

Maarten J. J. E. Loonen: 0000-0002-3426-4595; m.j.j.e.loonen@rug.nl

Margje E. de Jong: 0000-0002-4941-8323; me.d.jong@gmail.com

Acknowledgements

We are grateful to Børge Moe (Norwegian Institute for Nature Research, NINA) as the

project administrator of AJS' field grant, the staff of the French - German Arctic Research

Base at Ny-Ålesund, chiefly Bettina Haupt (station leader AWI 2020/21), and KingsBay AS,

Ny-Ålesund for logistic support. Funding was provided by the Austrian Science Fund

(P32216 to I. B. R. Scheiber) and an Arctic Field Grant (The Research Council of Norway) in

collaboration with NINA (ES676286 - to A. J. Slettenhaar).

Journal Section: Behavioural Note

Conflict of Interest Statement:

The authors declare no conflict of interest.

Data Availability Statement:

The data that support the findings of this study are provided within the article and

corresponding supplemental material.

Ethics Approval Statement:

The study conforms to Directive 2010/63/EU and was conducted under FOTS ID 23358 from

the Norwegian Animal Research Authority and approved by the Governor of Svalbard (RIS

ID 11237).

Author Contributions

Isabella B. R. Scheiber: Conceptualization; formal analysis; funding acquisition;

methodology; project administration; supervision; verification; visualization; writing –

original draft; writing – review and editing. **Annabel J. Slettenhaar:** Data curation; funding

acquisition, investigation; software; visualization; writing – review and editing. **M. J. J. E.**

Loonen: Resources; writing – review and editing. **Margje E. de Jong**: conceptualization;

data curation; formal analysis; investigation; methodology; software; supervision;

verification; visualization; writing – original draft; review and editing.

Abstract

In most Anseriformes (ducks, geese and swans) only females are known to incubate. The few exceptions where males have been described to incubate regularly are several species of swans (Cygninae) and whistling ducks (Dendrocygninae). Here we describe, unprecedented in any goose species, indicidents of male nest sitting in barnacle geese (*Branta leucopsis*) as a form of paternal care of nest attendance. Based on pictures from wildlife cameras we identified males, which sat on their nests when their mates took incubation recesses. Wildlife cameras were placed at nests of which either the male or female was fitted with a GPS neck collar in the year prior to this study. To attach transmitters, some geese were caught while defending their nests, thus we may have unintentionally selected bolder males which defended their nests more aggressively and were easier to catch. Nest sitting occurred relatively frequent, *i.e.* in 6/15 individuals. Our results show that males with collars were more likely to nest sit, but this does not deflect from the fact this behaviour is novel in geese. Nest sitting may even be more common as collared birds are more easily recognised, but also uncollared males have been observed to nest sit. We discuss several possible functions of this behaviour, *i.e.* against raiding of nests by aerial predators, thermal control of nest temperature, and intraspecific brood parasitism. At this time we cannot demonstrate a possible function, as chances of successful hatching were not increased in nests of males, which were nest sitting. At this time, we lack sample size for more in depth analyses. Lastly, we discuss whether 'male incubation' is misleading in the waterfowl literature, as it is truly justified for only two species, the black swan and black-bellied whistling duck.

Keywords:
Branta leucopsis, geese, waterfowl, paternal care, male incubation, incubation behaviour

1 INTRODUCTION

Nest defence by parents is considered a risky behaviour, as they may endure injury or even
death when trying to ward off predators [1], thus the risk of clutch protection has to be gauged
against its benefits [2]. In birds, there is large variation in how much males and females
contribute to parental care and in more than half of all species, males participate in incubation
[3]. In most waterfowl species, *i.e.* swans, geese and ducks, however, only females incubate.
Thus far, it was never observed in geese (Anserinae). Therefore, the purpose of this paper is
threefold: First, we revive the term '*nest sitting*' [4] and describe it for the first time in geese,
i.e. barnacle ganders (*Branta leucopsis*). Nest sitting refers to cases, where the non-incubating
parent attends to or is present directly on the nest and differs from incubation, which entails
the transfer of heat to the developing eggs [5]. Secondly, we will discuss possible functions of
nest sitting in barnacle geese, and, finally, we question whether previously reported studies in
waterfowl, which refer to 'male incubation', are actually misinterpreted.

Barnacle geese are an Arctic breeding species, well-known for nesting high on steep
mountain cliffs and islets as a protection against predators [6]. Our study population breeds
contemporaneously on islands with common eiders (*Somateria mollisma*), glaucous gulls
(*Larus hyperboreus*), Arctic (*Stercorarius parasiticus*) and great skuas (*S. skua*), the latter
three being the main avian predators of barnacle geese [7, 8]. Just as other geese (*e.g.* lesser
snow goose, *Anser caerulescens caerulescens;* pink-footed goose, *A. brachyrhynchus;*
Canada goose, *Branta canadensis),* barnacle geese, protect their nests against avian predators
through a high rate of nest attendance [9-11] and active defence [1, 12-14].

Nests on islets seem relatively safe from terrestrial predators. Islets in the inner fjord,
however, can stay connected with the mainland via ice-bridges when there is a late break-up
of fjord ice. This allows Arctic foxes (*Vulpes lagopus*) not only to reach the islands, but may
also confine them to stay there, once the ice breaks, at the time when the geese breed [15].
However, as the fjord ice has diminished in recent years [*e.g.* 16, 17], the risks of
encountering foxes on the breeding islands has gone down [18]. Polar bears (*Ursus
maritimus*), which are known to prey on eggs and goslings [19], have increased substantially
in numbers [8, 20]. They now visit islands more and more during the breeding season to raid
nests [8, 21, 22]. This results in complete nest failures in some years [8].

Barnacle geese are socially monogamous and biparental with precocial offspring where
parental roles before and after hatching are clearly defined. Before hatching, males normally
do not come straight up to the nest, but stay in close vicinity, maintaining contact with their

incubating mates visually and vocally [9, 23]. In case of an intruder coming close, most males

actively protect and defend the nest and incubating female [1, 13].

Just as in all other geese, only females incubate. During nest construction, they develop

thermosensitive brood patches, which guarantee an efficient heat transfer from the body to the

egg [24]. Female geese actively pluck feathers, which they add as an insulating inner layer of

nest lining. They then begin to lay one egg per day until a clutch is completed with a mean

clutch size of about five eggs [25]. During this time, barnacle ganders closely guard their

receptive mates [26]. Incubation in barnacle geese lasts for 25 days on average [27].

Incubation starts after the penultimate egg is laid [28]. Females only take brief incubation

recesses for feeding and preening, as each time they leave the nest, the eggs are at risk of

dropping below an optimal temperature, possibly overheating, or being discovered by a

predator [29, 30]. In our study population, geese take five 20-min incubation recesses per day

on average, yielding an incubation constancy of 93% [31]. Before leaving the nest, females

use the down lining of the nest feathers to cover the eggs for insulation and/or hide them from

predators. Common comprehension is that males either join their mates during incubation

recesses, leaving the nest unattended, or stay behind close to the nest [32]. To the best of our

knowledge, we now report, for the first time in geese, that some males nest sit, while females

are away.

2 METHODS

Our study population resides on the west coast of Spitsbergen, the largest island of the

Svalbard archipelago. In Kongsfjorden an established breeding colony nests on a group of

several islets [15]. These islets are characterised by exposed ridges and flat stretches of tundra

[7]. Two are monitored in detail; the larger main breeding islet, Storholmen (ca. 30 ha, 259

nests in 2021) and Prins Heinrichøya (ca. 3 ha, 29 nests in 2021) located offshore of the

village Ny-Ålesund (78°55'N, 11°56'E), Svalbard, Norway.

Within the scope of a larger study on circadian and circannual rhythmicity, we fitted 24

adults (½ males, ½ females of established pairs) in 2020 with solar-powered GPS-GSM

transmitters attached with neckbands (OrniTrack-NL40 3G, Ornitela, UAB, Lithuania) in

order to locate geese during migrations and on the wintering grounds. Those 24 individuals

were chosen following a suite of criteria, including that it would be desirable if both pair

partners were marked with unique colour rings already (for details of criteria see

supplemental material). To attach transmitters, geese were caught in the vicinity of the nest

either by hand, with a small hand-net, or a fishing rod with an attached nylon snare ['nest

catches', 33], or else during the annually performed mass captures of moulting geese, where considerable numbers of geese are funnelled into a trap ['moult catches', 34]. Only males, which remained close to the nest when a human observer approached, could be caught by hand, using the fishing rod or hand-net. We are aware that these methods might comprise a bias in trappability *sensu* STRANGE [35], as these males show high nest defence and risk-taking behaviour [1, 36]. From here on we consider them 'bold'.

In 2021, we were able to locate 15 nests of the initial 24 pairs, where either the male (n=7) or female (n=8) was fitted with a transmitter in the previous year. During transmitter attachment, four males were caught at the nest, while three were caught during the moult catch. The neck collar allows a definite recognition of the pair partners on and close by the nest, even when colour rings are not visible. In the following we refer to colour ring codes, when we mention certain individuals.

Near the 15 nests, we set up wildlife cameras (Usogood TC30 Trail Camera) to get a more detailed picture of behaviours of males and females during incubation. Cameras were used in the past and do not seem to disturb the geese [*e.g.* 37]. At each nest, we placed the camera, set in time-lapse mode, in one-metre distance of the nest to monitor primarily the behaviour of the incubating female. Cameras took two pictures every five minutes to detect any movement of the geese. The time in which cameras provided reliable pictures varied between nests ($\bar{x}$ days ± SD: 12 ± 6.1; for details see supplement, Table S1)

When scanning the photos, we noticed that in several instances males, rather than females, were sitting on the nest (Figure 1). In the goose data base, first sporadic notes of this behaviour stem from 2006 (M. E. de Jong & M. J. J. E. Loonen, pers. obs.). We thus decided to investigate this phenomenon further. In particular, we wanted to determine how often nest sitting of males occurred, whether only certain individuals did so, if male age played a role, and whether individuals, which performed nest sitting did so multiple times or only once. As successive photos in a series are contingent, we only counted male nest sitting bouts as independent if the female returned to the nest and incubated the clutch for a pronounced period of time. As occurrence of male nest sitting might have been a consequence of our selection of males [35], we also took the method of catching into account. Finally, we investigated if eggs, were more likely to hatch if males were nest sitting.

2.1 Statistical Analysis

Using the R programming environment [38], we (1) applied a Student's *t*-test to determine whether male age influenced the propensity to nest sit. We applied Fisher's Exact Tests and

give odds ratios and their 95% confidence intervals to determine, (2) if males, which sat on

the nest, were more likely to have been fitted with transmitters, (3) whether the method of

catching, *i.e.* snare/net vs. moult catch, was correlated with male nest sitting, and (4) if male

nest sitting was associated with hatching probability of the clutch, *i.e.* at least one egg hatched

(for details on counts of the three queries see 2 x 2 contingency tables, supplement Table S2).

2.2 Ethics Approval Statement:

The study conforms to Directive 2010/63/EU and was conducted under FOTS ID 23358 from

the Norwegian Animal Research Authority and approved by the Governor of Svalbard (RIS

ID 11237).

3 RESULTS

Observations of the 15 nests where cameras were placed revealed that six males sat on the

nest, whereas nine did not (Table 1). Age of males had no effect on nest sitting [$n_{males\ not\ nest}$

$_{sitting}$ = 9, $\bar{x}$ years ± SD: 8.33 ± 3.16; $n_{males\ nest\ sitting}$ = 6, $\bar{x}$ years ± SD: 11.00 ± 3.1; $t_{(13)}$ =

1.612, p = 0.131, Figure 2a)]. Of the six nest sitting males, two were observed on the nest

only once (gZUN, oFIJ), whereas the others four males were observed multiple times (oFIA

twice, oFIY and yBYI four times, oFZY five times). Males with neck collars were more likely

to sit on nests than males without [Fisher's Exact Test: p = 0.041, $OR_{2\text{-sided}}$ = 13.59 with 95%

CI of [0.86, 934.01], Figure 2b)]. The likelihood of males sitting on the nest was independent

of whether they were caught near the nest during incubation or during the moult catches

[Fisher's Exact Test: p = 0.143, $OR_{2\text{-sided}}$ = 0.00 with 95% CI of [0.00, 3,53], Figure 2c)].

Whether males nest sat had no effect on whether eggs hatched or else were abandoned/

depredated [Fisher's Exact Test: p = 0.329, $OR_{2\text{-sided}}$ = 0.3 with 95% CI of [0.02, 4.30], Figure

2d)]. However, throughout this study sample size is small and these results should be taken

with caution.

4 DISCUSSION AND CONCLUSION

4.1 Nest sitting in barnacle ganders – possible functions

To the best of our knowledge, this is the first confirmation of males sitting on nests in any

goose species. Although we may have introduced a bias to catch predominantly bold

individuals [35], this does not alter the fact that we have shown for the first time that some

ganders perform this behaviour. We oppose to calling this behaviour 'male incubation', as

termed repeatedly in the waterfowl literature [*e.g.* 39-42], because there is no active transfer

of heat from the male to the eggs as ganders do not form brood patches. Rather, we revive the

term 'nest sitting' [4] as the appropriate label. Before female geese start actual incubation,

they adopt a posture when on the nest, in which the brood patch is not brought into contact

with the eggs, so there is no effective heat transfer to start embryonic development [43]. This

behaviour may serve to shield the nest and eggs from *e.g.* predation, inclement weather and/or

intraspecific brood parasitism. We suggest that male nest sitting is equivalent to this posture

at times when the female is on incubation recesses. To date, we cannot provide a definite

explanation of why this behaviour occurs, as neither the age of males seems to play a role, nor

could we show an increase in hatching success of nests, where males nest sat. At this moment,

we lack sample size for other more in depth analyses, such as when nest sitting occurs over

the incubation period or if males compensate with nest sitting for females taking more

incubation recesses. In the following we suggest three possible, not mutually exclusive,

functions of male nest sitting, to initiate future studies not only in barnacle geese, but also

other species.

The primary cause of hatching failure in ground nesting waterfowl is predation [*e.g.* 11,

44], thus several different strategies are employed for nest protection. These range from high

nest attendance [11], selecting sites where nests can either be successfully concealed or are

difficult for predators to reach [45], breeding in close proximity to more aggressive species to

take advantage of their nest defence [46] to actively put predators to rout [11]. Barnacle geese

historically nested high up on steep rock faces, particularly suitable to avoid terrestrial

predators. There, however, food is scarce for incubating females [9]. Breeding islands off the

coast used to be fairly safe from polar bears and Arctic foxes, but this turns out to now also be

changing [15, 20]. Furthermore, breeding on islands forces geese to nest in close proximity to

their main aerial predators, thus nest sitting of barnacle ganders might function in shielding

the nest from them. This is backed by our choosing the most aggressive males when attaching

transmitters, which might be bold enough to attack predators actively. What needs to be

investigated, is whether males adjust their behaviour to certain environmental conditions, *i.e.*

through behavioural plasticity, or whether nest sitting confers fitness benefits as an adaptive

response to breeding on islands [47, 48]. In the future, attention should be paid to how often

nest sitting of males occurs, how long the incubation recesses of the female last, what the

predation pressure is, and at which stage during incubation the behaviour is performed.

Another possible function of nest sitting by males is to control the temperature in the nest.

To some extent, nest insulation can reduce both the degree of egg cooling when the female is

absent and the time spent to rewarm the eggs after she returns from an incubation recess [49].

Avoidance of egg cooling is particularly important for Arctic breeding birds, where the

difference between optimal temperature for developing embryos (in geese: 37-38°C) and the

average air temperature during incubation (Ny-Ålesund average air temperature 2021: June

2.6°C, July 5.5°C, Norwegian Meteorological Institute) is striking [50]. Thus, eggs should be

covered more or less constantly. Indeed, Arctic geese adjust their incubation behaviour in

response to prevailing weather conditions, such as wind velocity, air temperature and solar

radiation [50, 51], remaining on the nest more continuously during inclement weather [51,

52]. Yet, egg cooling in the absence of the female might be overrated. In greater snow geese,

egg temperature throughout incubation remained high with only slight drops during the

frequent incubation recesses of females [43]. This might be, in part, ascribed to the female's

covering the nest with down [49]. Even under unfavourable weather conditions, egg

temperature drops during incubation recesses are probably nowhere near the minimal

temperatures, which endanger an embryo's development or survival. Rather, this will only

lengthen the incubation period until the eggs hatch [7, 53].

Overheating of unprotected eggs in the nest might become an issue in the Arctic in the

future. Also in Ny-Ålesund temperature has risen steadily since the 1970s [*e.g.* 54]. During

continuous incubation in climate control chambers, the critical temperature, at which no

embryos will survive, is 40.5°C [53]. Even if barnacle nests are exposed to direct sun, they

probably will, at present, not reach this temperature limit long enough for embryos to die in

the Arctic. Yet, common guillemots (*Uria aalge*), a high-latitude seabird, already showed an

increased probability of egg loss at higher temperatures in a colony in the Baltic Sea [55].

Whether nest sitting in barnacle ganders occurs predominantly on extremely warm or cold

days could be subject to further investigations. In sum, whereas nest sitting of barnacle

ganders undoubtedly will assist in temperature control of eggs in the nest, we suggest it to

only play a minor role, at least here and now.

If male nest sitting occurs during the period of egg laying, another possible function

might be to prevent intraspecific brood parasitism, which frequently occurs in barnacle geese

[27]. When conspecific barnacle geese attempt to dump eggs, females apply one of three

strategies. They either lay eggs in unguarded nests, leave parasitic eggs in close proximity to

the nest for the host female to retrieve, or attempt to lay an egg into the nest despite vigorous

defence by the host female. In contrast, males let those instances happen, behaving only

mildly aggressive, if at all, towards the parasitic female [56]. The tameness of the male

towards the parasitic female, when the host female is present, leads us to suggest that during

egg laying his propensity to nest sit in the absence of his mate is probably low. We cannot

answer this question adequately, however, because we placed cameras not until the host

female had started incubation, therefore lack photos from the laying period.

A question is whether nest sitting of barnacle ganders is a behaviour that has been

overlooked in the past or is a response to a changing environment. Although adult males are

slightly bigger than the females, barnacle geese are monomorphic. It is difficult to determine

the sex of an individual, even if it is marked, as rings are not visible when a bird sits on the

nest. It is comprehensible that a bird on the nest was assumed to be female, just as the

literature suggests. The relatively frequent occurrence of male nest sitting, *i.e.* 40% in our

limited number of individuals was unexpected to us. Even more, considering that we

synergise many years of detailed observation of goose behaviour in various species, we feel

confident that we are capable of distinguishing between the sexes convincingly.

4.2 'Male incubation' in waterfowl?

Found throughout the literature is the general assumption that in swans, and to some extent

whistling ducks, deviate from the female only incubation pattern in waterfowl, as here 'males

incubate' [39, 42, 57]. But are these males indeed incubating or is it a term that should be

avoided, as is leads to a misconception when compared with species, where incubation is truly

shared [3]?

In an early study, Hawkins already concluded that the well-developed male incubation

behaviour in tundra swans (*Cygnus columbianus columbianus*.) protected against egg

predation, provided some egg cooling benefits, but was not essential for successful embryo

development [42]. This led Henson & Cooper to apply the term 'nest sitting' in the case of

trumpeter swans *(C. buccinator*). They state that the behaviour was not 'as well developed as

in tundra swans', because it was independent of predator presence or adverse weather

conditions, despite males occasionally exhibiting some of the nest-settling motions

characteristic of incubating females [4]. Our extensive search of the literature revealed that in

most waterfowl, including three single observations in wood duck (*Aix sponsa*), mandarin

duck (*A. galericulata*) and fulvous whistling duck [*Dendrocygna bicolor*, 40, 41, 58], males

nest sit rather than incubate.

The two notable exceptions, for which the term incubation is warranted, are the black

swan (*C. atratus*, BLSW) and black-bellied whistling duck (*D. autumnalis*, BBWD*)*. Whereas

BLSW exhibit the Cygnidae typical long-term monogamy and biparental care, this is also true

for BBWD, unusual for Anatidae, although in some species males participate in attending

broods after hatching [59]. Besides male incubation, there are some features, which BLSW

and BBWD share. In both species, neither females nor males develop brood patches [60, 61],

thus either parent of each species can incubate in a similar manner. Also, both species are

known to re-nest after successful breeding attempts [39, 60, 62-64], whereas most other

waterfowl may attempt a second clutch only if the first one failed particularly during early

stages of incubation. We speculate that re-nesting might be the reason of both parents

incubating. Near constant incubation minimizes the time it takes for eggs to develop and

hatch, thereby possibly shortening the inter-clutch interval, but this would need to be tested in

mate removal experiments. In waterfowl with uniparental female care, mate removal did not

affect hatching success or body condition of the female [65, 66].

In conclusion, this is the first report in any goose species, where males expand paternal

duties beyond the established view, which during the breeding period is to protect the nest and

female in close vicinity of the nest. We have pictorial evidence that several individuals

displayed nest sitting behaviour. Although we may have unintentionally selected bolder

individuals, this should not divert from the fact that the behaviour exists. Both predation and

thermal risks are increased for eggs, which are not protected by parents, thus minimising

those risks are the main causes for successful hatching. In barnacle geese, we suggest nest

sitting of males to be a response to avian predator presence. We cannot exclude that

preventing egg cooling (and overheating) in the absence of the female may also play a role,

but consider this unlikely. Our final point is that in most waterfowl species, males do not

participate in incubation. Nest sitting is an unbiased term and should be used for cases, in

which no active warming of eggs has (yet) been demonstrated.

REFERENCES

[1] de Jong, M.E., Nicolaus, M., Fokkema, R.W.*, et al.* 2021. State dependence explains individual variation in nest defence behaviour in a long-lived bird. *J. Anim. Ecol.* **90**, 809-819. (doi:10.1111/1365-2656.13411).

[2] Ringelman, K.M. & Stupaczuk, M.J. 2013. Dabbling ducks increase nest defense after partial clutch loss. *Condor* **115**, 290-297. (doi:10.1525/cond.2013.120096).

[3] Deeming, D.C. & Reynolds, S.J. 2015. *Nests, Eggs, and Incubation*. Oxford, UK, Oxford University Press.

[4] Henson, P. & Cooper, J.A. 1992. Divison of labour in breeding trumpeter swans *Cygnus buccinator*. *Wildfowl* **43**, 40-48.

[5] Wang, J.M. & Beissinger, S.R. 2011. Partial incubation in birds: Its occurrence, function, and quantification. *The Auk* **128**, 454-466. (doi:10.1525/auk.2011.10208).

[6] Mitchell, C., Black, J.M. & Evans, M. 1998. Breeding success of cliff-nesting and island-nesting barnacle geese in Svalbard. *Norsk Polar. Skrift.* **200**, 141-146.

[7] Tombre, I.M. & Erikstad, E. 1996. An experimental study of incubation effort in high-Arctic barnacle geese. *J. Anim. Ecol.* **65**, 325-331.

[8] Prop, J., Aars, J., Bårdson, B.-J.*, et al.* 2015. Climate change and the increasing impact of polar bears on bird populations. *Frontier in Ecology and Evolution* **3**, 33. (doi:10.3389/fevo.2015.00033).

[9] Prop, J., Eeerden, M.R. & van Drent, R.H. 1984. Reproductive success in the barnacle goose *Branta leucopsis* in relation to food exploitation on the breeding grounds, western Spitsbergen. *Norsk Polar. Skrift.* **181**, 87-117.

[10] Schreven, K.H.T., Stolz, C., Madsen, J.*, et al.* 2021. Nesting attempts and success of Arctic-breeding geese can be derived with high precision from accelerometry and GPS-tracking. *Anim. Biotelem.* **9**, 25. (doi:10.1186/s40317-021-00249-9).

[11] Samelius, G. & Alisauskas, R.T. 2001. Deterring arctic fox predation: The role of parental nest attendance by lesser snow geese. *Can. J. Zool.* **79**, 861-866.

[12] Dittami, J.P., Kennedy, S. & Thomforde, C. 1979. Observations on barnacle goose breeding, *Branta leucopsis*, in Spitsbergen 1975. *J. Ornithol.* **120**, 188-195.

[13] Speelman, F.J.D., Hammers, M., Komdeur, J.*, et al.* 2022. Nest defence behaviour is similar between pair members but only male behaviour predicts nest survival in barnacle geese. *J. Avian Biol.* **2022**, e02982. (doi:10.1111/jav.02982).

[14] Clermont, J., Réale, D. & Giroux, J.-F. 2019. Similarity in nest defense intensity in Canada goose pairs. *Behav. Ecol. Sociobiol.* **73**. (doi:10.1007/s00265-019-2719-3).

[15] Tombre, I.M., Mehlum, F. & Loonen, M.J.J.E. 1998. The Kongsfjorden colony of
barnacle geese: Nest distribution and the use of breeding islands 1980-1997. *Norsk Polar.*
Skrift. **200**, 57-65.

[16] Pavlova, O., Gerland, S. & Hop, H. 2019. Changes in sea-ice extent and thickness in
Kongsfjorden, Svalbard (2003-2016). In *The Ecosystem of Kongsfjorden, Svalbard. Advances*
in Polar Ecology (eds. H. Hop & C. Wiencke), pp. 105-136, Springer.

[17] Gilg, O., Istomina, L., Heygster, G.*, et al.* 2016. Living on the edge of a shrinking
habitat: The ivory gull, *Pagophila eburnea*, an endangered sea-ice specialist. *Biol. Lett.* **12**,
20160277. (doi:10.1098/rsbl.2016.0277).

[18] Dørum, H. 2019. Effects of climate change and diminishing sea ice on common eider
breeding distribution in Kongsfjorden, Svalbard [Master]. Trondheim, Norway, Norwegian
University of Science and Technology.

[19] Smith, P.A., Elliott, K.H., Gaston, A.J.*, et al.* 2010. Has early ice clearance increased
predation on breeding birds by polar bears? *Polar Biol.* **33**, 1149-1153. (doi:10.1007/s00300-
010-0791-2).

[20] Andersen, M. & Aars, J. 2016. Barents Sea polar bears (*Ursus maritimus*): Population
biology and anthropogenic threats. *Polar Research* **35**, 26029. (doi:10.3402/polar.v35.26029
).

[21] Fuglei, E., Øritsland, N.A. & Prestrud, P. 2003. Local variation in Arctic fox abundance
on Svalbard, Norway. *Polar Biol.* **26**, 93-98. (doi:10.1007/s00300-002-0458-8).

[22] Hanssen, S.A., Moe, B., Bårdsen, B.J.*, et al.* 2013. A natural antipredation experiment:
Predator control and reduced sea ice increases colony size in a long-lived duck. *Ecol. Evol.* **3**,
3554-3564. (doi:10.1002/ece3.735).

[23] Lorenz, K. 1991. *Here I Am - Where Are You?* New York, NY, USA, Hartcourt Brace
Jovanovich.

[24] Jones, R.E. 1971. The incubation patch of birds. *Biol. Rev.* **46**, 315-339.
(doi:10.1111/j.1469-185X.1971.tb01048.x).

[25] Forslund, P. & Larsson, K. 1992. Age-related reproductive success in the barnacle goose.
The Journal of Animal Ecology **61**, 195. (doi:10.2307/5522).

[26] Lamprecht, J. 1989. Mate guarding in geese: Awaiting female receptivity, protection of
paternity or support of female feeding. In *The Sociobiology of Sexual and Reproductive*
Strategies (eds. A.E. Rasa, C. Vogel & E. Voland). Dordrecht, NL, Springer Science and
Business Media.

[27] Black, J.M., Prop, J. & Larsson, K. 2014. *The Barnacle Goose*. London, UK,
Bloomsbury Publishing.

[28] Hübner, C.E., Tombre, I.M. & Erikstad, K.E. 2002. Adaptive aspects of intraclutch egg-
size variation in the high Arctic barnacle goose (*Branta leucopsis*). *Can. J. Zool.* **80**, 1180-
1188.

[29] Alsos, I.G., Elvebakk, A. & Gabrielsen, G.W. 1998. Vegetation exploitation by barnacle
geese *Branta leucopsis* during incubation on Svalbard. *Polar Research* **17**, 1-14.

[30] Ahmad, I.M. & Li, D. 2023. More than a simple egg: Underlying mechanisms of cold
tolerance in avian embryos. *Avian Res.* **14**, 100104. (doi:10.1016/j.avrs.2023.100104).

[31] Tombre, I.M., Erikstad, E. & Bunes, V. 2012. State-dependent incubation behaviour in
the high Arctic barnacle geese *Polar Biol.* **35**, 985-992. (doi:10.1007/s00300-011-1145-4).

[32] Owen, M. & Wells, R.L. 1979. Territorial behaviour in breeding geese - A re-
examination of Ryder's hypothesis. *Wildfowl* **30**, 20-26.

[33] Fenstad, A.A., Bustnes, J., Lierhagen, S.*, et al.* 2017. Blood and feather concentrations of
toxic elements in a Baltic and an Arctic seabird population. *Marine Pollution* **114**, 1152-1158.
(doi:j.marpolbul.2016.10.034).

[34] Loonen, M.J.J.E., Oosterbeek, K. & Drent, R.H. 1997. Variation in growth of young and
adult size barnacle geese *Branta leucopsis*: Evidence for density dependence. *Ardea* **85**, 177-
192.

[35] Webster, M.M. & Rutz, C. 2020. How STRANGE are your study animals? *Nature* **582**,
337-340. (doi:10.1038/d41586-020-01751-5).

[36] Blumstein, D.T., Samia, D.S.M. & Cooper, W.E. 2016. Escape behavior: Dynamic
decisions and a growing consensus. *Curr. Opin. Behav. Sci.* **12**, 24-29.
(doi:10.1016/j.cobeha.2016.08.006).

[37] de Jong, M.E., Wetherbee, R. & Loonen, M.J.J.E. 2019. Effects of fleas on nest success
of Arctic barnacle geese: Experimentally testing the mechanism. *J. Avian Biol.* **50**, e01944.
(doi:10.1111/jav.01944).

[38] R Core Team. 2021. R: A language and environment for statistical computing. (1.13 ed.
Vienna, Austria, R Foundation for Statistical Computing.

[39] Brugger, C. & Taborsky, M. 1994. Male incubation and its effect on reproductive
success in the black swan, *Cygnus atratus*. *Ethology* **96**, 138-146. (doi:10.1111/j.1439-
0310.1994.tb00889.x).

[40] Bruggers, R.L. 1979. Nesting patterns of captive mandarin ducks. *Wildfowl* **30**, 45-54.

[41] Rollin, N. 1957. Incubation by drake wood duck in eclipse plumage. *The Condor* **59**,
263-265. (doi:10.2307/1364656).

[42] Hawkins, L.L. 1986. Nesting behaviour of male and female whistling swans and
implications of male incubation. *Wildfowl* **37**, 5-27.

[43] Poussart, C., Larochelle, J. & Gauthier, G. 2000. The thermal regime of eggs during
laying and incubation in greater snow geese. *The Condor* **102**, 292-300.
(doi:10.1093/condor/102.2.292).

[44] Peterson, S.H., Ackerman, J.T., Keating, M.P.*, et al.* 2022. Predator movements in
relation to habitat features reveal vulnerability of duck nests to predation. *Ecol. Evol.* **12**.
(doi:10.1002/ece3.9329).

[45] Schranck, B.W. 1972. Waterfowl nest cover and some predation relationships. *J. Wildl.*
Manag. **36**, 182. (doi:10.2307/3799210).

[46] Quinn, J.L., Prop, J., Kokorev, Y.*, et al.* 2003. Predator protection or similar habitat
selection in red-breasted goose nesting associations: Extremes along a continuum. *Anim.*
Behav. **65**, 297-307. (doi:10.1006/anbe.2003.2063).

[47] Madsen, J., Jaspers, C., Frikke, J.*, et al.* 2019. A gloomy future for light-bellied brent
geese in Tusenøyane, Svalbard, under a changing predator regime. *Polar Research* **38**, 3393.
(doi:10.33265/polar.v38.3393).

[48] Mery, F. & Burns, J.G. 2010. Behavioural plasticity: An interaction between evolution
and experience. *Evolutionary Ecology* **24**, 571-583. (doi:10.1007/s10682-009-9336-y).

[49] Thompson, S.C. & Raveling, D.G. 1988. Nest insulation and incubation constancy of
Arctic geese. *Wildfowl* **39**, 124-132.

[50] Poussart, C., Gauthier, G. & Jacques, L. 2001. Incubation behaviour of greater snow
geese in relation to weather conditions. *Can. J. Zool.* **79**, 671-678.

[51] Harvey, J.M. 1971. Factors affecting blue goose nesting success. *Can. J. Zool.* **49**, 223-
234. (doi:10.1139/z71-03).

[52] Elkins, N. 2004. Breeding: Incubation. In *Weather and Bird Behaviour* (ed. N. Elkins),
pp. 101-104. London, UK, T.& A.D. Poyser.

[53] White, F.N. & Kinney, J.L. 1974. Avian incubation: Interactions among behavior,
environment, nest, and eggs result in regulation of egg temperature. *Science* **186**, 197-115.
(doi:10.1126/science.186.4159.1).

[54] Wei, T., Ding, M., Wu, B.*, et al.* 2015. Variations in temperature-related extreme events
(1975–2014) in Ny-Ålesund, Svalbard. *Atmos. Sci. Let.* **17**, 102-108. (doi:10.1002/asl.632).

[55] Olin, A.B., Dück, L., Berglund, P.A.*, et al.* 2024. Breeding failures and reduced nest
attendance in response to heat stress in a high-latitude seabird. *Mar. Ecol. Prog. Ser.* **737**,
147-160. (doi:10.3354/meps14244).

[56] Forslund, P. & Larsson, K. 1995. Intraspecific nest parasitism in the barnacle goose:
Behavioural tactics of parasites and hosts. *Anim. Behav.* **50**, 509-517.

[57] Scott, D. 1977. Breeding behaviour of wild whistling swans. *Wildfowl* **28**, 101-106.

[58] Flickinger, E.L. 1975. Incubation by a male fulvous tree duck. *Wilson Bull.* **87**, 106-107.

[59] McKinney, F. & Brewer, G.L. 1989. Parental attendance and brood care in four species
of Argentine dabbling ducks. *Condor* **91**, 131-138. (doi:10.2307/1368156).

[60] Bolen, E.G. & Smith, E.N. 1979. Notes on the incubation behavior of black-bellied
whistling ducks. *Prairie Nat.* **11**, 119-123.

[61] Skutch, A.F. 1976. *Parent Birds And Their Young*, University of Texas Press; 503 p.

[62] Delnicki, D.E. 1973. Renesting, incubation behavior, and compound clutches of the
black-bellied tree duck in southern Texas. Lubbock, TX, USA, Texas Tech University.

[63] James, J.D., Thompson, J.E. & Ballard, B.M. 2012. Evidence of double brooding by
black-bellied whistling-ducks. *Wilson J. Ornithol.* **124**, 183-185. (doi:10.1676/11-082.1).

[64] Coleman, J.T. 2014. Breeding biology of the black swan *Cygnus atratus* in southeast
Queensland, Australia. *Wildfowl* **64**, 217-230.

[65] Manlove, C.A. & Hepp, G.R. 1998. Effects of mate removal on incubation behavior and
reproductive success of female wood ducks. *Condor* **100**, 688-693.

[66] LeSchack, C.R., Afton, A.D. & Alisauskas, R.T. 1998. Effects of male removal on
female reproductive biology in Ross' and lesser snow goose. *Wilson Bull.* **110**, 56-64.
489

491
492 **Table 1:** Identities of 15 focal pairs, whose nests were monitored with wildlife cameras in 2021. For each
493 individual, we provide codes of unique colour and aluminium rings, and if they were fitted with a transmitter, the
494 neck collar ID. Except two individuals, which were ringed as goslings (*), the minimum age is given ($\bar{x}$ age $\pm$
495 SD: males 9.4 $\pm$ 3.31; females 10.4 $\pm$ 4.46). Whether or not males sat on nests is also indicated.
496

Male colour ID	Male ID	Male neck collar ID	Age (years)	Male nest sitting	Female colour ID	Female ID	Female neck collar ID	Age (years)
oFYJ	CA46944	201913	16	yes	oFVD	CA46857		10
gZUN	CA45605	201915	9	yes	oFLC	CA45810		12
alur	CA47072	201917	8	no	yBSV	CA45997		4
oFZY	CA46942	201918	10	yes	gSNF	CA36931		19
oFIJ	CA45800	201924	12	yes	yBHJ	CA45891		5
oFIA	CA45795	201930	12	yes	gLDC	CA41238		14
yBID	CA46840	201932	7	no	yBIB	CA44228		14
oFZX	CA46941		10	no	gTTL	CA41113	201912	16*
yBYI	CA45746		7	yes	yAUF	CA45744	201919	7
yAXY	CA45703		8	no	yBHY	CA46848	201925	5*
gLHX			2	no	yAVV	CA45736	201926	7
yAZT	CA45715		8	no	oFXV	CA45632	201927	10
yAXC	CA45942		8	no	yAPH	CA45929	201931	8
gTCZ	CA46957		10	no	yAXV	CA45701	201934	8
yBYX	CA41397		14	no	yBYV	CA41400	201935	14

Figure 1: An example of one barnacle gander nest sitting. Photos taken on June 18th, 2021

between 15:48 and 16:28. Panel A) Female oFVD without a neck collar is incubating. Panel

B) The female left the nest. Panel C) Male alur with neck collar approaches the nest. Panel D)

He sits on the nest with the neck collar clearly visible. Panel E) The female returns to the nest.

Panel F) She resumes incubation. Temperature and time are shown in the lower right hand

corner of each photo.

Figure 2: Features, which might influence nest sitting in barnacle ganders. a) Age did not

affect the likelihood of male to nest sit (Student's *t*-test: $p > 0.05$). b) Males, which were fitted

with a neck collar, were more likely to nest sit than males without (Fisher's Exact Test: $p =$

0.041). c) The method of catching for transmitter deployment, either near the nest or the

during moult catch, had no effect on nest sitting propensity of males (Fisher's Exact Test: $p >$

0.05). d) Whether eggs hatched or did not hatch was independent of males' nest sitting

(Fisher's Exact Test: $p > 0.05$).

Figure 1

518
519   Figure 2