Power-law adaptation in the presynaptic vesicle cycle

Fabian A. Mikulasch[1], Svilen V. Georgiev[2,3], Lucas Rudelt[1,6],
Silvio Rizzoli[2,4,5], Viola Priesemann[1,5,6]

1 Max-Planck-Institute for Dynamics and Self-Organization, Göttingen, Germany

2 University Medical Center Göttingen, Institute for Neuro- and Sensory Physiology, Germany

3 International Max Planck Research School for Neuroscience, Göttingen, Germany

4 Biostructural Imaging of Neurodegeneration (BIN) Center, Göttingen, Germany

5 Excellence Cluster Multiscale Bioimaging, Göttingen, Germany

6 Institute for the Dynamics of Complex Systems, University of Göttingen, Göttingen, Germany

After synaptic transmission, fused synaptic vesicles are recycled, enabling the synapse to recover its capacity for renewed release. The recovery steps, which range from endocytosis to vesicle docking and priming, have been studied individually, but it is not clear what their impact on the overall dynamics of synaptic recycling is, and how they influence signal transmission. Here we model the dynamics of vesicle recycling and find that the multiple timescales of the recycling steps are reflected in synaptic recovery. This leads to multi-timescale synapse dynamics, which can be described by a simplified synaptic model with 'power-law' adaptation. Using cultured hippocampal neurons, we test this model experimentally, and show that the duration of synaptic exhaustion changes the effective synaptic recovery timescale, as predicted by the model. Finally, we show that this adaptation could implement a specific function in the hippocampus, namely enabling efficient communication between neurons through the temporal whitening of hippocampal spike trains.

Introduction

One of the main bottlenecks of synaptic transmission is the recovery of synaptic vesicles after vesicle release, since it determines the availability of releasable vesicles. By now there is ample data on the underlying biochemical processes that proceed through several stages of vesicle recovery (Fig 1A). After release, vesicles are taken back up through endocytosis, reloaded with neurotransmitters, docked to the membrane, and finally primed for repeated release. The dynamics of these recovery processes have been measured using a variety of experimental setups that elucidated their molecular pathways, their speed, and the numbers of vesicles involved [1, 2]. Yet, it remains poorly understood how these individual stages of recovery interact, and ultimately affect synaptic signal transmission and synaptic computation.

An important feature of these recovery processes is that they operate on vastly different timescales, and thus might affect synaptic recovery in different ways. These timescales range from the orders of $100\,\mathrm{ms}$ (vesicle priming), over $1\,\mathrm{s}$ (docking and loading), to $10\,\mathrm{s}$ (endocytosis). While synapses typically recover within seconds [3], previous research has shown that synaptic recovery can slow down under strong stimulation [4, 5]. For example, in hippocampal neurons, this slow form of recovery can take tens of seconds, up to minutes [5]. One explanation of this slowing down of recovery is the depletion of vesicles that are available for release and a fallback on the slowest recovery process (endocytosis) [4]. Thus, there is evidence that both fast and slow recovery timescales play a role under different

conditions.

In contrast to these findings, models of the function of synaptic adaptation typically simplify the recovery process to a single recovery timescale [6, 7, 8, 9]. This simplification of the recovery dynamics allowed for important insights into how synaptic depression could shape neural computations, for example, through temporal filtering of spike trains, adaptation, and gain control [10]. Yet, the presence of processes with timescales across several orders of magnitude in the recovery pathway could lead to different implications for synaptic function, which are not captured by these simplified concepts.

One challenge in achieving an understanding of the functional aspects of multiple timescales in vesicle recovery is that the dynamics of their interaction are not well explored. Previous studies typically modeled vesicle pool dynamics only with (up to) two recovery timescales and found that the overall recovery in different conditions might be described by a single or double exponential function [3, 4, 11, 12]. However, this leaves open the possibility that more processes, with different timescales, could shape synaptic dynamics. Furthermore, it is not clear whether the parameters of the vesicle dynamics, which were measured in isolation [13, 14, 15], are consistent with the overall recovery dynamics that can be measured experimentally.

To address the question of whether the diverse timescales of different vesicle recovery steps play a substantial role in synaptic transmission, we employ here both theoretical modeling and an analysis of cultured neurons. First, we analyze a multistage model of synaptic recovery in hippocampal synapses, which is based on timescales of recovery processes that were measured previously [13]. We show that, in this model, each of the timescales plays a role in shaping the overall recovery from release, leading to multi-timescale adaptation dynamics. We furthermore show that this form of depression can be described by a simplified synaptic model with effective 'power-law' adaptation. We then employ an experimental setup that allows us to measure vesicle release in hundreds of synapses in parallel. With this setup, we verify that the measured parameters in our model are close to the parameters that fit the experimental data.

Finally, to elucidate how power-law adaptation could contribute to neural information processing, we show that it specifically tunes down synaptic responses to strong, slow frequency fluctuations of presynaptic spiking. Our results suggest that hippocampal synapses are designed to efficiently transmit hippocampal activity, where such long timescale dependencies can be found [16, 17].

Results

Multi-stage model of synaptic recovery   To simulate the presynaptic vesicle cycle, we employed the model proposed in [13]. This model incorporates the five most prominent stages of vesicle recycling in the presynapse as discrete vesicle pools: Previously released vesicles are recovered by endocytosis, loaded with neurotransmitters, docked to the membrane, and ultimately primed for release (Fig 1A). Upon a spike, primed vesicles are released with a certain probability p_{fuse} (in the main results $p_{\text{fuse}} = 0.2$). In between spikes, vesicles transition stochastically from pool P_i to pool P_j with a filling level-dependent transition rate

$$r_{ij} = \frac{c}{\tau_{ij}} \frac{P_i}{P_i^{max}} \left(1 - \frac{P_j}{P_j^{max}} \right),$$

where $c = 4$ ensures that $r_{ij} = 1/\tau_{ij}$ if pools are half full, and P_i^{max} is the maximum capacity of pool P_i. An exception is the transition from the fused to the recovery pool via endocytosis, which happens with a fixed rate r_{reform} as long as the fused pool is not empty and the recovery pool is not full.

Fig 1. Model of the presynaptic vesicle cycle. A Model of the presynaptic vesicle cycle, as proposed in [13]. Vesicles transition stochastically from pool to pool with a given timescale. Timescales of transitions are taken from experimental measurements [13] (Appendix) and become orders of magnitudes larger the further they are away from the release stage (Primed pool). Similarly, the maximum number of vesicles in pools (numbers in circles) is larger for pools farther away from release. Upon an input spike, vesicles in the primed pool are emitted with probability p_{fuse}. Subsequently, vesicles can age with probability p_{age} and are dynamically resupplied if the synapse lacks vesicles. We would like to point out that these very slow processes have limited significance for the analysis in this work. **B,C** Example of how spikes result in the release of vesicles (top) and how vesicle pool levels change over time (bottom). Pools close to release operate on a fast timescale (B), whereas pools far away operate on slow timescales (C).

Intriguingly, the parameters of this model, which have been determined experimentally from cultured hippocampal neurons [13], have a very particular ordering on the cycle (Fig 1A). Transition timescales between pools tend to increase with distance from release, from vesicle priming, which can be as fast as 25 ms, up to vesicle endocytosis and reformation, which takes several seconds for a single vesicle. Similarly, the upper limit for the sizes of pools increases with the distance to release. Together, these parameters result in very different recovery speeds of vesicle pools, where pools close to release recover on fast timescales in the order of 100 ms, whereas pools further away operate on timescales as slow as 10 s up to minutes (Fig 1B,C).

Power-law adaptation in the vesicle cycle   To investigate whether adaptation is dominated by one of these recovery timescales, or if all timescales play a role in adaptation, we set up several simulation experiments. First, we tested how the model synapse depressed under constant strong stimulation. We stimulated the model with a regular 50 Hz spike train (Fig 2A) and measured how the average number of released vesicles per spike changes over time. To observe these changes on both short and long timescales, averages were taken over spikes grouped in log-spaced intervals. We found that the average of released vesicles decreased rapidly at the beginning of stimulation, but later this decrease slowed down and lasted up until about 100 s (Fig 2B). In comparison, models with a single timescale of vesicle recovery showed only a very short phase of adaptation, after which vesicle release stayed constant. This indicates that, in our model of vesicle recovery, processes operating on long, medium, and short timescales interact to shape synaptic depression.

To reach a better understanding of how the release of a vesicle impacts the probability of release in the future, we fitted a generalized linear model (**GLM**) to our multi-stage model of vesicle recovery (Fig 2D, see Methods). The GLM is an effective model of vesicle release, in which every spike triggers a vesicle release with a certain probability, but previous releases reduce this probability depending on an effective kernel $\kappa(\Delta t)$ (given the release was Δt in the past). This adaptation kernel thus captures how releases reduce the future release probability on average. For example, a model similar to the classical Tsodyks-Markram model [18, 19] without facilitation would employ an exponential kernel $\kappa(\Delta t) \propto \exp(-\Delta t/\tau)$. To estimate $\kappa(\Delta t)$, we stimulated the vesicle cycle model with highly variable spike trains (with 1/f spectral power, see Methods) and measured the resulting releases. We then fitted this input-output behavior with the GLM using maximum likelihood estimation (SI Fig S15). The resulting effective kernel $\kappa(\Delta t)$ showed that previous vesicle releases have a measurable effect on release probability on a wide range of timescales, and could be well approximated by a power law $\kappa(\Delta t) \propto \Delta t^{-0.3}$, up to about 120 s (Fig 2E).

Experimental validation of model dynamics   Our simulation experiments indicated that, under strong stimulation, vesicle releases can significantly impact the efficacy of the synapse, even minutes into the future. To test this prediction, we set up an *in-vitro* experiment with rat hippocampal cultures to measure the long timescale dependence of synaptic exhaustion on vesicle releases. In our experimental protocol we first aimed to partially exhaust synaptic vesicle pools by strong stimulation, let the synapse recover while pausing stimulation, and then tested the final efficacy of the synapse (Fig 3A). By varying the time of exhaustion and pausing over several orders of magnitude, we could then get an understanding of the dependence of synaptic depression over a range of timescales (Fig 3B).

As a measure of synaptic release, we turned to super-ecliptic pHluorin, a pH-sensitive fluorophore that is introduced within the lumen of synaptic vesicles, where it is quenched by the acidic milieu, and reports their exocytosis by un-quenching upon reaching the neutral extracellular buffer [21]. Typical pHluorin experiments involve the expression of synaptic vesicle proteins that carry this molecule on their intravesicular chains. To avoid any effects of genetic manipulation and protein expression, we turned to antibodies against the intravesicular domain of synaptotagmin 1, which carried secondary nanobodies [22] conjugated to pHluorin molecules. The synaptotagmin antibodies were loaded within

Fig 2. Power-law adaptation resulting from multi timescale dynamics of the presynaptic vesicle cycle. A Reaction of the vesicle cycle when stimulated with a 50 Hz signal (Color code as in Fig 1B,C). **B** The average number of vesicles emitted per spike decreases approximately linearly in log-log scale up until $\sim 10 - 100$ s, which corresponds to a power-law decay. In comparison, models with a single timescale of vesicle recovery show a rapid decay that levels out after 1 s. Inset shows the data in linear scale from 0 to 5 s. Error bars denote 95% bootstrapping confidence intervals estimated from 100 experiments; inset shows the same data but linear axes (also in C and E). **C** The resources used (average number of vesicles emitted up to that time) increase rapidly for the single timescale model with fast recovery and slowly for the single timescale model with slow recovery. The full model strikes a balance between coding fidelity on short timescales and resource efficiency for strong stimulation on long timescales. **D** We used a simple response model to fit the full vesicle cycle model. An effective kernel κ mediates negative feedback on the release probability after a vesicle release. **E** When fitted to responses to simulated spike trains (see Methods) the kernel can be approximated well by a power law $\kappa(\Delta t) \propto \Delta t^{\alpha}$ with exponent $\alpha \approx -0.3$ up to ~ 120 s. This means that a single vesicle release has a measurable effect on future releases more than 2 minutes into the future. For the single timescale model with fast recovery (gray), this decay is much more rapid. The inset shows the kernels in linear scale up to $\Delta t = 8$ s. Kernel bins for very short and long Δt that are difficult to estimate from the data have been excluded in this plot.

Fig 3. Hippocampal synapses in vitro show stimulus-dependent recovery timescales.
A Cultured synapses were first exhausted using 20 Hz stimulation, recovered for a variable time,
after which the synaptic efficacy was tested. **B** From our model, we expected that longer exhaustion
times result in longer recovery timescales. **C** Fluorescence images of synaptic activity recorded
with super-ecliptic pHluorin during baseline activity (left), strong stimulation during the exhaustion
period (middle) and test stimulation in the test period (right). For display purposes, the images
were deconvolved using the Richardson-Lucy algorithm from the DeconvolutionLab2 [20]. **D** Example
fluorescence traces recorded from a single region of interest as an average across multiple synapses for
an exhaustion time of 4 s, and a pause time of 10 s (top) or 100 s (bottom). **E** The recordings indeed
show that longer exhaustion time leads to slower recovery. This can be fitted well by the full vesicle
cycle model. **F** A single timescale recovery model cannot fit this difference. The effective recovery
timescale of the fitted model is ≈ 17 s, which does not change depending on the exhaustion time.
E,F show mean change in $\Delta F/F$ during the test stimulus. Shaded areas denote 95% bootstrapping
confidence intervals, bars denote 95% confidence intervals based on the posterior distribution of the
Bayesian models.

all of the active vesicles, by prolonged incubations, before the analysis [23]. This procedure enables the analysis of synaptic release dynamics and should provide virtually identical results to conventional pHluorin measurements [24].

Our experiments indeed showed that synaptic depression is captured well by the model (Fig 3E). As predicted from our model, the speed of recovery changed based on the duration of stimulation, with long stimulation drastically slowing down the recovery. This behavior could not be reproduced by a model with single timescale recovery (Fig 3F). We showed this by fitting the model parameters to the experimental data using Bayesian inference with MCMC (see Methods). The fitted parameters of the full model remained close to the previously measured parameters (see SI Fig S7), showing that only minor adjustments are needed to be consistent with the data. To fit the data, changes in the mean parameters occurred mainly in vesicle pool sizes (primed pool from 4 to 5.5 vesicles, docked pool from 20 to 14 vesicles) and release probability (from 0.2 to 0.37). All of these changes are well within the measured parameters for cultured hippocampal synapses (e.g., [25, 26]).

Temporal whitening of hippocampal spike trains through power-law adaptation   Finally, we considered what functional advantage power-law adaptation in synapses could bring for synaptic information transmission. To increase the efficiency of neural communication, it has long been proposed that redundant information should be discarded, which could be implemented by synaptic adaptation [7]. Natural signals often contain correlations on many timescales, in which case redundancy reduction, i.e., neural adaptation, should also operate on many timescales [29].

Formally, for temporal signals, the goal of efficient coding is to reduce the auto-correlation in the encoded signal. This is commonly illustrated by showing that the encoding flattens the power spectral density (**PSD**) of the signal (i.e., achieving the PSD of a temporally uncorrelated white noise signal, which is also called temporal whitening) [29]. To test whether the vesicle cycle model could implement this coding strategy, we stimulated it with simulated spike trains exhibiting correlations on long and short timescales, that is, spike trains with 1/f PSD (Fig 4A, see Methods). Our simulations showed that the depression dynamics of the full vesicle cycle model partially flattened the PSD of the signal, meaning it partially removed correlations over several orders of magnitude of timescales (Fig 4B). In comparison, the single timescale vesicle cycle model only flattened the PSD for high frequencies. One intuitive way to understand whitening is that it dampens fluctuations in the input rate on slow timescales, which can be extremely large (Fig 4C). During periods of high activity, where spikes are very predictable, average releases per spike are dynamically tuned down, which leaves more resources for periods of low activity, with less predictable spikes (Fig 4C). Thus, power-law adaptation in the presynaptic vesicle cycle could enable efficient synaptic information transmission for spike trains with long-tail auto-correlation functions.

Based on these results we wondered whether this form of efficient coding could be relevant in the rat hippocampus. To test this, we analyzed long recordings of CA1 activity of freely behaving rats (see Methods), and simulated the responses of the vesicle cycle under this stimulation (Fig 4E,F). We observed that hippocampal spike trains have approximately $f^{-0.5}$ PSD in the range of frequencies in which the synapse operates, which, similarly to $1/f$ PSD, is associated with long tail autocorrelations [30]. In comparison, the vesicle release train generated by the response of the vesicle cycle model is nearly flat (Fig 4F). This effect can also be seen in the measured coefficient of variation (**CV**) of the activities (where CV=1 is associated with Poisson activity and maximum information rate). Measured spike trains had a mean CV of 3.4 (1.4 to 15), while simulated releases had mean CVs of 1.8 (0.9 to 3.3) and 2.7 (1.6 to 7.7) for the full and the single timescale model, respectively. Overall, this shows that, not only is the vesicle cycle able to remove long tail correlations from the input spike trains, but it is also approximately tuned to the type of correlations found in hippocampal activity.

Fig 4. Power-law adaptation enables temporal whitening of hippocampal spike trains.
A Reaction of the vesicle cycle when stimulated with a spike train with 1/f power spectrum and an average rate of 1 Hz (Color code as in Fig 1B,C). **B** The response of the vesicle cycle model shows a flatter spectrum when compared to the original 1/f spectrum of the input. In comparison, the model with a single vesicle recovery timescale only attenuates high-frequency fluctuations. Power spectra are plotted as normalized densities. The full and single timescale models have comparable average power, with average release rates of 0.26 Hz and 0.29 Hz, respectively. **C,D** Power-law adaptation dynamically adjusts the releases per spike to the input rate. Here we demonstrate this behavior in the strong input regime (~ 4 Hz), which allows for clear visualization. During high activity, releases per spike are selectively reduced, mainly mediated by slow endocytosis. During low activity, they are enhanced when compared to a single timescale model with similar average release rate. The single timescale model closely follows the linear prediction from input spikes (thin red line). Rates were estimated by convolving the spike and release trains with kernels (see Methods). **E** Reaction of the vesicle cycle when stimulated with a spike train recorded from hippocampus CA1 [27]. **F** Hippocampal spike trains show an approximate $f^{-0.5}$ power spectrum over the relevant range. Here the full model nearly flattens the spectrum. Spectra for individual spike trains (transparent lines) behave similarly to their mean, showing that the power law is not a result of averaging spectra of spike trains with different single timescales [28].

Discussion

Here we analyzed a model of synaptic vesicle release based on experimentally determined parameters of the most prominent steps of the vesicle recycling machinery, which allowed us to generate large amounts of data for the input-output function of the presynapse. Our analysis showed that the interplay of the different recycling steps can lead to multi-timescale adaptation, which can be described by a synaptic model with an effective power-law depression kernel up to about 120 s. These results motivated us to specifically test for the presence of both, short and long timescales of synaptic recovery in cultured hippocampal neurons. Our experiments demonstrated that synaptic recovery can indeed operate on a range of timescales, even up to minutes, depending on the stimulation protocol. Finally, we showed that synaptic multi-scale adaptation is effective in attenuating high-amplitude, low-frequency fluctuations in hippocampal spike trains, which enables an efficient allocation of synaptic resources for synaptic transmission.

Power-law adaptation through the interaction of different recovery processes   The multi-timescale depression we described is a result of the particular ordering of vesicle pool sizes and timescales in the vesicle cycle. Under strong stimulation, smaller vesicle pools close to the release site are depleted first, whereas larger pools further away can continue to supply vesicles for a longer time. This leads to a gradual shift of the recovery bottleneck from later (faster) to earlier (slower) stages of recovery (Fig 2, SI Fig S5). The interaction of these processes in our model leads to effective depression and recovery timescales up to the order of minutes (Fig 2B,E)—nearly an order of magnitude slower than the slowest recovery rate in the model.

Previously, several proposals have been made to explain why recovery slows down under strong stimulation. Candidate mechanisms include the long-lasting depletion of vesicle pools [3, 4, 31], the blocking of release sites [5], or the replacement of tethered vesicles on the presynaptic membrane [32]. Our results corroborate the idea that the depletion of vesicle pools is a central limiting factor of synapses under strong stimulation. Vesicle cycle parameters only had to be slightly tuned to fit the experimental data (SI Fig S7), meaning that the timescales and sizes of vesicle pools that were measured previously yield a plausible explanation of the overall timescales of synaptic recovery.

In comparison to the full vesicle cycle model, a single timescale recovery model was not able to fit the experimental data as well (Fig 3). This is also the case when comparing the models using Bayesian model comparison, which takes the number of fitted parameters of the models into account (SI Fig S10). It has to be noted that a model with two timescales of vesicle recovery fits the data only slightly worse than the full model, and is not significantly worse under Bayesian model comparison (SI Fig S9,S10). The caveat of such a model, however, is that the fitted parameters can not easily be related to vesicle dynamics in the presynapse, which is possible for the full model. Furthermore, since dynamics over multiple orders of magnitude can be difficult to measure in a single experimental setup, we here specifically tested for medium and long timescales of synaptic recovery. The two timescales model succeeded in recovering the medium and long timescales (SI Fig S9), but it would fail to also incorporate the faster recovery timescales that have been observed in other experiments [11, 33].

Functional implications   Synaptic transmission is the most energy-intensive process in neural computation and is estimated to consume about 40% of the total cerebral energy budget [34, 35]. Consequently, synapses face a strong energy/information trade-off, which often means that synapses should actively reduce the number of uninformative releases. Previous studies have shown that the depression of synaptic release can improve the amount of information transmitted per release [6, 7, 8, 36, 37], by dynamically lowering synaptic responsiveness when current spikes can be predicted from past spikes. This is a general mechanism that enables the efficient encoding of events, which, for

example, also applies to neural spike-frequency adaptation [29, 38]. Our results extend these previous ideas and suggest that synaptic depression improves information transfer not only on fast timescales, as suggested before [6, 7, 8, 36, 37], but rather on a broad range of fast to slow timescales.

More specifically, we showed that power-law adaptation is effective in decorrelating synaptic transmission events, even when driven by input spike trains with correlations on several orders of magnitude (Fig 4). During extended periods of high activity, where spikes become highly predictable, power-law adaptation dynamically tunes down synaptic responsiveness in order to conserve energy (Fig 2C,D; Fig 4C). A similar attenuation of slow rate fluctuations is observed in a wide range of systems in the form of firing-rate adaptation [39, 40, 41]. An interesting feature of this adaptation is that it changes dynamically depending on the timescale of fluctuations in the stimulus history. Here, we found a surprisingly simple mechanism for history-dependent adaptation in the synaptic vesicle cycle: The existence of multiple stages of recovery leads to a history-dependent depletion of vesicle pools, which tunes the timescale of adaptation to the timescale of rate fluctuations in the input spike train (SI Fig S16).

Not all spike trains, however, are encoded efficiently by power-law adaptation, and we therefore wondered whether this adaptation is matched to the correlations found in hippocampus. Consider, for example, a neural code that consists of bursts of spikes separated by unpredictable intervals of silence, which only has correlations on the length of bursts. In this case, synaptic adaptation should only operate on the timescale of bursts, but not longer timescales, as this would reduce responsiveness to the unpredicted (not transmitted) events [7]. In contrast, we found that spike trains measured in the rat hippocampus (CA1) show slow rate fluctuations that follow a power-law over almost three orders of magnitude (similar to findings of previous research [16, 17], SI Fig S14). These fluctuations are whitened by encoding them via multi-stage recovery, which means that the strong correlations on long timescales are effectively removed in the encoded signal (Fig 4F). This indicates that synaptic power-law adaptation is tuned to the statistics of spike trains in hippocampus, and thus could be an important contribution to its efficient functioning.

Other functions of presynaptic dynamics Our model focused on the dynamics of vesicle recovery, although other processes are also affecting presynaptic adaptation. Most importantly, this includes the presynaptic calcium dynamics which effect synaptic facilitation and the speeding up of vesicle recovery, but possibly also other factors [3, 42]. These processes critically determine how synapses transmit signals on shorter timescales, and thus should become important when considering information transmission on a finer temporal scale than we did here. For example, naive temporal whitening is only useful for high signal-to-noise ratios, where redundancy reduction directly leads to an efficient code. However, in the face of a more noisy signal, an optimal signal encoding would require a more robust code that exploits redundancies [38, 43, 44]. At the synapse, such a robust code could for example be achieved through short-term facilitation [45]. For the future, it will be important to study how such facilitation processes interact with multi-stage vesicle dynamics.

Beyond the efficient transmission of information, synaptic adaptation has also been suggested to have other important purposes for neural computations. For example, neural adaptation could allow to transmit only certain aspects of multiplexed neural codes [46]. On the network level synaptic adaptation can take a central role in shaping dynamics, for example through the induction of synchronization [47], the alteration of attractor dynamics [48, 49, 50], or memory effects [51, 52]. Especially on the network level power-law adaptation could have a wide range of important implications that extend these previous concepts, for example by contributing to homeostatic regulation across a wide range of activity levels [53].

Conclusion and outlook   Multi-stage vesicle recovery is a plausible mechanism for long timescales of synaptic depression. Aside from efficiently transmitting signals, long timescales of synaptic adaptation could have a wide range of implications for neural function, especially on the network level. The effective description of multi-stage vesicle recovery through a power-law adaptation kernel will allow to test this in large-scale simulations.

Our results also raise the question of whether synapses could be tuned to their presynaptic firing statistics more generally. For example, hair cell synapses, which code for auditory stimuli that commonly have 1/f spectrum [54], have been implicated to employ power-law adaptation [55], and some cortical synapses are also known to possess at least two timescales of adaptation [11]. Given the variety of functional subtypes of synaptic dynamics [56], which can depend systematically on the spikes they encode [57], it would be interesting to investigate the relationship between spiking statistics and the temporal dynamics of adaptation in the corresponding synapses—not only in hippocampus but also in other neural subsystems.

Methods

Simplified response model of vesicle cycle

To obtain a simplified description of the vesicle cycle, we modeled the input-output response using a generalized linear model (GLM), similar to what has been used to model neural responses [58]. In this model, every incoming spike has the chance to trigger a release event ($r(t) = 1$), or not ($r(t) = 0$), with a probability that depends on a base release probability and a reduction based on past releases

$$p\left(r(t)|\vec{t}_{\text{past}}, \theta\right) = \frac{e^{r(t)a(t)}}{1 + e^{a(t)}},$$

$$a(t) = b - \sum_{t' \in \vec{t}_{\text{past}}} r(t')\kappa(t - t').$$

We parameterized the kernel κ through 40 values in log-spaced bins. Maximum-likelihood estimates for parameters $\hat{\theta} = \{\hat{b}, \hat{\kappa}\}$ that describe the simulated data (vesicle releases in response to 1/f spike-trains) were then obtained via gradient ascent on the log-likelihood (SI Fig S15).

Bayesian modeling of experimental data

Parameters of the models were fitted to the experimental data using Bayesian inference with MCMC sampling. To make sampling tractable, we first switched the stochastic description of vesicle dynamics to a deterministic description. This amounts to simply replacing stochastic vesicle transitions between pools through deterministic differential equations, such that for pool P_i

$$\frac{\partial P_i}{\partial t} = r_{i-1,i} - r_{i,i+1},$$

where r_{ij} are transition rates from pools i to j as before. Similarly, vesicle release was replaced by a continuous release rate $r_{\text{primed,fused}} = p_{\text{fuse}}r_{\text{stim}}P_{\text{primed}}$, where r_{stim} is the stimulation rate given by presynaptic spiking. This model shows comparable dynamics to the stochastic model (Fig S5).

Using MCMC, we then estimated the model parameters $\theta = \{\alpha_{\text{obs}}, \sigma_{\text{obs}}\} \cup \theta_{\text{obs}}$, where α_{obs} is a factor that scales the number of released vesicles (model) to the increase in dF/F (experiment), σ_{obs} is the

width of the likelihood (see below), and $\theta_{\mathrm{obs}} = \{p_{\mathrm{fuse}}, r_{ij}, P_i^{\max} : i,j \in \mathrm{Pools}\}$ are the vesicle cycle parameters. For computation, we employed the Python package PyMC3 [59] with NUTS (No-U-Turn Sampling) [60] using multiple, independent Markov chains.

Formally, we computed the posterior probability of the parameters θ given the experimental data $D = \{D_i : i \in \mathrm{conditions}\}$: $p(\theta|D) \propto p(D|\theta)p(\theta)$. Here, the likelihood was defined as a product of Gaussian likelihoods for each experimental condition

$$p(D|\theta) = \prod_{i \in \mathrm{cond.}} \mathcal{N}(\alpha_{\mathrm{obs}} O_i(\theta_{\mathrm{obs}}); \mu = \mathbb{E}[D_i], \sigma = \sigma_{\mathrm{obs}} + \mathrm{SEM}(D_i)),$$

where $\mathrm{SEM}(D_i)$ is the error of the mean, and $O_i(\theta_{\mathrm{obs}})$ is the number of releases measured in the simulation under stimulation condition i and parameters θ_{obs}. Details about the prior distributions $p(\theta)$ can be found in Supplementary Table 1.

Creation of 1/f spike trains

To simulate a point process exhibiting a 1/f power spectrum we employed the intermittent Poisson process proposed in [61], which allowed us to tune average firing rates and slopes of the power spectrum. The intermittent Poisson process models an emitter with 'on' and 'off' phases, where it generates bursts of events with a rate $1/\tau_0$ or is silent respectively. The exponent of the spectrum can be controlled by tuning the distribution of the length of 'on' and 'off' phases, which we chose such as to obtain a 1/f spectrum (for details see [61]). To tune the firing rate we scaled all emission times with a constant factor. 'On' phase burst rates of $1/\tau_0 = 25\,\mathrm{Hz}$, $37.5\,\mathrm{Hz}$ and $50\,\mathrm{Hz}$ resulted in total rates of the process of $0.5\,\mathrm{Hz}$, $0.75\,\mathrm{Hz}$ and $1\,\mathrm{Hz}$, respectively. See also the simulation code for details.

Data of hippocampal neuron spiking

Estimating the power spectra of recorded spike trains requires long recordings under natural conditions. We here used the CRCNS hc-11 dataset [27, 62], which is composed of eight bilateral silicon-probe multi-cellular electrophysiological recordings performed in CA1 of four freely behaving male Long-Evans rats. The recordings were performed during 6–8 h of rest/sleep time in cages, and 45 min of a maze running task, which we considered representative of typical rat behavior. We found that the power spectra exhibited a slight mean rate dependence, with low rate units having a flatter spectrum (SI Fig S14). For the results in the main paper, we, therefore, excluded high-rate units ($> 5\,\mathrm{Hz}$), which likely are interneurons or multi-units, and low-rate units ($< 0.7\,\mathrm{Hz}$), for which estimated power spectra become more difficult to estimate, and which we considered less relevant for efficient coding. Nevertheless, the results for low-rate units are qualitatively the same (SI Fig S14).

Estimation of power spectra

Power spectra of point processes can be estimated directly from the spike times, without binning the data. This brings the advantage that sample frequencies ω can in principle be chosen freely and are not constrained by the bin size or the length of the data, as they would be for the Fast Fourier Transform that is commonly used to compute the spectrum. Note that, for slower frequencies, bins should still be chosen as multiples of $1/T$, where T is the data length, in order to avoid resonance effects in the estimation. However, results in the main paper (Fig 4) are in the frequency regime where these can be neglected (see SI Fig S14).

An estimate for the power-spectrum $S[\omega]$ of the data (sampled from time $t = 0$ to T) can be obtained via [63]

$$S[\omega] = \left\langle \frac{1}{T} \left\| \sum_{t_{\text{event}}} \exp\left(-i2\pi\omega t_{\text{event}}\right) \right\|^2 \right\rangle,$$

where $\langle \cdot \rangle$ is an average over n realizations. For simulation experiments (Fig 4B) we chose $n = 100$ and $T \approx 1000\,\text{h}$, computed $S[\omega]$ for ω in 100 log-spaced bins, and lastly normalized the integral of the obtained spectrum to 1. For experimental data (Fig 4E) the same procedure was applied, but we split the data into parts of equal length before averaging, in order to reduce noise in the estimation. Thus, the number of realizations was $n = (\#\text{sorted units})(\#\text{splits})$, where $\#\text{sorted units} = 49$ and $\#\text{splits} = 5$, as well as $T \approx 6\text{h}/5 \sim 10\text{h}/5$, depending on the recording.

Estimation of release rates

To illustrate the effect of power-law adaptation, we computed the dynamical release rate of the models (Fig 4C,D). To this end we convolved the release trains $r(t) = \sum_{t_{\text{event}}} \delta(t - t_{\text{event}})$ with a kernel $k(t, t')$

$$\text{rate}(t) = \int_{t'=0}^{t_{max}} r(t')k(t, t')dt' = \sum_{t_{\text{event}}} k(t, t_{\text{event}}).$$

For illustration purposes, in Figure 4C we chose a Gaussian kernel with width $\sigma = 300\,\text{s}$. To compute the rate-rate plot in Figure 4D, we used a window kernel $k(t, t') = 1/W$ if $|t - t'| < W/2$ else 0, with $W = 50\,\text{s}$. Results do not critically depend on the widths of the kernels, although relatively wide kernels are required to show the effects of fluctuations on slow timescales.

Animals

Animals were handled according to the regulations of the local authorities, the University of Göttingen, and the State of Lower Saxony (Landesamt für Verbraucherschutz, LAVES, Braunschweig, Germany). All animal experiments were approved by the local authority, the Lower Saxony State Office for Consumer Protection and Food Safety (Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit), and performed in accordance with the European Communities Council Directive (2010/63/EU).

Rat dissociated hippocampal cultures

Newborn rats (*Rattus norvegicus*) were used for the preparation of dissociated primary hippocampal cultures, following established procedures [64, 65]. Shortly, hippocampi of newborn rat pups (wild-type, Wistar) were dissected in Hank's Buffered Salt Solution (HBSS, 140 mM NaCl, 5 mM KCl, 4 mM NaHCO$_3$, 6 mM glucose, 0.4 mM KH$_2$PO$_4$, and 0.3 mM Na$_2$HPO$_4$). Then the tissues were incubated for 60 min in enzyme solution (Dulbecco's Modified Eagle Medium (DMEM, #D5671, Sigma-Aldrich, Germany), containing 50 mM EDTA, 100 mM CaCl$_2$, 0.5 mg/mL cysteine, and 2.5 U/mL papain, saturated with carbogen for 10 min). Subsequently, the dissected hippocampi were incubated for 15 min in a deactivating solution (DMEM containing 0.2 mg/mL bovine serum albumin, 0.2 mg/mL trypsin inhibitor, and 5% fetal calf serum). The cells were then triturated and seeded on circular glass coverslips with a diameter of 18 mm at a density of about 80,000 cells per coverslip. Before seeding, all coverslips

underwent treatment with nitric acid, sterilization, and coating overnight (ON) with 1 mg/mL poly-L-lysine. The neurons were allowed to adhere to the coverslips for 1 h to 4 h at 37 °C in plating medium (DMEM containing 10% horse serum, 2 mM glutamine, and 3.3 mM glucose). Subsequently, the medium was switched to Neurobasal-A medium (Life Technologies, Carlsbad, CA, USA) containing 2% B27 (Gibco, Thermo Fisher Scientific, USA) supplement, 1% GlutaMax (Gibco, Thermo Fisher Scientific, USA) and 0.2% penicillin/streptomycin mixture (Biozym Scientific, Germany). The cultures were then incubated at 37 °C, and 5% CO_2 for 13-15 days before use. Percentages represent volume/volume.

Labeling

Before labeling, the primary mouse anti-synaptotagmin1 antibody (Cat# 105 311, Synaptic Systems, Göttingen, Germany) at a dilution of 1:500 and the secondary anti-mouse nanobody, conjugated to superecliptic pHluorin (custom made, NanoTag, Göttingen, Germany) at a dilution of 1:250, were preincubated in neuronal culture medium (constituting 10% of the final volume for labeling) for 40 min at room temperature (RT). The pre-incubation was performed to ensure the formation of a stable complex between the primary antibody and the secondary nanobody. The labeling solution's volume was then increased to the final volume needed for the labeling procedure (300 µL per coverslip). Following a brief vortexing, 300 µL of labeling solution was pipetted to the wells of a new 12-well plate (Cat# 7696791, TheGeyer, Renningen, Germany). Subsequently, the coverslips were transferred to the well plate. The neuronal cultures were then incubated for 90 min at 37 °C. After incubation, the cell cultures were washed 3 times in pre-heated Tyrode's solution (containing 30 mM glucose, 5 mM KCl, 2 mM $CaCl_2$, 124 mM NaCl, 25 mM HEPES, 1 mM $MgCl_2$, pH 7.4) and returned to their initial well plate, containing their own conditioned media. After an additional brief period of incubation (15 min to 20 min), the cells were ready for imaging.

Stimulation

To block neuronal activity, 10 µM CNQX (Tocris Bioscience, Bristol, UK; Abcam, Cambridge, UK) and 50 µM AP5 (Tocris Bioscience, Bristol, UK; Abcam, Cambridge, UK) were added to the imaging solution (Tyrode's buffer). Electrical stimulation of the neuronal cultures was performed with field pulses at a frequency of 20 Hz at 20 mA. This stimulation was achieved with 385 Stimulus Isolator (both, World Precision Instruments, Sarasota, FL, USA) and A310 Accupulser Stimulator and with the help of a platinum custom-made plate field stimulator (with 8 mm distance between the plates).

Imaging

The cells were mounted on a custom-made chamber used for live imaging, containing a pre-warmed imaging solution (Tyrode's solution complemented with the aforementioned drugs). The neurons were then imaged with an inverted Nikon Ti mocroscope, equipped with a cage incubator system (OKOlab, Ottaviano, Italy), Plan Apochromat 60x 1.4NA oil objective (Nikon Corporation, Chiyoda, Tokyo, Japan), an Andor iXON 897 emCCCD Camera (Oxford Instruments, Andor), with a pixel size of 16×16 µm and Nikon D-LH Halogen 12 V 100 W Light Lamp House. A constant temperature of 37 °C was maintained throughout the imaging procedure. The illumination was 200 ms and the acquisition frequency was 1.7 frames per second (fps). The imaging protocols were conducted as follows: after an initial 10 s baseline, the cells were stimulated for 0.4, 4 or 40 s, followed by a recovery period of 1, 10 or 100 s. In each stimulation, the recovery period was followed by a 2 s test stimulus, followed by 30 s final recovery. Each stimulation was paired with each recovery period, resulting in a total of 9 stimulation/recovery paradigms. The 40 s stimulation set contains an additional longer recovery period

of 200 s.

Data Analysis

The resulting movies were analyzed as follows, using routines generated in Matlab (The Mathworks Inc., Natick MA, USA; version R2022b). The frames were first aligned, to avoid the effects of drift, and they were then summed, to obtain an overall image with a signal-to-noise ratio superior to that of the individual frames. Synapse positions and areas were determined automatically in this image, by a bandpass filtering procedure [66] followed by thresholding, using an empirically-determined threshold. The signal within each of the synapse areas was then determined, for every frame, was corrected for background signal, and was then analyzed by measuring the changes induced by stimulation (as fractional change in intensity, in comparison to the baseline).

Code Availability

Simulation code is available on github under

`github.com/Priesemann-Group/synaptic_power_law_adaptation`.

Acknowledgements

We want to thank the members of the Priesemann Lab, especially Jonas Dehning, for helpful discussions. F.A.M., L.R. S.V.G. and S.R. were funded by the German Research Foundation (DFG), SFB1286, Quantitative Synaptology. V.P. received support from the SFB1528, Cognition of Interaction. V.P. and L.R. were supported by the MBExC Excellence Cluster, Deutsche Forschungsgemeinschaft (DFG) under Germany's Excellence Strategy - EXC 2067/1- 390729940.

Declaration of interests

The authors declare no competing interests.

References

[1] Erwin Neher and Nils Brose. "Dynamically primed synaptic vesicle states: key to understand synaptic short-term plasticity". In: *Neuron* 100.6 (2018), pp. 1283–1291.

[2] Natali L Chanaday et al. "The synaptic vesicle cycle revisited: new insights into the modes and mechanisms". In: *Journal of Neuroscience* 39.42 (2019), pp. 8209–8216.

[3] AbdulRasheed A Alabi and Richard W Tsien. "Synaptic vesicle pools and dynamics". In: *Cold Spring Harbor perspectives in biology* 4.8 (2012), a013680.

[4] Jun Guo et al. "A three-pool model dissecting readily releasable pool replenishment at the calyx of held". In: *Scientific reports* 5.1 (2015), p. 9517.

[5] Charles F Stevens and John F Wesseling. "Identification of a novel process limiting the rate of synaptic vesicle cycling at hippocampal synapses". In: *Neuron* 24.4 (1999), pp. 1017–1028.

[6]   Larry F Abbott et al. "Synaptic depression and cortical gain control". In: *Science* 275.5297 (1997), pp. 221–224.

[7]   Mark S Goldman, Pedro Maldonado, and LF Abbott. "Redundancy reduction and sustained firing with stochastic depressing synapses". In: *Journal of Neuroscience* 22.2 (2002), pp. 584–591.

[8]   Hagar Lavian and Alon Korngreen. "Short-term depression shapes information transmission in a constitutively active GABAergic synapse". In: *Scientific Reports* 9.1 (2019), p. 18092.

[9]   Umasankar Kandaswamy et al. "The role of presynaptic dynamics in processing of natural spike trains in hippocampal synapses". In: *Journal of Neuroscience* 30.47 (2010), pp. 15904–15914.

[10]   Haroon Anwar et al. "Functional roles of short-term synaptic plasticity with an emphasis on inhibition". In: *Current opinion in neurobiology* 43 (2017), pp. 71–78.

[11]   Juan A Varela et al. "A quantitative description of short-term plasticity at excitatory synapses in layer 2/3 of rat primary visual cortex". In: *Journal of Neuroscience* 17.20 (1997), pp. 7926–7940.

[12]   Björn Granseth and Leon Lagnado. "The role of endocytosis in regulating the strength of hippocampal synapses". In: *The Journal of physiology* 586.24 (2008), pp. 5969–5982.

[13]   Sebastian Jähne et al. "Presynaptic activity and protein turnover are correlated at the single-synapse level". In: *Cell Reports* 34.11 (2021), p. 108841.

[14]   Erwin Neher. "Merits and limitations of vesicle pool models in view of heterogeneous populations of synaptic vesicles". In: *Neuron* 87.6 (2015), pp. 1131–1142.

[15]   Silvio O Rizzoli. "Synaptic vesicle recycling: steps and principles". In: *The EMBO journal* 33.8 (2014), pp. 788–822.

[16]   William Mau et al. "The same hippocampal CA1 population simultaneously codes temporal information over multiple timescales". In: *Current Biology* 28.10 (2018), pp. 1499–1508.

[17]   Ilya Kolb et al. "Evidence for long-timescale patterns of synaptic inputs in CA1 of awake behaving mice". In: *Journal of Neuroscience* 38.7 (2018), pp. 1821–1834.

[18]   Misha Tsodyks, Klaus Pawelzik, and Henry Markram. "Neural networks with dynamic synapses". In: *Neural computation* 10.4 (1998), pp. 821–835.

[19]   Rui P Costa, P Jesper Sjöström, and Mark CW Van Rossum. "Probabilistic inference of short-term synaptic plasticity in neocortical microcircuits". In: *Frontiers in computational neuroscience* 7 (2013), p. 75.

[20]   Daniel Sage et al. "DeconvolutionLab2: An open-source software for deconvolution microscopy". In: *Methods* 115 (2017), pp. 28–41.

[21]   Gero Miesenböck, Dino A De Angelis, and James E Rothman. "Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins". In: *Nature* 394.6689 (1998), pp. 192–195.

[22]   Shama Sograte-Idrissi et al. "Circumvention of common labelling artefacts using secondary nanobodies". In: *Nanoscale* 12.18 (2020), pp. 10226–10239.

[23]   Sven Truckenbrodt et al. "Newly produced synaptic vesicle proteins are preferentially used in synaptic transmission". In: *The EMBO journal* 37.15 (2018), e98044.

[24]   Yunfeng Hua et al. "A readily retrievable pool of synaptic vesicles". In: *Nature neuroscience* 14.7 (2011), pp. 833–839.

[25]   Sunil P Gandhi and Charles F Stevens. "Three modes of synaptic vesicular recycling revealed by single-vesicle imaging". In: *Nature* 423.6940 (2003), pp. 607–613.

[26]   Thomas Schikorski and Charles F Stevens. "Morphological correlates of functionally defined synaptic vesicle populations". In: *Nature neuroscience* 4.4 (2001), pp. 391–395.

[27] Andres D Grosmark, John Long, and György Buzsáki. "Recordings from hippocampal area CA1, PRE, during and POST novel spatial learning". In: *CRCNS. org* 10 (2016), K0862DC5.

[28] Viola Priesemann and Oren Shriki. "Can a time varying external drive give rise to apparent criticality in neural systems?" In: *PLoS computational biology* 14.5 (2018), e1006081.

[29] Christian Pozzorini et al. "Temporal whitening by power-law adaptation in neocortical neurons". In: *Nature neuroscience* 16.7 (2013), pp. 942–948.

[30] Pedro Carpena and Ana V Coronado. "On the autocorrelation function of 1/f noises". In: *Mathematics* 10.9 (2022), p. 1416.

[31] Guosong Liu and Richard W Tsien. "Properties of synaptic transmission at single hippocampal synaptic boutons". In: *Nature* 375.6530 (1995), pp. 404–408.

[32] Thomas Gabriel et al. "A new kinetic framework for synaptic vesicle trafficking tested in synapsin knock-outs". In: *Journal of Neuroscience* 31.32 (2011), pp. 11563–11577.

[33] Eric Hanse and Bengt Gustafsson. "Vesicle release probability and pre-primed pool at gluta-matergic synapses in area CA1 of the rat neonatal hippocampus". In: *The Journal of Physiology* 531.2 (2001), pp. 481–493.

[34] Julia J Harris, Renaud Jolivet, and David Attwell. "Synaptic energy use and supply". In: *Neuron* 75.5 (2012), pp. 762–777.

[35] Clare Howarth, Padraig Gleeson, and David Attwell. "Updated energy budgets for neural computation in the neocortex and cerebellum". In: *Journal of Cerebral Blood Flow & Metabolism* 32.7 (2012), pp. 1222–1232.

[36] Mehrdad Salmasi et al. "Short-term synaptic depression can increase the rate of information transfer at a release site". In: *PLoS computational biology* 15.1 (2019), e1006666.

[37] Ziv Rotman, Pan-Yue Deng, and Vitaly A Klyachko. "Short-term plasticity optimizes synaptic information transmission". In: *Journal of Neuroscience* 31.41 (2011), pp. 14800–14809.

[38] Dawei W. Dong and Joseph J. Atick. "Temporal Decorrelation: A Theory of Lagged and Nonlagged Responses in the Lateral Geniculate Nucleus". In: *Network: Computation in Neural Systems* 6.2 (1995), pp. 159–178.

[39] Adrienne L Fairhall et al. "Efficiency and ambiguity in an adaptive neural code". In: *Nature* 412.6849 (2001), pp. 787–792.

[40] Brian N Lundstrom et al. "Fractional differentiation by neocortical pyramidal neurons". In: *Nature neuroscience* 11.11 (2008), pp. 1335–1342.

[41] Barry Wark, Adrienne Fairhall, and Fred Rieke. "Timescales of inference in visual adaptation". In: *Neuron* 61.5 (2009), pp. 750–761.

[42] Robert S Zucker and Wade G Regehr. "Short-term synaptic plasticity". In: *Annual review of physiology* 64.1 (2002), pp. 355–405.

[43] Joseph J. Atick. "Could Information Theory Provide an Ecological Theory of Sensory Processing?" In: *Network: Computation in Neural Systems* 3.2 (1992), pp. 213–251.

[44] Matthew Chalk, Olivier Marre, and Gašper Tkačik. "Toward a Unified Theory of Efficient, Predictive, and Sparse Coding". In: *Proceedings of the National Academy of Sciences* 115.1 (2018), pp. 186–191.

[45] Gaurang Mahajan and Suhita Nadkarni. "Local design principles at hippocampal synapses revealed by an energy-information trade-off". In: *Eneuro* 7.5 (2020).

[46] Richard Naud and Henning Sprekeler. "Sparse bursts optimize information transmission in a multiplexed neural code". In: *Proceedings of the National Academy of Sciences* 115.27 (2018), E6329–E6338.

[47] Alex Loebel and Misha Tsodyks. "Computation by ensemble synchronization in recurrent networks with synaptic depression". In: *Journal of computational neuroscience* 13 (2002), pp. 111–124.

[48] Lawrence Christopher York and Mark CW van Rossum. "Recurrent networks with short term synaptic depression". In: *Journal of computational neuroscience* 27.3 (2009), pp. 607–620.

[49] Yuichi Katori et al. "Representational switching by dynamical reorganization of attractor structure in a network model of the prefrontal cortex". In: *PLoS Computational Biology* 7.11 (2011), e1002266.

[50] Joaquin J Torres et al. "Competition between synaptic depression and facilitation in attractor neural networks". In: *Neural Computation* 19.10 (2007), pp. 2739–2755.

[51] Dean V Buonomano and Wolfgang Maass. "State-dependent computations: spatiotemporal processing in cortical networks". In: *Nature Reviews Neuroscience* 10.2 (2009), pp. 113–125.

[52] Uma R Karmarkar and Dean V Buonomano. "Timing in the absence of clocks: encoding time in neural network states". In: *Neuron* 53.3 (2007), pp. 427–438.

[53] Johannes Zierenberg, Jens Wilting, and Viola Priesemann. "Homeostatic plasticity and external input shape neural network dynamics". In: *Physical Review X* 8.3 (2018), p. 031018.

[54] Bert De Coensel, Dick Botteldooren, and Tom De Muer. "1/f noise in rural and urban soundscapes". In: *Acta acustica united with acustica* 89.2 (2003), pp. 287–295.

[55] Ian C Bruce, Yousof Erfani, and Muhammad SA Zilany. "A phenomenological model of the synapse between the inner hair cell and auditory nerve: Implications of limited neurotransmitter release sites". In: *Hearing research* 360 (2018), pp. 40–54.

[56] John Beninger et al. "Functional subtypes of synaptic dynamics in mouse and human". In: *Cell Reports* 43.2 (2024).

[57] Stefan N Oline and R Michael Burger. "Short-term synaptic depression is topographically distributed in the cochlear nucleus of the chicken". In: *Journal of Neuroscience* 34.4 (2014), pp. 1314–1324.

[58] Jonathan W Pillow et al. "Spatio-temporal correlations and visual signalling in a complete neuronal population". In: *Nature* 454.7207 (2008), pp. 995–999.

[59] John Salvatier, Thomas V Wiecki, and Christopher Fonnesbeck. "Probabilistic programming in Python using PyMC3". In: *PeerJ Computer Science* 2 (2016), e55.

[60] Matthew D Hoffman, Andrew Gelman, et al. "The No-U-Turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo." In: *J. Mach. Learn. Res.* 15.1 (2014), pp. 1593–1623.

[61] Ferdinand Grüneis. "An intermittent poisson process generating 1/f noise with possible application to fluorescence intermittency". In: *Fluctuation and Noise Letters* 13.02 (2014), p. 1450015.

[62] Andres D Grosmark and György Buzsáki. "Diversity in neural firing dynamics supports both rigid and learned hippocampal sequences". In: *Science* 351.6280 (2016), pp. 1440–1443.

[63] Bijan Pesaran. "Spectral analysis for neural signals". In: *Short Course III* 1 (2008).

[64] Gary A Banker and W Maxwell Cowan. "Rat hippocampal neurons in dispersed cell culture". In: *Brain research* 126.3 (1977), pp. 397–425.

[65] Stefanie Kaech and Gary Banker. "Culturing hippocampal neurons". In: *Nature protocols* 1.5 (2006), pp. 2406–2415.

[66] John C Crocker and David G Grier. "Methods of digital video microscopy for colloidal studies". In: *Journal of colloid and interface science* 179.1 (1996), pp. 298–310.