Early and Delayed STAT1-Dependent Responses Drive Local Trained Immunity of Macrophages in the Spleen

Aryeh Solomon[1], Noa Bossel Ben-Moshe[1], Dotan Hoffman[1], Sebastien Trzebanski[1], Dror Yehezkel[1], Leia Vainman[1], Mihai Netea[2,3], Roi Avraham[1]

1. Department of Immunology and Regenerative Biology, Weizmann Institute of Science, 7610001 Rehovot, Israel.
2. Department of Internal Medicine and Radboud Center for Infectious Diseases, Radboud University Medical Center, Nijmegen, Netherlands.
3. Department of Immunology and Metabolism, Life and Medical Sciences Institute, University of Bonn, Bonn, Germany.

Abstract

Trained immunity (TI) is the process wherein innate immune cells gain functional memory upon exposure to specific ligands or pathogens, leading to augmented inflammatory responses and pathogen clearance upon secondary exposure. While the differentiation of hematopoietic stem cells (HSCs) and reprogramming of bone marrow (BM) progenitors are well-established mechanisms underpinning durable TI protection, remodeling of the cellular architecture within the tissue during TI remains underexplored. Here, we study the effects of peritoneal Bacillus Calmette–Guérin (BCG) administration to find TI-mediated protection in the spleen against a subsequent heterologous infection by the Gram-negative pathogen *Salmonella* Typhimurium (*S*.Tm). Utilizing single cell RNA-sequencing and flow cytometry, we discerned STAT1-regulated genes in TI-associated resident and recruited splenic myeloid populations. The temporal dynamics of TI were further elucidated, revealing both early and delayed myeloid subsets with time-dependent, cell type-specific STAT1 signatures. Using lineage tracing, we find that tissue-resident red pulp macrophages (RPM), initially depleted by BCG exposure, are restored from both tissue-trained, self-renewing macrophages and from bone marrow-derived progenitors, fostering long lasting local defense. Early inhibition of STAT1 activation, using specific JAK-STAT inhibitors, reduces both RPM loss and recruitment of trained monocytes. Our study suggests a temporal window soon after BCG vaccination, in which STAT1-dependent activation of long-lived resident cells in the tissue mediates localized protection.

Introduction

Trained immunity (TI) is defined as the capacity of innate immune cells to recall and modulate their subsequent responsiveness following prior exposure to a diverse array of stimuli[1]. Unlike the adaptive immune system, these changes are antigen-agnostic and are more reflective of sustained alterations in cellular and systemic states. TI has been demonstrated to directly reprogram the metabolic, transcriptional, and epigenetic state of monocytes and macrophages, generating heightened inflammatory capabilities in-vitro[1,2]. As the transient lifespans of these cells are weeks only, this alone is insufficient to explain the sustained protection generated by TI that can last from months to years. To address this limitation, current research has been focused on long-lived multipotent progenitor immune cells.

BCG vaccine (Bacillus Calmette-Guérin), an attenuated strain of *Mycobacterium bovis*, is a potent inducer of TI, which facilitate nonspecific protection against heterologous pathogens in mice, primates, and humans[3,4]. The effects of intravenous (i.v.) BCG vaccination, where bacilli persist within the bone marrow (BM), drive the expansion of LSK+ hematopoietic stem cells (HSCs) and multipotent progenitors (MPPs)[4]. In conjunction, MPPs undergo biased differentiation towards myelopoiesis, increasing the proliferation of myeloid cells. These effects are driven primarily by interferon activation, particularly that of interferon-gamma (IFNγ) and its downstream transcription factors, STAT1 and IRF1, in MPPs and short-term HSCs, resulting in a transcriptional program of cell cycling (Cdk1, Cdk4, Ccnf) and histone modifications. Furthermore, these newly differentiated circulating cells have enhanced microbicidal properties[5,6]. Upon infection, in order to elicit protection against invading agents, trained progenitor immune cells from the BM interact within local immune cell compartments in the tissue. Understanding these mechanisms is particularly relevant given emerging challenges in infectious disease control. The rise of antimicrobial-resistant bacteria demand the development of attenuated strains-specific vaccines[5], such as the typhoid conjugate vaccine (TCV) for the gram negative *Salmonella* Typhi[6]. The premise of BCG and trained immunity is providing global heterologous protection against a range of pathogens. However, the broader potential of TI-mediated heterologous protection requires understanding of TI-mediated protection as it translates within the tissue and the local cellular architectures is not completely understood.

Most tissues harbor long-lived resident macrophages capable of self-renewal, independent of BM-
derived precursors. These resident populations, including pulmonary alveolar macrophages, Kupffer
cells in the liver and red pulp macrophages in the spleen, originate from embryonic erythro-myeloid
progenitors[7]. Antimicrobial activity of resident populations control early stages of invasion by pathogens
but are than depleted, and the niche is repopulated either by self-renewal[8] or by progenitors derived
from the BM to maintain local functions[9]. Intriguingly, resident alveolar macrophages (AM) have been
shown to undergo in-situ training, maintaining an altered phenotype over an extended period[10–12]. One
mechanism driving this training in-situ in AMs involves CX3CR1[+] T cells, which have been implicated in
local reprogramming via IFNγ release. This process has been demonstrated to occur around three weeks
after initial BCG vaccination, after which interferon-driven responses occur across the organism,
including lung and BM, resulting in viral and mycobacterial protection. It remains unclear however, how
during TI, a bridge is formed between peripheral, resident cells and BM-derived recruitment of trained
myeloid cells.

In this study, we set out to investigate the interplay of circulating and tissue-level immune cells that
mediate protection and remodeling during TI. To induce tissue specific TI, we used a model of
intraperitoneal (i.p.) injection of BCG, which rapidly delivers the bacteria to target lymphatic organs
(*e.g.,* spleen), and measured protection against subsequent *Salmonella* Typhimurium (*S*.Tm) challenge.
Within the spleen's unique structure and cellular composition, we characterized STAT1-mediated, cell
type-specific TI signatures. We demonstrate that an initial depletion of resident red-pulp macrophages is
followed by repopulation by recruited trained monocytes and a local self-renewing population,
contributing to the maintenance of STAT1 signatures and long-lasting protection within the tissue.

Results

Intraperitoneal BCG results in heterologous *S*.Tm protection and a distinct myeloid subsets with
signatures driven by STAT1.

To establish an in-vivo BCG training model targeted to the splenic tissue, and assess the extent of cross
pathogen tissue protection conferred, we administered BCG-Pasteur[10] (5×10^6 colony forming units
(CFU)) or PBS as a control via i.p injection to 8-week-old female C57BL/6J mice. Following a two-week
training period, mice were challenged with *S*.Tm (5x10^5 CFU) through i.p inoculation (**Fig. 1A**). Mice were

sacrificed after twenty-four hours, and spleen homogenates were cultured on LB agar medium to

quantify *S*.Tm load using CFU (**Fig. 1B**). Relative to the control group, mice that received BCG exhibited

enhanced protection against *S*.Tm, with a four- to five-fold decrease in CFU, indicating TI-mediated

splenic protection.

For identification of transcriptional changes in splenic myeloid populations related to TI, we isolated

splenocytes and performed staining with CD11b, Ly6C, and F4/80 gating for myeloid mononuclear

phagocytes (MPs), including classical monocytes (CM) (CD11b⁺Ly6C⁺), and CD11b⁺ Ly6C⁻ MPs comprising

non-classical monocytes (NCM) and conventional dendritic cells (cDC), and also resident red pulp

macrophages (RPM) (CD11b⁻ F4/80⁺) (**Fig. 1C**, **Figure 1 — figure supplement 1A**). We observed an

overall expansion of the myeloid compartment due to training, with an increase in the CD11b⁺ subset,

and a pronounced reduction of resident RPM (**Fig. 1D**). This observed loss of RPM is common during

infection and inflammation and has been previously described as the resident macrophage

disappearance reaction[11].

We next determined transcriptional alterations caused by TI across the myeloid compartment. CD11b⁺

cells from trained and naïve mice were isolated by sorting and processed for single-cell RNA-sequencing

(scRNA-seq)[12]. K-nearest neighbor (K-NN) clustering differentiated six distinct cell types, designated as

CM, NCM, immature DC2 (cDC2), mature DC2 (mcDC2), neutrophils, and NK cells (**Fig. 1E**). The identity

of each cell type cluster was based on established transcriptional markers (**Fig. 1F**). Additional analysis

was performed to identify differentially expressed genes (DEG) unique to training (**Fig. 1G-I; Table S1**).

While most populations derived from control or trained mice clustered together by cell type, we

identified a subset of trained CM (CM-T) particular to BCG (**Fig. 1G**). This subset represents an enhanced

state of the STAT1-regulated IFNγ response that characterizes trained CM (*e.g., Gbp2, Ly6a, Cxcl9*, and

Irg1) (**Fig. 1J**). Notably, these STAT1 activated genes are known to drive maturation and activation of

monocytes and macrophages, enhancing their antimicrobial response[13].

Stat1 and its downstream targets (*e.g., Gbp2, Irf1,* and *Ly6a*), exhibited a pronounced elevation also in

other myeloid cells (**Figure 1 — figure supplement 1B**), aligning with an earlier report that BCG induces

BM and HSC remodeling via interferon signaling, resulting in STAT1 upregulation in progenitors[14].

However, not all populations were similarly activated in conjunction with STAT1-regulated genes, with

gene set enrichment analysis (GSEA) demonstrating differential pathway upregulation across cell types

(**Fig. 1I**). CM and mcDC2 primarily activated the IFNγ response, while complement associated genes

were enriched in mcDC2. NK cells alone showed minimal activation due to training across all detected

pathways (**Figure 1 — figure supplement 1B**).

We also observed increased SCA-1 (*Ly6a/e*) expression across the monocyte subsets and mcDC2, most

notably in NCM (**Fig. 1F**; **Figure 1— figure supplement 1B**). SCA-1 (stem cell antigen-1) is a

glycosylphosphatidylinositol-anchored cell surface protein commonly used as a marker for murine

hematopoietic stem and progenitor cells within the BM. The STAT1 signaling pathway has been shown

to be involved in the regulation of SCA-1 expression[15], suggesting a role for STAT1 in the modulation of

SCA-1 during development of TI. Recent studies have also demonstrated a link between SCA-1 and

inflammation, with its expression found to be upregulated on a specific subset of Ly6C$^+$ monocytes

during infection[16]. These SCA-1$^+$ monocytes exhibited a pro-inflammatory phenotype, characterized by

the production of inflammatory cytokines and chemokines, and were implicated in the amplification of

these responses. For Ly6C$^-$ NCM, the role of SCA-1 has not yet been studied.

In addition to the upregulation of inflammatory and antimicrobial genes, we also observed a significant

downregulation of specific genes in CM-T as a result of training, including *Ccr2*, *S100A8/9*, and *Ngp* (**Fig.**

1J). CCR2, a chemokine receptor crucial for monocyte recruitment, was found to be downregulated in

response to IFNγ. This reduction in CCR2 expression is mediated by IFNγ-induced mRNA instability,

potentially serving to retain monocytes at the site of recruitment and dampen a positive feedback

loop[17]. Similarly, BCG has been shown in other contexts to reduce the expression of S100A8 and S100A9,

two calcium-binding proteins that can heterodimerize and stimulate IFNγ production in CD4+ T-cells via

an IL-10 dependent mechanism[18]. By suppressing their activity, runaway signaling and exhaustion is

avoided. Lastly, NGP (neutrophilic granule protein), has been implicated in the regulation of

inflammation through its ability to block NF-κB signaling[19]. The downregulation of these genes suggests

that trained immunity not only enhances pro-inflammatory responses but also modulates the

expression of key regulators to maintain a balanced immune response and prevent excessive

inflammation.

Dynamics of TI-associated subsets and signatures indicates early and delayed kinetics.

To gain insight into dynamic processes underlying splenic cell type-specific TI, we conducted a two-

month experiment, sacrificing mice at days 3, 14, 30, 45, and 60 post-vaccination with BCG, and

challenging with S.Tm at 14 and 60 days post training (**Fig. 2A**). We assessed BCG growth in BM and

spleen, resistance to *S*.Tm infection, flow cytometry analysis with cell type-specific training markers (CXCL9 and SCA-1), and bulk RNA-seq. Notably, protection against *S*.Tm persisted for two months post-vaccination, albeit with waning resistance over time (**Fig. 2B**). While intravenous (i.v) BCG delivery and subsequent localization of the bacterium to the BM were previously noted as crucial for robust training[14,20], our results indicate that during i.p administration this process is dispensable, as no BCG were detected in the BM (**Fig. 2C**). Conversely, in the spleen, we isolated BCG at all time points, with CFU declining sharply by day 14, reaching the limit of detection by day 30 with minimal bacterium remaining (**Fig. 2C**).

Flow cytometry analysis revealed a dynamic process during trained immunity (**Figure 2— figure supplement 1A**). Compared to mean control values, we observed rapid recruitment of CMs in the spleen, starting already at day three after BCG, peaking at day 14 before returning to baseline (**Fig. 2D**, **Figure 2—figure supplement 2B**). We also stained cells for CXCL9 expression, as a marker gene of STAT1-mediated TI that defines the CM-T subset, and revealed that this population reaches peak levels at day 14 (**Figure 2— figure supplement 1C**). In a separate experiment to assess early kinetics of CM-Ts, we measured a significant increase of this subset already at five days post BCG (**Figure 2— figure supplement 1D**), indicating an early recruitment of these cells due to the initial response to BCG. NCMs exhibit a different TI signature and kinetic pattern. While their ratio initially declines, they return to steady-state levels by day 30 (**Fig. 2D**). NCMs express SCA-1 (*Ly6a*), another STAT1-regulated gene, which persists and remains highly elevated at all subsequent time points (**Figure 2— figure supplement 1C**). Challenge with S.Tm at day 14 triggered an expansion of CM-Ts, but this response was lost by day 60, coinciding with the diminished protection observed at this later timepoint (**Figure 2— figure supplement 1E**). In contrast, SCA-1 expression in NCMs remained upregulated (**Figure 2— figure supplement 1E**).

Notably, the expression of STAT1-mediated TI markers is not solely restricted to BM-derived myeloid cells but also occurs in tissue-resident RPM. RPM loss is evident after 14 days of training (**Fig. 2D**), together with increased expression of the STAT1-regulated CXCL9 (**Figure 2— figure supplement 1C, D**). SCA-1$^+$ RPM levels fluctuate, elevating at day 14 and increasing again at day 60. Given their pivotal role in curbing the early replication of intracellular pathogens[21], such as *S*.Tm, and their long life span, as opposed to CM and NCM[22], trained RPM could be an additional determinant in generating long-lasting, localized tissue protection.

From the splenocytes collected during the kinetics experiment, we generated and analyzed bulk RNA-
seq data. Principal component analysis (PCA) displayed a substantial shift along PC1 due to BCG training
(**Figure 2— figure supplement 1F**). Examination of all differentially expressed genes (DEGs) between
BCG and PBS across these time points revealed the greatest transcriptional changes at day 14 and 30
(**Fig. 2E; Table S2**).

To investigate the biological processes represented at these time points, GSEA was performed. At day
14, we detected enrichment for IFNγ response, IFNα response, and IL6-JAK-STAT signaling, with many
genes shared across these pathways (**Fig. 2F**). This STAT1-dependent program includes several
interferon-regulated genes upregulated at day 14, including the *Gbp* gene family, *Ifitm3*, and *Socs3* (**Fig.
2G**). GBPs are guanylate-binding proteins involved in antimicrobial activity and inflammasome
activation[23], while IFITM3 is an interferon-induced transmembrane protein that plays a role in the
antiviral response[24]. SOCS3, a suppressor of cytokine signaling, particularly via IL-6/STAT3, is a necessary
component for mycobacterial infection control. By blocking IL-6 signaling, SOCS3 allows for TNF and IL-
12 signaling to occur, initiating CD4-dependent IFNγ release[25].

Within the IFNγ response pathway, we noted that associated genes display distinct temporal expression
patterns. For example, *Irg1* upregulation is limited to day 14 (**Figure 2— figure supplement 1G**)
returning to control values for all subsequent time points. This may reflect that IRG1 induction requires
bacterial phagocytosis and activation of the TLR-2/MYD88/NFKB axis[26]. Conversely, an increase in *Cxcl9*
expression is already observed by day 3, plateauing at day 14, and decreasing by day 30, mirroring its
induction in CM-Ts in the flow cytometry analysis. Curiously, despite its loss in CM-Ts, the bulk
transcriptional expression of *Cxcl9* is still sustained after 2 weeks, though at reduced levels. This may be
attributed to its activity in other immune cells or regulatory mechanisms preventing further translation.
Importantly, the IFNγ response mirrored STAT1 activity, showing initial reduction after 2 weeks followed
by sustained upregulation for all subsequent time points (**Fig. 2H**), reinforcing their coupled role in
mediating the effects of TI. Other downstream targets of STAT1, including *Ly6a* (SCA-1), maintain high
expression 2 months post vaccination (**Figure 2— figure supplement 1G**).

Distinct from these immune signatures, day 30 was characterized by enrichment of E2F targets, G2M
checkpoint, and heme metabolism (**Fig. 2E, F**). Interestingly, RPM play a crucial role in iron homeostasis,
recycling and storing iron from senescent erythrocytes, preventing accumulation of free iron that can
lead to ROS production, oxidative damage, and pathogen utilization[27]. Given the dynamics of RPM loss
by day 14, it is tempting to speculate that their subsequent repopulation is reflected by these processes.

We then investigated whether TI signatures could be maintained through progenitor programming

alone. BCG or PBS were injected i.p into mice with either a CD45.1 or CD45.2 background, respectively

(**Figure 2— figure supplement 1H**). After two weeks, BM was harvested from both, HSCs mixed 1:1, and

injected into irradiated mice for BM transplant. Six weeks post transfer, mice were sacrificed and

myeloid cells in the spleen were assessed (**Figure 2— figure supplement 1I**). Within each mouse, we

found in the spleen a greater fraction of NCMs expressed SCA-1 from trained donors relative to the

naïve control (**Figure 2— figure supplement 1J**). This persistence of the training signature in a naive host

environment demonstrates that SCA-1 expression in NCMs can be maintained through progenitor

programming alone, though it does not exclude potential contributions of local tissue signals during

normal BCG training. CM-Ts however, were undetectable, consistent with the transient nature of this

subset observed in our kinetics data.

RPM niche is replenished by recruited trained monocytes and by local training of tissue-resident

populations.

We observed a substantial reduction in RPM numbers upon BCG, followed by the expression of STAT1-

mediated TI markers in RPMs. We hypothesized that upon BCG, two possible scenarios for training and

replenishment of open niche are possible. The first is self-renewal by the remaining local tissue-resident

macrophages who are trained within the tissue, while the other is that BM-derived trained monocytes

differentiate and repopulate the open niche. While BM-derived macrophages may adopt signatures and

function of their local counterparts, they may also retain aspects of their origin, particularly enhanced

inflammatory capacities (**Fig. 3A**). To investigate this, we employed MS4A3^{Tdtm};CX3CR1^{GFP} reporter

mice[28] with a knock-in flox-cre system to selectively label BM-derived monocytes with TdTomato

fluorescence. As MS4A3 is distinctly expressed in granulocyte-monocyte progenitors (GMP), only this

lineage will be TdTomato positive. These mice were administered BCG or PBS-i.p following the training

protocol and sacrificed two weeks later (**Fig. 3B**). Effective labeling was determined by flow cytometry,

measuring the fraction of MP expressing TdTomato and/or CX3CR1 (**Figure 3— figure supplement 1A**).

As expected, the monocyte population were primarily double positive (**Figure 3— figure supplement**

1B). Intriguingly, training increased the percentage of labeled cells across monocytes, suggesting a

potential lineage bias towards granulocyte-monocyte progenitors (GMPs). This observation is consistent

with prior studies demonstrating that various microbial components can induce short-term

differentiation biases in monocytes derived from GMPs or MDPs (monocyte-dendritic progenitors),

endowing them with neutrophil- or dendritic cell-like properties[29].

For the majority of RPM, we expected minimal TdTomato expression, representative of self-
replenishment during homeostasis. However, already within the control we observed that ~16% of the
population were labeled (**Fig. 3C**). When this population was depleted during training, an even greater
fraction was positively labeled. Accordingly, even under steady-state conditions, BM-derived cells
contribute to the resident niche, but when sufficiently diminished post vaccination, active
replenishment from the BM does occur.

To determine whether Tdtm⁺ BM-derived RPM acquired a distinct transcriptional profile due to training
compared to the native Tdtm⁻ population, we sorted Tdtm⁺ RPMs, Tdtm⁻ RPMs, and Tdtm⁺ CM and NCM,
from both trained and naïve conditions. These sorted cell populations were then subjected to bulk RNA-
seq and subsequent analysis. Crucially, training associated genes were differentially expressed across all
sorted cell types (**Fig. 3D**; **Figure 3— figure supplement 1C; Table S3**). When evaluating specific DEGs
upregulated across trained RPM, we identified five gene clusters that varied between CM, NCM, and
RPM, as well as genes primarily upregulated in Tdtm⁺ or Tdtm⁻ RPM (**Fig. 3E**). Cluster I is composed of
MHCII-associated genes (*H2-Aa*, *H2-Eb1*, CD72, etc.), and is enriched in NCM and RPM cells, indicating
enhanced antigen presentation that may facilitate the activation of humoral immunity. Cluster II consists
of the STAT-1 regulated TI hallmark genes, including *Cxcl9*, *Gbp2* and *Stat1* and is upregulated in all
trained subsets. Genes in Cluster III, such as *Aif1*, *Fpr1*, and *Hk3*, are primarily observed in trained RPM
and are involved in response to tissue damage/disruption and immune infiltration. AIF1 is an established
marker of macrophage activation and is functionally involved in phagocytosis and membrane ruffling[30].
Similarly, FPR1, a formyl peptide receptor, induces chemotaxis, phagocytic uptake, and reactive oxygen
species (ROS) production[31]. Cluster IV is upregulated in BCG Tdtm⁻ RPM, though the role of most genes
detected in this cluster remains unknown in regard to their tissue-specific function and activation.
Interestingly, *CD8A* expression, typically relegated to lymphocytes, was observed in this cluster. In
monocytes, CD8A can co-engage with FcR, resulting in TNF release[32]. Lastly, cluster V, which is primarily
activated in CM regardless of training, was also upregulated in RPM due to BCG. Three of the genes
within the cluster, *Anxa1*, *Vim*, and *Wfdc17*, have all been shown to dampen excess inflammatory
responses through various mechanisms, including suppressing oxidative stress and promoting local
resolution[33–35].

GSEA performed across all upregulated genes revealed that the IFNγ response was enriched, followed by
IFNα response, allograft rejection, and JAK-STAT signaling (**Fig. 3F**). Although the IFNγ signature was
most prominent in CM, it was also observed in NCM and in the Tdtm⁺ RPM. This finding suggests that

engrafted monocytes differentiating within the niche may maintain a more inflammatory phenotype

and a heightened sensitivity to IFNγ activation. However, the BM-derived RPM are not the sole

population responsive to training, as the local fraction also upregulates the same genes in clusters I-III.

Notably, RPM, as a whole, demonstrate a greater capacity to upregulate the expression of many

interferon genes, including *Cxcl9* and *Stat1* (**Figure 3— figure supplement 1D**). Taken together, our

results indicate that BCG can reprogram populations and generate training via two separate routes.

First, the recruitment of trained progenitors and monocytes within the spleen, which differentiate

within a vacant niche, retaining their trained identity. Second, activation directly within the spleen in the

context of native tissue-resident RPM, generating tissue-specific protection.

Transient IFNγ-STAT1 inhibition prevents TI signatures and splenic infection resistance.

We observed that STAT1 signaling holds a critical role in training, with its regulated gene expression

elevated across the myeloid population. However, BCG is an intact attenuated bacterium that can

activate numerous PRRs. To evaluate whether STAT1 is necessary for TI signatures and protection, we

vaccinated STAT1-KO mice[36] with either PBS or BCG-i.p (**Figure 4— figure supplement 1A**). After a two-

week period, we assessed the myeloid population and STAT1 regulated genes. We observed a complete

absence of expression for both CXCL9 and SCA-1 across myeloid populations (**Figure 4— figure**

supplement 1B). This aligns with previous findings that demonstrated compromised acquisition of

trained immunity in IFNγR$^{-/-}$ mice[14]. Interestingly, RPM, typically depleted following BCG inoculation,

remain preserved in STAT1-KO mice (**Figure 4— figure supplement 1C**), suggesting that STAT1-mediated

pathways are involved in triggering the cellular death processes that occur during BCG interaction

and/or engulfment.

There is however a significant limitation in this mouse model, as STAT1$^{-/-}$ leave mice highly susceptible to

infection due to a severe compromise of immune homeostasis, limiting our ability to assess TI-mediated

protection upon *S*.Tm challenge. In light of our findings that STAT1 signaling is activated shortly after

BCG administration, we sought to transiently restrict STAT1 activity at these early time points, also

enabling us to investigate the effect of inhibition of STAT1 signaling on training and protection, without

affecting STAT1 activation during a secondary *S*.Tm challenge. To accomplish this, we used Fedratinib, a

specific inhibitor of JAK2 activation of STAT1 through IFNγ signaling[34], and Deucravacitinib, a specific

inhibitor of TYK2 activation of STAT1/STAT2 through IFNα/β signaling[37]. First, we administered DMSO

(control), Deucravacitinib, and Fedratinib i.p followed by BCG or PBS-i.p four hours after. These
inhibitors were then injected daily via i.p for the following four days, followed by a subsequent nine-day
rest period (**Figure 4— figure supplement 1D**). At the two-week mark, splenocytes were extracted from
all mice and analyzed by bulk RNA-seq. Only Fedratinib, not Deucravacitinib, resulted in inhibition of
STAT1-mediated TI signatures (**Figure 4— figure supplement 1E; Table S4**).

We then repeated this experiment, focusing on training phenotypes in the spleen and BM. At the two-
week mark, splenocytes were extracted from all mice for flow cytometry analysis, with CXCL9 utilized as
a marker for STAT1-mediated TI signature in CM-Ts. As observed in the transcriptional response, only
Fedratinib ablated the CXCL9$^+$ CM-Ts (**Figure 4— figure supplement 1F**). We also sought to determine if
these effects were localized solely to the tissue, or if they extended to progenitors in the BM, which
expand upon BCG exposure[14]. To ascertain this, we isolated BM from the femur, measuring the percent
of LSK$^+$ HSCs (**Figure 4— figure supplement 1G**). Here too, only Fedratinib resulted in suppressing their
expansion to levels comparable to the control (**Figure 4— figure supplement 1H**). Conversely,
perturbation of IFNα/β signaling with Deucravacitinib lead to no observable changes on trained subsets,
suggesting that it is not involved in our BCG-i.p model. Type-I interferon has been established as a
training signaling pathway in other contexts, as observed with β-glucan[38], *Candida Albicans*[39] and LPS[40].

In order to prove that early STAT1 inhibition is sufficient to block the TI protective phenotype, and not
just downstream markers, we repeated the inhibitor regime with control or trained mice receiving
either Fedratinib or DMSO, with or without *S*.Tm infection after 2 weeks (**Fig. 4A**). We then extracted
spleens to measure splenic expansion, splenoctye population levels and marker expression with flow
cytometry, bulk splenocyte transcription, and *S*.Tm susceptibility. Importantly, while we observed no
differences in CFU between control mice with or without Fedratinib, trained mice receiving Fedratinib
were significantly more susceptible to infection (**Fig. 4B**). Treatment with Fedratinib resulted in
diminished recruitment and splenocyte expansion, causing an appreciable reduction in spleen size
comparable to the control (**Figure 4— figure supplement 1I**). Accordingly, the balance of monocytes
ratios, particularly NCM, was shifted to levels similar to those observed in the DMSO control (**Fig. 4C**). In
conjunction, the frequency of CXCL9$^+$ CM-Ts was diminished, reflecting a reduction in the subset, while
in RPMs, CXCL9 showed decreased expression (**Fig. 4D**). Finally, RPM, which typically undergo depletion
after training, exhibited significantly enhanced survival, similar to the observations in STAT1-KO mice
(**Fig. 4C**).

To probe the effects of Fedratinib inhibition beyond myeloid expansion and marker acquisition, we

conducted bulk RNA sequencing on total splenocytes isolated from all experimental conditions. PCA of

the resulting data revealed two major axes of divergence among the populations: PC1, associated with

training, and PC2, associated with the response to *S*.Tm (**Figure 4— figure supplement 1J**). While the

PBS-i.p mice treated with either DMSO or Fedratinib were grouped together, the BCG-i.p samples

treated with Fedratinib clustered distinctly, shifting closer to the PBS control group. Analysis of

downregulated DEGs identified the IFNγ response as the most significantly affected pathway due to

JAK2 inhibition by Fedratinib. This included its downstream effector *Stat1*, and other key STAT1-

regulated trained immunity genes, such as *Cxcl9/10*, *Irf1*, *Gbp2*, and *Irg1*, across all treated mice (**Fig. 4E,**

F; Table S5). To further validate our findings and ensure that the loss of the trained immunity signature

was not solely a result of blocking other JAK2-associated pathways, we repeated the inhibitor

experiment using recombinant α-IFNγ. Mice were given either α-IFNγ or an isotype control after BCG or

PBS vaccination, with injections on days 0, 2, and 4, and assayed two weeks post-vaccination. Upon

sorting and sequencing CM and RPM from these treated mice, we found that the STAT1 regulated TI

signatures were completely ablated with early inhibition (**Figure 4— figure supplement 1K-L**). Although

we measured no changes in viable BCG in cultured spleen homogenates from control and treated mice

at the two-week interval (**Figure 4— figure supplement 1M**), early α-IFNγ was sufficient in blocking

training signatures. Interestingly, previous studies show that while removal of BCG by long term

antibiotic treatment decreases expansion of progenitor cells, protection against secondary ex-vivo

challenges remains[14].

Thus, the impact of Fedratinib and α-IFNγ treatment on the TI phenotype further underscores the

pivotal role of IFNγ and the JAK2-STAT1 axis in orchestrating the early programs and signatures required

for long-term local and recruited myeloid populations within the tissue. When compromised,

remodeling is suppressed and so too protection. Crucially, even though its administration was early and

removed after only five days administration, inhibition occurred. This indicates that the acquisition of TI

in the tissue occurs within a critically narrow temporal window.

Discussion

TI, or innate immune memory, embodies the capacity of innate immune cells to remember past

encounters and modify their subsequent responses, triggered by diverse conditions and stimuli. In-vivo

these processes are sustained in progenitor cells within the BM to provide long term TI, with

differentiated cells inheriting this memory[14,41]. In this study, we set out to demonstrate how long-term

protection events in the BM affords training within the tissue through local interactions with resident

populations. We demonstrated how the immune effects of BCG vaccination are differentially imparted

across recruited trained monocytes from the BM and diverse MP populations within the spleen, the

temporal dynamics governing these processes, and the mechanisms necessary for them to be sustained

long-term.

We verified that BCG-i.p inoculation stimulates an expansion of myeloid MP, notably a subset of classical

monocytes marked by CXCL9 expression (CM-T) within the spleen. This CM-T population, which peaks at

day 14 before declining, exemplifies the broader training response we observed, as our dissection of

single-cell training signatures uncovered a fundamental program dominated by IFNγ response and

STAT1 regulated genes across diverse populations. Among these, GBP2, CXCL9, and IRG1 represent

some of the pathways involved in the protective response: GBP2, a member of the GTPase family, drives

macrophages toward an inflammatory state[42] while combating bacterial infection through vacuole

disruption and pyroptosis induction[43]. CXCL9 orchestrates immune cell recruitment and activation at

infection sites[44], supporting trained immunity by establishing a pro-inflammatory environment

conducive to pathogen clearance[45]. Meanwhile, IRG1 produces the metabolite itaconate, which serves

dual functions in both directly inhibiting pathogen growth and modulating immune responses to prevent

excessive inflammation[46,47]. However, IRG1's role in training appears context-specific, as exposure to

training stimuli like β-glucan can actually block its activity[48].

By monitoring the kinetics of training over an extended duration, we observed that the initial surge in

CM recruitment is confined to the first two weeks of training, potentially signaling inflammatory

recruitment triggered by BCG exposure and RPM loss. Concurrently, the presence of CM-Ts starts to

diminish at a similar rate and is eventually lost, suggesting that these cells may be differentiated to

trained MPs in the tissue.

The depletion of RPMs at 14 days post-vaccination sets in motion a series of complex cellular responses.

Notably, we observed an enrichment of heme metabolism at day 30, which may indicate an ongoing

process of RPM replenishment. This process typically involves two key mechanisms: the local expansion

and repopulation by native populations, and the recruitment of erythroid and myeloid progenitors,

including classical and non-classical monocytes, to the organ[49]. These recruited cells, particularly

monocytes, can differentiate within the niche, transitioning through a pre-RPM state via heme-mediated

pathways[50]. Interestingly, heme metabolism has also been implicated in the control and pathogenicity of

mycobacterial species, particularly *Mycobacterium tuberculosis* (MtB). During systemic infection, MtB

activates type I interferon (IFN-I) signaling, which, along with its own virulence factors, disrupts iron

transport and uptake[20].

The presence of TI markers not only in BM-derived myeloid cells such as CM and NCM, but also in RPM,

prompted us to uncover how these cells acquired this phenotype. Utilizing Ms4a3^{TdTm} labeled mice, we

investigated whether trained RPM were being derived from cells originating in the BM. Unlike CM and

NCM, which are predominantly TdTm$^+$, RPM express this lineage tracing marker at steady state around

16%, which was further increased with training. Probing the transcriptional profile of these cells also

revealed differences due to origin, with Tdtm$^+$ RPM demonstrating a heightened response to IFNγ

stimulation and elevated expression of STAT1. However, both populations exhibit significant

upregulation of TI-associated genes post-vaccination. These findings indicate that TI in RPM is not solely

due to BM-derived precursors filling a depleted niche but suggests a capacity for both resident and BM-

derived RPM to undergo training within the tissue itself. Considering the importance of these cells in

restricting early infection events[21], their capacity for training reveals an additional factor contributing to

the TI phenotype. While our examination was limited to a singular tissue-resident macrophage subset,

other splenic macrophages like marginal zone macrophages (MZMs) and marginal metallophilic

macrophages (MMMs), known to have immunological roles, may similarly exhibit a capacity for being

trained. This research can be further broadened to include other resident myeloid populations, to

explore how BCG can bestow localized protection independent of central HSC reprogramming.

When investigating the longevity of the trained transcriptional response within the spleen, we noted

that *Stat-1* and other interferon response genes remained upregulated even two months after

vaccination. We found one such gene SCA-1 (*Ly6A*), was continuously expressed in NCM due to training,

and was still identifiable after a BM transplant. Although SCA-1 is traditionally a marker for upstream

progenitors, recent studies link its expression to inflammation, with upregulation noted in B/T cells

during *Mycobacterium Tuberculosis* (Mtb) infection[51]. Further exploration is warranted to investigate the

mechanisms that enable such enduring expression, particularly in this cell type. Also, while *Ly6a* (SCA-1)

is unique to mice, a recent study identified a human equivalent termed *Ly6s*, primarily expressed in

splenic NCM and regulated by interferon signaling, that was linked to an inflammatory cell phenotype

with resistance to viral infections[52].

Prior publications have demonstrated the importance of BCG localization in generating local vs. systemic

changes[14,53], and that BCG's initial presence within the BM is associated with HSC reprogramming and

long-term TI. Despite this, while we were able to isolate BCG from splenic tissue, not BM, from all

sampled time points, we could still achieve training and protection, including LSK+ subset expansion in

the BM. This leads us to hypothesize that the signaling initiated due to BCG can act in trans, affecting

system wide changes, and is not solely acquired due to local pathogen interactions in the BM.

Supporting this, recent findings suggest that training in alveolar macrophages can also occur through

subcutaneous BCG administration, potentially acting via the gut-lung axis[54].

Finally, given STAT1's ubiquity among all trained cells, we hypothesized that this transcription factor and

its upstream activation via interferon signaling were instrumental in driving BCG-induced trained

immunity. We observed that early transient inhibition, specifically targeting IFNγ-mediated STAT1

activation, effectively negated the hallmarks of trained immunity. These include myeloid recruitment,

disappearance of RPM, the expression of training markers, LSK+ expansion, transcriptional alterations,

and, crucially, heterologous *S*. Tm protection - highlighting STAT1's particular role in orchestrating a

protective response against this pathogen. Furthermore, while it is generally known that BCG can reside

within the tissue for weeks after inoculation, our findings suggest that even brief STAT1 and IFNγ

inhibition is sufficient to disrupt the development of trained immunity, regardless of the pathogen's

presence. It is plausible that during the initial stages of BCG exposure and inflammation, IFNγ-secreting

cells such as T/NKTs initiate the immunological remodeling needed for training. Should these processes

be obstructed at this critical juncture, the opportunity for subsequent training is lost. While our inhibitor

protocol was applied for a period of five days, the minimal duration required for effective inhibition and

comparable outcomes may be even shorter.

In summary, our study emphasizes that examining the tissue, specifically the spleen, as a tissue for

probing TI offers valuable insights into the temporal dynamics and signaling cascades that instigate and

sustain TI locally, in parallel with established systemic effects. Central to these processes is the IFNγ-

STAT1 pathway, which we identified as a key driver in establishing TI, by replenishment of resident naïve

MPs with trained recruited and local immune populations. We further delineated that during

intraperitoneal vaccination the resultant immune interactions limit BCG dissemination, while still

effectively eliciting training. Our findings open new avenues to harness STAT1 pathway induction for

optimized training and the importance of vaccines that can induce robust cross-domain protection.

Methods

Experimental Methods

Mice and bacteria strains

C57BL/6J mice, 7-9 weeks old, were purchased from ENVIGO, housed at the Weizmann Institute pathogen-free facility, and provided with standard food and water ad libitum. The Nr4a1 super-enhancer sub-domain E2-KO (E2$^{-/-}$ or ^{C57BL}/6-$Rr39^{em1Ched}$/J) mice were purchased from The Jackson Laboratory (#030204)[55]. The mice strains below were kindly provided by the following investigators:

- STAT1-KO mice by Prof. Dr. Mathias Müller[36].
- Ms4a3Cre^{TdTomato}-CX3CR1^{GFP} and CD45.1 mice by Prof. Steffen Jung.

All experiments were performed in accordance with the guidelines outlined by the Weizmann Institute Committee on Animal Care.

For in-vivo training, BCG-Pasteur, generously donated by Dr. Daniel Barkan, was utilized. The *Salmonella enterica* serovar Typhimurium strain SL1344 was used exclusively for all infection challenge experiments.

Mice training and infection

BCG were grown in Middlebrook 7H9 media (BD) supplemented with Middlebrook OADC (BD) at 37°C for 1 week to stationary phase. Bacterial aliquots of 1mL were dispensed to 2mL cryotubes (Simport) and frozen at -80C for long term storage. Prior to inoculation, tubes were thawed, centrifuged (10,000g, 2 min, RT), with pellet resuspended in 1 mL phosphate-buffered saline (PBS) (Sartorius). Bacterial concentration was calculated based on optical density at 600nm (OD$_{600}$) assuming a concentration of 5×10^8 CFU/OD, with BCG diluted to 25×10^6 CFU/mL in PBS. Mice were injected intraperitoneally (i.p) with 200µl containing 5×10^6 CFU or PBS (as controls). At given time points, mice were euthanized by CO$_2$, spleens and/or BM from the femur were harvested, and CFU numbers were evaluated by plating serial 10-100-fold dilutions of homogenized spleens or BM suspension on selective 7H9-Middlebrook agar plates.

For the initial challenge as observed in Fig.1A, cultures of S.*Tm* were grown in Luria-Bertani (LB) medium (BD) at 37°C for 16 hours to stationary phase. For all subsequent experiments, S.*Tm* were grown at 37°C for 16 hours to stationary phase in SPI-2 inducing media[56]: MgMES media (170 mM 2-(N-morpholino) ethanesulfonic acid (MES) at pH 5.0, 5 mM KCl, 7.5 mM (NH4)2SO4, 0.5 mM K2SO4, 1 mM KH2PO4, 8

mM MgCl2, 38 mM glycerol, and 0.1% casamino acids. Cultures were centrifuged (10,000g, 2 min, RT),

with pellet resuspended in PBS. Bacteria were diluted 10-fold in PBS and concentration calculated based

on optical density at 600nm (OD_{600}). Assuming a concentration of $1x10^9$ CFU/OD, S.*Tm* was diluted to

$2.5x10^6$ CFU/mL in PBS. Mice were injected intraperitoneally (i.p) with 200 µl containing $1x10^5$ CFU of

bacteria or PBS (as controls). Injected bacterial load was verified by CFU. 24 hours post infection, mice

were euthanized by CO_2, spleens were harvested, and CFU numbers were evaluated by plating serial 10-

100-fold dilutions of homogenized spleens on streptomycin LB agar plates.

BCG CFU

Spleens or BM were homogenized and serially diluted in PBS + 0.1% Triton on 7H9 Middlebrook media +

OADC with Zeocin and Kanamycin. Plates were incubated for three weeks in a humidified 5% CO^2

incubator, with CFU determined using an automated colony counter.

In-vivo interferon inhibition

JAK-STAT inhibitors Fedratinib (cat #202893), and Deucravacitinib (cat #555349) (MedKoo Biosciences)

were resuspended in DMSO, aliquoted, and stored at -80C for later use. For injection, a mixture of PEG-

300 (Sigma):Tween-80 (Sigma) was prepared at a ratio of 18:1 and filtered using a 0.22 µm filter. For

each injection, 10µl of DMSO with or without the inhibitor was added to 105 µl of the PEG:Tween mix,

followed by 180µl of PBS for a total of 30µl and injected i.p. Inhibitor concentrations are 1mg and 0.5mg

per mouse for Fedratinib and Deucravacitinib respectively. Four hours post inhibitor, BCG or PBS-i.p was

injected for training. Daily repeat injections of the inhibitors were repeated for four additional days.

For the antibody inhibition experiment, 1mg of monoclonal αIFNγ antibody (clone: XMG1.2) or isotype

control (clone: Rat IgG1) was injected i.p. four hours pre BCG vaccination (day 0), then every other day

(day 2 and 4).

Splenocytes and BM isolation and flow cytometry preparation

Spleens and BM from femurs were extracted and stored in cooled FACS Buffer (PBS, 10 mM EDTA, 2%

FBS) until further extraction.

For BM extraction, femurs were cut at both ends and placed in 0.5mL microfuge tube with a small hole

in the bottom cut out using an 18G needle. This tube was then placed in a 1.5mL microfuge tube and

centrifuged (3 min, 500g, 4°C). The pellet was resuspended in red blood cell (RBC) lysis buffer for 4

minutes at room temperature, centrifuged (3 min, 500g, 4°C) and re-suspended with FACS buffer. For

CFU determination, suspension is used directly for serial dilution. For further flow cytometry processing,

FACS buffer containing CD16/CD32 blocking antibodies (BioLegend) is added for a 15-minute incubation

on ice. All subsequent processing is identical to splenocytes.

The spleens were dissected, mashed against a 70μm cell strainer (Falcon) and washed with 5mL of cold

FACS buffer. 1mL of splenocytes were aliquoted to microfuge tubes and centrifuged twice (3 min, 500g,

4C). Pellets were re-suspended with RBC lysis buffer (Sigma), incubated for 4 minutes at room

temperature, centrifuged (3 min, 500g, 4C) and re-suspended with FACS buffer containing CD16/CD32

blocking antibodies (BioLegend) for 15 minutes on ice. Cells were centrifuged once more, and pellets

were transferred to wells of a 96-well low attachment plate for multi-sample preparation. Subsequently,

fluorophore-conjugated antibodies cocktails (listed below) in Brilliant Stain Buffer (BD) were used to

resuspend pellets, followed by a 30-minute incubation on ice. Cells were washed, re-suspended with

500-1000μl FACS buffer and passed through a 35μm cell strainer (Falcon). For absolute quantification of

cell populations, 10-50μl of Precision Count beads (BioLegend) were added to the samples.

Antibodies used in this study for splenocyte and BM staining

Epitope	Conjugation	Clone	Company
CD16/CD32	NA	93	BioLegend
NK1.1	APC	PK136	BioLegend
CD19	APC	6D5	BioLegend
CD3	APC	17A2	BioLegend
Ly6G	APC	1A8	BioLegend
Ly6C	Alexa Fluor® 700™	HK1.4	BioLegend
Ly6C	Brilliant Violet 605™	HK1.4	BioLegend
CD11b	APC/Cy7	M1/70	BioLegend
CD11c	FITC	N418	BioLegend
CXCL9	PE	MIG-2F5.5	BioLegend
F4/80	PE	BM8	BioLegend
F4/80	Brilliant Violet 421™	BM8	BioLegend
I-A/I-E (MHC-II)	Brilliant Violet 605™	M5/114/15.2	BioLegend

Ly-6A/E (Sca-1)	PE/Cyanine7	E13-161.7	BioLegend
CD11c	PerCP/Cyanine5.5	N418	BioLegend
Ter-119	FITC	TER-119	BioLegend
CD4	FITC	GK1.5	BioLegend
CD8a	FITC	53-6.7	BioLegend
Gr-1	FITC	RB6-8C5	BioLegend
CD45R/B220	FITC	RA3-6B2	BioLegend
CD11b	FITC	M1/70	BioLegend
CD117/c-Kit	APC	2B8	BioLegend
Ly-6A/E (Sca-1)	PE-Vio 770™	REA422	Miltenyi

Flow cytometry and sorting for RNA sequencing

Flow cytometry and sorting was performed using the BD FACSAria III (BD). Single cells were sorted into 384-well plates (Eppendorf) containing 2μl of a solution containing barcoded poly-T primers for reverse transcription (Sigma, Israel) according to the MARS-seq v2.0 protocol[57]. For bulk cell capture, 5-10x10^3 cells from each population were sorted into tubes containing 300μl RLT buffer (Qiagen) with β-mercaptoethanol (BME). Immediately after sorting, plates or tubes were spun down, flash-frozen in a mixture of dry ice and ethanol and stored in -80C until processing.

Single-cell RNA-seq library preparation

Single-cell libraries were prepared as described[57]. Briefly, mRNA from cells was converted to cDNA alongside barcoding and UMI addition. The cDNA of each plate was pooled followed by second DNA strand synthesis and T7 in vitro transcription. Amplified RNA was fragmented, followed by ligation of partial P5 Illumina sequence, and converted to cDNA. Full sequence of barcoded P5 and P7 of P5 were added by PCR for a sequence ready library. Final libraries were quantified for peak size and concentration using the Agilent TapeStation and Qubit HS DNA Assay kit (Invitrogen), respectively.

Bulk RNA-seq library preparation

RNA was extracted and cleaned using the RNeasy mini kit (Qiagen) with DNaseI digestion. Libraries were then prepared according to an in-house MARs-seq or CEL-seq protocol optimized for bulk RNA samples.

543   Final libraries were quantified for peak size and concentration using the TapeStation 4200 (Agilent) and

544   Qubit HS DNA Assay kit (Invitrogen), respectively.

Library Sequencing

546   Bulk and single cell libraries were diluted to a concentration of 1.8pM and run on the NextSeq platform

547   (Illumina) according to Illumina guidelines, with 75 reads for read1, and 15 reads for read2. A mean of

548   6M reads per library for the kinetics data; a mean of 12M reads per library for IFNγ inhibitor data; and a

549   mean of 3M reads per library for the MS4a3^{TdTm} bulk sorted population data.

CD45.1/CD45.2 adoptive transfer

C57BL/6J mice, expressing CD45.1 or CD45.2, were trained according to our standard protocol using BCG

or PBS, respectively. After two weeks, BM was isolated from the femur from both mice, resuspended in

PBS, and mixed in a 1:1 ratio. Recipient mice (WT C57/BL6J) were irradiated with a single dose of 950

cGy using an XRAD 320 machine (Precision X-Ray [PXI]) and reconstituted the next day via retro-orbital

injection of 5x10^6 mixed donor BM cells/mouse in 200μl PBS. Mice were given 6 weeks to allow for

reconstitution and repopulation of the hematopoietic system.

Flow cytometry, CFU, and spleen size analysis and quantification

558   Flow cytometry data was analyzed using the FlowJo software.

559   For size quantification, spleens were imaged against a contrasting background, and two-dimensional

560   area was calculated using the ImageJ software.

561   All graphs quantifying the results from flow cytometry and CFU results were performed using R on

562   RStudio with the Tidyverse package[58].

Bioinformatics Analysis

scRNA-seq data analysis

Data preprocessing

MARS-seq pipeline[57] was used for demultiplexing, alignment to the genome (mm9), and gene counting

by unique molecular identifier (UMI). Overall, we sequenced 1536 cells (768 from control mouse and

768 from trained mouse), with 1474 median UMI count per cell and 668 median genes per cell.

Data normalization and gene filtration

Only genes with at least one UMI count detected in at least one cell were used. Data was normalized to

a library size factor. Factors were calculated by dividing total UMI counts in each cell to the median of

the total UMI counts across all cells. Data was transformed to log10 scale (log10(UMI count+1)). Cells

with less than 200 UMIs were excluded due to low coverage (24 cells, 9 from naïve mouse and 15 from

trained mouse). We filtered out cell cycle and ribosomal genes and selected the top 425 most variable

genes for further analysis. Variable genes were selected based on fitting of the data to a simple noise

model based on the genes mean expression and dispersion (coefficient of variance).

Data clustering and annotations

Principal component analysis (PCA) was performed on the variable genes, and the first 40 PCs were used

for downstream analysis for k-nearest neighbor (KNN)-graph, based on Euclidian distance in PC space.

Clustering was performed using Louvain community detection on the KNN-graph (k=20). Overall, we

obtained 7 clusters. Cluster identity was inferred using cluster-specific and manually selected genes

based on cell classification literature.

Bulk RNA-seq data processing and normalization

MARS-seq pipeline was used for samples demultiplexing, alignment to the genome (mm9), and gene

counting. Data was normalized to a library size factor. Factors were calculated by dividing the total

number of reads from each sample to the median total number of reads across all samples. These

procedures were done for each dataset alone.

Kinetics data

Data was transformed to log2 scale, and minimal expression threshold was set to 3. Replicate samples of

each condition were averaged, except for 1 sample that was excluded due to low coverage (<100k

reads; BCG 30d replicate 3). Preceding PCA analysis genes were centered and normalized to a mean of 0

and a standard deviation of 1. To identify genes that were up-regulated due to training we calculated

the differences between the integrals of each gene in BCG relative to PBS along time. The differences

across all genes were approximately normally distributed, with a mean of 0.2 and a standard deviation

of 10.11. Genes with more than 3 standard deviations above the mean were defined as up-regulated

due to training.

Bulk inhibitor data

Data was transformed to log2 scale, and minimal expression threshold was set to 3. Two Fedratinib samples were excluded from analysis due to technical issues during injection that resulted in a lack of inhibition. Heatmap was generated using DEGs calculated by ANOVA (5% FDR and a minimal 2-fold; 167 genes).

Fedratinib inhibitor data

Data was transformed to log2 scale, and minimal expression threshold was set to 4. One sample was excluded due to low coverage (<100k reads; PBS +S.*Tm* + Fedratinib replicate 3). Preceding PCA analysis genes were centered and normalized to a mean of 0 and a standard deviation of 1. PCA analysis was performed on DEGs calculated using two-sided t-tests between all relevant conditions: control vs. trained samples, trained with or without inhibitor, control with or without inhibitor, uninfected vs. infected, infected with or without training, infected vs. infected with training with inhibitor, and infected with training vs. infected with training with inhibitor (5% FDR; 453 genes). Heatmap was generated using two-sample t-test between control and BCG-trained samples (1% FDR).

αIFNγ inhibitor data

Data was transformed to log2 scale, and minimal expression threshold was set to 3. Two sample was excluded due to low coverage (<250k reads CM BCG + Isotype replicate 1 and RPM BCG + Isotype replicate 4). Heatmap genes were selected from cluster I and II from the lineage tracing experiment representing shared interferon/STAT-1 upregulated genes.

MS4A3^Tdtm bulk population data

Data was transformed to log2 scale, and minimal expression threshold was set to 3. One sample was excluded due to low coverage (<150k reads; control NCM-Tdtm⁺ replicate 4). Preceding PCA analysis genes were centered and normalized to a mean of 0 and a standard deviation of 1. DEGs between control and trained mice were calculated using two-sided t-test between all control samples versus all trained samples (from all sorted populations together; 10% FDR, 382 genes).

Acknowledgments

This research was supported by the Israel Science Foundation (ISF grant No. 1289/22), the Minerva Foundation with funding from the Federal Ministry for Education and Research, the Dr. Barry Sherman Institute for Medicinal Chemistry, and the Shimon and Golde Picker Weizmann Annual Grant.

Author contributions

AS and RA designed the study. AS, DH, ST and LV performed the experiments. DY and NBBM analyzed the data. AS and RA wrote the manuscript.

Competing Interests

The authors declare no competing interests

Data and materials availability

All RNA-seq data have been deposited in NCBI's Gene Expression Omnibus (GEO) under the super-series accession number GSE252014.

References

1. Arts, R. J. W. *et al.* Long-term in vitro and in vivo effects of γ-irradiated BCG on innate and adaptive immunity. *Journal of Leukocyte Biology* **98**, 995–1001 (2015).
2. Saeed, S. *et al.* Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science* **345**, 1251086 (2014).
3. Darrah, P. A. *et al.* Prevention of tuberculosis in macaques after intravenous BCG immunization. *Nature* **577**, 95–102 (2020).
4. Moorlag, S. J. C. F. M., Arts, R. J. W., van Crevel, R. & Netea, M. G. Non-specific effects of BCG vaccine on viral infections. *Clinical Microbiology and Infection* **25**, 1473–1478 (2019).
5. Hasso-Agopsowicz, M. *et al.* Identifying WHO global priority endemic pathogens for vaccine research and development (R&D) using multi-criteria decision analysis (MCDA): an objective of the Immunization Agenda 2030. *eBioMedicine* **0**, (2024).
6. Qadri, F. *et al.* 5-year vaccine protection following a single dose of Vi-tetanus toxoid conjugate vaccine in Bangladeshi children (TyVOID): a cluster randomised trial. *The Lancet* **404**, 1419–1429 (2024).
7. Kurotaki, D., Uede, T. & Tamura, T. Functions and development of red pulp macrophages. *Microbiology and Immunology* **59**, 55–62 (2015).
8. Hoeffel, G. *et al.* C-Myb(+) erythro-myeloid progenitor-derived fetal monocytes give rise to adult tissue-resident macrophages. *Immunity* **42**, 665–678 (2015).
9. Lai, S. M. *et al.* Organ-Specific Fate, Recruitment, and Refilling Dynamics of Tissue-Resident Macrophages during Blood-Stage Malaria. *Cell Reports* **25**, 3099-3109.e3 (2018).
10. Zhang, L. *et al.* Variable Virulence and Efficacy of BCG Vaccine Strains in Mice and Correlation With Genome Polymorphisms. *Molecular Therapy* **24**, 398–405 (2016).
11. Review of the macrophage disappearance reaction - Barth - 1995 - Journal of Leukocyte Biology - Wiley Online Library. https://jlb.onlinelibrary.wiley.com/doi/abs/10.1002/jlb.57.3.361.
12. Keren-Shaul, H. *et al.* MARS-seq2.0: an experimental and analytical pipeline for indexed sorting combined with single-cell RNA sequencing. *Nat Protoc* **14**, 1841–1862 (2019).
13. Hu, X. & Ivashkiv, L. B. Cross-regulation of Signaling and Immune Responses by IFN-γ and STAT1. *Immunity* **31**, 539–550 (2009).
14. Kaufmann, E. *et al.* BCG Educates Hematopoietic Stem Cells to Generate Protective Innate Immunity against Tuberculosis. *Cell* **172**, 176-190.e19 (2018).
15. Yang, S.-H. *et al.* IFN-γ–STAT1–iNOS Induces Myeloid Progenitors to Acquire Immunosuppressive Activity. *Frontiers in Immunology* **8**, (2017).
16. Biram, A. *et al.* Bacterial infection disrupts established germinal center reactions through monocyte recruitment and impaired metabolic adaptation. *Immunity* **55**, 442-458.e8 (2022).
17. Penton-Rol, G. *et al.* Selective Inhibition of Expression of the Chemokine Receptor CCR2 in Human Monocytes by IFN-γ1. *The Journal of Immunology* **160**, 3869–3873 (1998).
18. Wang, Y. *et al.* Bacillus Calmette–Guérin–induced interleukin-10 inhibits S100A8/A9 production and hinders development of T helper type 1 memory in mice. *European Journal of Immunology* **53**, 2250204 (2023).
19. Liu, K. *et al.* Neutrophilic granule protein (NGP) attenuates lipopolysaccharide-induced inflammatory responses and enhances phagocytosis of bacteria by macrophages. *Cytokine* **128**, 155001 (2020).
20. Khan, N. *et al.* M. tuberculosis Reprograms Hematopoietic Stem Cells to Limit Myelopoiesis and Impair Trained Immunity. *Cell* **183**, 752-770.e22 (2020).

21. Hoffman, D. *et al.* A non-classical monocyte-derived macrophage subset provides a splenic
replication niche for intracellular Salmonella. *Immunity* **54**, 2712-2723.e6 (2021).
22. Yona, S. *et al.* Fate mapping reveals origins and dynamics of monocytes and tissue macrophages
under homeostasis. *Immunity* **38**, 79–91 (2013).
23. Tretina, K., Park, E.-S., Maminska, A. & MacMicking, J. D. Interferon-induced guanylate-binding
proteins: Guardians of host defense in health and disease. *J Exp Med* **216**, 482–500 (2019).
24. Clement, M. *et al.* IFITM3 restricts virus-induced inflammatory cytokine production by limiting
Nogo-B mediated TLR responses. *Nat Commun* **13**, 5294 (2022).
25. Carow, B. *et al.* Critical and independent role for SOCS3 in either myeloid or T cells in resistance to
Mycobacterium tuberculosis. *PLoS Pathog* **9**, e1003442 (2013).
26. Bomfim, C. C. B. *et al.* Mycobacterium tuberculosis Induces Irg1 in Murine Macrophages by a
Pathway Involving Both TLR-2 and STING/IFNAR Signaling and Requiring Bacterial Phagocytosis.
Front Cell Infect Microbiol **12**, 862582 (2022).
27. Vogt, A.-C. S. *et al.* On Iron Metabolism and Its Regulation. *International Journal of Molecular*
Sciences **22**, 4591 (2021).
28. Liu, Z. *et al.* Fate Mapping via Ms4a3-Expression History Traces Monocyte-Derived Cells. *Cell* **178**,
1509-1525.e19 (2019).
29. Yáñez, A. *et al.* Granulocyte-Monocyte Progenitors and Monocyte-Dendritic Cell Progenitors
Independently Produce Functionally Distinct Monocytes. *Immunity* **47**, 890-902.e4 (2017).
30. De Leon-Oliva, D. *et al.* AIF1: Function and Connection with Inflammatory Diseases. *Biology (Basel)*
12, 694 (2023).
31. Prevete, N., Liotti, F., Marone, G., Melillo, R. M. & de Paulis, A. Formyl peptide receptors at the
interface of inflammation, angiogenesis and tumor growth. *Pharmacol Res* **102**, 184–191 (2015).
32. Gibbings, D. J., Marcet-Palacios, M., Sekar, Y., Ng, M. C. Y. & Befus, A. D. CD8 alpha is expressed by
human monocytes and enhances Fc gamma R-dependent responses. *BMC Immunol* **8**, 12 (2007).
33. Sugimoto, M. A., Vago, J. P., Teixeira, M. M. & Sousa, L. P. Annexin A1 and the Resolution of
Inflammation: Modulation of Neutrophil Recruitment, Apoptosis, and Clearance. *J Immunol Res*
2016, 8239258 (2016).
34. Håversen, L. *et al.* Vimentin deficiency in macrophages induces increased oxidative stress and
vascular inflammation but attenuates atherosclerosis in mice. *Sci Rep* **8**, 16973 (2018).
35. Karlstetter, M. *et al.* The Novel Activated Microglia/Macrophage WAP Domain Protein, AMWAP,
Acts as a Counter-Regulator of Proinflammatory Response. *The Journal of Immunology* **185**, 3379–
3390 (2010).
36. Kernbauer, E. *et al.* Conditional Stat1 Ablation Reveals the Importance of Interferon Signaling for
Immunity to Listeria monocytogenes Infection. *PLoS Pathog* **8**, e1002763 (2012).
37. Lé, A. M., Puig, L. & Torres, T. Deucravacitinib for the Treatment of Psoriatic Disease. *Am J Clin*
Dermatol **23**, 813–822 (2022).
38. Kalafati, L. *et al.* Innate Immune Training of Granulopoiesis Promotes Anti-tumor Activity. *Cell* **183**,
771-785.e12 (2020).
39. Huijser, E. *et al.* Trained Immunity in Primary Sjögren's Syndrome: Linking Type I Interferons to a
Pro-Atherogenic Phenotype. *Frontiers in Immunology* **13**, (2022).
40. Zahalka, S. *et al.* Trained immunity of alveolar macrophages requires metabolic rewiring and type 1
interferon signaling. *Mucosal Immunol* **15**, 896–907 (2022).
41. Moorlag, S. J. C. F. M. *et al.* β-Glucan Induces Protective Trained Immunity against Mycobacterium
tuberculosis Infection: A Key Role for IL-1. *Cell Reports* **31**, 107634 (2020).
42. Li, X. *et al.* GBP2 promotes M1 macrophage polarization by activating the notch1 signaling pathway
in diabetic nephropathy. *Front. Immunol.* **14**, (2023).

43. Meunier, E. *et al.* Caspase-11 activation requires lysis of pathogen-containing vacuoles by IFN-
induced GTPases. *Nature* **509**, 366–370 (2014).
44. Groom, J. R. & Luster, A. D. CXCR3 ligands: redundant, collaborative and antagonistic functions.
Immunol Cell Biol **89**, 207–215 (2011).
45. Joosten, S. A. *et al.* Mycobacterial growth inhibition is associated with trained innate immunity. *The*
Journal of Clinical Investigation **128**, 1837 (2018).
46. Lampropoulou, V. *et al.* Itaconate Links Inhibition of Succinate Dehydrogenase with Macrophage
Metabolic Remodeling and Regulation of Inflammation. *Cell Metab* **24**, 158–166 (2016).
47. Michelucci, A. *et al.* Immune-responsive gene 1 protein links metabolism to immunity by catalyzing
itaconic acid production. *Proc Natl Acad Sci U S A* **110**, 7820–7825 (2013).
48. Domínguez-Andrés, J. *et al.* The Itaconate Pathway Is a Central Regulatory Node Linking Innate
Immune Tolerance and Trained Immunity. *Cell Metabolism* **29**, 211-220.e5 (2019).
49. Liao, C., Prabhu, K. S. & Paulson, R. F. Monocyte-derived macrophages expand the murine stress
erythropoietic niche during the recovery from anemia. *Blood* **132**, 2580–2593 (2018).
50. Haldar, M. *et al.* Heme-mediated SPI-C induction promotes monocyte differentiation into iron-
recycling macrophages. *Cell* **156**, 1223–1234 (2014).
51. Akter, S. *et al.* Mycobacterium tuberculosis infection drives a type I IFN signature in lung
lymphocytes. *Cell Reports* **39**, 110983 (2022).
52. Shmerling, M. *et al.* LY6S, a New IFN-Inducible Human Member of the Ly6a Subfamily Expressed by
Spleen Cells and Associated with Inflammation and Viral Resistance. *ImmunoHorizons* **6**, 253–272
(2022).
53. Vierboom, M. P. M. *et al.* Stronger induction of trained immunity by mucosal BCG or MTBVAC
vaccination compared to standard intradermal vaccination. *Cell Rep Med* **2**, 100185 (2021).
54. Jeyanathan, M. *et al.* Parenteral BCG vaccine induces lung-resident memory macrophages and
trained immunity via the gut–lung axis. *Nat Immunol* **23**, 1687–1702 (2022).
55. Thomas, G. D. *et al.* Deleting an Nr4a1 Super-Enhancer Subdomain Ablates Ly6Clow Monocytes
while Preserving Macrophage Gene Function. *Immunity* **45**, 975–987 (2016).
56. Stapels, D. A. C. *et al.* Salmonella persisters undermine host immune defenses during antibiotic
treatment. *Science* **362**, 1156–1160 (2018).
57. MARS-seq2.0: an experimental and analytical pipeline for indexed sorting combined with single-cell
RNA sequencing | Nature Protocols. https://www.nature.com/articles/s41596-019-0164-4.
58. Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).
762
763

Figure legends

Figure 1. Intraperitoneal BCG results in heterologous *S*.Tm protection and a distinct myeloid subsets with signatures driven by STAT1. (A) Mouse model of BCG vaccination and S.Tm challenge. (**B**) Splenic *S.Tm* CFU 24 hours post infection between control (n=11) and trained mice (n=11). (**C-D**) Flow cytometry plots of myeloid populations two weeks post vaccination (C) and mean percentage fold change of BCG over control for each given gated population percent from the Lin⁻ population (D). (**E**) K-nearest neighbors (KNN) plot for total CD11b⁺ single cells sorted from control and BCG mice. Color is based on conditions, or cluster identity. (**F**) Cell markers and training induced genes for each subset. Size and color intensity indicates percentage of cells within a given cluster expressing the gene and average expression. (**G**) Proportions of monocyte subsets based on classifications in E. (**H-I**) Number of DEGs in each cell subset (H), and their corresponding gene set enrichment analysis (I). (**J**) Volcano plot of DEGs in CM subset. Data in bar graphs are presented as mean±SEM, with each individual point in **B** a biological repeat. Two-tailed *t*-test used for data in **B** and **D** (*P<0.05, **P<0.01).

Figure 2. Dynamics of TI-associated subsets and signatures indicates early and delayed kinetics. (A) Experimental setup tracking TI kinetics over a two-month interval, including S.Tm challenge at days 14 and 60. (**B**) Splenic *S.Tm* CFU at 24 hours post infection for control and BCG mice at 14- and 60- days after vaccination (n=5-6). (**C**) BCG CFU from spleen and BM of BCG vaccinated mice (n=4) across time points. Red-dotted line indicates limit of detection. (**D**) Contribution of MP populations across time points from control (PBS) and BCG mice (days post injection). Percentage of CM, NCM, and dendritic cells calculated from flow cytometry analysis of CD11b⁺ population. Percentage of RPM calculated from Lin⁻ population (control: n=3, BCG: n=4 in each time point). PBS values are the mean of all time points. (**E**) Heatmap of upregulated genes due to training and relative gene expression ordered according to peak expression time. (**F**) Gene set enrichment analysis of DEGs in days 14 and 30. (**G-H**) Heatmap of IFNγ response genes (G) and their average expression dynamics compared to STAT1 expression (H). Data in bar and line graphs are presented as mean±SEM. For bar graph **B** and **C**, each individual point is a biological repeat. For line graph **H**, significance represents comparison between day 60 control and BCG. Heatmap rows in **E** and **G** indicate biological replicates. Two-tailed *t*-test used for data in **B**, **C**, and **H** (*P<0.05, **P<0.01, ***P<0.005, ****P<0.001).

Figure 3. RPM niche is replenished by recruited trained monocytes and by local training of tissue-resident populations. (**A**) Scheme representing known myeloid differentiation pathways and potential trans-differentiation of trained CM to RPM. (**B**) Mouse model to track contribution of TI-associated signatures in local and recruited MP populations with lineage tracing. (**C-D**) Flow cytometry analysis of TdTm⁺ or Tdtm⁻ Ly6C⁺ MPs and RPM and quantification of RPM TdTm⁺ subset (n=3). (**D-F**) Number of DEGs of each sorted population (D), heatmap of normalized log2 expression from TI-associated DEGs specific to trained RPM populations and gene set enrichment analysis of DEGs in each sorted population (F). Data in bar graphs are presented as mean±SEM. Heatmap rows in **E** indicate biological replicates. Two-tailed *t*-test used for data in **C** (*P<0.05).

Figure 4. Transient IFNγ-STAT1 inhibition prevents TI signatures and splenic infection resistance. (A)
Mouse model of BCG vaccination with early interferon inhibition using the Fedratinib inhibitor. (**B**)
Splenic S.*Tm* CFU for control and BCG mice, with and without Fedratinib inhibitor, 24h post infection
(n=2-6).(**C**) MP populations from control (gray) and BCG (black) mice, with or without Fedratinib
inhibition. Percentage of CM and NCM cells calculated from CD11b$^+$ population. Percentage of RPM
calculated from Lin$^-$ population. (**D**) Percentage of CXCL9$^+$CM-T, NCM, and RPM populations from
control (gray) and BCG (black) mice, with or without Fedratinib inhibition (control: n=3, BCG: n=4,
control+Fedratinib: n=3, BCG+Fedratinib: n=6). (**E**) Heatmap of normalized log2 expression of DEGs
across naïve and training conditions. (**F**) Gene set enrichment analysis of DEGs from E. Data in bar graphs
are presented as mean±SEM. For bar graph **B** each individual point is a biological repeat. Heatmap rows
in E indicate biological replicates. Two-tailed *t*-test used for data in **B**, **C**, and **D** (*P<0.05, **P<0.01,
***P<0.005, ****P<0.001).

Figure 1

Figure 2

Figure 3

Figure 4