Genetic Foundations
of Inter-individual Neurophysiological Variability.

Jason da Silva Castanheira[1], Jonathan Poli[1,2], Justine Y. Hansen[1],
Bratislav Misic[1], Sylvain Baillet[1, †]

Affiliations:
[1] Montreal Neurological Institute, McGill University, Montreal QC, Canada
[2] CentraleSupélec, Université Paris-Saclay, Paris, France

[†] Corresponding author: sylvain.baillet@mcgill.ca

Abstract: Neurophysiological brain activity shapes cognitive functions and individual traits. Here, we investigated the extent to which individual neurophysiological properties are genetically determined and how these adult traits align with cortical gene expression patterns across development. Using task-free magnetoencephalography in monozygotic and dizygotic twins, as well as unrelated individuals, we found that neurophysiological traits were significantly more similar between monozygotic twins, indicating a strong genetic influence. These heritable brain dynamics were predominantly associated with genes involved in neurotransmission, expressed along a topographical gradient that mirrors major cognitive and psychological functions. Furthermore, the cortical expression patterns of genes that contribute to individual differentiation showed progressive changes throughout development. These findings underscore a persistent genetic influence on neurophysiological activity, supporting individual cognitive and behavioral variability.

Keywords

Neurophysiology, Heritability, Genetic determinants, Brain-fingerprints, Magnetoencephalography (MEG), Monozygotic and dizygotic twins, Neurotransmission, Neurodevelopment, Cognitive and affective brain functions.

Short title: Genetics of Individual Brain Activity Patterns.

Teaser: Genetic influences shape lifelong brain activity profiles, particularly in cognitive regions.

Main Text:

Several recent neuroimaging studies have shown that ongoing brain activity at rest, without performing a specific task, defines a neurophysiological profile unique to each person. Unlike hand fingerprints, these *brain-fingerprints* are associated with individual cognitive traits and are altered by pathology (*1–10*), forming a distinct personal neurophysiological profile. Whether a person's genotype is associated with their neurophysiological profile is currently unknown. Heritability studies of inter-individual variability (*11*) have reported some genetic associations with brain structures (*12–15*) and activity (*16–21*). Genetic factors determine, to some extent, inter-individual variations in broad cognitive domains such as attentional and general intellectual abilities (*22–26*). Here, we studied how both individual neurophysiological and cognitive profiles relate to the spatial organization of gene expression patterns.

We used task-free magnetoencephalographic (MEG) imaging (*27*) to derive the individual neurophysiological profiles (*2*) of monozygotic (MZ) and dizygotic (DZ) twins, along with unrelated individuals. We hypothesized that if neurophysiological brain activity is determined by genetic factors, then the neurophysiological profiles of MZ twins, who share nearly identical genomes, would be nearly identical, unlike those of DZ twins (*28*). We then identified which genes are related to the features that differentiate individuals based on their neurophysiological profiles. Additionally, we investigated whether the genetic influence on these neurophysiological traits is linked to major psychological processes that characterize individual traits and how this influence aligns with cortical gene expression patterns throughout development.

Results:

Individual Neurophysiological Profiling and Differentiation

We derived the neurophysiological profile of 89 individuals (17 pairs of monozygotic twins, 11 pairs of dizygotic twins, and 33 unrelated individuals; 22-35 years old) from three 6-minute task-free MEG recordings provided by the Human Connectome Project(*29*). We derived individual profiles from the distribution of neurophysiological signal power of MEG activity across the cortex for each recording (*2*) (see Methods).

We first assessed the accuracy of inter-individual differentiation based on the neurophysiological profiles obtained from the three recordings of each participant (Figure 1A). The accuracy of inter-individual differentiation from these neurophysiological profiles was 83.4% (95% bootstrapped CI [73.8, 90.0]) across a broad frequency spectrum of brain activity ([1-150 Hz]). The high temporal resolution of the data enabled to study how the accuracy of neurophysiological profiling varied between the prototypical frequency bands of electrophysiology. Inter-individual differentiation varied substantially across these frequency ranges, from 59.7% in the delta band (1-4 Hz; [57.5, 73.8]) to 87.4% in the high-gamma band (50-150 Hz; [80.0, 92.5]) as detailed Figure 1B.

We then evaluated the similarity between the neurophysiological profiles of siblings in a twin pair (Figure 1A), finding that those of monozygotic twins matched with 61.5% accuracy ([46.7, 76.7]). In contrast, the profiles of dizygotic twins showed a considerably lower match accuracy of only 5.2% ([0.0, 15.0]; Figure 1B). These differences in matching accuracies between monozygotic and dizygotic twins varied across the frequency bands of neurophysiological brain activity as shown in Figure 1B. The discrepancies were particularly pronounced in the alpha band (8-13 Hz), where MZ twins matched with an accuracy of 47.8% ([33.3, 66.7]) compared to only 1.2% for DZ twins ([0.0, 5.0]). In the beta band (13-30 Hz), accuracies were 52.1% for MZ twins ([33.3, 70.0]) versus 7.1% for DZ twins ([0.0, 20.0]). Refer to Supplemental Figure S2 for the distributions of self-, other-, and twin-pair similarity between neurophysiological profiles. While differentiation accuracy remains our primary metric, we provide additional analyses illustrating that MZ twins exhibit higher neurophysiological profile similarity than DZ twins, while unrelated individuals do not significantly differ from either group. These findings reinforce that differentiation accuracy is a robust measure of individual distinctiveness in neurophysiology.

To ensure the robustness of our results, we computed chance-level neurophysiological profiling and twin pair matching based on environmental and equipment noise using empty-room recordings taken before each MEG recording session (Figure 1B). We observed above-chance differentiation across all frequency bands, including high frequencies, confirming that differentiation effects were not driven by SNR biases. We removed the variance associated with physiological artifacts from the neurophysiological profiles and found that our results remained robust (Figure S1).

Heritability of Neurophysiological Traits
Although we reported above that the neurophysiological profiles of MZ twins match each other more closely than those of the general population (Figure 1B, yellow bars), the neurophysiological profile of each MZ twin remains distinguishable from their sibling such that we can still correctly differentiate individual twins from a cohort that includes their sibling (Figure 1B, purple bars).
This suggests that individual-specific features stand apart from the heritable aspects of their neurophysiological profile. However, our results show that the most differentiable features of neurophysiological profiles tend to be heritable.

To assess this, we measured the spatial alignment between the most salient features for participant differentiation (Figure S4A-B) and the heritability of neurophysiological traits (Figure 1C & S3C). While absolute heritability values provide an upper bound on genetic influence, our primary focus was on spatial patterns of heritability across frequency bands and cortical regions.

First, we leveraged intra-class correlation (ICC) statistics (*2*, *3*, *30*) to quantify which cortical parcel and frequency band contributed the most toward participant

differentiation (see Methods). ICC measures the ratio of within-participant to between-participant variance, where higher ICC values indicate that a neurophysiological trait can robustly differentiate individuals across multiple recordings. Our analysis revealed that posterior cortical activity in the theta (4-8 Hz; ICC = 0.74), alpha (8-13 Hz; ICC = 0.83), and beta (13-30 Hz; ICC = 0.74) frequency bands were the most distinctive traits for individual differentiation (Figure S4B).

Next, we quantified the heritability of these neurophysiological traits using Falconer's method. We observed mean heritability (H²) values of 0.85 for theta-band traits, 0.76 for alpha-band traits, and 0.77 for beta-band traits across the temporal, frontal, and occipital cortices (Figure 1C). The occipital visual regions showed the highest heritability, whereas the limbic network showed the lowest (Figure 1C).

Permutation tests confirmed significant cortical-wide heritability in the alpha band, as well as regionally significant heritability in ventral and medial areas for beta-band traits and frontal regions for theta-band traits (Figure S5).

Finally, we observed significant spatial correlations between ICC-derived differentiation maps and heritability maps, confirming that the most distinctive features of neurophysiological profiles tend to be highly heritable. These correlations were significant across broadband cortical signals (1-150 Hz; r = 0.28, pspin = 0.026), with particularly strong relationships in the alpha (r = 0.62, p_{spin} = 0.0009) and beta (r = 0.58, p_{spin} = 0.0009) bands (Figure S6B). These findings confirm that the most distinctive features of neurophysiological profiles tend to be heritable, further emphasizing the genetic basis for the neurophysiological characteristics that distinguish individuals.

Fig. 1. Neurophysiological Profiling.

a) The color-coded array reports the similarity between neurophysiological profiles within individuals (self-similarity, diagonal elements) and between individuals (similarity with others, off-diagonal elements). Similarity is measured using cross-correlation coefficients derived between the neurophysiological profiles of each individual across all three datasets. A participant is correctly differentiated if their neurophysiological profile remains most similar across their own recordings. Using these statistics, we also assessed how well a twin's neurophysiological profile matched their sibling's profile compared to others in the cohort.

b) The purple bar graphs report inter-individual differentiation accuracy across all individuals in the cohort. The shaded orange bar graphs indicate twin pair matching accuracy. Data are reported with 95% confidence intervals for broadband (1-150 Hz) and specific frequency bands: delta (1-4 Hz), theta (4-8 Hz), alpha (8-13 Hz), beta (13-30 Hz), gamma (30-50 Hz), and high gamma (50-150 Hz). Chance-level accuracies (grey bars) are presented to rule out the possibility that environmental conditions

around each participant's recording and head geometry biased inter-individual differentiation accuracy (see Methods for details). Chance-level estimates were computed using 'mock brain-fingerprints' derived from environmental and instrument noise recorded during MEG sessions.

c) Cortical maps depicting the heritability of neurophysiological traits, computed using Falconer's equation. Regions with high heritability values indicate areas where neurophysiological activity is most strongly influenced by genetics (see Main Text and Methods).

Legend: b-fp1: first brain-fingerprint; b-fp2: second brain-fingerprint; MZ: monozygotic twins; DZ: dizygotic twins.

Person-Specific Neurophysiological Signals are Aligned with Cortical Gene Expression

We then assessed whether neurophysiological profiles for participant differentiation (full ICC maps from Figure 1C) also align topographically with genetic cortical expressions. To do this, we studied the spatial covariation of the ICC of neurophysiological profiles with maps of cortical gene expressions with greater differential stability (>0.1) (31–35), retrieved from the microarray Allen Human Brain Atlas (33), using Partial Least Squares (PLS) correlation (Figure S7A). We found a single PLS component significantly accounting for 85.2% of the covariance (CI [73.4, 90.1], p_{spin} = 0.01; Figure S7B). This analysis revealed a topographical pattern where visual and somatomotor regions exhibited positive covariance with genetic expressions, while limbic regions exhibited negative covariance (Figure S7B). This indicates that the most differentiable traits of individual neurophysiological profiles are spatially aligned with specific gene expression patterns along the cortical surface. We cross-validated this observation with 1,000 permutations corrected for spatial autocorrelation, resulting in a median out-of-sample correlation of r = 0.64 (p_{spin} = 0.002; Figure S7C).

We then measured how each frequency range of cortical activity contributed to this alignment between cortical gene expression and the salient traits of neurophysiological profiles. To do so, we computed the loadings for each frequency band, which correspond to the Pearson correlation coefficients between ICC data and the observed cortical score pattern (see Methods). We found that all frequency ranges contributed exclusively positively to this association: delta (1-4 Hz; r = 0.52, 95% CI [0.45, 0.61]), alpha (8-13 Hz; r = 0.63, 95% CI [0.58, 0.69]), beta (13-30 Hz; r = 0.52, 95% CI [0.42, 0.62]), gamma (30-50 Hz; r = 0.71, 95% CI [0.66, 0.76]), and high gamma bands (50-150 Hz; r = 0.43, 95% CI [0.34, 0.53]) with the exception of the theta band (4-8 Hz; r = 0.07, 95% CI [-0.09, 0.22]).

Genes and Cell Types Associated with Neurophysiological Traits

We then investigated which genes' expressions contribute the most to the reported association with neurophysiological traits, aiming to identify the biological functions

associated with these genes and the specific cell types involved. We selected the top 2,208 genes based on their highest positive loadings in the PLS analysis and the top 2,344 genes based on their highest negative loadings. We performed a gene ontology (GO) analysis using the ShinyGO pipeline and resources from the GO database (*36*), which revealed distinct biological processes linked to these sets of genes (see Methods). Genes with positive loadings were enriched in biological processes such as ion transport, synaptic functioning, and neurotransmitter release, while genes with negative loadings were associated with processes like development, neurogenesis, and cell morphogenesis (Figure 2A, bottom panel; complete gene list in Supplemental Information). Additionally, we verified that the spatial alignment between gene-ontological categories was not driven by the spatial autocorrelation of cortical maps through SPIN tests (colours of points in Figure 2A; see Methods).

To determine the types of cells corresponding to these genes, we analyzed gene sets that are preferentially expressed in seven cell types as determined by RNA sequencing studies(*37–42*). We found an overrepresentation of genes with positive loadings in excitatory neurons (p_{FDR}= 0.002; 1,000 gene permutations, two-tailed, FDR: corrected for false discovery rate) and inhibitory neurons (p_{FDR} = 0.006). Conversely, genes with positive loadings were underrepresented in astrocytes (p_{FDR} = 0.002) and oligodendrocyte precursor cells (OPCs; p_{FDR} = 0.03). Genes with negative loadings were predominantly represented in astrocytes (p_{FDR} = 0.002), microglia (p_{FDR} = 0.004), and oligodendrocytes (p_{FDR} = 0.002), but were less represented in OPCs (p_{FDR} = 0.002; Figure 2B left panel).

These findings suggest a clear dichotomy: genes that show positive loadings and thus positively correlate with the distinctive traits of neurophysiological profiles are predominantly expressed in neurons, whereas genes with negative loadings, indicating a negative correlation with these traits, are more frequently expressed in neuron-supporting cells such as astrocytes and microglia. Our results are consistent with existing models of the physiological origins of MEG signals (*27*, *43*, *44*).

Association with Neuropsychological Processes

Building on prior work reporting associations between brain-fingerprint features and cognitive traits (*1–3*, *45*, *46*) as well as genetic influences on inter-individual variations in cognitive domains (*22–26*), we investigated how the gene-neurophysiological associations discovered in our data may relate to neuropsychological processes.

To investigate the relationship between neurophysiological differentiation and cognitive function, we utilized Neurosynth, a meta-analytic tool that aggregates fMRI activation data from over 15,000 studies (*47*). Neurosynth generates probabilistic cortical maps indicating the likelihood that activity in a given brain region is associated with a specific cognitive term (e.g., 'memory,' 'attention'). Importantly, these maps do not distinguish between task-related activations and deactivations, instead reflecting the overall spatial distribution of reported cognitive associations.

To assess whether cortical gene expression aligns with cognitive function, we performed a Partial Least Squares (PLS) analysis comparing cortical expression patterns with Neurosynth-derived cognitive term maps. This allowed us to test whether the genes linked to neurophysiological differentiation also show spatial correspondence with cognitive processes.

A single significant component accounted for 67.2% of this covariance (CI [54.7, 72.2%], p_{spin} = 0.002), highlighting a distinction between cognitive and emotional domains (Figure 2C): negative loadings were associated with processes related to emotions, mood, and arousal, whereas positive loadings were linked to attentional, planning, and multimodal sensory processes.

We found that these patterns of covariance between gene expression and psychological processes were strongly aligned with those linking gene expression to individual neurophysiological traits (r=0.99, p_{spin} < 0.001; Figure S8, middle and right panels; see also Supplemental Information).

Figure 2: Associations Between Neurophysiological Traits, Cortical Gene Expression, and Neuropsychological Processes.

(a) Gene-differentiation Partial Least Squares (PLS) analysis. The top panel shows neurophysiological brain score patterns for positive and negative loadings, indicating which cortical parcels align with the observed covariance pattern.

The bottom panel presents the results of the gene ontology analysis. Each point represents an enriched biological process within the corresponding gene set. The size of each point denotes the associated p-value, while color intensity reflects the p-

value after spatial autocorrelation correction (pSPIN). For clarity, related terms have been grouped together using horizontal bars.

(b) Cell-type deconvolution analysis illustrating the proportion of genes (both positive and negative) preferentially expressed in seven distinct cell types, based on prior single-cell and single-nucleus RNA sequencing studies (*37–41, 48*). The significance of these ratios was assessed via permutation testing (*p < 0.05*). Points represent observed ratios, while box plots show the distribution of permuted gene set ratios. Key: 'neuron I' refers to inhibitory neurons, 'neuron E' to excitatory neurons, 'endo' to endothelial cells, 'oligo' to oligodendrocytes, 'OPC' to oligodendrocyte precursor cells, 'micro' to microglia, and 'astro' to astrocytes.

(c) Gene-Neuropsychological Processes PLS analysis. This panel illustrates the top 10 psychological terms that most significantly contribute negatively (cyan) and positively (pink) to the latent component identified in the gene-neuropsychological processes PLS analysis. The bar graphs display the loadings of neuropsychological terms. Confidence intervals were computed via bootstrapping and are not necessarily symmetric. Terms in the top quartile of negative and positive loadings delineate a cognitive-affective gradient.

Developmental Trajectory of the Gene-Differentiation Signature

Previous work has established that genetic influences on neuropsychological processes become more pronounced with development (*24, 26, 49*). Following our identification of the cortical topography of a gene-differentiation signature that aligns with brain activations associated with neuropsychological processes, we tested whether informative neurophysiological profiles in adulthood relate to cortical gradients of gene expression throughout development. To test this hypothesis, we assessed the topographical alignment between the cortical expression of genes across life stages in 12 cortical regions (*50*) and that of the gene-differentiation signature measured in adulthood. We found that this alignment (i.e., slope) was stronger in later life stages in all tested cortical regions, except for the hippocampus (Figure 3). We also report this alignment separately for genes with positive and negative loading: while genes with positive loadings were more strongly expressed in the cortex across developmental stages, genes with negative loadings were less expressed with age (see Figure S9). These findings suggest that individual neurophysiological profiles in adulthood are most aligned to patterns of gene expression in adulthood, which become increasingly pronounced throughout development.

Figure 3. Strengthening of the Gene-Differentiation Signature Across Development.
The left panel illustrates the progressive increase in gene-differentiation scores across developmental stages (prenatal through adulthood) for 12 brain regions, based on the BrainSpan data (*50*). The solid black line with grey shading indicates the trajectory of gene scores derived from random auto-correlation preserving permutations of gene expression data. The right panel shows regional slopes of these gene-differentiation scores, with vertical lines representing empirical slopes and asterisks indicating brain regions where the gene-differentiation signature significantly strengthens with age (*p_{FDR} < 0.05). Note that this analysis is based on adult neurophysiological profiles and does not assume longitudinal changes; it reflects how these profiles align with age-related changes in gene expression patterns.

Discussion:
Patterns of brain activity can differentiate between individuals, akin to hand fingerprints (*1–3*, *51*). However, these brain-fingerprints are associated with neuropsychological traits (*1*, *2*, *45*, *46*) and pathophysiology (*6*, *7*, *30*). We investigated the genetic bases of individual neurophysiological profiles.

Heritability of Neurophysiological Traits
Our data show that some of the neurophysiological traits that shape individual profiles are heritable. The neurophysiological profiles of monozygotic twin pairs match each other beyond what can be explainable by mere neuroanatomical resemblances (Figure 1B & Figure S3). Conversely, our ability to match the neurophysiological profiles of dizygotic twins was at random chance, achieving similar accuracy to noise recordings (Figure 1B). This suggests that the similarity between the neurophysiological profiles of dizygotic twin pairs is comparable to that of two unrelated individuals. These observations underscore that neurophysiological traits are in part shaped by genetics.

The neurophysiological traits of alpha and beta-band activity in parietal cortical regions showed particularly strong heritability (Figure S6), confirming previous observations of

the heritability of the individual frequency of alpha activity in humans (*16, 17*). We observed a connection between these brain rhythms and brain transcriptome gradients. These gradients correlate with brain activations during cognitive tasks, suggesting a deeper genetic-based influence on neurophysiological variability.

The neurophysiological profiles most informative for participant differentiation followed similar spatial distributions across alpha, beta, and gamma bands (Figure S4B). These findings align with prior studies on MEG brain fingerprinting, which emphasize posterior sensorimotor regions—particularly in the beta band—as key for individual differentiation (*2, 51, 52*). Prior work has also shown that differentiation is driven by multiple MEG signal features, including power spectra, phase-based connectivity, and amplitude-based functional connectivity. Despite methodological differences, posterior cortex consistently emerges as a central region for neurophysiological individuality. Our own prior work using a large MEG dataset across the lifespan (N > 1000) similarly highlights sensorimotor regions as crucial for participant differentiation in early-to-mid adulthood (*52*).

The spatial alignment between cortical gene expression and Neurosynth-derived cognitive maps reflects an organization pattern consistent with previously reported functional gradients (*53–56*). Specifically, the observed differentiation aligns with a cognitive-affective gradient, where sensory-driven cognitive processes (e.g., attention, perception) are spatially distinct from affective and emotion-related processes. These results suggest that individual differences in neurophysiology are embedded within this large-scale functional architecture

The cortical gradients identified herein—including inter-individual neurophysiological differentiation, gene expression patterns, and cognitive functional associations—closely resemble the well-documented unimodal-to-transmodal axis of cortical organization (*53–56*). This large-scale gradient is a fundamental organizing principle of the brain, observed across multiple levels of neurobiology, including structural connectivity, functional networks, and transcriptomic variation (*54, 56*)

Our findings suggest that inter-individual variability in neurophysiological traits follows this primary axis of cortical organization. Sensory areas, which exhibit the highest neurophysiological differentiation, are functionally and molecularly distinct from transmodal association areas. These results extend prior literature by demonstrating that this organizational framework also applies to individual differences in large-scale neurophysiology. Future studies should explore the genetic and developmental factors shaping these gradients and their role in cognitive and behavioral variability.

A Gene-Differentiation Signature

Gene expression profoundly impacts brain structures and functions (*32, 48, 57–61*), such as cortical folding and connectivity within brain networks (*13, 62, 63*). Our study extends these findings by demonstrating that, in addition to anatomical and brain-network properties, genetics also shape task-free, ongoing neurophysiological brain activity. This activity reflects a set of distinctive neurophysiological traits, uniquely

defining an individual's profile in relation to the cortical expression of specific genes, forming a gene-differentiation signature.

Using gene expression atlases, we found that these traits are related to the expression of genes regulating ion transport and neurotransmission. Genetic variants and variations in gene expression levels likely influence the function of these genes, contributing to the observed inter-individual differences in neurophysiological profiles. Notable genes previously reported as major elements of neurochemical signalling include the catechol-O-methyltransferase (COMT) enzyme, crucial for dopamine metabolism(*64*); the adenosine A2A receptor gene (*65*); adenosine deaminase gene; regulators of glutamatergic receptor channels (*17*); and the GABAB receptor (GABABR1) (*66*), which are major elements of neurochemical signaling. This aligns with previous research that identified related genetic variants modulating alpha rhythms and other components of cortical neurodynamics (*16*, *18*, *66*). Our findings indicate that this gene signature is primarily active in both excitatory and inhibitory neurons (Figure 2B). Genetic influences on these cells likely manifest as observable differences in overall brain activity patterns, leading to distinct variations in macroscale brain signaling across individuals.

Conversely, our study revealed that the cortical expression of genes involved in cell morphogenesis and neurogenesis, particularly in limbic regions, exhibits a negative correlation with individual neurophysiological differentiation (Figure 2B). In contrast to positively-loaded genes, which relate to neurochemical interactions between neurons, these negatively-loaded genes are mainly expressed in supportive cells such as astrocytes and microglia. This suggests that genetic influences on neurophysiological individual traits are more related to neuronal communication than to inherited structural brain features, consistent with existing models of the physiological origins of MEG signals(*27*, *43*, *44*).

Our research highlights potential pathways for future studies on individual neurophysiology by providing a biologically grounded framework to understand behavioural variations. Animal models could be instrumental in manipulating alleles of the genes identified in our study to assess their impact on gene product functioning, large-scale brain signal characteristics, and ultimately, behaviour. This approach would provide further insights into how genetic variations influence both neurophysiological traits and behavioural differences.

Our study demonstrates that genetic factors play a significant role in defining the uniqueness of neurophysiological profiles, yet they do not account for all observed variations. While the differentiation accuracy for monozygotic twins is substantial, it does not reach the same level as that for single individuals. This discrepancy suggests an explanatory gap likely attributable to environmental influences. Environmental factors affect gene expression and behavior(*25*, *67*, *68*) and may also modulate certain neurophysiological traits. The inability to differentiate twin pairs with absolute precision

underscores the need for further research with larger twin cohorts to fully elucidate the interplay between genetics and environment in shaping individual brain activity.

Alignment of the Gene-Differentiation Signature with Neuropsychological Processes

We found that, across the cortex, the gene-differentiation signature of neurophysiological profiles aligns with maps of cortical activations related to specific cognitive and emotional processes (Figure 2C). This alignment was pronounced for cognitive functions such as attention and, unexpectedly, for affective processes such as emotion regulation, challenging the traditional separation between cognition and emotions (*69*). This suggests that both cognitive abilities and emotional regulation are linked to gene expression related to neuronal communication. As shown above, genes involved in neuronal communication exhibited the strongest cortical expression alignment with the most differentiating traits of neurophysiological profiles. This linkage between cortical mapping of cognitive abilities and emotional regulation with the gene-differentiation signature supports a more integrated model of neuropsychological processes, bridging cognitive and affective functions.

Our research also highlights the importance of limbic brain regions, including the anterior cingulate cortex (Figure 2C), which are often linked to mental health disorders (*70–73*). We found that limbic regions involved in affective processes are associated with gene expression related to cell morphogenesis, primarily expressed in astrocytes and microglia (Figure 2B & Figure S8). This relationship, indicated by negative gene scores (Figure 2B & C), underscores the potential role of neuron-support cells in mental health. Furthermore, our data suggest that the neurophysiological signals in limbic regions, particularly those related to emotional processes, may not be distinct enough to differentiate individuals effectively (Figure 2A). This insight is particularly relevant for the application of personalized medicine in mental health.

Trajectory of the Gene-Differentiation Signature Throughout Development

Prior research has shown that genetic influences on neuropsychological processes increase across development (*24*, *26*, *49*) and that individual brain-fingerprint features become more stable and unique with age (*5*, *74*). To examine whether neurophysiological differentiation aligns with gene expression across the lifespan, we assessed the spatial alignment between adult neurophysiological differentiation maps and cortical gene expression gradients in 12 cortical regions (51). We found that this alignment (i.e., slope) was strongest in later developmental stages in all regions except for the hippocampus (Figure 3).

Further, we observed a differential expression trajectory: Genes with positive loadings in the gene-differentiation signature showed stronger cortical expression in adulthood. Genes with negative loadings showed higher expression in early life and decreased with age (Figure S9).

These findings suggest that the molecular landscape supporting neurophysiological differentiation in adulthood becomes increasingly structured throughout development, but do not imply that we directly tracked individual-level gene expression changes over time. These findings provide a preliminary molecular framework for understanding age-related changes in neurophysiological variability (*75–80*). Future research should examine the causal mechanisms linking neurophysiology, gene expression, and behavioral variability across the lifespan.

In light of these results, we propose three hypotheses for further investigation: i) Individual neurophysiological profiles become increasingly distinct with age, ii) The specific neurophysiological traits most relevant for differentiation evolve across the lifespan, and iii) Developmental shifts in participant differentiation mirror changes in cortical gene expression patterns.

The interpretation of these developmental findings should be considered in light of the limitations of available gene expression data. The BrainSpan dataset provides a valuable, albeit limited, cross-sectional window into developmental gene expression trajectories. While our analysis demonstrates a spatial alignment between informative neurophysiological profiles and gene expression gradients across development, this does not imply direct longitudinal tracking of gene expression within individuals.

Our findings are supported by independent evidence demonstrating increasing genetic influences on cognition and brain activity across the lifespan (*24*, *26*, *49*). Additionally, a recent complementary study on a lifespan dataset (>1000 individuals, ages 4–89) suggests a progressive shift from sensory to transmodal regions in individual differentiation, paralleling the developmental trajectory of gene expression gradients (*52*). However, future studies leveraging highly sampled developmental cohorts with multimodal neurophysiological and transcriptomic data will be essential to confirm and refine these interpretations.

Methodological Considerations and Future Directions
Previous neuroimaging studies with fMRI have shown how gene expression modulates functional connectivity in the frontoparietal network, a key feature of inter-individual differentiation in fMRI brain-fingerprints (*81*). In contrast, our study highlights the role of posterior unimodal sensory cortical regions in driving inter-individual differentiation based on neurophysiological traits. This disparity between neuroimaging modalities underscores the distinct biological underpinnings of hemodynamic fMRI and electrophysiological signals (*1*, *2*, *51*, *82*). Specifically, while fMRI connectome-based individual profiles partially reflect the structural connections between network nodes (*83*, *84*), our findings indicate that individual neurophysiological profiles are predominantly shaped by processes of neuronal communication, particularly through the expression of genes involved in ion transport and neurotransmission. This distinction underscores the

critical role of these genes in mediating neuronal signaling and brain activity patterns, providing novel insight into the mechanisms underlying neurophysiological individuality.

We must also underscore potential limitations in the interpretation of our findings. Heritability estimates were derived using Falconer's formula, which does not account for variance due to environmental factors. Moreover, due to our limited sample size, we do not interpret heritability estimates in terms of absolute variance explained by genetics alone. Instead, we emphasize the spatial distribution of heritable neurophysiological traits and their alignment with neurophysiological differentiation patterns (Figures S4 & S6). Future studies with larger twin cohorts and comprehensive demographic documentation are warranted to refine these estimates further.". Our genomics analysis is based on rare, albeit limited, data from a small sample of post-mortem brain tissue. The tissue sampling process has inherent biases, such as a focus on the left hemisphere and sex imbalance. Future research should aim to mitigate these biases. Additionally, we acknowledge that post-mortem gene expression measures may not accurately reflect in vivo conditions (*85*).

A specific strength and weakness of our approach lies in aggregating data from multiple sources. While this integration uniquely enables addressing multiscale neuroscientific questions, it also presents inherent methodological challenges. To mitigate concerns regarding data aggregation (e.g., alignment between cortical maps from the Allen Human Brain Atlas and the Human Connectome Project), we cross-validated our PLS results on a held-out sample of cortical parcels (Figure S7C) and employed spatial autocorrelation-preserving permutations. Although our findings robustly demonstrate covariance between neurophysiological profiles and average cortical gene-expression patterns, future studies are needed to examine whether deviations from normative neurophysiological profiles predict inter-individual deviations in cortical gene expression. This will require extensive, coordinated data collection from highly sampled individuals

Although we observed a significant spatial alignment between neurophysiological differentiation (ICC) and cortical gene expression, it is important to recognize that these measures originate from different sources. Our findings do not imply that individual deviations in gene expression predict deviations in neurophysiology. Instead, they suggest that neurophysiological differentiation follows an organizational pattern that aligns with normative cortical gene expression. Testing causal relationships between gene expression and neurophysiological traits requires within-participant gene-neurophysiology datasets, which are currently rare. Future studies with highly sampled individuals or animal models will be critical to establishing causality.

The correlational nature of our findings highlights the need for further experimental validation, potentially through animal models, to establish causative links between gene expression and neurophysiological and behavioural traits. The gene-differentiation signature and its relationship to neuropsychological processes highlighted in our study were derived from meta-analytic findings (*47*). While we believe these findings are novel and biologically meaningful, they do not provide a complete picture of the neural basis

of behaviour. The inherent limitations of meta-analytic approaches and the limited scope of behaviours analyzed here will need to be expanded upon in future studies.

In conclusion, our research elucidates the relationship between molecular variations, brain activity, and individual differences. Using a multiscale, data-driven approach, the present study suggests new avenues for understanding the biological foundations of individual variability. Our findings lay the groundwork for future studies to further explore these complex interconnections, thereby enriching our understanding of the neural underpinnings of human behaviour and cognition. We hope that our work will inspire continued exploration and innovation in the field, ultimately advancing our knowledge of how genetic and neurophysiological factors shape the human experience.

Methods:

Participants: MRI and MEG data from 89 healthy young adults (22-35 years old; mean= 28.6, SD= 3.8 years; see Table S1) were collected from the Human Connectome Project (HCP)(*29*). Among these 89 participants, 34 were monozygotic twins, and 22 were dizygotic twins. The zygosity of the participants was confirmed with genotyping tests. All participants underwent three approximately six-minute resting-state eyes-open MEG recordings using a 248-magnetometer whole-head Magnes 3600 system (4DNeuroimaging, San Diego, CA). All sessions were conducted at the same location with a sampling rate of 2034.5 Hz, as detailed in HCP protocols(*29*). Data from twin pairs included in the Human Connectome Project were sex-matched and reared similarly (*29, 86*). This ensures that observed differences in neurophysiological differentiation between MZ and DZ twins are not attributable to differences in early-life environmental exposure.

Ethics: The procedures for the curation and analysis were reviewed and approved according to the institutional ethics policies of McGill University 's and the Montreal Neurological Institute's Research Ethics Boards (ref no. 22-06-079).

MEG Data Preprocessing & Source Mapping: MEG data were preprocessed following good practice guidelines(*87*) using Brainstorm(*88*). Source maps for each participant's recordings were computed using a linearly-constrained minimum-variance (LCMV) beamformer and were clustered into 200 cortical parcels of the Schaefer atlas (*89*), as detailed in Supplemental Information Methods.

Neurophysiological Profiles: Power spectrum density (PSD) estimates at each cortical parcel were derived using Welch's method (sliding window of 2 s, 50% overlap). The neurophysiological profile (or *brain-fingerprint*) of each participant consisted of PSD values defined at 301 frequency bins (range: 0-150Hz; ½ Hz resolution) for each of the 200 cortical parcels. Neurophysiological profiles were generated for each of the three MEG recordings per participant.

Individual Differentiation: Individual neurophysiological profiling was conducted following our previous work (Figure 1A)(*2*). We assessed the correlational similarity between

participants' neurophysiological profiles across recordings. For each probe participant, we computed Pearson's correlation coefficients between their neurophysiological profile from one their three recordings available and a test set consisting of the neurophysiological profiles of all participants derived from another one of the other two recordings (between-participants similarity), including the probe participant's profile (within-participant similarity). A participant was correctly differentiated if the highest correlation coefficient between their neurophysiological profile and the test set was obtained from their own neurophysiological profile from the other recording. This procedure was repeated for all participants. We then computed the percentage of correctly differentiated participants across the cohort, yielding a score of differentiation accuracy for the neurophysiological profiling approach. This procedure was repeated for all possible pairs of data recordings from the three available for each participant, and the mean differentiation accuracy was reported.

Matching Neurophysiological Profiles Between Twin Pairs: We declared that the neurophysiological profiles of twin siblings matched with one another if their Pearson's correlation coefficients were higher than with any other participant. This matching procedure was repeated for all twin pairs in the cohort, and we reported the percentage of correctly matched pairs separately for monozygotic and dizygotic twin pairs. The results of this analysis are presented in Figure 1b.

Band-limited Neurophysiological Profiles: We replicated the individual and twin pair neurophysiological profiling analyses, restricting the PSD features to those averaged over the typical electrophysiological frequency bands: delta (1–4 Hz), theta (4–8 Hz), alpha (8–13 Hz), beta (13–30 Hz), gamma (30–50 Hz), and high-gamma (50–150 Hz). For example, the alpha-band neurophysiological profile consists of power spectral density estimates at each frequency within 8–13 Hz, rather than an average across this range.

Bootstrapping of Differentiation Accuracy Scores: To derive confidence intervals for the reported differentiation accuracies, we employed a bootstrapping method. We randomly selected a subset of participants representing approximately 90% of the tested cohort (i.e., 30 MZ, 20 DZ, and 30 non-twins), derived a differentiation accuracy score, and repeated this procedure 1000 times with random subsamples of participants. We report 95% confidence intervals from the 2.5^{th} and 97.5^{th} percentiles of the resulting empirical distribution of differentiation accuracies. For deriving confidence intervals for the differentiation of twin pairs, we randomly subsampled 15 MZ twin pairs and 10 DZ twin pairs for each iteration of the random subsampling.

Saliency of Neurophysiological Traits: We calculated intraclass correlations (ICC) using a one-way random effects model to quantify the contribution of each cortical parcel and frequency band toward differentiating between individuals across the cohort (*3, 90*). This approach follows prior work introducing ICC for characterizing topological fingerprints in neuroimaging (*3, 90*). ICC quantifies the ratio of within-participant to between-participant variance. High ICC values indicate the saliency of a feature of the

neurophysiological profile (a neurophysiological *trait*) in distinguishing individuals, as it reflects high within-participant consistency and low between-participant variability. To avoid potential bias due to twin pairs, we computed ICC across all individuals in the cohort and across 100 random subsamples, ensuring only one twin from each pair was included in each subsample (i.e., for each subsample, we randomly selected either twin A or twin B to include in the calculation of ICC). The ICC values obtained from bootstrapping were nearly identical to those obtained from the entire cohort (98.6% correlation). We proceeded with the ICC values averaged across bootstraps for all analyses. The results of this analysis are presented in Figure S4b.

Chance-level participant differentiation: To rule out the possibility that environmental factors influenced participant differentiation and twin matching accuracies, we computed a measure of chance-level differentiation and matching accuracy using mock neurophysiological profiles derived from empty-room noise recordings collected during each MEG session. These recordings capture instrumental and environmental noise such as: Time of day effects (e.g., variations in background electromagnetic activity); scanner-related variability (e.g., fluctuations in baseline signal sensitivity); external noise sources (e.g., minor variations in shielding effectiveness or room conditions).

From these empty-room noise recordings, we derived a single 'mock brain-fingerprint' for each participant, based on our previous work (*2*, *8*, *52*). The noise time series was projected onto each participant's cortical source maps, and differentiation accuracy was recalculated using these mock profiles to test whether chance-level accuracy varied systematically across frequency bands. These were projected onto participant cortical source maps, and differentiation was recalculated using the same pipeline. Importantly, chance-level differentiation remained low (<15%) across all frequency bands, and the alpha band did not show disproportionate differentiation accuracy, ruling out SNR-driven bias.

Heritability of Neurophysiological Traits: We calculated the heritability of individual neurophysiological profiles, considered as phenotypes, using the Falconer formula(*11*). This method estimates the relative contribution of genetics versus environmental factors in determining a phenotype. A phenotype in this context refers to the overall neurophysiological profile of an individual, while a trait refers to specific aspects or features within this profile, such as power spectrum density in a particular frequency band or cortical region. If the similarity in a phenotype between monozygotic (MZ) twins is greater than that between dizygotic (DZ) twins, the trait is considered heritable:

$$H^2 = 2(r_{MZ} - r_{DZ}),$$

where r_{MZ} (and r_{DZ}, respectivey) is the intraclass correlation between MZ (and DZ, respectively) twin pairs for a given neurophysiological trait. It is important to note that heritability reflects the similarity within twin pairs for a given phenotype, whereas ICC reflects the stability of a trait within a person relative to others in the cohort.

Heritability estimates were computed using three recordings per twin, resulting in 9 observations per twin pair (i.e., 3 × 3 comparisons). To confirm the robustness of our results, we tested whether heritability remained stable across different sample sizes and correlation metrics. Subsampling MZ twins to match DZ sample sizes resulted in highly similar heritability estimates (r = 0.90). Additionally, using Pearson's correlation instead of ICC produced near-identical estimates (r = 0.98). The results of these analyses are presented in Figure S4.

Gene Expression Data: Gene expression data was obtained from the six postmortem brains provided by the Allen Human Brain Atlas (AHBA; http://human.brain-map.org/)(*33*)using the *abagen* python package(*35*). Our analyses followed a similar pipeline to prior studies(*32*). Gene expression was obtained by averaging across donors. We computed differential stability [Δ_S(p)] for every gene probe. Differential stability is defined as the consistency of gene expression patterns across different donor brains, quantified using Spearman rank correlation. Higher differential stability indicates that a gene exhibits spatially stable expression patterns across individuals, making it more suitable for population-level analyses. We retained 9,104 genes with a differential stability above 0.1 for all subsequent analyses. We retained 9104 genes with a differential stability above 0.1 for all future analyses, following good-practice guidelines and previous literature(*32*, *34*, *35*, *91*). See Supplemental Information Methods for further details.

PLS Derivation of a Gene-Differentiation Signature: We related features for participant differentiation to gene expression gradients using partial least squares (PLS) analysis (*92–94*). We z-scored the columns of two data arrays: one containing the full neurophysiological trait matrix (all ICC values across frequency bands) and the other containing gene expression levels. The neurophysiological traits array (denoted as *Y*) had 6 columns representing each frequency band of interest and 200 rows representing the cortical parcels of the Schaefer atlas. The gene expression array (denoted as *X*) had 9104 columns representing genes and 200 rows representing the same cortical parcels. We applied singular value decomposition (SVD) to the covariation matrix of *X* and *Y* such that:

$$\mathrm{Cov}(X,Y) = USV^T,$$

where *U* and *V* are the left and right singular vectors, and *S* is the diagonal matrix of singular values. As typical with PLS, this decomposition allowed us to identify the latent variables that maximally covary between the gene expression data and the neurophysiological traits. Each singular value indicates the amount of covariance explained by its corresponding latent component. The vectors *U* and *V* provide the weights for the genes and frequency bands, respectively, for each latent component. High weights in *U* correspond to genes that strongly covary with high weights in *V*, which correspond to specific frequency bands. Positively weighted genes covary with positively weighted frequency bands, elucidating the relationship between genetic and neurophysiological variability.

Gene expression and ICC scores were computed for each cortical parcel by projecting the original data matrices *X* and *Y* onto the singular vector weights obtained from the PLS analysis. Specifically, these scores represent the covariance between the gene expression data and the neurophysiological traits. For example, cortical parcels with positive scores indicate covariance between positively weighted genes and positively weighted frequency bands, which are important for participant differentiation.

The contribution of each frequency band to the observed latent variable (i.e., loadings) was computed using Pearson correlation between ICC values and the cortical score pattern obtained from PLS analysis. These correlations represent the loadings for each frequency band, quantifying their alignment with the gene-differentiation signature. Loadings were computed as Pearson's correlation coefficients between each variable's regional spatial distribution over the cortex (i.e., gene expression and ICC data) and the corresponding cortical score pattern (i.e., correlating gene expression with ICC scores). We used Pearson's correlation coefficients for our loadings because they provide a standardized measure of the strength and direction of the linear relationship between variables, facilitating interpretation. Variables with large absolute loadings are highly correlated with the observed score pattern and strongly relate to the latent component of covariance.

To assess the significance of the latent components, we conducted permutation tests that preserved the spatial autocorrelation of cortical maps (see below). Specifically, we performed 1,000 spin tests and computed a null distribution of singular values. P-values were computed as the proportion of null singular values that achieved a greater magnitude than the empirical singular values. Additionally, we computed bootstrapped confidence intervals for the singular values by randomly resampling the rows (corresponding to the cortical parcels) of both data matrices 1,000 times. We report the 2.5^{th} and 97.5^{th} percentiles of the resulting distribution of singular values. The results of this analysis are presented in Figure 2a.

Gene Ontology Analysis: To determine the biological processes that strongly contributed to the set of positively and negatively loaded genes, we conducted an enrichment analysis using gene ontology(*95*, *96*), a framework for categorizing gene products based on their molecular function and associated biological processes(*36*, *95*).

For negatively and positively loaded genes, we separately selected the 50% with the largest absolute loadings (e.g., genes with the 50% most negative or positive loadings) and input these genes into the *ShinyGO v.0.77* gene ontology tool(*36*) using the GO biological processes pathway databases(*96*). Genes without Enrtez IDs were excluded from the analysis. P-values associated with fold enrichment for all terms were FDR corrected. See Supplemental Data for a comprehensive list of all biological processes and their corresponding fold enrichment values. The results of this analysis are presented in Figure 2c.

Previous research has shown that standard gene ontology analyses may inflate significance estimates due to spatial autocorrelation in cortical maps (*97*). To mitigate

this issue, we employed spatial autocorrelation-preserving permutation tests to rigorously evaluate statistical significance (see Methods: **Correction for Spatial Autocorrelation of Cortical Maps**). Specifically, for each gene ontology category, we computed the spatial alignment between gene expression topographies and neurophysiological differentiation maps, then assessed significance against 1000 iterations of spatially permuted cortical maps. This approach ensures that our findings exceed what would be expected under a random spatially autocorrelated null model.

Critically, all previously reported gene ontology categories remained significant following these rigorous control analyses, confirming the robustness of our findings. These results are now illustrated in Figure 2A (see pSPIN values).

Cell-type Deconvolution: We aggregated cell-specific gene sets for seven cell types using data from five human adult postmortem single-cell and single-nucleus RNA sequencing studies(*37–41*, *48*). The seven cell classes were determined based on hierarchical clustering, resulting in the following cell types: astrocytes, endothelial cells, microglia, excitatory neurons, inhibitory neurons, oligodendrocytes, and oligodendrocyte precursor cells (OPC).

We assessed the preferential expression of cell-specific gene sets by 1) computing the ratio of positively loaded genes that overlapped with the cell-specific gene set; and 2) permuting gene sets 1,000 times to assess statistical significance.

This approach allowed us to determine the statistical significance of the overlap between the loaded genes and the cell-specific gene sets, providing insights into the cell-type-specific expression patterns of the genes contributing to the neurophysiological traits. The results of this analysis are presented in Figure 2b.

Gene Expression & Neuropsychological Processes PLS Analysis: We repeated the above-described PLS analysis to relate gene expression to neuropsychological processes, as indexed by brain activation maps obtained from *Neurosynth*(*47*). This analysis replicated the approach of Hansen and colleagues(*32*), using the Schaefer-200 atlas(*89*). For detailed methodology, see Supplemental Information Methods.

Gene-Signature Evolution Across Developmental Stages: We used brain gene expression data available from BrainSpan(*50*) ,which features gene expression levels from different developmental stages ranging from 8 post-conception weeks to 40 years of age. We computed gene scores for 12 cortical regions across neurodevelopmental stages by multiplying the gene expression matrix obtained from BrainSpan with the PLS-derived gene weights (columns of *U* above).

We fitted linear slopes—using the MATLAB *polyfit()* function—to the gene scores across neurodevelopment for each cortical parcel separately. These slopes were then compared to statistical null slope values obtained by performing spatially autocorrelation-preserving permutations, then running the PLS analysis pipeline, and multiplying the null gene weights with BrainSpan gene expression data. This resulted in

a null distribution of slopes (1,000 permutations). See Supplemental Information Methods for further details. The results of this analysis are presented in Figure 3 and S7.

Visualization: We plotted brain maps of ICC, heritability, and PLS brain scores using the *ggSchaefer* and *ggseg* R packages. All other plots were generated using the *ggplot2* package in R(*98*).

Correction for Spatial Autocorrelation of Cortical Maps: We corrected for spatial autocorrelation of cortical map data, where applicable, using SPIN tests. SPIN tests preserve the spatial autocorrelation of cortical topographies by rotating the cortex surface data, effectively permuting the spatial positions while maintaining the spatial structure. This generates a null distribution that accounts for spatial autocorrelation. We conducted 1,000 spin permutations of our brain maps using the Hungarian method(*99, 100*).

References:

1. E. S. Finn, X. Shen, D. Scheinost, M. D. Rosenberg, J. Huang, M. M. Chun, X. Papademetris, R. T. Constable, Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. Nat. Neurosci. 18, 1664–1671 (2015).
2. J. da Silva Castanheira, H. D. Orozco Perez, B. Misic, S. Baillet, Brief segments of neurophysiological activity enable individual differentiation. Nat. Commun. 12, 5713 (2021).
3. E. Amico, J. Goñi, The quest for identifiability in human functional connectomes. Sci. Rep. 8, 8254 (2018).
4. T. Kaufmann, D. Alnæs, C. L. Brandt, F. Bettella, S. Djurovic, O. A. Andreassen, L. T. Westlye, Stability of the Brain Functional Connectome Fingerprint in Individuals With Schizophrenia. JAMA Psychiatry 75, 749–751 (2018).
5. T. Kaufmann, D. Alnæs, N. T. Doan, C. L. Brandt, O. A. Andreassen, L. T. Westlye, Delayed stabilization and individualization in connectome development are related to psychiatric disorders. Nat. Neurosci. 20, 513–515 (2017).
6. P. Sorrentino, R. Rucco, A. Lardone, M. Liparoti, E. Troisi Lopez, C. Cavaliere, A. Soricelli, V. Jirsa, G. Sorrentino, E. Amico, Clinical connectome fingerprints of cognitive decline. NeuroImage 238, 118253 (2021).
7. E. Troisi Lopez, R. Minino, M. Liparoti, A. Polverino, A. Romano, R. De Micco, F. Lucidi, A. Tessitore, E. Amico, G. Sorrentino, V. Jirsa, P. Sorrentino, Fading of brain network fingerprint in Parkinson's disease predicts motor clinical impairment. Hum. Brain Mapp. 44, 1239–1250 (2023).
8. J. da Silva Castanheira, A. I. Wiesman, J. Y. Hansen, B. Misic, S. Baillet, J. Breitner, J. Poirier, P. Bellec, V. Bohbot, M. Chakravarty, The neurophysiological brain-fingerprint of Parkinson's disease. EBioMedicine 105 (2024).
9. Z. Fu, J. Liu, M. S. Salman, J. Sui, V. D. Calhoun, Functional connectivity uniqueness and variability? Linkages with cognitive and psychiatric problems in children. Nat. Ment. Health 1, 956–970 (2023).

10. D. S. Falconer, The inheritance of liability to certain diseases, estimated from the incidence among relatives. Ann. Hum. Genet. 29, 51–76 (1965).

11. F. Pizzagalli, G. Auzias, Q. Yang, S. R. Mathias, J. Faskowitz, J. D. Boyd, A. Amini, D. Rivière, K. L. McMahon, G. I. de Zubicaray, N. G. Martin, J.-F. Mangin, D. C. Glahn, J. Blangero, M. J. Wright, P. M. Thompson, P. Kochunov, N. Jahanshad, The reliability and heritability of cortical folds and their genetic correlations across hemispheres. Commun. Biol. 3, 1–12 (2020).

12. D. Posthuma, E. J. C. de Geus, E. J. C. M. Mulder, D. J. A. Smit, D. I. Boomsma, C. J. Stam, Genetic components of functional connectivity in the brain: The heritability of synchronization likelihood. Hum. Brain Mapp. 26, 191–198 (2005).

13. J. E. Schmitt, A. Raznahan, S. Liu, M. C. Neale, The Heritability of Cortical Folding: Evidence from the Human Connectome Project. Cereb. Cortex N. Y. NY 31, 702–715 (2020).

14. P. H. Lee, J. T. Baker, A. J. Holmes, N. Jahanshad, T. Ge, J.-Y. Jung, Y. Cruz, D. S. Manoach, D. P. Hibar, J. Faskowitz, K. L. McMahon, G. I. de Zubicaray, N. H. Martin, M. J. Wright, D. Öngür, R. Buckner, J. Roffman, P. M. Thompson, J. W. Smoller, Partitioning heritability analysis reveals a shared genetic basis of brain anatomy and schizophrenia. Mol. Psychiatry 21, 1680–1689 (2016).

15. C. M. Smit, M. J. Wright, N. K. Hansell, G. M. Geffen, N. G. Martin, Genetic variation of individual alpha frequency (IAF) and alpha power in a large adolescent twin sample. Int. J. Psychophysiol. Off. J. Int. Organ. Psychophysiol. 61, 235–243 (2006).

16. E. Salmela, H. Renvall, J. Kujala, O. Hakosalo, M. Illman, M. Vihla, E. Leinonen, R. Salmelin, J. Kere, Evidence for genetic regulation of the human parieto-occipital 10-Hz rhythmic activity. Eur. J. Neurosci. 44, 1963–1971 (2016).

17. B. P. Zietsch, J. L. Hansen, N. K. Hansell, G. M. Geffen, N. G. Martin, M. J. Wright, Common and specific genetic influences on EEG power bands delta, theta, alpha, and beta. Biol. Psychol. 75, 154–164 (2007).

18. R. F. Betzel, J. D. Medaglia, A. E. Kahn, J. Soffer, D. R. Schonhaut, D. S. Bassett, Structural, geometric and genetic factors predict interregional brain connectivity patterns probed by electrocorticography. Nat. Biomed. Eng. 3, 902–916 (2019).

19. J. Richiardi, A. Altmann, A.-C. Milazzo, C. Chang, M. M. Chakravarty, T. Banaschewski, G. J. Barker, A. L. W. Bokde, U. Bromberg, C. Büchel, P. Conrod, M. Fauth-Bühler, H. Flor, V. Frouin, J. Gallinat, H. Garavan, P. Gowland, A. Heinz, H. Lemaître, K. F. Mann, J.-L. Martinot, F. Nees, T. Paus, Z. Pausova, M. Rietschel, T. W. Robbins, M. N. Smolka, R. Spanagel, A. Ströhle, G. Schumann, M. Hawrylycz, J.-B. Poline, M. D. Greicius, IMAGEN CONSORTIUM, Correlated gene expression supports synchronous activity in brain networks. Science 348, 1241–1244 (2015).

20. B. D. Fulcher, J. D. Murray, V. Zerbi, X.-J. Wang, Multimodal gradients across mouse cortex. Proc. Natl. Acad. Sci. 116, 4689–4695 (2019).

21. I. Zwir, J. Arnedo, C. Del-Val, L. Pulkki-Råback, B. Konte, S. S. Yang, R. Romero-Zaliz, M. Hintsanen, K. M. Cloninger, D. Garcia, D. M. Svrakic, S. Rozsa, M. Martinez, L.-P. Lyytikäinen, I. Giegling, M. Kähönen, H. Hernandez-Cuervo, I. Seppälä, E. Raitoharju, G. A. de Erausquin, O. Raitakari, D. Rujescu,

T. T. Postolache, J. Sung, L. Keltikangas-Järvinen, T. Lehtimäki, C. R. Cloninger, Uncovering the complex genetics of human character. Mol. Psychiatry 25, 2295–2312 (2020).

22. R. A. Power, M. Pluess, Heritability estimates of the Big Five personality traits based on common genetic variants. Transl. Psychiatry 5, e604 (2015).

23. J. Mollon, E. E. M. Knowles, S. R. Mathias, R. Gur, J. M. Peralta, D. J. Weiner, E. B. Robinson, R. E. Gur, J. Blangero, L. Almasy, D. C. Glahn, Genetic influence on cognitive development between childhood and adulthood. Mol. Psychiatry 26, 656–665 (2021).

24. T. J. Bouchard, M. McGue, Genetic and environmental influences on human psychological differences. J. Neurobiol. 54, 4–45 (2003).

25. C. Haworth, M. Wright, M. Luciano, N. Martin, E. de Geus, C. van Beijsterveldt, M. Bartels, D. Posthuma, D. Boomsma, O. Davis, Y. Kovas, R. Corley, J. DeFries, J. Hewitt, R. Olson, S.-A. Rhea, S. Wadsworth, W. Iacono, M. McGue, L. Thompson, S. Hart, S. Petrill, D. Lubinski, R. Plomin, The heritability of general cognitive ability increases linearly from childhood to young adulthood. Mol. Psychiatry 15, 1112–1120 (2010).

26. S. Baillet, Magnetoencephalography for brain electrophysiology and imaging. Nat. Neurosci. 20, 327–339 (2017).

27. A. J. Mayhew, D. Meyre, Assessing the Heritability of Complex Traits in Humans: Methodological Challenges and Opportunities. Curr. Genomics 18, 332–340 (2017).

28. D. C. Van Essen, K. Ugurbil, E. Auerbach, D. Barch, T. E. J. Behrens, R. Bucholz, A. Chang, L. Chen, M. Corbetta, S. W. Curtiss, S. Della Penna, D. Feinberg, M. F. Glasser, N. Harel, A. C. Heath, L. Larson-Prior, D. Marcus, G. Michalareas, S. Moeller, R. Oostenveld, S. E. Petersen, F. Prior, B. L. Schlaggar, S. M. Smith, A. Z. Snyder, J. Xu, E. Yacoub, The Human Connectome Project: A data acquisition perspective. NeuroImage 62, 2222–2231 (2012).

29. J. da Silva Castanheira, A. I. Wiesman, J. Y. Hansen, B. Misic, S. Baillet, PREVENT-AD Research Group, Quebec Parkinson Network, Neurophysiological brain-fingerprints of motor and cognitive decline in Parkinson's disease. MedRxiv Prepr. Serv. Health Sci., 2023.02.03.23285441 (2023).

30. B. T. T. Yeo, F. M. Krienen, J. Sepulcre, M. R. Sabuncu, D. Lashkari, M. Hollinshead, J. L. Roffman, J. W. Smoller, L. Zöllei, J. R. Polimeni, B. Fischl, H. Liu, R. L. Buckner, The organization of the human cerebral cortex estimated by intrinsic functional connectivity. J. Neurophysiol. 106, 1125–1165 (2011).

31. J. B. Burt, M. Demirtaş, W. J. Eckner, N. M. Navejar, J. L. Ji, W. J. Martin, A. Bernacchia, A. Anticevic, J. D. Murray, Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. Nat. Neurosci. 21, 1251–1259 (2018).

32. J. Y. Hansen, R. D. Markello, J. W. Vogel, J. Seidlitz, D. Bzdok, B. Misic, Mapping gene transcription and neurocognition across human neocortex. Nat. Hum. Behav. 5, 1240–1250 (2021).

33. M. J. Hawrylycz, E. S. Lein, A. L. Guillozet-Bongaarts, E. H. Shen, L. Ng, J. A. Miller, L. N. van de Lagemaat, K. A. Smith, A. Ebbert, Z. L. Riley, C. Abajian, C. F. Beckmann, A. Bernard, D. Bertagnolli, A. F. Boe, P. M. Cartagena, M. M.

Chakravarty, M. Chapin, J. Chong, R. A. Dalley, B. D. Daly, C. Dang, S. Datta, N. Dee, T. A. Dolbeare, V. Faber, D. Feng, D. R. Fowler, J. Goldy, B. W. Gregor, Z. Haradon, D. R. Haynor, J. G. Hohmann, S. Horvath, R. E. Howard, A. Jeromin, J. M. Jochim, M. Kinnunen, C. Lau, E. T. Lazarz, C. Lee, T. A. Lemon, L. Li, Y. Li, J. A. Morris, C. C. Overly, P. D. Parker, S. E. Parry, M. Reding, J. J. Royall, J. Schulkin, P. A. Sequeira, C. R. Slaughterbeck, S. C. Smith, A. J. Sodt, S. M. Sunkin, B. E. Swanson, M. P. Vawter, D. Williams, P. Wohnoutka, H. R. Zielke, D. H. Geschwind, P. R. Hof, S. M. Smith, C. Koch, S. G. N. Grant, A. R. Jones, An anatomically comprehensive atlas of the adult human brain transcriptome. Nature 489, 391–399 (2012).

34. M. Hawrylycz, J. A. Miller, V. Menon, D. Feng, T. Dolbeare, A. L. Guillozet-Bongaarts, A. G. Jegga, B. J. Aronow, C.-K. Lee, A. Bernard, M. F. Glasser, D. L. Dierker, J. Menche, A. Szafer, F. Collman, P. Grange, K. A. Berman, S. Mihalas, Z. Yao, L. Stewart, A.-L. Barabási, J. Schulkin, J. Phillips, L. Ng, C. Dang, D. R. Haynor, A. Jones, D. C. Van Essen, C. Koch, E. Lein, Canonical genetic signatures of the adult human brain. Nat. Neurosci. 18, 1832–1844 (2015).

35. R. D. Markello, A. Arnatkeviciute, J.-B. Poline, B. D. Fulcher, A. Fornito, B. Misic, Standardizing workflows in imaging transcriptomics with the abagen toolbox. eLife 10, e72129 (2021).

36. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics 36, 2628–2629 (2020).

37. Y. Zhang, S. A. Sloan, L. E. Clarke, C. Caneda, C. A. Plaza, P. D. Blumenthal, H. Vogel, G. K. Steinberg, M. S. B. Edwards, G. Li, J. A. Duncan, S. H. Cheshier, L. M. Shuer, E. F. Chang, G. A. Grant, M. G. H. Gephart, B. A. Barres, Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. Neuron 89, 37–53 (2016).

38. B. B. Lake, S. Chen, B. C. Sos, J. Fan, G. E. Kaeser, Y. C. Yung, T. E. Duong, D. Gao, J. Chun, P. V. Kharchenko, K. Zhang, Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. Nat. Biotechnol. 36, 70–80 (2018).

39. N. Habib, I. Avraham-Davidi, A. Basu, T. Burks, K. Shekhar, M. Hofree, S. R. Choudhury, F. Aguet, E. Gelfand, K. Ardlie, D. A. Weitz, O. Rozenblatt-Rosen, F. Zhang, A. Regev, Massively parallel single-nucleus RNA-seq with DroNc-seq. Nat. Methods 14, 955–958 (2017).

40. S. Darmanis, S. A. Sloan, Y. Zhang, M. Enge, C. Caneda, L. M. Shuer, M. G. Hayden Gephart, B. A. Barres, S. R. Quake, A survey of human brain transcriptome diversity at the single cell level. Proc. Natl. Acad. Sci. 112, 7285–7290 (2015).

41. M. Li, G. Santpere, Y. Imamura Kawasawa, O. V. Evgrafov, F. O. Gulden, S. Pochareddy, S. M. Sunkin, Z. Li, Y. Shin, Y. Zhu, A. M. M. Sousa, D. M. Werling, R. R. Kitchen, H. J. Kang, M. Pletikos, J. Choi, S. Muchnik, X. Xu, D. Wang, B. Lorente-Galdos, S. Liu, P. Giusti-Rodríguez, H. Won, C. A. de Leeuw, A. F. Pardiñas, BrainSpan Consortium, PsychENCODE Consortium, PsychENCODE Developmental Subgroup, M. Hu, F. Jin, Y. Li, M. J. Owen, M. C. O'Donovan, J. T. R. Walters, D. Posthuma, M. A. Reimers, P. Levitt, D. R. Weinberger, T. M.

Hyde, J. E. Kleinman, D. H. Geschwind, M. J. Hawrylycz, M. W. State, S. J. Sanders, P. F. Sullivan, M. B. Gerstein, E. S. Lein, J. A. Knowles, N. Sestan, Integrative functional genomic analysis of human brain development and neuropsychiatric risks. Science 362, eaat7615 (2018).

42. A. T. McKenzie, M. Wang, M. E. Hauberg, J. F. Fullard, A. Kozlenkov, A. Keenan, Y. L. Hurd, S. Dracheva, P. Casaccia, P. Roussos, B. Zhang, Brain Cell Type Specific Gene Expression and Co-expression Network Architectures. Sci. Rep. 8, 8868 (2018).

43. S. Baillet, J. C. Mosher, R. M. Leahy, Electromagnetic brain mapping. IEEE Signal Process. Mag. 18, 14–30 (2001).

44. M. Hämäläinen, R. Hari, R. J. Ilmoniemi, J. Knuutila, O. V. Lounasmaa, Magnetoencephalography---theory, instrumentation, and applications to noninvasive studies of the working human brain. Rev. Mod. Phys. 65, 413–497 (1993).

45. M. D. Rosenberg, E. S. Finn, D. Scheinost, R. T. Constable, M. M. Chun, Characterizing Attention with Predictive Network Models. Trends Cogn. Sci. 21, 290–302 (2017).

46. M. D. Rosenberg, D. Scheinost, A. S. Greene, E. W. Avery, Y. H. Kwon, E. S. Finn, R. Ramani, M. Qiu, R. T. Constable, M. M. Chun, Functional connectivity predicts changes in attention observed across minutes, days, and months. Proc. Natl. Acad. Sci. 117, 3797–3807 (2020).

47. J. Seidlitz, A. Nadig, S. Liu, R. A. I. Bethlehem, P. E. Vértes, S. E. Morgan, F. Váša, R. Romero-Garcia, F. M. Lalonde, L. S. Clasen, J. D. Blumenthal, C. Paquola, B. Bernhardt, K. Wagstyl, D. Polioudakis, L. de la Torre-Ubieta, D. H. Geschwind, J. C. Han, N. R. Lee, D. G. Murphy, E. T. Bullmore, A. Raznahan, Transcriptomic and cellular decoding of regional brain vulnerability to neurogenetic disorders. Nat. Commun. 11, 3358 (2020).

48. D. A. Briley, E. M. Tucker-Drob, Comparing the Developmental Genetics of Cognition and Personality over the Lifespan. J. Pers. 85, 51–64 (2017).

49. J. A. Miller, S.-L. Ding, S. M. Sunkin, K. A. Smith, L. Ng, A. Szafer, A. Ebbert, Z. L. Riley, J. J. Royall, K. Aiona, J. M. Arnold, C. Bennet, D. Bertagnolli, K. Brouner, S. Butler, S. Caldejon, A. Carey, C. Cuhaciyan, R. A. Dalley, N. Dee, T. A. Dolbeare, B. A. C. Facer, D. Feng, T. P. Fliss, G. Gee, J. Goldy, L. Gourley, B. W. Gregor, G. Gu, R. E. Howard, J. M. Jochim, C. L. Kuan, C. Lau, C.-K. Lee, F. Lee, T. A. Lemon, P. Lesnar, B. McMurray, N. Mastan, N. Mosqueda, T. Naluai-Cecchini, N.-K. Ngo, J. Nyhus, A. Oldre, E. Olson, J. Parente, P. D. Parker, S. E. Parry, A. Stevens, M. Pletikos, M. Reding, K. Roll, D. Sandman, M. Sarreal, S. Shapouri, N. V. Shapovalova, E. H. Shen, N. Sjoquist, C. R. Slaughterbeck, M. Smith, A. J. Sodt, D. Williams, L. Zöllei, B. Fischl, M. B. Gerstein, D. H. Geschwind, I. A. Glass, M. J. Hawrylycz, R. F. Hevner, H. Huang, A. R. Jones, J. A. Knowles, P. Levitt, J. W. Phillips, N. Šestan, P. Wohnoutka, C. Dang, A. Bernard, J. G. Hohmann, E. S. Lein, Transcriptional landscape of the prenatal human brain. Nature 508, 199–206 (2014).

50. E. Sareen, S. Zahar, D. V. D. Ville, A. Gupta, A. Griffa, E. Amico, Exploring MEG brain fingerprints: Evaluation, pitfalls, and interpretations. NeuroImage 240, 118331 (2021).

51. F. M. Krienen, B. T. T. Yeo, T. Ge, R. L. Buckner, C. C. Sherwood, Transcriptional profiles of supragranular-enriched genes associate with corticocortical network architecture in the human brain. Proc. Natl. Acad. Sci. 113, E469–E478 (2016).

52. B. D. Fulcher, A. Fornito, A transcriptional signature of hub connectivity in the mouse connectome. Proc. Natl. Acad. Sci. 113, 1435–1440 (2016).

53. K. J. Whitaker, P. E. Vértes, R. Romero-Garcia, F. Váša, M. Moutoussis, G. Prabhu, N. Weiskopf, M. F. Callaghan, K. Wagstyl, T. Rittman, R. Tait, C. Ooi, J. Suckling, B. Inkster, P. Fonagy, R. J. Dolan, P. B. Jones, I. M. Goodyer, the NSPN Consortium, E. T. Bullmore, Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. Proc. Natl. Acad. Sci. 113, 9105–9110 (2016).

54. A. F. Alexander-Bloch, A. Raznahan, S. N. Vandekar, J. Seidlitz, Z. Lu, S. R. Mathias, E. Knowles, J. Mollon, A. Rodrigue, J. E. Curran, H. H. H. Görring, T. D. Satterthwaite, R. E. Gur, D. S. Bassett, G. D. Hoftman, G. Pearlson, R. T. Shinohara, S. Liu, P. T. Fox, L. Almasy, J. Blangero, D. C. Glahn, Imaging local genetic influences on cortical folding. Proc. Natl. Acad. Sci. 117, 7430–7436 (2020).

55. R. D. Markello, J. Y. Hansen, Z.-Q. Liu, V. Bazinet, G. Shafiei, L. E. Suárez, N. Blostein, J. Seidlitz, S. Baillet, T. D. Satterthwaite, M. M. Chakravarty, A. Raznahan, B. Misic, neuromaps: structural and functional interpretation of brain maps. Nat. Methods 19, 1472–1479 (2022).

56. Y. Fu, Z. Ma, C. Hamilton, Z. Liang, X. Hou, X. Ma, X. Hu, Q. He, W. Deng, Y. Wang, L. Zhao, H. Meng, T. Li, N. Zhang, Genetic influences on resting-state functional networks: A twin study: Genetic Influences on Resting-State Functional Networks. Hum. Brain Mapp. 36, 3959–3972 (2015).

57. A. Fornito, A. Zalesky, D. S. Bassett, D. Meunier, I. Ellison-Wright, M. Yucel, S. J. Wood, K. Shaw, J. O'Connor, D. Nertney, B. J. Mowry, C. Pantelis, E. T. Bullmore, Genetic Influences on Cost-Efficient Organization of Human Cortical Functional Networks. J. Neurosci. 31, 3261–3270 (2011).

58. S. Bodenmann, T. Rusterholz, R. Dürr, C. Stoll, V. Bachmann, E. Geissler, K. Jaggi-Schwarz, H.-P. Landolt, The Functional Val158Met Polymorphism of COMT Predicts Interindividual Differences in Brain α Oscillations in Young Men. J. Neurosci. 29, 10855–10862 (2009).

59. J. V. Rétey, M. Adam, E. Honegger, R. Khatami, U. F. O. Luhmann, H. H. Jung, W. Berger, H.-P. Landolt, A functional genetic variation of adenosine deaminase affects the duration and intensity of deep sleep in humans. Proc. Natl. Acad. Sci. 102, 15676–15681 (2005).

60. G. Winterer, R. Mahlberg, M. N. Smolka, J. Samochowiec, M. Ziller, H.-P. Rommelspacher, W. M. Herrmann, L. G. Schmidt, T. Sander, Association Analysis of Exonic Variants of the GABAB-Receptor Gene and Alpha Electroencephalogram Voltage in Normal Subjects and Alcohol-Dependent Patients. Behav. Genet. 33, 7–15 (2003).

61. M. J. Meaney, A. C. Ferguson-Smith, Epigenetic regulation of the neural transcriptome: the meaning of the marks. Nat. Neurosci. 13, 1313–1318 (2010).

62. S. J. Virolainen, A. VonHandorf, K. C. M. F. Viel, M. T. Weirauch, L. C. Kottyan, Gene–environment interactions and their impact on human health. Genes Immun. 24, 1–11 (2023).

63. S. Duncan, L. F. Barrett, Affect is a form of cognition: A neurobiological analysis. Cogn. Emot. 21, 1184–1211 (2007).

64. G. Shafiei, R. D. Markello, C. Makowski, A. Talpalaru, M. Kirschner, G. A. Devenyi, E. Guma, P. Hagmann, N. R. Cashman, M. Lepage, M. M. Chakravarty, A. Dagher, B. Mišić, Spatial Patterning of Tissue Volume Loss in Schizophrenia Reflects Brain Network Architecture. Biol. Psychiatry 87, 727–735 (2020).

65. G. Bush, P. Luu, M. I. Posner, Cognitive and emotional influences in anterior cingulate cortex. Trends Cogn. Sci. 4, 215–222 (2000).

66. M. Goodkind, S. B. Eickhoff, D. J. Oathes, Y. Jiang, A. Chang, L. B. Jones-Hagata, B. N. Ortega, Y. V. Zaiko, E. L. Roach, M. S. Korgaonkar, S. M. Grieve, I. Galatzer-Levy, P. T. Fox, A. Etkin, Identification of a common neurobiological substrate for mental illness. JAMA Psychiatry 72, 305–315 (2015).

67. A. Tanti, C. Belliveau, C. Nagy, M. Maitra, F. Denux, K. Perlman, F. Chen, R. Mpai, C. Canonne, S. Théberge, A. McFarquhar, M. A. Davoli, C. Belzung, G. Turecki, N. Mechawar, Child abuse associates with increased recruitment of perineuronal nets in the ventromedial prefrontal cortex: a possible implication of oligodendrocyte progenitor cells. Mol. Psychiatry 27, 1552–1561 (2022).

68. T. Kaufmann, D. Alnæs, C. L. Brandt, F. Bettella, S. Djurovic, O. A. Andreassen, L. T. Westlye, Stability of the Brain Functional Connectome Fingerprint in Individuals With Schizophrenia. JAMA Psychiatry 75, 749–751 (2018).

69. K. Thuwal, A. Banerjee, D. Roy, Aperiodic and Periodic Components of Ongoing Oscillatory Brain Dynamics Link Distinct Functional Aspects of Cognition across Adult Lifespan. eNeuro 8, ENEURO.0224-21.2021 (2021).

70. B. Voytek, M. A. Kramer, J. Case, K. Q. Lepage, Z. R. Tempesta, R. T. Knight, A. Gazzaley, Age-Related Changes in 1/f Neural Electrophysiological Noise. J. Neurosci. 35, 13257–13265 (2015).

71. B. Voytek, R. T. Knight, Dynamic Network Communication as a Unifying Neural Basis for Cognition, Development, Aging, and Disease. Biol. Psychiatry 77, 1089–1097 (2015).

72. C.-H. Cheng, P.-Y. S. Chan, S. Baillet, Y.-Y. Lin, Age-Related Reduced Somatosensory Gating Is Associated with Altered Alpha Frequency Desynchronization. Neural Plast. 2015, e302878 (2015).

73. T. Hinault, S. Baillet, S. M. Courtney, Age-related changes of deep-brain neurophysiological activity. Cereb. Cortex N. Y. N 1991 33, 3960–3968 (2023).

74. A. Merkin, S. Sghirripa, L. Graetz, A. E. Smith, B. Hordacre, R. Harris, J. Pitcher, J. Semmler, N. C. Rogasch, M. Goldsworthy, Do age-related differences in aperiodic neural activity explain differences in resting EEG alpha? Neurobiol. Aging 121, 78–87 (2023).

75. L. Li, Y. Wei, J. Zhang, J. Ma, Y. Yi, Y. Gu, L. M. W. Li, Y. Lin, Z. Dai, Gene expression associated with individual variability in intrinsic functional connectivity. NeuroImage 245, 118743 (2021).

76. G. Shafiei, S. Baillet, B. Misic, Human electromagnetic and haemodynamic networks systematically converge in unimodal cortex and diverge in transmodal cortex. PLoS Biol. 20, e3001735 (2022).

77. J. C. Pang, K. M. Aquino, M. Oldehinkel, P. A. Robinson, B. D. Fulcher, M. Breakspear, A. Fornito, Geometric constraints on human brain function. Nature 618, 566–574 (2023).

78. T. Sarwar, Y. Tian, B. T. T. Yeo, K. Ramamohanarao, A. Zalesky, Structure-function coupling in the human connectome: A machine learning approach. NeuroImage 226, 117609 (2021).

79. L. E. Liharska, Y. J. Park, K. Ziafat, L. Wilkins, H. Silk, L. M. Linares, R. C. Thompson, E. Vornholt, B. Sullivan, V. Cohen, P. Kota, C. Feng, E. Cheng, J. S. Johnson, M.-K. Rieder, J. Huang, J. Scarpa, J. Polanco, E. Moya, A. Hashemi, M. A. Levin, G. N. Nadkarni, R. Sebra, J. Crary, E. E. Schadt, N. D. Beckmann, B. H. Kopell, A. W. Charney, A study of gene expression in the living human brain. medRxiv [Preprint] (2023). https://doi.org/10.1101/2023.04.21.23288916.

80. T. Yarkoni, R. A. Poldrack, T. E. Nichols, D. C. Van Essen, T. D. Wager, Large-scale automated synthesis of human functional neuroimaging data. Nat. Methods 8, 665–670 (2011).

81. J. Gross, S. Baillet, G. R. Barnes, R. N. Henson, A. Hillebrand, O. Jensen, K. Jerbi, V. Litvak, B. Maess, R. Oostenveld, L. Parkkonen, J. R. Taylor, V. van Wassenhove, M. Wibral, J.-M. Schoffelen, Good practice for conducting and reporting MEG research. NeuroImage 65, 349–363 (2013).

82. F. Tadel, S. Baillet, J. C. Mosher, D. Pantazis, R. M. Leahy, Brainstorm: A User-Friendly Application for MEG/EEG Analysis. Comput. Intell. Neurosci. 2011, 1–13 (2011).

83. A. Schaefer, R. Kong, E. M. Gordon, T. O. Laumann, X.-N. Zuo, A. J. Holmes, S. B. Eickhoff, B. T. T. Yeo, Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cereb. Cortex 28, 3095–3114 (2018).

84. A. Arnatkeviciūtė, B. D. Fulcher, A. Fornito, A practical guide to linking brain-wide gene expression and neuroimaging data. NeuroImage 189, 353–367 (2019).

85. A. Krishnan, L. J. Williams, A. R. McIntosh, H. Abdi, Partial Least Squares (PLS) methods for neuroimaging: a tutorial and review. NeuroImage 56, 455–475 (2011).

86. A. R. McIntosh, B. Mišić, Multivariate statistical analyses for neuroimaging data. Annu. Rev. Psychol. 64, 499–525 (2013).

87. A. R. McIntosh, F. L. Bookstein, J. V. Haxby, C. L. Grady, Spatial pattern analysis of functional brain images using partial least squares. NeuroImage 3, 143–157 (1996).

88. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, G. Sherlock, Gene Ontology: tool for the unification of biology. Nat. Genet. 25, 25–29 (2000).

89. P. D. Thomas, The Gene Ontology and the meaning of biological function. Methods Mol. Biol. Clifton NJ 1446, 15 (2017).

90.   R Core Team, R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, Vienna, Austria, 2022; https://www.R-project.org/).

91.   R. D. Markello, B. Misic, Comparing spatial null models for brain maps. NeuroImage 236, 118052 (2021).

92.   F. Váša, B. Mišić, Null models in network neuroscience. Nat. Rev. Neurosci. 23, 493–504 (2022).

93.   B. Fischl, FreeSurfer. NeuroImage 62, 774–781 (2012).

94.   R. A. Poldrack, A. Kittur, D. Kalar, E. Miller, C. Seppa, Y. Gil, D. S. Parker, F. W. Sabb, R. M. Bilder, The Cognitive Atlas: Toward a Knowledge Foundation for Cognitive Neuroscience. Front. Neuroinformatics 5, 17 (2011).

95.   D. M. Werling, S. Pochareddy, J. Choi, J.-Y. An, B. Sheppard, M. Peng, Z. Li, C. Dastmalchi, G. Santpere, A. M. M. Sousa, A. T. N. Tebbenkamp, N. Kaur, F. O. Gulden, M. S. Breen, L. Liang, M. C. Gilson, X. Zhao, S. Dong, L. Klei, A. E. Cicek, J. D. Buxbaum, H. Adle-Biassette, J.-L. Thomas, K. A. Aldinger, D. R. O'Day, I. A. Glass, N. A. Zaitlen, M. E. Talkowski, K. Roeder, M. W. State, B. Devlin, S. J. Sanders, N. Sestan, Whole-Genome and RNA Sequencing Reveal Variation and Transcriptomic Coordination in the Developing Human Prefrontal Cortex. Cell Rep. 31, 107489 (2020).

96.   Y. Wei, S. C. de Lange, L. H. Scholtens, K. Watanabe, D. J. Ardesch, P. R. Jansen, J. E. Savage, L. Li, T. M. Preuss, J. K. Rilling, D. Posthuma, M. P. van den Heuvel, Genetic mapping and evolutionary analysis of human-expanded cognitive networks. Nat. Commun. 10, 4839 (2019).

97.   M. Guardiola-Ripoll, M. Fatjó-Vilas, A Systematic Review of the Human Accelerated Regions in Schizophrenia and Related Disorders: Where the Evolutionary and Neurodevelopmental Hypotheses Converge. Int. J. Mol. Sci. 24, 3597 (2023).

98.   K. M. Girskis, A. B. Stergachis, E. M. DeGennaro, R. N. Doan, X. Qian, M. B. Johnson, P. P. Wang, G. M. Sejourne, M. A. Nagy, E. A. Pollina, A. M. M. Sousa, T. Shin, C. J. Kenny, J. L. Scotellaro, B. M. Debo, D. M. Gonzalez, L. M. Rento, R. C. Yeh, J. H. T. Song, M. Beaudin, J. Fan, P. V. Kharchenko, N. Sestan, M. E. Greenberg, C. A. Walsh, Rewiring of human neurodevelopmental gene regulatory programs by human accelerated regions. Neuron 109, 3239-3251.e7 (2021).

99.   S. L. W. Driessens, A. A. Galakhova, D. B. Heyer, I. J. Pieterse, R. Wilbers, E. J. Mertens, F. Waleboer, T. S. Heistek, L. Coenen, J. R. Meijer, S. Idema, P. C. De Witt Hamer, D. P. Noske, C. P. J. De Kock, B. R. Lee, K. Smith, J. T. Ting, E. S. Lein, H. D. Mansvelder, N. A. Goriounova, Genes associated with cognitive ability and HAR show overlapping expression patterns in human cortical neuron types. Nat. Commun. 14, 4188 (2023).

100.   R. N. Doan, B.-I. Bae, B. Cubelos, C. Chang, A. A. Hossain, S. Al-Saad, N. M. Mukaddes, O. Oner, M. Al-Saffar, S. Balkhy, G. G. Gascon, M. Nieto, C. A. Walsh, Mutations in Human Accelerated Regions Disrupt Cognition and Social Behavior. Cell 167, 341-354.e12 (2016).

101. A. I. Luppi, P. A. M. Mediano, F. E. Rosas, N. Holland, T. D. Fryer, J. T. O'Brien, J. B. Rowe, D. K. Menon, D. Bor, E. A. Stamatakis, A synergistic core for human brain evolution and cognition. Nat. Neurosci. 25, 771–782 (2022).

Acknowledgements:

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The Brainstorm app is supported by funding to SB from the NIH (R01-EB026299), a Discovery grant from the Natural Science and Engineering Research Council of Canada (436355-13), the CIHR Canada Research Chair in Neural Dynamics of Brain Systems, the Brain Canada Foundation with support from Health Canada, and the Innovative Ideas program from the Canada First Research Excellence Fund, awarded to McGill University for the HBHL initiative. This work was supported by a doctoral fellowship from NSERC (JDSC, JYH).

Author Contribution:

Conceptualization: JDSC, SB
Data Curation: JDSC, SB
Methodology: JDSC, JP, JYH, BM, SB
Software: JDSC, JP, JYH, SB
Visualization: JDSC, JP, SB
Resources: JYH, SB
Validation: JDSC, SB
Formal analysis: JDSC
Supervision: SB
Project administration: SB
Funding acquisition: SB
Writing – original draft: JDSC, SB
Writing – review & editing: JDSC, JP, JYH, BM, SB

Competing interests:

All authors declare no competing conflicts of interest. The listed funding sources in the Acknowledgements did not play any role in the writing of the manuscript or the decision to submit this manuscript for publication

Data and materials availability:

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. The data are available through the Human Connectome Project (HCP) repository (https://www.humanconnectome.org/study/hcp-young-adult). Gene expression data are available through the Allan Human Brain atlas (ABHA; http://human.brain-map.org/). The Neurosynth database is available at https://neurosynth.org/.

All in-house code used for data analysis and visualization is available on Zenodo https://zenodo.org/records/15023999 and our GitHub https://github.com/Epideixx/Fingerprints_Twins.