Developmental changes in mouse motor skill learning and cortical circuitry

Abbreviated title: Developmental plasticity in the mouse motor system

Taehyeon Kim[1] and Bryan M. Hooks[1]

[1]Center for Neuroscience University of Pittsburgh (CNUP) and Department of

Neurobiology, University of Pittsburgh School of Medicine, Pittsburgh, PA

Correspondence: Dr. Bryan M. Hooks, 200 Lothrop Street BSTWR Suite W1458

Pittsburgh, PA 15213; 412-624-8465; hooksm@pitt.edu (B.M.H.) and TAK127@pitt.edu

(T.K.)

Number of pages: 42

Number of figures: 7 (5 Supplementary Data figures)

Number of tables: 0

Words for abstract (196), significance (132), introduction (890), and discussion (2374)

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: We thank Quincy Erickson-Oberg for assistance with the beam

break experiment, Tom Deakin for assistance with rotarod experiment, and Kyle

Bannerman for assistance with the gait analysis. We thank Omar Gharbawie, Chinfei

Chen, Ghanshyam Sinha, and other members of the Hooks lab for feedback on the

manuscript. This work was supported by a NARSAD Young Investigator Award (BMH),

NIH NINDS R01 NS103993 (BMH), a Whitehall Foundation award (BMH), and an

American Heart Association Predoctoral Fellowship (TK).

Abstract

Learning motor skills requires plasticity in the primary motor cortex (M1), including changes in inhibitory circuitry. But how inhibitory synaptic connections change during skill acquisition and whether this varies over development is not fully understood. This study assesses the normal developmental trajectory of motor learning and then addresses inhibitory connectivity changes after motor learning. We trained mice of both sexes to run on a custom accelerating rotarod at ages from postnatal day (P) 20 to P120, tracking paw position and quantifying time to fall and changes in gait pattern. Performance improved most rapidly between P30-60, while paw position and gait patterns change with learning, though differently between age groups. To address circuit changes, we labeled task-active and task-inactive pyramidal cells with CaMPARI2, a genetically encoded activity marker. We then evoked inhibitory responses (IPSCs) from two major interneuron types: parvalbumin-expressing (PV+) interneurons and somatostatin-expressing (SOM+) interneurons. After one training day, PV-mediated inhibition is greater in task-active cells, while SOM-mediated inhibition is not different. These results suggest early changes in PV-mediated inhibition may support motor skill acquisition in mice. Whether PV-mediated inhibitory changes persist or changes in SOM+ interneuron connections arise later in training remains to be tested.

Highlights

*Motor learning aptitude is developmentally regulated, peaking at P30-P45 for the cortically-dependent accelerating rotarod task

*Mice learn by shifting from hopping to walking gait, with the gait shift in forelimbs preceding the shift in hindlimbs

*Inhibitory responses in task-active neurons are stronger than task-inactive neurons from parvalbumin-expressing fast-spiking neurons on the first training day

*Other interneuron connections from somatostatin-expressing interneurons are unchanged

Significance Statement

Plasticity in sensory cortex is restricted to a limited developmental window. Learning motor skills requires motor cortex plasticity, but it is unknown whether there are changes in learning aptitude over development and whether the synaptic mechanisms of plasticity vary across cortical areas. Here, we define the developmental trajectory of motor learning aptitude for the accelerating rotarod task in mice. We find that learning peaks at P30-P45, with mice learning to shift from hopping to walking gait to stay on the rotarod longer. Further, the gait shift in forelimbs precedes hindlimbs. We then find, using recordings targeted to task-active neurons, that inhibitory responses from specific subtypes of interneurons (parvalbumin-expressing) are stronger in active cells, while other interneuron connections (somatostatin-expressing) are unchanged. These results suggest early changes in PV-mediated inhibition support motor skill acquisition.

Introduction

The critical period is a time during early development when neural circuits exhibit heightened plasticity in response to sensory experience[1–4]. The same sensory experience after the critical period does not cause the same magnitude of changes in the circuitry. Sensory areas have been extensively studied to define the critical period and its underlying circuit mechanisms. In both the visual and somatosensory systems, critical period timing is defined by deprivation of the corresponding sensation, such as incoming visual or whisker stimuli, and the resulting effects in the nervous system.

Plasticity in primary motor cortex (M1) is vital for the learning and executing of complex motor skills[5–7]. Studies of cortical plasticity have generally investigated sensory critical period plasticity, where manipulations at defined early ages can cause long-lasting changes in cortical connectivity and responsiveness, while plasticity is reduced or absent at later ages[8–10]. For visual areas, for example, critical period plasticity is limited to ~P18-P28 in mice[2,4,9]. Fewer studies have characterized plasticity during the corresponding period in M1 using motor skill acquisition. Most motor skill acquisition studies use young adults >1 month old. Motor skill acquisition across developmental ages in younger animals, closer to ages equivalent to the sensory critical period, is less unexplored. However, this requires addressing learning in a task that is cortex-dependent as well as one that animals of all ages can learn. Further, because skill acquisition can continue throughout life, it is worth exploring the degree to which plasticity is altered in later ages. Investigating age-related changes in motor learning will determine whether a period of heightened plasticity for motor learning exists, and how circuits mediate motor learning.

Voluntary movements and motor skill acquisition involves activity of subset of cortical cells, forming local circuit ensembles that may contain from 10-50% of local neurons depending on the task[11–14]. Motor skill acquisition is associated with structural and functional changes in these specific ensembles of excitatory neurons[13,15,16]. Neurons not related to the task may not undergo the same changes in connectivity and excitability. Because excitatory cells are the vast majority of cortical cells, much prior work, especially before transgenic approaches, has focused on local and long-range excitatory connectivity. M1 shares a number of local connectivity rules with S1 and other

sensory cortical areas[17–19], making a comparison to sensory plasticity of interest,

especially as regards plasticity mechanisms. Prior work has suggested changes in the

strength of corticocortical connectivity, potentially restricted to the upper cortical layers

during motor learning[20–22]. This includes reweighting of long-range corticocortical and

corticothalamic inputs[22]. Thus, plasticity in upper layers is likely to be involved in motor

learning.

110   Cortical inhibitory circuitry is crucial for regulating plasticity. Inhibitory neurons

are key circuit nodes in triggering in cortical plasticity in sensory areas. The opening of

the critical period is caused in part by the maturation of GABAergic inhibition[2]. Further,

transplantation of the inhibitory neurons can induce a period of heightened plasticity

later in life[23,24]. Similarly, in motor areas, inhibition also regulates plasticity. Reduced

inhibition is inversely correlated with motor learning performance in normal and post-

injury conditions in humans and mice[25–30]. Still, the mechanisms governing plasticity for

motor learning are not fully known. Many, but not all, features of circuit connectivity and

plasticity are conserved across cortical areas. About 30% of cortical neurons are

inhibitory interneurons[31,32]. Subtypes of inhibitory neurons can be further categorized

based on molecular markers, electrophysiological characteristics and morphology[31,32].

Two major inhibitory interneuron types are parvalbumin-expressing (PV+) and

somatostatin-expressing (SOM+) interneurons [31]. PV+ and SOM+ interneurons connect

in the circuit differently. PV+ interneurons inhibit excitatory neurons, synapsing near the

soma, while SOM+ interneurons synapse on distal dendrites of excitatory cells[33–35]. PV+

inhibition can be reweighted rapidly in an activity-dependent manner, providing greater

inhibition to more active pyramidal cells[36,37].

127   How does the role of interneuron plasticity in sensory critical periods compare to

that in M1? It is not as straightforward to fully deprive M1 of activity, though it has been

attempted[38,39]. Thus, we chose skill acquisition to observe changes during development.

Similar to excitatory neurons, inhibitory neurons go through both structural and

functional changes. SOM+ interneurons show a temporal shift in their activity with

learning[40]. PV+ interneurons in particular have a transient increase in activity in motor

cortex and retrosplenial cortex[41]. Due to their axodendritic targeting[34,35], SOM+

Figure 1 | Rotarod skill acquisition plateaus around P30-P60 on training day 1.
(A) Schematic of rotarod task with high speed videography and barriers to create the running lane during a trial. (B) Video frame of mouse during running. Paw positions (LFP, left front paw; LHP, left hind paw; RFP, right front paw; RHP, right hind paw) and the nose and tail base are tracked using DeepLabCut. (C) Learning curve during 20 trials on training day 1 (TD1). Six colors represent the six ages tested (P15-20, P30, P45, P60, P90, and P120). Gray bars represent time windows used for early and late performance to assess TD1 improvement. (D) Improvement (in seconds) in time to fall during TD1 for six age groups. Mean+/- SEM. N=63, 92, 60 63, 27, 12 . (E) Improvement (in seconds) in time to fall during TD1 for six age groups, subdivided by male (blue) and female (red). Mean+/- SEM.(F) Improvement (in seconds) in time to fall during five training days. Gray vertical lines divide the days. Mean+/- SEM. (G) The effect of M1 inactivation by bilateral muscimol injection on rotarod skill acquisition compared to saline injected controls. N=5 (control); N=9 (muscimol). Mean+/- SEM. Significance: *, p<0.05; **, p<0.01; ***, p<0.001.

interneurons are well-situated to mediate the changes in dendritic spines with motor

learning. Transient silencing of SOM+ cells leading impairment in motor learning further

implicate SOM+ cells' role in skill acquisition[1,40], but exact synaptic changes, including

synaptic strength, are unclear. Learning a lever pressing task, another type of motor

learning paradigm, induces structural changes in axonal boutons of both PV+ and

SOM+ populations[1]. While PV+ axonal boutons transiently increased, SOM axonal

boutons decreased, but more slowly over several days of learning[1]. Similar interneuron-

dependent changes likely occur in M1 during rotarod training. It is unknown whether the

postsynaptic targets of these axons that undergo dynamic changes are task-related

cells. The experiments here test whether changes in cortical inhibition regulate M1

plasticity. Below, we directly test whether the observed structural changes in axonal

boutons lead to functional changes of inhibitory synapses by defining ages with peak

plasticity and addressing whether changes in interneuron-mediated inhibition strength

occurs during motor learning.

Results

Rotarod performance peaks and plateaus around P30-60.

Figure 2 | Tracking step kinematics during rotarod training.
(A) Schematic of rotarod task with video frame of tracked mouse. (B) Cartoon of tracked image frame illustrating location of the mouse paws (LFP, left front paw; LHP, left hind paw; RFP, right front paw; RHP, right hind paw) and the nose and tail base using DeepLabCut. The rotarod and direction of motion are indicated, with the front and rear ends of the rod serving as 0.0 and 1.0 in normalized coordinates. At bottom, coordinates for the step start, step end, and step length are plotted to illustrate how they are determined. (C) Plot of the x-position of a single paw (LHP, left) or all paws (right) during a trial. Time given in frames (converted to seconds with example timebar). (D) Expansion of paw tracker showing step detection. Steps (red) are automatically detected (see methods). Step start (green), step end (black), and step length are then calculated. (E) Step length versus time to fall for the left front paw, plotted for all ages combined (red) and individual ages (20 and above). Data are fit to a line and the linear fit compared at right. Summary plots include step length versus time to fall for the right front paw, left hind paw, and right hind paw (far right). (F) Step end point presented similarly to step length.

Mice become ambulatory by P15. Walking is part of natural locomotion that can
be tracked in mice at a range of ages and sizes. To test motor learning, we used the
accelerating rotarod task, which takes advantage of walking on an accelerating rod.
Furthermore, improvement in the task can be observed within a single training day. The
time to fall and changes in foot position were measured to determine the learning curve
at different ages. To monitor changes in the gait, we built a custom rotarod made of
clear plexiglass and a high-speed camera was mounted at the bottom to record the
behavior (Figure S1A-B). Six age groups were used to test the age-related differences
in the aptitude to learn a motor task: P15-20, P30, P45, P60, P90 and P120 (Figure 1A-
D). Typical behavior during the first training day for these ages tended to improve during
the first half and reach a plateau within the first session, with some reduction in
performance in later training day 1 trials. We wanted to measure the improvement by

comparing the time to fall when the animal was at beginner level (early phase of
learning) and when the animal reached an expert level (when they reached plateau).
Thus, improvement was calculated by subtracting the average time to fall measured in
trials 11-13 (when the performance tends to reach plateau) from the average time to fall
measured in trials 1-3 (beginner; Figure 1B-D). Within the first two months of age, the
improvement was greater for the animals in P30-60 compared to those in the P15-20
group (Figure 1D, $p<0.001$). Compared to P90-120 animals, the animals in P45-60
showed significantly greater improvement (Figure 1D, $p<0.01$ (P45) and $p<0.05$ (P60)).
Testing a behavioral task across different age groups came with some
confounds. Although this task used no external rewards to exploit the natural walking
behavior in mice, it was possible that age-dependent differences in the time spent
walking in the home cage might contribute to rotarod performance. To assess if there
was a difference in baseline locomotion, animals were tracked in a standard cage with
infrared beam breaks to determine the number of times they crossed the length of the
cage. P15 mice crossed the least number of beam breaks compared to the other age
groups (Figure S2, $p<0.001$). P45 animals exhibited a greater number of beam breaks
compared to P30 (Figure S2A, $p<0.001$), but there were no other significant differences
within these older ages.
We next considered sex as a variable. Significant sex differences in improvement
were observed in the P30, P45, and P60 groups (Figure S3, $p<0.05$), with young adult
female mice outperforming males in these age groups. Because adult male mice tend to
weigh more than female mice, we considered that the sex difference in performance
might be due to weight. From P20 to P120, each age groups gained significant weight
(Figure S2C). Male mice significantly outweighed female mice at ages above P30
(Figure S2D). Across all ages, there was no correlation between weight and
improvement, nor was there a correlation in a specific sex (Figure S2E). P60 had a
significant correlation between improvement and weight ($p=0.0496$). However, no
particular age group showed sex-specific correlation with performance. Therefore,
weight alone did not explain differences in performance, since there were not significant
differences between heavier and lighter mice at P30-P60.

Figure 3 | Gait pattern changes with age and training.
(A) Schematic of rotarod task with video frame of tracked mouse. (B) Cartoon of tracked image frame illustrating location of the mouse paws (LFP, left front paw; LHP, left hind paw; RFP, right front paw; RHP, right hind paw) and the nose and tail base using DeepLabCut. The rotarod and direction of motion are indicated, with the front and rear ends of the rod serving as 0.0 and 1.0 in normalized coordinates. At bottom, coordinates for the step start, step end, and step length are plotted to illustrate how they are determined. (C) Plot of the x-position of a single paw (LHP, left) or all paws (right) during a trial. Time given in frames (converted to seconds with example timebar). (D) Expansion of paw tracker showing step detection. Steps (red) are automatically detected (see methods). Step start (green), step end (black), and step length are then calculated. (E) Step length versus time to fall for the left front paw, plotted for all ages combined (red) and individual ages (20 and above). Data are fit to a line and the linear fit compared at right. Summary plots include step length versus time to fall for the right front paw, left hind paw, and right hind paw (far right). (F) Step end point presented similarly to step length. Significance: *, p<0.05; **, p<0.01; ***, p<0.001.

193   To determine whether learning on the accelerating rotarod is M1-dependent task,

194 M1 was pharmacologically inactivated by bilaterally injecting muscimol in P30-45

195 animals. Mice that received either muscimol or saline bilateral injections were permitted

196 to recover from the procedure for 40 minutes and showed normal ambulation in the

197 home cage after recovery. When tested on the accelerating rotarod, muscimol-injected

198 animals did not show improvement compared to the saline-injected littermates,

199 confirming that the task is an M1-dependent task (Figure 1G).

201 **Changes in paw kinematics are correlated with better performance**

202   Although time to fall was an effective measure of overall task performance, a

203 more detailed understanding of how animals improved during the task was achieved by

204 high-speed videography to quantify changes in limb movements. This approach tracked

205 limb kinematics by acquiring video of paw position through the clear rod. During rotarod

206 training, four paws, the nose, and the base of the tail were tracked with DeepLabCut

(Figure 2A-B). Estimated positions of these parts were tracked in x- and y-coordinates
in the video, with the x-axis aligned to the direction of rod rotation. Paw positions were
filtered by likelihood (>0.95) and steps were automatically detected based on the
prominence of the peak in the trajectory with a minimum interstep interval using custom
software in Matlab. Step end and step start positions thus extracted were then
normalized to the width of the rod (Figure 2B). Step length was the normalized distance
between the step end and the start (Figure 2B). Raw position in pixels was plotted to
demonstrate accuracy of the step detection, while the paw position for all paws can be
simultaneously tracked (Figure 2C-D).
We then examined how paw kinematics changed during learning by comparing
the normalized step length and step end point with performance (Figure 2E-F and
Figure S4), measured as time to fall. When pooling data across ages, a few general
trends emerged. First, longer step length was associated with longer times on the
rotarod. This was consistent across all ages measured (P20-P60). The increase in step
length was similar for both front and hind paws. Second, the step end point moved
closer to the front edge of the rotarod. This effect was strongest for front paws but was
present to a lesser degree for hind paws (as expected if step size also increases). This
effect, however, varied with developmental age, being present and pronounced in P60
animals, shifting the location of the front paw to end steps from about 0.4 (closer to the
middle) to about 0.2 (close to the edge of the rod). At other ages, the step end position
is static.
228
Changes in gait patterns are correlated with better performance
The rotarod task involves not only kinematic changes in individual paws, but also
the coordination of left and right paws. Animals engaged in different types of gait
patterns during the task – simultaneously moving both paws (hopping; Figure 3A) or
alternating the left and right paws (walking; Figure 3A). To determine how the gait
pattern changed, we needed to consider the phases of steps between left and right
paws. We compared the step times of the paws and assigned each step a score (range
0.0 to 1.0). The step time of the left paw (Figure 3C, green for the left hind paw) was
marked. The time between two left steps was normalized from 0.0 to 1.0. The time of

Figure 4 | P-values for the gait analysis.
(A) Due to variations in the number of steps, the number of steps in a trial were divided into deciles to facilitate the comparison of average step phases across trials and age groups. The average step phase value for each age for front and hind limbs is plotted. (B) The P-value plot of deciles compares the first decile of trials 1-20 to rest of the deciles (2-9) for each trial. The top row compares front paws at four ages and the bottom row compares hind paws. Non-significant p-values are purple; p-values < 0.05 are as shown in the lookup table. (C) The P-value plot comparing across trials compares each decile from trials 2-20 with the corresponding decile from trial 1. The top row compares front paws at four ages and the bottom row compares hind paws. For instance, the front paw in P20 Decile 3 from trial 3 is compared to Decile 3 from trial 1. Non-significant p-values are purple; p-values < 0.05 are as shown in the lookup table. (D) The P-value plot of deciles compared across ages assess P20 versus P30, P45, and P60 groups for 20 trials. Each individual graph represents the comparison between the four age groups for a given trial. The X-axis denotes deciles, with a specific trial's decile being compared to the corresponding decile of the P20 group. For example, the front paw in P30 Decile 9 is compared to P20 Decile 9 in the trial 1 graph, and if significantly different, it is indicated by a bump in the y-axis. Significance: *; p<0.05. **;p<0.01, ***;p<0.001.

the right hind paw step (orange) was then computed as a fraction of the relative

distance between left steps. Alternating gait occurred when phase difference of the left
and right paws was closer to 0.5 while the simultaneous or hopping gait was closer to
0.0 or 1.0 (Figure 3C). Note the color map at the bottom that signifies the alternating
gait as red and the hopping gait as blue. The scale was reflected because we treated
step phase scores from 0.5 to 1.0 as equivalent to 0.5 to 0.0 to assess alternating gait.
To determine changes in gait pattern on the first training day and how these differed
with limb, age, and trial number, the data were plotted as follows. For each step taken,
such as the first step on the first trial (of 20), the mean step phase score was computed
across all mice tracked, and the corresponding color plotted as a tick mark (Figure 3D).
These were generally blue (simultaneous/hopping gait). This was repeated for each
subsequent step and for each trial. The number of steps included in the average varied
across the row for each trial, reflecting the variability in performance (as observed in
time to fall) within and across trials. Qualitatively, both front limb and hind limb and
within each age, the gait pattern generally shifts from hopping to walking (alternating
gait). This shift seemed to occur more rapidly for the front paws than the hind paws,
suggesting that the transition to a walking gait can occur at different time for the two
pairs of limbs.

To determine whether the gait pattern shifted significantly on the first training
day, the cumulative frequencies of the step phases at four different stages of the
training were plotted (Figure 3E): beginner (trials 1-5), early learning (trials 6-10), mid-
learning (trials 11-15), and late learning (trials 16-20). The offset in step phase between
front limbs (dotted lines, lower right) and hind limbs (solid line, upper left) was evident in
the data, suggesting that the front and hindlimbs employ different strategies during
rotarod training. Cumulative frequencies of the phases in beginner stage were
compared to the other three stages. Across all age groups, the front paws shifted to
alternating gait starting from early learning at all ages (trials 6-10) (Figure 3E, left). On
the contrary, the shift from simultaneous hopping to alternating in the hind paws was
delayed in the P20 group (Figure 3E, left). While animals in P30-60 groups transitioned
to alternating by early learning (trial 6-10), P20 groups did not start alternating until late
learning (trials 16-20).

Figure 5 | Schematic of CaMPARI2 manipulation.
(A) AAV is used to express CaMPARI2 in M1 in PVcre+/-;Ai32+/- or SOMcre+/-;Ai32+/- mice. (B) CaMPARI2-expressing cells are photoconverted to red by a skull-mounted UV LED. (C) Example images of a slice after photoconversion during rotarod. (E) Example traces from GFP+(green) and RFP+ (red) neurons after optogenetic stimulation of PV+ cells. (F) The photoconversion ratio is calculated by dividing red fluorescence by green fluorescence. Animals showed a greater photoconversion ratio after the rotarod task (grey) than the control animals that freely moved in the chamber (white). Significance: *; p<0.05.

However, the change in the distribution of the gait patterns over time within a trial
were not well-quantified in this manner. We wanted to compare whether there was a
difference in the emergence of a gait pattern and compare between age groups and
trials. To do so, the entire length of the steps in a trial were averaged into deciles which
were then compared between deciles, trials and age groups (Figure 4A). This appeared
similar to the raw data (Figure 3D), with all trials now scaled to the same length. This
approach allowed us to look at the changes in gait pattern despite the difference in the
number of steps in each trial. First, to compare whether gait pattern evolved within a
trial, we compared the first decile of a given trial to the rest of the deciles in the same
trial. This showed the strongest effect: The phase of the front paws, particularly in P30-
60 groups, showed significant differences compared to the first deciles across many
trials (Figure 4B). P30 groups showed most difference in the phase of the front paw
compared to the first decile. Hindpaws seemed to show few significant differences
across deciles. P30 and P60 showed some difference in late deciles while P45 showed
a few early-mid late deciles difference along with later decile differences in mid learning.

P20 group showed minimal difference in phase of hind paws. Second, to test whether

there was a difference in gait across trials, the decile of the first trial was compared to

the corresponding decile of the rest of the trials (trial 2-20). This effect was considerably

weaker (fewer significant comparisons). In the P20 group, both front and hind paws

showed little difference across trials (Figure 4C). In the P30 group, the difference in

front paw occurred early-mid learning (as early as trial 5) in late deciles (Figure 4C). In

the P45 group, the difference in the front paw appeared later and became more evident

around mid-learning, particularly in early-mid deciles. In the P60 group, differences in

front paw started to predominantly appear in late trials on the first training day (Figure

4C). As before, the difference in hind paw gait across the age groups was minimal and

inconsistent (Figure 4C). Lastly, to compare between age groups, we compared a decile

in a given trial of P20 to a corresponding decile in the same trial of P30, P45 or P60

(Figure 4D). Front paws of P30 started to show differences in early trials and differed

from P20 throughout many of the trials (Figure 4D). On the other hand, P45 and P60

showed difference to P20 later trials (Figure 4D). Trial 14 seemed to show very different

gait pattern in P30-60 compared to P20(Figure 4D). Hind paws P30-60 showed

differences compared to P20 as early as Trial 3 during early learning and this persisted

throughout many trials (Figure 4D).

PV+ interneurons provide greater inhibition to task-active cells following one

training day

When the cells are active during rotarod performance, CaMPARI2 is

photoconverted from green to red fluorescence following UV-light delivery, while the

CaMPARI2 in inactive cells remains green (Figure 5A-D)[42]. To ensure that

photoconversion was specific to activity during the rotarod task, we performed a control

experiment. Animals aged from P20-60 received UV-light while freely moving at the

bottom of the behavior chamber under the rotarod to simulate the same environment

without motor learning. The duration for which the animals stayed in the chamber was

determined by the average time to fall in training day 1 for the age- and sex-matched

group. Photoconversion (red/green) ratio was significantly less than those measured

from animals that completed the rotarod task (Figure 5F). To assess changes in

Figure 6 | IPSCs recorded in task-active (RFP+) vs. task-inactive cells (GFP+) after stimulating PV+ interneurons
(A-E) Scatter plot of IPSC responses in pairs of task-active (red) and task-inactive (green) cells in P20 (A), P30 (B), P45 (C), P60(D) and across all age groups (E). Solid line is the geometric mean of the pair responses. Dotted line is the line of equality (x=y). (F) Slopes of geometric mean across all ages plotted together. (G-K) Ratio of responses of task-active cells to their partner task-inactive cells after PV+ stimulation in P20 (G), P30 (H), P45 (I), P60 (J) and across all age groups (K). Significance: *, p<0.05; **, p<0.01; ***, p<0.001. (Also indicated with red arrows in (A-E). (L) Schematic of the genetic strategy for testing PV output to M1 pyramidal neurons.

inhibitory synaptic connectivity, we made whole-cell recordings in M1 layer 5 excitatory

neurons that were either red or green to determine whether the inhibition provided by

either PV+ or SOM+ interneurons differs in an activity-dependent manner. Inhibitory

responses (IPSCs) were optogenetically evoked in either PV+ or SOM+ neurons to

measure synaptic strength (Figure 5E).

Figures 6 and 7 show the scatter plot of peak IPSC recorded in pairs of task-

active (red) and task-inactive (green) cells (same slice) after evoking either PV+ or

SOM+ interneurons in animals. The dotted line on the plot is unity line where the

responses of task-active and -inactive cells would be equal, and the slope of the solid

line is a geometric mean. Across all ages, the geometric mean of the pair responses

lied left of a line of equality for PV+ stimulation groups (Figure 6). Thus, a shift to the left

of the dotted line represents input favoring task-active neurons, while a shift to the right

favors task-inactive neurons. Across all ages, PV-mediated inhibition was greater in red

(task-active) than those in green (task-inactive) neurons (Figure 6E, p<0.05).

Furthermore, responses recorded in RFP+ cells were normalized to their partner GFP+

cells (Figure 6G-K and 7G-K). After normalization, PV-mediated inhibition in red (task-

active) neurons was greater than those in green (task-inactive) neurons in all ages

combined (Figure K, p<0.01). P30 and P45 groups showed a trend in greater inhibition

on red (task-active) compared to green (task-inactive) cells (Figure 6H and I). On the

contrary, when SOM+ cells were optogenetically stimulated, IPSCs recorded in red

Figure 7 I IPSCs recorded in task-active (RFP+) vs. task-inactive cells (GFP+) after stimulating SOM+ interneurons
(A-E) Scatter plot of IPSC responses in pairs of task-active (red) and task-inactive (green) cells in P20 (A), P30 (B), P45 (C), P60(D) and across all age groups (E). Solid line is the geometric mean of the pair responses. Dotted line is the line of equality (x=y). (F) Schematic of the genetic strategy for testing SOM output to M1 pyramidal neurons. (G-K) Ratio of responses of task-active cells to their partner task-inactive cells after SOM+ stimulation in P20 (G), P30 (H), P45 (I), P60 (J) and across all age groups (K).

(task-active) cells were not significantly different from those recorded in green (task-

inactive) cells (Figure 7G-K). Thus, our data suggest that motor learning induced cell-

type-specific changes in inhibition depend on the activity of the postsynaptic target on

the first training day, but only for PV+ and not SOM+ interneuron connections.

Discussion

Rotarod Motor Skill Learning is Fastest in a Given Developmental Window

To address age-dependent differences in motor learning, we first used a walking

task that was consistent with the range of natural behaviors in mice. Walking is a

common behavior in vertebrates that relies, in part, on central pattern generators in the

spinal cord[43]. For mice, walking begins naturally prior to the ages we test. To assess

whether there is a sensitive (critical) period in the ability to acquire a motor skill, we

tested mice as young as P15 (who could still execute the task) up to age P120 on the

accelerating rotarod. Using time to fall as a measure of learning, these data show that

first day improvement in performance on the rotarod varies with age, plateauing at P30-

60. This suggests that there are differences in the aptitude for motor learning across

developmental time. This is unlikely to be due to difference in the baseline locomotion,

which does not show a heightened activity selectively at P30-60 (Figure S2A-B).

Previous experiments had also shown that prior exposure to exercise did not affect

learning[44]. Control experiments also ruled out effects of weight within each age cohort

(Figure S2E). Since improvement does occur over five days of training for all age

groups of mice (Figure 1F), the animals are physically capable of executing the task at

all ages. Although using a skilled reaching task to explore age-dependent differences in

skill learning is possible, it was not as clear that P15-P20 mice could execute this task.

The rotarod task was unrewarded, as mice walk naturally and running is an

intrinsically motivated behavior[45,46]. This removes the confound of reward circuitry with

motor learning. Although all ages learn, and some learn faster, it is not clear whether

other aspects of the animal's motivational state might differ across age and contribute to

task performance. If low motivation accounted for poor TD1 performance in P20, P90,

and P120, this does not explain how motivation changes on subsequent days of training

as behavior improves, while potentially differently motivated P30, P45, and P60 mice do

not have large swings in motivation. It is possible that instead of motivation, differences

in anxiety contribute to performance differences at different ages. An assay for fear

might clarify this, though a nonlinear age-dependent variation in fear would be

interesting to explore in other circuits.

It is difficult to assess whether learning is a shift in mindset where the animal

understands the goal of the task (walk so as to stay on the rod), or whether the goal is

clear and learning is a shift in precision of skilled movement. Both shifts could result in a

successful increase in time to fall. Although these experiments do not clearly distinguish

between these alternatives, tracking the kinematics of paw movement suggests that, at

least in part, better performance is associated with changes in motor control. This

seems to be the case because the step lengths for all paws evolve during the task

(Figure 2E), front paw step end points are targeted closer to the end of the rod, at least

for P60 (Figure 2F), and gait pattern evolves with time during the task (Figure 3 and

4B). Differences in sex did account for differences in performance, with female mice

performing better during P30-P60 (also corresponding to the peak period of learning-

related plasticity). Beyond ruling out weight as a factor, the basis for improved female

performance during this time window was not explored further, though there is evidence

of mechanism differences in other systems[47] that invite future efforts to probe M1

circuitry in a similar way.

Because spinal circuits also play a role in control of walking, it is important to test
whether rotarod skill learning depends on motor cortex. This task has been used in
previous studies of motor cortex to assess circuit changes during learning. Specifically,
rates of spine formation and retention in forelimb M1 change during rotarod learning[15,16].
Not only are there structural changes, but the firing patterns of neural ensembles show
task-specific modulation during training and reorganize over several sessions to support
motor learning task[11]. Together these imply that circuit changes assist in acquiring a
new skill. Consistent with this, our data shows that bilateral inhibition of forelimb M1 with
muscimol during rotarod training blocks motor learning relative to saline injected control
mice (Figure 1G). Normal walking (in the home cage), however, was not affected and
these mice exhibited other non-learned forelimb behaviors, such as jumping.
Collectively, this suggests a necessary role for forelimb M1 in rotarod learning.
398
Transition from hopping to walking gait facilitates motor learning best in P30-P60
mice
Markerless tracking[48] enabled tracking paw position and segmenting paw
trajectories into steps (Figure 2). This was essential to understanding changes in paw
kinematics as well as comparing steps to understand gait changes. Step length was
tracked for all four paws and showed a consistent effect across age groups. For both
forepaw and hindpaw, steps became longer as mice were more successful at remaining
on the rotarod. Further, while the trend was similar, P20 mice took shorter steps
compared to the other age groups, consistent with smaller body size. Targeting of step
endpoint relative to the end of the rod was more complex. In general, the endpoint
location was relatively stable during learning across ages, with the exception of the P60
mice. P60 animals developed a strategy of placing the forelimb step closer to the front
edge of the rotarod. The effect was smaller but present for the hind limb because the
step length became longer. This suggested that mice were able to change some of the
kinematic parameters of their steps, including targeting a specific location on the rod.
Whether this step location targeting requires somatosensory[49] or visual feedback is not
clear, although the experiments were run in the dark with infrared light.

More interesting than step kinematics is gait pattern. Overall, gait shifts from
hopping to walking as individual trials proceed. But the shift is more pronounced for
certain trials and at certain ages. Further, the shift from hopping to walking appears to
happen at different times for forelimbs and hindlimbs, suggesting that the circuitry
controlling the mode of locomotion can operate independently for each pair of limbs.
The shift from hopping to walking strategy enables animals to stay on the rod, though it
is unclear whether a split in gait pattern between forelimb and hindlimb would occur in
other more naturalistic conditions. Notably, humans are able to walk at different speeds
between the left and right leg on a split treadmill[50], but interpreting this adaption is
difficult. That the targeting of the paws closer to the front edge of the rod and the
change in gait occur in a more pronounced way for the forelimb suggests that the motor
learning may involve this limb more than the hindlimb. If the learning were simply a
response to movement of the rod, sensory feedback and balance, this might be more
likely to involve both limbs, while specific involvement of the forelimbs might imply a
more a deliberative or conscious effort that involves frontal cortex.
What about differences in the gait shift? These data can be compared across
bins within the same trial, for the same bin across trials, and across ages. Within the
first training session, forelimb step phase shifts significantly towards walking for ages
after trial 5 (Figure 3E). However, the shift for hindlimbs in the same comparisons is
strongest only for P30 and P45 (Figure 3E). This suggests that forelimb performance is
more plastic in this assay, and that the hindlimb plasticity is also enhanced at certain
ages (P30 and P45). When comparing across ages at the same trial number, older
animals perform better than P20 animals for all ages compared (Figure 3F). This shift
appears larger and has more statistical support for later trials (Figure 3F, Trials 11-15
and Trials 16-20). These plots, however, do not account perfectly for shifts in gait since
they compare steps across the entire trial (but see Figure 4D). Subdividing the steps
into bins by fraction of steps taken on a given trial, a more detailed comparison with
fewer steps per bin can be made, though this reduced the statistical power. In this
approach, the strongest effects were observed across deciles in a given trial for the
front limbs, further supporting the plasticity of the front limb.
446

Cell type specific changes in cortical inhibition for task-active networks

Previous reports demonstrated that motor learning was accompanied with reduction in inhibition. In our experiments, PV+ cells provide greater inhibition to the active cells while SOM+ cells provide similar amount of inhibition. The discrepancy could arise from our approach and experimental design that not only distinguishes the interneuron subtype but also its postsynaptic target by their activity during learning. We directly probed the changes in inhibition mediated by PV+ and SOM+ interneurons to the pyramidal cells depending on their task involvement.

Similar to our observation, learning a lever pressing task, another form of motor learning paradigm, induces transient increase in PV+ axonal boutons, their activity, and immunoreactivity of PV+ which positively correlate GABAergic activity[1,41,51]. This suggests that PV+ may regulate motor learning, particularly in early phases, by transient increases in its connection strength to excitatory neurons. This increased inhibition by PV+ could be a homeostatic mechanism to balance the increased activity in task-active pyramidal cells by increasing inhibition[36,37]. PV+, but not SOM+, interneurons modulate their inhibition depending on the relative activity of the post-synaptic pyramidal cell in V1 [36]. This homeostatic effect may be part of forming specialized subnetworks[52,53] to recruit M1 neurons in a specific spatiotemporal pattern that emerges with training[13,40,54]. In contrast to a homeostatic change, perhaps PV+ synaptic changes occur first and pyramidal neuron activity is altered following this change. Another possibility is that the PV+ interneurons may be involved in shift in the timing of pyramidal cell activity with motor learning. PV+ neurons affect the oscillation of pyramidal cells, particularly in the gamma band[55,56]. Gamma oscillation can be observed during cognitive tasks and recently with motor learning[57–59]. Timing of inhibition can modulate spike timing, potentially influencing the temporal shift in excitatory activity. During reaching task, the activity of PV+ interneurons precede the activity of putative excitatory neurons[60]. During the rotarod task, their timing of activity in relation to their post-synaptic excitatory partner may modulate the firing of excitatory cells in an activity-dependent manner. We observed a trend in increase in inhibition in P30-45 groups, but not in P20 and 60. This suggests that the circuit mechanisms of motor learning can vary with age in motor cortex. Even at postnatal ages, PV+ interneurons go through significant maturation in

both sensory and frontal areas[61–63]. Even though P60 animals are capable of achieving

similar level of motor learning as P30-45 animals, the circuit changes that support

learning may take a different form once the circuitry has matured after P60 and onward.

The synaptic changes in PV+ input are detectable on the first training day. It is

unknown if the synaptic changes by PV+ neurons are persistent or transient (thus will

be eliminated with further training as the task becomes consolidated). PV+ axonal

bouton and activity changes, for example, are transient[1,41] and with enough training,

inhibitory activity to perform the task and potential mechanism to refine the cortical

circuit may no longer be needed and move out of cortex [5,64].

Even though our study did not observe difference in inhibition mediated by SOM+

interneurons, synaptic changes in these neurons may occur with a longer time

course[1,41,65]. Additional factors that may explain our results. First, SOM+ interneuron

postsynaptic targets include other inhibitory neurons, along with pyramidal cells[33]. Thus,

changes in their activity could influence inhibition onto other inhibitory interneurons. A

second potential factor is that due to their subcellular targeting on dendrites and our

recordings done on soma, changes in SOM+ mediated inhibition on distal dendrites may

not have been detected with our approach. Dendritic recordings might elucidate whether

SOM+ inhibition also differs depending on the activity of the postsynaptic targets. A third

potential explanation is that the changes in SOM+ cells may take longer to occur

following learning. Unlike the changes in PV+ interneurons, changes in SOM+ cells are

more persistent[1,41,66]. For future studies, it would be useful to follow SOM+ connectivity

changes in later training days, while learning plateaus and animals reach expert level.

We targeted our recordings to task-active neurons in L5. This can be further

divided into sublaminae: L5A (upper) and L5B (lower). These sublaminae contain

different types of long-range projecting excitatory neurons: IntraTelencephalic (IT-) type

neurons (whose projections target ipsi- and contralateral cortex and striatum) and

Pyramidal Tract (PT-) type neurons (whose projects target ipsilateral subcortical

structures, including striatum, thalamus, superior colliculus, brainstem, and spinal cord)

[67]. IT- and PT-type neurons are the major output neurons of the motor cortex and are

critical in the preparation and execution of a movement in the motor cortex[68–71]. Their

involvement in preparation and execution of voluntary movement is different[68–72]. With

our current approach, however, we cannot distinguish between these two excitatory cell

types while maintaining the task-involvement through CaMPARI2; thus, we cannot

conclude if PV+ or SOM+ differentially inhibit these neurons in an activity-dependent

manner.

Our study has demonstrated that the aptitude of animals to perform on an

accelerating rotarod task differs depending on the age. Animals between P30-60

perform better than the younger age groups (P20) and older age groups (P90-120). In

addition, learning is accompanied with changes in the paw kinematics and gait. Animals

tend to take larger steps and shift the gait from hopping to walking, which is more

pronounced and happens earlier in the trial in forelimbs. After a single training day, PV+,

but not SOM+, induced greater inhibition in task-active postsynaptic cells than its

neighboring task-inactive cells, suggesting a role of activity dependent PV+ inhibition

during early learning. However, this effect was greatest at P30, with reduced changes in

PV+ inhibition at later ages, suggesting a shift in the circuit mechanisms of learning

related plasticity later in life.

Methods

Stereotactic injections

Animal protocols were approved by the Institutional Animal Care and Use Committee at the University of Pittsburgh. Mice of both sexes were used. Mice were held in standard housing cages with ad libitum access to food and water. Mice were housed in standard mouse caging without running wheels to minimize prior exposure to a treadmill, as running wheel exposure might shift the learning curve (Buitrago 2004). Injections were made with glass pipette (Drummond) using a custom-made positive displacement system (Narashige). In order to label and optogenetically stimulate specific interneurons, Cre driver mouse lines (PV-Cre, JAX 008069; and SOM-Cre, JAX 013044) were crossed with Ai32(lsl-ChR2-EYFP, JAX 012569). lsl-ReaChR-mCitrine mice (JAX 026294) were used in some pilot experiments. Offspring were then injected with CaMPARI2 (pAAV_hsyn_NES-his-CaMPARI2-WPRE-SV40, Addgene 101060) perinatally at postnatal day (P) 0-3. Due to the age of the animals, pups were cryoanesthetized on ice and placed in a custom clay mold to position the animal during the injection while maintaining low temperature. Stereotaxic coordinates for M1 differ from the adult and Bregma is invisible above the skin. To overcome this, confluence of sinuses, which approximately corresponds to Lambda, was used for alignment. The injection was made at 2100 µm anterior and 1000 µm lateral at two depths – 695 µm and 665 µm – 50 nl each.

Clear Skull and LED mounting

Once animals approach the developmental age for testing, the skull was cleared 3-5 days prior to rotarod training to minimize tissue damage. The mice were anesthetized with a O_2 rate of 1 L/min and 2.5%-3.0% isoflurane. During surgery, Buprenorphine (0.1mg/kg) and Ketoprofen (4 mg/kg) were administered intraperitoneally for analgesia. The scalp and periosterum of the skull were removed. The CaMPARI2 injection site was visible through the skull by illuminating the exposed skull with a Nightsea system (SFA Stereo Microscope Fluorescence Adapter) and the boundary of the injection site was delineated using a marker. The skull was scored with scalpel to enhance the adherence of the cap. A layer of cyanoacrylate (~1-2 mm) was applied and a headplate was

positioned above the cerebellum. Clear dental cement was applied over the skull before

the cyanoacrylate was fully dry. The day before the training day, clear nail polish was

applied and a high-power UV-LED (405 nm) was placed on top of the injection site

(Mouser, 604-ATDS3534UV405B). The rest of the skull cap was sealed with black

silicone. Because the headcap came unattached more readily in young mice, the

younger age group was shifted from P15 to P20 because the LED adhered better at this

age. Mice where the headcap was removed were taken out of the study.

Mice and Behavior

Animals were trained on an accelerating rotarod. The rotation accelerated uniformly

from 0 to 90 RPM in 3 minutes. Animals performed 20 trials in a session and only one

session per day. Animals were trained for 5 consecutive days, except those sacrificed

earlier for brain slice experiments. The custom-designed rotarod (Figure S1) was built

from clear plexiglass. A stepper motor (uxcell) was connected to a clear rod and was

controlled by an Arduino board. Panels dividing the lanes of the rotarod were able to

slide along the rod to adjust the width of the running lane to the size of the animal

running on the rotarod. At the bottom, a high-speed camera (FLIR Blackfly S USB3)

was mounted to record the animal during the task. Six age groups were tested:

postnatal day(P) 15-20, P30, P45, P60, P90, and P120. For the animals expressing

CaMPARI2, UV-LED light was delivered (0.1s ON, 0.9s OFF) during rotarod training.

Once the animal fell from the rod, LED stimulation was terminated. Time to fall was

measured throughout all trials. The animal's performance was recorded using high-

speed camera at 200 frames per second. The youngest age group includes P15-20.

While animals around P15 become ambulatory and can stay on the rotarod, this age

was not ideal for rotarod training. First, the skull at P15 is still not mature enough to

mount LED. Second, because animals at this age are not weaned from the parents,

oftentimes the LED mount was damaged by the parents. Therefore, P20 animals

became more suitable to test in both the rotarod task and electrophysiology.

Behavioral Analysis

Improvement measured in time to fall was calculated by taking the average of trials 1-3
and subtracting it from the average of trials 11-13 (Figure 1C). A one-way ANOVA was
performed to compare the improvement in six age groups. Tukey's multiple comparison
test was used after the ANOVA. For single comparisons (muscimol), t-test was used.
The same analysis was used to compare the baseline locomotion. To determine the
difference in weights between male and female within an age group, t-test was used.
We focused on P20-60 animals, which correspond with best ages of the animals
recorded for slice electrophysiology. The body positions of the animals were estimated
using DeepLabCut [48]. The nose, right front paw, left front paw, right hind paw, left hind
paw and base of the tail were annotated manually for training and subsequently tracked
for analysis. Estimated data points with likelihood less than 0.95 were excluded from
analysis. To analyze the changes in the position of a single paw, steps were detected
using custom software. The start and end of a step were converted to location relative
to the front (1.0) and rear (0.0) edge of the rotarod (Figure 2). The step length was
calculated by measuring the distance between the step start and end. To analyze the
changes in the gait pattern, the phase of the left and right paws (Figure 3) was
determined by comparing location and timing of one step between the corresponding
steps with the opposite paw, for example where a left hind paw (LHP) step occurred
between two right hind paw (RHP) steps. For the gait analysis, one-way ANOVA was
performed to compare the gait phases between the deciles and the trials. Tukey's
multiple comparison test was used after the ANOVA. Comparison between a decile and
trial of P20 to the corresponding decile of P30, P45 and P60 were made using t-test.
609
Muscimol inactivation
To determine whether M1 is necessary to learn the accelerating rotarod task, M1 was
pharmacologically inactivated. Muscimol (5 mg/ml, 40 nL per hemisphere) or saline was
injected into forelimb M1 bilaterally in animals P30-45 (600 μm anterior, 1500 μm
lateral, 500 and 800 μm depth, with 20 nL at each depth). 40 minutes following removal
from anesthesia, the animals were tested on the accelerating rotarod.
616
Fluorescence Analysis

Using ImageJ, red and green fluorescence were measured for the injection site and the

recorded cells. Mean fluorescence was calculated by dividing the sum of the count

multiplied by their histogram value with the sum of total counts. Photoconversion ratio

was calculated by dividing the red fluorescence mean by the green fluorescence mean.

For the pair of cells recorded, photoconversion ratio between the cells were compared

and the cells with higher photoconversion ratio was considered as task-active while the

other cells in each pair with lower photoconversion ratio was considered as task-

inactive.

Electrophysiology

After completing rotarod training, mice were anesthetized with isoflurane, and the brains

were extracted in ice-cold choline chloride based cutting solution (in mM: 25 $NaHCO_3$,

25 D-glucose, 1.25 NaH_2PO_4, 2.5 KCl, 7 $MgCl_2$, 0.5 $CaCl_2$, 3.1 Sodium Pyruvate, 11.6

Sodium Ascorbate, 110 Choline Chloride). Then the brains were sectioned coronally at

300 μm, using a vibratome in choline chloride based cutting solution. Sections were

moved to artificial cerebrospinal fluid (aCSF in mM: 127 NaCl, 25 $NaHCO_3$, 25 D-

Glucose, 2.5 KCl, 2 $CaCl_2$, 1.25 NaH_2PO_4, 1 $MgCl_2$), then transferred into aCSF at

38°C for 30 minutes then at room temperature for 30 minutes. Patch pipettes were

pulled from borosilicate glass. The pipette was filled with cesium based intracellular

solution (in mM: 35 CsF, 100 CsCl, 10 EGTA, and 10 HEPES). Recordings were

performed at room temperature (22 °C), amplified by an Axopatch 700B amplifier

(Molecular devices) and acquired at 10KHz in WaveSurfer, open-source software,

(https://wavesurfer.janelia.org). Neurons were selected across the thickness of L5. The

cell was voltage clamped at +40 mV. To elicit IPSCs, blue LED stimulation (470 nm)

was delivered for 1, 2, and 5 ms with 20 second intervals in between each sweep for

recovery. The cycle was repeated three times.

Electrophysiology Analysis

Electrophysiology data were low-pass filtered (1kHz). The three traces were averaged.

IPSCs were measured when they exceeded a threshold of greater than 5x standard

deviation from the baseline. Cells with membrane capacitance (Cm) below 70 pF were

excluded from analysis. Neurons expressing CaMPARI2 (red, task-active cells; green,

task-inactive cells) were targeted for recordings. To compare neurons within a slice,

nearby task-active (red) and task-inactive cells in the same high-power (60x field of

view) were recorded. The peak IPSC amplitude of task-active and task-inactive cells

were compared using Wilcoxon rank sum test. IPSCs of task-active cells were

normalized to their partner inactive cells and the ratio and the ratio was analyzed using

Wilcoxon rank sum test.

Figure Legends

Figure 1 | Rotarod skill acquisition plateaus around P30-P60 on training day 1.
(A) Schematic of rotarod task with high speed videography and barriers to create the
running lane during a trial. (B) Video frame of mouse during running. Paw positions
(LFP, left front paw; LHP, left hind paw; RFP, right front paw; RHP, right hind paw) and
the nose and tail base are tracked using DeepLabCut. (C) Learning curve during 20
trials on training day 1 (TD1). Six colors represent the six ages tested (P15-20, P30,
P45, P60, P90, and P120). Gray bars represent time windows used for early and late
performance to assess TD1 improvement. (D) Improvement (in seconds) in time to fall
during TD1 for six age groups. Mean+/- SEM. N=63, 92, 60 63, 27, 12. (E) Improvement
(in seconds) in time to fall during TD1 for six age groups, subdivided by male (blue) and
female (red). Mean+/- SEM.(F) Improvement (in seconds) in time to fall during five
training days. Gray vertical lines divide the days. Mean+/- SEM. (G) The effect of M1
inactivation by bilateral muscimol injection on rotarod skill acquisition compared to
saline injected controls. N=5 (control); N=9 (muscimol). Mean+/- SEM. Significance: *,
$p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Figure 2 | Tracking step kinematics during rotarod training.
(A) Schematic of rotarod task with video frame of tracked mouse. (B) Cartoon of tracked
image frame illustrating location of the mouse paws (LFP, left front paw; LHP, left hind
paw; RFP, right front paw; RHP, right hind paw) and the nose and tail base using
DeepLabCut. The rotarod and direction of motion are indicated, with the front and rear
ends of the rod serving as 0.0 and 1.0 in normalized coordinates. At bottom,
coordinates for the step start, step end, and step length are plotted to illustrate how they
are determined. (C) Plot of the x-position of a single paw (LHP, left) or all paws (right)
during a trial. Time given in frames (converted to seconds with example timebar). (D)
Expansion of paw tracker showing step detection. Steps (red) are automatically
detected (see methods). Step start (green), step end (black), and step length are then
calculated. (E) Step length versus time to fall for the left front paw, plotted for all ages
combined (red) and individual ages (20 and above). Data are fit to a line and the linear

fit compared at right. Summary plots include step length versus time to fall for the right

front paw, left hind paw, and right hind paw (far right). (F) Step end point presented

similarly to step length.

Figure 3 | Gait pattern changes with age and training.

(A) Schematic of rotarod task with video frame of tracked mouse. (B) Cartoon of tracked

image frame illustrating location of the mouse paws (LFP, left front paw; LHP, left hind

paw; RFP, right front paw; RHP, right hind paw) and the nose and tail base using

DeepLabCut. The rotarod and direction of motion are indicated, with the front and rear

ends of the rod serving as 0.0 and 1.0 in normalized coordinates. At bottom,

coordinates for the step start, step end, and step length are plotted to illustrate how they

are determined. (C) Plot of the x-position of a single paw (LHP, left) or all paws (right)

during a trial. Time given in frames (converted to seconds with example timebar). (D)

Expansion of paw tracker showing step detection. Steps (red) are automatically

detected (see methods). Step start (green), step end (black), and step length are then

calculated. (E) Step length versus time to fall for the left front paw, plotted for all ages

combined (red) and individual ages (20 and above). Data are fit to a line and the linear

fit compared at right. Summary plots include step length versus time to fall for the right

front paw, left hind paw, and right hind paw (far right). (F) Step end point presented

similarly to step length. Significance: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Figure 4 | P-values for the gait analysis.

(A) Due to variations in the number of steps, the number of steps in a trial were divided

into deciles to facilitate the comparison of average step phases across trials and age

groups. The average step phase value for each age for front and hind limbs is plotted.

(B) The P-value plot of deciles compares the first decile of trials 1-20 to rest of the

deciles (2-9) for each trial. The top row compares front paws at four ages and the

bottom row compares hind paws. Non-significant p-values are purple; p-values < 0.05

are as shown in the lookup table. (C) The P-value plot comparing across trials

compares each decile from trials 2-20 with the corresponding decile from trial 1. The top

row compares front paws at four ages and the bottom row compares hind paws. For

instance, the front paw in P20 Decile 3 from trial 3 is compared to Decile 3 from trial 1.

Non-significant p-values are purple; p-values < 0.05 are as shown in the lookup table.

(D) The P-value plot of deciles compared across ages assess P20 versus P30, P45,

and P60 groups for 20 trials. Each individual graph represents the comparison between

the four age groups for a given trial. The X-axis denotes deciles, with a specific trial's

decile being compared to the corresponding decile of the P20 group. For example, the

front paw in P30 Decile 9 is compared to P20 Decile 9 in the trial 1 graph, and if

significantly different, it is indicated by a bump in the y-axis. Significance: *; p<0.05.

;p<0.01, *;p<0.001.

Figure 5 | Schematic of CaMPARI2 manipulation.

(A) AAV expresses CaMPARI2 in M1 in PVcre+/-;Ai32+/- or SOMcre+/-;Ai32+/- mice.

(B,C) CaMPARI2-expressing cells are photoconverted to red by a skull mounted UV

LED. (D) Example images of a slice after photoconversion during rotarod. Green

arrowheads mark green neurons not photoconverted (task-inactive) to red; red

arrowheads mark red neurons (task-active). (E) Example traces from task-inactive

(green) and task-active (red) neurons after LED stimulation of PV+ cells. (F)

Photoconversion ratio is calculated by dividing red fluorescence over green

fluorescence. Animals showed greater photoconversion ratio after rotarod training (grey)

compared to the control animals that were permitted to move freely in the rotarod

chamber floor (white). Significance: *; p<0.05.

Figure 6 l IPSCs recorded in task-active (RFP+) vs. task-inactive cells (GFP+) after

stimulating PV+ interneurons

(A-E) Scatter plot of IPSC responses in pairs of task-active (red) and task-inactive

(green) cells in P20 (A), P30 (B), P45 (C), P60(D) and across all age groups (E). Solid

line is the geometric mean of the pair responses. Dotted line is the line of equality (x=y).

(F) Slopes of geometric mean across all ages plotted together. (G-K) Ratio of responses

of task-active cells to their partner task-inactive cells after PV+ stimulation in P20 (G),

P30 (H), P45 (I), P60 (J) and across all age groups (K). Significance: *, p<0.05; **,

p<0.01; ***, p<0.001. (Also indicated with red arrows in (A-E). (L) Schematic of the

genetic strategy for testing PV output to M1 pyramidal neurons

Figure 7 I IPSCs recorded in task-active (RFP+) vs. task-inactive cells (GFP+) after

stimulating SOM+ interneurons

(A-E) Scatter plot of IPSC responses in pairs of task-active (red) and task-inactive

(green) cells in P20 (A), P30 (B), P45 (C), P60(D) and across all age groups (E). Solid

line is the geometric mean of the pair responses. Dotted line is the line of equality (x=y).

(F) Schematic of the genetic strategy for testing SOM output to M1 pyramidal neurons.

(G-K) Ratio of responses of task-active cells to their partner task-inactive cells after

SOM+ stimulation in P20 (G), P30 (H), P45 (I), P60 (J) and across all age groups (K).

Supplementary Information

Figure S1 | Rotarod design

(A) Image of a clear plexiglass rotarod with high speed video. Steeper motor and

camera run by computer adjacent to the operator. Multiple lanes are possible (divided

by moveable lane dividers) to simultaneously run multiple mice (two lanes in the camera

field of view). Infrared (IR) LEDs used for illumination during behavioral training. (B)

Overhead view of the apparatus. Lane dividers opened further than during trials to

illustrate capacity for sliding.

Figure S2 | Developmental changes in mobility and weight do not account for rotarod performance

(A) Measurement of home cage walking (beam breaks) used to assess motility across developmental ages. (B) Home cage motility quantified by sex across ages. Red/blue is used to designate sex (red, female; blue, male), with brightness used to indicate age. The same color scheme is used in D-E. (C-D) Changes in weight across developmental ages, quantified for all mice or by sex. Male mice weigh more than female mice at ages older than P30. (E) Plot comparing the first training day (TD1) time-to-fall improvement versus weight. Red/blue color code is used to indicate age and sex. Linear fit is relatively poor (R^2=-0.04). Significance: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Figure S3 | Developmental changes in rotarod performance plotted for male and

female

(A) Measurement of training day 1 (TD1) performance, separated by male (blue) and

female (red). (B) Summary of time-to-fall improvement, plotted as in Figure 1. (C)

Measurement of rotarod performance across all training days, separated by male (blue)

and female (red). Significance: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Figure S4 | Step kinematics across ages and limbs

(A) Step length versus time to fall for the left front paw, plotted for all ages combined (red) and individual ages (20 and above). Data are fit to a line and the linear fit compared at right. (B-D) Step length versus time to fall for the right front paw (B), left hind paw (C), and right hind paw (D). (E-H) Step end point presented similarly to step length.

Figure S5 I Additional factors considered for the pair recordings

To determine whether other factors could potentially affect the pair recordings, we

replotted Figure 5 and 6, using the color of the data representing a third variable. The

degree of photoconversion was assessed by plotting by the red/green fluorescence ratio

for the recordings from PV+ (A) and SOM+ (D) stimulation. Higher the ratio is, the

greater the difference in photoconversion between the pair. In addition, four different

improvement magnitudes were plotted by four different colors for the recordings from

PV+ (B) and SOM+ (E) stimulation. The geometric mean of the respective improvement

groups was plotted in solid line of the same color scheme. Lastly, male and female were

plotted in red and blue for the recordings from PV+ (C) and SOM+ (F) stimulation. The

geometric mean of the respective sex was plotted in solid line of the same color

scheme.

References

1. Chen, S. X., Kim, A. N., Peters, A. J. & Komiyama, T. Subtype-specific plasticity of
inhibitory circuits in motor cortex during motor learning. *Nat Neurosci* **18**, 1109–15
(2015).

2. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews*
Neuroscience vol. 6 877–888 Preprint at https://doi.org/10.1038/nrn1787 (2005).

3. Erzurumlu, R. S. & Gaspar, P. Development and critical period plasticity of the
barrel cortex. *European Journal of Neuroscience* (2012) doi:10.1111/j.1460-
9568.2012.08075.x.

4. Hooks, B. M. & Chen, C. Circuitry Underlying Experience-Dependent Plasticity in
the Mouse Visual System. *Neuron* (2020) doi:10.1016/j.neuron.2020.08.004.

5. Kawai, R. *et al.* Motor Cortex Is Required for Learning but Not for Executing a
Motor Skill. *Neuron* **86**, 800–812 (2015).

6. Sauerbrei, B. A. *et al.* Cortical pattern generation during dexterous movement is
input-driven. *Nature* **577**, 386–391 (2020).

7. Papale, A. E. & Hooks, B. M. Circuit Changes in Motor Cortex During Motor Skill
Learning. *Neuroscience* **368**, 283–297 (2018).

8. Knudsen, E. I. & Knudsen, P. F. Sensitive and critical periods for visual calibration
of sound localization by barn owls. *Journal of Neuroscience* **10**, 222–232 (1990).

9. Gordon, J. A. & Stryker, M. P. Experience-dependent plasticity of binocular
responses in the primary visual cortex of the mouse. *J Neurosci* **16**, 3274–3286
(1996).

10. Van Der Loos, H. & Woolsey, T. A. Somatosensory cortex: structural alterations
following early injury to sense organs. *Science* **179**, 395–398 (1973).

11. Costa, R. M., Cohen, D. & Nicolelis, M. A. L. Differential Corticostriatal Plasticity
during Fast and Slow Motor Skill Learning in Mice. *Current Biology* **14**, 1124–1134
(2004).

12. Dombeck, D. A., Graziano, M. S. & Tank, D. W. Functional Clustering of Neurons
in Motor Cortex Determined by Cellular Resolution Imaging in Awake Behaving
Mice. *Journal of Neuroscience* **29**, 13751–13760 (2009).

13. Peters, A. J., Chen, S. X. & Komiyama, T. Emergence of reproducible
spatiotemporal activity during motor learning. *Nature* **510**, 263–267 (2014).

14.   Hira, R. *et al.* Spatiotemporal dynamics of functional clusters of neurons in the
844   mouse motor cortex during a voluntary movement. *J Neurosci* **33**, 1377–1390
845   (2013).

846   15.   Yang, G., Pan, F. & Gan, W.-B. Stably maintained dendritic spines are associated
847   with lifelong memories. *Nature* **462**, 920–4 (2009).

848   16.   Yang, G. *et al.* Sleep promotes branch-specific formation of dendritic spines after
849   learning. *Science* **344**, 1173–8 (2014).

850   17.   Weiler, N., Wood, L., Yu, J., Solla, S. A. & Shepherd, G. M. G. Top-down laminar
851   organization of the excitatory network in motor cortex. *Nat Neurosci* **11**, 360–366
852   (2008).

853   18.   Mao, T. *et al.* Long-Range Neuronal Circuits Underlying the Interaction between
854   Sensory and Motor Cortex. *Neuron* **72**, 111–123 (2011).

855   19.   Hooks, B. M. *et al.* Organization of cortical and thalamic input to pyramidal neurons
856   in mouse motor cortex. *Journal of Neuroscience* (2013)
857   doi:10.1523/JNEUROSCI.4338-12.2013.

858   20.   Jacobs, K. M. & Donoghue, J. P. Reshaping the cortical motor map by unmasking
859   latent intracortical connections. *Science* **251**, 944–947 (1991).

860   21.   Rioult-Pedotti, M. S., Friedman, D. & Donoghue, J. P. Learning-induced LTP in
861   neocortex. *Science* **290**, 533–536 (2000).

862   22.   Kaneko, T., Caria, M. A. & Asanuma, H. Information processing within the motor
863   cortex. II. Intracortical connections between neurons receiving somatosensory
864   cortical input and motor output neurons of the cortex. *J Comp Neurol* **345**, 172–
865   184 (1994).

866   23.   Southwell, D. G., Froemke, R. C., Alvarez-Buylla, A., Stryker, M. P. & Gandhi, S.
867   P. Cortical plasticity induced by inhibitory neuron transplantation. *Science (1979)*
868   (2010) doi:10.1126/science.1183962.

869   24.   Tang, Y., Stryker, M. P., Alvarez-Buylla, A. & Espinosa, J. S. Cortical plasticity
870   induced by transplantation of embryonic somatostatin or parvalbumin interneurons.
871   *Proc Natl Acad Sci U S A* (2014) doi:10.1073/pnas.1421844112.

872   25.   Bachtiar, V. & Stagg, C. J. The role of inhibition in human motor cortical plasticity.
873   *Neuroscience* **278**, 93–104 (2014).

874   26.   Kolasinski, J. *et al.* The dynamics of cortical GABA in human motor learning. *J*
875   *Physiol* **597**, 271–282 (2019).

876   27.   Blicher, J. U. *et al.* GABA levels are decreased after stroke and GABA changes
877   during rehabilitation correlate with motor improvement. *Neurorehabil Neural Repair*
878   **29**, 278–286 (2015).

879   28.   Alia, C. *et al.* Reducing GABA A-mediated inhibition improves forelimb motor
880   function after focal cortical stroke in mice. *Sci Rep* (2016) doi:10.1038/srep37823.

881   29.   Kida, H. *et al.* Motor training promotes both synaptic and intrinsic plasticity of layer
882   V pyramidal neurons in the primary motor cortex. *J Physiol* **601**, 335–353 (2023).

883   30.   Kida, H. *et al.* Motor Training Promotes Both Synaptic and Intrinsic Plasticity of
884   Layer II/III Pyramidal Neurons in the Primary Motor Cortex. *Cerebral Cortex (New*
885   *York, NY)* **26**, 3494 (2016).

886   31.   Rudy, B., Fishell, G., Lee, S. & Hjerling-Leffler, J. Three groups of interneurons
887   account for nearly 100% of neocortical GABAergic neurons. *Dev Neurobiol* **71**, 45–
888   61 (2011).

889   32.   Tremblay, R., Lee, S. & Rudy, B. GABAergic Interneurons in the Neocortex: From
890   Cellular Properties to Circuits. *Neuron* **91**, 260–92 (2016).

891   33.   Pfeffer, C. K., Xue, M., He, M., Huang, Z. J. & Scanziani, M. Inhibition of inhibition
892   in visual cortex: the logic of connections between molecularly distinct interneurons.
893   *Nat Neurosci* **16**, 1068–1076 (2013).

894   34.   Kawaguchi, Y. & Kubota, Y. Neurochemical features and synaptic connections of
895   large physiologically-identified GABAergic cells in the rat frontal cortex.
896   *Neuroscience* **85**, 677–701 (1998).

897   35.   Somogyi, P., Tamás, G., Lujan, R. & Buhl, E. H. Salient features of synaptic
898   organisation in the cerebral cortex. *Brain Res Rev* **26**, 113–135 (1998).

899   36.   Xue, M., Atallah, B. V. & Scanziani, M. Equalizing excitation–inhibition ratios
900   across visual cortical neurons. *Nature* **511**, 596–600 (2014).

901   37.   Kuhlman, S. J. *et al.* A disinhibitory microcircuit initiates critical-period plasticity in
902   the visual cortex. *Nature* **501**, 543–546 (2013).

903   38.   Friel, K., Chakrabarty, S., Kuo, H. C. & Martin, J. Using Motor Behavior during an
904   Early Critical Period to Restore Skilled Limb Movement after Damage to the
905   Corticospinal System during Development. *Journal of Neuroscience* **32**, 9265–
906   9276 (2012).

907   39.   Walton, K. D., Lieberman, D., Llinás, A., Begin, M. & Llinás, R. R. Identification of a
908   critical period for motor development in neonatal rats. *Neuroscience* **51**, 763–767
909   (1992).

910   40.   Adler, A., Zhao, R., Shin, M. E., Yasuda, R. & Gan, W.-B. Somatostatin-
911   Expressing Interneurons Enable and Maintain Learning-Dependent Sequential
912   Activation of Pyramidal Neurons. *Neuron* **102**, 202-216.e7 (2019).

913   41.   Ren, C. *et al.* Global and subtype-specific modulation of cortical inhibitory neurons
914   regulated by acetylcholine during motor learning. *Neuron* **110**, 2334-2350.e8
915   (2022).

916   42.   Moeyaert, B. *et al.* Improved methods for marking active neuron populations. *Nat*
917   *Commun* **9**, 4440 (2018).

918   43.   Grillner, S. & Wallén, P. Central pattern generators for locomotion, with special
919   reference to vertebrates. *Annu Rev Neurosci* **8**, 233–261 (1985).

920   44.   Buitrago, M. M., Schulz, J. B., Dichgans, J. & Luft, A. R. Short and long-term motor
921   skill learning in an accelerated rotarod training paradigm. *Neurobiol Learn Mem* **81**,
922   211–216 (2004).

923   45.   Greenwood, B. N. & Fleshner, M. Voluntary Wheel Running: A Useful Rodent
924   Model for Investigating the Mechanisms of Stress Robustness and Neural Circuits
925   of Exercise Motivation. *Curr Opin Behav Sci* **28**, 78 (2019).

926   46.   Sherwin, C. M. Voluntary wheel running: a review and novel interpretation. *Anim*
927   *Behav* **56**, 11–27 (1998).

928   47.   Jain, A., Huang, G. Z. & Woolley, C. S. Latent Sex Differences in Molecular
929   Signaling That Underlies Excitatory Synaptic Potentiation in the Hippocampus.
930   *Journal of Neuroscience* **39**, 1552–1565 (2019).

931   48.   Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body
932   parts with deep learning. *Nat Neurosci* **21**, 1281–1289 (2018).

933   49.   Huang, Y., Zhang, X. & Li, W. Involvement of primary somatosensory cortex in
934   motor learning and task execution. *Neurosci Lett* **828**, 137753 (2024).

935   50.   Iturralde, P. A. & Torres-Oviedo, G. Corrective Muscle Activity Reveals Subject-
936   Specific Sensorimotor Recalibration. *eNeuro* **6**, (2019).

937   51.   Donato, F., Rompani, S. B. & Caroni, P. Parvalbumin-expressing basket-cell
938   network plasticity induced by experience regulates adult learning. *Nature* (2013)
939   doi:10.1038/nature12866.

940   52.   Goz, R. U. & Hooks, B. M. Correlated Somatosensory Input in
941   Parvalbumin/Pyramidal Cells in Mouse Motor Cortex. *eNeuro* **10**, 1–18 (2023).

942   53.   Znamenskiy, P. *et al.* Functional specificity of recurrent inhibition in visual cortex.
943   (2024) doi:10.1016/j.neuron.2023.12.013.

944   54.   Isomura, Y., Harukuni, R., Takekawa, T., Aizawa, H. & Fukai, T. Microcircuitry
945   coordination of cortical motor information in self-initiation of voluntary movements.
946   *Nature Neuroscience 2009 12:12* **12**, 1586–1593 (2009).

947   55.   Sohal, V. S., Zhang, F., Yizhar, O. & Deisseroth, K. Parvalbumin neurons and
948   gamma rhythms enhance cortical circuit performance. *Nature* **459**, 698 (2009).

949   56.   Cardin, J. A. *et al.* Driving fast-spiking cells induces gamma rhythm and controls
950   sensory responses. *Nature* **459**, 663 (2009).

951   57.   Howard, M. W. *et al.* Gamma oscillations correlate with working memory load in
952   humans. *Cereb Cortex* **13**, 1369–1374 (2003).

953   58.   Lewis, D. A., Hashimoto, T. & Volk, D. W. Cortical inhibitory neurons and
954   schizophrenia. *Nature Reviews Neuroscience 2005 6:4* **6**, 312–324 (2005).

955   59.   Nowak, M. *et al.* Driving Human Motor Cortical Oscillations Leads to Behaviorally
956   Relevant Changes in Local GABAA Inhibition: A tACS-TMS Study. *Journal of*
957   *Neuroscience* **37**, 4481–4492 (2017).

958   60.   Estebanez, L., Hoffmann, D., Voigt, B. C. & Poulet, J. F. A. Parvalbumin-
959   Expressing GABAergic Neurons in Primary Motor Cortex Signal Reaching. *Cell*
960   *Rep* **20**, 308–318 (2017).

961   61.   Chattopadhyaya, B. *et al.* Experience and Activity-Dependent Maturation of
962   Perisomatic GABAergic Innervation in Primary Visual Cortex during a Postnatal
963   Critical Period. *Journal of Neuroscience* **24**, 9598–9611 (2004).

964   62.   Miyamae, T., Chen, K., Lewis, D. A. & Gonzalez-Burgos, G. Distinct Physiological
965   Maturation of Parvalbumin-Positive Neuron Subtypes in Mouse Prefrontal Cortex.
966   *The Journal of Neuroscience* **37**, 4883 (2017).

967   63.   Canetta, S. E. *et al.* Mature parvalbumin interneuron function in prefrontal cortex
968   requires activity during a postnatal sensitive period. *Elife* **11**, (2022).

969   64.   Hwang, E. J., Dahlen, P. J. E., Mukundan, P. M. & Komiyama, T. Disengagement
970   of Motor Cortex during Long-Term Learning Tracks the Performance Level of
971   Learned Movements. *Journal of Neuroscience* **41**, 7029–7047 (2021).

972   65.   Yang, J. *et al.* Functionally distinct NPAS4-expressing somatostatin interneuron
973   ensembles critical for motor skill learning. *Neuron* **110**, 3339-3355.e8 (2022).

974   66.   Yang, J. *et al.* Functionally distinct NPAS4-expressing somatostatin interneuron
975   ensembles critical for motor skill learning Highlights d Motor learning selectively
976   induces NPAS4 expression in SST-INs in M1 d Cell-type specific KO of Npas4 in
977   SST-INs of M1 in mice impairs motor learning d NPAS4 + SST-INs exhibited lower

978   activity during task-related movements d NPAS4 re-expression maintains the

979   functionally distinct SST-INs throughout learning.

980   doi:10.1016/j.neuron.2022.08.018.

981   67.   Shepherd, G. M. G. Corticostriatal connectivity and its role in disease. *Nature*

982   *Reviews Neuroscience* vol. 14 278–291 Preprint at https://doi.org/10.1038/nrn3469

983   (2013).

984   68.   Park, J. *et al.* Motor cortical output for skilled forelimb movement is selectively

985   distributed across projection neuron classes. *Sci Adv* **8**, (2022).

986   69.   Currie, S. P. *et al.* Movement-specific signaling is differentially distributed across

987   motor cortex layer 5 projection neuron classes. (2022)

988   doi:10.1016/j.celrep.2022.110801.

989   70.   Li, N., Chen, T. W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit

990   for motor planning and movement. *Nature* **519**, 51–56 (2015).

991   71.   Soma, S. *et al.* Systems/Circuits Distinct Laterality in Forelimb-Movement

992   Representations of Rat Primary and Secondary Motor Cortical Neurons with

993   Intratelencephalic and Pyramidal Tract Projections. (2017)

994   doi:10.1523/JNEUROSCI.1188-17.2017.

995   72.   Shinotsuka, T., Tanaka, Y. R., Terada, S. I., Hatano, N. & Matsuzaki, M. Layer 5

996   Intratelencephalic Neurons in the Motor Cortex Stably Encode Skilled Movement. *J*

997   *Neurosci* **43**, 7130–7148 (2023).

998