Effects of *Bacillus thuringiensis* subsp. *kurstaki*

application on non-target nocturnal macromoth

biodiversity in the eastern boreal forest, Canada.

Jodi O. Young[1], Joseph J. Bowden[1,2], Eric R. D. Moise[1,2],

Robert Scott[1], B. Christian Schmidt[3]

1. School of Science and the Environment, Memorial

University of Newfoundland, Corner Brook, NL, Canada.

2. Natural Resources Canada, Canadian Forest Service, Corner

Brook, NL, Canada.

3. Agriculture and Agri-Food Canada, Canadian National

Collection of Insects, Arachnids and Nematodes, Ottawa, ON,

Canada

Corresponding author: Jodi O. Young, (709)660-6613,

joyoung@mun.ca, 20 University Dr, Corner Brook, NL A2H

5G5

Running Title: *Btk* & Nocturnal Macromoth Biodiversity

Abstract:

1. Insect biodiversity is crucial for resilient ecosystems,

supporting essential services. Moths play significant roles as

herbivores, pollinators, decomposers, and as prey for birds,

bats, and predatory invertebrates, emphasizing their

conservation importance. However, large-scale studies on the

non-target effects of *Bacillus thuringiensis* subs. *kurstaki* (Btk)

on moth communities are limited, particularly in northern

forests where diversity is lower. Btk is commonly used to

control pest insect populations, with forest managers in eastern

Canada applying it to manage eastern spruce budworm.

2. We established a replicated two-year study in the North

American Boreal Forest of western Newfoundland, Canada,

using paired Btk-treated and control (untreated) sites.

4. No significant differences in total abundance or richness

were found between treated and control sites. In 2021, Hill

numbers only differed between our northern treatment and

control sites, which may reflect those stands having already

received multiple years of treatment compared to just one year

in the southern sites. In 2022, control sites showed higher

diversity (Shannon and Simpson diversity metrics) compared to

treated sites, extending to all locations.

5. Multiple years of Btk treatment led to shifts in community

composition and the relative abundance of some common

species, without affecting total richness or abundance. Species

responses varied, likely due to Btk sensitivity, application

timing, and differences in phenology and voltinism, making it

difficult to generalize the effects of Btk on moth communities.

Keywords: Lepidoptera, forest pests, forest management,

biological control, Spruce Budworm Early Intervention

Strategy

Introduction

Insect biodiversity is vital to all ecosystem services such

as supporting services (e.g., decomposers), provisioning

services (e.g., pollinators), regulation services (e.g., serving as

a food source), and cultural services (e.g., ecotourism)

(Sanmartín-Villar & Cordero-Rivera, 2024). The order

Lepidoptera (moths and butterflies) is highly diverse and well

represented in boreal forests. Investigating the biodiversity

patterns and dynamics of Lepidoptera species offers crucial

insights into the complexity of ecosystem health and resilience.

While butterflies are more well known, moths are 10 times

more taxonomically diverse (Wagner *et al.*, 2021). They

provide numerous important ecosystem services such as

defoliation, pollination, nutrient cycling, plant decomposition,

and prey for other animals (Holmes, Schultz & Nothnagle,

1979; Hammond & Miller, 1998; Wagner, 2005). Nocturnal

moths have been shown to be more efficient pollinators of

certain plants, such as bramble (*Rubus fruticosus* agg.)

(Anderson, Rotheray & Mathews, 2023). Moths also differ

substantially in their flight phenology, meaning that early

season flyers are likely to pollinate a different cohort of plants

than those that emerge later. Moths are an important food

source for many animals such as predatory invertebrates, birds

and bats (Wagner *et al.*, 2021) Still, relatively little is known

about the status of this largely nocturnal insect group, and to

conserve the species and the important ecosystem services they

provide we must have data on the current state of moth

biodiversity and the potential impacts of forest management

techniques.

The Tortricidae is a family of micromoths that contains

numerous species of economic and ecological importance in

forests. One of them, the spruce budworm (*Choristoneura*

fumiferana, Clem., henceforth SBW), has a particularly large

impact on fir and spruce forests across Canada and the United

States (Maclean & Ostaff, 1989; Chang *et al.*, 2012). SBW

outbreaks occur cyclically approximately every 30-40 years

and often last for many years (Blais, 1983). During these

outbreaks, stands are heavily defoliated which leads to reduced

growth and eventual tree mortality (MacLean, 1984). Within

the 20[th] century, there were three major outbreaks in Canada,

resulting in tens of millions of hectares of damage (Blais,

1983). These outbreaks have both economic (e.g., timber

losses, Chang *et al.*, 2012) and ecological consequences (e.g.,

changes to wildlife habitat and carbon sequestering ability;

Dymond *et al.*, 2010; Chagnon, Bouchard and Pothier, 2022),

therefore appropriate management strategies are imperative for

forest health.

Bacillus thuringiensis subsp. *kurstaki* (*Btk*) has been

widely used in Canada to manage this irruptive pest since the

mid-1980s (van Frankenhuyzen, 2000). *Btk* is a naturally

occurring soil bacterium that produces proteinaceous crystals,

endowing it with insecticidal properties(Lambert & Peferoen,

1992). *Bacillus thuringiensis* subsp. *kurstaki* is a bacteria

comprised of spores that produce a crystalline toxin specifically

harmful to Lepidoptera (Schünemann, Knaak & Fiuza, 2014).

Once consumed by larval stages, the toxins contained within

the crystals are released in the gut of the caterpillar and bind

with specific receptors in the lining, ultimately resulting in their

death by cell lysis (Knowles & Ellar, 1987; Lambert &

Peferoen, 1992). *Btk* is applied aerially onto forest stands

intending to protect the current year's foliage and/or reduce the

outbreak spread and magnitude (Fuentealba et al., 2019; Johns

et al., 2019). Previous studies have shown that *Btk* is an

effective insecticide for lepidopterous pests (Bauce *et al.*, 2004;

Sanahuja *et al.*, 2011). With *Btk* being Lepidoptera specific, it

can negatively impact non-target moth species.

Previous studies have shown an overall negative effect

on abundance and richness of non-target species (Boulton et al.,

2002; Boulton, 2004; Miller, 1990; Sample et al., 1996).

However, other research has shown that the impact on non-

target species is not linear and taxa are not equally affected,

with differences in phenology and reproductive traits playing a

role (Boulton et al., 2002; Boulton & Otvos, 2004; Leza et al.,

2021; Manderino et al., 2014; Miller, 1990, 2000; Peacock et

al., 1998; Strazanac et al., 2005; Wagner et al., 1996).

Moreover, there is evidence to suggest that *Btk* may have a

positive effect on some genera due to decreased competition in

sprayed areas (Sample *et al.*, 1996; Manderino, Crist &

Haynes, 2014). The variation in susceptibility, in combination

with the rarity of large-scale field experiments, makes it

difficult to predict the non-target effects of *Btk* application *in*

vivo and highlights the knowledge gaps that still exist.

The objective of this study was to quantify *Btk* effects

on non-target macromoth species in the context of a SBW

management program. Forest management practices that

negatively impact lepidopteran communities can have

significant trophic implications across all levels of the food

chain, by either decreasing prey availability to higher trophic

levels or modifying the richness of consumers of primary

production (Summerville, 2011), all of which pose ecosystem

consequences. As one of the most functionally significant taxa

in forest ecosystems, nocturnal macromoths were selected for

this study due to their diversity, abundance, and ease of capture

and identification (Holmes, Schultz & Nothnagle, 1979;

Burford, Lacki & Covell, 1999; Young, 2005; Alison *et al.*,

2022).

Using a replicated 'treatment vs. control' design, we

made use of an ongoing "Early Intervention Strategy" (Johns et

al., 2019) established to manage SBW in western

Newfoundland, Canada, to investigate how the aerial

application of *Btk* in forests affects the community structure

(i.e., diversity, abundance, and composition) of nocturnal

macromoths.

Materials and Methods

Study Location

We conducted this study over two years (2021 and

2022) on the west coast of Newfoundland, Canada. The study

sites were divided into two locations: north and south of Gros

Morne National Park, Newfoundland and Labrador, Canada

(Figure 1). The northern sites were located in the Northern

Peninsula Forest ecoregion (Protected Areas Association of

Newfoundland and Labrador, 2008a). These sites experience

cool summers and prolonged winters, characterized by some of

the coldest temperatures on the island. The annual rainfall in

these regions ranges between 1300-1500mm. Climate modeling

using the CanESM2 model revealed average temperatures for

June, July, and August to be 11.2°C, 16.8°C, and 17°C,

respectively (Chylek *et al*., 2011). The forested areas in which

we sampled are dominated by balsam fir (*Abies balsamea*, (L.)

Mill).

The southern sites were located in the Western

Newfoundland Forest ecoregion (Protected Areas Association

of Newfoundland and Labrador, 2008b). These sites experience

warm summers and cold winters with an annual rainfall of

1200mm. Based on CanESM2 climate modelling, the average

temperatures for June, July, and August are 12°C, 17.1°C, and

16.9°C, respectively. Consistent with the northern sites, all

collections took place in balsam fir dominated boreal forest

stands.

We sampled a total of 13 sites. In 2021, we collected

moths from 12 sites in a paired design of treatment and control

sites. In the south, there were three sites that had been sprayed

with *Btk* for one year (2021) and three control sites that had not

been sprayed. In the north, there were three sites that had been

sprayed for two consecutive years (2020 and 2021) and three

control sites that were not sprayed. In 2022, we sampled most

of these sites again, with an extra consecutive year of *Btk*

treatment: in the south, the treatment sites had been sprayed for

two consecutive years (2021 and 2022) and in the north, the

treatment sites had been sprayed for three consecutive years

(2020, 2021, and 2022). The control sites remained the same. In

2022, one northern treatment site had to be shifted to a nearby

stand and replaced as it was no longer included in the 2022

spray program, and we wanted to maintain consistent spray

parameters across all sites. We chose a new site approximately

4.5 kilometers away from the previous site. As with all northern

treatment sites, this one had been sprayed for three consecutive

years (2020, 2021, and 2022).

Spray Program

204   The Spruce Budworm Early Intervention Strategy is a

management strategy adopted by the Healthy Forest

Partnership, a group of federal, provincial, academic, and

industry stakeholders. The goal of this program is to identify

potential SBW outbreak "hotspot" locations by employing

landscape-level monitoring, then proactively curtailing the

growth of SBW populations with application of *Btk* before

insects surpassing outbreak thresholds. The province of

Newfoundland and Labrador began their treatment in 2020,

actioned by their Department of Fisheries, Forestry, and

Agriculture (Government of Newfoundland and Labrador,

2020).

216   *Btk* was used in 2020, 2021, and 2022 to protect 35 583,

138 950, and 142 474 hectares of forested land, respectively

(Government of Newfoundland and Labrador, 2020;

https://www.gov.nl.ca/ffa/programs-and-funding/forestry-

programs-and-funding/idc/controlprogram/). This largely took

place in the Great Northern Peninsula using one or more

applications at a concentration of 30 billion international units

per liter (BIU/L) at a standard rate of 1.5 liters per hectare

(L/ha) for a total of 45 BIU/ha. All aerial applications were

applied when weather conditions and larval development were

optimal for *Btk* treatment. These optimal weather conditions

were considered when: wind speeds were between 2 and

10km/h, air temperatures were below 25°C, relative humidity

was above 50%, and it was not raining or expected to rain

within 2 hours of treatment (Government of Newfoundland and

Labrador, 2019). In 2020, spraying took place between June

26th and July 16th. In 2021, all spraying took place between

June 7th and July 12th. Finally, in 2022, all spraying took place

between June 13th and July 8th. In 2020, all spraying was

limited to areas north of Gros Morne National Park;

consequently, our study sites south of the park received one

year less treatment than the northern sites.

Moth collection

At each site we used one bucket light trap to capture

adult nocturnal macromoths. Each trap was equipped with a

12W black-light fluorescent tube light that was powered by a

deep-cycle lead-acid battery (Figure 1C). 5-gallon buckets were

attached to hold collected moths and three insecticide strips as a

killing agent (Hercon® Vaportape II™). The traps were also

equipped with photocell light sensors that turned the tube

lighting on at dusk and off at dawn.

We placed the traps approximately two meters off the

ground by tying them to nearby trees. Sampling took place over

two summers (2021 and 2022), with three collections (June,

July, and August) per year. Although this does not encompass

the breeding times of all species, this is when the majority are

active (Pinksen *et al.*, 2021). We chose sampling dates based

on the lunar cycle, with the new moon being targeted due to a

reduced influence of moonlight (Yela & Holyoak, 1997).

Sampling in 2021 took place June 15-17, July 13-15, and

August 10-12, and in 2022 from June 28-30, July 27-29, and

August 23-25 for a total of 108 trap-nights per year. There was

no trap disturbance during sampling. During each sampling

event, the traps were set up for three days and emptied daily

into collection boxes that allowed for adequate air flow to

promote drying and limit mold growth. Once collected, samples

were sorted to retain only moths that belonged to families that

are considered macromoths (i.e., Drepanidae, Geometridae,

Saturniidae, Sphingidae, Notodontidae, Erebidae, and

Noctuidae). Moths were collected and identified to the lowest

possible taxon using a variety of resources (Beadle & Leckie,

2012; *Moth Photographers Group*, 2022; John, 2022) A

voucher collection was created and is stored with Natural

Resources Canada, Canadian Forest Service, Corner Brook,

Newfoundland and Labrador, Canada. This voucher collection

was confirmed by B. C. S.

Statistical analyses

We completed all data analysis using R (version 4.2.2;

R Core Team, 2023). We separated the data by stand type

(North Control, North Treatment, South Control, South

Treatment). Separate analyses were conducted in 2021 and

2022.

To assess changes in moth abundance, we used a

generalized linear model using the *glm.nb* function in the R

package *MASS* (Ripley *et al.*, 2023), fitted with a negative

binomial distribution. The model assessed the impact of stand

type on absolute moth abundance per site, pooling data across

months to reduce the effect of seasonal variation. Singletons

were excluded, and post hoc analyses using pairwise Tukey's

tests were conducted for significant findings between treatment

types. The top five most abundant species per year, constituting

$\geq 35\%$ of the collection each year, were similarly analyzed. For

each stand, diversity indices, encompassing species richness,

Shannon diversity, and Simpson diversity, were calculated

using Hill numbers ($q = 0$; species richness, $q = 1$: Shannon

diversity, $q = 2$; Simpson diversity) with the *iNext* package

(Chao *et al.*, 2014; Hsieh, Ma & Chao, 2016). To visualize

differences in species composition between stand types,

nonmetric multidimensional scaling (NMDS) ordination was

calculated with Bray-Curtis dissimilarity scores using the

function *MetaMDS* in the R package *Vegan* (Oksanen *et al.*,

2019). A species-by-site matrix was created using transformed

counts ($x'=\log(x+1)$). Data was pooled by month and

singletons were removed to eliminate the influence of rare

species. The ordination was plotted using the function *ggplot* in

the R package *ggplot2* (Wickham, Navarro & Pedersen, 2016).

Two dimensions were chosen as it was promoted visualization

and interpretation. To assess differences between stand types,

we performed a permutational multivariate analysis of variance

(PERMANOVA) using the function *adonis2* in the R package

Vegan (Oksanen *et al.*, 2019).

Results

We collected and identified 11,802 macromoths,

representing 174 species that belonged to seven families

(Drepanidae, Erebidae, Geometridae, Noctuidae, Notodontidae,

Nolidae, and Sphingidae). Of these, 95.7% were identified to

species level with the remaining unidentifiable.

Abundance

In 2021 and 2022, overall moth abundance was

significantly influenced by stand type ($p<0.001$ and $p=0.0208$,

respectively; Table 1, Figure 2). In 2021, no significant

differences were observed between treatment and control

groups within the north or south regions, the south control sites

had higher abundance than the north control or north treatment

sites (p<0.001 and p=0.0121, respectively). Although not

statistically significant, the mean abundance trended higher in

the north treatment group than the north control group and in

the south control group than the south treatment group. Like

the previous year, we detected no significant difference

between the treatment and control groups within the north and

south regions in 2022. The only significant difference was

between the north control sites and south control sites, with

more individuals in the south control sites (p=0.0467).

However, the mean abundance was higher in the north

treatment group than in the north control group and higher in

the south control group than in the south treatment group.

We assessed any changes in abundance in the most

common species across both years. In 2021, the five most

abundant species made up 39% of the collection, with

Dysstroma citrata (Linnaeus; Geometridae) (n=1183), *Macaria*

signaria (Hübner; Geometridae) (n=613), *Caripeta divisata*

(Walker; Geometridae) (n=381), *Syngrapha viridisigma* (Grote;

Noctuidae) (n=178), and *Anaplectoides pressus* (Grote;

Noctuidae) (n=178). In 2022, the five most abundant species

accounted for 35% of the collection for that year. Two species

remained the same as 2021 (*Macaria signaria* (n=333) and

Dysstroma citrata (n=869)), while three were different (*Xestia*

badicollis (Grote; Noctuidae) (n=152), *Syngrapha rectangula*

(Kirby; Noctuidae) (n=94), and *Phlogophora periculosa*

(Guenée; Noctuidae) (n=133)). We detected significant

differences in the abundance of these species among stand

types in both 2021 and 2022.

In 2021, significant differences were detected in the

abundance of *Syngrapha viridisigma, Macaria signaria,*

Caripeta divisata, Phlogophora periculosa, Dysstroma

citrata,and Xestia badicollis among stand types ($p < 0.001$

each). *Syngrapha viridisigma* were more abundant in the south

control sites than the south treatment sites ($p = 0.0273$), with no

significant difference observed between treatment and control

in the north. *Macaria signaria* was significantly more abundant

in north treatment sites than in north control sites ($p = 0.0221$),

while no significant differences were found in the south.

Caripeta divisata was more abundant in north treatment than

north control ($p < 0.001$), and *Phlogophora periculosa* was

more abundant in south treatment than south control ($p <$

0.001). *Dysstroma citrata* and *Xestia badicollis* had an overall

significant difference in abundance detected but no pairwise

difference was detected. In 2022, *Xestia badicollis*, *Syngrapha*

rectangula, Caripeta divisata, *Dysstroma citrata*, and

Phlogophora periculosa showed significant differences in

abundance based on stand type (p < 0.001 for all except p =

0.0213 for *S. syngrapha*), with controls having higher

abundance. More *Xestia badicollis* individuals were collected

in south controls than south treatments (p = 0.0487). Although

not significant, there were more individuals in north treatment

than north control. For *Syngrapha rectangula*, a significant

difference was observed between south control and south

treatment sites (p = 0.013), with higher numbers in controls.

Despite not being statistically significant, north control had

more individuals compared to north treatment. The other three

species did not have significant pairwise results.

Diversity

In 2021, we observed significant differences in Shannon

diversity (q=1) and Simpson diversity (q=2) between the north

control and treatment sites (Figure 3), but not in species

richness (q=0). There were no differences between the south

control and treatment sites. Despite higher species richness in

the controls, confidence intervals overlapped between north

control and treatment, as well as between south control and

treatment, indicating no significant differences.

Shannon diversity was similar in the south control and

south treatment sites, while confidence intervals did not overlap

between the north control and north treatment sites, suggesting

higher diversity in the north control. Simpson diversity was

similar in the south control, south treatment, and north

treatment sites, but higher in the north control sites.

In 2022, Shannon diversity (q=1) and Simpson diversity

(q=2) differed between stand types, with non-overlapping

confidence intervals between the north treatment and controls,

and the south treatment and controls. Although species richness

(q=0) had overlapping confidence intervals, it was higher in the

north controls than the north treatments and higher in the south

controls than the south treatments.

For Shannon diversity, there was no overlap in

confidence intervals between the south control and south

treatment sites, as well as between the north control and north

treatment sites, indicating higher diversity in the controls in

both regions. Simpson diversity was higher in the north control

group than the north treatment group and higher in the south

control group than the south treatment group.

Between years, we note differences in Simpson and

Shannon diversity, but not in species richness based on

overlapping confidence intervals. Simpson diversity increased

in the north treatment sites and decreased in the control sites

from 2021 to 2022, with no differences found in the southern

sites. Shannon diversity increased in the north treatment sites

but decreased in the south treatment and north control sites. No

difference between years was observed in the south control

sites.

Species composition

NMDS ordination indicated that species composition

varied among stand types (Figure 4). The visualization of the

ordination was recapitulated statistically by the PERMANOVA

results (2021: F=2.1505, p=0.002; 2022: F=2.7568, p=0.001;

stress < 0.1 for all). However, pairwise post-hoc testing did not

identify significant differences between any specific pairs of

groups. The 2021 ordination plot revealed distinct clustering

patterns, with the south treatment group located in the bottom

right quadrant, the south control group located in the top right

quadrant. The northern treatment and control groups

overlapped in the upper and lower left quadrants; however, they

were distinct from the southern groups. Similarly, the 2022

ordination revealed distinct clustering, with the south control

group being in the lower right quadrant, the south treatment

group being in the upper right quadrant, the north control being

in the lower left quadrant, and the north treatment being in the

upper left quadrant.

Discussion

Our results show that overall macromoth abundance and

richness was not higher in the control sites, but community

composition did differ between sites treated with Btk and those

437 that were not (controls). Patterns of changing relative

438 abundance among species were reflected in both the Shannon

439 and Simpson diversity indices. Although species richness did

440 not differ significantly, it was consistently higher in control

441 sites. These complex results point to the need for a more

442 nuanced and thorough assessment of interspecific variation in

443 *Btk* impacts on non-target Lepidoptera, with particular

444 emphasis on understanding individual species responses to *Btk*

445 treatment that could reveal deeper phylogenetic patterns of

446 susceptibility.

447 ***Abundance***

448  We found that the overall abundance of macromoths did

449 not differ between *Btk* treated and control (untreated) sites.

450 These findings are consistent with Glaus *et al.* (2023), who

451 employed similar *Btk* application rates. However, while both of

452 our studies found that the overall abundance of non-target

453 Lepidoptera did not differ between *Btk-treated* areas and non-

454 treated areas, our present study found, although not always

455 significant, there is evidence of interspecific variation in

456 treatment effects. For instance, in 2021 we found that six of the

457 eight most common species across both years (*Dysstroma*

458 *citrata, Macaria signaria*, *Caripeta divisata, Syngrapha*

459 *viridisigma*, *Xestia badicollis* and *Syngrapha rectangula*) were

460 more abundant in the north treatment than the north control.

461 This pattern then carried over to the 2022 season where seven

of the eight most common species were more abundant

(*Caripeta divisata*, *Dysstroma citrata*, *Phlogophora periculosa*,

Syngrapha rectangula Xestia badicollis). In the south sites,

only one of the most abundant species (*Phlogophora*

periculosa) was higher in the treated sites in 2021, but this

increased to three species (*Anaplectoides pressus*, *Dysstroma*

citrata, and *Phlogophora periculosa)* in 2022.

Our findings are in contrast to those of Boulton (2004)

who observed a decline in non-target caterpillar abundance

after *Btk* application. In their study, their *Btk* application was

only for one year, but the spray rate was higher than in our

current study. Sample et al. (1996) also found a decrease in

abundance of both adult and larvae non-target macromoths

following *Btk* application. However, their application was

significantly less than ours: further highlighting the complex

susceptibility of non-targets. Our results were influenced

heavily by the five most abundant species in each year, which

accounted for 39% and 35% of total macromoths in 2021 and

2022, respectively. These species were the primary drivers of

total macromoth abundance. This is in contrast to the findings

of Wagner *et al*. (1996) who found the abundance of the most

common species to be decreased in the treated sites, although

his findings were not statistically significant. Moreover, they

treated their plots with one application of 90 BIU/ha while our

sites were treated with 45 BIU/ha. This discrepancy between

the findings of Wagner et al. (1996) and our study highlights

the potential importance of the concentration (and possibly

frequency) of *Btk* application. The applications of lower doses

may have allowed for reduced immediate impacts and potential

recovery periods, contributing to differing outcomes in our

study. These distinctions underscore the importance of

considering dosage frequency and intensity in understanding

nuanced effects of *Btk* application on non-target moth

populations. While our findings reveal no significant

differences in overall abundance, this is attributed to the

notable increase observed in less than 40% of our entire

collection each year; therefore, the remaining 60% offsets any

potential discernible variations in the total moth population.

Because species differ in phenology, their respective

exposure to Btk can differ substantially. Among Lepidoptera

taxa, sensitivity to *Btk* application often aligns with life history

characteristics (Peacock et al., 1998). Species with an early

phenology, are multivoltine, or univoltine with a long larval

period appear to be less sensitive to *Btk* (Boulton, Otvos &

Ring, 2002; Boulton, 2004). Although comprehensive

phenological data are lacking for all prevalent species in this

study, some patterns emerge. For instance, the activity window

of *Macaria signaria*, which spans from May to August,

suggests that it would have surpassed its larval stage prior to

Btk application. *Anaplectoides pressus*, *Caripeta divisata*, and

Syngrapha rectangula's emergence around June aligns with

their pupal stage occurring prior to *Btk* application. In contrast,

the flight times of *Dysstroma citrata*, *Syngrapha viridisigma*,

Phlogophora periculosa, and *Xestia badicollis* begin in July,

potentially coinciding with the SBW's flight. The overlap in

flight times between these and the SBW could suggest that they

have overlapping larval stages, meaning they likely would be

feeding during *Btk* applications. However, we did not find the

abundance of these species to be lower in the treated sites. This

opens another avenue of investigation for other possible

explanations of why some species may be more-or-less

susceptible to *Btk*.

Certain species may inherently exhibit reduced

sensitivity to *Btk* (Peacock *et al.*, 1998), potentially

contributing to the absence of significant differences in

abundance between treatment and control sites. This may

explain why *Dysstroma citrata*, *Syngrapha viridisigma*,

Phlogophora periculosa, and *Xestia badicollis* were slightly

more abundant in the treatment sites. Given the limited

understanding of these dynamics, this presents an interesting

avenue for future investigation.

Diversity

In the north sites, our findings indicate different

outcomes between 2021 and 2022. Both years showed greater

Shannon and Simpson diversity in control groups compared to

treatment groups, with consistently higher richness in controls,

though not significantly different. This suggests a shift in the

relative abundance of different species within the community in

treated sites, leading to uneven species distribution and

ultimately creating diversity differences. The analysis of

species with low and intermediate abundance revealed minimal

differences between treatment and control groups. However,

focusing on dominant species, including *Anaplectoides pressus*,

Caripeta divisata, *Dysstroma citrata*, *Macaria signaria*,

Phlogophora periculosa, Syngrapha rectangula, Syngrapha

viridisigma, and *Xestia badicollis* were more abundant in the

north treatment than the north control, influencing evenness

and explaining the diversity difference. The control sites

exhibited a higher diversity due to a more equitable distribution

of abundance across species.

In the south, few differences in richness and diversity

between treatment and control groups were observed in 2021.

However, in 2022, diversity decreased in treatment sites. In

2021, only one dominant species (*Phlogophora periculosa*) was

more abundant in treatment sites, but in 2022, three of eight

dominant species (*Dysstroma citrata*, *Anaplectoides pressus*,

and *Phlogophora periculosa*) showed higher abundance in

treatment sites. Richness remained similar among stand types

in 2021 but was slightly higher in control sites. In 2022,

Shannon and Simpson diversity were greater in the control

group than the treatment group, while richness was similar.

This suggests that shifts in relative abundance, especially

among dominant species in treatment sites, account for

variation in diversity.

2021 was the first year of treatment in the southern

sites, followed by a second application in 2022. Lack of

diversity difference in 2021, followed by a decrease in 2022,

suggests a lagged effect of adult macromoth responses to *Btk*

application. This aligns with previous studies noting significant

impacts occurring 12 to 14 months post-spray (Sample *et al.*,

1996; Boulton *et al.*, 2007).

Species composition

We anticipated differences in species composition

between treated and control sites due to documented changes in

relative abundance following exposure to *Btk* (Wagner et al.,

1996). Our ordination analysis revealed distinct clustering

patterns, indicating differences in species composition between

treated and untreated stands.

While PERMANOVA analysis revealed significant

differences in species composition among stand types, there

were no significant pairwise differences. This suggests that

while overall differences exist in moth community structure in

response to *Btk* treatment, these differences may not manifest

uniformly at the level of individual pairwise comparisons

between treated and untreated (control) sites.

The lack of significant pairwise differences underscore

the complexity of community responses to *Btk* treatment and

suggests that the impacts may vary across different species or

sites. Despite the non-significance of the pairwise comparisons,

the significant overall differences indicated by PERMANOVA

suggest that *Btk* treatment exerts discernible effects on the

composition of moth communities, with potential implications

for ecosystem dynamics.

Conclusions

It is a rare and unique opportunity to study the impacts

of insect management at regional spatial scales in forests. Here,

we have shown that the application of *Btk* to control the current

SBW outbreak in western Newfoundland significantly affected

moth species community structure in the first few years of

SBW management. While the overall differences in moth

community composition were significant, the impacts of

treatment may not be consistent across all regions. Moreover,

macromoth responses to *Btk* are somewhat idiosyncratic in that

interspecific variation in developmental phenology and

treatment sensitivity is likely to influence overall impacts;

however, understanding such mechanisms will require further

research.

609   Although significant, impacts observed in our study

610 were much less than those reported for treatments of longer

611 durations and higher application rates (Wagner *et al.*, 1996;

612 Boulton, 2004). Minimizing such effects will require a balance

613 between SBW outbreak control while minimizing *Btk*

614 application, which has been successfully achieved elsewhere

615 (MacLean *et al.*, 2019). The current treatment program was

616 initiated in the context of the Spruce Budworm Early

617 Intervention Strategy framework (Johns *et al.*, 2019), and non-

618 target impacts will ultimately depend on successful execution

619 of this proactive program. With Lepidoptera providing

620 important ecosystem services such as herbivory, pollination,

621 and a food source for other species (Hammond & Miller, 1998),

622 it is imperative to understand the non-target impacts of forest

623 pest control programs. Such information will allow landscape

624 managers to develop strategies that effectively balance pest

625 control measures with the preservation of biodiversity in boreal

626 forest ecosystems.

628 **Acknowledgements**

629   We would like to thank Memorial University – Grenfell

630 Campus for financial support. We would also like to thank

631 Parks Canada, Gros Morne National Park for access to some

632 sites for data collection (permit: GMP-2022-42098 to JJB). We

would also like to acknowledge the collaborative efforts of the

Dept. of Fisheries, Forestry and Agriculture, Newfoundland and

Labrador as we would not be able to conduct this large-scale

field experiment without their cooperation. Lastly, we would

like to thank Jamie Warren for his technical help.

Conflicts of Interest

The authors declare that they have no competing interests.

References

Alison, J. *et al.* (2022) Moths complement bumblebee

pollination of red clover: A case for day-And-night insect

surveillance. *Biology Letters*, **18**(7).

https://doi.org/10.1098/rsbl.2022.0187.

Anderson, M., Rotheray, E.L. and Mathews, F. (2023)

Marvellous moths! pollen deposition rate of bramble (Rubus

futicosus L. agg.) is greater at night than day. *PloS one*, **18**(3).

https://doi.org/10.1371/journal.pone.0281810.

Bauce, E. *et al.* (2004) Bacillus thuringiensis subsp. kurstaki

aerial spray prescriptions for balsam fir stand protection against

spruce budworm (Lepidoptera: Tortricidae). *Journal of*

Economic Entomology, **97**, 1624–1634. Available at:

https://cfs.nrcan.gc.ca/publications?id=25129.

Beadle, D. and Leckie, S. (2012) *Peterson Field Guide to Moth*

of Northeastern North America.

Blais, J.R. (1983) Trends in the frequency, extent, and severtiy

of spruce budworm outbreaks in eastern Canada. *Laurentian*

Forest Research Center [Preprint].

Boulton, T.J. (2004) Responses of nontarget Lepidoptera to

foray 48b® Bacillus thuringiensis var. kurstaki on Vancouver

Island, British Columbia, Canada. *Environmental Toxicology*

and Chemistry, **23**(5), 1297–1304. https://doi.org/10.1897/03-

88.

Boulton, T.J. *et al.* (2007) Recovery of nontarget lepidoptera on

Vancouver Island, Canada: One and four years after a gypsy

moth eradication program. *Environmental Toxicology and*

Chemistry, **26**(4), 738–748. https://doi.org/10.1897/06-

079R1.1.

Boulton, T.J. and Otvos, I.S. (2004) Monitoring native non-

target Lepidoptera for three years following a high dose and

volume application of Bacillus thuringiensis subsp. kurstaki.

International Journal of Pest Management, **50**(4), 297–305.

https://doi.org/10.1080/09670870412331286058.

Boulton, T.J., Otvos, I.S. and Ring, R.A. (2002) Monitoring

nontarget Lepidoptera on Ribes cereum to investigate side

effects of an operational application of Bacillus thuringiensis

subsp. kurstaki. *Environmental Entomology*, **31**(5), 903–913.

https://doi.org/10.1603/0046-225X-31.5.903.

680 Burford, L.S., Lacki, M.J. and Covell, C. V. (1999) Occurrence

681 of moths among habitats in a mixed mesophytic forest:

682 Implications for management of forest bats. *Forest Science*,

683 **45**(3), 323–332.

684 Chagnon, C., Bouchard, M. and Pothier, D. (2022) Impacts of

685 spruce budworm defoliation on the habitat of woodland

686 caribou, moose, and their main predators. *Ecology and*

687 *Evolution*, **12**(3), 1–14. https://doi.org/10.1002/ece3.8695.

688 Chang, W. *et al.* (2012) Economic impacts of forest pests: a

689 case study of spruce budworm outbreaks and control in New

690 Brunswick, Canada, **505**(February), 490–505.

691 https://doi.org/10.1139/X11-190.

692 Chao, A. *et al.* (2014) Rarefaction and extrapolation with Hill

693 numbers: A framework for sampling and estimation in species

694 diversity studies. *Ecological Monographs*, **84**(1), 45–67.

695 https://doi.org/10.1890/13-0133.1.

696 Dymond, C.C. *et al.* (2010) Future spruce budworm outbreak

697 may create a carbon source in Eastern Canadian forests.

698 *Ecosystems*, **13**(6), 917–931. https://doi.org/10.1007/s10021-

699 010-9364-z.

700 van Frankenhuyzen, K. (2000) Application of Bacillus

701 thuringiensis in forestry. in J.-F. Charles, A. Delécluse, and

702 C.N.-L. Roux (eds) *Entomopathogenic Bacteria: from*

703 *Laboratory to Field Application*. Dordrecht: Springer

Netherlands, 371–382. https://doi.org/10.1007/978-94-017-

1429-7_20.

Fuentealba, A. *et al.* (2019) Comparing the efficacy of various

aerial spraying scenarios using Bacillus thuringiensis to protect

trees from spruce budworm defoliation. *Forest Ecology and*

Management, **432**(October 2018), 1013–1021.

https://doi.org/10.1016/j.foreco.2018.10.034.

Glaus, V. *et al.* (2023) Nontarget impacts of insecticide-based

population control of eastern spruce budworm (Lepidoptera:

Tortricidae) on nontarget caterpillar communities and

parasitism. *Canadian Entomologist*, **155**, 1–17.

https://doi.org/10.4039/tce.2022.47.

Government of Newfoundland and Labrador (2019) Pesticide

Operator Licence - Terms and Conditions.

Government of Newfoundland and Labrador (2020) *Forest*

insect disease control. Available at:

https://www.gov.nl.ca/ffa/programs-and-funding/forestry-

programs-and-funding/idc/2022-insect-control-program/.

Hammond, P.C. and Miller, J.C. (1998) Comparison of the

Biodiversity of Lepidoptera Within Three Forested

Ecosystems. *Annals of the Entomological Society of America*,

91(3), 323–328.

Holmes, R.T., Schultz, J.C. and Nothnagle, P. (1979) Bird

Predation on Forest Insects : An Exclosure Experiment,

206(4417), 462–463.

Hsieh, T.C., Ma, K.H. and Chao, A. (2016) iNEXT: an R

package for rarefaction and extrapolation of species diversity

(Hill numbers). *Methods in Ecology and Evolution*, **7**(12),

1451–1456. https://doi.org/10.1111/2041-210X.12613.

John, V. (2022) *Bug Guide*. Available at: www.bugguide.net.

Johns, R.C. *et al.* (2019) A conceptual framework for the

spruce budworm Early Intervention Strategy: Can outbreaks be

stopped? *Forests*, **10**(10), 1–19.

https://doi.org/10.3390/f10100910.

Knowles, B.H. and Ellar, D.J. (1987) Colloid-osmotic lysis is a

general feature of the mechanism of action of Bacillus

thuringiensis δ-endotoxins with different insect specificity.

BBA - General Subjects, **924**(3), 509–518.

https://doi.org/10.1016/0304-4165(87)90167-X.

Lambert, B. and Peferoen, M. (1992) Insecticidal Promise of

Bacillus thuringiensisFacts and mysteries about a successful

biopesticide. *BioScience*, **42**(2), 112–122. Available at:

https://academic.oup.com/bioscience/article/42/2/112/259045.

Leza, M., Gonzalez-Ruiz, R. and Alemany, A. (2021) Short

term effects of Bacillus thuringiensis and diflubenzuron aerial

applications on non-target arthropods in a Mediterranean

Forest. *International Journal of Pest Management*, **67**(2), 139–

146. https://doi.org/10.1080/09670874.2019.1710619.

MacLean, D.A. (1984) Effects of Spruce Budworm Outbreaks

on the Productivity and Stability of Balsam Fir Forests. *The*

Forestry Chronicle, **60**(5), 273–279.

https://doi.org/10.5558/tfc60273-5.

MacLean, D.A. *et al.* (2019) Positive results of an early

intervention strategy to suppress a spruce budworm outbreak

after five years of trials. *Forests*, **10**(5).

https://doi.org/10.3390/f10050448.

Maclean, D.A. and Ostaff, D.P. (1989) Patterns of balsam fir

mortality caused by an uncontrolled spruce budworm outbreak.

Canadian Journal of Forest Research, **19**(9), 1087–1095.

Manderino, R., Crist, T.O. and Haynes, K.J. (2014)

Lepidoptera-specific insecticide used to suppress gypsy moth

outbreaks may benefit non-target forest Lepidoptera.

Agricultural and Forest Entomology, **16**(4), 359–368.

https://doi.org/10.1111/afe.12066.

Miller, J.C. (1990) Field Assessment of the Effects of a

Microbial Pest Control Agent on Nontarget Lepidoptera.

American Entomologist, **36**(2), 135–140.

https://doi.org/10.1093/ae/36.2.135.

Miller, J.C. (2000) Monitoring the Effects of Bacillus

thuringiensis kurstaki on Nontarget Lepidoptera in Woodlands

and Forests of Western Oregon. in *Nontarget Effects of*

Biological Control, 277–286. https://doi.org/10.2307/177420.

Moth Photographers Group (2022). Available at:

https://mothphotographersgroup.msstate.edu/.

Oksanen, A.J. *et al.* (2019) Package 'Vegan'. *Encyclopedia of*

Food and Agricultural Ethics, 2395–2396.

https://doi.org/10.1007/978-94-024-1179-9_301576.

Peacock, J.W. *et al.* (1998) Laboratory Assessment of the

Effects of Bacillus thuringiensis on Native Lepidoptera.

Environmental Entomology, **27**(2), 450–457.

https://doi.org/10.1093/ee/27.2.450.

Pinksen, J. *et al.* (2021) Living on the edge: Effects of clear-cut

created ecotones on nocturnal macromoth assemblages in the

eastern boreal forest, Canada. *Forest Ecology and*

Management, **494**(April), 119309.

https://doi.org/10.1016/j.foreco.2021.119309.

Protected Areas Association of Newfoundland and Labrador

(2008a) Northern Peninsula Forest.

Protected Areas Association of Newfoundland and Labrador

(2008b) Western Newfoundland Forest. Available at:

https://www.gov.nl.ca/eccm/files/publications-parks-

ecoregions-island-1a-serpentine-range.pdf.

R Core Team (2023) R: A Language and Environment for

Statistical Computing. Vienna, Austria. Available at:

https://www.r-project.org/.

Ripley, B. *et al.* (2023) Package 'MASS'.

Sample, B.E. *et al.* (1996) Effects of Bacillus thuringiensis

berliner var. kurstaki and defoliation by the gypsy moth

[Lymantria dispar (L.) (Lepidoptera:Lymantriidae)] on native

arthropods in West Virginia. *The Canadian Entomologist*, **128**,

573–592.

Sanahuja, G. *et al.* (2011) Bacillus thuringiensis: A century of

research, development and commercial applications. *Plant*

Biotechnology Journal, **9**(3), 283–300.

https://doi.org/10.1111/j.1467-7652.2011.00595.x.

Sanmartín-Villar, I. and Cordero-Rivera, A. (2024)

Biodiversity and the importance of insect diversity. *Biological*

Invasions and Global Insect Decline, 19–46.

https://doi.org/10.1016/B978-0-323-99918-2.00002-1.

Schünemann, R., Knaak, N. and Fiuza, L.M. (2014) Mode of

Action and Specificity of Bacillus thuringiensis Toxins in the

Control of Caterpillars and Stink Bugs in Soybean Culture .

ISRN Microbiology, **2014**, 1–12.

https://doi.org/10.1155/2014/135675.

Strazanac, J.S. *et al.* (2005) Long Term Evaluation of the

Effects of Bacillus thuringiensis kurstaki , Gypsy Moth

Nucleopolyhedrosis Virus Product Gypchek ® , and

Entomophaga maimaiga on Nontarget Organisms in Mixed

Broadleaf-Pine Forests in the Central Appalachians. *US*

Department of Agriculture, Forest Service, FHTET, **122**(4), 1–

402.

Summerville, K.S. (2011) Managing the forest for more than

the trees: Effects of experimental timber harvest on forest

Lepidoptera. *Ecological Applications*, **21**(3), 806–816.

https://doi.org/10.1890/10-0715.1.

Wagner, D.L. *et al.* (1996) Field Assessment of Bacillus

thuringiensis on Nontarget Lepidoptera. *Environmental*

Entomology, **25**(6), 1444–1454.

https://doi.org/10.1093/ee/25.6.1444.

Wagner, D.L. (2005) *Caterpillars of Eastern North America*.

Princeton University Press, Princeton & Oxford.

Wagner, D.L. *et al.* (2021) A window to the world of global

insect declines: Moth biodiversity trends are complex and

heterogeneous. *Proceedings of the National Academy of*

Sciences of the United States of America, **118**(2), 1–8.

https://doi.org/10.1073/PNAS.2002549117.

Wickham, H., Navarro, D. and Pedersen, T.L. (2016) *ggplot2:*

Elegant Graphics for Data Analysis. Springer-Verlag New

York. Available at: https://ggplot2.tidyverse.org.

Yela, J.L. and Holyoak, M. (1997) Effects of Moonlight and

Meteorological Factors on Light and Bait Trap Catches of

Noctuid Moths (Lepidoptera: Noctuidae). *Environmental*

Entomology, **26**(6), 1283–1290.

https://doi.org/10.1093/ee/26.6.1283.

Young, M. (2005) Insects in flight. in *Insect Sampling in forest*

Ecosystems, 116–146.

Table 1. Total raw moth abundance for the five most abundant species in 2021 and 2022. Collection days amalgamated by year and stand type.

Control/treatment pairs that appear in a box are significantly different. NC = North control, NT = North treatment, SC = South control, ST =

	2021	2022

854   South treatment

Species	Top 5 year	N	NC	NT	SC	ST	Overall sig.	N	NC	NT	SC	ST	Overall sig.
Anaplectoides pressus	2021	178	37	26	67	48	NS	118	41	23	25	29	NS
Caripeta divisata	2021	381	36	173	115	57	*	90	17	51	11	11	*
Dysstroma citrata	2021/22	1183	70	123	585	405	*	967	98	255	260	354	*
Macaria signaria	2021/22	613	61	271	154	127	*	356	47	169	78	62	NS
Phlogophora periculosa	2022	135	21	20	24	70	*	158	8	48	23	79	*
Syngrapha rectangula	2022	94	9	14	62	9	NS	204	43	59	80	22	*
Syngrapha viridisigma	2021	178	18	26	109	25	*	74	24	25	19	6	NS
Xestia badicollis	2022	152	1	5	135	11	*	312	4	20	207	81	*

Figure 1. a) Newfoundland, Canada. b) Region of study in western Newfoundland. White

circles represent control (non-Btk treated) sites, orange triangles represent Btk treated sites,

and the hollow orange triangle represents the replaced site. c) Black-light fluorescent trap set

up for moth collection at one of the boreal forest sites.

Figure 2. Box and whisker plots of moth abundance from four stand types, across two years,

a) 2021 and b) 2022. 2021 treatment sites had been sprayed with *Btk* for one (South

Treatment) and two (North Treatment) consecutive year(s). 2022 treatment sites had been

sprayed with *Btk* for two (South Treatment) and three (North Treatment) consecutive years.

Comparisons tested using a generalized linear model (GLM), fitted with a negative binomial

distribution. Letters a, b, and c indicate significant differences between stand types based on

post-hoc testing (95% confidence).

Figure 3. Diversity indices, characterized by effective number of species ($\pm$ 95% confidence

interval), for each treatment group for a) 2021 and b) 2022. Values calculated from Hill

number order $q = 0$ (species richness), $q = 1$ (Shannon diversity), and $q = 2$ (Simpson

diversity), for each stand type.

Figure 4. Nonmetric multidimensional scaling of species composition across stand types in a)

2021(Stress=0.119) and b) 2022 (Stress=0.093). Species abundance pooled across all dates

and log transformed ($x' = \log(x+1)$). Bray-Curtis dissimilarity scores were used. Each

individual treatment plot represents a collection month for north control, north treatment,

south control, and south treatment.

2021

2022

North Control   North Treatment   South Control   South Treatment