Transcriptional shifts and microbiome dynamics in sorghum roots during vegetative growth under drought stress

Ahmad H. Kabir[1*], Philip Brailey-Jones[2], Mostafa Abdelrahman[3], Jean Legeay[4], Bulbul Ahmed[4],
Lam-Son Phan Tran[3], Jeffrey L. Bennetzen[2*]

[1]School of Sciences, University of Louisiana at Monroe, LA 71209, USA

[2]Department of Genetics, University of Georgia, Athens, GA 30602, USA

[3]Institute of Genomics for Crop Abiotic Stress Tolerance, Department of Plant and Soil Science, Texas Tech University, TX 79409, USA

[4]African Genome Center, University Mohamed VI Polytechnic (UM6P), Ben Guerir 43150, Morocco

[*]e-mail: kabir@ulm.edu, maize@uga.edu

Abstract

Drought stress significantly impairs growth, nutrient uptake, and microbial interactions in sorghum, posing a critical challenge to its productivity. This study explores the transcriptional and microbial shifts in 6-week-old sorghum under drought conditions, revealing key adaptations to water deficit. Our LC-MS (Liquid Chromatography-Mass Spectrometry) analyses revealed that drought stress induced abscisic acid while significantly reducing jasmonic acid levels in sorghum roots, likely due to resource conservation strategies during drought. Transcriptional reprogramming highlighted the upregulation of genes in the roots involved in mineral homeostasis (*Ferritin 1*, *Iron dehydrogenase*, *Nitrate transporter 1*), hormone signaling (*Ethylene-insensitive protein 3*, *Gibberellin 2-oxidase*), and osmotic regulation (*Aquaporin*, *Dehydrin*), underscoring their roles in maintaining nutrient uptake, mitigating oxidative stress, and preserving cellular turgor. These transcriptional shifts align with the adaptive responses of sorghum plants to optimize their water use, thereby sustaining their growth under drought. Drought reduced root-associated bacterial diversity and richness while enriching drought tolerance-associated genera, such as *Burkholderia, Caballeronia* and *Paraburkholderia*, known for promoting plant growth through auxin production, phosphate solubilization, and siderophore-mediated iron acquisition. In contrast, fungal diversity and richness remained unchanged, dominated by *Talaromyces*, which showed a statistically non-significant increase under drought. Random forest models could not identify functional predictors for fungi but revealed a shift in bacterial functional groups under drought, with enrichment in phototrophy, methylotrophy, and nitrate reduction, traits emphasizing microbial roles in nutrient cycling and drought adaptation of sorghum. This study provides insights into the transcriptional and microbial responses underlying drought tolerance and identifies potential bacterial bioinoculants that could enhance sorghum resilience. Future research should validate these findings in field conditions to develop sustainable strategies for improving drought tolerance in sorghum, with broader implications for climate-resilient agriculture.

Keywords: Bioenergy crop, jasmonic acid, microbial alternations, osmotic adjustment, photosynthetic efficiency.

47
48
49 **Running title:** Responses in drought-exposed sorghum
50
Introduction
Drought stress, leading to adverse impacts on growth, development, and productivity (Lamaoui
et al., 2018), is ubiquitous across crops and natural plant populations but varies dramatically in
intensity and frequency. Primary drought stress symptoms, especially wilting and leaf curling,
are reflective of reduced turgor pressure and water loss reduction strategies (Ruehr et al., 2019;
Zoghi et al., 2019). Additional drought stress manifestations include leaf yellowing, stunted
growth, and premature leaf senescence (Khan et al., 2018). Water deficiency disrupts essential
physiological processes like photosynthesis, nutrient uptake, and hormone regulation, ultimately
compromising the plant's ability to thrive (Hussain et al., 2018). Additionally, crops subjected to
drought stress often show increased susceptibility to pests and diseases (Duveiller et al., 2007).
This major agricultural problem is expected to be more prevalent under current climate change
scenarios. As global temperatures rise and weather patterns shift, many regions are experiencing
unusually long periods of low precipitation and resultant water shortages. Sorghum (*Sorghum
bicolor*) is known for its exceptional drought tolerance, but its yields are affected by prolonged
periods of water scarcity (Rajarajan et al., 2021). Studies predict that drought, associated with
temperature increases, could cause up to a 44% yield loss in US sorghum production in the near
future, emphasizing the need for additional efforts to improve resilience, especially during the
pre-flowering phase (Tack et al., 2017).
69
Plants employ diverse strategies to regulate water loss, including stomatal and osmotic
adjustments through the accumulation of compatible solutes and the enhanced production of
antioxidant enzymes and metabolites to counteract reactive oxygen species (ROS) generated
under drought conditions (Seleiman et al., 2021; Takahashi et al., 2020). Hormones, particularly
abscisic acid (ABA) and auxin, play a pivotal role in orchestrating these responses by
modulating stomatal closure and activating stress-responsive genes, ultimately enhancing a
plant's ability to withstand periods of water shortage (Iqbal et al., 2022; Sadok and Schoppach,
2019). Minerals are also essential for plant drought tolerance, as they regulate osmotic
adjustment, stomatal control, and stress signaling, which help plants maintain water balance
(Khan et al., 2023). Particularly, calcium is vital for maintaining cell wall integrity and
membrane stability, which can be compromised during drought conditions (Feng et al., 2023).
81
Microbes can improve plant stress resistance, including drought tolerance, by instigating a range
of processes that alter plant biochemistry and physiology (Poudel et al., 2021; Timmusk et al.,
2013). Fungal communities, including AMF (arbuscular mycorrhizal fungi), are sometimes
associated with drought stress tolerance in some crops (Chen et al., 2021; Ferrol et al., 2019).
Studies suggest that AMF can directly influence plant performance by modifying the plant–water
relationship, leading to increased plant productivity during periods of drought stress (Ma et al.,
2020; Rapparini and Peñuelas, 2014). *Trichoderma*, a fungal genus, has gained attention for its
potential role in enhancing drought tolerance in plants. Two ways in which *Trichoderma* are
believed to contribute to drought tolerance are by inducing a systemic drought stress response in
plants and by participating in the production of specific metabolites that positively impact plant
responses to drought (Scudeletti et al., 2021; Mona et al., 2017). Monoderm bacterial lineages,

such as *Actinobacteria*, *Chloroflexi*, and *Firmicutes*, induce increased desiccation tolerance in
plants, associated with thicker plant cell walls, and these microbes' increased abundances are
driven by drought-responsive plant metabolism (Xu and Coleman-Derr, 2019). Inoculation of
Bacillus amyloliquefaciens S-134 on wheat plants increased the production of auxins, stimulated
root growth, and assisted plants in adapting to water stress conditions. Particularly, *Streptomyces*
are reported to produce auxin and siderophores that can promote mineral availability, stress
tolerance, and growth in plants (Anwar et al., 2016; Palaniyandi et al., 2014).
100
Sorghum is a promising bioenergy crop because of its high biomass yield and suitability for
producing biofuels (Mathur et al., 2017; Prasad et al., 2021). Given the increasing frequency of
drought events caused by climate change, understanding the molecular and microbial
mechanisms underlying drought tolerance in sorghum is essential. Previous studies have
documented individual mechanisms of drought tolerance, such as hormonal regulation or
bacterial interactions (Qi et al., 2022; Rajarajan et al., 2021). However, the interconnected
dynamics of plant transcriptomics and root-associated bacterial and fungal microbiomes remain
underexplored, particularly in a bioenergy crop like sorghum at vegetative stage. Therefore, this
research seeks to elucidate the interplay between plant gene expression and microbial
interactions, identifying key gene networks and microbial taxa and their functional groups that
may contribute to drought tolerance in sorghum.
112
Materials and methods
Plant cultivation and growth conditions
Seeds of sorghum (USDA-GRIN accession PI 591038) were surface sterilized using a 1%
sodium hypochlorite bleach solution for 5 min. Following sterilization, the seeds were rinsed
three times with sterile water and subsequently placed in a tray for germination at 25°C in an
incubator for a duration of 2 d. Plants were cultivated in the greenhouse in pots in a mixture of
field soil and SunGro potting mix (1:3) in two different conditions: control and drought. Drought
conditions were initiated by reducing water availability to 75% compared to the controls (well-
watered) two weeks after transplanting the seedlings into the soil pots. Daily monitoring of soil
moisture content was conducted to maintain 75% and 25% soil moisture levels for control and
drought-stressed plants, respectively. Plants were cultivated following a randomized complete
block design (RCBD), with 9 replications (3 plants/pot) per treatment in a growth chamber (10 h
light/14 h darkness, 250 μmol m^{-2} s^{-1}) at 25°C for 6 weeks before measuring physiological
properties and harvesting samples for other physiological and nucleic acid analyses, for all
experiments.
128
Determination of morphological and gas-exchange attributes
The height of each plant was measured using a measuring tape. The aboveground portions of the
plant were harvested and dried for 3 days at 75°C in an incubator before obtaining the dry
weight. Furthermore, the relative water content (%) in the leaf was measured as previously
described (Soltys-Kalina et al., 2016). Relative water content (RWC) of leaves was measured on
the fully expanded trifoliate leaves at the tip of the main stem as previously described (Schonfeld
et al., 1988). Briefly, the fresh weights (FW) of leaves from each treatment were recorded
immediately after removal. The turgid weight (TW) was measured after soaking the leaves in
distilled water in beakers for 24 h at room temperature. Following the soaking period, the leaves
were quickly and carefully blotted dry with tissue paper to prepare for turgid weight

determination. The dry weight (DW) of the leaves was obtained by oven-drying the leaf samples for 72 h at 70ºC. RWC was calculated using the formula: RWC (%) = (FW – DW) / (TW – DW) x 100.

Gas exchange and photosynthetic parameters of the leaves were measured using the LI-6800 portable Photosynthesis System (LI-COR Biosciences, USA). The photosynthetic rate, transpiration rate, stomatal conductance, and intrinsic water-use-efficiency (iWUE) were assessed on the central portion of the fourth fully expanded leaf under the following conditions: 2 cm² leaf area, flow rate of 600 $\mu mol\ s^{-1}$, relative humidity of 65%, CO_2 reference of 400 $\mu mol\ mol^{-1}$, leaf vapor pressure deficit of 1.5 kPa, leaf temperature of 25°C, and leaf-level light intensity of 250 μmol photons $m^{-2}\ s^{-1}$. Measurements were conducted three times, averaged for statistical analyses, and calculated as previously described (de Oliveira et al., 2022).

Elemental analysis of plant tissues
After harvesting, the roots of 6-week-old sorghum plants were excised and rinsed with running tap water, followed by immersion in 0.1 mM $CaSO_4$ for 10 min, and a final wash with deionized water. Subsequently, the roots and leaves were dried in an incubator for 3d at 75°C. Elemental concentrations were analyzed using ICP-MS (inductively coupled plasma mass spectrometry) at the Agricultural & Environmental Services Laboratories at the University of Georgia. The concentration of elements in the samples was determined by comparing the intensity of ion peaks in the spectrum to calibration standards.

Determination of H_2O_2 in roots
For H_2O_2 determination, root samples were mixed in a solution of 0.1% trichloroacetic acid, as previously described (Alexieva et al., 2001). Briefly, the samples were finely ground into a powder using a mortar and pestle. The extracted fluid underwent centrifugation for 15 min at 10,000 rpm to separate cellular residues. The resulting upper aqueous phase was supplemented with potassium iodide (1 M) and phosphate buffer (10 mM, pH adjusted to 7.0), and then left in darkness for 60 min. Finally, the optical density of the solution was measured at 390 nm using a spectrophotometer (ThermoFisher, Waltham, USA), and the concentration of H_2O_2 was calculated based on the standard curve.

RNA-seq analysis
Before RNA extraction, roots were meticulously cleansed twice in sterile phosphate-buffered saline (PBS) through vortexing for 10 sec, followed by two rinses with sterile water to minimize surface contaminants. Subsequently, samples were pulverized into a powder using a pre-cooled pestle and mortar with liquid nitrogen. The resulting powdered material was then utilized for RNA extraction. Total RNA extraction was performed using the SV Total RNA Isolation System (Promega, USA), following the manufacturer's instructions. A total of 1 µg of RNA per sample, possessing a RIN value >8, was employed as input material for RNA-seq library preparations. Sequencing libraries were constructed using the KAPA HiFi HotStart Library Amplification Kit (Kapa Biosystems, USA), per the manufacturer's instructions. The resulting libraries were sequenced on the Illumina NextSeq SE 100bp platform. Overall, >80% of reads passed filters with Phred score values exceeding 33% per sample.

Adaptor sequences and low-quality sequences were respectively removed and filtered from raw reads using Trimmomatic v0.38 (Bolger et al., 2014). The sorghum reference genome (Sbicolor_454_v3.0.1) was obtained from Phytozome (McCormick et al., 2017) and the filtered reads were aligned to the reference genome using HISAT2 (v2.0.5) software. HT-Seq was utilized to count the read numbers mapped to each gene (Anders et al., 2014). Differentially expressed genes (DEGs) were identified using the Limma R package v3.19 (Law et al., 2016) and read counts were normalized by using the FPKM (fragments per kilobase of exon per million mapped fragments) method. Genes were identified as up-regulated or down-regulated using a cut-off threshold of $\log_2$ fold change (Log_2FC) ≥ 1 and ≤ -1, respectively, with a false discovery rate (FDR) ≤ 0.05. Venn diagrams and heatmaps were generated using the VennDetail R package v1.20 (Hinder et al., 2017) and the ClustVis (https://biit.cs.ut.ee/clustvis/) tool (Metsalu et al., 2015), respectively. Gene ontology for sorghum was retrieved using Phytozome (Goodstein et al., 2012) resources. Gene annotation for molecular function (MF), biological process (BP), and cellular component (CC) was performed through the g:Profiler web tool (Kolberg et al., 2023).

Phytohormone analysis in roots

Phytohormone concentrations in sorghum roots were determined using LC-MS (Liquid Chromatography-Mass Spectrometry) at the Michigan State University Mass Spectrometry and Metabolomics Core. In brief, approximately 100 mg of fresh root material was homogenized in liquid nitrogen and extracted using a solution of 80:20 (v/v) methanol: water, 0.1% formic acid, and 0.1 g/L butylated hydroxytoluene. The samples were then placed in pre-frozen bead beater adaptors and beaten (30/s for 30 s) in a Tissue Lyser II (Qiagen, USA). Following a brief centrifugation (20 s) at 4°C to settle ground tissue at the tube bottom, the samples were further incubated on a shaker at 5,000 rpm at 4°C for 16 h. Subsequently, the samples were vortexed and centrifuged at 12,000 rpm for 10 min at 4°C, and the supernatant was transferred to autosampler vials. Standards for each phytohormone (Sigma-Aldrich, USA) were prepared in 80% methanol and 0.1% formic acid.

Amplicon sequencing targeting fungi and bacteria

Harvested roots were cleaned by vortexing in sterile phosphate-buffered saline (PBS) for 10 seconds, followed by two rinses with sterile water to remove surface contaminants. Approximately 0.2 g aliquots of cleaned root samples were used for DNA extraction following the procedure outlined by Clarke (2009), along with RNase and Proteinase K treatments. Four biological replicates were used for microbiome analysis. Amplicon libraries were prepared for sequencing using a two-step PCR approach (Kozich et al., 2013). Two primer sets targeting regions of the ribosomal DNA gene were used to generate the libraries. The internal transcribed spacer (ITS) was used to target fungal communities (fITS7: GTGARTCATCGAATCTTTG and ITS4: TCCTCCGCTTATTGATATGC), whereas the small sub-unit encoding region (16S) was used to target bacterial communities (515F: TGTGYCAGCMGCCGCGGTAA and 806R: GGACTACNVGGGTWTCTAAT). Primers were adapted to include a 12N UDI sequence immediately prior to the target gene sequence to aid in cluster delineation when sequencing and Illumina P5/P7 complementary overhangs to facilitate barcode index addition in Step 2 PCR. Step 1 PCRs were carried out using Hot Start GoTaq master mixes (Promega, Madison, WI USA) in 12.5 ul volumes containing 2 ul of DNA, 2 mM dNTPs, and 10 pmol of each primer. PCR conditions were the same for both 16S and ITS amplicon generation: 95C for 5 min, 35 cycles at 94C for 30 s, 55C for 30s, 72C 30s, and 72C for 5 min. PCR steps were carried out

using a BioRad T100 thermal cycler (Bio-Rad, Hercules, CA USA). Amplicons were dual barcoded with Illumina i7/i5 style barcodes in step 2 PCR using Phusion Hot Start II master mix (Thermo Fisher, Waltham, MA USA) to ensure high-fidelity barcode additions. Amplicons were sequenced using the NextSeq2000 platform on a P2 flow cell run in paired-end 300 bp mode (Illumina, San Diego, USA) by the Georgia Genomics and Bioinformatics Core (Athens, GA USA).

Adapter and primer sequences were trimmed from the raw sequencing reads using *Cutadapt v. 4.5* (Martin 2011). Next, low-quality reads were removed using the filterAndTrim function of *dada2* (*v. 1.28.0*) implemented in R (*v. 4.3.0*, Callahan et al., 2016), followed by error model estimation, dereplication, sample inference, and nonchimeric removal to generate a table of amplicon sequence variants (ASVs). Prior to taxonomy assignment, ASVs were pre-filtered by BLAST-matching against the UNITE database for ITS (dynamic_s_all_25.07.2023, Abarenkov et al., 2024) and the SILVA database for 16S (v 138.2_SSURef_NR99, Quast et al., 2013). Any sequences with a query coverage < 80% were removed. Both ITS and 16S taxonomy was assigned using CONSTAX2 (Liber *et al.,* 2021), which carries out consensus taxonomy assignment from multiple approaches, using the same databases. Further taxonomic curation was performed using *phyloseq* implemented in R (Callahan et al., 2016). 16S OTUs were filtered to include only OTUs assigned to the Bacterial kingdom as the 'higher level taxonomy' assignment, which excludes chloroplast and mitochondrial sequences. ITS OTUs were filtered to remove any OTUs where the 'higher level taxonomy' assignment through CONSTAX2 was not to the Fungal kingdom. Alpha diversity metrics (observed species richness and Shannon index) were calculated using *phyloseq*. Alpha diversity metrics were compared between treatment groups using ANOVA, followed by post-hoc testing where appropriate with 'hsd' multiple testing corrections employed. Beta diversity was assessed through Bray-Curtis dissimilarities calculated between samples and statistically compared between treatments with PERMANOVA analysis using vegan (*v. 2.6.4*). Community dissimilarity between samples and treatments was visualized through principal co-ordinate analysis (PCoA). Two methods were employed to determine which taxa were more associated with different treatments at the genus level. Indicator species analysis was conducted using the R package indicspecies (*v. 1.7.14*) on a core set of genera (>0.001 RA in > 10% of samples) consisting of 80 out of 306 genera. The resulting *p*-values were adjusted for multiple testing using "fdr" correction. The relative abundances of the top ten most abundant genera of bacteria and fungi were also compared between treatments using ANOVA, again followed by post-hoc testing where appropriate.

Furthermore, the dataset was trimmed to retain only taxa accounting for more than 0.1% of the total reads before conducting co-occurrence network analyses and random forest modeling. Co-occurrence networks were calculated and plotted using SpiecEasi (Kurtz et al., 2015) and igraph (Antonov et al., 2024) packages in R, with betweenness centrality values computed for all ASVs using the betweenness function of igraph. Taxa in the top 5% of betweenness values were designated as hub taxa. Random forest models, implemented using the randomForest (Breiman 2001) package in R with 100 iterations of decision trees, were used to identify ASVs predictive of drought conditions, with ASVs ranked by their GINI index, where a lower index indicates higher predictive importance. Functional pathways for bacterial 16S rRNA ASVs were attributed using the PICRUSt2 (Langille et al., 2013) package and the MetaCyc (Karp et al., 2002) database, while bacterial functional traits were assigned using the FAPROTAX (Louca et al.,

276   2016) package. Similarly, fungal ITS ASVs were annotated with functional groups using the
277   FunFun (Krivonos et al., 2023) package and the KEGG (Kanehisa et al., 2000) database.
278
279   **Statistical analysis of plant characteristics**
280   All measurements were conducted with at least three independent biological replicates for each
281   treatment. To identify significant differences between treatment groups ($P < 0.05$), student t-test
282   was carried out with treatment as an independent variable using SPSS Statistics 20.0 (IBM).
283   Graphical representation was performed using GraphPad Prism 6.0.
284
Results
Changes in morphological and physiological traits under drought stress
Drought stress caused a substantial decline in shoot and root systems compared to controls in
these growth chamber experiments, with soils containing natural field microbiomes (Fig. 1A and
1B). The root length, root dry weight, shoot height, shoot dry weight, leaf relative water content,
and leaf chlorophyll score showed a significant decrease under drought compared to controls
(Fig. 1C-1H). Furthermore, we found a significant increase in H_2O_2 concentrations in the roots of
sorghum exposed to drought relative to controls (Fig. 1I).
293
ICP-MS was employed to investigate changes in elemental concentrations in the roots and shoots
of sorghum plants. Fe concentration was significantly different between treatments, exhibiting a
significant decline in the roots and shoots of drought-exposed sorghum (Table 1). Zn level
significantly reduced in both root and shoot with drought. Mn levels significantly declined in the
roots but showed no significant alternations in the shoot with drought compared to controls. S
levels significantly declined in both root and shoot with drought compared to controls. Ca
concentration significantly increased in the roots of drought-imposed plants relative to controls
(Table 1).
302
Changes in leaf gas-exchange parameters and root phytohormone
We measured the gas-exchange parameters in the leaves of sorghum cultivated under different
treatments. In this study, drought caused a significant decrease in the photosynthetic rate,
transpiration rate, stomatal conductance, and water use efficiency (intrinsic) compared to well-
watered controls (Table 2). Furthermore, drought-exposed plants showed a significant decrease
in IAA concentrations in roots relative to controls (Table 2). Sorghum plants exposed to drought
showed a significant increase in ABA concentrations in roots in contrast to control plants with
added Fe boosting the ABA level even higher. Relative to well-watered controls, drought caused
a significant decline in JA concentrations in the roots of sorghum (Table 2). In this study, GA3
levels showed no significant changes in the roots of sorghum cultivated under different
treatments of drought and Fe supplementation (Table 2).
314
RNA-seq analysis of differentially expressed genes in roots of sorghum
Transcript abundances in sorghum roots were seen to be quite different for the two treatments
(Fig. 2). Principal component analysis (PCA) based on three biological replicates revealed that
the first two principal components, PC1 and PC2, contributed 51.5% and 19.2% of the total
variance, respectively (Fig. 2A). Control plants clustered far from the drought-exposed samples.
The analysis of differentially expressed genes (DEGs) in sorghum roots showed significantly
different gene activity in response to two different conditions (Fig. 2B). There were 640

upregulated (2B) and 744 downregulated (2C) genes in 'drought vs. control' comparison (Fig. 2B).

We further annotated the functions of DEGs between control and drought using the Phytozome (V. 13) database for sorghum. We have categorized several differentially expressed ($P < 0.05$) genes known to be associated with drought and stress responses in earlier plant studies for more detailed investigation. Drought imposed to plants caused a significant upregulation of several crucial genes related to mineral homeostasis (*Sobic.005G004500 - Ferritin 1*; *Sobic.008G190200 - Nitrate transporter 1*; *Sobic.002G188600 - Potassium transporter 12-related*; *Sobic.004G093800 - Iron dehydrogenase*; *Sobic.009G168700 - Ammonium transporter*), osmotic adjustment (*Sobic.007G152800 - Glutamate decarboxylase 2*; *Sobic.009G075900 - Aquaporin transporter*; *Sobic.009G116700 - Dehydrin*; *Sobic.002G075800 - Raffinose synthase*; *Sobic.001G357800 - Amino acid transporter*), hormone signaling (*Sobic.003G203500 - Aldehyde dehydrogenase-related*; *Sobic.004G189100 - Ethylene-insensitive protein 3*; *Sobic.003G300800 - Gibberellin 2-oxidase*; *Sobic.001G508200 - ABC transporter C family member 5*; *Sobic.004G345600 - Auxin-responsive protein IAA10-related*) and *redox regulation (Sobic.003G250600 - Glutaredoxin*; *Sobic.010G092500 - Glutathione S-transferase*; *Sobic.006G216300 - Glutathione S-transferase U9*; *Sobic.002G188600 - Potassium transporter 12-related*; *Sobic.003G104200 - Oxidoreductase*) in the roots (Fig. 2C). In addition, several genes associated with mineral uptake (*Sobic.001G523300 - Heavy metal-associated protein*; *Sobic.007G075700 - Zinc/iron transporter*; *Sobic.005G183900 - Sulfate transporting ATPase*; *Sobic.006G244000 - Sulfate transporter*; *Sobic.001G167100 - Cation transporting ATPase*), hormone signaling (*Sobic.003G379500 - Gibberellin 20 oxidase 2*; *Sobic.004G051900 - Auxin response factor*; *Sobic.009G085100 - Auxin-responsive protein IAA*; *Sobic.001G481400 - Ethylene-responsive TF0171*; *Sobic.003G385900 - Lipooxygenase*), osmotic and redox regulation (*Sobic.002G159900 - Solute carried family 35*; *Sobic.003G031900 - Glutaredoxin*; *Sobic.006G150100 - Aquaporin transporter*; *Sobic.004G295100 - Aquaporin transporter*; *Sobic.004G351200 - Aquaporin transporter*), and symbiotic signaling (*Sobic.004G220300 - Phenylalanine ammonia-lyase*; *Sobic.004G220400 - Phenylalanine ammonia-lyase*; *Sobic.001G360700 - Flavonoid 3',5'-hydroxylase*; *Sobic.002G000400 - Isoflavonoid synthase*; *Sobic.008G030100 - Chalcone-flavonone isomerase*) were found to be downregulated in response to drought (Fig. 2D).

We have analyzed the biological significance of DEGs in roots of sorghum in 'control vs drought' conditions. Several pathways with genes that were highly enriched in representation within the upregulated DEGs included, but were not limited to, those involved in various biological processes (response to hydrogen peroxide, protein complex oligomerization, response to heat, response to ROS, response to osmotic stress, response to abscisic acid, response to abiotic stimulus, response to oxygen-containing compound) and KEGG pathways (galactose metabolism, carotenoid biosynthesis, MAPK signaling pathway, amino sugar and nucleotide sugar metabolism, metabolic pathways) in sorghum (Fig. 3A). Conversely, downregulated genes were enriched in pathways related to biological processes such as nucleosome assembly, chromatin organization, cellular protein-containing complex assembly, peptide biosynthetic process, cellular component biogenesis, organic substance biosynthetic process) and various KEGG pathways (cyanoamino acid metabolism, ribosome, phenylalanine metabolism,

367 glutathione metabolism, phenylpropanoid biosynthesis, biosynthesis of amino acid, biosynthesis
368 of secondary metabolites) (Fig. 3B).
369
Changes in diversity and composition of bacterial and fungal communities
Total bacterial and fungal communities in sorghum roots were analyzed by ribosomal DNA
amplicon sequencing. PERMANOVA analysis of bacterial community composition revealed that
drought significantly altered the composition of the bacterial ($F = 6.3682$, df = 1, p = 0.03) and
fungal ($F = 7.7593$, df = 1, p = 0.03) communities (Fig. 4A). The experimental drought treatment
was found to significantly alter the observed OTU richness for bacteria ($F = 72.28$, $df = 1$, $p =$
0.000145) but not for fungi ($F = 3.667$, $df = 1$, $p = 0.104$) in the roots (Fig 4B). Similarly, the
Shannon evenness of bacterial ($F = 324$, $df = 1$, $p = 1.89e-06$) communities showed a significant
change, while fungi ($F = 1.829$, $df = 1$, $p = 0.225$) remained unchanged (Fig. 4B).
379
Differential abundance tests were performed at the genus level for the top ten most abundant
genera to balance the specificity of taxa interactions, statistical power of the analysis, and
redundancy, as the majority of OTUs in both datasets are single representatives of their genera.
Four of the ten most abundant bacterial genera were found to be differentially abundant between
treatments. In particular, *Burkholderia-Caballeronia-Paraburkholderia* was in very low
abundance in the control samples and increased through drought. However, the relative
abundance of *Herbaspirillum*, *Rhizobium,* and *Sphingomonas* were negatively affected by the
drought treatment, exhibiting lower relative abundances than control samples (Fig. 4C). We
found that *Azospirillum*, *Streptomyces*, *Asticcacaulis*, *Massilia*, *Dyella,* and *Ralstonia* remained
unchanged due to drought. In the case of fungi, only *Malassezia* showed a significant increase
while the other nine (*Exidia*, *Sarocladium*, *Bjerkandera*, *Paecilomyces*, *Scedosporium*,
Talaromyces, *Thermothielavioides*, *Chromelosporium* and *Fusarium*) among the top ten
representative genera showed no significant shift in the roots due to drought (Fig. 4C).
393
There was no effect of drought on the general functional composition of either bacteria ($p =$
0.058) or fungi (p = 0.942). Random forest models were not able to find functional predictors for
fungi (an error rate of 88%) but were able to find functional predictors for bacteria (an error rate
of 12.5%). We also identified the functional groups associated with bacterial communities under
control or drought conditions, predicted using either the MetaCyc (M) database or the
FAPROTAX (F) annotation (Table 3). Most groups, such as chorismate metabolism, glycolysis,
L-lysine biosynthesis II, and protochatechuate degradation I, were associated with the control
condition. In contrast, groups like clavulanate biosynthesis, phototrophy, and photosynthetic
cyanobacteria were specifically linked to drought conditions. While the majority of functional
predictions were derived from the MetaCyc database, FAPROTAX contributed to identifying
additional drought-associated groups, including methyltrophy and nitrate reduction. This
highlights the differential functional responses of bacterial communities under varying
environmental conditions (Table 3).
407
408 **Discussion**
409 Drought stress can significantly impact sorghum growth and development. Understanding
410 transcriptional and microbial shifts during the vegetative stage is crucial to conceptualizing
411 strategies to enhance drought resilience in this vital crop. In this study, sorghum plants cultivated
412 under drought conditions exhibited a significant improvement in plant biomass, leaf chlorophyll

score, and relative water content when supplemented with exogenous Fe in the soil. Drought stress causes considerable damage to photosynthetic pigments, leading to chlorophyll degradation, which is considered a symptom of oxidative stress that causes growth retardation in plants (Ma et al., 2020; Rahdari et al., 2012). The reduced chlorophyll content under drought conditions may trigger the inactivation of photosynthesis and decreased transpiration efficiency in plants (Fahad et al., 2017). We also observed a decline in gas exchange parameters in sorghum exposed to drought. Of the gas-exchange parameters, stomatal conductance is most significantly affected by low water availability, because drought-stressed plants typically close their stomata to conserve water (Marchin et al., 2022; Farooq et al., 2009). While this response helps conserve water, it can also affect other physiological processes, such as photosynthesis and overall plant growth. In addition, drought conditions lead to a reduction in water use efficiency and transpiration rate, ultimately resulting in increased leaf and canopy temperatures (Turner et al., 2001). The morphological and physiological changes are coupled with the nutrient relations of plants exposed to drought stress (Ge et al., 2012). Furthermore, the composition and activity of soil microbial colonies are adversely impacted by soil water deficit, ultimately disrupting plant nutrient relations (Schimel et al., 2007). Previous studies showed that plants modify the length and surface area of their roots, as well as alter their architectural structure, to enhance the capture of less mobile nutrients (Lynch and Brown 2001). In this study, the decline of several nutrient elements in the roots of drought-exposed sorghum, particularly Fe and Mn may aggravate the reduction in bacterial richness as evident by the Observed richness in ASV. Previous studies have also shown that drought delays microbiome development and alters root metabolite profiles (Xu et al., 2018; Caddell et al., 2023).

The differential expression of genes associated with mineral homeostasis, hormone signaling, redox regulation, and osmotic adjustment in sorghum under drought stress reflects the plant's multifaceted response to mitigate stress-induced damage and maintain growth. The upregulation of *Ferritin 1* (*Sobic.005G004500*), *Nitrate transporter 1* (*Sobic.009G119200*), and *Potassium transporter 12-related* (*Sobic.002G168800*) under drought conditions indicates that sorghum actively manages nutrient homeostasis to sustain metabolic activity and protect cellular structures. Ferritin, a key Fe-storage protein, reduces oxidative stress by sequestering excess Fe, thereby decreasing ROS formation (Ravet et al., 2009). Similarly, enhanced nitrate and potassium transporter expression suggests increased nutrient acquisition to counteract drought-induced nutrient limitations (Xu et al., 2021). Furthermore, drought significantly altered hormone signaling, as evidenced by the upregulation of *Ethylene-insensitive protein 3* (*Sobic.004G181900*) and genes involved in gibberellin and auxin pathways, such as *Gibberellin 2-oxidase* (*Sobic.003G309800*) and *Auxin-responsive protein* (*Sobic.004G345600*). In addition, the modulation of auxin-related genes further highlights the importance of maintaining root architecture to optimize water and nutrient uptake (Sharma et al., 2020). Ethylene-insensitive proteins may also play a role in modulating stress responses by altering downstream transcription factors, as demonstrated in Arabidopsis under abiotic stress (Sakamoto et al., 2008). The interaction of auxin with ABA and ethylene often shifts the response toward stomatal closure as drought stress progresses (Lu et al., 2019). Consistently, our LC-MS analyses revealed the induction of ABA in the roots of sorghum supplemented under drought conditions. In the apoplast of *Arabidopsis thaliana*, ABA acts as a signal to induce stomatal closure by binding to receptors on guard cells, triggering a cascade of ion fluxes and turgor loss that minimize water loss during drought (Kuromori et al., in 2010 and 2014). Furthermore, *gibberellin 2-oxidase*,

responsible for gibberellin degradation, may suppress growth-related processes to conserve resources, consistent with findings from drought studies on cereal crops (Shohat et al., 2021a). Drought-induced deactivation of gibberellins in guard cells promotes stomatal closure during the initial stages of drought in tomatoes (Shohat et al., 2021b). Interestingly, the levels of jasmonic acid significantly decreased in the roots of sorghum due to drought. Since plants prioritize drought survival strategies, such as maintaining water balance and minimizing cellular damage, the biosynthesis of jasmonic acid, which requires significant metabolic energy and carbon skeletons, might be downregulated to conserve resources in drought-exposed sorghum. Furthermore, drought stress strongly induces ABA in plants, as evidenced by our study. However, ABA often antagonizes JA signaling, potentially reducing jasmonic acid levels in plants, in response to biotic and abiotic stresses (Anderson et al., 2004). Our findings are consistent with the role of ABA as the primary drought-responsive hormone, though JA's role in modulating root architecture and defense mechanisms under limited water availability warrants further exploration in sorghum.

Genes involved in redox regulation, including *Glutaredoxin* (*Sobic.003G259600*) and multiple *Glutathione S-transferases* (GSTs), were highly upregulated under drought stress. These genes play crucial roles in mitigating oxidative damage caused by ROS accumulation during water deficit (Chen et al., 2012). GSTs are known to detoxify harmful by-products of oxidative stress, such as lipid peroxidation products, and their induction is a hallmark of drought tolerance (Nahar et al., 2016). The expression of redox-regulating genes demonstrates the plant's investment in maintaining cellular redox balance under drought. We also found the regulation of several genes related to water adjustment in the roots of sorghum exposed to drought. The upregulation of *Aquaporin* (*Sobic.009G075500*), *Dehydrin* (*Sobic.009G161700*), and *Glutamate decarboxylase 2* (*Sobic.007G152800*) suggests sorghum's osmotic adjustment mechanisms to maintain cellular turgor. Aquaporins facilitate water transport across membranes, aiding in water retention during drought (Maurel et al., 2015). Dehydrins, known as late embryogenesis abundant (LEA) proteins, stabilize membranes and proteins under water-deficit conditions (Kosová et al., 2014). The induction of these genes highlights the importance of osmotic balance in maintaining physiological function during stress.

The downregulation of key genes in sorghum under drought stress may highlight the potential vulnerabilities in critical biological pathways, but also points out the possible ways in which an unproductive root (i.e., one that is not taking up water or other nutrients) might be shut down or even sacrificed by the plant in order to concentrate nutrient discovery by the most productive roots. In the category of mineral uptake, the repression of *Zinc transporter* (*Sobic.007G075700*) and *Sulfate transporter ATPase* (*Sobic.006G124400*) indicates impaired nutrient acquisition, which may exacerbate stress-induced metabolic imbalances (Marschner, 2012). Similarly, the downregulation of hormone signaling genes, such as *Gibberellin 20-oxidase 2* (*Sobic.003G375900*) and *Auxin-responsive protein IAA6* (*Sobic.009G051600*), suggests a suppression of growth-related hormonal pathways, potentially leading to inhibited root and shoot development under water deficit conditions (Nelissen et al., 2018; Sharma et al., 2020), which may be vital in order to abandon these roots. Furthermore, the reduced expression of aquaporin transporter genes (*Sobic.003G105100* and *Sobic.004G251600*) likely compromises water transport and cellular turgor maintenance, further contributing to drought-mediated wilting and reduced physiological activity (Maurel et al., 2015). Additionally, the suppression of

Glutaredoxin (*Sobic.003G259500*), a critical redox regulator, implies diminished capacity to mitigate oxidative stress caused by reactive oxygen species (ROS), exacerbating cellular damage (Noctor et al., 2012). Furthermore, the downregulation of genes associated with symbiotic signaling, such as *Phenylalanine ammonia-lyase* (*Sobic.004G220300*) and *Chalcone synthase* (*Sobic.006G103100*), may indicate a disruption in the plant's ability to establish or maintain effective symbiotic relationships under stress conditions, potentially leading to reduced synthesis of key metabolites required for signaling pathways and mutualistic interaction dynamics in drought-exposed sorghum. Studies indicate reduced production of signaling compounds vital for microbe-plant interactions, potentially impairing nutrient acquisition through microbial symbiosis (Mitra et al., 2021; Mhlongo et al. 2018). These findings highlight how drought modulates key genes in symbiotic support, stress regulation, and nutrient relations, emphasizing the context-dependent roles of gene expression in response to drought in sorghum.

Drought stress causes a major compositional change in the rhizosphere and endophytic communities (Xu et al., 2021; Ma et al., 2020). In this study, amplicon sequencing data showed a reduction in both the diversity and richness of bacterial microbial communities, but not of fungi, in the roots of sorghum under drought conditions. In our study, fungal communities were dominated by *Talaromyces* under both experimental treatments. *Talaromyces* are primarily known for their role in the decomposition of organic matter (Heo et al., 2019). We found that *Talaromyces* were substantially enriched under drought, though these conclusions lack firm statistical support. The greenhouse conditions of this study were apparently selected for a narrower range of fungal taxa that had variable abundance responses to drought than seen for bacteria. Fungi are generally agreed to be more drought resistant than bacteria (Sun et al., 2020), and the stochasticity between replicates likely points to their being little selection pressure within this experiment from either the soil environment caused by drought, or from the plant in response to any beneficial associations with this narrow set of fungal taxa. In a recent study (Kaur and Saxena, 2023), *Talaromyces purpureogenus* has been identified as a drought-tolerant endophyte with significant plant growth-promoting attributes. When applied as a bioinoculant to wheat seedlings, *T. purpureogenus* significantly improved various physiological and biochemical growth parameters under both normal and drought-stressed conditions. Similarly, research by Pang et al. (2020) highlighted the drought tolerance of *Talaromyces purpureogenus*, which exhibited improved growth in rice plants under 10% polyethylene glycol (PEG)-induced drought stress (Pang et al., 2020). In our ITS analysis, we found a significant decrease in the relative abundance of *Malassezia* which is not commonly recognized as a typical endophytic fungus. The imposition of drought is a highly selective force on the fungal microbial community, possibly through encouraging the proliferation of a few putatively tolerant species at the expense of other members of the community or a reduction in those intolerant members leaving only tolerant ones surviving without associated abundance shifts. The nature of amplicon sequencing cannot readily unpick the difference between these scenarios. It should be noted, however, that severe drought is likely to kill some root tissues, and that this necrotic process might be enhanced by a plant looking to concentrate on its most productive roots (Gill and Jackson, 2000). Hence, fungi that remove unproductive roots as part of a general decomposition process might have more available root tissue to decompose, and thus become more prominent, under drought conditions.

When considering the bacterial community drought response, drought conditions appear to favor the proliferation of *Burkholderia-Caballeronia-Paraburkholderia* in the roots of sorghum. The

taxonomy assignment "*Burkholderia-Caballeronia-Paraburkholderia*" reflects the complexities of microbial taxonomy, particularly within the *Burkholderia sensu lato* group, which has undergone significant reclassification in recent years (Bach et al., 2023). However, the genera *Burkholderia*, *Caballeronia*, and *Paraburkholderia* encompass a diverse group of bacteria known for their plant growth-promoting properties, including enhancing drought tolerance (Ahmad et al., 2022). *Paraburkholderia phytofirmans* strain PsJN produces phytohormones such as indole-3-acetic acid (IAA), which stimulate root elongation and branching, thereby improving water uptake under drought conditions (Donoso et al., 2016). Furthermore, auxin is found to be a key determinant in mycorrhizal symbiotic association with plants (Kabir and Bennetzen, 2024; Ludwig-Müller and Güther, 2007). The genera belonging to the Burkholderiaceae family are known for containing a diverse set of taxa capable of plant symbiosis and functions in plant growth, stress resilience, and nitrogen fixation (Pal et al., 2022; Esmaeel et al., 2020). *Burkholderia* species have been well documented to be associated with an increase in the drought stress resilience of host plants in inoculation studies (Naveed et al., 2014; Naveed et al., 2014; Huang et al., 2017; Tallapragada et al., 2016). Studies have also reported that switchgrass inoculated with *Burkholderia* were more photosynthetically active, allocated more biomass, and produced more tillers than uninoculated plants (Wang et al., 2015; Kim et al., 2012). *Burkholderia* may alleviate drought stress by producing Fe-chelating compounds (siderophores) that help plants survive in environments with low Fe levels (Pal et al., 2022; Barrera-Galicia et al., 2021). Both *Paraburkholderia* and *Burkholderia* have also been identified as inorganic phosphate-solubilizing bacteria (Damo et al., 2022). In light of this evidence, our findings underscore the potential application of *Burkholderia-Caballeronia-Paraburkholderia* species as bioinoculants to enhance drought tolerance in sorghum, offering a sustainable solution to mitigate the adverse impacts of water deficit in soil.

The decline in *Rhizobium*, *Herbaspirillum*, and *Sphingomonas* that we observed under drought conditions is likely due to reduced soil moisture, altered root exudation patterns, and/or oxidative stress, which collectively create a less hospitable environment for these potentially beneficial microbes (Jones et al., 2019; Ricks et al., 2023). Although sorghum roots do not host *Rhizobium* for nitrogen fixation, sorghum may interact with other beneficial microbes in the rhizosphere, such as diazotrophic bacteria that can fix nitrogen and contribute to plant growth. Furthermore, drought-driven decreased accumulation of soil iron by sorghum may disrupt bacterial metabolic processes. These stresses may alter sorghum root exudation patterns, making them less favorable for sustaining beneficial microbes, as we observed in drought-exposed sorghum. *Herbaspirillum* and *Sphingomonas*, recognized for their roles in phytohormone production and nutrient solubilization (Grillo-Puertas et al., 2021; Yang et al., 2014), also exhibited reduced population density in response to drought, thereby potentially impairing their plant growth-promoting functions. Metze et al. (2023) showed that drought inhibited the growth of over 90% of bacterial taxa, but favored the growth of specialized taxa, including Actinomycetota such as *Streptomyces*. In this study, we observed a less significant increase in *Streptomyces* abundance in response to drought stress than seen in previous studies conducted in soil systems (Metze et al., 2023; Xu et al., 2021; Fitzpatrick et al., 2018). As our experiment was conducted in greenhouse pots, and not in the field as per the other referenced studies, some differences in outcomes would be expected because our observed initial community is likely a subset of the field community (Kushwaha et al., 2024). For example, Forero et al. (2019) showed that different plant-soil feedback is observed between greenhouse and field trials even when greenhouse pots are inoculated with field soil.

Our findings suggest that the drought stress we imposed had a minimal impact on the overall functional composition of bacterial and fungal communities, compared to the major physiological and transcriptional changes observed. This aligns with studies that propose microbial functional stability as a mechanism for ecosystem resilience to environmental stress (Evans & Wallenstein, 2014; Griffiths and Philippot, 2013). However, the divergent capacities of bacterial communities to respond functionally under drought were evident in the predictive accuracy of the random forest models. Bacterial functions were effectively predicted (error rate of 12.5%), indicating that bacterial communities may exhibit a more defined or deterministic functional response to drought in sorghum.

While bacterial communities exhibited clear functional shifts under drought conditions, the fungal communities showed minimal changes in diversity and composition, with their functional responses being less distinct. The limited functional predictability for fungal communities, as evidenced by the high error rate in random forest models, may be attributed to their lower diversity under drought conditions and potential functional redundancy among dominant taxa, which could mask distinct functional shifts in response to drought stress. Identifying functional groups associated with bacterial communities highlights their distinct responses under drought and control conditions, showcasing microbial adaptation strategies. Functional groups such as chorismate metabolism, glycolysis, and protochatechuate degradation I was more active in the control conditions, reflecting bacterial contributions to primary metabolism and carbon cycling under stable conditions. Conversely, functional groups like clavulanate biosynthesis, phototrophy, and photosynthetic cyanobacteria were associated with drought, highlighting shifts in bacterial functional strategies to adapt to water scarcity. These results are consistent with findings by Lennon et al. (2012), which suggest that bacterial communities can reconfigure their functional potential to maintain ecosystem processes under drought stress. The integration of the MetaCyc and FAPROTAX databases provided complementary insights into these functional dynamics. While MetaCyc predicted most groups, FAPROTAX identified additional drought-associated functions such as methyltrophy and nitrate reduction. These functions are critical for nutrient cycling and drought resilience, as methylotrophic activity and nitrate reduction can mitigate resource limitations (Torres Vera et al., 2024; Deng et al., 2024). These results highlight the importance of combining multiple annotation databases to obtain a comprehensive understanding of microbial functional shifts under environmental stress. Overall, the results suggest that bacterial communities exhibit greater functional plasticity in response to drought stress compared to fungal communities. This finding emphasizes the need for future studies to explore the underlying mechanisms driving these functional responses, such as shifts in microbial taxa, substrate availability, and resource competition.

Conclusion
This study underscores the complex responses of sorghum exposed to drought stress at both the transcriptional and microbial levels. The upregulation of genes related to nutrient homeostasis, hormone signaling, and redox balance demonstrates the plant's ability to activate pathways essential for maintaining physiological processes under stress. Drought-induced genes such as *Ferritin 1*, *Fe dehydrogenase* and *Nitrate transporter 1* play critical roles in mitigating oxidative damage and ensuring nutrient uptake, while aquaporins and dehydrins highlight the importance of osmotic regulation in sustaining cellular function during water deficit. These molecular responses reflect the adaptive strategies of sorghum to survive under drought conditions. In

parallel, drought significantly shaped the root-associated microbiome, favoring specific bacterial taxa like *Burkholderia-Caballeronia-Paraburkholderia*, known for their contributions to plant growth through auxin production, phosphate solubilization, and Fe chelation. The enrichment of functional groups such as phototrophy and nitrate reduction further illustrates the role of microbiome dynamics in supporting nutrient availability and plant resilience. These findings highlight the potential of leveraging both plant genetics and beneficial microbial consortia to enhance sorghum's drought tolerance. Future efforts should prioritize field-based validation and the development of bioinoculants to support sustainable agricultural practices in water-limited environments.

Acknowledgments

We express our gratitude to the Georgia Genomics and Bioinformatics Core (GGBC) and the Mass Spectrometry and Metabolomics Core of Michigan State University. This research received support from the US Department of Energy (grant DE-SC0021386), Georgia Research Alliance, and a startup grant (5SFAES-293007) from the University of Louisiana at Monroe.

Competing interest

We have no conflict of interest.

Author contributions

AHK conceived the idea for the study, conducted physiological assays, RNA-seq analysis, split-root assay, and prepared the draft manuscript. PBJ prepared the DNA library, analyzed the amplicon sequencing data, and wrote the microbial part of the results. MA crosschecked the RNA-seq data, performed gene enrichment analysis, and reviewed the discussion. JL carried out the bioinformatics analysis of microbial networks and functional groups. BA interpreted the microbial functional analysis section and critically reviewed the manuscript. SPLT provided significant contributions to the manuscript. JLB supervised the work, provided intellectual input, and critically revised the manuscript.

Data availability

Illumina sequencing data was submitted to NCBI under the following accession numbers: PRJNA1128172 (RNA-seq), PRJNA1128337 (16S) and PRJNA1128338 (ITS).

References

Abarenkov, Kessy, R. Henrik Nilsson, Karl-Henrik Larsson, Andy F. S. Taylor, Tom W. May, Tobias Guldberg Frøslev, Julia Pawlowska. "The UNITE Database for Molecular Identification and Taxonomic Communication of Fungi and Other Eukaryotes: Sequences, Taxa and Classifications Reconsidered. *Nucleic Acids Res.* 2024;52 (D1):D791–97.

Ahmad, H. M., Fiaz, S., Hafeez, S., Zahra, S., Shah, A. N., Gul, B., Aziz, O., Mahmood-Ur-Rahman, Fakhar, A., Rafique, M., Chen, Y., Yang, S. H., & Wang, X. (2022). Plant growth-promoting Rhizobacteria eliminate the effect of drought stress in plants: A Review. Frontiers in plant science, 13, 875774.

Alexieva V, Sergiev, Mapelli S, Karanov E. The effect of drought and ultraviolet radiation on growth and stress markers in pea and wheat. *Plant Cell Env.* 2001:24:1337–1344.

Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics.* 2015:31(2):166–169.

Anderson, J. P., Badruzsaufari, E., Schenk, P. M., Manners, J. M., Desmond, O. J., Ehlert, C., Maclean, D. J., Ebert, P. R., & Kazan, K. (2004). Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in Arabidopsis. The Plant cell, 16(12), 3460–3479.

Antonov M, Csárdi G, Horvát S, Müller K, Nepusz T, Noom D, et al., igraph enables fast and robust network analysis across programming languages [Internet]. arXiv; 2023 [cité 18 sept 2024]. Disponible sur: https://arxiv.org/abs/2311.10260

Anwar S, Ali B, Sajid I. Screening of rhizospheric Actinomycetes for various *in vitro* and *in-vivo* plant growth promoting (PGP) traits and for agroactive compounds. *Front. Microbiol.* 2016:7:1334.

Bach, E., Volpiano, C. G., Sant'Anna, F. H., & Passaglia, L. M. P. (2023). Genome-based taxonomy of Burkholderia sensu lato: Distinguishing closely related species. Genetics and molecular biology, 46(3 Suppl 1), e20230122.

Barrera-Galicia GC, Peniche-Pavía HA, Peña-Cabriales JJ, Covarrubias SA, Vera-Núñez JA, Délano-Frier JP. Metabolic footprints of *Burkholderia Sensu* Lato rhizosphere bacteria active against maize *Fusarium* pathogens. *Microorganisms* 2021:9(10):2061.

Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics* 2014:btu170.

Breiman, L. (2001). Random forests. Machine Learning, 45(1), 5–32. https://doi.org/10.1023/A:1010933404324

Caddell, D.F., Pettinga, D., Louie, K., Bowen, B.P., Sievert, J.A., Hollingsworth, J., Rubanowitz, R., Dahlberg, J., Purdom, E., Northen, T., Coleman-Derr, D.A. 2023. Drought shifts sorghum root metabolite and microbiome profiles and enriches for pipecolic acid. Phytobiomes Journal. 7(4):449-463.

Cantalapiedra, C. P., García-Pereira, M. J., Gracia, M. P., Igartua, E., Casas, A. M., & Contreras-Moreira, B. (2017). Large Differences in Gene Expression Responses to Drought and Heat Stress between Elite Barley Cultivar Scarlett and a Spanish Landrace. Frontiers in plant science, 8, 647.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Met.* 2016:13(7):581–583.

Cheng S, Zou YN, Kuča K, Hashem A, Abd Allah EF, Wu QS. Elucidating the mechanisms underlying enhanced drought tolerance in plants mediated by arbuscular mycorrhizal fungi. *Front Microbiol.* 2021:12:809473.

Chen, J. H., Jiang, H. W., Hsieh, E. J., Chen, H. Y., Chien, C. T., Hsieh, H. L., & Lin, T. P. (2012). Drought and salt stress tolerance of an Arabidopsis glutathione S-transferase U17 knockout mutant are attributed to the combined effect of glutathione and abscisic acid. Plant physiology, 158(1), 340–351.

Coombs JT, Franco CM. Isolation and identification of actinobacteria from surface-sterilized wheat roots. *Appl Environ Microbiol.* 2003:69:5603–5608.

Cortés-Patiño S, Vargas C, Álvarez-Flórez F, Bonilla R, Estrada-Bonilla G. Potential of *Herbaspirillum* and *Azospirillum* consortium to promote growth of perennial ryegrass under water deficit. *Microorganisms.* 2021:9(1):91.

Curá JA, Franz DR, Filosofía JE, Balestrasse KB, Burgueño LE. Inoculation with *Azospirillum* sp. and *Herbaspirillum* sp. bacteria increases the tolerance of maize to drought stress. *Microorganisms.* 2017:5(3):41.

Damo JLC, Ramirez MDA, Agake SI, Pedro M, Brown M, Sekimoto H, Yokoyama T, Sugihara S, Okazaki S, Ohkama-Ohtsu N. Isolation and characterization of phosphate solubilizing bacteria from paddy field soils in Japan. *Microb Environ.* 2022:37(2):ME21085.

Dao TT, Linthorst HJ, Verpoorte R. Chalcone synthase and its functions in plant resistance. *Phytochem Rev.* 2011:10(3):397–412.

de Oliveira JPV, Duarte VP, de Castro EM, Magalhães PC, Pereira FJ. Stomatal cavity modulates the gas exchange of *Sorghum bicolor* (L.) Moench. grown under different water levels. *Protoplasma.* 2022:259(4):1081–1097.

Deng, D., He, G., Ding, B., Liu, W., Yang, Z., & Ma, L. (2024). Denitrification dominates dissimilatory nitrate reduction across global natural ecosystems. Global change biology, 30(3), e17256.

Donoso, R., Leiva-Novoa, P., Zúñiga, A., Timmermann, T., Recabarren-Gajardo, G., & González, B. (2016). Biochemical and Genetic Bases of Indole-3-Acetic Acid (Auxin Phytohormone) Degradation by the Plant-Growth-Promoting Rhizobacterium Paraburkholderia phytofirmans PsJN. Applied and environmental microbiology, 83(1), e01991-16.

Duveiller E, Singh RP, Nicol J.M. The challenges of maintaining wheat productivity: Pests, diseases, and potential epidemics. *Euphytica.* 2007:157:417–430.

Esmaeel Q, Jacquard C, Sanchez L, Clément C, Ait Barka E. The mode of action of plant associated Burkholderia against grey mould disease in grapevine revealed through traits and genomic analyses. *Sci Rep.* 2020:10(1) :19393.

Evans, S. E., & Wallenstein, M. D. (2014). Climate change alters ecological strategies of soil bacteria. Ecology letters, 17(2), 155–164.

Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins S, Saud S, Ihsan MZ, Alharby H, Wu C, Wang D, Huang J. Crop production under drought and heat stress: plant responses and management options. *Front Plant Sci.* 2017:8:1147.

Farooq M, Wahid A, Kobayashi N, Fujita D, Basra SMA. Plant drought stress: effects, mechanisms and management. *Agron Sustain Dev.* 2009:29:185–212.

Forero, Leslie E, Grenzer J, Heinze J, Schittko C, Kulmatiski A. Greenhouse- and field-measured plant-soil feedbacks are not correlated. *Front Environ Sci Eng China* 2019:7.

Feng D, Wang X, Gao J, Zhang C, Liu H, Liu P, Sun X. Exogenous calcium: Its mechanisms and research advances involved in plant stress tolerance. Front Plant Sci. 2023 Mar 21;14:1143963.

Ferrol N, Azcon-Aguilar C, Pérez-Tienda J. Arbuscular mycorrhizas as key players in sustainable plant phosphorus acquisition: an overview on the mechanisms involved. *Plant Sci.* 2019:280:441–447.

Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. Assembly and ecological function of the root microbiome across angiosperm plant species. *PNAS.* 2018:115(6:E1157–E1165.

Fu Y, Yan R, Liu D, Jiang S, Cui L, Guo X, Wang X, Zhang J, Xiang W. *Trinickia diaoshuihuensis* sp. nov., a plant growth promoting bacterium isolated from soil. *Int J Syst Evol Microbiol.* 2019:69(1):291–296.

Ge TD, Sun NB, Bai LP, Tong CL, Sui FG. Effects of drought stress on phosphorus and potassium uptake dynamics in summer maize (*Zea mays*) throughout the growth cycle. *Acta Physiol Plant.* 2012:34:2179–2186.

Gill, R. A., & Jackson, R. B. (2000). Global patterns of root turnover for terrestrial ecosystems. New Phytologist, 147(1), 13–31.

Goodstei DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, Rokhsar DS. Phytozome: a comparative platform for green plant genomics. *Nucleic Acid Res.* 2012:D1:D1178-D1186.

Griffiths, B. S., & Philippot, L. (2013). Insights into the resistance and resilience of the soil microbial community. FEMS microbiology reviews, 37(2), 112–129.

Grillo-Puertas, M., Villegas, J. M., Pankievicz, V. C. S., Tadra-Sfeir, M. Z., Teles Mota, F. J., Hebert, E. M., Brusamarello-Santos, L., Pedraza, R. O., Pedrosa, F. O., Rapisarda, V. A., & Souza, E. M. (2021). Transcriptional responses of *Herbaspirillum seropedicae* to environmental phosphate concentration. Frontiers in microbiology, 12, 666277.

Heo I, Hong K, Yang H, Lee HB, Choi YJ, Hong SB. Diversity of *Aspergillus*, *Penicillium*, and *Talaromyces* species isolated from freshwater environments in Korea. *Mycobiol.* 2019:47(1):12–19.

Hinder LM, Park M, Rumora AE, Hur J, Eichinger F, Pennathur S, Kretzler M, Brosius FC, Feldma EL. Comparative RNA-Seq transcriptome analyses reveal distinct metabolic pathways in diabetic nerve and kidney disease. *J Cell Mol Med.* 2017:21(9):2140–2152.

Huang X, Zhou D, Lapsansky ER, Reardon KF, Guo J, Andales MJ, Vivanco JM, Manter DK. *Mitsuaria* Sp. and *Burkholderia* Sp. from Arabidopsis rhizosphere enhance drought tolerance in *Arabidopsis thaliana* and maize (*Zea mays* L.). *Plant Soil.* 2017:419(1):523–39.

Hussain HA, Hussain S, Khaliq A, Ashraf U, Anjum SA, Men S, Wang L. Chilling and drought stresses in crop plants: implications, cross talk, and potential management opportunities. *Front Plant Sci.* 2018:9:393.

Iqbal S, Wang X, Mubeen I, Kamran M, Kanwal I, Díaz GA, Abbas A, Parveen A, Atiq MN, Alshaya H, Zin El-Abedin TK, Fahad S. Phytohormones trigger drought tolerance in crop plants: outlook and future perspectives. *Front Plant Sci.* 2022:12:799318.

Iturrieta-González I, Gené J, Wiederhold N, García D. Three new *Curvularia* species from clinical and environmental sources. *MycoKeys.* 2020:68:1–21.

Janků M, Luhová L, Petřivalský M. On the Origin and Fate of Reactive Oxygen Species in Plant Cell Compartments. *Antioxidants.* 2019:8(4):105.

Jones P, Garcia BJ, Furches A, Tuskan GA, Jacobson D. Plant Host-Associated Mechanisms for Microbial Selection. Front Plant Sci. 2019;10:862.

Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, et al., Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. Nat Biotechnol. sept 2013;31(9):814 21.

Liber JA, Bonito G, Benucci GMN. CONSTAX2: improved taxonomic classification of environmental DNA markers. *Bioinformatics.* 2021:37(21):3941-3943.

Nahar K, Hasanuzzaman M, Rahman A, et al., Polyamines Confer Salt Tolerance in Mung Bean (Vigna radiata L.) by Reducing Sodium Uptake, Improving Nutrient Homeostasis, Antioxidant Defense, and Methylglyoxal Detoxification Systems. Front Plant Sci. 2016;7:1104.

Kabir AH, Bennetzen JL. Molecular insights into the mutualism that induces iron deficiency tolerance in sorghum inoculated with *Trichoderma harzianum*. *Microbiol Res.* 2024:281:127630.

Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1 janv 2000;28(1):27 30.

Karp PD. The MetaCyc Database. Nucleic Acids Res. 1 janv 2002;30(1):59 61.

Kaur, R., & Saxena, S. (2023). Evaluation of drought-tolerant endophytic fungus Talaromyces purpureogenus as a bioinoculant for wheat seedlings under normal and drought-stressed circumstances. Folia microbiologica, 68(5), 781–799.

Khan A, Pan X, Najeeb U, Tan DK, Fahad S, Zahoor R, Luo H. Coping with drought: Stress and adaptive mechanisms, and management through cultural and molecular alternatives in cotton as vital constituents for plant stress resilience and fitness. *Biol Res.* 2018:51:47.

Khan, M. I. R., Nazir, F., Maheshwari, C., Chopra, P., Chhillar, H., & Sreenivasulu, N. (2023). Mineral nutrients in plants under changing environments: A road to future food and nutrition security. The plant genome, 16(4), e20362.

Kim S, Lowman S, Hou G, Nowak J, Flinn B, Mei C. Growth promotion and colonization of switchgrass (*Panicum virgatum*) cv. Alamo by bacterial endophyte *Burkholderia phytofirmans* strain PsJN. *Biotechnol Biofuels.* 2012:5(1):37.

King, E., Wallner, A., Guigard, L., Rimbault, I., Parrinello, H., Klonowska, A., Moulin, L., & Czernic, P. (2023). Paraburkholderia phytofirmans PsJN colonization of rice endosphere triggers an atypical transcriptomic response compared to rice native Burkholderia s.l. endophytes. Scientific reports, 13(1), 10696.

Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g: Profiler—interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). *Nucl Acids Res.* 2023:gkad347.

Krivonos DV, Konanov DN, Ilina EN. FUNFUN☐: ITS ☐based functional annotator of fungal communities. Ecol Evol. mars 2023;13(3):e9874.

Kuromori T, Shinozaki K. ABA transport factors found in Arabidopsis ABC transporters. Plant *Signal Behav.* 2010:5:1124–1126.

Kuromori T, Sugimoto E, Shinozaki K. Intertissue signal transfer of abscisic acid from vascular cells to guard cells. *Plant Physiol.* 2014:164:1587–1592.

Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. Sparse and Compositionally Robust Inference of Microbial Ecological Networks. Von Mering C, éditeur. PLOS Comput Biol. 7 mai 2015;11(5):e1004226.

Kushwaha P, Soto Velázquez AL, McMahan C, Neilson JW. Field to Greenhouse: How Stable Is the Soil Microbiome after Removal from the Field? *Microorganisms.* 2024:12(1):110.

Lamaoui M, Jemo M, Datla R, Bekkaoui F. Heat and drought stresses in crops and approaches for their mitigation. *Front Chem.* 2018:6:26.

Law CW, Alhamdoosh M, Su S, Dong X, Tian L, Smyth GK, Ritchie ME. RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Research* 2016:5:J-1408.

Lennon, J. T., Aanderud, Z. T., Lehmkuhl, B. K., & Schoolmaster, D. R., Jr (2012). Mapping the niche space of soil microorganisms using taxonomy and traits. Ecology, 93(8), 1867–1879. https://doi.org/10.1890/11-1745.1

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2-\Delta\Delta CT$ method. *Methods* 2001:25:402–408.

Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean microbiome. Science. 16 sept 2016;353(6305):1272 7.

Lozano YM, Aguilar-Trigueros CA, Roy J, Rillig MC. Drought induces shifts in soil fungal communities that can be linked to root traits across 24 plant species. *New Phytol.* 2021:232(5):1917–1929.

Lu, C., Chen, M. X., Liu, R., Zhang, L., Hou, X., Liu, S., Ding, X., Jiang, Y., Xu, J., Zhang, J., Zhao, X., & Liu, Y. G. (2019). Abscisic Acid Regulates Auxin Distribution to Mediate Maize Lateral Root Development Under Salt Stress. Frontiers in plant science, 10, 716.

Ludwig-Müller J, Güther M. Auxins as signals in arbuscular mycorrhiza formation. *Plant Signal Behav.* 2007:2(3):194-6.

Ma Y, Dias MC, Freitas H. Drought and salinity stress responses and microbe-induced tolerance in plants. *Front Plant Sci.* 2020:11:591911.

Marchin RM, Backes D, Ossola A, Leishman MR, Tjoelker MG, Ellsworth DS. Extreme heat increases stomatal conductance and drought-induced mortality risk in vulnerable plant species. *Glob Chang Biol.* 2022:28(3):1133-1146.

Mathur, S., Umakanth, A. V., Tonapi, V. A., Sharma, R., & Sharma, M. K. (2017). Sweet sorghum as biofuel feedstock: recent advances and available resources. Biotechnology for biofuels, 10, 146.

Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 2011:17(1):10–12.

McCormick RF, Truong SK, Sreedasyam A, Jenkins J, Shu S, Sims D, Kennedy M, Amirebrahimi M, Weers BD, McKinley B, Mattison A, Morishige DT, Grimwood J, Schmutz J, Mullet JE. The Sorghum bicolor reference genome: improved assembly, gene annotations, a transcriptome atlas, and signatures of genome organization. *Plant J.* 2018:93(2) :338–354.

McDonald D, Jiang Y, Balaban M, Cantrell K, Zhu Q, Gonzalez A, Morton JT, Nicolaou G, Parks DH, Karst SM, Albertsen M, Hugenholtz P, DeSantis T, Song SJ, Bartko A, Havulinna AS, Jousilahti P, Cheng S, Inouye M, Niiranen T, Jain M, Salomaa V, Lahti L, Mirarab S, Knight R. Greengenes2 unifies microbial data in a single reference tree. *Nat Biotechnol.* 2024:42(5):715-718..

McMurdie P.J, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS One* 2013:8(4):e61217.

Metsalu T, Vilo J. ClustVis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Res.* 2015:43(W1):W566-70.

Metze D, Schnecker J, Canarini A, Fuchslueger L, Koch BJ, Stone BW, Hungate BA, Hausmann B, Schmidt H, Schaumberger A, Bahn M, Kaiser C, Richter A. Microbial growth under drought is confined to distinct taxa and modified by potential future climate conditions. *Nat Commun.* 2023:14(1):5895.

Mhlongo, M. I., Piater, L. A., Madala, N. E., Labuschagne, N., & Dubery, I. A. (2018). The Chemistry of Plant-Microbe Interactions in the Rhizosphere and the Potential for Metabolomics to Reveal Signaling Related to Defense Priming and Induced Systemic Resistance. Frontiers in plant science, 9, 112.

Mona SA, Hashem A, Allah EFA, Alqarawi AA, Soliman DWK, Wirth S, Egamberdieva D.. Increased resistance of drought by *Trichoderma harzianum* fungal treatment correlates with increased secondary metabolites and proline content. Journal of Integrative *Agriculture.* 2017:16:1751-1757.

Naveed M, Hussain MB, Zahir ZA, Mitter B, Sessitsch A. Drought stress amelioration in wheat through inoculation with *Burkholderia phytofirmans* Strain PsJN. *Plant Growth Regul.* 2014:73(2):121–31.

Naveed M, Mitter B, Reichenauer TG, Wieczorek K, Sessitsch A. Increased drought stress resilience of maize through endophytic colonization by *Burkholderia phytofirmans* PsJN and *Enterobacter* Sp. FD17." *Environ Exp Bot.* 2014:97:30–39.

Nilsson RH, Larsson KH, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, Kennedy P, Picard K, Glöckner FO, Tedersoo L, Saar I, Kõljalg U, Abarenkov K. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* 2019:47(D1):D259-D264.

Pal G, Saxena S, Kumar K, Verma A, Sahu PK, Pandey A, White JF, Verma SK. Endophytic Burkholderia: Multifunctional roles in plant growth promotion and stress tolerance. *Microbiol Res.* 2022:265:127201.

Pang, Z., Zhao, Y., Xu, P., & Yu, D. (2020). Microbial diversity of upland rice roots and their influence on rice growth and drought tolerance. Microorganisms, 8(9), 1329.

Palaniyandi SA, Damodharan K, Yang SH, Suh JW. *Streptomyces* sp. strain PGPA39 alleviates salt stress and promotes growth of' Micro Tom' tomato plants. *J. Appl. Microbiol.* 2014:117:766–773.

Pekker I, Tel-Or E, Mittler R. Reactive oxygen intermediates and glutathione regulate the expression of cytosolic ascorbate peroxidase during iron-mediated oxidative stress in bean. *Plant Mol Biol.* 2002:49:429–438.

Pinto MR, Rodrigues ML, Travassos LR, Haido RM, Wait R, Barreto-Bergter E. Characterization of glucosylceramides in *Pseudallescheria boydii* and their involvement in fungal differentiation. *Glycobiol.* 2002:12(4) :251–260.

Poudel M, Mendes R, Costa LAS, Bueno CG, Meng Y, Folimonova SY, Garrett KA, Martins S. J. The role of plant-associated bacteria, fungi, and viruses in drought stress mitigation. *Front Microb.* 2021:12:743512.

Prasad, Jay Kishor, Priyanka Pandey, Richa Anand, and Richa Raghuwanshi. Drought exposed *Burkholderia seminalis* JRBHU6 exhibits antimicrobial potential through pyrazine-1,4-dione derivatives targeting multiple bacterial and fungal proteins. *Front Microbiol.* 2021:12:633036.

Qi, M., Berry, J. C., Veley, K. M., O'Connor, L., Finkel, O. M., Salas-González, I., Kuhs, M., Jupe, J., Holcomb, E., Glavina Del Rio, T., Creech, C., Liu, P., Tringe, S. G., Dangl, J. L., Schachtman, D. P., & Bart, R. S. (2022). Identification of beneficial and detrimental bacteria impacting sorghum responses to drought using multi-scale and multi-system microbiome comparisons. The ISME journal, 16(8), 1957–1969.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 2012:41(D1) :D590–D596.

Rahdari P, Hoseini SM, Tavakoli S. The studying effect of drought stress on germination, proline, sugar, lipid, protein and chlorophyll content in purslane (*Portulaca oleraceae* L.) leaves. *J Med Plant Res.* 2012:6:1539–1547.

Raheem A, Shaposhnikov A, Belimov AA, Dodd IC, Ali B. Auxin production by rhizobacteria was associated with improved yield of wheat (*Triticum aestivum* L.) under drought stress. *Arch Agron Soil Sci.* 2018):64, 574–587.

Rajarajan, K., Ganesamurthy, K., Raveendran, M., Jeyakumar, P., Yuvaraja, A., Sampath, P., Prathima, P. T., & Senthilraja, C. (2021). Differential responses of sorghum genotypes to drought stress revealed by physio-chemical and transcriptional analysis. Molecular biology reports, 48(3), 2453–2462.

Rasheed, S., Bashir, K., Matsui, A., Tanaka, M., & Seki, M. (2016). Transcriptomic analysis of soil-grown *Arabidopsis thaliana* roots and shoots in response to a drought stress. Frontiers in plant science, 7, 180.

Rajarajan K, Ganesamurthy K, Raveendran M, Jeyakumar P, Yuvaraja A, Sampath P, Prathima P. T, Senthilraja C. Differential responses of sorghum genotypes to drought stress revealed by physio-chemical and transcriptional analysis. *Mol Biol Rep.* 2021:48(3) :2453–2462.

Rapparini F, Peñuelas J. Mycorrhizal fungi to alleviate drought stress on plant growth," in use of microbes for the alleviation of soil stresses, ed. M. Miransari (New York, NY: Springer), 2014:21–42.

Ricks, K. D., & Yannarell, A. C. (2023). Soil moisture incidentally selects for microbes that facilitate locally adaptive plant response. Proceedings. Biological sciences, 290(2001), 20230469.

Ravet K, Touraine B, Boucherez J, Briat JF, Gaymard F, Cellier F. Ferritins control interaction between iron homeostasis and oxidative stress in *Arabidopsis*. *Plant J.* 2009:57:400–412.

Ruehr NK, Grote R, Mayr S, Arneth A. Beyond the extreme: Recovery of carbon and water relations in woody plants following heat and drought stress. *Tree Physiol.* 2019:39:1285–1299.

Sakamoto W., Miyagishima S.-Y., Jarvis P. Chloroplast Biogenesis, Control of plastid development, protein import, division and inheritance. The Arabidopsis book. 0110. 2008.

Sadok W, Schoppach R. Potential involvement of root auxins in drought tolerance by modulating nocturnal and daytime water use in wheat. *Ann Bot.* 2019:124(6):969–978.

Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its implications for ecosystem function. *Ecol.* 2007:88:1386–1394.

Schonfeld MA, Johnson RC, Carver BF, Mornhinweg DW. Water relations in winter wheat as drought resistance indicators. *Crop Sci.* 1988:28:526-531.

Scudeletti D, Crusciol CAC, Bossolani JW, Moretti LG, Momesso L, Servaz TB, de Castro SGQ, De Oliveira EF, Hungria M. *Trichoderma asperellum* inoculation as a tool for attenuating drought stress in sugarcane. *Front Plant Sci.* 2021:12:645542.

Seleiman MF, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid HH, Battaglia ML. Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants.* 2021:10(2):259.

Shohat H, Eliaz NI, Weiss D. Gibberellin in tomato: metabolism, signaling and role in drought responses. Mol Hortic. 2021a;1(1):15.

Shohat, H., Cheriker, H., Kilambi, H. V., Illouz Eliaz, N., Blum, S., Amsellem, Z., Tarkowská, D., Aharoni, A., Eshed, Y., & Weiss, D. (2021). Inhibition of gibberellin accumulation by water deficiency promotes fast and long-term 'drought avoidance' responses in tomato. The New phytologist, 232(5), 1985–1998.

Soltys-Kalina D, Plich J, Strzelczyk-Żyta D, Śliwka J, Marczewski W. The effect of drought stress on the leaf relative water content and tuber yield of a half-sib family of 'Katahdin'-derived potato cultivars. *Breeding Sci.* 2016:66(2) :328–331.

Sinha R, Irulappan V, Mohan-Raju B, Suganthi A, Senthil-Kumar M. Impact of drought stress on simultaneously occurring pathogen infection in field-grown chickpea. *Sci Rep.* 2019:9(1):5577.

Stres B, Danevcic T, Pal L, Fuka MM, Resman L, Leskovec S, Hacin J, Stopar D, Mahne I, Mandic-Mulec I. Influence of temperature and soil water content on bacterial, archaeal and denitrifying microbial communities in drained fen grassland soil microcosms. *FEMS Microbiol Ecol.* 2008:66(1):110–122.

Sun Y, Han YHC, Long J, Cuiting W, Ruiting Z, Honghua R, Jinyan Y. Drought stress induced increase of fungi: bacteria ratio in a poplar plantation. *Catena* 2020:193:104607.

Tack J, Lingenfelser J, Jagadish SVK. Disaggregating sorghum yield reductions under warming scenarios exposes narrow genetic diversity in US breeding programs. *PNAS.* 2017:114(35), 9296–9301.

Tallapragada P, Rashmi D, Swetha S. Influence of *Rhizophagus* Spp. and *Burkholderia seminalis* on the growth of tomato (*Lycopersicon Esculatum*) and bell pepper (*Capsicum Annuum*) under drought stress. *Commun Soil Sci Plant Anal.* 2016:(17): 975–84.

Takahashi F, Kuromori T, Urano K, Yamaguchi-Shinozaki K, Shinozaki K. Drought stress responses and resistance in plants: from cellular responses to long-distance intercellular communication. *Front Plant Sci.* 2020:11:556972.

Torres Vera R, Bernabé García AJ, Carmona Álvarez FJ, Martínez Ruiz J, Fernández Martín F. Application and effectiveness of Methylobacterium symbioticum as a biological inoculant in maize and strawberry crops. Folia Microbiol (Praha). 2024;69(1):121-131.

Theil EC, Matzapetakis M, Liu X. Ferritins: iron/oxygen biominerals in protein nanocages. *J Biol Inorg Chem.* 2006:11:803–810.

Timmusk S, Timmusk K, Behers L. Rhizobacterial plant drought stress tolerance enhancement: towards sustainable water resource management and food security. *J Food Secur.* 2013:1:6–9.

Turner NC, Wright GC, Siddique KHM. Adaptation of grain legumes (pulses) to water-limited environments. *Adv Agron.* 2001:71 193–231.

Varoquaux, N., Cole, B., Gao, C., Pierroz, G., Baker, C. R., Patel, D., Madera, M., Jeffers, T., Hollingsworth, J., Sievert, J., Yoshinaga, Y., Owiti, J. A., Singan, V. R., DeGraaf, S., Xu, L., Blow, M. J., Harrison, M. J., Visel, A., Jansson, C., Niyogi, K. K., … Purdom, E. (2019). Transcriptomic analysis of field-droughted sorghum from seedling to maturity reveals biotic and metabolic responses. Proceedings of the National Academy of Sciences of the United States of America, 116(52), 27124–27132.

Waadt R, Seller CA, Hsu PK, Takahashi Y, Munemasa S, Schroeder JI. Plant hormone regulation of abiotic stress responses. *Nat Rev Mol Cell Biol.* 2022:23(10) :680–694.

Wang B, Mei C, Seiler JR. Early growth promotion and leaf level physiology changes in *Burkholderia phytofirmans* strain PsJN inoculated switchgrass. *Plant Physiol Biochem.* 2015:86:16–23.

Xu L, Coleman-Derr D. Causes and consequences of a conserved bacterial root microbiome response to drought stress. *Curr. Opin. Microbiol.* 2019:49:1–6.

Xu L, Dong Z, Chiniquy D, Pierroz G, Deng S, Gao C, Diamond S, Simmons T, Wipf HM, Caddell D, Varoquaux N, Madera MA, Hutmacher R, Deutschbauer A, Dahlberg JA, Guerinot ML, Purdom E, Banfield JF, Taylor JW, Lemaux PG, Coleman-Derr D. Genome-resolved metagenomics reveals role of iron metabolism in drought-induced rhizosphere microbiome dynamics. *Nat Commun.* 2021:12(1):3209.

Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, Kim YM, Zink EM, Engbrecht KM, Wang Y, Gao C, DeGraaf S, Madera MA, Sievert JA, Hollingsworth J, Birdseye D, Scheller HV, Hutmacher R, Dahlberg J, Jansson C, Taylor JW, Lemaux PG, Coleman-Derr D. Drought delays development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc Natl Acad Sci US.* 2018:115(18): E4284-E4293.

Yang Z, Qiao Y, Konakalla NC, Strøbech E, Harris P, Peschel G, Agler-Rosenbaum M, Weber T, Andreasson E, Ding L. *Streptomyces* alleviate abiotic stress in plant by producing pteridic acids. *Nat Comm.* 2023:14(1) :7398.

Yang, S., Zhang, X., Cao, Z., Zhao, K., Wang, S., Chen, M., & Hu, X. (2014). Growth-promoting Sphingomonas paucimobilis ZJSH1 associated with Dendrobium officinale through phytohormone production and nitrogen fixation. Microbial biotechnology, 7(6), 611–620.

Zboralski A, Filion M. Genetic factors involved in rhizosphere colonization by phytobeneficial *Pseudomonas* spp. *Comp Struc Biotech J.* 2020:18:3539–3554.

Zhang R, Yang P, Liu S, Wang C, Liu J. Evaluation of the Methods for Estimating Leaf Chlorophyll Content with SPAD Chlorophyll Meters. *Remote Sens.* 2022:14:5144.

Zoghi Z, Hosseini SM, Kouchaksaraei MT, Kooch Y, Guidi L. The effect of biochar amendment on the growth, morphology and physiology of *Quercus castaneifolia* seedlings under water-deficit stress. *Eur J For Res.* 2019:138:967–979.

Fig. 1. Changes in morphological and physiological attributes in sorghum cultivated with or without drought and Fe supplementation. Shoot phenotype (A), root phenotype (B), root length (C), root dry weight (D), shoot height (E), shoot dry weight (F), leaf relative water content (G), leaf SPAD score (H) and root H_2O_2 concentration (I) in sorghum cultivated in the absence (control) or presence of drought. The data presented are means with standard deviations ($n = 3$ of individual replicates). Statistically significant differences among the treatments at a $p < 0.05$ level are indicated by different alphabetical letters (a, b) above the bars, where applicable. Plants were cultivated in a growth chamber for 6 weeks.

Fig. 2. Principal component analysis of samples (A), Pie chart of differentially expressed genes (> Log2 fold, p value < 0.05) (B), top upregulated genes (C) and top downregulated genes (D) in the roots of sorghum plants cultivated in controla and drought conditions. A minimum of 20 million reads on average, with more than 33% read quality per sample, were utilized per treatment. Plants were cultivated in a growth chamber for 6 weeks.

Fig. 3. Gene ontology and enrichment analysis of pathways classified as biological process (BP), and KEGG pathways. These pathways or terms within the list of genes differentially (>Log2 fold, p value <0.05) upregulated (A) and downregulated (B) in 'drought vs control' conditions in roots of sorghum.

1088

1089
1090

Fig. 4. Panel A) Non-metric dimensional scaling (NMDS) ordination that displays community
dissimilarity between samples for bacterial and fungal communities among the treatments. Mean
centroids are displayed as diamonds ± standard errors in the NMDS1 and NMDS2 axes to
demonstrate the dispersal of data points around the mean within each treatment group. Individual
sample shapes and colors are associated with experimental treatment (green / circle = control,
yellow / triangle = drought). Panel B) Observed OTU richness and Shannon index values for

bacterial and fungal communities. Mean values ± SE for each treatment is displayed in black, and individual sample values are colored by treatment. Letters denote distinct statistical groupings of treatments determined through post-hoc pairwise testing. Panel C) Stacked barplots displaying the mean relative abundance of the top 10 most abundant genera for bacteria and fungi. The 'Other' category includes both lower abundance genera and taxa that were unclassified at the genus level.

Fig. 5. Shift in hormonal regulation, transcriptome and microbiome dynamics in sorghum exposed to drought. Drought stress enhances abscisic acid (ABA) levels, leading to osmotic regulation through the induction of *Aquaporin* and *Dehydrin*. Simultaneously, jasmonic acid (JA) levels are suppressed, possibly due to its antagonistic interaction with ABA. In response to drought, symbiotic and nutrient-related processes are upregulated, as evidenced by the induction of genes such as *Phenylalanine ammonia-lyase*, *Isoflavonoid synthase*, and *Chalcone-flavonone isomerase* (symbiosis-related), along with *Ferritin 1*, *Fe dehydrogenase*, and *Nitrate transporter 1* (nutrient acquisition-related). The root microbiome, including species such as members of the *Burkholderia-Caballeronia-Paraburkholderia*, potentially contributes to these adaptive responses by supporting nutrient uptake and stress resilience.

Table 1. ICP-MS analysis of nutrients in the root and shoot of sorghum grown in control and drought conditions. Different letters indicate significant differences between the control and high pH conditions at a $P <0.05$ level, based on t-test. The data represents the means ± SD of three independent biological samples ($n = 3$). Plants were cultivated in a growth chamber for 6 weeks.

Elements	Root		Shoot	
	Control	Drought	Control	Drought
Fe (ppm)	395±75[a]	305±61[b]	156±31[a]	96±7.76[b]
Zn (ppm)	47±5.6[a]	40±4.5[a]	62±9.9[a]	59±17.1[a]
Mn (ppm)	52±19[a]	24.1±7.2[b]	40±2.8[a]	47.1±11.9[a]
S (%)	0.05±0.015[a]	0.025±0.005[b]	0.034±0.007[a]	0.024±0.003[b]
Ca (%)	0.16±0.04[a]	0.33±0.06[b]	0.82±0.34[a]	0.92±0.05[a]

Table 2. Changes in leaf gas-exchange parameters and root phytohormone in sorghum cultivated in control and drought conditions. Different letters in superscript within a row (i.e. phenotype) indicate significant differences between means ± SD of three independent biological replicates ($n = 3$) at $p < 0.05$ based on t-test. Plants were cultivated in a growth chamber for 6 weeks.

Parameters	Control	Drought
Photosynthetic rate (μmol m^{-2} s^{-1})	19±1.3[a]	5±0.52[b]
Transpiration rate (μmol m^{-2} s^{-1})	0.002±0.0009[a]	0.0005±0.0001[b]
Stomatal conductance (μmol m^{-2} s^{-1})	0.09±0.01[a]	0.005±0.003[b]
Water use efficiency (intrinsic) (μmol mol^{-1})	195±12.8[a]	72±2.9[b]
Plant hormone	Control	Drought
IAA (ng g^{-1})	8.3±1.5[a]	4.4±1.0[b]
ABA (ng g^{-1})	3.7±1.4[a]	78.6±28.9[b]
JA (ng g^{-1})	34.5±1.4[a]	9.2±2.5[b]
GA3 (ng g^{-1})	10.5±2.4[a]	9.3±2.2[a]

Table 3. The best functional predictors associated with either drought or control were assigned using the MetaCyc pathway database or the FAPROTAX functional annotation.

Metacyc label (M) or Faptroax prediction (F)	Pathway	Treatment group
ALL-CHORISMATE-PWY(M)	Chorismate metabolism	Control
GLYCOLYSIS(M)	Glycolysis	Control
P184-PWY(M)	Protocatechuate degradation I	Control
P341-PWY(M)	Pyrococcus	Control
PWY-2941(M)	L-lysine biosynthesis II	Control
PWY-5121(M)	Geranylgeranyl diphosphate biosynthesis II	Control
PWY-5896(M)	Menaquinol-10 biosynthesis	Control
PWY-5679(M)	Clavulanate biosynthesis	Drought
PWY-6071(M)	Phenylethylamine degradation	Control

(F)	Nitrate reduction	Control
(F)	Methylotrophy	Control
(F)	Phototrophy	Drought
(F)	Photosynthetic Cyanobacteria	Drought

1133
1134
1135