A transposable element insertion in *IAA16* disrupts splicing and causes dicamba resistance in *Bassia scoparia*

Jacob S. Montgomery[1], Neeta Soni[1], Sofia Marques Hill[1], Sarah Morran[1], Eric L. Patterson[1,2],
Seth A. Edwards[3,4], Sandaruwan Ratnayake[3], Yu-Hung Hung[3], Pratheek H. Pandesha[3,5], R. Keith
Slotkin[3,4], Richard Napier[6], Franck Dayan[1], Todd A. Gaines[1]

[1]Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado, USA.

[2]Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,
Michigan, USA.

[3]Donald Danforth Plant Science Center, St. Louis, Missouri, USA.

[4]Division of Biological Sciences, University of Missouri, Columbia, Missouri, USA

[5]Division of Biology & Biomedical Sciences, Washington University in St. Louis, St. Louis,
Missouri, USA

[6]School of Life Sciences, University of Warwick, Coventry, UK

Running title: Transposon-induced herbicide resistance

Author for correspondence: Todd Gaines, Professor, Colorado State University, 1177 Campus
Delivery, Fort Collins, CO 80523, USA. (Email: todd.gaines@colostate.edu)

22   Keywords: Herbicide Resistance, Auxin, Evolution, Genetic Mapping, Transposable Elements,

23   Weed Science

Abstract

A dicamba-resistant population of kochia (*Bassia scoparia*) identified in Colorado, USA in 2012

was used to generate a synthetic mapping population that segregated for dicamba resistance.

Linkage mapping associating dicamba injury with genotype derived from restriction-site-

associated DNA sequencing identified a single locus in the kochia genome associated with

resistance on chromosome 4. A mutant version of *Auxin/Indole-3-Acetic Acid 16* (*AUX/IAA16*; a

gene previously implicated in dicamba resistance in kochia) was found near the middle of this

locus in resistant plants. Long read sequencing of dicamba-resistant plants identified a recently

inserted Ty1/Copia retrotransposon near the beginning of the second exon of *AUX/IAA16*,

leading to disruption of normal splicing. A molecular marker for this insertion allows for rapid

detection of resistance. Stable transgenic lines of *Arabidopsis thaliana* ectopically expressing the

mutant and wildtype alleles of *AUX/IAA16* were developed. *Arabidopsis thaliana* plants

expressing the mutant *AUX/IAA16* allele grew shorter roots on control media. However,

transgenic root growth was less inhibited on media containing either dicamba (5 μM) or IAA (0.5

μM) when compared to non-transgenic plants or those expressing the wildtype allele of

AUX/IAA16. In vitro assays indicate reduced binding affinity and more rapid dissociation of the

mutant AUX/IAA with TIR1 in the presence of several auxins, and protein modeling suggests the

substitution of the glycine residue in the degron domain of AUX/IAA16 is especially important

for resistance. A fitness cost associated with the mutant allele of *AUX/IAA16* has implications for

resistance evolution and management of kochia populations with this resistance mechanism.

Significance

Auxin mimics are amongst the most important herbicides in modern agriculture. Evolution of

weeds that are resistant to these herbicides threatens sustainable crop production. Understanding

the basis of auxin herbicide resistance informs the development of improved weed control

technologies. Additionally, auxin-resistant mutations and their pleotropic effects help us

understand auxin perception and signalling. We describe a transposable element insertion within

an herbicide target site gene that alters splicing and reduces synthetic and natural auxin

perception.

Introduction

Weed control in modern agricultural production systems relies largely on herbicides. The

first synthetic herbicides to be commercialized (e.g. 2,4-D; described by Quastel, 1950) were

eventually determined to mimic the endogenous plant hormone auxin. The low cost of these

synthetic auxin herbicides, matched with their ability to selectively control broadleaf weeds in

cereal production, led to rapid adoption of herbicide-based weed management. As new herbicides

were discovered and commercialized, mixing multiple herbicides in a single spray mixture has

allowed for robust control of many of the most important weeds in diverse agricultural settings.

While evolution of herbicide resistant weed populations has reduced the efficacy of many

herbicides (Gaines et al., 2020), evolution of resistance to synthetic auxin herbicides has been

slower and less frequent than for other modes of action (Busi et al., 2018; Heap, 2024). Thus,

some of the oldest herbicides (e.g., 2,4-D and dicamba) remain some of the most widely used

across many agricultural settings. Indeed, auxin resistance traits were engineered into dicot crops

(e.g. soybean and cotton) to provide farmers more control options against weed populations that

had evolved glyphosate resistance (Behrens et al., 2007; Wright et al., 2010). These traits

expanded the potential use and value of two of the most used synthetic auxin herbicides, dicamba

and 2,4-D. This expansion has had the unintended effect of increasing selection pressure for

resistance alleles and is correlated with a recent increase in the number of resistance reports to

these herbicides. Beyond weed control, these synthetic auxin compounds have been crucial for

experiments to understand auxin perception and signaling, perhaps the most complex of all plant

hormone signaling pathways (Ma et al., 2017).

While the physiological mechanism of action for many herbicides has been well

understood for decades, the cascade of cellular responses following the application of a synthetic

auxin herbicide is convoluted and is the subject of many recent studies. The topic has been

reviewed several times, with new experimental results incrementally advancing our knowledge

of the key factors leading to plant death (Grossmann, 2010; Song, 2014; Christoffoleti et al.,

2015; McCauley et al., 2020). Auxin, whether synthetic or natural, first binds to one or more of

the isoforms of its receptor protein, Transport Inhibitor Response 1/Auxin-Signaling F-Box

(TIR1/AFB) (Dharmasiri et al., 2005; Kepinski & Leyser, 2005). Once bound to TIR1/AFB,

auxin acts as a "molecular glue", mediating the interaction between TIR1/AFB and the co-

receptor Auxin/Indole-3-Acetic Acid (AUX/IAA) proteins. Identification of conserved domains

and experimentation have produced a list of residues in TIR1/AFB and AUX/IAA that are vital

for this interaction (Tan et al., 2007; Uzunova et al., 2016; Ramans-Harborough et al., 2023).

This interaction leads to the ubiquitination and rapid degradation of AUX/IAA proteins (Gray et

al., 2001; Zenser et al., 2001). Because AUX/IAA proteins act as transcriptional regulators,

AUX/IAA degradation leads to the modulation of auxin-responsive gene expression (Grossmann,

2010). TIR1/AFB and AUX/IAA are each multi-gene families with varying numbers of

homologs. The diversity in gene members of these families allow for many possible

combinations of TIR1/AFB and AUX/IAA, which in turn allow for tissue, temporal and

environmental regulation of auxin responses. Different synthetic auxin herbicides likely promote

the interaction of certain AUX/IAA proteins with certain TIR1/AFB proteins more efficiently,

but this network of interactions is still not well understood. For example, LeClere et al. (2018)

show that AUX/IAA16 preferentially binds with TIR1 and AFB6 (a homolog of AFB found in

members of amaranthaceae, but not brassicaceae) in the presence of some auxin analogs, but not

others.

A disruption in any step of this auxin signaling system (e.g. TIR1 binding auxin,

AUX/IAA interacting with TIR1/AFB, or the ubiquitination and degradation pathway of

AUX/IAA) could cause reduced activity of synthetic auxin herbicides (Ruegger et al., 1998;

LeClere et al., 2018; Todd et al., 2020). Thus, evolution of synthetic auxin resistance should be

relatively simple and common across weed species in the face of incredibly pervasive and strong

selection pressure. However, mutants causing resistance to synthetic auxin herbicides are likely

also to cause a reduction in fitness compared to the wildtype due to the conserved function of

these proteins (Roux & Reboud, 2005; Wu et al., 2021), indicating that resistance mutations are

likely to be purged during temporal absence of synthetic auxin herbicide selection pressure

during rotations in land and herbicide use. Nevertheless, there are several reported cases of

synthetic auxin herbicide resistance becoming established in some weed species, including

examples of enhanced metabolic detoxification in synthetic auxin herbicide resistant weeds

similar to the mechanism of selectivity in tolerant crops (Torra et al., 2024). Interestingly, in

cases in which the genetic basis of resistance has been determined, all of these reports pertain to

mutations in or near the degron domain of Aux/IAA genes. LeClere et al. (2018) and Ghanizadeh

et al. (2024) report, in closely related species, different mutations of a specific glycine residue in

AUX/IAA16 that is known to be vital to normal auxin signaling. In unrelated species, de

Figueiredo et al. (2022b) report a 27-bp deletion that shortens the distance between the degron

and PB1 domains of IAA2, and Krishnan et al. (2024) report a double deletion mutation flanking

the degron region of IAA20.

Here, we seek to understand the genetic and molecular basis of dicamba resistance in a

population of kochia (*Bassia scoparia*) collected from Colorado, USA. We use linkage mapping

to identify a region of the genome that segregates with resistance and contains a mutant version

of *AUX/IAA16*. Long read sequencing allowed us to identify a transposable element insertion in

the coding region of *AUX/IAA16* that disrupts normal splicing and changes several amino acids

in the protein product relative to the wildtype. Ectopic expression in *Arabidopsis thaliana* proves

this mutant *AUX/IAA16* is sufficient to cause resistance to dicamba. Surface plasmon resonance

assays detect a reduction in binding affinity and more rapid dissociation for the mutant

AUX/IAA protein, and protein modeling points to the importance of a glycine-to-threonine

substitution in the degron domain of this mutant. This mutation also seems to confer a fitness

cost associated with this mechanism of dicamba resistance, with implications for agricultural

management of dicamba-resistant weed populations.

Results and Discussion

Characterization of herbicide resistance

Dose response experiments were conducted to confirm and quantify the level of dicamba

and 2,4-D resistance in a population of kochia (named M32) collected from Akron, Colorado,

USA (Figure 1A and 1B). Parameter estimates for two-parameter log-logistic models for each

population used in these experiments are shown in Table S1 and Table S2. In the dicamba

experiment, populations M32 and 9425 (a known dicamba-resistant kochia population previously

characterized by LeClere et al. (2018) and Pettinga et al. (2018)) had ED50 estimates 7.6

(p<0.0001) and 10.5 (p<0.0001) times greater than an herbicide-sensitive kochia population

(7710; described by Preston et al. (2009)). The ED50 parameter represents the dose of herbicide

needed to cause 50% injury to that specific population. These estimates were significantly

different from the herbicide-sensitive control and from each other (p<0.0001), indicating

dicamba resistance in M32 is not as high as in 9425. Nevertheless, the ED50 estimate for both

resistant populations were greater than the recommended field rate of dicamba (560g ha^{-1}),

indicating neither population would be controlled by dicamba in a field setting. Similarly, in the

2,4-D experiment, the ED50 estimate for M32 was 5.0 (p<0.0001) times greater than that of the

herbicide-sensitive control and greater than the recommended field rate for 2,4-D (approximately

560 g ha^{-1}). It is worth noting that the ED50 estimate for the herbicide-sensitive population (543

g ha^{-1}) is nearly equal to the recommended field rate for 2,4-D application. Thus, 2,4-D is not

likely to provide good control of kochia in general.

While herbicide-sensitive plants show epinasty starting approximately 5 days after

treatment with dicamba or 2,4-D, resistant plants in our study do not show any symptoms when

treated with a field rate of either herbicide. This lack of initial injury suggests that resistant plants

either rapidly metabolize/sequester the herbicides before they cause injury (Behrens et al., 2007;

Ge et al., 2010) or possess mutant target site(s) that reduce their interaction with their molecular

target(s) (Walsh et al., 2006; LeClere et al., 2018; de Figueiredo et al., 2022b). Highly efficient

resistance has been engineered into some crops using genes coding for metabolic enzymes from

bacteria (Behrens et al., 2007; Wright et al., 2010). In these cases, no injury symptoms are

observed at field rates. However, in documented cases of metabolic auxin herbicide resistance,

resistant plants often do show symptoms of herbicide damage but eventually recover (de

Figueiredo et al., 2022a; Todd et al., 2024).

To understand if physiological mechanisms prevent dicamba from being absorbed or

translocated within resistant plants, radiolabeled dicamba was used to track its uptake and

movement in dicamba-resistant and -sensitive plants. Significantly less dicamba was absorbed by

herbicide-sensitive plants over 192 h compared to the M32 and 9425 populations ($p=0.036$ and

$p=0.019$ respectively), but the rate of absorption (time required for 90% absorption) was not

significantly different between populations (Figure 1C, Table S3). Therefore, reduced dicamba

absorption is not responsible for resistance. Time series data for the translocation of radiolabeled

dicamba out of the treated leaf did not fit well to any tested models, so t-tests were used to

determine if means differed between resistant and sensitive populations at each time point

(Figure 1D). Dicamba-resistant plants translocated less dicamba out of the treated leaf compared

to the sensitive plants. These results suggest that the dicamba resistance mechanism in the

experimental population is caused by a mutation in *BsAUX/IAA16*, which is similar to what has

been described by Pettinga et al. (2018) and LeClere et al. (2018) and that is caused by a

mutation in *BsAUX/IAA16*.

A single locus on chromosome 4 is associated with dicamba resistance

A quantitative trait loci (QTL) mapping approach was taken to identify regions of the

genome associated with dicamba resistance. Segregating F_3 populations were developed from

multiple independent biparental crosses using a dicamba-resistant plant as the male parent and a

dicamba-sensitive plant as the female parent. We expected to find a single dominant locus

controlling dicamba-resistance because the ratio of alive:dead F_3 plants following dicamba

treatment at a rate of 560 g ha^{-1} (200:85) did not significantly differ from 3:1 (p = 0.06). We

hypothesize that the over-abundance of dead (herbicide-sensitive) plants may be caused by a

fitness penalty associated with the dicamba resistance mechanism that reduces germination

frequency (see section on fitness cost below). A novel double-digest restriction site associated

DNA sequencing (ddRADseq) protocol was used to determine genotypes of 103 segregating F_3

plants for 1,592 variant loci across the genome. A genome scan associating this genotype

information to visual injury rating identified one significant QTL that peaks at 88 Mbp on

chromosome 4 (Figure 2A). This interval contains *BsAUX/IAA16* at 87.4 Mbp, a gene that is

important for auxin perception and has been implicated in auxin herbicide resistance in kochia

and a closely related species (LeClere et al., 2018; Ghanizadeh et al., 2024). In plants from an

independent segregating F_3 line, the genotype of *BsAUX/IAA16* explained 43% of variation in

visual injury following dicamba treatment at 560 g ha^{-1} (Figure 2B). While survival at this rate of

dicamba may be dominant, visual injury is incompletely dominant (Figure 2B) with a calculated

degree of dominance of -0.58 (Falconer, 1964). This dominance has important implications for

the spread of herbicide resistance alleles. For instance, a dominant trait is likely to spread more

quickly throughout a population, but is less likely to be driven to fixation under selection

(Holmes et al., 2022).

A transposable element insertion in BsIAA16 causes a change in exon splicing

Sequencing transcripts of *BsAUX/IAA16* from plants homozygous for either parental

allele of the QTL discovered on chromosome 4 revealed near identical translated protein

sequences except for the region around the degron domain (Figure 3B; Table S4). The gene

model for *BsAUX/IAA16* predicts 5 exons in wild type plants with the degron of *BsAUX/IAA16*

located near the splice junction of exons one and two (Figure 3A). To understand the sequence of

genomic DNA in and around *BsAUX/IAA16*, a plant homozygous for the resistant parent allele of

BsAUX/IAA16 was used for PacBio HiFi sequencing. After *de novo* assembly, we discovered a

3,466 bp insertion at the beginning of exon two of *BsAUX/IAA16* (Figure 3A). The insertion

disrupts splicing of *BsAUX/IAA16*, replacing 16 base-pairs at the beginning of exon two in the

wildtype with the final 19 bases of the insertion in the mutant (Figure 3A). The mutant allele will

herein be referred to as *BsAUX/IAA16$_{Mut}$*, while the wildtype allele will be referred to as

BsAUX/IAA16$_{WT}$. This insertion results in the addition of one amino acid and the replacement of

five more in the final protein product (Figure 3B; Table S4). The new splice sites contain

canonical GT and AG motifs. This change includes a substitution of a glycine to threonine at the

N-terminal end of degron domain (Figure 3B; Table S4). LeClere et al. (2018) showed that a

substitution of this glycine to asparagine in BsAUX/IAA16 is sufficient for dicamba resistance in

kochia. Similarly, Ghanizadeh et al. (2024) associate a substitution of this glycine to aspartic acid

with dicamba resistance in *Chenopodium album*, a closely related species. Finally, NMR

experiments by Ramans-Harborough et al. (2023) show that this glycine is involved in the

interaction between AUX/IAA and TIR1. Given the demonstrated importance of this specific

glycine residue, we hypothesize that the glycine to threonine substitution is the major driver of

dicamba resistance. Protein modeling shows that the larger side group of threonine contacts the

surface of the TIR1 binding pocket and causes steric hindrance of the IAA/TIR1 interaction

(Figure 3C). The rest of the mutant residues do not interact with the binding pocket of TIR1 and

are within a disordered domain of AUX/IAA; thus, they are unlikely to affect IAA/TIR1

interactions (Tan et al., 2007; Ramans-Harborough et al., 2023).

The retrotransposon insertion in *BsAUX/IAA16$_{Mut}$* has 5 bp target site duplications

flanking 429 bp long terminal repeats (LTRs), identifying it as a Class I LTR element. No protein

coding sequences were found within this insertion, suggesting this element is non-autonomous

and relies on another transposable element for transposition. In order to search for the

corresponding autonomous element, we aligned the insertion to the *de novo* assembly built from

a dicamba-resistant plant and the reference genome built from a herbicide-sensitive plant. This

identified a near perfect match (99.9% identity) 1 Mbp away on chromosome 4 and another

match (96.7% identity) on chromosome 2 in both assemblies (Figure 4B, Table S5). Because the

site on chromosome 4 is the most similar across the genome and retrotransposons use a "copy

and paste" mechanism of transposition, we hypothesize this to be the donor location for the

retrotransposon insertion into in *BsAUX/IAA16$_{Mut}$*. A maximum likelihood phylogenetic tree

supports this idea, as the sequences from both assemblies form a distinct clade from those on

chromosome 2 (Figure S2). When comparing the insertion in *BsAUX/IAA16$_{Mut}$* with its donor site

on chromosome 4, there is one single nucleotide polymorphism (SNP) and two single base pair

insertions (Table S5). These insertions are found in homopolymer runs of at least 9 bp and are

likely artifacts of the sequencing technology (Espinosa et al., 2024). This similarity suggests very

recent transposition. In fact, the 429 bp LTR sequences of the insertion in *AUX/IAA16* are

identical. Given the approximated rate of mutation in *Arabidopsis thaliana* of 7×10^{-9} base

substitutions per site per generation (Ossowski et al., 2010), we estimate a new single base

substitution to be generated in either one of these 429 bp regions approximately every 166,500

generations. This estimation is very crude, but it suggests that the insertion is extremely young in

the context of evolutionary time. Future research using collections of kochia in its native range

could test whether this adaptation is the consequence of selection on standing genetic variation or

de novo genetic diversity that was generated in the time of modern agriculture (Kreiner et al.,

2022).

Since we had identified a likely donor retrotransposon and had not detected any protein

coding sequences within this element, there must have been an autonomous element that

retrotransposed the donor from chromosome 4 into *BsAUX/IAA16*. To identify the autonomous

version(s) of the retrotransposon within *BsAUX/IAA16*, BLAST was first used to find similar

LTRs in the previously assembled reference genome (Hall et al., 2023). Putative LTRs were first

filtered by sequence similarity and distance between other putative LTRs. Matches were only

considered if the putative LTRs were in the same orientation. Once matches were found,

identification of required retrotransposon domains was performed for each match. This process

led to the identification of a single autonomous element that contained all domains required for

transposition (Table S6). This 13 kb putative autonomous retrotransposon, named gi_5_13kb,

was found on chromosome 5 (Figure 4C). There are two known superfamilies of

retrotransposons, Ty1/Copia and Ty3/Gypsy. To determine which gi_5_13kb belongs to, we first

built a curated list of high-confidence LTR transposons in Arabidopsis which contained all

required domains. Then, we used the predicted amino acid sequence of these high-confidence

LTR transposons to build a phylogenetic tree with the predicted amino-acid sequence of

gi_5_13kb. This element clustered within the Ty1/Copia branch of the tree (Figure 4A). We have

subsequently named this novel Ty1/Copia element *Outlaw*. This one protein-coding version of

the *Outlaw* seems to be driving retrotransposition of the non-autonomous element that generated

herbicide resistance in this population of kochia.

BsIAA16$_{Mut}$ expression is sufficient for dicamba resistance in Arabidopsis thaliana

To demonstrate the role of *BsAUX/IAA16$_{Mut}$* in dicamba resistance, stable transgenic lines

of *Arabidopsis thaliana* expressing *BsAUX/IAA16$_{WT}$* or *BsAUX/IAA16$_{Mut}$* were generated through

the floral dip method. When seeds were grown on phytoagar plates without herbicide,

Arabidopsis expressing *BsAUX/IAA16$_{Mut}$* grew shorter roots compared to those expressing

BsAUX/IAA16$_{WT}$ or with no transgene (Figure 5A). This is evidence of a likely fitness cost

associated with disrupting normal auxin signaling in roots due to mutant Aux/IAA over-

expression as seen by de Figueiredo et al. (2022b) and LeClere et al. (2018). When grown on

media containing 5 μM dicamba, only seedlings expressing *BsAUX/IAA16$_{Mut}$* were able to grow

and had significantly longer roots than seedlings expressing *BsAUX/IAA16$_{WT}$* or no transgene

when normalized to the length of roots grown on solvent-only media (Figure 5B). A similar

result was observed when seedlings were grown on media containing 0.5 μM IAA (Figure 5C),

suggesting this variant of AUX/IAA16 also reduces the perception of natural auxin. No seeds

grew roots on media containing 5 μM or 0.5 μM 2,4-D.

To confirm the results from root growth assays that indicate *BsAUX/IAA16$_{Mut}$* is sufficient

for dicamba resistance, plants of select T$_3$ lines either expressing *BsAUX/IAA16$_{WT}$*,

BsAUX/IAA16$_{Mut}$, or no transgene were grown in the greenhouse and treated with 140 g dicamba

ha^{-1}. Only one of the lines expressing *BsAUX/IAA16$_{Mut}$* showed less visual injury than the other

lines tested and appeared stunted and darker green than the other lines even when no dicamba

treatment was administered (Figure S3). Quantitative PCR indicated similar expression levels of

BsAUX/IAA16 between lines expressing *BsAUX/IAA16$_{WT}$* or *BsAUX/IAA16$_{Mut}$* (Figure 5D).

However, plants expressing *BsAUX/IAA16$_{Mut}$* had no detected induction of auxin response genes

AtIAA19 or *AtGH3.3* six hours after dicamba treatment (Figure 5E and 5F). These genes are

known to have increased expression following exposure to natural or synthetic auxins (Tatematsu

et al., 2004; Gleason et al., 2011; Pettinga et al., 2018; de Figueiredo et al., 2022b).

Reduced binding of BsAUX/IAA16$_{Mut}$ to auxin receptor protein TIR1 provides a molecular

explanation of resistance

305   To provide a molecular explanation of the role of *BsAUX/IAA16$_{Mut}$* in dicamba

resistance, we measured binding of the Arabidopsis auxin receptor protein AtTIR1 to the WT and

mutant BsAUX/IAA16 degron sequences using surface plasmon resonance (SPR; de Figueiredo

et al., 2022b; Prusinska et al., 2023). Binding was auxin-dependent. In the presence of indole-3

acetic acid (IAA), binding of the WT peptide was strong and persistent (dissociation of the co-

receptor complex was slow; Fig3D). Binding of the WT peptide in the presence of 2,4-D and

dicamba was somewhat lower than with IAA as reported previously (Prusinska et al., 2023) with

binding amplitudes at the end of the binding phase 56% and 21% of the IAA value, respectively.

Binding to the mutant degron peptide was poorer in all respects. The binding amplitudes were

lower than for the WT degron (IAA = 37%, 2,4-D 18% and dicamba 5%), and the complexes

dissociated more rapidly. In summary, the assembly of the co-receptor complex with the mutant

degron was greatly reduced, and when formed, its lifetime was shorter. This would likely lead to

far less ubiquitination, higher concentrations of BsAux/IAA16$_{Mut}$ remaining in the cell and

reduced auxin signal strength. The greatly reduced signals in the presence of the synthetic auxins

2,4-D and dicamba probably account for resistance as recorded in this line.

The BsIAA16$_{Mut}$ allele is associated with a reduction in fitness

Because we observe lower binding of *BsAUX/IAA16$_{Mut}$* with TIR1 in the presence of natural auxin, we expect this change to also confer a fitness penalty in the absence of synthetic auxin application (LeClere et al., 2018; Wu et al., 2021; de Figueiredo et al., 2022b). To formally test this, kochia plants were grown from an F$_3$ family that is segregating for the *BsAUX/IAA16* locus and fitness traits were measured without the application of auxin. The ratio of genotypes significantly differed from the expected 1:2:1 (p<0.0001), with fewer than expected plants with the *BsAUX/IAA16$_{Mut}$* allele (Figure 6). All seedlings that emerged were used in the experiment, so we hypothesize that *BsAUX/IAA16$_{Mut}$* causes reduced germination. Further, *BsAUX/IAA16$_{Mut}$* was associated with shorter plants, both at transplanting and at maturity (Figures 6A and 6B). However, *BsAUX/IAA16* did not have a significant effect on above ground biomass (Figure 6C) or average seed weight (Figure 6D). Thus, while the *BsAUX/IAA16$_{Mut}$* allele is associated with reduced plant height, plants with this allele accumulate similar biomass and produce similarly sized seeds. Taken together, these results suggest this herbicide resistance mechanism does slightly reduce plant fitness under normal (non-selective) conditions, but not enough to cause it to be completely purged from the population. Nevertheless, evidence of a fitness cost associated with herbicide resistance has implications for kochia management. For instance, if dicamba is not used for several years, the frequency of *BsAUX/IAA16$_{Mut}$* is expected to decrease. Future experiments to investigate allele frequency across generations under competition or abiotic stress will help clarify the real-world ecological consequences of the *BsAUX/IAA16$_{Mut}$* allele (Wu et al., 2018). Hopefully, the negative pleotropic effect of *BsAUX/IAA16$_{Mut}$* may help limit the spread of this resistance allele and allow producers to better manage resistant populations that have already

established. This will be especially important as high gene flow in kochia has facilitated rapid

spread of other herbicide resistance alleles and mitigated negative effects of genetic

bottlenecking following selection for resistant phenotypes (Martin et al., 2020; Ravet et al.,

2021).

Taken together, our results further emphasize the importance of the glycine residue within

the degron domain of AUX/IAA proteins in auxin signaling. They also reinforce the importance

of *BsAUX/IAA16*, specifically, in the action of synthetic auxin herbicides. We describe the first

instance of the insertion of a mobile genetic element into the coding sequence of a gene to create

an herbicide resistance allele through gain-of-function. The genome of kochia may be especially

active for transposable element activity, as the insertion of a mobile genetic element is also

associated with gene duplication resulting in glyphosate resistance in this species (Patterson et

al., 2019; Ravet et al., 2021; Hall et al., 2023). These results highlight the amazing adaptability

of weedy genomes, and the remarkable ways genomes evolve in the face of strong selection

pressures (Montgomery et al., 2024).

Experimental Procedures

Plant growth and herbicide treatment conditions

Unless otherwise noted, all plants were grown in greenhouses at Colorado State

University in 3.8-cm by 3.8-cm by 5.8-cm pots containing fine-grade potting mix (Fafard #2-SV;

American Clay Works, Denver, CO). Temperatures were held between 22C and 24C, and

supplemental light was provided by liquid halogen grow lights to ensure a photoperiod of

14h/10h. Herbicide applications were made using a moving overhead single-nozzle sprayer

(DeVries Manufacturing, Hollandale, MN) calibrated to deliver 187 L ha^{-1}. Engenia (BASF,

Research Triangle Park, NC) was used for dicamba applications, and Clean Amine (Loveland

Products, Inc., Loveland, CO) was used for 2,4-D applications. For kochia, herbicide

applications took place when plants were 10-15 cm tall, and for Arabidopsis, plants were sprayed

when they had 10-12 leaves and before bolting.

Plant material

The M32 population of kochia was collected near Akron, Colorado as part of an herbicide

resistance survey conducted between 2012 and 2014 (Westra et al., 2019). Seeds from this field

collection were grown, and the resulting plants were treated with 560 g dicamba ha^{-1}. Survivors

were open-pollinated, and their seed was combined to form a composite dicamba-resistant seed

lot. Approximately 50 seeds from this combined lot were grown and treated with 560 dicamba

ha^{-1}. The individual with the least herbicide injury was used as a male parent in a biparental cross

with an individual from the population 7710, an inbred, herbicide-susceptible population

(Preston et al., 2009) that was used to generate the reference genome assembly for kochia

(Patterson et al., 2019; Hall et al., 2023). To make the cross, immature flowers of the dicamba-

sensitive plant were emasculated using forceps under a microscope, and the emasculated flowers

were labeled. The two plants were then grown to maturity together in a pollen-exclusion tent.

Seeds from the dicamba-sensitive plant were collected and grown; the resulting plants were

treated with 560 g dicamba ha^{-1}. A Kompetative Allele Specific PCR (KASP) assay that detects

BsAUX/IAA16$_{Mut}$ was used to confirm survivors as hybrids (primer sequence and thermocycler

conditions found in Table S7). Several confirmed hybrid plants were grown in pollen exclusion

tents and self-pollinated to create several F_2 families. Plants from two F_2 families were grown,

but because of low seed production in the F_1 generation, plants were allowed to self-pollinate

with no selection to produce F_3 families.

Dose response

Plants were grown from the 7710 (Preston et al., 2009) and 9425 (LeClere et al., 2018)

populations as well as from the M32 composite resistant seed lot. Plants were treated with one of

several rates of dicamba. These rates included 0, 8.75, 17.5, 35, 70, 140, 280, 560, and 1120 g

dicamba ha^{-1} for the herbicide-sensitive population and 0, 70, 140, 280, 560, 1120, 2240, and

4480 g dicamba ha^{-1} for the herbicide-resistant populations. Six uniform plants were used for

each population at each herbicide rate before being returned to the greenhouse. Visual injury and

survival were rated 21 days after treatment (DAT). The drc package v3.0-1 (Ritz et al., 2015) in

R v4.0.2 (R-Core-Team, 2021) was used to fit the injury data to a 2-parameter log-logistic dose

response curve. Significant difference is in the ED50 parameter were determined using the

compParm function from the drc package (Ritz et al., 2015).

Herbicide absorption and translocation profiles

Kochia plants from M32, 7710, and 9425 were germinated in a growth chamber (60%

relative humidity, 21/18□C, and 16/8 h photoperiod) in potting soil and transplanted to fine sand

when they reached ~3 true leaves. The plants were irrigated with fertilizer until the plants

reached 10 cm in height. At this point, aluminum foil was used to cover the second youngest

fully expanded leaf while the plants were sprayed with 560 g dicamba ha^{-1}. After spraying, the

aluminum foil was removed, the covered leaf was then marked and treated with 10 μL of a ^{14}C-

labeled dicamba solution (total radioactivity of 3.33 KBq or 200,000 dpm per plant). Plants were

returned to the growth chamber until sampling at 3, 6, 12, 24, 48, 96, and 192 h after treatment.

The treated leaf, remaining above ground tissue, and below ground tissue of three biological

replicates were separated for each population at each time point. The treated leaf was washed in

10 mL 10% methanol + 1% NIS, and radioactivity in this wash solution was quantified in 10 mL

of scintillation mixture (Ecoscint XR, National Diagnostics, Atlanta, GA) using liquid

scintillation spectrometry (Packard Tricarb 2300TR, Packard Instrument Co., Meriden, CT).

Plant tissue was dried in an oven at 60 C for at least 14 d before oxidation in a biological

oxidizer (OX500; RJ Harvey Instrument Co., Tappan, NY) followed by radioactivity

measurement by liquid scintillation spectrometry. Absorption time series data was fitted to a

rectangular hyperbolic model, and the parameters "absorption max" (*Amax*) and "time to 90

percent absorption" (*t90*) were estimated and compared in R (Kniss et al., 2011; Ritz et al., 2015;

R-Core-Team, 2021). Because the time series data for translocation of dicamba outside of the

treated leaf did not fit any of the models proposed by Kniss et al. (2011), t-tests were used to

determine if the amount of dicamba retained in the treated leaf was different between the M32

and 7710 populations at each time point.

QTL mapping

Plants from several F_3 families derived from a biparental cross described above were

grown and treated with 560 g dicamba ha[-1]. Two families that segregated for survival following

dicamba treatment (4-1-1 and 4-5-10, each from a different F_1 hybridization event) were used for

QTL mapping. Approximately 150 plants from the F_3 family 4-5-10 were grown, and a single

young leaf was sampled and frozen in liquid nitrogen before treatment with 560 g dicamba ha[-1].

Visual injury on a scale from 0-100 was rated for each plant 21 DAT, and DNA was extracted

from each plant (including the original parents of the cross) using the CTAB method (Doyle &

Doyle, 1990). The concentration of each DNA sample was quantified using the Qubit™ dsDNA

BR kit (Thermo Scientific, Waltham, MA) and diluted to 20 ng/μl. A double-digest restriction-

site associated DNA sequencing (ddRADseq) protocol was developed in conjunction with the

University of Minnesota Genomics Center (Minneapolis, MN) by estimating the number of RAD

sites across the reference genome (Patterson et al., 2019) for several enzyme combinations and

then empirically testing the BtgI-TaqI combination on several samples at multiple sequencing

depths. DNA from the two original parents and 103 plants from the 4-5-10 F_3 family were used

for ddRADseq with BtgI and TaqI as the restriction enzymes with a target of 4 million 150 bp

single end reads per sample. Library prep and sequencing was completed by the University of

Minnesota Genomics Center.

Reads from each sample were passed through a variant calling pipeline that can be found

at github.com/JMont12/dicamba_kochia. Briefly, raw reads were trimmed with Trimmomatic

v0.36 (Bolger et al., 2014) and aligned to the reference genome of kochia (Hall et al., 2023)

using the burrows-wheeler aligner v0.7.17 (Li & Durbin, 2009). The alignments were processed

with samtools v1.15 (Li et al., 2009) and passed to GATK v4.2.0 (Poplin et al., 2018; Van der

Auwera & O'Connor, 2020) to call variants and filter them based on depth, quality, and strand

bias. Variants were further filtered using custom scripts to only include biallelic single nucleotide

polymorphisms that were homozygous and different between the two original parents. Filtered

variants and phenotype data were used to conduct a genome scan within the qtl2 package in R

v0.32 (Broman et al., 2019). Because a permutation test for significance was not possible, a

significance threshold was established through a Bonferroni correction of alpha=0.05

considering the number of tests to equal the number of filtered markers used in the scan then

converting the adjusted p-value to LOD as described by Nyholt (2000). The genotype around the

QTL discovered on chr4 was determined for 210 plants from another F_3 family (4-1-1) using the

KASP assay developed for the *BsAUX/IAA16* gene (primer sequence and thermocycler

conditions in Table S6). A simple linear model was fitted using visual injury 21 DAT as the

dependent variable and *BsAUX/IAA16* genotype as the independent variable and plotted using

ggplot2 v3.4.4 (Wickham, 2016) in R (R-Core-Team, 2021).

Similarly, to quantify the effect of *BsAUX/IAA16* genotype on fitness traits, ~300 seeds of

an F3 family segregating for the *BsAUX/IAA16* locus were sown into damp soil. Three weeks

after planting, all emerged seedlings were transplanted into individual pots (10.5 x 10.5 x 12 cm)

filled with potting soil and irrigated evenly throughout their lifecycle. Plants were spaced evenly,

watered sufficiently, and not fertilized to reduce competition and environmental variation. Plant

height was measured at transplanting and maturity. After allowing the plants to senesce naturally,

above ground biomass was collected and weighed. For a subset of plants, seeds were harvested

and a random sample of 50 seeds was used to determine average seed weight for each sample.

During development, a small amount of leaf tissue was collected from each plant and used to

determine the genotype of the *BsAUX/IAA16* locus using the KASP assay described above.

Genotype and phenotype information was used to fit linear models and plotted using ggplot in R.

BsIAA16$_{M32}$ sequencing

Total RNA was extracted using the Direct-zol RNA Microprep kit (Zymo Research,

Irvine, CA) from individuals of the 4-5-10 selected line that were homozygous for either parental

allele of *BsAUX/IAA16* based on the results of KASP testing. This RNA was used to generate

cDNA libraries for each sample using the ProtoScript® II First Strand cDNA Synthesis Kit (New

England Biolabs, Ipswich, MA). Primers that bind to the beginning and end of the *BsAUX/IAA16*

gene (Table S7) were used with the EconoTaq PLUS Green kit (LGC, Teddington, Middlesex,

UK) to amplify the full-length coding sequence. The product of each PCR reaction was run on a

1% agarose gel to ensure a single product, then sent to Azenta (Burlington, MA) for Sanger

sequencing using the forward and reverse primers.

To determine the genomic sequence around the *BsAUX/IAA16$_{Mut}$*, 2 g of dark-treated leaf

tissue from a plant that was homozygous for *BsAUX/IAA16$_{Mut}$* was sampled and flash frozen in

liquid nitrogen. This tissue was sent to Corteva Agriscience Center for Genome Excellence

(Johnston, IA) for whole genome PacBio HiFi sequencing to a depth of ~30X. The resulting

reads were assembled with HiFiasm v0.19.5 (Cheng et al., 2021), and *BsAUX/IAA16* was located

by BLAST v2.6.0 (Camacho et al., 2009).

Characterization of transposable element insertion

To find similar elements to the one inserted in *BsAUX/IAA16$_{Mut}$*, the sequence of the 429

bp LTRs of the insertion were aligned to the kochia reference genome assembly (Hall et al.,

2023) and to the newly created assembly described above using BLAST (Camacho et al., 2009).

Sequential filtering was used to identify the autonomous version of the element. First, alignments

of >85% identity and >100bp alignment length were identified as putative LTRs. Next, only

putative LTRs that were between 25 kb and 4 kb apart and in the direct orientation were

considered potential *Outlaw* transposon family members. Finally, of the elements of the proper

length and LTR orientation, we then only considered putative elements that contained all

required retrotransposon domains as potential autonomous elements. Necessary retrotransposon

domains were identified manually and with TESorter (Zhang et al., 2022). Specifically, the

required domains that we used were as follows: gag, pol (RNAseH, Integrase, Protease, Reverse

Transcriptase), tRNA Primer Binding Site, and Polypurine Tract.

Autonomous transposable element characterization

To classify the retro element into one of two possible superfamilies (Ty1/Copia or

Ty3/Gypsy), we used the *Arabidopsis thaliana* transposable element annotation dataset from

RepetDBv2 (Amselem et al., 2019). We procured all Ty1/Copia and Ty3/Gypsy nucleotide

elements and predicted the protein domain sequences from this dataset using TESorter (Zhang et

al., 2022). We only used TEs that had all required retrotransposon domains (both pol and gag).

We then manually concatenated all pol domain proteins (RT, RNaseH, Protease, Integrase) as a

single polypeptide for each transposable element and then used *Outlaw* predicted pol polypeptide

sequence to create a phylogenetic tree. MegaX was used to align all of the proteins using

MUSCLE to create a maximum likelihood tree with 500 bootstraps (Kumar et al., 2018).

Development of transgenic Arabidopsis thaliana lines

The coding sequence of *BsAUX/IAA16$_{Mut}$* and *BsAUX/IAA16$_{WT}$* were amplified from

cDNA libraries discussed above using the PrimeSTAR MAX DNA Polymerase kit (Takara Bio

USA, Inc., San Jose, CA) and primers with tails that complement restriction sites in the

expression vector *pFGC5941* (primer sequences and thermocycler conditions in Table S8).

Empty *pFGC5941* was digested with AscI and BamHI (New England Biolabs, Ipswich, MA),

and the *BsAUX/IAA16* amplicons were ligated into the digested vector using the In-Fusion

Cloning kit (Takara Bio USA, Inc., San Jose, CA). Plasmid sequencing ensured correct assembly

of expression vectors (Plasmidsaurus, Eugene, OR). Vectors containing *BsAUX/IAA16$_{Mut}$* or

BsAUX/IAA16$_{WT}$ were transformed into the GV3101 strain of *Agrobacterium tumefaciens*, and

subsequently transfected into the Col-0 genotype of *Arabidopsis thaliana* using the floral dip

method (Clough & Bent, 1998). Transformants were selected by spraying T$_1$ seedlings with

BASTA herbicide (BASF, Research Triangle Park, NC) and confirmed with PCR for the

transgene (primers and thermocycler conditions found in Table S9). Confirmed transformants

were inbred for two generations, and T$_3$ populations fixed for the transgene were identified by

spraying seedlings with BASTA herbicide and PCR for the transgene.

Effect of dicamba on transgenic Arabidopsis lines

Seeds of Col-0 and of fixed T$_3$ lines of Arabidopsis, each from a different transformation

event, were gas sterilized and plated on media plates containing synthetic or natural auxin as

described by de Figueiredo et al. (2022b). Plates were incubated in a growth chamber with a

yellow light filter to prevent indole-3 acetic acid degradation. Ten seeds were used per

population for each treatment. Root growth was measured 7 d after moving plates to growing

conditions. T-tests were used to detect significant differences in root length between lines

expressing *BsAUX/IAA16$_{Mut}$* compared to the Col-0 control. Results were plotted with ggplot2 in

R.

Plants from select T$_3$ lines were grown to the 10-leaf stage and either untreated or treated

with 140 g dicamba ha^{-1}. Four plants of each population were used for each treatment. Photos

were taken 21 DAT to illustrate visual injury. Young leaf tissue was collected from treated and

untreated plants 6 h after treatment. Total RNA was extracted from these leaf samples using the

Direct-zol RNA Microprep kit (Zymo Research, Irvine, CA), and cDNA libraries were generated

using the ProtoScript® II First Strand cDNA Synthesis Kit (New England Biolabs, Ipswich,

MA). Following the methods of de Figueiredo et al. (2022b), relative expression was quantified

for the *BsAUX/IAA16* transgene (primers and thermocycler conditions available in Table S10)

and the auxin response genes *AtIAA19* and *AtGH3.3* using *AtCyclophilin* as a reference gene and

the $2^{-\Delta\Delta Ct}$ method.

In silico mutagenesis of IAA variants

The crystal structure of the degron of IAA7 bound to TIR1 in the presence of 2,4-D

(2p1n) was downloaded from the RCSB Protein Data Bank (Tan et al., 2007). Mutagenesis of the

G127 residue was conducted and visualized in PyMOL.

Reeceptor protein preparation and Surface Plasmon Resonance measurement of auxin binding

Receptor protein (AtTIR1) was purified and the SPR assays performed following the

methods described in Prusinska et al., 2023. The SPR chip (Series S SA) was loaded with our

reference negative control degron peptide mIAA7 (mutated Arabidopsis Aux/IAA7); channel 2

with positive control degron peptide IAA7; channel 3 with BsAUX/IAA16$_{WT}$ degron peptide

(biotin-AKTQVVGWPPVRAFRKN-amide) and channel 4 with BsAUX/IAA16$_{Mut}$ degron

peptide (biotin-AKSIKFPTWPPVRAFRKN-amide). Similar peptide loads were recorded for all

four channels. Purified receptor was passed over all four channels simultaneously in parallel in

PBS buffer + 0.01% Tween20, with or without test auxin (IAA; 2,4-D; or dicamba) at a

concentration of 50 µM. Binding was recorded for 120 s, and dissociation for 180 s. Data were

visualized in BiaEvaluation software.

Data Availability

The RAD sequencing data from the segregating F_3 line used in the mapping project is available

through NCBI, BioSample IDs SAMN42890277-SAMN42890594 and SRA accession numbers

SRR30019316-SRR30019633. Phenotypic data can be found in Supplemental Table S11. PacBio

HiFi sequencing used to generate the de novo genome assembly is available through NCBI under

BioSample ID SAMN43525227 and SRA accession number SRR30575711. All sequence data is

associated with NCBI BioProject PRJNA1141446.

Acknowledgments

This research was supported in part by BASF SE, by the Colorado Wheat Administrative

Committee, and by the USDA National Institute of Food and Agriculture, Hatch project

COL00783 to the Colorado State University Agricultural Experiment Station. The authors have

no conflict of interest to declare.

Figures

Figure 1. Dose response curves showing magnitude of resistance towards dicamba (A) and 2,4-D

(B) for the M32 population of *Bassia scoparia* (Experimental Resistant) compared to a known

dicamba-sensitive population and known dicamba-resistant population (LeClere et al., 2018;

Pettinga et al., 2018). C) Time course of ^{14}C dicamba absorption showing lack of herbicide

absorption is not associated with dicamba resistance in dicamba-resistant populations of *Bassia*

scoparia. D) Proportion of absorbed ^{14}C dicamba retained in the treated leaf showing less

dicamba exits the treated leaf in dicamba-resistant populations compared to a dicamba-sensitive

population. Error bars represent standard error. *=p<0.05, **=p<0.01. Populations are

represented by the same colors and symbols as in A-C.

Figure 2. A) Quantitative trait loci (QTL) genome scan from a segregating F$_3$ population derived

from a biparental cross of herbicide-resistant and -sensitive *Bassia scoparia* plants. Red line

represents a significance threshold determined by Bonferroni-adjustment of alpha=0.05 followed

by conversion to LOD as described by Nyholt (2000). B) Violin plot displaying the effect of

BsAUX/IAA16 genotype on visual injury 21 days after treatment with dicamba in an independent

segregating F$_3$ population of *Bassia scoparia*. Red line represents a linear model fitted to all

points with equation listed in black. Number of observations for each genotype listed at the base

of the graph.

Figure 3. Characterization of a transposable element insertion in *BsAUX/IAA16* and its effect on

protein structure and function. A) Genomic and coding sequence for *BsAUX/IAA16* alleles

detected in a mapping population segregating for dicamba resistance. B) Amino acid sequence of

mutant (Mut) and wildtype (WT) isoforms of BsAUX/IAA16 around the degron domain. C)

Crystal structure of IAA7 (gold) bound to TIR1 (blue) showing the glycine 127 residue (red)

superimposed over the predicted structure if this residue is substituted for threonine, colored by

element (carbon=grey, oxygen=red, nitrogen=blue, chlorine=green). Residues of TIR1 that are

within 4Å of the threonine residue are colored by element, and 2,4-D is in the background,

colored by element. Steric clashes are shown as red discs. D) Binding/dissociation of BsIAA16

peptides with AtTIR1 in the presence of various auxins as determined by surface plasmon

resonance experiments.

Figure 4. Characterization of autonomous element driving transposition of the retrotransposon

that was inserted in *BsAUX/IAA16*. A) Maximum likelihood phylogenetic tree showing

evolutionary relationships between autonomous LTR elements from Arabidopsis and the

autonomous *Outlaw* element discovered in *Bassia scoparia*. B) Alignment of *Outlaw* elements

similar to the one inserted in *BsAUX/IAA16*. C) Comparison of nonautonomous (top) and

autonomous (bottom) versions of *Outlaw*. Colored blocks represent regions necessary for

retrotransposition.

Figure 5. Evaluation of *BsAUX/IAA16* alleles in *Arabidopsis thaliana*. A) Root length of plants

expressing either *BsAUX/IAA16$_{Mut}$* (red), *BsIAA16$_{WT}$* (blue), or no transgene (grey) grown on

phytoagar media. Proportional root length (compared to ethanol-only control) of the same

populations when grown on media containing 5 μM dicamba (B) or 0.5 μM IAA (C). P values in

A-C are from t-tests comparing means to the Col0 population. Relative expression of the

BsAUX/IAA16 transgene (D) and the auxin response genes *AtAUX/IAA19* (E) and *AtGH3.3* (F) in

plants expressing *BsAUX/IAA16$_{Mut}$* (Mut), *BsAUX/IAA16$_{WT}$* (WT), or no transgene (Col0) that

were either treated (T) or untreated (U) with 140 g dicamba ha^{-1} six hours after treatment. Each

observation is represented by a black dot. **=p<0.01, ***=p<0.001.

Figure 6. Effect of *BsAUX/IAA16* genotype on plant height at transplanting (A) or maturity (B),

dry biomass accumulation at maturity (C), or weight of 50 seeds (D) for kochia plants from an F_3

population segregating for dicamba resistance. Number of mutant alleles indicate homozygous

wildtype (0), heterozygous (1), or homozygous mutant (2). Linear models fitted to the data are

plotted in red with equations listed at the top right of each pane. Replicate number is listed at the

base of each violin with either boxplots or data plotted as dots within each violin.

References

Amselem, J., Cornut, G., Choisne, N., Alaux, M., Alfama-Depauw, F., Jamilloux, V. et al. (2019) RepetDB: a unified resource for transposable element references. *Mob DNA*, **10**, 6.

Behrens, M.R., Mutlu, N., Chakraborty, S., Dumitru, R., Jiang, W.Z., Lavallee, B.J. et al. (2007) Dicamba resistance: enlarging and preserving biotechnology-based weed management strategies. *Science*, **316**, 1185-8.

Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, **30**, 2114-20.

Broman, K.W., Gatti, D.M., Simecek, P., Furlotte, N.A., Prins, P., Sen, Ś. et al. (2019) R/qtl2: software for mapping quantitative trait loci with high-dimensional data and multiparent populations. *Genetics*, **211**, 495-502.

Busi, R., Goggin, D.E., Heap, I.M., Horak, M.J., Jugulam, M., Masters, R.A. et al. (2018) Weed resistance to synthetic auxin herbicides. *Pest Management Science*, **74**, 2265-2276.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. et al. (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421.

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H. & Li, H. (2021) Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, **18**, 170-175.

Christoffoleti, P., Figueiredo, M., Peres, L., Nissen, S. & Gaines, T. (2015) Auxinic herbicides, mechanisms of action, and weed resistance: A look into recent. *Scientia Agricola*, **72**, 356-362.

Clough, S.J. & Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal*, **16**, 735-43.

de Figueiredo, M.R.A., Barnes, H., Boot, C.M., de Figueiredo, A., Nissen, S.J., Dayan, F.E. et al.

(2022a) Identification of a novel 2,4-D metabolic detoxification pathway in 2,4-D-resistant

waterhemp (*Amaranthus tuberculatus*). *J Agric Food Chem*, **70**, 15380-15389.

de Figueiredo, M.R.A., Küpper, A., Malone, J.M., Petrovic, T., de Figueiredo, A.B.T.B.,

Campagnola, G. et al. (2022b) An in-frame deletion mutation in the degron tail of auxin

coreceptor IAA2 confers resistance to the herbicide 2,4-D in *Sisymbrium orientale*.

Proceedings of the National Academy of Sciences, **119**, e2105819119.

Dharmasiri, N., Dharmasiri, S. & Estelle, M. (2005) The F-box protein TIR1 is an auxin receptor.

Nature, **435**, 441-5.

Doyle, J. & Doyle, J. (1990) A rapid total DNA preparation procedure for fresh plant tissue.

Focus, **12**, 13-15.

Espinosa, E., Bautista, R., Larrosa, R. & Plata, O. (2024) Advancements in long-read genome

sequencing technologies and algorithms. *Genomics*, **116**, 110842.

Gaines, T.A., Duke, S.O., Morran, S, Rigon, C.A.G., Tranel, P.J., Küpper, A. et al. (2020)

Mechanisms of evolved herbicide resistance. *J Biol Chem*, **295**, 10307-30

Ge, X., d'Avignon, D.A., Ackerman, J.J. & Sammons, R.D. (2010) Rapid vacuolar sequestration:

the horseweed glyphosate resistance mechanism. *Pest Manag Sci*, **66**, 345-8.

Ghanizadeh, H., He, L., Griffiths, A.G., Harrington, K.C., Carbone, V., Wu, H. et al. (2024) A

novel mutation in *IAA16* is associated with dicamba resistance in *Chenopodium album*. *Pest

Manag Sci*, 10.1002/ps.8071

Gleason, C., Foley, R.C. & Singh, K.B. (2011) Mutant analysis in Arabidopsis provides insight

into the molecular mode of action of the auxinic herbicide dicamba. *PLOS ONE*, **6**, e17245.

Gray, W.M., Kepinski, S., Rouse, D., Leyser, O. & Estelle, M. (2001) Auxin regulates

SCF(TIR1)-dependent degradation of AUX/IAA proteins. *Nature*, **414**, 271-6.

Grossmann, K. (2010) Auxin herbicides: current status of mechanism and mode of action. *Pest

Manag Sci*, **66**, 113-20.

Hall, N., Chen, J., Matzrafi, M., Saski, C.A., Westra, P., Gaines, T.A. et al. (2023) *FHY3/FAR1*

transposable elements generate adaptive genetic variation in the *Bassia scoparia* genome.

bioRxiv, **DOI**, 10.1101/2023.05.26.542497.

Heap, I., The International Herbicide-Resistant Weed Database. www.weedscience.org [accessed

20 June 2024].

Holmes, K.H., Lindquist, J.L., Rebarber, R., Werle, R., Yerka, M. & Tenhumberg, B. (2022)

Modeling the evolution of herbicide resistance in weed species with a complex life cycle.

Ecological Applications, **32**, e02473.

Kepinski, S. & Leyser, O. (2005) The Arabidopsis F-box protein TIR1 is an auxin receptor.

Nature, **435**, 446-451.

Kniss, A.R., Vassios, J.D., Nissen, S.J. & Ritz, C. (2011) Nonlinear regression analysis of

herbicide absorption studies. *Weed Science*, **59**, 601-610.

Kreiner, J.M., Latorre, S.M., Burbano, H.A., Stinchcombe, J.R., Otto, S.P., Weigel, D. et al.

(2022) Rapid weed adaptation and range expansion in response to agriculture over the past

two centuries. *Science*, **378**, 1079-1085.

Krishnan, M., Petrovic, T., Schwerdt, J.G., Merriam, A.B., Hereward, J.P. & Preston, C. (2024) A

novel mutation in *SoIAA20* confers cross-resistance to 2,4-Dichlorophenoxyacetic acid and

other auxinic herbicides in *Sonchus oleraceus*. *Pest Manag Sci*, 10.1002/ps.8413

Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. (2018) MEGA X: Molecular

evolutionary genetics analysis across computing platforms. *Mol Biol Evol*, **35**, 1547-1549.

LeClere, S., Wu, C., Westra, P. & Sammons, R.D. (2018) Cross-resistance to dicamba, 2,4-D,

and fluroxypyr in *Kochia scoparia* is endowed by a mutation in an *AUX/IAA* gene.

Proceedings of the National Academy of Sciences, **115**, E2911-E2920.

Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics*, **25**, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N. et al. (2009) The Sequence

Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078-2079.

Ma, Q., Grones, P. & Robert, S. (2017) Auxin signaling: a big question to be addressed by small

molecules. *Journal of Experimental Botany*, **69**, 313-328.

Martin, S.L., Benedict, L., Wei, W., Sauder, C.A., Beckie, H.J. & Hall, L.M. (2020) High gene

flow maintains genetic diversity following selection for high *EPSPS* copy number in the

weed kochia (Amaranthaceae). *Scientific Reports*, **10**, 18864.

McCauley, C.L., McAdam, S.A.M., Bhide, K., Thimmapuram, J., Banks, J.A. & Young, B.G.

(2020) Transcriptomics in *Erigeron canadensis* reveals rapid photosynthetic and hormonal

responses to auxin herbicide application. *Journal of Experimental Botany*, **71**, 3701-3709.

Montgomery, J., Morran, S., MacGregor, D.R., McElroy, J.S., Neve, P., Neto, C. et al. (2024)

Current status of community resources and priorities for weed genomics research. *Genome*

Biology, **25**, 139.

Nyholt, D.R. (2000) All LODs are not created equal. *Am J Hum Genet*, **67**, 282-8.

Ossowski, S., Schneeberger, K., Lucas-Lledó, J.I., Warthmann, N., Clark, R.M., Shaw, R.G. et al.

(2010) The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*.

Science, **327**, 92-4.

Patterson, E.L., Saski, C.A., Sloan, D.B., Tranel, P.J., Westra, P. & Gaines, T.A. (2019) The draft

genome of *Kochia scoparia* and the mechanism of glyphosate resistance via transposon-

mediated *EPSPS* tandem gene duplication. *Genome Biol Evol*, **11**, 2927-2940.

Pettinga, D.J., Ou, J., Patterson, E.L., Jugulam, M., Westra, P. & Gaines, T.A. (2018) Increased

chalcone synthase (*CHS*) expression is associated with dicamba resistance in *Kochia*

scoparia. *Pest Management Science*, **74**, 2306-2315.

Poplin, R., Ruano-Rubio, V., DePristo, M.A., Fennell, T.J., Carneiro, M.O., Van der Auwera,

G.A. et al. (2018) Scaling accurate genetic variant discovery to tens of thousands of samples.

bioRxiv, 10.1101/201178201178.

Preston, C., Belles, D.S., Westra, P.H., Nissen, S.J. & Ward, S.M. (2009) Inheritance of

resistance to the auxinic herbicide dicamba in kochia (*Kochia scoparia*). *Weed Science*, **57**,

43-47.

Prusinska, J., Uzunova, V., Schmitzer, P., Weimer, M., Bell, J. & Napier, R.M. (2023) The

differential binding and biological efficacy of auxin herbicides. *Pest Management Science*,

79, 1305-1315.

Quastel, J.H. (1950) 2,4-Dichlorophenoxyacetic acid (2,4-D) as a selective herbicide. In

Agricultural Control Chemicals eds). American Chemical Society, pp. 244-249.

R-Core-Team, R: A Language and Environment for Statistical Computing. https://www.R-

project.org/ [accessed

Ramans-Harborough, S., Kalverda, A.P., Manfield, I.W., Thompson, G.S., Kieffer, M., Uzunova,

V. et al. (2023) Intrinsic disorder and conformational coexistence in auxin coreceptors. *Proc*

Natl Acad Sci U S A, **120**, e2221286120.

Ravet, K., Sparks, C.D., Dixon, A.L., Küpper, A., Westra, E.P., Pettinga, D.J. et al. (2021)

Genomic-based epidemiology reveals independent origins and gene flow of glyphosate

resistance in *Bassia scoparia* populations across North America. *Mol Ecol*, **30**, 5343-5359.

Ritz, C., Baty, F., Streibig, J.C. & Gerhard, D. (2015) Dose-response analysis using R. *PLOS*

ONE, **10**, e0146021.

Roux, F. & Reboud, X. (2005) Is the cost of herbicide resistance expressed in the breakdown of

the relationships between characters? A case study using synthetic-auxin-resistant

Arabidopsis thaliana mutants. *Genet Res*, **85**, 101-10.

Ruegger, M., Dewey, E., Gray, W.M., Hobbie, L., Turner, J. & Estelle, M. (1998) The TIR1

protein of *Arabidopsis* functions in auxin response and is related to human SKP2 and yeast

grr1p. *Genes Dev*, **12**, 198-207.

Song, Y. (2014) Insight into the mode of action of 2,4-dichlorophenoxyacetic acid (2,4-D) as an

herbicide. *Journal of Integrative Plant Biology*, **56**, 106-113.

Tan, X., Calderon-Villalobos, L.I., Sharon, M., Zheng, C., Robinson, C.V., Estelle, M. et al.

(2007) Mechanism of auxin perception by the TIR1 ubiquitin ligase. *Nature*, **446**, 640-5.

Tatematsu, K., Kumagai, S., Muto, H., Sato, A., Watahiki, M.K., Harper, R.M. et al. (2004)

MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with

the transcriptional activator NPH4/ARF7 to regulate differential growth responses of

hypocotyl and formation of lateral roots in *Arabidopsis thaliana*. *Plant Cell*, **16**, 379-93.

Todd, O.E., Figueiredo, M.R.A., Morran, S., Soni, N., Preston, C., Kubeš, M.F. et al. (2020)

Synthetic auxin herbicides: finding the lock and key to weed resistance. *Plant Sci*, **300**,

110631.

Todd, O.E., Patterson, E.L., Westra, E.P., Nissen, S.J., Araujo, A.L.S., Kramer, W.B. et al. (2024)

Enhanced metabolic detoxification is associated with fluroxypyr resistance in *Bassia*

scoparia. *Plant Direct*, **8**, e560.

Torra, J., Alcántara-de la Cruz, R., de Figueiredo, M.R.A., Gaines, T.A., Jugulam, M., Merotto,

Jr A. et al. (2024) Metabolism of 2,4-D in plants: comparative analysis of metabolic

detoxification pathways in tolerant crops and resistant weeds. *Pest Management Science*,

10.1002/ps.8373.

Uzunova, V.V., Quareshy, M., Del Genio, C.I. & Napier, R.M. (2016) Tomographic docking

suggests the mechanism of auxin receptor TIR1 selectivity. *Open Biol*, **6**, 160139.

Van der Auwera, G. & O'Connor, B. (2020) *Genomics in the Cloud: Using Docker, GATK, and*

WDL in Terra (1st Edition). O'Reilly Media

Walsh, T.A., Neal, R., Merlo, A.O., Honma, M., Hicks, G.R., Wolff, K. et al. (2006) Mutations in

an auxin receptor homolog *AFB5* and in *SGT1b* confer resistance to synthetic picolinate

auxins and not to 2,4-dichlorophenoxyacetic acid or indole-3-acetic acid in *Arabidopsis*.

Plant Physiol, **142**, 542-52.

Westra, E.P., Nissen, S.J., Getts, T.J., Westra, P. & Gaines, T.A. (2019) Survey reveals frequency

of multiple resistance to glyphosate and dicamba in kochia (*Bassia scoparia*). *Weed*

Technology, **33**, 664-672.

Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York

Wright, T.R., Shan, G., Walsh, T.A., Lira, J.M., Cui, C., Song, P. et al. (2010) Robust crop

resistance to broadleaf and grass herbicides provided by *aryloxyalkanoate dioxygenase*

transgenes. *Proc Natl Acad Sci U S A*, **107**, 20240-5.

Wu, C., Davis, A.S. & Tranel, P.J. (2018) Limited fitness costs of herbicide-resistance traits in

Amaranthus tuberculatus facilitate resistance evolution. *Pest Manag Sci*, **74**, 293-301.

Wu, C., LeClere, S., Liu, K., Paciorek, M., Perez-Jones, A., Westra, P. et al. (2021) A dicamba

resistance-endowing *IAA16* mutation leads to significant vegetative growth defects and

impaired competitiveness in kochia (*Bassia scoparia*). *Pest Manag Sci*, **77**, 795-804.

Zenser, N., Ellsmore, A., Leasure, C. & Callis, J. (2001) Auxin modulates the degradation rate of

Aux/IAA proteins. *Proceedings of the National Academy of Sciences of the United States of*

America, **98**, 11795-11800.

Zhang, R.-G., Li, G.-Y., Wang, X.-L., Dainat, J., Wang, Z.-X., Ou, S. et al. (2022) TEsorter: An

accurate and fast method to classify LTR-retrotransposons in plant genomes. *Horticulture*

Research, **9**, uhac017.

826