Hierarchical Bayesian modeling of multi-region brain cell count data

Sydney Dimmock[1]*, **Benjamin M. S. Exley[2]**, **Gerald Moore[3]**, **Lucy Menage[4]**, **Alessio Delogu[4]**, **Simon R. Schultz[3]**, **E. Clea Warburton[2]**, **Conor Houghton[1]**, **Cian O'Donnell[1,5]**

*For correspondence:
sydney.dimmock@bristol.ac.uk (SD);
conor.houghton@bristol.ac.uk (CH);
c.odonnell2@ulster.ac.uk (CO'D)

[1]School of Engineering Mathematics and Technology, University of Bristol, Michael Ventris Building, Woodland Road, Bristol, BS8 1UB, United Kingdom; [2]School of Physiology, Pharmacology and Neuroscience, University of Bristol, Biomedical Sciences Building, University Walk, Bristol, BS8 1TD, United Kingdom; [3]Centre for Neurotechnology and Dept of Bioengineering, Imperial College London, South Kensington, London SW7 2AZ, United Kingdom; [4]Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE5 9NU, United Kingdom; [5]School of Computing, Engineering & Intelligent Systems, Ulster University, Derry/Londonderry, BT48 7JL, Northern Ireland, United Kingdom

Abstract   We can now collect cell-count data across whole animal brains quantifying recent neuronal activity, gene expression, or anatomical connectivity. This is a powerful approach since it is a multi-region measurement, but because the imaging is done post-mortem, each animal only provides one set of counts. Experiments are expensive and since cells are counted by imaging and aligning a large number of brain sections, they are time-intensive. The resulting datasets tend to be under-sampled with fewer animals than brain regions. As a consequence, these data are a challenge for traditional statistical approaches. We present a 'standard' partially-pooled Bayesian model for multi-region cell-count data and apply it to two example datasets. These examples demonstrate that hierarchical Bayesian methods are well suited to these data. In both cases the Bayesian model outperformed standard parallel t-tests. Overall, inference for cell-count data is substantially improved by the ability of the Bayesian approach to capture nested data and by its rigorous handling of uncertainty in under-sampled data.

In studying the brain we are often confronted with phenomena that involve specific subsets of neurons distributed across many brain regions. Computations, for example, are performed by neuronal networks connecting cells in different parts of the brain. As another example, from development, neurons in different anatomical regions of the brain share the same lineage. Data for each of these types of experiment will be considered here, but the challenge is very general: how to measure and analyze multi-region neuronal data with cellular resolution.

In a typical cell-count experiment, gene expression is used to tag the specific cells of interest with a targeted indicator *Kawashima et al.* (*2014*). The brain is sliced, an entire stack of brain sections from a single animal is imaged and the images are aligned and registered to a standardized brain atlas such as the Allen mouse atlas (*Lein et al., 2007*; *Oh et al., 2014*; *Daigle et al., 2018*; *Harris et al., 2019*), the images are segmented into anatomical regions and the labeled cells in each region are counted. The resulting dataset consists of labeled cell counts across each of ~10–100 brain

regions. This technology is being deployed to address questions in a broad range of neuroscience subfields, for example: memory (*Kim and Cho, 2017*; *Haubrich and Nader, 2023*; *Dorst et al., 2024*), neurodegenerative disorders (*Liebmann et al., 2016*), social behavior *Kim et al.* (*2015*), and stress (*Bonapersona et al., 2022*).

Cell-counts are often compared across groups of animals which differ by an experimental condition such as drug treatment, genotype or behavioural manipulation. However, the expense and difficulty of the experiment means that the number of animals in each group is often small. Ten is a typical number of sample for these experiments, but fewer is not uncommon. This means that these data are undersampled: the dimensionality of the data, which corresponds to the number of brain regions, is much larger than the number of samples, which usually corresponds to the number of animals (*Figure 1*A). Current statistical methods are not well-suited to these nested wide-but-shallow datasets. Furthermore, because of the complicated preparation and imaging procedure, there is often missing data along with variability derived from experimental artifacts.

In cell count data there are two obvious sources of noise. The first of these is easy to describe: if a region has a rate that determines how likely a cell is to be marked for counting, then the actual number of marked cells is sampled from a Poisson distribution. The second source of noise is the animal to animal variability of the rate itself and this depends on diverse features of the individual animal and the experiment that are often unrelated to the phenomenon of interest. The challenge is to control for outliers and 'poor' data points whose rate is noisy, while extracting as much information as possible about the underlying process. Dealing with outliers is often an opaque and *ad hoc* procedure. It is also a binary decision, a point is either excluded so it does not contribute, or included, noise and all. This is where partial pooling helps. Partial pooling allows for the simultaneous estimation of parameters describing individual data points and parameters describing populations. This helps the data to self-regularize and elegantly balances the contribution of informative and weak observations to parameter values.

In recent years Bayesian approaches to data analysis have become powerful alternatives to classical frequentist approaches (*Gelman et al., 2013*; *McElreath, 2018*; *van de Schoot et al., 2021*). They have been applied to some types of neuroscience data including neurolinguistics (*Dimmock et al., 2023*), neural coding (*Brown et al., 1998*; *Zhang et al., 1998*), synaptic parameters (*Moran et al., 2008*; *Costa et al., 2013*; *Bird et al., 2016*; *Bykowska et al., 2019*) and neuronal-circuit connectivity (*Mishchencko et al., 2011*; *Cinotti and Humphries, 2022*). A Bayesian approach is particularly well suited to cell-count data but has not previously been applied to this problem.

A Bayesian approach formalizes the process of scientific inference, it distinguishes the data and a probabilistic mathematical model of the data. This model has a likelihood which gives the probability of the observed data for a given set of model parameters. The model often has a hierarchical structure which we compose to reflect the structure of the experiment and the investigators hypothesis of how the data depends on experimental condition. This hierarchy determines a set of *a priori* probabilities for the parameter values. The result of Bayesian inference is a probability distribution for these model parameters given the data, termed the posterior.

There are three advantages of a Bayesian approach that we want to emphasize, (1) while traditional multi-level models also allow a hierarchy (*Aarts et al., 2014*), Bayesian models are more flexible and the role of the model is clearer, (2) since the result of Bayesian inference is a probability distribution over model parameters, it indicates not just the fitted value of a parameter but the uncertainty of the parameter value. Finally (3) Bayesian models tend to make more efficient use of data and therefore improve statistical power.

A Bayesian model also includes a set of probability distributions, referred to as the prior, which represent those beliefs it is reasonable to hold about the statistical model parameters before actually doing the experiment. The prior can be thought of as an advantage, it allows us to include in our analysis our understanding of the data based on previous experiments. The prior also makes explicit in a Bayesian model assumptions that are often implicit in other approaches. However, having to design priors is often considered a challenge and here we hope to make this more straight-

Figure 1. Introduction. **A**: Each of N animals produce a cell-count from a total of R brain regions of interest. Cell-count data is typically undersampled with $N \ll R$. Scientists analyse the brain sections from experiment for positive signals. Here an example section is shown where teal points mark cells expressing the immediate early gene *c-Fos* (green and red lines indicate regions labeled as damaged). The final cell-count is equal to the sum of these individual items (sagittal brain map taken from the Allen mouse brain atlas: https://mouse.brain-map.org. **B**: Partial pooling is a hierarchical structure that jointly models observations from some shared population distribution. It is a continuum that depends on the value of the population variance τ. When $\tau = 0$ there is no variation in the population and each individual observation is modeled as a conditionally independent estimate of some fixed population mean θ (complete pooling). As τ tends to infinity observations do not combine inferential strength but inform an independent estimate γ_i (no pooling). In between the two extremes combine. Each observation can contribute to the population estimate while simultaneously supporting a local one to effectively model the variance in the data. The observed data quantities, y_i to y_n, are highlighted with a thick line in the model diagrams. **C**: An example of partial pooling on simulated count data. As the population standard deviation increases on the x-axis the individual estimates $\exp(\gamma_i)$ trace a path from a completely pooled estimate to an unpooled estimate. Circular points give the raw data values. Parameters are exponentiated because the outcomes are Poisson and so parameters are fit on the $\log$ scale.

forward by suggesting priors that are suitable for this class of data.

Here, our aim is to introduce a 'standard' Bayesian model for cell count data. We illustrate the application of this model to two datasets, one related to neural activation and the other to developmental lineage. For the second dataset, we also demonstrate a second example extension Bayesian model. In all cases, the Bayesian models produce clearer results than the classical frequentist approach.

Materials and Methods

Data

To illustrate our approach we consider two example applications, one which counts cells active in regions of the recognition memory circuit of rat during a familiarity discrimination task, and the other which examines the distribution of a specific interneuron type in the mouse thalamus.

Case study one - transient neural activity in the recognition memory circuit

The recognition memory network, *Figure 3*, is a distributed network which has been well studied using a variety of behavioural tasks. It includes the hippocampus (HPC) and perirhinal cortex (PRH), shown to deal with object spatial recognition and familiarity discrimination respectively (***Barker***

Figure 2. Methods. A table of partial pooling behaviour for different likelihood and prior combinations. Rows are the two prior choices for the population distribution, and columns the two distributions for the data. Within each cell the expectation of the marginal posterior $p(\exp(\gamma_i)|\theta, \tau, y)$ is plotted as a function of τ. The solid black line is the expectation of the marginal posterior $p(\theta|\tau, y)$ with one standard deviation highlighted in grey. **Top left**: Combining a normal prior for the population with a Poisson likelihood is unsatisfactory in the presence of a zero observation. The zero observations influence the population mean in an extreme way owing to their high importance under the Poisson likelihood. **Bottom left**: By changing to a horseshoe prior the problematic zero observations can escape the regularization machinery. However, regularization of the estimates with positive observations is much less impactful. **Top right**: A zero-inflated Poisson likelihood accounts for the zero observations with an added process, reducing the burden on the population estimate to compromise between extreme values. **Bottom right**: No model.

and Warburton, 2011; *Ennaceur et al., 1996*; *Norman and Eacott, 2004*); medial prefrontal cortex
(MPFC), concerned with executive functions such as decision making but also with working memory,
and the temporal association cortex (TE2) used for acquisition and retrieval of long-term object-
recognition memories (*Ho et al., 2011*). The nucleus reuniens (NRe) has reciprocal connectivity to
both MPFC and HPC (*Hoover and Vertes, 2012*) and for this reason it is also believed to be an im-
portant component of the circuit (*Barker and Warburton, 2018*, *2011*). In previous studies, lesions
of the NRe have been shown to significantly impair long-term but not short-term object-in-place
recognition memory (*Barker and Warburton, 2018*).

The data analysed in this case study were collected to investigate the role of the NRe in the
recognition memory circuit through contrasting the neural activation for animals with a lesion in
the NRe with neural activation for animals with a sham surgery. The immediate early gene *c-Fos* is
rapidly expressed following strong neural activation and is useful as a marker of transient neural
activity. Animals in the experiment performed a familiarity discrimination task (single-item recog-
nition memory), discriminating novel or familiar objects with or without an NRe lesion and the
number of cells that expressed *c-Fos* was counted in regions across the recognition memory cir-
cuit. The two-by-two experimental design allocated ten animals to each of the four experiment
groups {sham, lesion} × {novel, familiar} and cell counts were recorded from a total of 23 brain
regions. The visual cortex V2C and the motor cortex M2C were taken as control regions as they
were not expected to show differential *c-Fos* expression in response to novel or familiar objects
(*Exley, Masters Thesis, 2019*).

Figure 3. Recognition memory circuit. Schematic of the recognition memory network adapted from (*Exley, Masters Thesis, 2019*). Bold arrows show the assumed two-way connection between the medial prefrontal cortex and the hippocampus facilitated by the NRe. Colours highlight the HPC (red), MPC (blue) and specific areas of the rhinal cortex (yellow). The NRe was lesioned in the experiment.

Case study two - Ontogeny of inhibitory neurons in mouse thalamus and hypothalamus
The second dataset comes from a study *Jager et al.* (*2021*) that, in part, counted the number of
inhibitory interneurons in the thalamocortical regions of mouse. *Sox14* is a gene associated with
inhibitory neurons in subcortical areas. It is required for the development and migration of lo-
cal inhibitory interneurons in the dorsal lateral geniculate nucleus (LGd) of the thalamus *Jager*
et al. (*2016*, 2021). Consequently, *Sox14* is useful for identifying discrete neuronal populations in
the thalamus and hypothalamus (*Golding et al., 2014*; *Jager et al., 2016*, *2021*). The experiment
compared heterozygous (HET) and knock-out (KO) mouse lines. The HET knock-in mouse line
Sox14[GFP/+] marked *Sox14* expressing neurons with green fluorescent protein (GFP), the homozy-
gous KO mouse line *Sox14*[GFP/GFP], in contrast, was engineered to block expression of the endoge-
nous *Sox14* coding sequence *Delogu et al.* (*2012*). Each animal produced two samples, one for each

hemisphere. In total there are ten data points, six belonging to HET (three animals), and four to KO
(two animals). Each observation is 50 dimensional, corresponding to fifty individual brain regions
in each hemisphere.

Hierarchical modeling

Our goal in both cases is to quantify group differences in the data. We present a 'standard' hier-
archical model. This model reflects the experimental features common to cell count experiments
and reflects the hierarchical structure of cell count data; the standard model is designed to deal
robustly and efficiently with noise. On some occasions, to reflect a specific hypotheses, the struc-
ture of a particular experiment or an observed source of noise, this model can be further refined
or changed to target the analysis. We will give an example of this for our second dataset.

At the bottom of the model are the data themselves, the cell counts y_i. The index i runs over
the full set of samples, which in this case comprises 23 brain regions $\times$ 10 animals $\times$ 4 groups $\approx$ 920
datapoints in the first study, and 50 brain regions $\times$ 6 HET animals + 50 brain regions $\times$ 4 KO animals
$\approx$ 500 datapoints in the second. The basic assumption the model makes is that this count is derived
from an underlying propensity, $\lambda_i > 0$, which depends on brain region and, potentially, group:

$$y_i \sim \text{Poisson}(\lambda_i) \tag{1}$$

154 Hence the propensity λ_i is the mean of the Poisson distribution, and a statistical model is used to
155 describe the dependence of this parameter on brain region and animal. Since λ_i is strictly positive
156 a log-link function is introduced:

$$\log \lambda_i = \theta_{r[i],g[i]} + \gamma_i + E_i \tag{2}$$

157 where we have used 'array notation' *Gelman and Hill* (*2006*), mapping the sample index i to prop-
158 erties of the sample, so $r[i]$ returns the region index of observation i, and similar for $g[i]$ but for
159 groups and animals. The sample-by-sample variability is given by γ_i, this is modelled as Gaussian
160 noise:

$$\gamma_i \sim \text{Normal}(0, \tau_{r[i],g[i]}) \tag{3}$$

whose size depends on region and group. This equation demonstrates a potentially surprising
aspect of partially-pooled models: the over-parameterization.

Ignoring E_i for now, the rate has been split between two terms, $\theta_{r[i],g[i]}$ is the *fixed effect* which
is constant across animals and $\gamma_{r[i],a[i]}$ is the *random effect* which captures the animal to animal
variability. While θ_{rg} models the mean for log-cell-count for each region, given the condition; γ_i
models variation around this mean. For this reason, γ_i is assumed to follow a normal distribution
with zero mean. The regression term may appear over-parameterized, without θ_{rg} the γ_i could 'do
the work' of matching the data. However, the model is regularized by a prior; observations with
a weak likelihood will have their random effect γ_i shrunk towards the population location. The
amount of regularization depends on the variation in the population, a quantity that is estimated
from each likelihood. This is how partial pooling works as an adaptive prior for 'similar' parameters
(*Figure 1***B**). The data 'pools' some evidence while still allowing for individual differences in samples.

The final term is the exposure E_i. Cell counts may be recorded from sections with different
areas. The exposure term scales the parameters in the linear model as the recording area increases
(*McElreath, 2018*). In our model, the exposure is equal to the logarithm of the recording area; this
value is available as part of the experimental data.

The set of parameters $\tau_{r,g}$ models the population standard deviations of the noise for each
region r and animal group g. When working on the log scale, priors for these parameters are
typically derived in terms of multiplicative increases. Since the parameters are positive they are
assigned a half-normal distribution

$$\tau_{r,g} \sim \text{HalfNormal}(\log(s)) \tag{4}$$

Table 1. Parameter table for the hierarchical model.

Parameter	Description
E_i	Exposure.
κ_i	Horseshoe inflation.
π	Zero inflation.
γ_i	Random effect for observation i.
θ_{rg}	Fixed effect for region r in group g.
τ_{rg}	Scale of random effects for region r in group g.

with an appropriately chosen scale $s > 1$. For our analyses we used $s = 1.05$ because this gives a HalfNormal distribution with 95% of its density in the interval $[0, \log(1.1)]$. This translates into an approximate 10% variation around $\exp(\theta)$ at the upper end, which is a moderately informative prior, reflecting our belief that within-group animal variability is small relative to between-group variability. This regularization also helps model inference when the datasets are undersampled.

Horseshoe prior

Cell count data often has outliers, for example due to experimental artifacts. Since by default the likelihood does not account for these outliers, they may cause substantial changes in fitted parameter values. This is demonstrated in *Figure 2*, where a careless application of the Poisson distribution on data with several zero counts has a large influence on the posterior distribution. There are two general options for dealing with outliers: either modeling them in the likelihood or in the prior. Although the likelihood option is preferred as it is more direct, see our zero-inflation model below, it can be hard to design because it requires knowledge of the outlier generation process. The alternative is via a flexible prior such as the horseshoe *Carvalho et al.* (*2010*); *Piironen and Vehtari* (*2017*). This more generic option may be suitable as a default 'standard' approach in the typical case where outliers are poorly understood.

The horseshoe prior is a hierarchical prior for sparsity. It introduces an auxiliary parameter κ_i that multiplies the population scale τ. This construction allows surprising observations far from the bulk of the population density to escape regularisation.

$$\gamma_i \sim \mathrm{Normal}(0, \tau_{r[i],g[i]} \times \kappa_i) \tag{5}$$

$$\kappa_i \sim \mathrm{HalfNormal}(1). \tag{6}$$

An example of this is given in *Figure 2* as the bottom left cell of the 2×2 table of models. The horseshoe prior often uses a Cauchy distribution but in our case the heavy tail causes problems for the sampling algorithm, see SI section 8.

Zero inflation

A particular trait of the second dataset is that there are a large number of zero data points ($\sim 6\%$). Although a zero observation is always possible for a Poisson distribution, for plausible values of the propensity, zeros should be rare. It is likely that for some regions the experiment has not worked as expected and the zeros show that something has 'gone wrong' and that the readings are not well described by a Poisson distribution. Here we extend the model to include this possibility. This is a useful elaboration of the standard model. In the standard model the horseshoe prior ensures that these anomalous readings only have a small effect on the result but it is more informative to extend the model to include them. While this particular extension is specific to these data, it also serves as an example of how a standard Bayesian model can serve as a starting for point for an iterative investigation of the data.

The zero-inflated Poisson model is intended to model a situation where there are zeros unrelated to the Poisson distribution, in this case this might, for example, be the result of an error in

the automated registration process that identifies regions and counts their cells. It is a mixture
model, if

$$y_i \sim \text{ZIPoisson}(\pi, \lambda_i) \tag{7}$$

there is a probability π that $y_i = 0$ and a $1 - \pi$ probability that y_i follows a Poisson distribution with
rate λ_i. Importantly this means there are two ways in which y_i can be zero, through the Bernoulli
process parameterized by π, or through the Poisson distribution. This has the effect of 'inflating'
the probability mass at zero with the additional parameter π giving the proportion of extra zeros
in the data that could not be explained by the standard Poisson distribution. This distribution can
be visualized in Figure 2 and further mathematical details are described in SI 2C.

Model inference

Posterior inference was performed with the probabilistic programming language Stan (*Carpenter
et al., 2017*), using its custom implementation of the No-U-Turn (NUTS) sampler *Betancourt* (*2016*);
Hoffman et al. (*2014*). For each model, the posterior was sampled using four chains for 8000 itera-
tions, with half of these being attributed to the warm-up phase. This gives a total of 16000 samples
from the posterior distribution.

Results

We describe differences in estimated counts between groups in terms of $\log_2$-fold changes. Fold
changes are useful because they prevent differences that are small in absolute magnitude from be-
ing masked by regions with high overall expression. Our results compare Bayesian highest density
intervals (HDI) with the confidence interval (CI) from an uncorrected Welch's t-test. The Bayesian
HDI is calculated from the posterior distribution and is the smallest width interval that includes a
chosen probability, here 0.95 (to correspond to $\alpha = 0.05$), and summarizes the meaningful uncer-
tainty over a parameter of interest.

Case study one - transient neural activity in the recognition memory circuit

Results for the first dataset are presented in *Figure 4*. *Figure 4***A** plots cell-count differences be-
tween the novel and familiar conditions without lesion and *Figure 4***B** with lesion. These data were
collected to investigate the role of different hippocampal and adjacent cortical regions in memory.
However, some regions of interest, such as the intermediate dentate gyrus (IDG), and the dorsal
subiculum (DSUB) look under-powered: for both regions there is a markedly non-zero difference in
expression between the novel and familiar conditions in the sham animals but a wide confidence
interval overlapping zero makes the evidence unreliable (orange bars *Figure 4***A**).

In contrast, the Bayesian estimates (green bars *Figure 4*) produce a clear result. For a number of
brain regions in *Figure 4***A**, sham-novel animals have higher expression than sham-familiar ones.
These differences disappear in *Figure 4***B** with lesion-novel and lesion-familiar animals showing
roughly equal cell counts. This indicates that the difference is only present when the NRe is intact.

Case study two - Ontogeny of inhibitory interneurons of the mouse thalamus

For each of the fifty brain regions the estimated $\log_2$-fold difference in GFP expressing cells between
the two genotypes is plotted in *Figure 5*. This includes the purple and pink 95% HDI from the
horseshoe and zero-inflated Poisson models along with the 95% confidence interval arising from
a t-test in orange. For most brain regions, the two Bayesian models gave narrower HDIs than the
t-test CI. Accordingly, the Bayesian models identified a greater number of brain regions that had
genotype differences in *Sox14*-positive cell count in the sense that it found more places where the
appropriate uncertainty interval does not overlap zero.

Despite the large difference in interval estimation between the Bayesian HDIs and t-test CI for
many brain regions, as the data becomes stronger from the perspective of the frequentist p-value
towards the right-hand side of the second row in *Figure 5*, the model results become much more

Figure 4. Results—Case study 1. **A:** Heatmap of the raw log cell count data. Each row corresponds to a single animal, columns correspond to brain regions. Animals are grouped into lesion-familiar (LF), lesion-novel (LN), sham-familiar (SF) and sham-novel (SN). **B,C:** $\log_2$-fold differences for each surgery type: B shows differences between SF and SN groups, C shows differences between LF and LN groups. The 95% Bayesian HDI is given in green, and the 95% confidence interval calculated from a Welch's *t*-test in orange. Horizontal lines within the intervals mark the posterior mean of the Bayesian results, and the raw data means in the *t*-test case. The *x*-axis is ordered in terms of decreasing *p*-value from the significance test and ticks have been color-paired with the nodes in the recognition memory circuit diagram *Figure 3*. Black ticks are not present in the circuit because they are the control regions in the experiment.

Figure 5. Results—Case study 2. **A:** Heatmap of the raw log cell count data. Each row corresponds to a single animal, columns correspond to brain regions. L and R denote left and right hemispheres, respectively. **B:** $\log_2$ fold differences in GFP positive cells between mouse genotypes, heterozygous (HET) and knockout (KO), for each of the fifty recorded brain regions spread across two rows. The 95% Bayesian HDI is given in purple and pink for the Bayesian horseshoe and zero-inflated model. The 95% confidence interval calculated from a Welch's t-test in orange. Horizontal lines within the intervals mark the posterior mean of the Bayesian results and the data estimate for the t-test. The x-axis is ordered in terms of decreasing p-value from the significance test.

Figure 6. Example data and inferences highlighting model discrepancies. On the left under 'data': boxplots with medians and interquartile ranges for the raw data for four example brain regions. The shape of each point pairs left and right hemisphere readings in each of the five animals. On the right under 'inference': HDIs and confidence intervals are plotted. Purple is the Bayesian horseshoe model, pink is the Bayesian ZIP model, and orange is the sample mean. The Bayesian estimates are not strongly influenced by the zero-valued observations (MPN, SCH, TMd) or large-valued outliers (MH) and have means close to the data median. This explains the advantage of the Bayesian results over the confidence interval.

compatible. The variation within groups is very small for these regions. Further regularization is not necessary and so the impact of partial pooling has been reduced. The sample estimate of the t-test has 'caught up' to the regularized estimate because the signal is strong.

The zero-inflated Poisson distribution sometimes differs with the t-test confidence intervals. One example of this is the result for dorsal tuberomammillary nucleus (TMd). *Figure 6*, bottom row, plots the raw cell count values for TMd alongside the inferred frequentist mean and two Bayesian model means. For this region the HET animals have high GFP expression across both hemispheres, yet animal three has a reading of zero for both hemispheres. This injects variability into the standard deviation of the HET group. Consequently, the pooled standard deviation used in the t-test is large and almost certainly guarantees a non-significant result. Furthermore, the sample mean of this region looks nothing like zero, but also nothing like the other two animals with positive counts. In addition to the wide interval, the sign of the difference does not agree with the data. The medial preoptic nucleus (MPN) also suffers from poor estimation. Once again, this region contains a single HET animal for which the reading from both hemispheres is zero. The zero-inflated Poisson produces a posterior distribution of the appropriate sign with small uncertainty.

The two Bayesian models did not always agree. In some cases, such as the Medial habenula (MH) and Suprachiasmatic nucleus (SCH), the 'standard' horseshoe model does not show a genotype difference in cell counts, while the ZIP model indicates that heterozygotes had higher cell counts than knock-outs (*Figure 5*, *Figure 6*). The opposite can be seen in the case of the parvicellular ventral posteromedial nucleus of the thalamus (VPMpc), the horseshoe model suggests a genotype difference where the ZIP model did not (*Figure 5*). Further examination of the data shows why this happens, for example for region MH (*Figure 6*, top row), the 'standard' horseshoe model sensibly ignores the large positive outlier value in the heterozygote data, while the ZIP model does not. As a result the ZIP model's estimate for the mean is pulled upward, leading to an inferred

difference in heterozygote versus the wild-type.

Discussion

We have presented a standard workflow for Bayesian analysis of multi-region cell-count data. We propose a likelihood and appropriate priors with a nested hierarchical structure reflecting the structure of the experiment. We applied this to two distinct example datasets and demonstrate that they capture more fruitfully the characteristics of the data when compared to field-standard frequentist analyses.

For both case studies, the Bayesian uncertainty intervals are more precise than the confidence intervals. These confidence intervals tend to be quite wide on these data because of the small sample size and because of violations to their parametric model assumptions. Our standard workflow uses a horseshoe prior, along with the partial pooling, this allows our model to deal effectively with outliers. Furthermore, for the data sizes presented here, a full Bayesian inference using Stan does not require long computation time, or even particularly high performance hardware. Modern multi-core laptop processors are quite sufficient for this task. Fitting a model typically takes less than an hour.

In our analysis we have noted examples where different Bayesian models give discrepant conclusions. The obvious question to ask is which should we trust? The disappointing but inevitable answer is that, as with more traditional methods, Bayesian analysis is only a tool useful for interpreting data and brings with it a set of assumptions and biases regarding the experiment and the data. A Bayesian analysis does not avoid inconsistent or inconclusive results but it usually makes the assumptions more explicit and transparent. Typically the solution these model inconsistencies is to inspect the raw data and ask which model better captures those aspects of the data we are most interested in. Overall, the lesson here is that Bayesian hierarchical modeling has greater flexibility and statistical power but all statistical analyses, even those claiming to 'test hypotheses', just support exploration and it is ultimately the researcher's responsibility to make sure that a model's assumptions are appropriate and its behavior is sensible for the target dataset.

The horseshoe prior model workflow we have exhibited here is intended as a standard approach. We believe that, without extension, it will provide a robust model for cell-count data. However, we also suggest that the standard workflow can be a useful first step for a more comprehensive, extended, model when one is required. We have given an example of this for the second dataset where the anomalous zeros prompted us to change the likelihood to a zero-inflated Poisson. There are other possibilities, for example, zero inflation is not the only way to handle an anomaly in the number of zeros: the hurdle model is an alternative, *Cragg* (*1971*). This is not a mixture model, instead it restricts the probability of zeros to some value π with the probabilities for the positive counts coming from a truncated Poisson distribution. The hurdle model can deflate as well as inflate the probability mass at zero. This did not match the situation in the data we considered but might for other datasets. Another extension might involve tighter priors based on previous experiments. This is likely to be very relevant for cell-count data since these experiments are rarely performed in isolation and so prior information can be leveraged from a history of empirical results.

One obvious elaboration of our model would replace normal distributions with multivariate normal distributions. This would have two advantages. Firstly, correlations are difficult to estimate for under-sampled data. Including correlation matrix priors provides extra information, for example, based on anatomical connectivity, that can aid the statistical estimation of other parameters. Secondly, it would more closely match our understanding of the experiment: we know that activity is likely to be correlated across regions and so it is apposite to include that directly in the model. Unfortunately the problem of finding a suitable prior for the correlation proved insurmountable: the standard Lewandowski-Kurowicka-Joe distribution *Lewandowski et al.* (*2009*) which has been useful in lower-dimensional situations is too regularizing here. This is an area where further work needs to be done.

It is important to highlight that a mixed effects model is not a uniquely Bayesian construction. Indeed, any model that tries to include more sophistication through hierarchical structures, Bayesian or otherwise, is useful. However, non-Bayesian models can be complicated and opaque, they are also often more restrictive, for example they often assume normal distributions, and circumventing these restrictions can make the models even less transparent. A Bayesian approach is, at first, unfamiliar; this can make it seem more obscure than better established methods, but, in the long run, Bayesian models are typically clearer and do not involve so many different assumptions and so many fine adjustments.

Code and data availability

The code necessary to run the models presented in this manuscript can be found at *Dimmock et al.* (*2025*) and on our Github https://BayesianCellCounts.github.io. The data for case study one on nucleus reuniens lesion are available from https://doi.org/10.5281/zenodo.12787211 *Exley et al.* (*2024*). The data from case study two on *Sox14* expressing neurons are available from https://doi. org/10.5281/zenodo.12787287 *Gerald et al.* (*2024*).

Glossary

Likelihood	The probability of the observed data, under a given statistical model and choice of parameters.
Mixture model	A complicated probability distribution that is composed of two or more simpler probability distributions.
Partial pooling	A hierarchical modeling method that links estimates for similar-but-not-identical parameters, for example mean cell counts from multiple animals within the same experimental group.
Prior	Data-independent probability distributions describing plausible values for the statistical model's parameters. These are typically set 'prior' to observing the data, based on the analyst's domain knowledge.
Posterior	The final probability distribution over model parameters, inferred from both the data-independent prior and data-dependent likelihood.
Regularization	A family of methods for reducing model overfitting to noise in limited data, by biasing parameter estimates towards sensible values.

Acknowledgments

We are grateful to Andrew Dowsey and Matthew Nolan for useful discussion and helpful suggestions.

References

Aarts E, Verhage M, Veenvliet JV, Dolan CV, Van Der Sluis S. A solution to dependency: using multilevel analysis to accommodate nested data. Nature Neuroscience. 2014; 17(4):491–496.

Barker GR, Warburton EC. When is the hippocampus involved in recognition memory? Journal of Neuroscience. 2011; 31(29):10721–10731.

Barker GRI, Warburton EC. A Critical Role for the Nucleus Reuniens in Long-Term, But Not Short-Term Associative Recognition Memory Formation. Journal of Neuroscience. 2018; 38(13):3208–3217. https://www.jneurosci.org/content/38/13/3208, doi: 10.1523/JNEUROSCI.1802-17.2017.

Betancourt M. Identifying the Optimal Integration Time in Hamiltonian Monte Carlo. arXiv:160100225. 2016; .

Bird AD, Wall MJ, Richardson MJ. Bayesian inference of synaptic quantal parameters from correlated vesicle release. Frontiers in Computational Neuroscience. 2016; 10:116.

Bonapersona V, Schuler H, Damsteegt R, Adolfs Y, Pasterkamp RJ, van den Heuvel MP, Jo"els M, Sarabdjitsingh RA. The mouse brain after foot shock in four dimensions: Temporal dynamics at a single-cell resolution. Proceedings of the National Academy of Sciences. 2022; 119(8):e2114002119.

Brown EN, Frank LM, Tang D, Quirk MC, Wilson MA. A statistical paradigm for neural spike train decoding applied to position prediction from ensemble firing patterns of rat hippocampal place cells. Journal of Neuroscience. 1998; 18(18):7411–7425.

Bykowska O, Gontier C, Sax AL, Jia DW, Montero ML, Bird AD, Houghton C, Pfister JP, Costa RP. Model-based inference of synaptic transmission. Frontiers in Synaptic Neuroscience. 2019; 11:21.

Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P, Riddell A. Stan: A probabilistic programming language. Journal of Statistical Software. 2017; 76(1).

Carvalho CM, Polson NG, Scott JG. The horseshoe estimator for sparse signals. Biometrika. 2010; 97(2):465–480.

Cinotti F, Humphries MD. Bayesian mapping of the striatal microcircuit reveals robust asymmetries in the probabilities and distances of connections. Journal of Neuroscience. 2022; 42(8):1417–1435.

Costa RP, Sj"ostr"om PJ, Van Rossum MC. Probabilistic inference of short-term synaptic plasticity in neocortical microcircuits. Frontiers in Computational Neuroscience. 2013; 7:75.

Cragg JG. Some statistical models for limited dependent variables with application to the demand for durable goods. Econometrica: Journal of the Econometric Society. 1971; p. 829–844.

Daigle TL, Madisen L, Hage TA, Valley MT, Knoblich U, Larsen RS, Takeno MM, Huang L, Gu H, Larsen R, et al. A suite of transgenic driver and reporter mouse lines with enhanced brain-cell-type targeting and functionality. Cell. 2018; 174(2):465–480.

Delogu A, Sellers K, Zagoraiou L, Bocianowska-Zbrog A, Mandal S, Guimera J, Rubenstein JL, Sugden D, Jessell T, Lumsden A. Subcortical visual shell nuclei targeted by ipRGCs develop from a Sox14+-GABAergic progenitor and require Sox14 to regulate daily activity rhythms. Neuron. 2012; 75(4):648–662.

Dimmock S, Houghton C, O'Donnell C, BayesianCellCounts/DimmockEtAl2025-v1.01. GitHub; 2025. doi: 10.5281/zenodo.16340993.

Dimmock S, O'Donnell C, Houghton CJ. Bayesian analysis of phase data in EEG and MEG. eLife. 2023 sep; 12:e84602. https://doi.org/10.7554/eLife.84602, doi: 10.7554/eLife.84602.

Dorst KE, Senne RA, Diep AH, de Boer AR, Suthard RL, Leblanc H, Ruesch EA, Pyo AY, Skelton S, Carstensen LC, et al. Hippocampal engrams generate variable behavioral responses and brain-wide network states. Journal of Neuroscience. 2024; 44(2).

Ennaceur A, Neave N, Aggleton JP. Neurotoxic lesions of the perirhinal cortex do not mimic the behavioural effects of fornix transection in the rat. Behavioural Brain Research. 1996; 80(1):9–25. https://www.sciencedirect.com/science/article/pii/016643289600006X, doi: https://doi.org/10.1016/0166-4328(96)00006-X.

Exley B, Dimmock S, Warburton C, Exley_Warburton_NRe_lesion_cell_count_data. Zenodo; 2024. doi: 10.5281/zenodo.12787211.

Exley BMS, The role of the nucleus reuniens of the thalamus in the recognition memory network; Masters Thesis, 2019.

Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian data analysis. CRC press; 2013.

Gelman A, Hill J. Data analysis using regression and multilevel/hierarchical models. Cambridge University Press; 2006.

Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. Statistical Science. 1992; p. 457–472.

Gerald M, Sydney D, Menage L, Delogu A, Schultz S, Moore_Schultz_Sox14_expressing_neurons. Zenodo; 2024. doi: 10.5281/zenodo.12787287.

Golding B, Pouchelon G, Bellone C, Murthy S, Di Nardo AA, Govindan S, Ogawa M, Shimogori T, L"uscher C, Dayer A, et al. Retinal input directs the recruitment of inhibitory interneurons into thalamic visual circuits. Neuron. 2014; 81(5):1057–1069.

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, Bohn P, Caldejon S, Casal L, Cho A, et al. Hierarchical organization of cortical and thalamic connectivity. Nature. 2019; 575(7781):195–202.

Haubrich J, Nader K. Network-level changes in the brain underlie fear memory strength. eLife. 2023;
12:RP88172.

Ho JWT, Narduzzo KE, Outram A, Tinsley CJ, Henley JM, Warburton EC, Brown MW. Contributions of area Te2
to rat recognition memory. Learning & Memory. 2011; 18(7):493–501.

Hoffman MD, Gelman A, et al. The No-U-Turn sampler: adaptively setting path lengths in Hamiltonian Monte
Carlo. Journal of Machine Learning Research. 2014; 15(1):1593–1623.

Hoover WB, Vertes RP. Collateral projections from nucleus reuniens of thalamus to hippocampus and medial
prefrontal cortex in the rat: a single and double retrograde fluorescent labeling study. Brain Structure and
Function. 2012; 217:191–209.

Jager P, Moore G, Calpin P, Durmishi X, Salgarella I, Menage L, Kita Y, Wang Y, Kim DW, Blackshaw S, Schultz SR,
Brickley S, Shimogori T, Delogu A. Dual midbrain and forebrain origins of thalamic inhibitory interneurons.
eLife. 2021; 10:e59272.

Jager P, Ye Z, Yu X, Zagoraiou L, Prekop HT, Partanen J, Jessell TM, Wisden W, Brickley SG, Delogu A. Tectal-
derived interneurons contribute to phasic and tonic inhibition in the visual thalamus. Nature Communica-
tions. 2016; 7(1):13579.

Kawashima T, Okuno H, Bito H. A new era for functional labeling of neurons: activity-dependent promoters
have come of age. Frontiers in Neural Circuits. 2014; 8:37.

Kim WB, Cho JH. Encoding of discriminative fear memory by input-specific LTP in the amygdala. Neuron. 2017;
95(5):1129–1146.

Kim Y, Venkataraju KU, Pradhan K, Mende C, Taranda J, Turaga SC, Arganda-Carreras I, Ng L, Hawrylycz MJ,
Rockland KS, et al. Mapping social behavior-induced brain activation at cellular resolution in the mouse. Cell
Reports. 2015; 10(2):292–305.

Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, Boe AF, Boguski MS, Brockway KS, Byrnes EJ, et al.
Genome-wide atlas of gene expression in the adult mouse brain. Nature. 2007; 445(7124):168–176.

Lewandowski D, Kurowicka D, Joe H. Generating random correlation matrices based on vines and extended
onion method. Journal of Multivariate Analysis. 2009; 100(9):1989–2001.

Liebmann T, Renier N, Bettayeb K, Greengard P, Tessier-Lavigne M, Flajolet M. Three-dimensional study of
Alzheimer's disease hallmarks using the iDISCO clearing method. Cell Reports. 2016; 16(4):1138–1152.

McElreath R. Statistical rethinking: A Bayesian course with examples in R and Stan. Chapman and Hall/CRC;
2018.

Mishchencko Y, Vogelstein JT, Paninski L. A Bayesian approach for inferring neuronal connectivity from calcium
fluorescent imaging data. The Annals of Applied Statistics. 2011; p. 1229–1261.

Moran RJ, Stephan KE, Kiebel SJ, Rombach N, O'connor W, Murphy K, Reilly RB, Friston KJ. Bayesian estimation
of synaptic physiology from the spectral responses of neural masses. Neuroimage. 2008; 42(1):272–284.

Norman G, Eacott M. Impaired object recognition with increasing levels of feature ambiguity in rats with perirhi-
nal cortex lesions. Behavioural Brain Research. 2004; 148(1-2):79–91.

Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, Wang Q, Lau C, Kuan L, Henry AM, et al. A mesoscale
connectome of the mouse brain. Nature. 2014; 508(7495):207–214.

Piironen J, Vehtari A. Sparsity information and regularization in the horseshoe and other shrinkage priors.
Electronic Journal of Statistics. 2017 Jan; 11(2). http://dx.doi.org/10.1214/17-EJS1337SI, doi: 10.1214/17-
ejs1337si.

van de Schoot R, Depaoli S, King R, Kramer B, M"artens K, Tadesse MG, Vannucci M, Gelman A, Veen D, Willem-
sen J, et al. Nature Reviews Methods Primers. 2021; 1(1):1–26.

Vehtari A, Gelman A, Simpson D, Carpenter B, B"urkner PC. Rank-normalization, folding, and localization: an
improved R for assessing convergence of MCMC (with discussion). Bayesian Analysis. 2021; 16(2):667–718.

Zhang K, Ginzburg I, McNaughton BL, Sejnowski TJ. Interpreting neuronal population activity by reconstruc-
tion: unified framework with application to hippocampal place cells. Journal of Neurophysiology. 1998;
79(2):1017–1044.

Supplementary information

Full model

Here we present the complete mathematical model for each of of the three models applied in the main text. In all cases, the exposure term is only necessary for the first case study. Area was not available for the second so the exposure term was left out.

Poisson model

$$y_i \sim \text{Poisson}(\lambda_i) \tag{8}$$

$$\log \lambda_i = E_i + \gamma_i \tag{9}$$

$$\theta_{rg} \sim \text{Normal}(5, 2) \tag{10}$$

$$\gamma_i \sim \text{Normal}\big(\theta_{r[i],g[i]}, \tau_{r[i],g[i]}\big) \tag{11}$$

$$\tau_{rg} \sim \text{HalfNormal}(\log(1.05)) \tag{12}$$

Horseshoe model

$$y_i \sim \text{Poisson}(\lambda_i) \tag{13}$$

$$\log \lambda_i = E_i + \gamma_i \tag{14}$$

$$\theta_{rg} \sim \text{Normal}(5, 2) \tag{15}$$

$$\gamma_i \sim \text{Normal}\big(\theta_{r[i],g[i]}, \kappa_i \times \tau_{r[i],g[i]}\big) \tag{16}$$

$$\tau_{rg} \sim \text{HalfNormal}(\log(1.05)) \tag{17}$$

$$\kappa_i \sim \text{HalfNormal}(1) \tag{18}$$

Zero-inflated model

$$y_i \sim \text{ZIPoisson}(\lambda_i, \pi) \tag{19}$$

$$\pi \sim \text{Beta}(1, 5) \tag{20}$$

$$\log \lambda_i = E_i + \gamma_i \tag{21}$$

$$\theta_{rg} \sim \text{Normal}(5, 2) \tag{22}$$

$$\gamma_i \sim \text{Normal}\big(\theta_{r[i],g[i]}, \tau_{r[i],g[i]}\big) \tag{23}$$

$$\tau_{rg} \sim \text{HalfNormal}(\log(1.05)) \tag{24}$$

Distributions

Zero inflated Poisson distribution

The zero inflated Poisson distribution is a mixture distribution with mixing parameter π. The distribution is formally defined below. If,

$$p_Y(y) \equiv \text{ZIPoisson}(\pi, \lambda) \tag{25}$$

then

$$p_Y(y) = \begin{cases} \pi + (1 - \pi)\exp(-\lambda) & \text{if} \quad y = 0 \\ (1 - \pi)f(y) & \text{otherwise} \end{cases} \tag{26}$$

where

$$f(y) \equiv \text{Poisson}(\lambda) \tag{27}$$

Equivalently, the mixture can be specified with an indicator function.

$$p_Y(y) = \mathbf{1}_0(y)\pi + (1 - \pi)f(y) \tag{28}$$

Half Normal distribution

The HalfNormal distribution coincides with a zero-mean normal distribution truncated at zero. It has a single scale parameter σ. If,

$$p_Y(y) \equiv \operatorname{HalfNormal}(\sigma) \tag{29}$$

then

$$p_Y(y) = \frac{\sqrt{2}}{\sigma \pi} \exp\left(-\frac{y^2}{2\sigma^2}\right) \tag{30}$$

is the probability density function.

Additional methods

Fold differences

In our results we present differences between experimental groups in terms of $\log_2$-fold differences. We calculate this as follows. The parameter of interest $\theta_{r,g=i}$ is modeled on the natural log scale owing to the log-link function necessary for the Poisson regression. At the average animal the difference,

$$\theta_{r,g=i} - \theta_{r,g=j} = \log \frac{\exp(\theta_{r,g=i})}{\exp(\theta_{r,g=j})} \tag{31}$$

$$= \log \Delta_{ij} \tag{32}$$

is the natural log of the ratio of the expected counts. To obtain $\log_2$-fold differences we simply change the base by multiplying $\log \Delta_{ij}$ by $\log_2(e)$.

Data transformations

To facilitate a comparison with the Bayesian intervals in terms of $\log_2$-fold differences, it was necessary to add one to any zero counts before applying the t-test.

Non-centered parameterisation

Hierarchical models can produce geometry that is difficult for the sampler to explore. Fortunately, there exists a simple reparameterisation known as non-centering that can remedy this problem. In our model, instead of sampling γ_i directly, we sample the parameter $\tilde{\gamma}_i$ instead, and use it to reconstruct γ_i. That is, sample $\tilde{\gamma}_i$ from a standard normal distribution,

$$\tilde{\gamma}_i \sim \text{Normal}(0, 1) \tag{33}$$

and reconstruct γ_i as a deterministic function of sampled values of $\tilde{\gamma}_i$ and $\tau_{r[i],g[i]}$.

$$\gamma_i = \theta_{r[i],g[i]} + \tau_{r[i],g[i]} \times \tilde{\gamma}_i \tag{34}$$

This removes the frustrating joint behaviour between of $(\gamma_i, \tau_{r[i],g[i]})$, and promotes efficient sampling.

Preprocessing - case study 1

In these data some animals produced more than on reading per brain region. Before fitting our model these were summed together to produce a single count; the exposure term was also properly adjusted to correctly reflect the area of the recording site.

Software, packages and libraries

R Libraries	Version	Description
rstan	2.26.3	complete Stan library
cmdstanr	0.5.2	lightweight Stan library
HDInterval	0.2.2	calculating HDI in R
ggplot2	3.4.1	plotting
bayesplot	1.9.0	plotting
tidyverse	1.3.1	tibble, tidyr, readr, purr, dplyr, stringr, forcats

- R version 4.2.1–"Funny-looking-kid".
- Computation was performed locally on a Dell XPS 13 7390 laptop[1][2]
- Panels composed using inkscape version 1.2.2.

---

[1]Intel i7-10510U @ 1.80 GHz, 16 GB of RAM.
[2]Ubuntu 20.04.4 LTS.

Brain-region name acronyms

Here we include a list of acronyms for the brain regions in each case study.

Term	Definition
ACC	Anterior cingulate cortex.
DCA1/3	Dorsal CA1/3
DDG	Dorsal dendate gyrus.
DPC	Dorsal peduncular cortex.
DSUB	Dorsal subiculum.
HPC	hippocampus.
ICA1/3	Intermediate CA1/3
IDG	Intermediate dendate gyrus.
IFC	Infralimbic cortex.
LENT	Lateral entorhinal cortex.
MOC	medial orbital cortex.
MPFC	Medial prefrontal cortex
M2C	Motor cortex M2
NRe	Nucleus reuniens
PRL	prelimbic cortex.
PRH	Perirhinal cortex
PSTC	postrhinal cortex.
TE2	Temporal association cortex
VCA1/3	Ventral CA1/3
VDG	Ventral dendate gyrus.
VOC	Ventral orbital cortex.
VSUB	Ventral subiculum.
V2C	Visual cortex V2

Term	Definition
AHN	Anterior hypothalamic nucleus
ARH	Arcuate hypothalamic nucleus
CL	Central lateral nucleus of the thalamus
DMH	Dorsomedial nucleus of the hypothalamus
FF	Fields of Forel
IGL	Intergeniculate leaflet of the lateral geniculate complex
LD	Lateral dorsal nucleus of thalamus
LM	Lateral mammillary nucleus
LGv	Ventral part of the lateral geniculate complex
LGd	Dorsal part of the lateral geniculate complex
LH	Lateral habenula
LHA	Lateral hypothalamic area
LP	Lateral posterior nucleus of the thalamus
MD	Mediodorsal nucleus of thalamus
MGd	Medial geniculate complex, dorsal part
MGv	Medial geniculate complex, ventral part
MGm	Medial geniculate complex, medial part
MH	Medial habenula
MMme	Medial mammillary nucleus, median part
MPN	Medial preoptic nucleus
PH	Posterior hypothalamic nucleus
PMd	Dorsal premammillary nucleus
PMv	Ventral premammillary nucleus
PO	Posterior complex of the thalamus
POL	Posterior limiting nucleus of the thalamus

Term	Definition
PP	Peripeduncular nucleus
PR	Perireunensis nucleus
PVa	Periventricular hypothalamic nucleus, anterior part
PVH	Paraventricular hypothalamic nucleus
PVHd	Paraventricular hypothalamic nucleus, descending division
PVi	Periventricular hypothalamic nucleus, intermediate part
PVp	Periventricular hypothalamic nucleus, posterior part
RCH	Retrochiasmatic area
RT	Reticular nucleus of the thalamus
SBPV	Subparaventricular zone
SCH	Suprachiasmatic nucleus
SGN	Suprageniculate nucleus
SPFm	Subparafascicular nucleus, magnocellular part
SPFp	Subparafascicular nucleus, parvicellular part
SUM	Supramammillary nucleus
TMd	Tuberomammillary nucleus, dorsal part
TMv	Tuberomammillary nucleus, ventral part
TU	Tuberal nucleus
VAL	Ventral anterior-lateral complex of the thalamus
VMH	Ventromedial hypothalamic nucleus
VM	Ventral medial nucleus of the thalamus
VPL	Ventral posterolateral nucleus of the thalamus
VPM	Ventral posteromedial nucleus of the thalamus
VPMpc	Ventral posteromedial nucleus of the thalamus, parvicellular part
ZI	Zona incerta

Sampler diagnostics

The basic Poisson model (*Equation 1*) was sampled excellently. Measures of sampling performance such as $\hat{R}$ *Gelman and Rubin* (*1992*); *Vehtari et al.* (*2021*), and effective sample size, were all satisfactory (SI figure 1). Similarly, for the zero-inflated model (*Equation 7*), no problems were observed for any of the diagnostics (SI figure 2). Contrasting with this, the horseshoe model (*Equation 6*), exhibited some signs of fitting problems (SI figure 3). Divergences were not observed, and given the longer chain length this is reassuring evidence against biased computation. However for many parameters the effective sample size is much lower than we would like to see. This is reflected in the trace plots: the sampler is not making large jumps across the parameter space implying high auto-correlation and low effective sample size. Unfortunately, the horseshoe is notoriously hard to fit and we resort to brute-force methods such as increasing the number of iteration and reducing the step size of the sampler to improve the inference. In the following three plots diagnostics have been summarised with the following three items:

- **A**: The performance of the sampler is illustrated by plotting $\hat{R}$ (R-hat, $\hat{R} \approx 1$ ideal) against the ratio of the effective number of samples (larger is better) for each parameter in the model. Points represent individual parameters in the model, and have further been colour coded by their type. For example, all $\theta_{r,g}$ are coloured in green. Points have also been scaled based on how numerous the parameters are so the more numerous parameters have smaller dots, the less numerous, larger.
- **B**: A histogram comparing the marginal energy distribution π_E, and the transitional energy distribution $\pi_{\Delta E}$ of the Hamiltonian. Ideally, these distributions should match each other closely if the posterior distribution has been properly explored by the sampler.
- **C**: For each parameter type the parameter with the 'poorest' mixing (largest $\hat{R}$) are presented with a post-warmup trace plot that overlays the ordered sequence of samples from each of the four chains. Corresponding points in **A** are marked with a black border and zero transparency.

Figure 7. Diagnostics - Poisson. Standard Poisson model, case study - 1.

Figure 8. Diagnostics - Horseshoe. Horseshoe model, case study - 2.

Figure 9. Diagnostics - ZIPoisson. Zero-inflated Poisson, case study - 2.

Posterior predictive checking

Posterior predictive checks use the posterior predictive distribution,

$$p(y^{\text{rep}}|y) = \int p(y^{\text{rep}}|\theta)p(\theta|y)\,d\theta \qquad (35)$$

where $p(\theta|y)$ is the posterior distribution, and $p(y^{\text{rep}}|\theta)$ is the data distribution for y^{rep} that follows the same form as the likelihood for y. If we can verify that the posterior predictive distribution can generate replicate datasets with similar statistics to the observed data, then we might conclude that our model is consistent with the observed data and useful for answering questions about it. In practice a Monte-carlo approach is used to approximate statistics of the posterior predictive distribution. For example, if T is the test statistic of interest such as the sample mean, then

1. For $1, \ldots, S$.
   (a) sample $\theta_s \sim p(\theta|y)$
   (b) sample $y_s^{\text{rep}} \sim p(y^{\text{rep}}|\theta_s)$
   (c) calculate $T(y_s^{\text{rep}})$, where T is the statistic of interest
2. return $\{T(y_1^{\text{rep}}), \ldots, T(y_S^{\text{rep}})\} \approx p(T(y^{\text{rep}})|y)$

The posterior predictive checks that follow examine two important statistics of count data. 1) The standard deviation of the data (dispersion), as figure **a**. 2) The proportion of zeroes in the data (zero-inflation), as figure **B**. The value of the test statistic applied to the observed data $T(y)$ is plotted as a solid purple line, and the distribution of test statistics $\{T(y_1^{\text{rep}}), \ldots, T(y_S^{\text{rep}})\}$ as a purple histogram.

Figure 10. PPC - Poisson. Posterior predictive check for the standard Poisson model in case study - 1. **A**: The proportion of zeroes in the data matches the proportion of zeroes in posterior predictive samples. This proportion is zero. **B**: The distribution of standard deviations computed over a number of posterior predictive datasets (histogram), aligns with the standard deviation of the data.

Figure 11. PPC - Horseshoe. Horseshoe model, case study - 2. Posterior predictive check for the standard horseshoe model in case study - 2. **A**: The proportion of zeroes in the data is larger than those found in posterior predictive datasets. This makes sense, because the likelihood is still a Poisson distribution. **B**: The distribution of standard deviations computed over a number of posterior predictive datasets (histogram), aligns with the standard deviation of the data.

Figure 12. PPC - ZIPoisson. Zero-inflated Poisson, case study - 2. **A**: The proportion of zeroes in the data matches the proportion of zeroes in posterior predictive samples. **B**: The distribution of standard deviations computed over a number of posterior predictive datasets (histogram), aligns with the standard deviation of the data.

Horseshoe densities

In our model a horseshoe prior was used to allow some γ_i, typically those informed by $y_i = 0$, to escape regularisation by partial pooling. However, we encountered many problems with the default parameterisation that assigns a HalfCauchy density to the individual inflation parameters κ_i. In *Figure 13*A, the proportional conditional posterior density

$$, p(\tilde{\gamma}_i, \kappa_i | \theta, \tau, y_i) \propto p(y_i | \tilde{\gamma}_i, \theta, \tau, \kappa_i) p(\tilde{\gamma}_i) p(\kappa_i) \tag{36}$$

where θ and τ have been fixed, is plotted when y_i lies close to the population mean (left), or when it equals zero (right). Note that the x-axis is $\tilde{\gamma}_i$ and not γ_i because the model samples a non-centered parameterisation. That is,

$$\tilde{\gamma}_i = \frac{\gamma_i - \theta}{\tau \kappa_i}, \quad \tilde{\gamma}_i \sim \mathrm{Normal}(0, 1) \tag{37}$$

captures the deviations of γ_i around the population mean θ.

*Figure 13*B plots samples from the marginal posterior $p(\tilde{\gamma}, \kappa_i | y_i)$, when fitting to the data $\mathbf{y} = \{770, 820, 713, 541, 0, 0\}$. Each data point corresponds to one of the six plots in *Figure 13*B. This small example dataset is in fact the cell-counts for region TMd recorded from the heterozygote group in case study 2. The similarities in geometry can be readily seen with *Figure 13*A. A large number of divergences were produced by the sampler as demonstrated by the number of pink points compared to the non-divergent transitions in blue. For fixed τ the values $\tilde{\gamma}$ can take increase or decrease with smaller or larger κ respectively. The HalfCauchy places an extremely long right tail over κ that frustrates this relationship resulting in a posterior density that is difficult to sample from.

Figure 13. Horseshoe densities. **A**: Conditional posterior. **B**: MCMC pair plots. Divergent samples are coloured in pink, non-divergent in blue.

Modified horseshoe

For a Poisson model with parameters modeled on the log scale we consider the Cauchy parameterisation to be too extreme. In light of this we opted for a pragmatic approach, a modification to the original horseshoe by replacing the HalfCauchy distribution with a HalfNormal distribution. The modified horseshoe cuts off the top of the funnels by restricting κ to produce pleasant posterior geometry *Figure 13***A**, **B**. The modified horseshoe is much easier to sample from, but with the cost of a much more constraining prior over γ_i.

Figure 14. Modified horseshoe densities. **A**: The conditional posterior $p(\tilde{\gamma}, \kappa | \theta, \tau, y)$ when $y = 0$ (left) and $y \neq 0$ (right). **B**: MCMC pair plots of samples from the marginal posterior density $p(\tilde{\gamma}, \kappa | \mathbf{y})$.