Rapalink-1 reveals novel mTOR-dependent genes and an agmatinergic axis-based metabolic

feedback regulating mTOR activity and lifespan

Juhi Kumar[1] and Charalampos Rallis[1,*]

[1]Research Centre for Molecular Cell Biology, School of Biological and Behavioural Sciences, Queen Mary,

University of London, Mile End Road, London E1 4NS, UK.

[*]Author for correspondence: c.rallis@qmul.ac.uk

Abstract

The mechanistic Target of Rapamycin, mTOR, is a conserved pro-ageing pathway with characterised

inhibitors such as rapamycin, rapalogues and torins. Bi-steric third-generation inhibitors, such as rapalink-1

have been developed, however, their effects on organismal gene expression and lifespan have not been

characterised. Here, we demonstrate that rapalink-1 affects fission yeast spatial and temporal growth and

prolongs chronological lifespan with a distinct TORC1 selectivity profile. Endosome and vesicle-mediated

transport and homeostasis processes related to autophagy and Pik3, the orthologue of human PI3K, render

cells resistant to rapalink-1. Our study reveals mTOR-regulated genes with unknown roles in ageing

including all fission yeast agmatinases, the enzymes that convert agmatine to putrescine and urea. Through

genome-wide screens, we identify sensitive and resistant mutants to agmatine and putrescine. Genetic

interactome assays for the agmatinase *agm1* and further cell and molecular analyses, demonstrate that

impairing the agmatinergic branch of arginine catabolism results in mTOR activity levels that are beneficial

for growth but detrimental for chronological ageing. Our study reveals the anti-ageing action of agmatinases

within a metabolic feedback circuit regulating mTOR activity levels and ageing and with possible

implications to other systems, including human cells.

Key words: Ageing, TORC1, TORC2, agmatine, putrescine, polyamines, rapamycin, torin1, nutrient

response, *S. pombe*, MLN0128, sapanisertib

Introduction

Pharmacological inhibition of the evolutionarily conserved, nutrient-responsive and pro-ageing mechanistic Target of Rapamycin signalling pathway presents great interest in disease and biogerontology [1,2]. The centrepieces of the pathway are the mTOR PI3-related kinases that operate within two structurally and functionally distinct protein complexes, TORC1 and TORC2. The complexes are reported to have roles in both spatial and temporal aspects of cellular growth [3]. While TORC1 is responsible for promoting protein translation, lipid and central carbon metabolism and inhibiting autophagy, TORC2 regulates cytoskeleton and is required for cell survival [3–5]. Rapamycin, a macrolide able to inhibit TORC1 in an allosteric manner through interaction with the FKBP12 protein (Fkh1 in fission yeast), is shown to extend lifespan in cellular and animal models [1,6–8]. Torin1, a dual ATP-competitive inhibitor of TORC1 and TORC2 has also been shown to be beneficial for lifespan in various models including fission yeast [9]. Both drugs together with several rapalogues are used in numerous clinical trials (www.clinicaltrials.org).

Rapalink-1 is a third generation bi-steric mTOR inhibitor that combines rapamycin and sapanisertib (MLN0128) [10]. Its structure does not disrupt the binding of rapamycin to FKBP12 or the FRB domain of mTOR. While MLN0128 has been shown to have poor *in vivo* efficacy, its link to rapamycin has shown great promise in targeted TORC1 inhibition and cancer regression [10,11]. Rapalink-1 has been reported to prevent ethanol-induced senescence in endothelial cells [12]. Nevertheless, its precise effects on chronological lifespan have not been addressed and it is not known whether they will have resemblance to rapamycin or an ATP-competitive inhibitor such as Torin1.

Here, we have investigated the effects of rapalink-1 through comparisons with rapamycin using fission yeast, a relevant model in cell biology and ageing studies [13–15]. Both drugs advance mitosis in fission yeast, albeit with different profiles and kinetics, without halting the cell cycle like the pan-mTOR inhibitor torin1 [9] and can prolong chronological lifespan. The effects observed on gene expression following rapalink-1 treatment, are consistent with TORC1 inhibition [7–9,16] although new mTOR-dependent genes are revealed. Using a combination of classical cell biology and genome-wide cell-based screens we uncover new mTOR-related genes and demonstrate that the enzymes converting agmatine to urea and the polyamine putrescine (agmatinases) are a pivotal pathway that positively impacts lifespan. Microbiota-derived agmatine has been previously reported to prolong host lifespan due to changes in lipid homeostasis. However, here, and for the first time we reveal that the function of enzymes breaking down agmatine is linked to ageing. Analysis of

genetic interactome of agmatinase *agm1* [17,18] together with cellular and molecular analyses, uncover a

metabolic regulatory feedback loop that impacts on the regulation of TORC1 activity itself. Our data, reveal

a more complex image around the agmatinergic function: while agmatine may affect metabolism and

lifespan, agmatinases are pivotal in maintaining low mTOR activity levels during TORC1-inhibition states,

such as starvation or dietary restriction, with clear positive consequences on lifespan. Our data have possible

implications to other organisms, including human cells and provide additional novel information on arginine

metabolism and the role of polyamines [19,20] and polyamine-generating enzymes in ageing.

Results

Rapalink-1 affects cell cycle progression, cell size and TORC1-dependent processes in fission yeast.

Rapalink-1 is reported to preferentially inhibit TORC1 [10,11] in human cells. We, therefore, wished to

investigate its effects through comparing it with rapamycin, a well-characterised and widely used TORC1

inhibitor and without the complications of dual (TORC1 and TORC2) inhibitors such as torin1 [21,22].

Rapamycin-induced inhibition of mTOR advances mitotic onset, mimicking the reduction in cell size at

division observed following shifts to poor nitrogen sources [16]. We, therefore, analysed related effects when

cells are treated with 100nM rapamycin and rapalink-1 as this concentration of rapamycin has been shown

previously to inhibit TORC1-related processes in fission yeast, [7,8]. In rich YES media, treatment with

rapamycin and rapalink-1 leads to increased mitotic index (49% and 37% respectively compared to 27% in

control cells), peaking at 30-minute post treatments (Figure 1A). Similarly, when cells are treated with

rapamycin in EMM2 the mitotic index peaks at 30 minutes post-treatment (44% compared to 20% in control

cells) returning to levels comparable with the untreated cells following this (Figure 1B). Treatment with

rapalink-1 leads to a milder but sustained increase in the mitotic index (27%, 26% and 32% at 30', 60' and

120' respectively). Along with mitotic onset, treatments with both rapamycin and rapalink-1 result in

decreased cell size upon division in rich YES media (see 30'-120' time course in Figure 1C). Rapamycin has

a more severe effect on cell size compared to rapalink-1, at least in these media (Figure 1C) with gradual

decrease over time (from 30 to 120 minutes) while cells following rapalink-1 treatment decrease their size in

30 minutes and remain in the same levels for the whole examined period of two hours (Figure 1C).

Likewise, in minimal EMM2 media, cell sizes upon division decrease for both rapamycin and rapalink-1

(Figure 1D). The patterns of cell sizes differ between the two drugs: while in rapamycin cells reach their

minimum size in 60 minutes (12.4 μm) and then start to gradually recover, in rapalink-1 they decrease in 90

minutes (13.5 μm) and remain at the same size range for the duration (120') of the assay (Figure 1D).

Our septation and cell size results indicate phenotypes related to TORC1 inhibition as previously

shown and described [7–9,16]. We, therefore, investigated whether drug treatments at these concentrations affect

multiple aspects of TORC1 functions [4]. Negative cross-regulation between the AMPK and TORC1 pathways

have been previously reported in fission yeast [23,24]. We, therefore, examined the phosphorylation status of the

fission yeast serine/threonine protein kinase AMPK catalytic (alpha) subunit Ssp2. Following two and five

hours of treatment with 100nM of rapamycin or rapalink-1, no observable changes in Ssp2 phosphorylation

are found (Figure 1E). TORC1 negatively regulates autophagy [4]. Examination of GFP-Atg8 patterns [25]

following drug treatment in the aforementioned time points, do not show a significant degree of processing [25]

indicating that autophagy may not be upregulated or that it may be upregulated (albeit only marginally) in

these conditions (Figure 1E). Contrary to autophagy, global protein translation might be suppressed five

hours post-treatment with both drugs, as seen through phosphorylation of eIF2α while its total levels remain

unchanged at both two-hour and five-hour time points examined. Finally, we wondered whether drug

treatment can affect the activity of Maf1, a Pol III repressor [26], through its TORC1-dependent

phosphorylation status [23]. Immunoblotting for Maf1-pK revealed that following treatment with both

rapamycin and rapalink-1, Maf1-pK becomes dephosphorylated, more evidently in five hours of treatment

(Figure 1E).

We have, previously, shown that the GATA transcription factor Gaf1 mediates Pol II- and Pol III-

mediated transcription downstream of TORC1 [9]. Gaf1 is localised mainly in the cytoplasm and upon TORC1

inhibition translocates to the nucleus and binds target genes [9]. We examined localisation of Gaf1-GFP

(controlled by the endogenous Gaf1 promoter as described in Rodríguez-López et al., 2020). Gaf1-GFP

signal is found throughout the cell in untreated cells (Figure 1F). Treatment with rapamycin and rapalink-1

results in nuclear localization of Gaf1-GFP within 5 minutes (Figure 1F) indicating that both drugs can

inhibit TORC1 at sufficient levels to allow Gaf1 dephosphorylation and translocation. One of the major

actions of Gaf1 is to repress all the Pol III transcribed tRNAs in fission yeast upon TORC1 inhibition. The

result of Gaf1-GFP localisation agrees with Maf1 activation seen through Maf1-pK dephosphorylation

(Figure 1E). Overall, our results show that rapalink-1 targets fission yeast TORC1 and, like rapamycin, can

inhibit several but not all aspects of TORC1-dependent functions in the examined concentrations.

We then assessed the chronological lifespan (CLS) of fission yeast cells when treated with

rapamycin and rapalink-1 in YES media and at OD_{600} 0.5 at 100nM concentration. As expected, rapamycin

induces CLS extension (Fig.1G, log-rank p=$1.25*10^{-12}$). Similarly, rapalink-1 prolongs lifespan in similar

fashion to rapamycin (Fig.1H, p=$2.2*10^{-12}$) demonstrating the potential of the drug and its action in ageing

through TORC1 inhibition. Rapalink-1 lifespan extension depends on *fkh1* (Fig. 1H, p=0.047) as has

previously been observed in the case of rapamycin [8] demonstrating that the effects of rapalink-1 in these

conditions are TORC1-dependent.

A genome-wide screen for rapalink-1 resistance highlights roles of endosome-related processes in

growth and lifespan

Previous data in fission yeast have shown that mTOR inhibition results in lifespan extension with (in

the case of torin1) or without (in the case of rapamycin) cell-cycle progress inhibition. Fast growing cultures

of wild-type fission yeast cells at OD_{600} 0.2 were treated either with DMSO (control) or rapamycin and

rapalink-1 at a 100nM concentration. Cultures were placed at a Biolector setup as previously described [8] and

growth was monitored continuously through light scattering measurements [7,8,27]. While no lag phase

following drug treatment is observed (Figure 2A), a mild growth effect is seen in this assay consistent with

results previously seen with rapamycin [8]. We then performed a genome-wide fitness screen for rapalink-1

using the Bioneer deletion library [28] at two rapalink-1 concentrations (100 nM and 300 nM, Figure 2B) and

at conditions that each mutant is assessed and measured in quadruplicate [7,9]. Results from the screens

correlate well (Pearson cor.coeff=0.73, Figure 2C, supplemental table 1). Sensitive strains are related to

protein translation, sulphur metabolism and mitochondrial membrane organisation among others (Figure 2D)

representing proteins and groups related to anabolic functions and cellular growth [29]. Resistant strains are

related to CAMKK-AMPK signalling, vacuole and vesicle-mediated transport and autophagy (Figure 2E).

The fitness signatures acquired are consistent with TORC1 inhibition with GO groups functionally linked to

transport of macromolecules, processing of proteins, membranes and autophagic functions (Figure 2F)

representing fundamental metabolic functions in the crossroads of catabolism and growth [30]. Most of these

mutants are required for normal lifespan [31] and they have been identified in previous mTOR inhibition

screens from our group [9] and others [32]. We wondered whether such factors are required for rapalink-1-

dependent lifespan extension. We have, therefore, performed CLS analysis for the related *pik3* mutant as a

proof of principle. *Pik3Δ* cells are short-lived compared to *wt* (Figure 2G, compare with Figure 1H)

demonstrating that it is required for normal chronological lifespan in fission yeast. Nevertheless, rapalink-1

prolongs lifespan of *pik3* mutant cells (log rank, p=10^{-5}). To quantify the role of Pik3 in rapalink-1-mediated

longevity and directly compare with the rapalink-1 effects on *wt* cells, we calculated the areas under the

curve (AUCs, measured as days × percentage of survival from lifespan assays). In wild-type cells, the

lifespan was prolonged from an average AUC of 692 to 2,351 (increase of 1,659), whereas in *pik3Δ* cells, the

lifespan was prolonged to a lesser extent, from an average AUC of 473 to 1431 (increase of 958) (Figure

2H). A major difference between the rapalink-1 lifespan increase between normal and *pik3Δ* mutant cells is

in the first day of CLS: rapalink-1 can prolong the healthspan of *wt* in this timeframe (cell population retains

viability close to 100%, Figure 1H).

Rapalink-1-dependent gene expression points to TORC1-related genes with both known and unknown

roles in ageing

159   Beyond proteostasis, autophagy and global translation levels, TORC1 activity controls gene

160   expression at the level of transcription through multiple transcription factors [5,33,34]. We therefore performed

161   RNA-seq analysis on fission yeast cells treated with 100nM rapamycin and rapalink-1 for 5 hrs. This

162   timepoint was chosen based on the observed changes in the relevant markers described at Figure 1E.

163   PCA analysis shows that Rapalink-1 induces a stronger effect on gene expression compared to

rapamycin with the PCA1 component accounting for 71.36% difference between control, rapamycin and

rapalink-1 datasets separating well rapalink-1-treated transcriptomes from untreated and rapamycin-treated

states (Figure 3A). In these conditions, 17 genes are 2-fold, 70 genes 1.5-fold and 277 genes 1.2-fold

upregulated in rapamycin-treated cells compared to untreated cells while 2, 4 and 87 genes are

downregulated at the 2-fold, 1.5-fold and 1.2-fold levels respectively (Figure S1A, supplemental tables 2 and

3). In the case of Rapalink-1-treated cells, 190 genes are 2-fold, 355 genes 1.5-fold and 701 genes 1.2-fold

upregulated while 4, 28 and 516 are downregulated (Figure S1B) compared to untreated cells (see

supplemental tables 4 and 5). Interestingly, our analyses reveal 277 genes upregulated and 87 downregulated

genes in rapalink-1 compared to rapamycin (Figure S1C). Following rapamycin treatment, the

downregulated gene list is enriched among others in cytoplasmic translation, tRNA metabolic process and

transcription by polymerase III (Figure 2B). Upregulated genes are involved in cell cycle and specifically in

mitotic prometaphase. On the other hand, the gene list of rapalink-1-upregulated genes are enriched in

vacuolar transport and organisation. This has been previously observed in the case of torin1-treated fission

yeast cells ([9]. The subcellular localisation of TORC1 in fission yeast [35] is predominantly on the vacuole. It is

interesting that factors related to biogenesis and dynamics of the organelle where TORC1 resides on,

transcriptionally depend on TORC1 activity. In addition, small molecule, amino acid metabolic processes

and transmembrane transport are also enriched categories, among others (Figure 2B). While both rapamycin

and rapalink-1 target TORC1 and modulate lifespan in a similar fashion (in terms of lifespan extension

patterns), it may be that they do so through the regulation of common as well as distinct genes, at least, in the

given analysed conditions. We, therefore, examined the overlaps of upregulated and downregulated genes for

the two drugs as well as genes that are apparently regulated differently between the two pharmacological

treatments. 541 upregulated (Figure 2C) and 475 downregulated (Figure 2D) genes are unique to rapalink-1.

Genes that are upregulated in rapalink-1 compared to rapamycin are related to small molecule catabolic

processes, arginine and proline metabolism, glycolysis, transmembrane transport and small molecule

catabolic processes (just to name a few, see Figure S1D) while downregulated genes are related to ribosome

biogenesis, translation, amino acid and tRNA metabolic processes (Figure S1E).

High-throughput phenotyping approaches have majorly contributed to characterising the

chronological lifespan of haploid fission yeast mutants [31,33]. TORC1 is a major pro-ageing pathway, and

TORC1-dependent genes are likely to be related in lifespan regulation. We, therefore, examined whether

differentially expressed genes from rapalink-1-treated cells are linked with long or short reported lifespans of

the corresponding mutants. While many rapalink-1 upregulated genes have reported lifespans, a significant

number (331 genes) have no lifespan annotations (Figure 3E). GO enrichment of these genes demonstrates

that they are related to amino acid transport, amide transport, galactose, tyrosine arginine and proline

metabolism (Figure S2A). Likewise, 150 genes that are found to be downregulated in rapalink-1 treated cells

have not been annotated for lifespan (Figure S2B) and these are related to L13a-mediated translational

silencing, ribonucleoside biosynthesis, response to stimulus as well as to alanine, aspartate and glutamate

metabolism (Figure S2C). Examination of upregulated genes not linked yet to lifespan, revealed that arginine

catabolism and specifically all the agmatinase enzymes involved in processing agmatine to putrescine are

boosted following TORC1 inhibition (Figure 3F). We therefore wondered whether agmatinases are required

for normal chronological lifespan. CLS assays reveal that *agm1Δ* (log rank p=9.7x10^{-5}, Figure 3G), *agm2Δ*

(log rank p=1.5x10^{-5}, Figure 3H) and *agm3Δ* (log rank, p=3.9x10^{-5}, Figure 3I) mutant cells are shortlived

compared to wild-type cells. In addition, cells mutant for both *agm1* and *agm3* are also short-lived (log rank

p=3.3x10^{-6}, Figure 3J).

Agmatine and putrescine affect growth, metabolism and lifespan

The requirement of agmatinases in normal chronological lifespan led us to wonder about the effects

of supplementing agmatine and putrescine on the fitness and lifespan of fission yeast cells. Towards this, we

have performed genome-wide screens for both agmatine and putrescine in concentrations that are not toxic

and only induce a very small delay in growth (15mM agmatine and 25mM putrescine, data not shown). Each

mutant has been assessed in quadruplicate and fitness ratios have been obtained through normalisations with

untreated copies of the deletion library [28,36]. Good quality data have been obtained for 3,208 mutants (Figure

4A). Fitness ratios of the mutant strains between the two screens are well correlated (Pearson cor.coeff=0.74,

Figure 4B) showcasing that they have similar metabolic and growth outcomes. Gene ontology enrichments

for sensitive and resistant strains (cutoffs set at >1.35 and <0.65 fitness ratios for resistant and sensitive,

respectively) show overlapping and unique enrichments (Figure 4C, Supplemental Tables 6 for agmatine and

putrescine fitness ratios of all mutants). Both compounds render cells with gene deletions related to arginine

metabolism, as sensitive. However, while putrescine affects more strongly ribosome-related mutants,

agmatine affects mutants related to cytosolic transport or response to oxygen containing compound (Figure

4C for a complete list of GOs). On the other hand, mutants related to the CAMKK-AMPK signaling cascade

(as in the case of rapalink-1) and intracellular transport and tRNA wobble uridine modification are resistant

to both agmatine and putrescine. The list of resistant strains to putrescine is enriched in categories related to

polyphosphate metabolic process, cellular respiration and chaperone-mediated folding among others (Figure

4C). Interestingly, GO enrichments for strains resistant to agmatine include vacuolar transport and

organisation and are reminiscent of lists of mutant strains resistant to rapalink-1 (genes coding for

endolysosomal-related proteins). This result together with the requirement of agmatinase enzymes in normal

fission yeast chronological lifespan led us to wonder whether agmatine and putrescine supplementation can

be beneficial to *S. pombe* CLS. Indeed, addition of both compounds to fast growing wild-type *S. pombe*

cultures at the same concentrations used in our genome-wide screens lead to lifespan extension with

agmatine having more profound effects (Figure 4D, log rank p=5.3x10^{-11} ; 4E p=9.4x10^{-10}). Our results are

consistent with previous observations on the positive effects of agmatine in lifespan and ageing [37]. We

examined whether agmatinases are required for the lifespan extension and repeated the experiments in the

genetic background of *agm1Δ* (Figure 4F, log rank p=4.4x10^{-13} ; 4G, p=4.5x10^{-10}), *agm2Δ* (Figure 4H, log

rank p=3.5x10^{-11} ; 4I, p=3.5x10^{-8}) *and agm3Δ* (Figure 4J log rank p=1.1x10^{-13} ; 4K, p=2x10^{-9}), gene

deletions. Lifespan extensions comparable with the wild-type cells are observed in all single mutant

backgrounds. The same beneficial effects are also observed in double *agm1Δ agm3Δ* mutant cells (Figure

4L log rank p=4.1x10^{-10} ; 4M, p=3x10^{-10}). This might be due to either functional compensation or beneficial

effects related to other pathways such as lipid metabolism [37] or through oxidative stress protection [38].

***Agm1* interactome reveals metabolism-mediated mTOR activity regulation with effects on growth and**

lifespan

To understand the mechanisms underlying the requirements of agmatinases' functions in ageing we

performed a Synthetic Genetic Array (SGA) for *agm1* using *ade6::natMX6* as a control SGA query. As the

deletion library strains carry *ade6* auxotrophy alleles [28,36], the control query/SGA does not affect the fitness

profiles of the deletion collection as previously reported in published methods [7]. Following normalisations

for plate positional effects, low fitness mutants, and linkage we have isolated data for 3,108 double mutants

with *agm1* (Supplemental Table 7). Physical mapping of the isolated genetic interactions is reassuring of the

successful double mutants' generation as it demonstrates the characteristic accumulation of negative values

around the *agm1* locus and the positive values around the control *ade6* locus (Figure 5A). These values have

been filtered out during our quality control steps. With interaction value cutoffs of -15 and +15 (see materials

and methods) we find 179 mutants to show negative interactions and 284 mutants exhibiting positive

interactions with *agm1* (Figure 5B). The negative interaction list is enriched for genes coding for post-

transcriptional regulation, carbon metabolism, and reticulophagy among other GO terms (Figure 5C,

Supplemental Table 7). Positive interactions are enriched in amide metabolism, stress response and nitrogen

metabolism (Figure 5D, Supplemental Table 7).

Close inspection of the interactors' list reveals that genes coding for stress and amino acid sensing

that inhibit TORC1 such as *gcn1* [39] and *fil1* [40] positively interact with *agm1* (Figure 5E, green fonts). On the

other hand, the TORC1 core component *tco89* [3], negatively interacts with *agm1* (Figure 5E, red fonts). These

results could mean that absence of *agm1* (and possibly the rest of the agmatinases) promotes growth that is

enhanced through the absence of stress response players. The cellular growth enhancement is TORC1-

dependent as absence of *tco89* results in reduced cellular fitness/growth. Therefore, in this scenario,

agmatinases may provide a metabolic regulatory feedback for TORC1 activity in fission yeast: when mTOR

is inhibited, cells process vacuole-stored amino acids such as arginine [41], thus, ensuring that mTOR

inhibition signals are maintained, relayed within the cell and preventing anabolism while preparing for

catabolic processes and autophagy. In this case, *agm* mutants would showcase phenotypes related to

increased TORC1 activity. Indeed, we have found that all agmatinase deletion mutants have short

chronological lifespans compared to *wt* cells (Figure 3G-J). *Agm1* mutants have already been reported to be

fast-growing [42] and to have decreased mating efficiency [43], both cellular characteristics of increased TORC1

activity. We have then assessed cell size upon division for normal (wt) and agmatinase mutants. Our results

demonstrate that *agm* mutant cells are larger (Figure 5F, ***: p<0.001, Wilcoxon testing) strengthening the

case of increased TORC1 levels as the latter controls both temporal and spatial aspect of fission yeast

growth. TORC1 activity regulates multiple targets related to growth that are mediated through several

downstream kinases such as Sck1, Sck2, Psk1 and Gad8. The activity of these downstream of mTOR can be

visualised using phospho-Akt target antibody. Although different targets seem to be phosphorylated within

the different mutants (Figure 5G) the overall activity seems is enhanced (Figure 5H). Interestingly, while

tco89 mutant cells are long-lived [8], *agm1Δ tco89Δ* are short-lived as single *agm1* mutants (Figure 5I). This

result shows that in the case of the chronological lifespan phenotype, *agm1* is epistatic of *tco89*. In addition,

while *tco89* negatively interacts with *agm1*, the effect on anabolism and growth is profound but does not

reflect, in this case, to the ageing phenotype. Our data reveal that the agmatinergic branch of arginine

catabolism is favoured during mTOR inhibition and forms a regulatory loop required for maintaining lower

mTOR level, according to what environmental pressures dictate (Figure 5J).

Discussion

In this study we have assessed the cellular and molecular effects of the bi-steric mTOR inhibitor

rapalink-1 using fission yeast, a genetically tractable eukaryotic model. Our results show that rapalink-1

targets TORC1, prolongs chronological ageing and demonstrate that mTOR activity negatively regulates

agmatinase genes. In fast growing conditions and nutrient availability, mTOR is highly active. The cells

retain mTOR activity, thus, enhancing anabolism and promoting growth and, therefore, repress agmatinase

genes and the catabolic processing of arginine pools. This metabolic circuit ensures that arginine, stored

within vacuoles [41], continues to provide mTOR activation input. In stress conditions (including stationary

phase and during chronological lifespan) and when mTOR activity is lower [44], cells catabolise arginine. Once

again, this ensure that in such conditions, cells upregulate defensive mechanisms, scavenge and recycle

nutrients and prevent anabolism. Fission yeast arginases Aru1 and Car1 process arginine to ornithine.

Interestingly, upon direct TORC1 inhibition through rapalink-1 the main arginase Aru1 is downregulated

while Car1 is upregulated 1.4-fold. Data from the orfeome and global localisation study in fission yeast [45]

indicate that Aru1 and Car1 are localised in the cytoplasm and the nucleus. Nevertheless, Agm1 and Agm3

(no relevant data on Agm2 are available) are found in the endoplasmic reticulum (note that reticulophagy is a

Gene Ontology term for genes interacting with *agm1*) and the vacuole. Therefore, agmatinase genes are

localised in the right compartment to control arginine pools crucial for TORC1 activity and fission yeast

cells' survival during nutritional and other stresses as well as in the context of lifespan. Taken together, these

data and circuits can explain the requirements of the agmatinases in chronological ageing.

Our gene expression analyses have revealed many TORC1-regulated genes with no annotations or

characterisation related to ageing and lifespan. Recent studies have focused on broad profiling of fission

yeast mutants [33] using non-competitive settings. In addition, we and others have performed genome-wide

assays [13] towards characterising mutants related to lifespan regulation [31,42]. Despite these efforts and due to

the competitive nature of pooled barcode-based assays, only a fraction of the available viable mutants has

been characterised. In addition, many medium-or high-throughput lifespan approaches have already been

described and conducted in fission yeast [27,31,46]. Nevertheless, this gap of knowledge remains. Uncovering

novel TORC1-regulated genes will focus current and future efforts in discovering new biology within the

biogerontology field and beyond.

Parallel to gene expression analyses we have performed cell-based genome-wide screens to reveal

mutants that are sensitive and resistant to rapalink-1 as we have previously reported for torin1 [9]. We have

concluded that Pik3 is required, at least partially, for healthspan and lifespan extension resulting from

rapalink-1-mediated mTOR inhibition during cell proliferation. Rapalink-1 prolongs lifespan considerably

without Pik3, as previously observed for gaf1 and torin1 [9]. This is anticipated due to the plethora of proteins

and compensation mechanisms that might operate downstream of mTOR (see next RNA-seq analysis below)

including factors related for endolysosomal processes. Our data showcase that beneficial effects of mTOR

inhibition in lifespan are related to this pathway and involve ESCRT, HOPS, CORVET, autophagy and the

PIK3 complex. Taken together, these results indicate that these complexes, as in the case of rapalink-1, may

be required for mTOR-dependent lifespan control through other inhibitors including rapamycin and torin1.

Recent work has linked the inhibition of S6K with lifespan increase through the endolysosomal system [47]

showcasing the universal roles of these functional connections in the ageing process.

Interest in agmatine as a nutraceutical is significant and it is reported as a promising therapeutic

agent for treating a broad spectrum of central nervous system-associated diseases [48]. Large portion of

agmatine in mammals is supplemented from diets and gut microbiota and long-term (5 years)

supplementation studies [49] have demonstrated its safety in the examined doses. Beneficial effects on the

host's lifespan through microbiome-derived agmatine induced by metformin treatment have been previously

reported [37,50]. Indeed, agmatine can augment metformin's therapeutic effects. However, it does not always

promote beneficial effects as it can contribute to pathology [51]. Here we show that a consequence of the direct

TORC1 inhibition, is the transcriptional activation of all the enzymes that catabolise agmatine to putrescine

and urea with this pathway being vital for viability in non-dividing states. We have performed genome-wide

phenomic screens revealing mutant backgrounds that benefit (or not) from agmatine and putrescine

treatments. Our results show that supplementations of agmatine and putrescine are beneficial for cellular

fitness only when arginine metabolism pathways are intact. This may explain diverse outcomes of agmatine

supplementation and possible contributions to pathological conditions that are observed in other studies.

There is a growing interest in polyamines and the how they regulate longevity. The triamine

spermidine, a natural polyamine, can be generated from both the ornithine and the agmatine axes. In

mammalian cells its biosynthesis starts from arginine with the enzyme arginase (ARG1) catalysing the

reaction from arginine to ornithine [53]. Spermidine is well-known to stimulate cytoprotective mechanisms and

autophagy [52]. Spermidine has been implicated with lifespan in yeast and we have previously uncovered

longevity alleles and revealed connections of Rim15/Ppk31 with spermidine metabolism [19]. Recently,

spermidine is shown to be pivotal in mediating fasting-induced autophagy in yeast, nematodes and human

cells [20]. Nevertheless, in our present study we show that agmatinases do not function to simply contribute to

putrescine and ultimately spermidine production. While spermidine induces autophagy and mediates

hypusination of the translation regulator eIF5A [20], agmatinases' induction due to lowering of mTORC1

activity levels contribute towards maintaining the low TORC1 activity. Our analyses, reveal a metabolic

feedback loop mediated through agmatinase enzymes ensuring that TORC1 activity levels are maintained as

required for physiological cell needs and according to the signals received from the environment (nutritional

or pharmacological). The revealed metabolic control is required for survival in non-dividing states and in

chronological lifespan and can have implications and be of importance for human cells. Understanding how

TORC1 activity is tuned may be beneficial in both normal ageing and also pathological states as well as in

cancer where mTOR plays important roles [2,54].

Methods

Strains and Media

972 h− was used as the *wild-type*. The strain containing C-terminally GFP-tagged Gaf1 (*gaf1-GFP kanMX6*)

has been generated according to [55] with *gaf1* being under the control of the endogenous promoter and as in

[7,9,56]. Strains Maf-1-pK and GFP-atg8 have been previously described [23,25]. We have also used *h+*

agm1::kanMX6; h+ agm2::kanMX6; h+ agm3::kanMX6; h+ fkh1::kanMX6; h+ pik3::kanMX6;

agm1::natMX6 agm3::kanMX6; agm1::natMX6 tco89::kanMX6 [28]. Microscopy was performed using an

EVOS M5000 and Leica DMRA2 epi-fluorescent microscope fitted with a monochrome Orca-ER camera

from Hamamatsu Photonics. YES and EMM2 media (Formedium) are used as indicated in figures and main

text. Drug treatments have been conducted in fast growing cultures at OD_{600} 0.5 (see main text for details).

Liquid cultures were grown at 32°C with shaking at 130 rpm.

Western blotting

Cells were treated with 100 nM rapamycin or 100 nM rapalink-1 for two or five hours as indicated in figures

and main text. Cells were disrupted using RIPA buffer and glass beads in a FastPrep 5G MP lysis system.

Antibodies directed against phospho-eIF2a (#9721, 1:1,000), eIF2a (#9722, 1:1,000), phospho-Ssp2

(#50081, 1:1,000), Phospho-(Ser/Thr) Akt Substrate Antibody (#9611, 1:2,000) were purchased from Cell

Signaling Technologies. The antibodies directed against V5 (sc-81594, 1:1,000) and HA (sc-7392, 1:1,000)

were purchased from Santa Cruz Biotechnology. Secondary antibodies Goat Anti-Mouse IgG H&L (HRP)

(ab205719) and Goat Anti-Rabbit IgG H&L (HRP) (ab205718) were purchased from Abcam while the ECL

Western Blotting Detection system was from Pierce™. α-tubulin (T5168, 1:1,000) and anti-GFP

(11814460001, 1:1,000) were purchased from Sigma-Aldrich. α-tubulin and Ponceau S staining have been

used as loading controls for western blots as and where indicated. Western blotting quantifications have been

performed using ImageJ/Fiji [57] as previously described [27].

Cell size and septation index determination

To measure septation index, cells were resuspended in calcofluor solution (18909, Sigma-Aldrich) and

incubated at room temperature for five minutes. Cells were visualized using an EVOS M5000 microscope.

Percentages of septation from 200 cells for each time point have been recorded. For cell length at division,

measurements of 50-100 septated cells were conducted using Fiji/ImageJ software [57] as previously described

[7].

Chronological Lifespan (CLS) Assays

CLS was determined as previously described in [8]. Fission yeast cells are grown to stationary phase.

However, in this study, cells are left for an additional 48 hours as seen in previous related studies [15] and then

the assay starts with this time point considered as the 100% survival for the time course. 2 or 3 independent

biological repeats with each repeat having 3 technical replicates have been used. Log rank tests are

performed for all CLS assays with CFUs and percentages used in the statistical testing. Statistical

significance has been also validated with AUC measurements [57] as previously described in [9] and where

indicated/shown. Lifespan lengths vary largely in yeasts depending on the time point used following

stationary phase entry and the media used. Different batches of YES media can generate different lifespan

curve patterns depending on the Yeast Extract used in the particular batch. We have used a unique YES

batch (Formedium) throughout the study.

RNA Sequencing data

Untreated as well as rapamycin and rapalink-1 treated cells (treatment was at OD_{600}=0.5) for 5 hrs were

harvested and processed for RNA isolation as previously described (ref) using acidic phenol-chloroform

extraction method followed by RNAeasy (Qiagen) cleanup with on column DNase treatments before double

elution. Followign appropriate quality control steps, RNA library formed by polyA capture and Illumina 150

bp paired-end sequencing followed by standard R-based bioinformatic analysis was performed. Differential

gene expression analysis of two conditions/groups (two biological replicates per condition) was performed

using DESeq2R package (1.20.0). Genes with an adjusted P-value <=0.05 found by DESeq2 and at various

fold cutoffs (see main text) were assigned as differentially expressed. Sequences are deposited at GEO

(accession number: GSE2722A).

High throughput phenomics screens

The deletion library [28] was arrayed on solid YES media at 1536-spot density, with each strain represented by

4 spots with various concentrations of drugs as indicated for each assay. Plates were incubated at 32°C and

high-resolution images of the plates were acquired. Colony size quantitations were then performed using the

R package Gitter [58]. Median colony sizes were calculated for each plate and replicate. Strain colony size data

per condition were normalized to the corresponding growth on YES.

Synthetic genetic arrays

SGAs were performed using the Bioneer v5.0 haploid library [28] essentially as described previously [7], with

each interaction examined in quadruplicate. Query strains were the following: *h− agm1::natMX6,* and

control query *h− ade6::natMX6.* Colony size was used as a proxy for double mutant fitness. Colony size

measurements were obtained using the gitter package [58]. Library mutants which fell within 30 kb distance of

query mutations were excluded from the respective datasets to avoid spurious interactions as well as mutants

for which the *ade6* double mutant colony size was less than 50 pixels as they were deemed to be present in

too little amount for accurate interaction values to be calculated. Colony sizes were normalised to the plate

median accounting for the effect of the query mutation and plate-specific effects and normalised for column-

or row-specific effects. Medians of colony sizes of the SGAs were normalised with respect to the *ade6* SGA

(which represents fitness of library single mutants as this query does not affect the library) to calculate the

value of genetic interactions between query and library mutants. Interactions with high within-replicate

variability were excluded. Finally, the logarithms in base 10 of the interaction were used for interaction

scores.

Enrichment analyses of interactions.

Enrichment analyses were performed using Metascape [59]. p-values were corrected for multiple tests

according to FDR. Enrichment analysis was conducted by comparing lists of interacting genes to all genes in

the dataset (Bioneer v5.0 collection strains [28]).

Funding

This work was supported by funding to C.R. from the the Biotechnology and Biological Sciences Research

Council [Research grant numbers: BB/V006916/1, BB/V006916/2]. C.R. also acknowledges support and

funding of the group from the Medical Research Council [Grant number: MR/W001462/1]).

Conflict of Interest

None declared.

Acknowledgments

We thank members of the Rallis Lab Rowshan Islan and John-Patrick Alao for critical reading and valuable

comments on the manuscript. We are grateful to the Bahler lab for the use of the Biolector micro-

fermentation system used in Fig.2A.

Authors' contributions

Conceptualization, C.R.; Methodology, C.R. Investigation, J.K., C.R., Formal Analysis, C.R., J.K.; Writing,

C.R., and J.K.; Funding Acquisition, C.R.; Supervision, C.R.

References

1. Bjedov, I. & Partridge, L. A longer and healthier life with TOR down-regulation: genetics and

drugs. *Biochem. Soc. Trans.* **39**, 460–465 (2011).

2. Laplante, M. & Sabatini, D. M. mTOR signaling in growth control and disease. *Cell* **149**, 274–

293 (2012).

3. Ikai, N., Nakazawa, N., Hayashi, T. & Yanagida, M. The reverse, but coordinated, roles of Tor2

(TORC1) and Tor1 (TORC2) kinases for growth, cell cycle and separase-mediated mitosis in

Schizosaccharomyces pombe. *Open Biol* **1**, 110007 (2011).

4. Gonzalez, S. & Rallis, C. The TOR Signaling Pathway in Spatial and Temporal Control of Cell

Size and Growth. *Frontiers in Cell and Developmental Biology* **5**, (2017).

5. Bjedov, I. & Rallis, C. The Target of Rapamycin Signalling Pathway in Ageing and Lifespan

Regulation. *Genes (Basel)* **11**, (2020).

6. Harrison, D. E. *et al.* Rapamycin fed late in life extends lifespan in genetically heterogeneous

mice. *Nature* **460**, 392–395 (2009).

7. Rallis, C., Townsend, S. & Bähler, J. Genetic interactions and functional analyses of the fission

yeast gsk3 and amk2 single and double mutants defective in TORC1-dependent processes.

Sci Rep **7**, 44257 (2017).

8. Rallis, C., Codlin, S. & Bähler, J. TORC1 signaling inhibition by rapamycin and caffeine affect

lifespan, global gene expression, and cell proliferation of fission yeast. *Aging Cell* **12**, 563–573

(2013).

9. Rodríguez-López, M. *et al.* The GATA Transcription Factor Gaf1 Represses tRNAs, Inhibits

Growth, and Extends Chronological Lifespan Downstream of Fission Yeast TORC1. *Cell Rep*

30, 3240-3249.e4 (2020).

10. Fan, Q. *et al.* A Kinase Inhibitor Targeted to mTORC1 Drives Regression in Glioblastoma.

Cancer Cell **31**, 424–435 (2017).

11. Kuroshima, K. *et al.* Potential new therapy of Rapalink-1, a new generation mammalian target

of rapamycin inhibitor, against sunitinib-resistant renal cell carcinoma. *Cancer Sci* **111**, 1607–

1618 (2020).

12. Zhou, H. *et al.* mTOR Inhibitor Rapalink-1 Prevents Ethanol-Induced Senescence in

Endothelial Cells. *Cells* **12**, 2609 (2023).

13. Legon, L. & Rallis, C. Genome-wide screens in yeast models towards understanding

chronological lifespan regulation. *Brief Funct Genomics* **21**, 4–12 (2022).

14. Roux, A. E., Arseneault, G., Chartrand, P., Ferbeyre, G. & Rokeach, L. A. A screen for genes

involved in respiration control and longevity in Schizosaccharomyces pombe. *Ann N Y Acad*

Sci **1197**, 19–27 (2010).

15. Roux, A. E. *et al.* Pro-aging effects of glucose signaling through a G protein-coupled glucose

receptor in fission yeast. *PLoS Genet* **5**, e1000408 (2009).

16. Petersen, J. & Nurse, P. TOR signalling regulates mitotic commitment through the stress MAP

kinase pathway and the Polo and Cdc2 kinases. *Nature Cell Biology* **9**, 1263–1272 (2007).

17. Lock, A., Harris, M. A., Rutherford, K., Hayles, J. & Wood, V. Community curation in PomBase:

enabling fission yeast experts to provide detailed, standardized, sharable annotation from

research publications. *Database (Oxford)* **2020**, baaa028 (2020).

18. Rutherford, K. M., Lera-Ramírez, M. & Wood, V. PomBase: a Global Core Biodata Resource-

growth, collaboration, and sustainability. *Genetics* **227**, iyae007 (2024).

19. Ellis, D. A. *et al.* Uncovering Natural Longevity Alleles from Intercrossed Pools of Aging Fission

Yeast Cells. *Genetics* **210**, 733–744 (2018).

20. Hofer, S. J. *et al.* Spermidine is essential for fasting-mediated autophagy and longevity. *Nat*

Cell Biol (2024) doi:10.1038/s41556-024-01468-x.

21. Thoreen, C. C. *et al.* An ATP-competitive mammalian target of rapamycin inhibitor reveals

rapamycin-resistant functions of mTORC1. *J Biol Chem* **284**, 8023–8032 (2009).

22. Thoreen, C. C. & Sabatini, D. M. Rapamycin inhibits mTORC1, but not completely. *Autophagy*

5, 725–726 (2009).

23. Davie, E., Forte, G. M. A. & Petersen, J. Nitrogen Regulates AMPK to Control TORC1

Signaling. *Current Biology* **25**, 445–454 (2015).

24. Ling, N. X. Y. *et al.* mTORC1 directly inhibits AMPK to promote cell proliferation under nutrient

stress. *Nat Metab* **2**, 41–49 (2020).

25. Mukaiyama, H., Nakase, M., Nakamura, T., Kakinuma, Y. & Takegawa, K. Autophagy in the

fission yeast *Schizosaccharomyces pombe*. *FEBS Letters* **584**, 1327–1334 (2010).

26. Noguchi, C. *et al.* Maf1 limits RNA polymerase III-directed transcription to preserve genomic

integrity and extend lifespan. *Cell Cycle* **20**, 247–255 (2021).

27. Rallis, C., López-Maury, L., Georgescu, T., Pancaldi, V. & Bähler, J. Systematic screen for

mutants resistant to TORC1 inhibition in fission yeast reveals genes involved in cellular ageing

and growth. *Biol Open* **3**, 161–171 (2014).

28. Kim, D.-U. *et al.* Analysis of a genome-wide set of gene deletions in the fission yeast

Schizosaccharomyces pombe. *Nat. Biotechnol.* **28**, 617–623 (2010).

29. Laxman, S. *et al.* Sulfur amino acids regulate translational capacity and metabolic homeostasis through modulation of tRNA thiolation. *Cell* **154**, 416–429 (2013).

30. Gilleron, J., Gerdes, J. M. & Zeigerer, A. Metabolic regulation through the endosomal system. *Traffic* **20**, 552–570 (2019).

31. Romila, C. A. *et al.* Barcode sequencing and a high-throughput assay for chronological lifespan uncover ageing-associated genes in fission yeast. *Microb Cell* **8**, 146–160 (2021).

32. Lie, S., Banks, P., Lawless, C., Lydall, D. & Petersen, J. The contribution of non-essential Schizosaccharomyces pombe genes to fitness in response to altered nutrient supply and target of rapamycin activity. *Open Biol* **8**, 180015 (2018).

33. Rodríguez-López, M. *et al.* Broad functional profiling of fission yeast proteins using phenomics and machine learning. *Elife* **12**, RP88229 (2023).

34. Schmeisser, K. & Parker, J. A. Pleiotropic Effects of mTOR and Autophagy During Development and Aging. *Front Cell Dev Biol* **7**, 192 (2019).

35. Valbuena, N., Guan, K.-L. & Moreno, S. The Vam6 and Gtr1-Gtr2 pathway activates TORC1 in response to amino acids in fission yeast. *J. Cell. Sci.* **125**, 1920–1928 (2012).

36. Hayles, J. *et al.* A genome-wide resource of cell cycle and cell shape genes of fission yeast. *Open Biol.* **3**, 130053 (2013).

37. Pryor, R. *et al.* Host-Microbe-Drug-Nutrient Screen Identifies Bacterial Effectors of Metformin Therapy. *Cell* **178**, 1299-1312.e29 (2019).

38. Milosevic, K. *et al.* Agmatine Mitigates Inflammation-Related Oxidative Stress in BV-2 Cells by Inducing a Pre-Adaptive Response. *Int J Mol Sci* **23**, 3561 (2022).

39. Valbuena, N., Rozalén, A. E. & Moreno, S. Fission yeast TORC1 prevents eIF2α phosphorylation in response to nitrogen and amino acids via Gcn2 kinase. *J Cell Sci* **125**, 5955–5959 (2012).

40. Duncan, C. D. S., Rodríguez-López, M., Ruis, P., Bähler, J. & Mata, J. General amino acid control in fission yeast is regulated by a nonconserved transcription factor, with functions analogous to Gcn4/Atf4. *Proc Natl Acad Sci U S A* **115**, E1829–E1838 (2018).

41. Mülleder, M. *et al.* Functional Metabolomics Describes the Yeast Biosynthetic Regulome. *Cell* **167**, 553-565.e12 (2016).

42. Sideri, T. *et al.* Parallel Profiling of Fission Yeast Deletion Mutants for Proliferation and for

Lifespan During Long-Term Quiescence. *G3* **5**, 145–155 (2015).

43. Dudin, O. *et al.* A systematic screen for morphological abnormalities during fission yeast

sexual reproduction identifies a mechanism of actin aster formation for cell fusion. *PLoS Genet*

13, e1006721 (2017).

44. López-Maury, L., Marguerat, S. & Bähler, J. Tuning gene expression to changing

environments: from rapid responses to evolutionary adaptation. *Nat Rev Genet* **9**, 583–593

(2008).

45. Matsuyama, A. *et al.* ORFeome cloning and global analysis of protein localization in the fission

yeast Schizosaccharomyces pombe. *Nat Biotechnol* **24**, 841–847 (2006).

46. Kamrad, S. *et al.* Pyphe, a python toolbox for assessing microbial growth and cell viability in

high-throughput colony screens. *Elife* **9**, (2020).

47. Zhang, P., Catterson, J. H., Grönke, S. & Partridge, L. Inhibition of S6K lowers age-related

inflammation and increases lifespan through the endolysosomal system. *Nat Aging* **4**, 491–509

(2024).

48. Akasaka, N. & Fujiwara, S. The therapeutic and nutraceutical potential of agmatine, and its

enhanced production using Aspergillus oryzae. *Amino Acids* **52**, 181–197 (2020).

49. Gilad, G. M. & Gilad, V. H. Long-term (5 years), high daily dosage of dietary agmatine--

evidence of safety: a case report. *J Med Food* **17**, 1256–1259 (2014).

50. Tassoulas, L. J. & Wackett, L. P. Insights into the action of the pharmaceutical metformin:

Targeted inhibition of the gut microbial enzyme agmatinase. *iScience* **27**, 108900 (2024).

51. Yun, C. *et al.* The microbial metabolite agmatine acts as an FXR agonist to promote polycystic

ovary syndrome in female mice. *Nat Metab* **6**, 947–962 (2024).

52. Madeo, F., Bauer, M. A., Carmona-Gutierrez, D. & Kroemer, G. Spermidine: a physiological

autophagy inducer acting as an anti-aging vitamin in humans? *Autophagy* **15**, 165–168 (2019).

53. Kaiser, A. The Role of Spermidine and Its Key Metabolites in Important, Pathogenic Human

Viruses and in Parasitic Infections Caused by Plasmodium falciparum and Trypanosoma

brucei. *Biomolecules* **13**, 803 (2023).

54. Marques-Ramos, A. & Cervantes, R. Expression of mTOR in normal and pathological

conditions. *Mol Cancer* **22**, 112 (2023).

55. Bähler, J. *et al.* Heterologous modules for efficient and versatile PCR-based gene targeting in

Schizosaccharomyces pombe. *Yeast* **14**, 943–951 (1998).

56. Rallis, C. *et al.* Amino Acids Whose Intracellular Levels Change Most During Aging Alter

Chronological Life Span of Fission Yeast. *J Gerontol A Biol Sci Med Sci* **76**, 205–210 (2021).

57. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods*

9, 676–682 (2012).

58. Wagih, O. & Parts, L. gitter: A Robust and Accurate Method for Quantification of Colony Sizes

From Plate Images. *G3 Genes|Genomes|Genetics* **4**, 547–552 (2014).

59. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-

level datasets. *Nat Commun* **10**, 1523 (2019).

Figure 1 (Kumar and Rallis)

Rapalink-1 affects TORC1-related processes in fission yeast and prolongs lifespan.

(A) Time course of septation index for untreated (control) and rapamycin and rapalink-1 treated cells (100

nM), as indicated, in YES media. Dotted lines represent average values while coloured ribbons standard

deviation from three independent counts.

(B) Same as in A in EMM2 minimal media.

(C) Time course of cell size upon division measurements following rapamycin and rapalink-1 treatments in

YES media. Asterisks indicate statistically significant differences between the compared groups (p<0.01,

wilcoxon testing).

(D) Same as described in panel C with the assay conducted in EMM2 media.

(E) Western blots for markers dependent on TORC1 activity and representative loading controls. Assays

have been conducted following 2 hrs and 5 hrs of treatments with 100 nM rapamycin (Rm) and rapalink-1

(Rl) together with untreated control cells (C).

(F) DAPI staining and Gaf1-GFP localisation panels in control untreated, rapamycin and rapalink-1 treated

cells as indicated.

(G) CLS assays for untreated (control) and rapamycin treated wild-type fission yeast cells (log-rank

$p=1.25*10^{-12}$, see materials and methods).

(H) CLS assays for untreated (control) and rapalink-1 treated wild-type fission yeast cells (log-rank

$p=2.2*10^{-12}$, see materials and methods).

(I) CLS assays for untreated (control) and rapalink-1 treated *fkh1Δ* mutant cells (log-rank p=0.047, see

materials and methods).

*** in lifespans indicate p values <0.001.

Figure 2 (Kumar and Rallis)

A genome-wide screen for rapalink-1 points to the endosome-vacuole-lysosome network as important in mTOR-dependent lifespan regulation.

(A) Growth curves for control untreated, rapamycin and rapalink-1 treated wild type fission yeast cells, as indicated. The assay records light scattering dependent on culture cell mass (see materials and methods).

(B) Fitness ratio overviews of genome-wide screens (data acquired for 3,271 deletion mutants) for rapalink-1

at the indicated concentrations. Vertical dotted lines indicate cutoffs for sensitive and resistant mutants

(Supplementary table 1).

(C) Fitness ratio correlation between the two conducted screens. Each point represents the average fitness

obtained for a single deletion mutant of the library.

(D) Gene ontology enrichment for mutants sensitive to rapalink-1 treatment.

(E) Gene ontology enrichment for mutants resistant to rapalink-1 treatment.

(F) Network of enriched terms colored by cluster ID, where nodes that share the same cluster ID are

typically close to each other.

(G) CLS assay for untreated and rapalink-1 treated *pik3Δ* cells as indicated (log rank $p=10^{-5}$).

(H) AUC for lifespan assays of *wt* and *pik3Δ* mutant cells without or with rapalink-1 treatment (*wt-RL* and

pik3Δ-RL respectively. Vertical bars: rapalink-1-mediated increase in average AUC values for *wt* (black) and

pik3Δ (red) for curves until day 6 (*wt* curve shown in Fig.1), with the p value reflecting the significantly

larger lifespan increase in *wt* than in *pik3Δ* cells.

*** in lifespans indicate p values <0.001.

Figure 3 (Kumar and Rallis)

Gene expression analysis following rapamycin and rapalink-1 treatment reveals novel genes involved in chronological ageing.

(A) PCA analysis of control untreated, rapamycin- and rapalink-1-treated cells as indicated.

(B) Heatmap of representative GO enrichments for gene lists of upregulated and downregulated

genes in the drug treatments used (R-Up: rapalink-1 upregulated; R-Down: rapalink-1

downregulated; M-up: rapamycin upregulated; M-down: rapamycin downregulated).

(C) Venn diagram for upregulated genes following rapamycin and rapalink-1 treatments.

(D) Venn diagram for downregulated genes following rapamycin and rapalink-1 treatments.

(E) Venn diagram showing overlaps of rapalink-1 upregulated genes (the portion that

corresponding mutants are viable) with genes that their deletions lead to long-lived, short-lived or

normal-lived mutants.

(F) Schematic of arginine metabolism to ornithine, agmatine and putrescine with genes coding for

responsible enzymes. In green: genes that are upregulated, red: downregulated, black: not

affected, following rapalink-1 treatment.

(G-J) CLS for normal (*wt*) and mutant *agm1Δ* (G, log rank p=9.7x10^{-5}), *agm2Δ* (H, log rank p=1.5x10^{-5}),

agm3Δ (I, log rank, p=3.9x10^{-5}), and *agm1Δ agm3Δ* (J, log rank p=3.3x10^{-6}) cells as indicated.

*** in lifespans indicate p values <0.001.

Supplemental Figure 1 (Kumar and Rallis)

RNAseq data analyses of the paper related to Figure 3.

(A-C) Volcano plots with upregulated (red), downregulated (blue) and unchanged (black) genes.

(A) Untreated versus rapamycin-treated cells.

(B) Untreated versus rapalink-1-treated cells.

(C) rapamycin versus rapalink-1-treated cells.

(D) Bar graph representation of representative GO ontology enrichments for genes upregulated in rapalink-1

treatment only (related to Figure 3E)

(E) Bar graph representation of representative GO ontology enrichments for genes downregulated in

rapalink-1 treatment only (related to Figure 3F).

Supplemental Figure 2 (Kumar and Rallis)

Gene expression analyses and enrichments related to rapalink-1.

(A) Bar graph schematic for representative GO ontology enrichments of rapalink-1-upregulated

genes not related to lifespan phenotype (related to Figure 2H, genes in red fonts).

(B) Venn diagram showing overlaps of rapalink-1 downregulated genes (the portion that

corresponding mutants are viable) with genes that their deletions lead to long-lived, short-lived or

normal-lived mutants.

(C) Bar graph schematic for representative GO ontology enrichments of the 150 rapalink-1-downregulated

genes not related to lifespan phenotype shown in (B).

Figure 4 (Kumar and Rallis)

Genome-wide fitness screens of agmatine and putrescine and effects on chronological lifespan of

fission yeast.

(A) Density profiles of fitness ratios for 3,208 *S. pombe* deletion mutants following agmatine and putrescine

treatments.

(B) Correlation of fitness ratios of deletion mutants in agmatine and putrescine screens. Each point

represents the average fitness value for each mutant.

(C) Heatmap of representative Gene Ontology enrichments for mutant strains resistant and sensitive to drug

treatments (agm_sens: sensitive to agmatine; putr_sens: sensitive to putrescine; agm_resist: resistant to

agmatine; putr_resist: resistant to putrescine).

(D-M) CLS for normal (*wt*) and agmatinase mutant cells following agmatine and putrescine supplementation

during growth phase (see material and methods) as indicated.

*** in lifespans indicate p values <0.001, see log rank p values in the main text.

Figure 5 (Kumar and Rallis)

Genetic interactomes of *agm1* reveal a metabolic feedback mechanism that tunes TORC1 activity.

(A) Physical mapping of *agm1* interaction values with 3,108 mutants.

(B) Density plot of the genome-wide *agm1* genetic interactions. Vertical dotted lines show the cutoffs set for

the interactome analysis (see materials and methods).

(C) Bar graph representation of GO enrichments for negative *agm1* interactions.

(D) Bar graph representation of GO enrichments for positive *agm1* interactions.

(E) Schematic showing a previously reported tripartite TORC1 regulation in fission yeast. Green fonts

signify positive while red fonts negative interactions.

(F) Cell size upon division measurements in normal (wt) and *agm* mutant cells (*agm1Δ, agm2Δ, agm3Δ*) in

YES media. Asterisks indicate statistically significant differences between the compared groups ($p<0.01$,

wilcoxon testing).

(G) Western blot for p-Akt targets in normal (wt) and *agm* mutant cells (*agm1Δ, agm2Δ, agm3Δ*) with

accompanied α-tubulin western blot as loading control. Side numbers (1-4) indicate various detected targets.

No 4 represents and used in literature [24] for pRps6.

(H) Quantification of -Akt target signal (whole lane) from two repeats of the blot shown in (G).

(I) CLS patterns of *agm1Δ and agm1Δ tco89Δ* mutants is indicated. N.s.: log rank $p>0.05$

(J) A schematic model for the revealed role of agmatinase enzymes in tuning TORC1 activity in fission

yeast.

SUPPLEMENTAL TABLES

Supplementary Table 1

List of rapalink-1 sensitive (<0.75, representing 25% smaller colony compared to untreated control) and

resistant (>1.25, 25% larger colony compared to untreated control) mutant strains.

Supplementary Table 2

List of genes upregulated in fast growing fission yeast cells (0.5 OD_{600}) following 5 hrs of 100nM

rapamycin treatment in YES media.

Supplementary Table 3

List of genes downregulated in fast growing fission yeast cells (0.5 OD_{600}) following 5 hrs of 100nM

rapamycin treatment in YES media.

Supplementary Table 4

List of genes upregulated in fast growing fission yeast cells (0.5 OD_{600}) following 5 hrs of 100nM

rapalink-1 treatment in YES media.

Supplementary Table 5

List of genes downregulated in fast growing fission yeast cells (0.5 OD_{600}) following 5 hrs of 100nM

rapalink-1 treatment in YES media.

Supplementary Table 6

Fitness ratio values of mutant strains following genome-wide phenomics screen with Agmatine and

Putrescine (see main text and materials and methods).

Supplementary Table 7

Interaction values for all mutant strains against *agm1* following normalisations (see Materials and Methods)

and filtering of absent colonies, control and experimental SGA linkage, high variability of colonies and low

fitness strains.