Bacillus toyonensis **biovar Thuringiensis: an overlooked entomopathogen?**

Diego Herman Sauka[1,2], Cecilia Peralta[3], Eleodoro E. Del Valle[4], Leopoldo Palma[3]*

[1]Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Microbiología y Zoología

Agrícola (IMYZA), Hurlingham, Buenos Aires, B1686IGC, Argentina.

[2]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Ciudad Autónoma de

Buenos Aires, C1033AAJ, Argentina.

[3]Instituto de Biotecnología y Biomedicina (BIOTECMED), Departamento de Genética, Universitat de

València, 46100 Burjassot, Spain.

[4]ICiagro Litoral, CONICET, Facultad de Ciencias Agrarias, Universidad Nacional del Litoral, Kreder

2805, Esperanza S3080, Argentina.

*Correspondence: leopoldo.palma@uv.es

Abstract

Horizontal gene transfer (HGT) significantly influences prokaryotic genome evolution. *Bacillus*

cereus and *Bacillus thuringiensis* are nearly identical at the chromosomal level, except for *B.*

thuringiensis producing parasporal crystals. The genes for these crystal proteins (e.g., *cry1A*), along

with other encoded insecticidal proteins (e.g., *vip3A*), are located on megaplasmids and can be

horizontally transferred. Recently, Sauka et al. (2022) reported a *Bacillus toyonensis* strain that

produces parasporal crystals with dual insecticidal activity. This strain was classified as *Bacillus*

toyonensis biovar Thuringiensis (NCBI: txid2923195) following Carroll et al.'s (2020) nomenclature.

Misclassified *B. toyonensis* strains, previously identified as *B. thuringiensis* (e.g., strain MC28), encode

cry and *cyt* genes toxic to lepidopteran and dipteran insects. Advances in genome sequencing and

bioinformatics tools now reduce misidentifications, enabling accurate reclassification in databases like

GenBank. These findings highlight the need for genome-based taxonomic reassessment within the

Bacillus cereus group and clarify the chromosomal placement of crystal-forming *B. toyonensis* strains.

Keywords: *Bacillus thuringiensis*, *Bacillus toyonensis* biovar Thuringiensis, genome-based taxonomy,

average nucleotide identity (ANI), insecticidal proteins.

Bacillus thuringiensis is a gram-positive, spore-forming bacterium known for producing various

proteins with specific biocidal activity against invertebrates. This characteristic has made *B. thurin-*

giensis one of the most studied and widely used biological control agents for sustainable pest man-

agement in agriculture (targeting insects and nematodes) and mosquito control (1). The repertoire of

pesticidal proteins includes parasporal crystal proteins such as Cry, Tpp, and Cyt (δ-endotoxins), pro-

duced during the stationary growth phase, and secreted proteins historically referred to as vegetative

insecticidal proteins (Vip), including the binary Vpb1/Vpa2 (formerly Vip1/Vip2), Vip3, and Vpb4

(formerly Vip4) (2, 3). Vip3 proteins can be produced from mid-log growth phase through sporulation,

with *vip3A* transcription activated at the onset of stationary phase by the regulator VipR (4).

Other gram-positive, spore-forming bacteria also produce biocidal proteins. For example, *Lysini-*

bacillus sphaericus produces Tpp proteins (formerly BinA/BinB proteins) (5), and *Brevibacillus lat-*

erosporus produces Cry proteins, both of which exhibit toxicity against mosquitoes (6).

In 2022, Sauka et al. isolated and preliminarily classified a *B. thuringiensis* strain with parasporal

crystals and insecticidal activity against *Cydia pomonella* (Lepidoptera) and *Anthonomus grandis*

(Coleoptera) (7). However, with an average nucleotide identity (ANI) of 98.6% against *Bacillus*

toyonensis, this novel strain was officially named *B. toyonensis* biovar Thuringiensis strain

Bto_UNVM-94 (7), following the nomenclature system proposed by Carroll et al. (2020) (8). Another

similar case is reported for *Bacillus wiedmannii* biovar thuringiensis, a strain harboring genes encoding

for parasporal crystal proteins with mosquitocidal activity (9). *B. toyonensis* was first described in 2013

by Jimenez et al. as a member of the *Bacillus cereus* group, which includes *B. cereus*, *B. thuringiensis*,

and other related pathogens. This species has been used as a probiotic in the swine, poultry, cattle,

rabbit, and aquaculture industries for over 30 years (10).

Samples were taken with a tubular soil sampler from different areas of Argentina, namely: a native

forest area from Jujuy province (strain Bto_UNVM-42), Tancacha town from province of Córdoba

(strain Bto_UNVM-100) and as previously described for strain Bto_UNVM-94 (7). After collection,

samples were placed in zip-lock bags and stored at 4 °C until bacterial isolation could be performed. For

this purpose, 3 grams of the soil were suspended in 10 ml of sterile distilled water and homogenized by

vortexing for 1 minute. The suspension was then subjected to a heat shock at 80 °C for 30 minutes to

select for spore-forming bacteria. Serial decimal dilutions (from 10^{-3} to 10^{-5}) were prepared, and 50 µl

aliquots from each were plated on nutrient agar medium composed of 0.5% peptone, 0.3% beef extract,

0.5% NaCl, and 1.5% agar. The plates were spread using a Drigalsky spatula and incubated at 28 °C for

48 to 72 hours. Colonies resembling *Bacillus thuringiensis*, characterized by a matte-white appearance

and irregular margins, were selected and further streaked for purification on fresh agar plates. These

were incubated under the same conditions until sporulation was observed. Once sporulated, cultures

were heat-fixed on glass microscope slides and stained with a Coomassie Blue solution (0.133% in 50%

acetic acid) for visualization in a light microscope (11). The presence of parasporal crystals, charac-

teristic of a *B. thuringiensis*-like phenotype, was later confirmed using scanning electron microscopy.

Genomic DNA, encompassing both chromosomal and plasmid components, was extracted

using the Wizard Genomic DNA Purification Kit (Promega). The procedure was carried out in

accordance with the manufacturer's protocol, specifically optimized for DNA isolation from

gram-positive bacterial strains. The extracted DNA was used subsequently for the preparation

of a pooled Illumina library at the Genomics Unit from the National Institute of Agricultural

Technology (INTA, Argentina) by using high-throughput Illumina sequencing technology for

strains Bto_UNVM-42 along with Bto_UNVM-100, and as previously described for strain

Bto_UNVM-94 (7). The raw Illumina reads obtained were first trimmed and assembled into

contigs by using Velvet plug in included in Geneious R11 (www.geneious.com). The resultant

contigs were then analyzed with BLAST (12) using a customized non-redundant insecticidal

protein database. Genome annotation was performed with the NCBI Prokaryotic Genome

Annotation Pipeline (2021 release) although it was also annotated with RAST (13). Species

delimitation was performed with the Type (Strain) Genome Server (14), FastANI (15) and with

PubMLST (16).

By using this combination of genome-genome hybridization analysis with TYGS server (14) and %

ANI calculations with FastANI v0.1.3 (15), we have detected the potential misclassification of two

Bacillus species stored in the GenBank database (17). Strains *B. thuringiensis* MC28 (RefSeq

GCF_000300475.1) (18) and *B. cereus* Rock 1-3 (RefSeq GCF 000161155.1) (19) are clustered in the

same branch along with type strain *B. toyonensis* P18 (RefSeq GCF_016605985.1) (Figure 1).

In addition, estimated % ANI and PubMLST support values of these strains, strongly suggests the

need of performing a reclassification of these *Bacillus* species by following the novel nomenclature

system proposed by Carroll et al., (2020). We also added in the calculations the genomic sequences

coming from novel *Bacillus* strains Bto_UNVM-42 and Bto_UNVM-100 (Table 1).

Strain MC28 can form *B. thuringiensis*-like parasporal crystals and harbor several genes encoding

for δ-endotoxins (Cry and Cyt), reported to be highly toxic to lepidopteran and dipteran insects, ac-

cording to the original strain description ([18]). However, this strain, which clusters with *B. toyonensis* in

genome-based phylogenetic analyses, has gone unnoticed due to its original classification as *B. thu-*

ringiensis MC28. We hypothesize here the existence of horizontal gene transfer (HGT) events. HGT is

one of the most significant sources of evolution that continuously causes major changes into prokaryotic

genomes and has been previously described within the *Bacillus* genus ([20]). One consistent example of

HGT within the *Bacillus cereus* group is the transfer of genes encoding insecticidal proteins, such as the

parasporal crystal proteins (Cry and Cyt), which confer toxicity against insect pests. These genes are

predominantly plasmid-borne and are frequently found in genomic regions associated with insertion

sequences and other mobile elements, which contribute to their horizontal dissemination ([3], [20-23]).

Several studies have documented the occurrence of HGT events involving insecticidal genes within the

B. cereus group. For instance, Liu et al. (2015) investigated the genomic diversity of *B. thuringiensis*

strains and identified multiple instances of HGT-mediated acquisition of insecticidal genes from other

*Bacillu*s species ([24]). Similarly, Geng et al. (2023) demonstrated HGT of the 144-kb plasmid

pTAND672-2, carrying mosquitocidal toxin genes, through conjugation experiments, between *B. thu-*

ringiensis serovar *israelensis* and *L. sphaericus* ([25]).

Overall, HGT serves as a key mechanism driving genetic diversity and adaptation within the *B.*

cereus group, particularly in the context of insecticidal properties.

In addition, the *B. cereus* strain Rock 1-3 has been also potentially misclassified and should be

renamed as belonging to *B. toyonensis* species.

In the case of *Bacillus* sp. strains Bto_UNVM-42 and Bto_UNVM-100, the same hypothesis in-

volving HGT events may apply. Strain Bto_UNVM-42 and Bto_UNVM-100 have been isolated from

provinces of Jujuy and Córdoba (Argentina), respectively, and they were able to produce parasporal

crystals with their genomic sequences harboring several genes with potential insecticidal activity (Table

2). However, both TYGS server and estimation of % ANI against type strain *B. toyonensis* P18 showed

these strains as belonging to *B. toyonensis* species. Following the nomenclature proposed by Carroll et

al., (2020), we propose to name these strains as belonging to *B. toyonensis* biovar Thuringiensis species.

Following the same approach, our results indicate that strain GM18 has been correctly classified as

B. toyonensis (26). Based on the reported production of parasporal inclusions and the presence of pes-

ticidal and parasporin-related genes in its genome, we propose that this strain be designated *B.*

toyonensis biovar Thuringiensis, consistent with the nomenclature framework proposed by Carroll et al.

(2020). Although GM18 was reported to lack insecticidal activity, solubilized parasporal inclusion

proteins were shown to display cytotoxic activity against human cancer cell lines in the original de-

scription (24) (Table 2).   Based on our phylogenetic analyses, the *B. toyonensis* biovar Thuringiensis

strains described here exhibit phenotypic characteristics like those of *B. thuringiensis*. These strains not

only produce parasporal crystals but also exhibit biocidal activity against insects and human cancer

cells.

In this work, we provide genomic and taxonomic evidence indicating that several *Bacillus* strains

previously classified as *B. thuringiensis* or *B. cereus* in fact belong to *B. toyonensis*, and that a subset of

strains of the latter species harbor genes encoding parasporal crystal proteins (Cry and Cyt) and other

pesticidal factors (Vpb, App, Mpp and Tpp proteins) (27). These findings support the recognition of

crystal-forming *B. toyonensis* strains as part of the taxonomic and phenotypic diversity within the *B.*

cereus group. Our hypothesis that HGT events have contributed to the acquisition of *cry*, *cyt* and other

insecticidal genes provides a plausible explanation for the emergence of a *B. thuringiensis*-like phe-

notype in strains that are genomically assigned to a different species. This observation is consistent with

the well-documented role of mobile genetic elements in shaping ecological traits within the *B. cereus*

group. Taken together, these observations highlight the importance of genome-based taxonomic

frameworks for resolving species boundaries within the *B. cereus* group. The identification of crys-

tal-forming strains that are chromosomally assigned to *B. toyonensis* illustrates how accessory gene

content may complicate phenotype-based classification and supports the use of genome-informed no-

menclature.

Future studies should aim to clarify the frequency, directionality and ecological drivers of these

HGT events in natural soil populations, as well as to determine how widespread *B. thuringiensis*-like

phenotypes are within *B. toyonensis* and related taxa. Such efforts will be essential to fully appreciate

the evolutionary dynamics, ecological significance and biotechnological potential of mobile pesticidal

gene reservoirs within the *B. cereus* group.

Acknowledgments

Leopoldo Palma would like to express gratitude to the Spanish Government-Universities Ministry,

the Next Generation EU and Recovery, Transformation, and Resilience plans for funding his awarded

María Zambrano contract (ref. ZA21-003) and the Ramón y Cajal programme (Ref.

RYC2023-043507-I).

Conflict of interests

The authors declare no conflicts of interest.

References

1.   Jouzani GS, Valijanian E, Sharafi R. 2017. *Bacillus thuringiensis*: a successful insecticide with
new environmental features and tidings. Appl Microbiol Biotechnol 101:2691–2711.

2.   Panneerselvam S, Mishra R, Berry C, Crickmore N, Bonning BC. 2022. BPPRC database: a
web-based tool to access and analyse bacterial pesticidal proteins. Database 2022.

3.   Schnepf E, Crickmore N, Van Rie J, Lereclus D, Baum J, Feitelson J, Zeigler DR, Dean DH.
1998. *Bacillus thuringiensis* and its pesticidal crystal proteins. Microbiol Mol Biol Rev 62:775–
806.

4.   Chen H, Verplaetse E, Slamti L, Lereclus D. 2022. Expression of the *Bacillus thuringiensis*
vip3A Insecticidal Toxin Gene Is Activated at the Onset of Stationary Phase by VipR, an
Autoregulated Transcription Factor. Microbiol Spectr 10:e0120522.

5.   Berry C. 2012. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. Journal of
Invertebrate Pathology 109:1–10.

6.   Ruiu L. 2013. *Brevibacillus laterosporus*, a Pathogen of Invertebrates and a Broad-Spectrum
Antimicrobial Species. Insects 4:476–92.

7.   Sauka DH, Peralta C, Pérez MP, Onco MI, Fiodor A, Caballero J, Caballero P, Berry C, Del
Valle EE, Palma L. 2022. *Bacillus toyonensis* biovar Thuringiensis: A novel entomopathogen
with insecticidal activity against lepidopteran and coleopteran pests. Biol Control 167:104838.

8.   Carroll LM, Wiedmann M, Kovac J. 2020. Proposal of a Taxonomic Nomenclature for the
Bacillus cereus Group Which Reconciles Genomic Definitions of Bacterial Species with
Clinical and Industrial Phenotypes. mBio 11.

9.   Lazarte JN, Lopez RP, Ghiringhelli PD, Berón CM. 2018. *Bacillus wiedmannii* biovar
thuringiensis: A Specialized Mosquitocidal Pathogen with Plasmids from Diverse Origins.
Genome Biol Evol 10:2823–2833.

10.   Jiménez G, Urdiain M, Cifuentes A, López-López A, Blanch AR, Tamames J, Kämpfer P,
Kolstø A-B, Ramón D, Martínez JF, Codoñer FM, Rosselló-Móra R. 2013. Description of
Bacillus toyonensis sp. *nov.*, a novel species of the Bacillus cereus group, and pairwise genome
comparisons of the species of the group by means of ANI calculations. Systematic and Applied
Microbiology 36:383–391.

11.   Ammons D, Rampersad J, Khan A. 2002. Usefulness of staining parasporal bodies when
screening for *Bacillus thuringiensis*. J Invertebr Pathol 79:203–4.

12.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search
tool. J Mol Biol 215:403–410.

13.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass
EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK,
Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V,
Wilke A, Zagnitko O. 2008. The RAST Server: rapid annotations using subsystems technology.
BMC Genomics 9:75.

14.   Meier-Kolthoff JP, Goker M. 2019. TYGS is an automated high-throughput platform for
state-of-the-art genome-based taxonomy. Nat Commun 10:2182.

15.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun
9:5114.

16.   Jolley KA, Bray JE, Maiden MCJ. 2018. Open-access bacterial population genomics: BIGSdb
software, the PubMLST.org website and their applications. Wellcome Open Res 3:124.

17.   Sayers EW, Cavanaugh M, Clark K, Pruitt KD, Schoch CL, Sherry ST, Karsch-Mizrachi I.
2021. GenBank. Nucleic Acids Res 49:D92–D96.

18.   Guan P, Ai P, Dai X, Zhang J, Xu L, Zhu J, Li Q, Deng Q, Li S, Wang S, Liu H, Wang L, Li P,
Zheng A. 2012. Complete Genome Sequence of *Bacillus thuringiensis* Serovar *sichuansis*
Strain MC28. J Bacteriol 194:6975–6975.

19.   Zwick ME, Joseph SJ, Didelot X, Chen PE, Bishop-Lilly KA, Stewart AC, Willner K, Nolan N,
Lentz S, Thomason MK, Sozhamannan S, Mateczun AJ, Du L, Read TD. 2012. Genomic
characterization of the *Bacillus cereus sensu lato* species: backdrop to the evolution of Bacillus
anthracis. Genome Res 22:1512–24.

20.   Hinnekens P, Fayad N, Gillis A, Mahillon J. 2022. Conjugation across *Bacillus cereus* and kin:
A review. Front Microbiol 13:1034440.

21.   González JM, Jr., Brown BJ, Carlton BC. 1982. Transfer of *Bacillus thuringiensis* plasmids
coding for delta-endotoxin among strains of *B. thuringiensis* and *B. cereus*. Proc Natl Acad Sci
U S A 79:6951–5.

22. Thomas DJ, Morgan JA, Whipps JM, Saunders JR. 2001. Plasmid transfer between *Bacillus*
thuringiensis subsp. i*sraelensis* strains in laboratory culture, river water, and dipteran larvae.
Appl Environ Microbiol 67:330–8.
23. Vilas-Bôas GFLT, Vilas-Bôas LA, Lereclus D, Arantes OMN. 1998. *Bacillus thuringiensis*
conjugation under environmental conditions. FEMS Microbiology Ecology 25:369–374.
24. Liu Y, Lai Q, Göker M, Meier-Kolthoff JP, Wang M, Sun Y, Wang L, Shao Z. 2015. Genomic
insights into the taxonomic status of the *Bacillus cereus* group. Sci Rep 5:14082.
25. Geng P, Zhao P, Wan X, Mahillon J, Hu Y, Gong Y, Hu X. 2023. Interspecies Horizontal
Transfer and Specific Integration of the Mosquitocidal Toxin-Encoding Plasmid pTAND672-2
from *Bacillus thuringiensis* subsp. *israelensis* to *Lysinibacillus sphaericus*. Appl Environ
Microbiol 89:e0165222.
26. Moreno-Peña Carlos J, García-García Jorge H, Hernández-Vásquez Cesar I, Galán-Wong Luis
J, Pereyra-Alférez B. 2021. Draft Genome Sequence of *Bacillus toyonensis* Strain GM18,
Isolated from Agricultural Soil. Microbiology Resource Announcements
10:10.1128/mra.00008–21.
27. Berry C, Valby V, Mishra R, Bonning BC, Palma L, Crickmore N. 2025. Specificity database
for bacterial pesticidal proteins against invertebrate targets. Journal of Invertebrate Pathology
211:108319.
28. Konstantinidis KT, Tiedje JM. 2005. Genomic insights that advance the species definition for
prokaryotes. Proc Natl Acad Sci U S A 102:2567–72.
234
235
236

 Figure Legends

 Figure 1. Genome BLAST Distance Phylogeny (GBDP) [18] based on genome data (whole-genome

 sequence-based) using TYGS server. The *B. toyonensis* cluster is highlighted in green color. The ex-

 clamation marks indicate potentially misclassified *B. toyonensis* strains.

 Tables

 Table 1. % ANI calculation estimated by comparison with type strain *B. toyonensis* P18, showing the

 reclassification proposal following the nomenclature system by Carroll et al., (2020). A 95-96% ANI is

 generally accepted as the species boundary (28), therefore, genomes that share 95-96% ANI or higher

 may be considered as belonging to the same species.

Species	Strain	Acc. Number	% ANI	PubMLST support	New species proposal
B. thuringiensis	MC28	CP003687	99.5	100%	*B. toyonensis* biovar Thuringiensis
B. cereus	Rock 1-3	ACMG00000000	98.7	100%	*B. toyonensis*
Bacillus sp.	Bto_UNVM-42	JAYEPS000000000	98.2	100%	*B. toyonensis* biovar Thuringiensis
B. thuringiensis*	Bto_UNVM-94	NZ_QGLX00000000	98.6	100%	*B. toyonensis* biovar Thuringiensis
Bacillus sp.	Bto_UNVM-100	JAYGOI000000000	98.6	100%	*B. toyonensis* biovar Thuringiensis
B. toyonensis	GM18	JADNYS000000000	98.5	100%	*B. toyonensis* biovar Thuringiensis

 *This strain has been already named as *B. toyonensis* biovar Thuringiensis (NCBI:txid2923195).

 Table 2. Putative insecticidal gene content of *B. toyonensis* strains.

Strain	Homolog Proteins	Potential targets	Parasporal crystals	References
MC28	Cry4C, Cry30F, Cry53A, Cry54A, Cry68A, Cry69A, Cry70B, Cyt1D, Cyt2A	Lepidoptera and Diptera	Spherical	(18)
Bto_UNVM-42	Cry32E, Cry32G, Cry32K, Cry32M, Cry32U, Cry69A, Cry73B, Cyt1A, Cyt1D, Mpp3A, Mpp23A	Lepidoptera, Coleoptera and Diptera	Bipyramidal	This work
Bto_UNVM-94	Cry7G, Mpp2A	Lepidoptera and Coleoptera	Bipyramidal	(7)
Bto_UNVM-100	Cry7B, Cry7G, Cry73A, Mpp2A, Xpp22A	Lepidoptera and Coleoptera	Bipyramidal	This work
GM18	Xpp22A, App4A, Tpp78A, Vpb1A, Mpp75A, Vpb1E	Human-cancer cells	Produced, morphology is not specified	(26)

B. cereus Rock 1-3: Warning: unverified source organism.
Status: RefSeq GCF_000161155.1 is suppressed. doi: 10.1101/gr.134437.111
B. thuringiensis MC28: Warning: unverified source organism.
Status: RefSeq GCF_000300475.1 is suppressed. doi: 10.1128/JB.01861-12