Title: Relative fire-proneness of land cover types in the Brazilian Atlantic Forest

Authors: Bruno FCB Adorno [1,2], Augusto AJ Piratelli [3], Erica Hasui[4], Milton C Ribeiro[5], Pedro G. Vaz* [2]

[1] Centro de Ciências Biológicas e da Saúde, Universidade Federal de São Carlos, Sorocaba-SP, Brazil

[2] CE3C-Centre for Ecology, Evolution and Environmental Changes & CHANGE, Global Change and Sustainability Institute, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, C2, Piso 5, 1749-016 Lisboa, Portugal

[3] Departamento de Ciências Ambientais/CCTS, Universidade Federal de São Carlos, Sorocaba-SP, Brazil

[4] Instituto de Ciências da Natureza, Universidade Federal de Alfenas, Alfenas-MG, Brazil

[5] Laboratório de Ecologia Espacial e Conservação (LEEC), Departamento de Ecologia, Instituto de Biociências, Universidade Estadual Paulista (UNESP), Rio Claro-SP, Brazil

*****Corresponding author**: pjvaz@fc.ul.pt

Highlights

- First evaluation of land cover fire-proneness in a tropical biome using selection ratios.

- Secondary forests burned 61% more; old-growth forests 57% less than expected by chance.

- Pastures in the Atlantic Forest seem less fire-prone than expected, despite frequent burns.

- Secondary forest proneness decreased, likely due to maturation and better management.

- Tailored fire management is essential to protect secondary forests in restoration efforts.

ABSTRACT

Fires have long posed a global problem, but they are now increasingly affecting tropical biomes where landscape-fire interactions remain understudied. We investigate the fire-proneness of different land use and land cover types (LULCs) within the Brazilian Atlantic Forest (AF), one of the world's most important biodiversity hotspots, experiencing increased fire frequency due to human activities and climate change. Using a selection ratio-based approach, we analyzed fire-LULC interactions in 40,128 fires over a 35-year period (1987-2022) across various vegetational regions in the AF, evaluating the relative fire-proneness of different LULCs in a tropical biome for the first time. Our findings revealed that secondary forests, forest areas that have regrown after major disturbances, burned 61% more than expected by chance, whereas old-growth forests, native forests that have developed over very long periods, burned 57% less than expected, highlighting a nearly inverse relationship in their fire-proneness. Interestingly, our data indicate that pastures in the AF are less prone to fire than expected, despite being considered among the land uses that burn the most in Brazil. Other LULCs showed variable fire-proneness, with some differences between vegetational regions. Over time, the fire-proneness of secondary forests decreased, likely due to forest maturation and changes in land management practices. We emphasize the necessity for tailored fire management strategies that address the unique vulnerabilities of secondary forests, particularly in the context of ongoing restoration efforts aimed at increasing native forests. Effective measures, including the implementation of 'fire-smart management' practices and enhancing the perceived value of secondary forests among local communities, are crucial for mitigating fire risks. Integrating these strategies with incentive-based approaches can bolster fire prevention, ensuring the long-term success of restoration programs. Our study provides a framework for understanding fire-landscape

dynamics in tropical forests and offers actionable insights for practitioners working to safeguard these biomes from the escalating threat of wildfires.

KEYWORDS: Wildfire; Fire-proneness; Land cover; Secondary forests; Old-growth forests; Fire management; Atlantic Forest

1. INTRODUCTION

Tropical forests are fire-sensitive biomes, with species lacking evolutionary adaptations for such disturbance (Hoffmann et al., 2012; Pivello et al., 2021). Historically, these biomes experienced fires every few hundred years (Bush et al., 2008), but climate change and human activities have altered the frequency and nature of these events (Cochrane, 2003; Fonseca et al., 2019; Le Page et al., 2017). Fires now occur every decade or even annually, often escalating in size and severity (Cochrane, 2003; Flannigan et al., 2009; Uhl & Kauffman, 1990). Fires are associated with biodiversity changes (Kelly et al., 2017; Kelly et al, 2020), soil degradation (Nadporozhskaya et al., 2018), erosion (Girona-García et al., 2024), impacts on freshwater ecosystems (Vaz et al., 2014, 2021), diminished water retention (Sansevero et al., 2017; Schmerbeck & Fiener, 2015), and landscape dynamics alterations (Silva et al., 2011; Moreira et al., 2011).

Fire growing impacts are particularly worrying in the Atlantic Forest (AF) biome, one of the world's most important biodiversity hotspots, with a high number of endemism (Myers et al., 2000). Originally, the AF covered 160 million hectares across coastal and inland portions of Brazil, Argentina, and Paraguay (Muylaert et al., 2018). This biome is the most degraded and deforested Brazilian biome, having suffered from intense land use and land cover (LULC) changes in recent centuries (Rezende et al., 2018). Most old-growth forests—native forests that have developed over long periods without significant disturbance—were cleared to make way for human activities. Today, the AF retains only about 36% of its original extent of natural vegetation. Most of this vegetation is highly fragmented, with 97% of forest patches being under 50 ha, isolated, and impacted by various human-induced disturbances (Ribeiro et al., 2009a; Vancine et al., 2024). Despite these challenges, recent decades have seen improvements due to

conservation efforts led by the government and environmental organizations, reducing deforestation rates (Piffer et al., 2022a). Also, land abandonment and the depopulation of rural areas contributed to the resurgence of natural vegetation, particularly in the form of secondary forests—areas of forest that have regrown after major disturbances such as fires, whose ecological value and recovery potential depend on the time since disturbance and the surrounding landscape context (Rezende et al., 2018; Souza et al., 2020; Vancine et al., 2024). Changes in Brazilian legislation, such as the Atlantic Forest Law in 2006 and the Brazilian Forest Code in 2012, are also expected to influence the fire-landscape dynamic over the years. However, despite strict regulations, fire is increasingly shaping the AF landscape (dos Santos et al., 2019), namely by resetting the gradual transitions from secondary forests to vegetation resembling old-growth forests. Fires disrupt regeneration processes and can shift the biome towards a savanna-like state (dos Santos et al., 2019; Sansevero et al., 2020).

Fires do not occur randomly across the LULC types of a given landscape; instead, their occurrence and spread are determined by a complex interplay of human actions and natural phenomena (Brando et al., 2016; Fonseca et al., 2019; Le Page et al., 2017; Pivello, 2011). In tropical forests, most fires are caused by humans as a result of using fire to clear or expand agricultural areas (da Silva Junior et al., 2020; Gois et al., 2020; Oliveira-Júnior et al., 2020). Fire propagation through the landscape is influenced by wind, topography, vegetation structure, moisture content, composition of fuel loads (Carmo et al., 2011; Moreira et al., 2009; Oliveira et al., 2014; Silva et al., 2009), and factors impacting firefighting efforts (de Assis Barros et al., 2022; Gutiérrez-Velez et al., 2014; Pivello et al., 2021). Consequently, some LULC types are more susceptible to fire occurrence than others (Abreu et al., 2022; Fonseca et al., 2019; Gutiérrez-Vélez et al., 2014). Under moderate weather conditions, LULCs that are more prone to fire are likely

to be disproportionately affected, beyond what would be expected by chance alone (Moreira et al., 2001).

In Brazilian tropical forests, fires disproportionately affect LULC types, with pastures and croplands being the most likely to burn (Abreu et al., 2022; Cano-Crespo et al., 2015; de Assis Barros et al., 2022; de Santana et al., 2020; Freitas et al., 2020; Gutiérrez-Vélez et al., 2014; Herrmann et al., 2023; Uriarte et al., 2016). Conversely, a negative correlation is expected between fire occurrence and spread and the cover of old-growth forests (de Assis Barros et al., 2022; Fonseca et al., 2019; Singh & Huang, 2022). This low fire-proneness among old-growth forests can be attributed to their remarkable retention of transpired moisture, which effectively reduces susceptibility to fire (Cochrane et al., 1999; Uhl et al., 1988). Unfortunately, ongoing transitions from old-growth forests to other types of LULC can often result in increased fire-proneness (de Assis Barros et al., 2022; Eva & Lambin, 2000; Fonseca et al., 2019; Singh & Huang, 2022). Particularly, changes from old-growth forests to secondary forests likely increase fire-proneness, as secondary forests are more vulnerable due to their differing species composition, lower moisture content, and elevated light and temperature conditions (Fonseca et al., 2019; Gutiérrez-Vélez et al., 2014). Secondary forests in the AF typically emerge in degraded, abandoned pasturelands unsuitable for agriculture and livestock, and may also regenerate on the edges of small forest fragments, thus increasing the risk of fire (Guedes et al., 2020). As secondary forests mature and develop characteristics typical of old-growth forests, their susceptibility to fire is expected to decrease (Lebrija-Trejos et al., 2011; Ray et al., 2010).

Despite the extensive research on landscape-wildfire dynamics, critical gaps remain in our understanding of how fire-proneness varies among different LULC within Tropical

Forests. In Brazil, the bulk of fire research has been conducted in the Amazon Forest, leading to a disproportionate focus that overlooks the distinct dynamics within the AF (dos Santos et al., 2019; Sansevero et al., 2020). Furthermore, no previous studies have comprehensively assessed the fire-proneness of different LULCs in Brazilian Tropical forests (see Carmo et al., 2011; Moreira et al., 2009; Oliveira et al., 2014). Instead, research has typically focused on examining the distribution of burned areas and the number of fire foci (Abreu et al., 2022; de Assis Barros et al., 2022; Gutiérrez-Vélez et al., 2014). Moreover, existing studies often overlook the heterogeneity of vegetational regions in the AF and their cultural and socio-economic differences, which likely impact fire dynamics. These regions, including moisture-dependent (ombrophilous) and seasonal forests, as well as transitional ecotone areas, are shaped by distinct temperature and rainfall patterns (Joly et al., 2014). Their distribution and fire susceptibility are also influenced by the AF's extensive longitudinal, latitudinal, and altitudinal variations (Marques et al., 2021). A lack of long-term studies further limits understanding of fire-LULC dynamics over time.

In this study, we used a selection ratio-based approach (e.g., Moreira et al., 2009) to analyze relative fire-LULC interactions across the vegetational regions of the AF, examining data at five-year intervals over a 35-year period. We tested the following hypotheses. 1) Old-growth forests have low fire-proneness, irrespective of vegetational region or year analyzed. 2) Secondary forests exhibit high fire-proneness in all vegetational regions compared to old-growth forests. 3) Secondary forests exhibit a decreasing trend in fire-proneness over the years as the average age of their patches increases, making them resemble old-growth forests more closely. 4) Croplands, pastures, and other human-impacted LULCs exhibit variable fire-proneness dependent on vegetational region and year.

2. METHODS

2.1 Vegetational regions

To assess how vegetational regions affect fire-LULC interactions in the Brazilian AF, we analyzed the five regions that cover over 90% of the biome (Fig. 1) and record the most fires (IBGE, 2012):

1) Semideciduous seasonal forest (SSF): Covering 37% of the biome, this largest vegetational region in the AF is marked by two distinct seasons—rainy summers and dry winters. During the dry season, up to 50% of tree species undergo leaf shuddering.

2) Decidual seasonal forest (DSF): Occupying 6% of the AF, these forests prevail at higher altitudes where temperatures are lower. Like the SSF, they experience two distinct seasons—rainy summers and dry winters. Over 50% of their tree species shed leaves during the dry season.

3) Dense ombrophilous forest (DOF): This region occupies 17% of the AF, forming a verdant belt along the Brazilian coast. Characterized by the absence of a dry season, it experiences consistently high precipitation year-round and maintains temperatures averaging above 25°C.

4) Mixed ombrophilous forest (MOF): Commonly known as 'Araucaria forests', these forests are notable for consistent high precipitation and a diverse mix of evergreen and deciduous species. Covering 15% of the AF, they are distinguished by a high prevalence of gymnosperm species, notably *Araucaria angustifolia*.

5) Contact areas (CA): Covering 15% of the AF, these undifferentiated plant communities are transitional ecotone areas between two or more vegetational regions.

To delineate each vegetational region, we utilized the vector files accessible from the Database of Environmental Information provided by the Brazilian Institute of Geography and Statistics (IBGE), which can be found at: https://bdiaweb.ibge.gov.br.

2.2 Land cover maps

To obtain maps of LULC types in the vegetational regions of the AF, we used maps from the MapBiomas platform - Collection 8.0 (available at https://brasil.mapbiomas.org). This project produces annual LULC maps with a 30m resolution covering the period from 1985 to 2022 (Alencar et al., 2022). The platform also provides a collection focused on deforestation and secondary vegetation, identifying transitions from anthropogenic to forest classes and providing information on the age of secondary forests (Silva Junior et al., 2020).

We selected eight LULC maps, one for each five-year interval, spanning from 1986 to 2021 (Vancine et al., 2024). This interval helps minimize temporal autocorrelation, enhancing the independence of subsequent assessments of fire-LULC interactions. Next, we simplified the 30 LULC classes from MapBiomas Collection 8.0 into seven categories (Table 1), plus a 'Noncombustible areas' type, which was omitted from further analysis. This reclassification was based on structural similarities and potential for comparable fire behavior, particularly in terms of fuel structure and composition (Nunes et al., 2005).

2.2 Fire data

To obtain maps of fires in the vegetational regions of the AF, we used the MapBiomas platform – Fire Collection 2.0, which includes annual records of fire scars from 1985 to 2022. To ensure greater accuracy in identifying the parts of the LULC that

burned, we aligned our fire maps with our LULC data collection, which spans from 1986

to 2021 with a five-year interval, selecting fire maps from years that corresponded to one

year after each LULC map. Consequently, our fire maps, also selected at five-year

intervals, span from 1987 to 2022. We analyzed fires of five hectares or larger (see

Pereira and Santos, 2003; Moreira et al., 2009; Silva et al., 2009; Carmo et al., 2011), and

assigned each fire to a specific vegetational region of the AF.

2.3 Data analysis

To determine the relative fire-proneness of each LULC type, we employed a

selection ratio-based approach. This method involved comparing, for each fire, the

proportion of each LULC type within the burned area to its proportion in the total area

(burned and unburned) available for burning. We defined the unburned area available for

burning as a buffer zone surrounding each fire's perimeter, sized to four times the area of

the fire itself (see similar procedures in Silva et al., 2009; Moreira et al., 2009; Oliveira et

al., 2014). Thus, the total area available to burn consisted of the fire plus this buffer. This

method was originally proposed for the study of resource selection by animals (Manly et

al., 1993) and has been applied in fire research studies (e.g., Moreira et al., 2009; Silva et

al., 2009; Carmo et al., 2011; Oliveira et al., 2014).

The selection ratio (w_i) for a given LULC type i is an index estimated as $w_i = o_i/\pi_i$

(Manly et al., 1993), where o_i is the proportion of a burned patch belonging to LULC

type i (estimated from the area consumed by fire) and π_i is the proportion of available

area belonging to LULC type i (estimated from both the burned patch and the

surrounding buffer). Selection ratios range from 0, if the LULC type is available but not

consumed by the fire ($o_i = 0$), to a large value (in theory ∞), in the situation where the fire

consumes the only tiny patch of a given LULC type that was available to burn (π_i

approaches 0) (Moreira et al., 2009). To avoid these extreme *w* values in very small areas, we only analyzed LULC patches larger than 1 ha. If a given LULC type is burned in proportion to its availability, then *w*=1. If *w*>1, the LULC type is considered more fire-prone than expected by chance. Conversely, if *w*<1, the LULC type is considered less fire-prone than expected by chance. Since we were interested in the fire-proneness of LULC types per vegetational region, we omitted fires that, with their buffers, overlapped more than one region. A total of 40,128 burned patches were considered for analysis.

Next, we averaged values for each LULC type across the fires within a given vegetational region and estimated 95% confidence intervals. We considered differences between selection ratios for different classes significant when their respective confidence intervals did not overlap. Additionally, if the confidence interval of *w* did not include 1, we considered the class significantly more or less prone to fire than expected by chance.

3. RESULTS

Of the 40,128 fires analyzed between 1987 and 2022, 44% occurred in the SSF vegetational region, 21% in CA, and the lowest percentages were in MOF (9%) and DSF (7%) (Table 2). The mean fire size was 23 ha, ranging from 15 ha in MOF to 27 ha in SSF. Mosaic of uses, secondary forests, and old-growth forests, in that order, were the LULC types with the highest available areas across regions. The region with the highest proportion of old-growth forests coincides with the area with the most fires, the SSF. CA had the highest proportion of secondary forests.

3.1 Fire-proneness across vegetational regions

Across the vegetational regions of the Brazilian AF, secondary forests were the most fire-prone type of LULC, while old-growth forests were the least fire-prone (Fig. 2).

Secondary forests burned 61% more than expected by chance ($w = 1.61$), whereas old-growth forests burned 57% less ($w = 0.43$). Besides secondary forests, only the mosaic of uses LULC type was significantly prone to fire, with a selection ratio 5% higher than expected. Like old-growth forests, the LULC types openlands, temporary crops, and planted forests were also significantly less prone to fire than expected, with selection ratios between 31% and 35% below expected. Perennial crops were the only type of LULC to burn according to availability across AF's vegetational regions (confidence interval of w included 1).

3.2 Fire-proneness by vegetational region

As hypothesized, old-growth forests had low fire proneness across vegetational regions (Fig. 3), burning between 33% and 66% less than expected by chance, depending on the region. Except for the DSF region, this LULC type had the lowest selection ratio in all other regions. In contrast, with selection ratios between 46% and 87% above expected, secondary forests tended to have the highest fire proneness across regions, except for the DOF. Openlands had low fire-proneness, burning 20% to 59% less than expected, except in MOF where they burned according to availability. As expected, other human-impacted LULC types showed variable fire proneness, depending on the vegetational region.

3.3 Fire-proneness over time

The fire-proneness of old-growth forests remained consistently low over the 35-year study period (Fig. 4), showing no pattern of increase or decrease. Likewise, openlands and temporary crops generally had low fire-proneness, with no clear change over time, although temporary crops burned according to availability in 1992. Perennial crops, which generally burned in proportion to availability until 2012, became less prone

to burning than expected from that year onwards. Forest plantations consistently had low fire-proneness, except in 1992 and 2002 when their fire-proneness was neutral. Mosaic of uses showed mixed fire-proneness over the years but shifted to a significantly lower than expected fire-proneness from 2012 onwards. It burned in proportion to its availability in 1992 and 2012, had high fire-proneness in 1987 and 1997–2007, but had low fire-proneness in 2017 and 2022. Secondary forests followed the hypothesized trend of decreasing fire-proneness, being significantly prone to fire until 2012, then burning in proportion to their availability from that year onwards.

4. DISCUSSION

Our analysis spanning over 35 years of fire-LULC interactions in the Brazilian AF vegetational regions provides insight into the relative fire-proneness of different LULCs in a tropical biome. This is critical as fire becomes a growing issue both worldwide and in these biomes. Across all vegetational regions, our results confirmed the low susceptibility of old-growth tropical forests to fire, in contrast with the high susceptibility of secondary forests to this disturbance. Other LULCs showed more variable fire-proneness, with some differences between vegetational regions. Furthermore, fire-proneness has changed over time in several LULCs, suggesting that these trends may be driven by natural ecological succession, climatic conditions, or human actions potentially influenced by changes in Brazilian environmental legislation. Notably, we found a decline in the fire-proneness of secondary forests over time, likely due to their lower susceptibility as the average age of these patches in the landscape increases, resulting in characteristics increasingly similar to old-growth forests.

4.1 Fire-proneness in the Brazilian Atlantic Forest

By confirming the low susceptibility of old-growth tropical forests to fire, our results are consistent with studies indicating a reduced risk and density of fires in undisturbed forests within the AF (de Assis Barros et al., 2022; Guedes et al., 2020; Singh & Huang, 2022). Despite the significant fragmentation and deforestation in the AF (Ribeiro et al., 2009; Vancine et al., 2024), we demonstrated that, in general, old-growth forest fragments still remain the LULC type least likely to burn compared to all others. The low fire susceptibility of tropical old-growth forests can be attributed to their retention of transpired moisture and the trees' substantial contribution to forest humidity, which limits fire ignition and spread (Cochrane, 2003a; Cochrane et al., 1999). Interestingly, our data showed that the likelihood of old-growth forests burning less than expected across regions was similar to that of secondary forests burning more than expected, both at approximately 60%.

The high susceptibility of secondary forests to fire was anticipated, as they exhibit high fuel load, low moisture content, and high luminosity and temperature in the understory (Gutiérrez-Vélez et al., 2014; Hasselquist et al., 2010; Kyereh et al., 2007). This is also consistent with previous works in the Amazon Forest that found higher fire activity in degraded forests (Balch et al., 2015; Brando et al., 2014, 2016; Cochrane et al., 1999). Research further suggests that, because secondary forests are often viewed as having low economic value, farmers may be less motivated to prevent them from burning when nearby fires pose a threat (Sorrensen, 2000). Other studies also show a link between deforestation fires to expand agricultural areas and the high susceptibility of secondary forests to fire (Pivello et al., 2021). Despite Brazilian legislation prohibiting fires, except in specific situations, farmers still burn secondary forests in pastures and

agricultural areas to prevent forest encroachment. Furthermore, these areas are often re-cleared within the first few years following forest establishment (Piffer et al., 2022).

Among the LULC types significantly less prone to fire, the results for openlands, where pastures are included, were noteworthy. Despite pastures having the largest burned areas in Brazil (Alencar et al., 2022; Moreira de Araújo et al., 2012) and being generally associated with fire risk (Guedes et al., 2020; Gutiérrez-Vélez et al., 2014; Herrmann et al., 2023), our findings indicate lower-than-expected fire proneness. This shows the advantage of our selection ratio-based approach, as this method allows a distinction to be made between when a LULC type burns extensively due to its abundance in the landscape and when it burns disproportionately more than its availability. On the contrary, previous works have generally highlighted the large representativeness of pastures within the burned areas, overlooking the total available area that could have potentially burned in the landscape, which is usually also substantial. Indeed, although pastures occupy around 30% of the AF (Dos Santos et al., 2022; Souza et al., 2020), our results show that their fire proneness is likely lower than expected when availability per fire is taken into account. The high association of fire with pastures may not be due to a positive selectivity compared to other land covers, but rather to the frequent illegal use of fire for cleaning and renewal pastures (Moreira de Araújo et al., 2012). It should be noted that although our results indicate lower fire selectivity for pastures than expected by chance, pasture fires still pose a threat to the AF by contributing to the spread of fires to other fire-prone LULCs (Guedes et al., 2020).

Management-related aspects of the other LULC types that were significantly less prone to fire, such as temporary crops and planted forests, or that burned according to availability, like perennial crops, should be stressed. Managed croplands usually exhibit low fuel load and high moisture content due to irrigation (Duguy et al., 2007). Planted

forests, on the other hand, are often intensively managed by private companies to avoid fire hazards (Guedes et al., 2020; Mirra et al., 2017). Additionally, these LULCs are typically closer to urban areas, roads, and scattered populations. This proximity may result in more frequent fire occurrences (dos Santos et al., 2019; Barros et al., 2022), but it also facilitates faster fire detection and easier firefighting efforts (Moreira et al., 2009).

Lastly, it was to be expected that mosaic of uses would be prone to fire across vegetational regions. These areas typically consist of small properties where slash-and-burn practices are employed to renew the land and eliminate waste. Subsistence farmers and indigenous populations often use these fires, confining them to small areas with localized impacts (Pivello et al., 2021). However, there is a significant risk that these fires could accidentally spread to nearby fire-prone LULCs. Under climate change scenarios, the risk of fire increases, especially when traditional agricultural fire practices result in accidents or loss of control (Pivello et al., 2021). It is crucial to develop strategies that balance traditional practices with modern fire management to mitigate these risks effectively.

4.2 Fire-proneness by vegetational region

Although it was hypothesized and therefore unsurprising that the old-growth forests of the AF exhibit consistently low fire-proneness with minimal variation between vegetational regions, it is nonetheless remarkable that this low fire-proneness remains uniform across regions characterized by dry seasons, such as SSF and DSF, as well as in regions with evergreen vegetation, like ODF. These results reinforce the reduced susceptibility of tropical forests to fires (Cochrane, 2003; Cochrane et al., 1999), even under disparate regional conditions. Yet, previous studies have also shown that altitude, mean annual temperature, and drought severity can be positively associated with fire

occurrence and lead to variations in the area burned in the AF (Abreu et al., 2022; de Assis Barros et al., 2022; de Santana et al., 2020; Herrmann et al., 2023; Singh & Huang, 2022).

In the AF, regions with varying economic development can show different success rates in restoration programs (Piffer et al., 2022), affecting fire prevention and control. In the Brazilian Amazon, studies show that federal and state agencies, with distinct institutional capacities, implement diverse fire management practices, contributing to regional differences in fire management (Fonseca-Morello et al., 2017). In Portugal, research also shows that regions with different climatic and environmental conditions present LULCs with different fire-proneness (Moreira et al., 2009). In our study, this is likely why the LULC openlands type, which generally had low fire-proneness in most AF vegetational regions, exhibited higher fire-proneness, burning according to its availability, in the MOF region, a region with high average altitudes (IBGE, 1992).

Despite the environmental, cultural, social, and economic differences between vegetational regions, the secondary forest LULC type burned more than expected by chance in all AF regions, as predicted. Not even the consistently high rainfall in evergreen vegetational regions such as ODF (Oliveira et al., 2000) reduces the susceptibility of secondary forests to fire, highlighting the vulnerability of this LULC to such disturbance. Still, our results show that secondary forests are almost twice as fire-prone in CA as in DSF, with selection ratios 87% and 46% higher than expected, respectively. This may be due to the fact that CAs are mainly transition zones between the AF biome and the Brazilian Cerrado. Unlike the AF, the Brazilian Cerrado is a fire-dependent system, with ecological communities adapted to this disturbance (Pivello et al., 2021). As such, the greater proximity of CA's secondary forests to the Cerrado is likely to explain their greater fire proneness. Likewise, as in the Cerrado, where cattle

ranching is the leading cause of fire (Pivello et al., 2011), secondary forests in the CA regions are also burned to expand pasturelands. Worryingly, fires in secondary forests can promote the growth of herbaceous species and lead to the replacement of forests by open physiognomies (Pivello & Coutinho, 1996; Sansevero et al., 2020).

4.3 Fire-proneness of the Brazilian Atlantic Forest over time

Our results also confirmed the hypothesis that AF old-growth forests would exhibit consistently low fire-proneness over time, with minimal variation across the 8 years sampled from 1987 to 2022. Not even the inclusion of a major El Niño year (1996–1997) among the years sampled altered this consistency, despite major El Niño years being associated with extreme droughts and a greater potential for fires to occur and spread (Aragão et al., 2018; Marengo et al., 2018). In fact, this El Niño event does not seem to have altered the fire-proneness in 1997 or the chronosequential fire-proneness pattern for any LULC type studied. However, the years sampled did not include other extreme climatic events, such as the major La Niña (characterized by high precipitation) in 1988–1989 or the major El Niño in 2015-2016. Therefore, it remains possible that some LULC types had unusual fire-proneness patterns during these events. Future studies on fire-landscape interactions in the AF should include diverse climatic situations, such as those with and without major El Niño and La Niña events.

The hypothesized decreasing fire-proneness trend of secondary forests over time was confirmed and the result can be discussed in light of two main reasons. First, the early data from the MapBiomas collection that we used to obtain the LULC maps only included young secondary forests. Over time, the collection encompassed a broader range of ages for this LULC type, leading to an increase in the average age of secondary forest patches included in the annual maps. Thus, the reduced fire-proneness of secondary

forests in more recent years, burning in proportion to availability from 2012 onwards, is likely partly due to the increased age of these forests. Forest succession leads to significant environmental changes, with early stages having higher temperatures and lower humidity in the understory, transitioning to cooler and more humid conditions in later stages (Ray et al., 2010; Lebrija-Trejos et al., 2011). However, the approximate maximum age of the secondary forest patches in our data in 2022, 35 years, combined with the existence of younger patches, means that this LULC type has not yet reached the tipping point of having characteristics close enough to old-growth forests to become equally less prone to fire. Future studies must analyze the age of each patch of secondary forest so that we can understand more precisely after how long this type of LULC becomes less prone to fire than expected by chance.

The second reason for the decreasing fire-proneness is that early-stage secondary forests are often burned to prevent encroachment on agricultural areas. In the AF, regenerated forests in pastures and areas of shifting cultivation—where land is alternately farmed and left to regenerate—are frequently re-cleared before reaching advanced succession stages (Piffer et al., 2022), thus avoiding stricter protection under the Atlantic Forest Law (Law No. 11.428/2006). This law highly restricts cutting native vegetation, allowing it only in initial and medium regeneration stages with appropriate authorization, provided it follows sustainable management criteria. Cutting vegetation over 5 years old is prohibited. On the other hand, legislation changes may also be behind the shift to low fire-proneness from 2012 onwards for the perennial crops and mosaic of uses LULC types, after the approval of the New Brazilian Forest Code (Law nº 12.651/2012). Firefighting may well have become more effective in these LULCs after the enactment of this law, which devotes a chapter to the prohibition and control of wildfires.

4.4 Management implications

Our results show that secondary forests have disproportionately burned the most across all regions of the AF in the past few decades, emphasizing the need for fire management strategies to carefully address this LULC. This is even more crucial considering that numerous restoration programs have restored millions of hectares of native vegetation, primarily as secondary forests, in recent decades (dos Santos et al., 2019; Brancalion et al., 2019; Vancine et al., 2024). It is clear that a trade-off must be established between the benefits of increasing secondary forest areas and the costs of protecting them, especially in the early years, to avoid a greater risk of fire in the landscape. Fire will lead to the interruption of forest succession to mature forests (Cochrane, 2001; Cochrane et al., 1999; Sansevero et al., 2020). Implementing effective fire prevention and management measures in secondary forests will be necessary to safeguard the restoration efforts and ensure their long-term success (dos Santos et al., 2019). Some authors simply argue that in the face of deforestation fires, which are illegal but still used in secondary forests, law enforcement should be the primary response, defending that fines should then be used to fund law enforcement (Pivello et al., 2021).

By showing such a large discrepancy in fire-proneness between secondary forests and other LULC types, our data also suggest incorporating integrated landscape approaches from fire-prone landscapes in Mediterranean regions and North America (Pivello et al., 2021). One such approach is 'fire-smart management', which involves managing the entire landscape to reduce fire risk. In this context, implementing low fire risk matrices composed of LULC types that our results have shown to be less prone to fire around secondary forests can be an effective strategy to prevent the spread of fire. For example, there is evidence that eucalyptus plantations provide some protection against fire, because plantation owners seek to suppress and control fires that could

otherwise harm their eucalyptus investments (Guedes et al., 2020). Using this type of LULC around more fire-prone patches, including mosaic of uses, may be an option to consider.

The fire-proneness of secondary forests in the AF also highlights the need to increase their perceived value by local communities to prevent them from burning (Guedes et al., 2014). This could involve enriching younger forest patches (e.g., up to 5 years old) with fruit or timber trees and offering payments for ecosystem services such as carbon sequestration, water regulation, and habitat restoration (Guedes et al., 2014). Some authors suggest that financial institutions could offer better loan conditions and payment rates to farmers with a good fire governance record (Nepstad et al., 2014). Similarly, commodity buyers may provide favorable terms to farmers that adhere to the rules (Pivello et al., 2021). Lastly, the AF should contain programs dedicated to forecasting and monitoring wildfires, like those in the Amazon Forest. In this scenario, plans to combat and prevent fires should be developed well in advance, taking into account alerts for areas with high deforestation rates, especially those with secondary forests (Pivello et al., 2021).

5. Conclusion

This study used a selection ratio-based approach in 40,128 fires spanning over 35 years to evaluate the relative fire-proneness of LULC types in a tropical biome for the first time. This method compares, per fire, the proportion of each burned land cover type with its availability nearby. Our findings quantified the variation in fire susceptibility across LULC types within the Brazilian AF, highlighting that the likelihood of secondary forests burning more than expected was approximately 60%, mirroring the 60% likelihood of old-growth forests burning less than expected. Interestingly, pastures,

included in openlands, exhibited lower-than-expected fire proneness despite their large burned areas in Brazil. Secondary forests burned more than expected in all AF regions, with a particularly high susceptibility in CA, transition zones between the AF and the Cerrado, a fire-dependent system. The observed trend of decreasing fire-proneness in secondary forests over time likely reflects both the increasing age and maturity of these forests and changes in land management practices.

We emphasize the necessity for tailored fire management strategies that address the unique vulnerabilities of secondary forests, particularly in the context of ongoing restoration efforts aimed at increasing forest cover. Effective measures, including the implementation of 'fire-smart management' practices and enhancing the perceived value of secondary forests among local communities, are crucial for mitigating fire risks. Integrating these strategies with robust law enforcement and incentive-based approaches can bolster fire prevention and control, ensuring the long-term success of restoration programs. Our analysis provides a framework for understanding fire-landscape dynamics in tropical forests and offers actionable insights for policymakers, conservationists, and land managers working to safeguard these biomes from the escalating threat of wildfires.

Acknowledgments

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 (Grant 88887.889095/2023-00). BA (Grant 140361/2021-9) and AJP (Grant 316032/2023-9) were funded by CNPq—Conselho Nacional de Desenvolvimento Científico e Tecnológico. PGV (CE3C DOI 10.54499/UIDB/00329/2020; CHANGE DOI 10.54499/LA/P/0121/2020) was funded by FCT—Portuguese Science and Technology Foundation.

Authors' contributions

Bruno Adorno: Conceptualization, Investigation, Formal analysis, Writing - Original Draft**,** Writing – Editing. **Augusto Piratelli:** Writing - Review & Editing, Supervision. **Erica Hasui:** Writing - Review & Editing. **Milton Ribeiro:** Writing - Review & Editing. **Pedro Vaz:** Conceptualization, Formal analysis, Writing - Original Draft, Writing - Review & Editing, Supervision.

References

Abreu, M. C., Lyra, G. B., de Oliveira-Júnior, J. F., Souza, A., Pobočíková, I., de Souza Fraga, M., & Abreu, R. C. R. (2022). Temporal and spatial patterns of fire activity in three biomes of Brazil. *Science of The Total Environment*, *844*, 157138. https://doi.org/10.1016/j.scitotenv.2022.157138

Alencar, A. A. C., Arruda, V. L. S., Silva, W. V. da, Conciani, D. E., Costa, D. P., Crusco, N., Duverger, S. G., Ferreira, N. C., Franca-Rocha, W., Hasenack, H., Martenexen, L. F. M., Piontekowski, V. J., Ribeiro, N. V., Rosa, E. R., Rosa, M. R., dos Santos, S. M. B., Shimbo, J. Z., & Vélez-Martin, E. (2022). Long-term landsat-based monthly burned area dataset for the Brazilian biomes using deep learning. *Remote Sensing*, *14*(11), 2510. https://doi.org/10.3390/rs14112510

Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., Silva, C. V. J., Silva Junior, C. H. L., Arai, E., Aguiar, A. P., Barlow, J., Berenguer, E., Deeter, M. N., Domingues, L. G., Gatti, L., Gloor, M., Malhi, Y., Marengo, J. A., Miller, J. B., … Saatchi, S. (2018). 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nature Communications*, *9*(1), 536. https://doi.org/10.1038/s41467-017-02771-y

Balch, J. K., Brando, P. M., Nepstad, D. C., Coe, M. T., Silvério, D., Massad, T. J., Davidson, E. A., Lefebvre, P., Oliveira-Santos, C., Rocha, W., Cury, R. T. S., Parsons, A., & Carvalho, K. S. (2015). The susceptibility of southeastern Amazon forests to fire: insights from a large-scale burn experiment. *BioScience*, *65*(9), 893–905. https://doi.org/10.1093/biosci/biv106

Brancalion, P. H. S., Niamir, A., Broadbent, E., Crouzeilles, R., Barros, F. S. M., Almeyda Zambrano, A. M., Baccini, A., Aronson, J., Goetz, S., Reid, J. L., Strassburg, B. B. N., Wilson, S., & Chazdon, R. L. (2019). Global restoration opportunities in tropical rainforest landscapes. *Science Advances*, *5*(7), 3223–3226. https://doi.org/10.1126/sciadv.aav3223

Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., Silvério, D., Macedo, M. N., Davidson, E. A., Nóbrega, C. C., Alencar, A., & Soares-Filho, B. S. (2014). Abrupt increases in Amazonian tree mortality due to drought–fire interactions. *Proceedings of the National Academy of Sciences*, *111*(17), 6347–6352. https://doi.org/10.1073/pnas.1305499111

Brando, P. M., Oliveria-Santos, C., Rocha, W., Cury, R., & Coe, M. T. (2016). Effects of experimental fuel additions on fire intensity and severity: unexpected carbon resilience of a neotropical forest. *Global Change Biology*, *22*(7), 2516–2525. https://doi.org/10.1111/gcb.13172

Bush, M. B., Silman, M. R., McMichael, C., & Saatchi, S. (2008). Fire, climate change and biodiversity in Amazonia: a Late-Holocene perspective. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1498), 1795–1802. https://doi.org/10.1098/rstb.2007.0014

Cano-Crespo, A., Oliveira, P. J. C., Boit, A., Cardoso, M., & Thonicke, K. (2015). Forest edge burning in the Brazilian Amazon promoted by escaping fires from managed pastures. *Journal of Geophysical Research: Biogeosciences*, *120*(10), 2095–2107. https://doi.org/10.1002/2015JG002914

Carmo, M., Moreira, F., Casimiro, P., & Vaz, P. (2011). Land use and topography influences on wildfire occurrence in northern Portugal. *Landscape and Urban*

Planning, *100*(1–2), 169–176.

https://doi.org/10.1016/j.landurbplan.2010.11.017

Cochrane, M. A. (2001). Synergistic interactions between habitat fragmentation and

fire in evergreen tropical forests. *Conservation Biology*, *15*(6), 1515–1521.

https://doi.org/10.1046/j.1523-1739.2001.01091.x

Cochrane, M. A. (2003a). Fire science for rainforests. *Nature*, *421*(6926), 913–919.

https://doi.org/10.1038/nature01437

Cochrane, M. A. (2003b). Fire science for rainforests. *Nature*, *421*(6926), 913–919.

https://doi.org/10.1038/nature01437

Cochrane, M. A., Alencar, A., Schulze, M. D., Souza, C. M., Nepstad, D. C.,

Lefebvre, P., & Davidson, E. A. (1999). Positive feedbacks in the fire dynamic

of closed canopy tropical forests. *Science*, *284*(5421), 1832–1835.

https://doi.org/10.1126/science.284.5421.1832

da Silva Junior, C. A., Teodoro, P. E., Delgado, R. C., Teodoro, L. P. R., Lima, M.,

de Andréa Pantaleão, A., Baio, F. H. R., de Azevedo, G. B., de Oliveira Sousa

Azevedo, G. T., Capristo-Silva, G. F., Arvor, D., & Facco, C. U. (2020).

Persistent fire foci in all biomes undermine the Paris Agreement in Brazil.

Scientific Reports, *10*(1), 16246. https://doi.org/10.1038/s41598-020-72571-w

de Assis Barros, L., Mendonça, B. A. F. de, Sothe, C., Fernandes Filho, E. I., &

Elkin, C. (2022). Fire in the Atlantic Rainforest: an analysis of 20 years of fire

foci distribution and their social-ecological drivers. *Geocarto International*,

37(16), 4737–4761. https://doi.org/10.1080/10106049.2021.1899301

de Santana, R. O., Delgado, R. C., & Schiavetti, A. (2020). The past, present and

future of vegetation in the Central Atlantic Forest Corridor, Brazil. *Remote*

Sensing Applications: Society and Environment, *20*, 100357.

https://doi.org/10.1016/j.rsase.2020.100357

dos Santos, C. O. dos, Mesquita, V. V., Parente, L. L., Pinto, A. de S., & Ferreira, L.

G. (2022). Assessing the wall-to-wall spatial and qualitative dynamics of the

brazilian pasturelands 2010–2018, based on the analysis of the Landsat data

archive. *Remote Sensing*, *14*(4), 1024. https://doi.org/10.3390/rs14041024

dos Santos, J. F. C., Gleriani, J. M., Velloso, S. G. S., de Souza, G. S. A., do

Amaral, C. H., Torres, F. T. P., Medeiros, N. D. G., & dos Reis, M. (2019).

Wildfires as a major challenge for natural regeneration in Atlantic Forest.

Science of The Total Environment, *650*, 809–821.

https://doi.org/10.1016/j.scitotenv.2018.09.016

Duguy, B., Alloza, J. A., Röder, A., Vallejo, R., & Pastor, F. (2007). Modelling the

effects of landscape fuel treatments on fire growth and behaviour in a

Mediterranean landscape (eastern Spain). *International Journal of Wildland*

Fire, *16*(5), 619. https://doi.org/10.1071/WF06101

Eva, H., & Lambin, E. F. (2000). Fires and land-cover change in the tropics:a

remote sensing analysis at the landscape scale. *Journal of Biogeography*, *27*(3),

765–776. https://doi.org/10.1046/j.1365-2699.2000.00441.x

Flannigan, M. D., Krawchuk, M. A., De Groot, W. J., Wotton, B. M., & Gowman,

L. M. (2009). Implications of changing climate for global wildland fire. In

International Journal of Wildland Fire (Vol. 18, Issue 5, pp. 483–507).

https://doi.org/10.1071/WF08187

Fonseca, M. G., Alves, L. M., Aguiar, A. P. D., Arai, E., Anderson, L. O., Rosan, T.

M., Shimabukuro, Y. E., & de Aragão, L. E. O. e C. (2019). Effects of climate

and land-use change scenarios on fire probability during the 21st century in the

Brazilian Amazon. *Global Change Biology*, *25*(9), 2931–2946.

https://doi.org/10.1111/gcb.14709

Freitas, W. K., Gois, G., Pereira, E. R., Oliveira Junior, J. F., Magalhães, L. M. S., Brasil, F. C., & Sobral, B. S. (2020). Influence of fire foci on forest cover in the Atlantic Forest in Rio de Janeiro, Brazil. *Ecological Indicators*, *115*, 106340. https://doi.org/10.1016/j.ecolind.2020.106340

Girona-García, A., Vieira, D., Doerr, S., Panagos, P., & Santín, C. (2024). Into the unknown: The role of post-fire soil erosion in the carbon cycle. *Global Change Biology*, *30*(6). https://doi.org/10.1111/gcb.17354

Gois, G. de, Freitas, W. K. de, & Oliveira Júnior, J. F. de. (2020). Spatial-temporal of fire foci in the state of Rio de Janeiro, Brazil. *Bioscience Journal*, *36*(3), 1008–1017. https://doi.org/10.14393/BJ-v36n3a2020-47769

Guedes, B. J., Massi, K. G., Evers, C., & Nielsen-Pincus, M. (2020). Vulnerability of small forest patches to fire in the Paraiba do Sul River Valley, southeast Brazil: implications for restoration of the Atlantic Forest biome. *Forest Ecology and Management*, *465*, 118095. https://doi.org/10.1016/j.foreco.2020.118095

Gutiérrez-Vélez, V. H., Uriarte, M., DeFries, R., Pinedo-Vásquez, M., Fernandes, K., Ceccato, P., Baethgen, W., & Padoch, C. (2014). Land cover change interacts with drought severity to change fire regimes in Western Amazonia. *Ecological Applications*, *24*(6), 1323–1340. https://doi.org/10.1890/13-2101.1

Hasselquist, N. J., Allen, M. F., & Santiago, L. S. (2010). Water relations of evergreen and drought-deciduous trees along a seasonally dry tropical forest chronosequence. *Oecologia*, *164*(4), 881–890. https://doi.org/10.1007/s00442-010-1725-y

Herrmann, P. B., Nascimento, V. F., Freitas, M. W. D. de, & Ometto, J. P. (2023).

    Fire in highland grasslands in the Atlantic Forest Biome, a burned areas time

    series analysis and its correlation with the legislation. *Frontiers in Remote*

    Sensing, *4*. https://doi.org/10.3389/frsen.2023.1099430

Hoffmann, W. A., Geiger, E. L., Gotsch, S. G., Rossatto, D. R., Silva, L. C. R., Lau,

    O. L., Haridasan, M., & Franco, A. C. (2012). Ecological thresholds at the

    savanna-forest boundary: how plant traits, resources and fire govern the

    distribution of tropical biomes. *Ecology Letters*, *15*(7), 759–768.

    https://doi.org/10.1111/j.1461-0248.2012.01789.x

Joly, C. A., Metzger, J. P., & Tabarelli, M. (2014). Experiences from the Brazilian

    Atlantic Forest: ecological findings and conservation initiatives. *New*

    Phytologist, *204*(3), 459–473. https://doi.org/10.1111/nph.12989

Kelly, L. T., & Brotons, L. (2017). Using fire to promote biodiversity. *Science*,

    355(6331), 1264-1265. https://doi.org/10.1126/science.aam7672.

    https://doi.org/10.1111/cobi.12861

Kelly, L. T., Giljohann, K. M., Duane, A., Aquilué, N., Archibald, S., Batllori, E.,

    Bennett, A. F., Buckland, S. T., Canelles, Q., Clarke, M. F., Fortin, M. J., Hermoso,

    V., Herrando, S., Keane, R. E., Lake, F. K., McCarthy, M. A., Morán-Ordóñez, A.,

    Parr, C. L., Pausas, J. G., … Brotons, L. (2020). Fire and biodiversity in the

    Anthropocene. In *Science* (Vol. 370, Issue 6519). American Association for the

    Advancement of Science. https://doi.org/10.1126/science.abb0355

Kyereh, B., Ninnoni, R., & Agyeman, V. (2007). Degraded forests are more

    susceptible to forest fires: Some possible ecological explanations. *Journal of*

    Science and Technology (Ghana), *26*(2).

    https://doi.org/10.4314/just.v26i2.32987

Le Page, Y., Morton, D., Hartin, C., Bond-Lamberty, B., Pereira, J. M. C., Hurtt, G., & Asrar, G. (2017). Synergy between land use and climate change increases future fire risk in Amazon forests. *Earth System Dynamics*, *8*(4), 1237–1246. https://doi.org/10.5194/esd-8-1237-2017

Lebrija-Trejos, E., Pérez-García, E. A., Meave, J. A., Poorter, L., & Bongers, F. (2011). Environmental changes during secondary succession in a tropical dry forest in Mexico. *Journal of Tropical Ecology*, *27*(5), 477–489. https://doi.org/10.1017/S0266467411000253

Manly, B. F. J., McDonald, L. L., & Thomas, D. L. (1993). *Resource selection by animals: Statistical design and analysis for field studies*. Springer Dordrecht. https://doi.org/10.1007/978-94-011-1558-2

Marengo, J. A., Souza, C. M., Thonicke, K., Burton, C., Halladay, K., Betts, R. A., Alves, L. M., & Soares, W. R. (2018). Changes in Climate and Land Use Over the Amazon Region: Current and Future Variability and Trends. In *Frontiers in Earth Science* (Vol. 6). Frontiers Media S.A. https://doi.org/10.3389/feart.2018.00228

Mirra, I. M., Oliveira, T. M., Barros, A. M. G., & Fernandes, P. M. (2017). Fuel dynamics following fire hazard reduction treatments in blue gum (Eucalyptus globulus) plantations in Portugal. *Forest Ecology and Management*, *398*, 185–195. https://doi.org/10.1016/j.foreco.2017.05.016

Moreira de Araújo, F., Ferreira, L. G., & Arantes, A. E. (2012). Distribution Patterns of Burned Areas in the Brazilian Biomes: An Analysis Based on Satellite Data for the 2002–2010 Period. *Remote Sensing*, *4*(7), 1929–1946. https://doi.org/10.3390/rs4071929

Moreira, F., Rego, F. C., & Ferreira, P. G. (2001). Temporal (1958-1995) pattern of

change in a cultural landscape of northwestern Portugal: implications for fire

occurrence. In *Landscape Ecology* (Vol. 16).

https://doi.org/10.1023/A:1013130528470

Moreira, F., Vaz, P., Catry, F., & Silva, J. S. (2009). Regional variations in wildfire

susceptibility of land-cover types in Portugal: implications for landscape

management to minimize fire hazard. *International Journal of Wildland Fire*,

18(5), 563. https://doi.org/10.1071/WF07098.

Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E.,

Barbati, A., Corona, P., Vaz, P., Xanthopoulos, G., Mouillot, F. & Bilgili, E.

(2011) Landscape - wildfire interactions in southern Europe: Implications for

landscape management. *Journal of Environmental Management*, 92, 2389-

2402. DOI:10.1016/j.jenvman.2011.06.028.

Muylaert, R. L., Vancine, M. H., Bernardo, R., Oshima, J. E. F., Sobral-Souza, T.,

Tonetti, V. R., Niebuhr, B. B., & Ribeiro, M. C. (2018). Uma nota sobre os limites

territoriais da Mata Atlântica. *Oecologia Australis*, *22*(03), 302–311.

https://doi.org/10.4257/oeco.2018.2203.09

Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J.

(2000). Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772),

853–858. https://doi.org/10.1038/35002501

Nadporozhskaya, M. A., Chertov, O. G., Bykhovets, S. S., Shaw, C. H., Maksimova, E.

Y., & Abakumov, E. V. (2018). Recurring surface fires cause soil degradation of

forest land: A simulation experiment with the EFIMOD model. *Land Degradation

and Development*, *29*(7), 2222–2232. https://doi.org/10.1002/ldr.3021

Nepstad, D., McGrath, D., Stickler, C., Alencar, A., Azevedo, A., Swette, B., Bezerra, T.,

    DiGiano, M., Shimada, J., Seroa da Motta, R., Armijo, E., Castello, L., Brando, P.,

    Hansen, M. C., McGrath-Horn, M., Carvalho, O., & Hess, L. (2014). Slowing

    Amazon deforestation through public policy and interventions in beef and soy

    supply chains. *Science*, 344(6188), 1118–1123.

    https://doi.org/10.1126/science.1248525

Nunes, M. C. S., Vasconcelos, M. J., Pereira, J. M. C., Dasgupta, N., Alldredge, R.

    J., & Rego, F. C. (2005). Land cover type and fire in Portugal: do fires burn

    land cover selectively? *Landscape Ecology*, *20*(6), 661–673.

    https://doi.org/10.1007/s10980-005-0070-8

Oliveira, S., Moreira, F., Boca, R., San-Miguel-Ayanz, J., & Pereira, J. M. C.

    (2014). Assessment of fire selectivity in relation to land cover and topography:

    a comparison between Southern European countries. *International Journal of*

    Wildland Fire, *23*(5), 620. https://doi.org/10.1071/WF12053

Oliveira-Júnior, J. F. de, Teodoro, P. E., Silva Junior, C. A. da, Baio, F. H. R., Gava,

    R., Capristo-Silva, G. F., Gois, G. de, Correia Filho, W. L. F., Lima, M.,

    Santiago, D. de B., Freitas, W. K., Santos, P. J. dos, & Costa, M. da S. (2020).

    Fire foci related to rainfall and biomes of the state of Mato Grosso do Sul,

    Brazil. *Agricultural and Forest Meteorology*, *282–283*, 107861.

    https://doi.org/10.1016/j.agrformet.2019.107861

Piffer, P. R., Rosa, M. R., Tambosi, L. R., Metzger, J. P., & Uriarte, M. (2022a).

    Turnover rates of regenerated forests challenge restoration efforts in the

    Brazilian Atlantic forest. *Environmental Research Letters*, *17*(4), 045009.

    https://doi.org/10.1088/1748-9326/ac5ae1

Piffer, P. R., Rosa, M. R., Tambosi, L. R., Metzger, J. P., & Uriarte, M. (2022b).

Turnover rates of regenerated forests challenge restoration efforts in the

Brazilian Atlantic forest. *Environmental Research Letters*, *17*(4), 045009.

https://doi.org/10.1088/1748-9326/ac5ae1

Pivello, V. R. (2011). The use of fire in the cerrado and Amazonian rainforests of

Brazil: Past and present. *Fire Ecology*, *7*(1), 24–39.

https://doi.org/10.4996/fireecology.0701024

Pivello, V. R., Vieira, I., Christianini, A. V., Ribeiro, D. B., da Silva Menezes, L.,

Berlinck, C. N., Melo, F. P. L., Marengo, J. A., Tornquist, C. G., Tomas, W.

M., & Overbeck, G. E. (2021). Understanding Brazil's catastrophic fires:

causes, consequences and policy needed to prevent future tragedies.

Perspectives in Ecology and Conservation, *19*(3), 233–255.

https://doi.org/10.1016/j.pecon.2021.06.005

Ray, D., Nepstad, D., & Brando, P. (2010). Predicting moisture dynamics of fine

understory fuels in a moist tropical rainforest system: results of a pilot study

undertaken to identify proxy variables useful for rating fire danger. *New

Phytologist*, *187*(3), 720–732. https://doi.org/10.1111/j.1469-

8137.2010.03358.x

Rezende, C. L., Scarano, F. R., Assad, E. D., Joly, C. A., Metzger, J. P., Strassburg,

B. B. N., Tabarelli, M., Fonseca, G. A., & Mittermeier, R. A. (2018). From

hotspot to hopespot: An opportunity for the Brazilian Atlantic Forest.

Perspectives in Ecology and Conservation, *16*(4), 208–214.

https://doi.org/10.1016/j.pecon.2018.10.002

Ribeiro, M. C., Metzger, J. P., Martensen, A. C., Ponzoni, F. J., & Hirota, M. M.

(2009a). The Brazilian Atlantic Forest: How much is left, and how is the

remaining forest distributed? Implications for conservation. *Biological Conservation*, *142*(6), 1141–1153. https://doi.org/10.1016/j.biocon.2009.02.021

Ribeiro, M. C., Metzger, J. P., Martensen, A. C., Ponzoni, F. J., & Hirota, M. M. (2009b). The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation*, *142*(6), 1141–1153. https://doi.org/10.1016/j.biocon.2009.02.021

Sansevero, J. B. B., Garbin, M. L., Sánchez-Tapia, A., Valladares, F., & Scarano, F. R. (2020). Fire drives abandoned pastures to a savanna-like state in the Brazilian Atlantic Forest. *Perspectives in Ecology and Conservation*, *18*(1), 31–36. https://doi.org/10.1016/j.pecon.2019.12.004

Sansevero, J. B. B., Prieto, P. V., Sánchez-Tapia, A., Braga, J. M. A., & Rodrigues, P. J. F. P. (2017). Past land-use and ecological resilience in a lowland Brazilian Atlantic Forest: implications for passive restoration. *New Forests*, *48*(5), 573–586. https://doi.org/10.1007/s11056-017-9586-4

Schmerbeck, J., & Fiener, P. (2015). Wildfires, ecosystem services, and biodiversity in tropical dry Forest in India. *Environmental Management*, *56*(2), 355–372. https://doi.org/10.1007/s00267-015-0502-4

Silva, J. S., Moreira, F., Vaz, P., Catry, F., & Godinho-Ferreira, P. (2009). Assessing the relative fire proneness of different forest types in Portugal. *Plant Biosystems - An International Journal Dealing with All Aspects of Plant Biology*, *143*(3), 597–608. https://doi.org/10.1080/11263500903233250

Silva, J.S., Vaz, P.G., Moreira, F., Catry, F. & Rego, F.C. (2011) Wildfires as a

 major driver of landscape dynamics in three fire-prone areas of Portugal.

 Landscape and Urban Planning, 101, 349-358.

Silva Junior, C. H. L., Heinrich, V. H. A., Freire, A. T. G., Broggio, I. S., Rosan, T.

 M., Doblas, J., Anderson, L. O., Rousseau, G. X., Shimabukuro, Y. E., Silva,

 C. A., House, J. I., & Aragão, L. E. O. C. (2020). Benchmark maps of 33 years

 of secondary forest age for Brazil. *Scientific Data*, *7*(1), 269.

 https://doi.org/10.1038/s41597-020-00600-4

Singh, M., & Huang, Z. (2022). Analysis of forest fire dynamics, distribution and

 main drivers in the Atlantic Forest. *Sustainability*, *14*(2), 992.

 https://doi.org/10.3390/su14020992

Sorrensen, C. L. (2000). Linking smallholder land use and fire activity: examining

 biomass burning in the Brazilian Lower Amazon. *Forest Ecology and*

 Management, *128*(1–2), 11–25. https://doi.org/10.1016/S0378-1127(99)00283-

 2

Souza, C. M., Z. Shimbo, J., Rosa, M. R., Parente, L. L., A. Alencar, A., Rudorff, B.

 F. T., Hasenack, H., Matsumoto, M., G. Ferreira, L., Souza-Filho, P. W. M., de

 Oliveira, S. W., Rocha, W. F., Fonseca, A. V., Marques, C. B., Diniz, C. G.,

 Costa, D., Monteiro, D., Rosa, E. R., Vélez-Martin, E., … Azevedo, T. (2020).

 Reconstructing Three Decades of Land Use and Land Cover Changes in

 Brazilian Biomes with Landsat Archive and Earth Engine. *Remote Sensing*,

 12(17), 2735. https://doi.org/10.3390/rs12172735

Uhl, C., & Kauffman, J. B. (1990). Deforestation, Fire Susceptibility, and Potential

 Tree Responses to Fire in the Eastern Amazon. *Ecology*, *71*(2), 437–449.

 https://doi.org/10.2307/1940299

Uhl, C., Kauffman, J. B., & Cummings, D. L. (1988). Fire in the Venezuelan

Amazon 2: environmental conditions necessary for forest fires in the evergreen

rainforest of Venezuela. *Oikos*, *53*(2), 176. https://doi.org/10.2307/3566060

Uriarte, M., Schwartz, N., Powers, J. S., Marín-Spiotta, E., Liao, W., & Werden, L.

K. (2016). Impacts of climate variability on tree demography in second growth

tropical forests: the importance of regional context for predicting successional

trajectories. *Biotropica*, *48*(6), 780–797. https://doi.org/10.1111/btp.12380

Vancine, M. H., Muylaert, R. L., Niebuhr, B. B., Oshima, J. E. de F., Tonetti, V.,

Bernardo, R., De Angelo, C., Rosa, M. R., Grohmann, C. H., & Ribeiro, M. C.

(2024). The Atlantic Forest of South America: Spatiotemporal dynamics of the

vegetation and implications for conservation. *Biological Conservation*, *291*,

110499. https://doi.org/10.1016/j.biocon.2024.110499

Vaz, P. G., Dias, S., Pinto, P., Merten, E. C., Robinson, C. T., Warren, D. R., & Rego, F.

C. (2014). Effects of burn status and conditioning on colonization of wood by

stream macroinvertebrates. *Freshwater Science*, *33*(3), 832–846.

https://doi.org/10.1086/676657

Vaz, P. G., Merten, E. C., Robinson, C. T., & Pinto, P. (2021). Severely burned wood

from wildfires has low functional potential in streams. *Journal of Applied Ecology*,

58(6), 1346–1356. https://doi.org/10.1111/1365-2664.13872

FIGURES

Figure 1. Study area. Vegetational regions within the Brazilian Atlantic Forest.

Figure 2. Mean selection ratios (*w*) with 95% confidence intervals for LULC types across vegetational regions of the Brazilian Atlantic Forest. *w* = 1 indicates the LULC type burns according to availability; *w* < 1 means it burns significantly less, and *w* > 1 means it burns significantly more than expected by chance.

Figure 3. Mean selection ratios (*w*) with 95% confidence intervals for LULC types in each vegetational region of the Brazilian Atlantic Forest. Refer to Fig. 2 for more explanations.

Figure 4. Mean selection ratios (*w*) with 95% confidence intervals for LULC types across vegetational regions of the Brazilian Atlantic Forest from 1987 to 2022. Refer to Fig. 2 for more explanations.

Table 1. Characterization of land use and land cover types.

Land cover type	Definition	Land cover type in MapBiomas
Old-growth forests	Areas with native forest cover since the beginning of the historical MapBiomas series.	Forest Formation; Floodable Forest
Secondary forests	Areas that have transitioned from anthropogenic use to native forest vegetation. The transition period (vegetation age) can vary from 1 to 35 years.	The captions for secondary forest areas provide information about the age of the vegetation. In this context, we grouped all age ranges into a single category corresponding to secondary forests.
Openlands	Areas characterized by ground and shrub vegetation.	Savanna Formation; Herbaceous Sandbank Vegetation; Grassland; Pasture; Other Non-forest Formations
Temporary crops	Areas that are both sown and harvested within the same agricultural year, sometimes more than once.	Soybean; Sugar cane; Rice; Cotton; Other Temporary Crops
Perennial crops	Areas with plant species that are cultivated and live for more than two years without needing to be replanted annually	Coffee; Citrus; Palm Oil; Other Perennial Crops
Planted forests	Areas dedicated to silviculture, primarily consisting of *Eucalyptus* sp. and *Pinus* sp.	Forest Plantation
Mosaic of uses	Mix or combination of different land uses, creating a diverse and complex pattern across the landscape. It generally represents small areas that might include household agriculture and traditional cattle ranching (Almeida et al., 2016).	Mosaic of uses
Noncombustible areas	Areas with low vegetation density (fuel accumulation) or high humidity and water levels. This class was excluded from the analysis.	Mangrove; Hypersaline Tidal Flat; Wetland; Rocky Outcrop; Non vegetated area; Water

Table 2. Number of fires larger than 5 ha studied by vegetational region and the percentage of area occupied by each type of LULC.

Vegetational region	no. of fires	LULC types							
		Old-growth forests	Openlands	Temporary crops	Perennial crops	Plantation forests	Mosaic of uses	Secondary forests	Noncombustible areas
Semideciduous seasonal forest	17587	27.3	11.0	2.3	1.4	0.6	33.3	19.3	4.9
Decidual seasonal forest	2923	18.8	25.4	0.1	0.5	0.9	26.9	26.3	1.2
Dense ombrophilous forest	7777	22.8	12.0	2.4	0.6	1.5	33.4	21.2	6.2
Mixed ombrophilous forest	3472	22.8	10.0	5.7	0.6	3.4	29.0	24.4	4.1
Contact areas	8369	15.9	13.8	3.5	1.5	1.0	21.1	38.5	4.7