AQuaRef: Machine learning accelerated quantum refinement of protein structures

Roman Zubatyuk[1,#], Malgorzata Biczysko[2,#], Kavindri Ranasinghe[3,#], Nigel W. Moriarty[4]

Hatice Gokcan[1], Holger Kruse[6], Billy K. Poon[4], Paul D. Adams[4,5] , Mark P. Waller[6],

Adrian E. Roitberg[3], Olexandr Isayev[1,*], Pavel V. Afonine[4, *]

[1] *Department of Chemistry, Carnegie Mellon University, Pittsburgh, PA 15213, USA*

[2] *Faculty of Chemistry, University of Wrocław, F. Joliot-Curie 14, 50-383 Wrocław, Poland*

[3] *Department of Chemistry, University of Florida, Gainesville, FL 32611, USA*

[4] *Molecular Biophysics & Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720-8235, USA*

[5] *Department of Bioengineering, University of California Berkeley, Berkeley, CA 94720, USA*

[6] *Pending.AI, Eveleigh, NSW 2015, Australia*

\# These authors contributed equally

*Corresponding authors: P.V.A. - PAfonine@LBL.Gov,

O.I. - olexandr@olexandrisayev.com

ABSTRACT

Cryo-EM and X-ray crystallography provide crucial experimental data for obtaining atomic-detail models of biomacromolecules. Refining these models relies on library-based stereochemical restraints, which, in addition to being limited to known chemical entities, do not include meaningful noncovalent interactions relying solely on nonbonded repulsions. Quantum mechanical (QM) calculations could alleviate these issues but are too expensive for large molecules. We present a novel AI-enabled Quantum Refinement (AQuaRef) based on AIMNet2 neural network potential mimicking QM at substantially lower computational costs. By refining 41 cryo-EM and 30 X-ray structures, we show that this approach yields atomic models with superior geometric quality compared to standard techniques, while maintaining an equal or better fit to experimental data. Notably, AQuaRef aids in determining proton positions, as illustrated in the challenging case of short hydrogen bonds in the parkinsonism-associated human protein DJ-1 and its bacterial homolog YajL.

INTRODUCTION

While advances in predictive modeling, such as AlphaFold3[1] or RoseTTAFold[2,3], have provided powerful tools for structural biology, they remain limited while experimental methods, including protein crystallography and cryo-EM, are still cornerstones of structural biology and drug development[4]. Experimental data allow for the discovery of new structures emerging in life evolution, potentially exhibiting previously unseen features. These discoveries require unbiased information provided by experiments to explore the unknown[5]. Atomic model refinement is a crucial near-final stage in crystallographic or cryo-EM structure determination aimed at producing molecular models that meet standard validation criteria while optimally fitting the experimental data[6]. Refinement heavily relies on stereochemical restraints to maintain the correct geometry of the atomic model while fitting to the experimental data[7]. These restraints originate from standard libraries that tabulate the topology and parameters of known chemical entities[8,9],

which are universally employed across popular software packages, such as CCP4[10] and

Phenix[11].

The limitations of library-based restraints are manifold. Firstly, they only include terms for

maintaining covalent bond lengths, bond angles, torsion angles, planes and chirality while

preventing clashes through non-bonded repulsion[12]. However, it has been demonstrated

that at low resolution, these restraints are insufficient to maintain realistic, chemically

meaningful macromolecular geometries, making it essential to include additional

restraints on protein main chain φ/ψ angles, side chain torsion χ angles, as well as

hydrogen bond parameters and π-stacking interactions to stabilize protein or nucleic acid

secondary structure[12–18]. These additional restraints cannot be reliably inferred from the

atomic model alone and thus require manual error-prone annotation and curation using

additional sources of information, such as homologous high-resolution models. Secondly,

library-based restraints parametrize only known chemical entities, such as standard

amino and nucleic acids as well as previously defined ligands. Consequently, any

nonstandard entities or interactions, such as novel ligands or covalent cross-chain links,

require manual annotation and definition, without which refinement may fail to proceed

correctly or at all. Finally, deviations from standard covalent geometry due to local

chemical interactions are not uncommon[19–21]. While these deviations are valid, restraints

may interpret them as violations requiring 'correction'.

The advantage of using simple restraints[7] is the minimal computational cost they add to

the refinement workflow. A possible next step is to use a classical force field to account

for geometric elements[22]. However, these force fields have their own set of limitations:

they require parametrization for new chemical species and cannot distinguish between

chemically equivalent bonds in different chemical environments.

Quantum refinement is a fundamentally different approach, balancing the fitting to

experimental data with a term related to the quantum mechanical energy of the

system[23,24]. It has been demonstrated that the entire atomic model can benefit from a full

QM treatment[25–27]. Figure 1 presents a timeline showcasing the evolution of quantum

mechanics calculations for proteins, highlighting four key stages of progress and

advancements in technology and methodology. Traditionally, quantum refinements were

deemed impractical for macromolecules due to the computational requirements. Methods

often focused solely on the macromolecular region of interest, such as a ligand-binding

pocket or enzyme active site, while employing a classical approach for the rest of the

molecule[28,29]. Numerous approaches and implementations have been reported over

time[30], with GPU-accelerated codes enabling QM calculations for peptides and small

proteins of a few hundred atoms being one of the most prominent milestones[31].

Interaction-based model partitioning into chemically meaningful fragments[32] solved the

scalability issue in quantum calculations[26], which in turn enabled the refinement of larger

proteins. However, this approach remained computationally demanding.

Figure 1. Timeline illustrating the progression of quantum mechanics calculations applied

to proteins, emphasizing four critical stages marked by advancements in technology and

methodology.

Refinement of selected cryo-EM and X-ray atomic models across various resolutions

demonstrates the AQuaRef's ability to produce atomic models with superior geometric

quality compared to conventional techniques while maintaining or improving agreement

with experimental data. This work represents the first example where machine learning

(ML) potentials have been adopted to perform quantum refinement of the entire protein,

in contrast with a recent approach where ML potentials have been combined with the

ONIOM-like QM/MM partitioning[38].

RESULTS

Conceptually, quantum-based atomic model refinement is very similar to classic

refinement wherein atomic model parameters are iteratively adjusted in order to minimize

the residual, $T = T_{data} + w * T_{restraints}$. Here, T_{data} describes the fit of the model to the data

and $T_{restraints}$ incorporates chemical restraints with an *a-priori* unknown weight, w[39].

However, there are four fundamental differences. First, in QM refinement, restraints are

derived from quantum-mechanical calculations for the specific macromolecule in

consideration. Second, the requirements for the initial atomic model in QM refinement are

stricter compared to standard refinement: the atomic model must be correctly protonated,

atom-complete and free of severe geometric violations such as steric clashes or broken

covalent bonds[23,24]. Third, while crystallographic software packages inherently account

for crystal symmetry, QM codes generally do not. Fourth, crystallographic software is

capable of handling static disorder that is modeled with alternative conformations,

whereas QM codes typically lack this capability. All these nuances specific to quantum

refinement (except handling static disorder, which is a current limitation) are addressed

in the Quantum Refinement package (Q|R)[23,26,27,40], which is being developed as part of

this work and provides the necessary procedures to enable quantum refinement within

the Phenix software.

Figure 2. Computational scaling of the AIMNet2 neural network model in AQuaRef. Time
to compute energy and forces (left axis) and peak GPU memory usage (right axis) versus
the number of atoms in the system. Calculations are performed on a single Nvidia H100
PCIE 80GB GPU.

Conventional QM methods like density functional theory (DFT) for N-electron systems
require $O(N^2)$ storage and $O(N^3)$ arithmetic operations. This $O(N^3)$ complexity is a critical
bottleneck that limits the ability to study large realistic biological systems like proteins.
Figure 2 shows the computational scaling of the AIMNet2 model, where both energy and
force calculations, as well as peak GPU memory usage, scale linearly ($O(N)$) with system
size. For a large protein system of 100,000 atoms, single-point energy and forces can be
computed in 0.5 seconds. Overall, an atomic model consisting of approximately 180,000
atoms can fit into the 80GB memory of a single NVIDIA H100 GPU.

We tested the new quantum refinement procedure on 41 cryo-EM atomic models, 20 low-
resolution and 10 very high-resolution X-ray atomic models. Standard
stereochemistry[41,42] and model-to-data fit criteria[43–45], MolProbity validation tools[46] along
with newly developed metrics to evaluate hydrogen bond quality[18] were used to assess
the atomic models. Typically, the time needed for quantum refinement is about twice as
long as standard refinement, and often shorter than the standard refinement with
additional restraints such as the Ramachandran plot, secondary structure and side-chain
rotamer restraints[47–50]. Quantum refinement takes under 20 minutes in about 70% of
models considered in this work, with a maximum of about 1 hour (Supplementary Data:
Table 6). These computations can be performed on GPU-equipped laptops, with the only
limitation being available GPU memory.

154 Quantum refinement

The AQuaRef refinement procedure begins with a check for the completeness of the
atomic model, followed by the addition of any missing atoms. This may result in steric
clashes, particularly if the model was previously refined without hydrogen atoms. Models
with missing atoms that cannot be trivially added (e.g., missing main chain atoms) cannot
be used for quantum refinement. If clashes or other severe geometric violations are
detected, quick geometry regularization is performed using standard restraints, ensuring
that atoms move as little as necessary to resolve the clashes. For crystallographic
refinement, to account for interactions arising from crystallographic symmetry and
periodicity of unit cells, the model is expanded into a supercell by applying appropriate
space group symmetry operators[25]. Subsequently, it is truncated to retain only parts of
the symmetry copies within a prescribed distance from atoms of the main copy[40]. This
step is unnecessary for refinement against cryo-EM data. The atom-completed and
expanded model then undergoes the standard atomic model refinement protocol as
implemented in Q|R package[23].

Figure 3. a-d: Summary of refinements of 41 low-resolution cryo-EM models and 20 low-

resolution X-ray models using standard stereochemistry (blue) and AQuaRef (orange)

restraints (Supplementary Data: Table 1). **a:** MolProbity score, Ramachandran plot Z-

score, CaBLAM disfavored and r.m.s. deviation of refined model from initial model. **b:**

cross-correlation between experimental and model-generated maps (CC_{mask}), and

EMRinger score for cryo-EM models. **c:** R_{free} and R_{free}-R_{work} for X-ray models

(Supplementary Data: Table 3). Green band indicates favored range of corresponding
values. **d:** skew-kurtosis plots for hydrogen bond parameters (Hydrogen(H)...Acceptor(A)
distances and Donor-H…A angles) for refinements using (left-to-right): standard
restraints; standard restraints with addition of Ramachandran plot, secondary-structure
and side-chain rotamer restraints; and AQuaRef restraints. **e:** r.m.s. deviations between
refined and high-resolution homology models, refinements using standard versus
AQuaRef restraints, calculated using matching Cartesian coordinates (blue, lower-left)
and matching torsion angles (red, upper-right) (Supplementary Data: Tables 2,4). **f:**
summary of mean values, for all test refined models: MolProbity score, Ramachandran
Z-score, CaBLAM outliers, r.m.s. deviation of matching torsion angles between refined
and high-resolution homologous models, as well as R_{free}-R_{work} and R_{free} for X-ray models
and CC_{mask} and EMRinger score for cryo-EM models for refined models with standard
restraints (blue rhombi), standard restraints with addition of Ramachandran plot,
secondary-structure and side-chain rotamer restraints (blue circles); and AQuaRef
restraints (red stars). Red bars show standard deviations for starred values.

Application of the new refinement procedure to a set of deposited atomic models

To evaluate the performance of the new QM-based refinement, we refined 41 low-
resolution cryo-EM atomic models, 20 low-resolution and 10 ultra-high-resolution X-ray
atomic models, which contain only proteins. All selected 61 low-resolution atomic models
have high-resolution homologs, which were used as the ground truth for comparison
(Supplementary Data: Tables 2,4). Refinements were carried out using three sets of
restraints: QM restraints from AIMNet2 (AQuaRef refinement); standard restraints; and
standard restraints plus additional restraints on hydrogen bonds and angles involved in
maintaining secondary structure, main-chain φ/ψ angles (Ramachandran plot restraints)
and side-chain torsion χ angles (rotamer restraints).

Overall, low-resolution atomic models after quantum refinement exhibit systematically
superior geometry quality compared to those obtained using standard restraints, as
indicated by their MolProbity scores[51], Ramachandran Z-scores[52], CaBLAM disfavored[46]
(Fig. 3a), and skew-kurtosis plots for hydrogen bond parameters[18] (Fig. 3d). They also

systematically deviate more from the initial coordinates. These atomic models
demonstrate a very similar fit to the experimental data (Fig. 3b,c), with slightly less data
overfitting for X-ray atomic models, as evidenced by a smaller R_{work}-R_{free} gap and similar
R_{free}[53,54]. Since there is no equally efficient control over overfitting in cryo-EM as there is
with R_{free} in crystallography, the slightly lower cross-correlation between experimental and
model-calculated masked maps (CC_{mask})[43] and essentially the same EMRinger scores[55],
together with significantly improved atomic model geometry, likely indicate a reduction in
overfitting. Augmenting standard restraints with secondary structure, Ramachandran plot
and side-chain rotamer restraints expectedly improves the geometry (Fig. 3d,f), yet using
AQuaRef still produces superior atomic model geometries. With a few exceptions, atomic
models refined with quantum restraints are systematically closer to their higher-resolution
homologs compared to those using standard restraints alone or complemented with
additional restraints (Fig. 3e,f). In some of the most remarkable cases, the local structure
obtained with AQuaRef restraints closely matches the high-resolution homologs and
differs from those obtained using standard restraints by up to two Angstroms (Fig. 9a-c).
223

Comparison with alternative state-of-the-art approaches

To further evaluate the performance of AQuaRef refinement compared to other major
refinement methods and software, we refined selected low-resolution X-ray models using
the AMBER force field as a source of geometric restraints[56], the Rosetta all-atom force
field combined with its powerful sampling methods[57], and standard refinement as
implemented in REFMAC5[58]. For cryo-EM, there are fewer refinement alternatives, with
Servalcat[59] being the most popular, which we also used in this analysis.

For X-ray models, AQuaRef produced slightly better overall R_{free} values (Fig. 4a) and
substantially less data overfitting, as indicated by the R_{free}-R_{work} gap (Fig. 4b). For cryo-
EM models Servalcat lead to notably better CC_{mask} (Fig. 4c) and both scored the same
by EMRinger method (Fig. 4d). Models refined using AQuaRef and Rosetta performed
similarly well in terms of Rama-Z scores, achieving excellent results in most cases, while
REFMAC5 and Servalcat had the worst scores, and AMBER fell somewhere in between
(Fig. 4e). In terms of MolProbity scores and CaBLAM outliers (Fig. 4f,g), AQuaRef and

Rosetta also performed similarly well, significantly outperforming REFMAC5 and

Servalcat. Rosetta-refined models were closest to the high-resolution reference models,

followed by AQuaRef (Fig. 4h). This is likely due to Rosetta's use of non-gradient

optimization techniques, such as sampling and local model repacking, which have a

larger convergence radius compared to the gradient-driven minimization used in other

programs. Finally, AQuaRef and Rosetta both produced models that fit the expected

distribution of hydrogen bond parameters (Fig. 4i, Fig. 3d), followed by AMBER.

REFMAC5 and Servalcat largely failed to produce models fitting this distribution, with

Servalcat performing the worst.

Figure 4. Summary of refinements for 41 low-resolution cryo-EM models using AQuaRef
and Servalcat, and 20 low-resolution X-ray models using AQuaRef, REFMAC5, AMBER,
and Rosetta. **a-e:** Distributions of R_{free}, R_{free}-R_{work}, CC_{mask}, EMRinger score, and Rama-
Z, respectively. **f-h:** Mean values of MolProbity score, CaBLAM outliers, and r.m.s.
deviation from the reference model, calculated across all refined models; gray bands
represent the standard deviation. **i:** Skew-kurtosis plots for hydrogen bond parameters

(Hydrogen(H)...Acceptor(A) distances and Donor-H…A angles) for refinements
performed using REFMAC5, AMBER, Rosetta and Servalcat.

Case study: Short hydrogen bonds in human DJ-1 and its bacterial homologue
YajL

Short hydrogen bonds play a key functional role in proteins, and determining the
protonation states of involved residues is critical. However, accurate location of proton
positions experimentally remains challenging at resolutions near 1 Å. Lin *et al* [60] analyzed
high-resolution X-ray crystal structures of human DJ-1 and its bacterial homolog YajL to
determine the protonation states of carboxylic acids involved in dimer-spanning hydrogen
bonds. Their approach combined bond length analysis, leveraging the distinct lengths of
C=O and C–OH bonds, with qualitative interpretation of difference map peaks to identify
potential evidence of protons.

This method is complicated by stereochemical restraints applied during coordinate
refinement, which can bias bond lengths. For example, in E/D residues, bond length
restraints for COOH groups depend on whether a hydrogen atom is explicitly modeled
(Fig. 5c). To minimize this bias, Lin et al. performed final rounds of conjugate gradient
least-squares refinement in SHELXL[61] without applying restraints to the residues of
interest. In contrast, QM-based refinement avoids such biases entirely.

Figure 5. Wild-type DJ-1 (PDB code: 5SY6). Bond distances in the moiety of hydrogen bond between Oε2 (E15) and Oδ2 (D24), **a:** as measured in downloaded from PDB model, **b:** starting geometry for all refinements (H is present only in AQuaRef refinement), **c:** ideal library values in Phenix; geometry of –COOH or –COO groups is the same for Asp and Glu residues, **d:** unrestrained and **e:** restrained refinement with phenix.refine, **f:** refinement with AQuaRef. Distances in parentheses correspond to refinement using resolution-truncated data at 2 Å. H atom is shown only if it was explicitly modelled (present in the PDB model file).

AQuaRef refinement of DJ-1, starting with E15/D24 in an unprotonated state (Fig. 5b), produced proton positions and bond geometries (Fig 5f) consistent with Lin et al.'s findings (Fig. 5a) and unrestrained refinement using phenix.refine (Fig. 5d). However, restrained refinement with phenix.refine (Fig. 5e) yielded bond geometries that matched

library values assuming no proton on either COO group, highlighting the impact of restraint bias.

To test the robustness of AIMNet2 restraints in preserving accurate geometries, the same refinements were performed using experimental data truncated at 2 Å resolution. This truncation removed atomic-level details that could resolve bond lengths and hydrogen positions. AQuaRef produced results nearly identical to those obtained using the original 1.1 Å atomic resolution data, whereas restrained refinements with phenix.refine further biased oxygen-carbon distances toward idealized values for the unprotonated state (Fig. 5e-f, values in parentheses).

Figure 6. AIMNet2 energy values relative to their minimum as a function of hydrogen position between corresponding oxygen atoms, **a:** Oδ2 (D24) and Oε2 (E15) in DJ-1, **b:** Oδ2 (D23) and Oε2 (E14) in YajL, and **c:** Oδ2 (D23) and Oγ1 (T16) in YajL. Solid and dashed lines represent two instances of the bond in the YajL model.

Starting from an idealized symmetric arrangement (Fig. 5b), the refinement could, in principle, place the proton on either E15 or D24. To explain why the proton ultimately settled on Oδ2 of D24, two independent sources of evidence were considered. First, sampling the hydrogen position along the Oδ2–Oε2 bond vector and computing the AIMNet2 energy profile revealed a slight preference for D24 protonation (Fig. 6a).

Second, while the resolution and R-factors were insufficient for definitive proton
identification in the difference map, the difference map values along the Oε2-Oδ2 axis
showed elevated positive values near Oδ2, close to the prospective hydrogen position
(Fig. 7a). This, together with the energetic preference from AIMNet2, may have guided
the refinement to move the hydrogen toward D24.

Figure 7. Mean values of the difference density map, shown in absolute units (e/Å³) and
as standard deviation values along the O-H vector for the analyzed bonds for: **(a)** DJ-1
and **(b-c)** E. coli YajL models. All peak centers are aligned to the origin. Atoms belonging
to chains A and B are shown in blue and orange, correspondingly.

Figure 8. E. coli YajL (PDB code: 5SY4). Bond distances in the moiety of hydrogen bond between $O\varepsilon2$ (E14) and $O\delta2$ (D23) across chains A (blue) and B (orange), **a:** as measured in downloaded from PDB model, **b:** unrestrained and **c:** restrained refinement using phenix.refine, **d:** refinement with AQuaRef. H atom is shown only if it was explicitly modelled (present in the PDB model file).

Bacterial DJ-1 homologue, the YajL structure, contains two copies of the molecule in the asymmetric unit, resulting in two instances of the E14/D23 interaction. Similar to DJ-1, unrestrained refinement of YajL (Fig. 8b) yielded results consistent with Lin et al. (Fig. 8a). As with DJ-1, restrained refinement introduced significant bias in bond lengths (Fig. 8c) for both instances of the E14/D23 interaction.

Results from AQuaRef refinement aligned with Lin et al. and unrestrained phenix.refine refinement, suggesting that both D23 and E14 are protonated. However, in contrast to DJ-1, the proton in YajL does not appear to be fully associated with either of the two oxygen atoms. Instead, it seems to be shared between $O\varepsilon2$ and $O\delta2$, consistent with a Low Barrier Hydrogen Bond.

The AIMNet2 energy profile between Oε2 and Oδ2 supports this interpretation, showing

a relatively flat energy landscape (Fig 6b). This indicates that the hydrogen's position

could be entirely guided by the experimental data while staying within the flat region of

the AIMNet2 energy well. Indeed, there is a significant difference map peaks above 3 s.d.

and well above mean solvent density of 0.25 e/Å^3 very close to the position of hydrogens

in the refined model in both instances of the E14/D23 interaction (Fig. 7b).

Further evidence that C-Oδ2 elongation is due to Oε2$\cdots$H$\cdots$Oδ2 LBHB is provided by the

analysis of another hydrogen bond involving D23 and T16. All three, AQuaRef refinement

(Fig 8d), the AIMNet2 energy profile (Fig. 6c), and difference map density values along

the Oδ2 of D23 and Oγ1 of T16 (Fig 7c), confirm the protonation of T16 and rule out the

D23 protonation in the "anti" configuration.

DISCUSSION

Here, we present AQuaRef, a novel approach to the quantum refinement of entire protein

structures, made possible by using ML-accelerated quantum mechanical calculations

with AIMNet2. For the first time, this allows for the refinement of full atomic models of

realistic protein structures using stereochemical restraints derived from quantum

mechanical calculations.

Test refinements using 61 low-resolution X-ray and cryo-EM atomic models show

systematic improvements in geometric validation criteria by using QM restraints while

maintaining a similar fit to the experimental data and reducing overfitting. The presence

of high-resolution homologous atomic models, which are expected to better represent the

actual true structures than low-resolution atomic models, allowed us to assess whether

these improvements are associated with refined structures becoming closer to the true

ones. With a few exceptions, we find that atomic models refined with AQuaRef restraints

are systematically closer to their high-resolution matches. This indicates that QM-based

refined atomic models not only improve standard validation metrics but also provide more

realistic representations of the true structures compared to atomic models refined with

standard restraints. Expectedly, refining 10 very high-resolution atomic models did not

significantly alter the atomic coordinates but did lead to improved R-factors for all ten
models (Supplementary Data: Table 5). The most notable differences compared to
refinement with standard restraints were observed in the position of hydrogen atoms,
specifically those with rotational degrees of freedom (Fig. 9 d-g), where some of these
atoms reoriented during refinement to better fit the data and, at the same time, form
favorable hydrogen bonds. Another notable difference is the increased r.m.s. deviations
from ideal (library) bond and angle values in the case of AQuaRef refinement
(Supplementary Data: Table 5), which together with improved hydrogen positions is likely
to contribute to improved R-factors.

An extended comparison with popular state-of-the-art software packages and refinement
methods, including the use of AMBER and Rosetta force fields as refinement restraints,
as well as REFMAC5 and Servalcat from the CCP4 software suite, shows that for crystal
structure refinement, only Rosetta approaches AQuaRef in terms of the quality of refined
atomic model geometries. However, AQuaRef produces slightly improved R_{free} values
and significantly better R_{free}-R_{work} gaps, indicating reduced data overfitting. It is also worth
noting that Rosetta-based refinement is only available for crystal structures using X-ray
data, and refinement times with Rosetta are up to an order of magnitude slower. Although
Servalcat achieved superior CC_{mask} values compared to AQuaRef (Fig. 4d), this suggests
that Servalcat overfits the map, producing higher CC_{mask} values at the cost of significantly
poorer model geometry.

The case study of short hydrogen bonds in human DJ-1 and its bacterial homolog YajL,
as well as the protonation states of carboxylic acids involved in these hydrogen bonds,
highlights the feasibility of AQuaRef in determining proton positions consistent with
experimental evidence across diverse scenarios. This process is fully automated and
unbiased by the choice of restraints. Additionally, AIMNet2 energy profiles provide further
information about the characteristics of hydrogen bonds and protonation states, which
can be used to support specific hypotheses.

The method has been implemented in the quantum refinement software (Q|R), which is
built upon the CCTBX library[62] and optionally utilizes tools from Phenix. Q|R is accessible

within Phenix, thereby making these methods readily available to the broader community

of structural biologists.

Currently, AQuaRef is trained using commonly known amino acid residues, which means

the method can only be applied to protein-only structures. Another main limitation is that,

at present, static disorder (alternate conformations) is not handled in Q|R. Removing both

limitations is the subject of future work.

Figure 9. **a-c:** Close-up showing models refined with standard restraints (blue) and AQuaRef restraints (orange) superposed onto their higher-resolution homologous models (green) with their corresponding 2mFo-DFc Fourier maps contoured at 2σ; for PDB 5YI5, 8R1G, and 6XMX, respectively. **d-g:** Refinement with standard AQuaRef restraints

(orange) overlaid with their corresponding 2mFo-DFc and mFo-DFc Fourier maps,
contoured at 5σ (blue) and ±2σ (green, red), respectively (PDB 4O8H). The focus is on
hydrogen atoms with rotational degrees of freedom that re-orient during refinement with
AQuaRef restraints to satisfy the residual map and participate in hydrogen bonding.

METHODS

AIMNet2 training dataset and AQuaRef model

Since our goal was the parametrization of ML potential for polypeptides, our training
dataset needed to cover chemical (amino acid sequence and protonation states),
conformational, and intermolecular degrees of freedom. We began by creating a library
of small peptides as SMILES strings. We used all 20 standard amino acids, 11 alternate
protonation forms, three options for sequence start (ACE, NH3+, NH2), and four options
for the end (NME, NHE, CBX, CBA). We enumerated all possible mono- and di-peptides
and selected a random subset for tri- and tetra-peptides. Additionally, we generated
SMILES for peptides linked by the cysteine-cysteine disulfide bond and their selenium
counterparts. Molecular conformations were generated with OpenEye Omega[63] software
using dense torsion sampling. No restrictions were applied to the configurations of the
chiral centers, ensuring that the dataset and resulting model should work equally well for
D-, L-, and mixed stereochemistry peptides. Intermolecular interactions were modeled by
generating intermolecular complexes of 2 to 4 peptides with random orientations. No prior
knowledge of preferred types of secondary structure for polypeptides was used. To
manage the size of the dataset and the training process, we limited the size of peptides
and complexes to less than 120 atoms, including hydrogens.

Non-equilibrium conformations of peptides and complexes were sampled with molecular
dynamics simulations using the GFN-FF[64] force field. Cartesian restraints were added to
keep structures near the input structure, with random torsion and intermolecular degrees
of freedom. Molecular configurations for labeling (DFT calculations) and inclusion into the
training dataset were selected using Query-By-Committee active learning (AL)

approach[35]. We started with a random selection of 500k samples, used an ensemble of
4 models, and performed a total of 4 iterations of AL adding new samples with high
uncertainty of energy and atomic forces prediction. In the final iteration of AL, we
performed uncertainty-guided optimization of the structures, minimizing the weighted
difference of energy prediction and its uncertainty. This type of active sampling finds
structures that balance low predicted forces and high energy uncertainty. The entire
procedure resulted in a training dataset containing about one million samples, with a
median number of 42 atoms per sample.

DFT calculations were performed with the B97M-D4/def2-QZVP[65-68] method using ORCA
5.0.3 software[69]. Since the Q|R does not use periodic boundary conditions, and usually
not all ions and solvent molecules are resolved in the refinement, we used implicit
treatment of solvent effects with CPCM[70] method using parameters for water as solvent.

The core architecture of the AQuaRef model matches the base AIMNet2 model[33], with
few modifications. First, we did not use explicit long-range Coulomb and dispersion
interactions, we trained to total DFT-D4 energy instead. With CPCM treatment, the
Coulomb term could not be estimated using interactions between partial atomic charges,
and also long-range interactions are effectively screened with a polarizable continuum.
Long range dispersion interactions beyond the local cutoff of 5 Å have little effect on
atomic forces, which are important in Q|R refinement. We also added explicit short-range
exponential repulsion terms to make the potential more robust for the structures with
clashes. The model was trained to reproduce DFT-D4 energies, forces, and Hirshfeld
partial atomic charges.

Experimental data and atomic models for test cases

Protein-only, single-conformation high-to-low resolution X-ray crystallography and Cryo-
EM models, along with their corresponding experimental datasets, were selected from
RCSB and EMDB based on multiple criteria. These criteria include model size (between
1,000 and 10,000 non-hydrogen atoms), resolution (between 2.5 and 4 Å), geometric
model quality (MolProbity clashscore better than 50, with no covalent bonds deviating by

more than 4 r.m.s.d. from ideal library values), goodness of fit between the model and the

experimental data (Cryo-EM: CC_{mask} > 0.6, X-ray: R_{work} < 0.3), and the availability of a

higher-resolution (better than 2 Å) homologous model (main chain superposition r.m.s.d.

< 1 Å, sequence identity greater than 95%) for each considered model. Additionally, 11

ultra-high resolution single-conformation X-ray models were selected that contained only

protein and ordered water atoms.

Comparison of models

All atoms were used to calculate coordinate r.m.s. deviations between models before and

after refinement, as shown in Figure 3a. Coordinate r.m.s. deviations between models

used for test refinements and their high-resolution homologues were calculated using the

Phenix tool phenix.superpose_pdbs, which included all non-hydrogen backbone atoms

plus Cβ and Cγ atoms where present. R.m.s. deviations in torsion angle space were

calculated using CCTBX[62], with matching torsion angles selected as described by Headd

et al.[15].

Atomic model preparation for refinement

485   Model preparation for refinement (e.g., adding any missing atoms) was done using

486   qr.finalise program of Q|R, which uses the Reduce program[71] to add hydrogen atoms at

487   geometrically predicted positions. Model geometry regularization was done using the

488   Phenix tool phenix.geometry_minimization.

Model refinement

The exact same input models were used for all trial refinements. Real-space refinement

in Phenix was performed using the phenix.real_space_refine program[12]. Four refinement

runs were performed independently, starting with the same input maps (cryo-EM) or

reflection data (X-ray) and models. The runs included: 1) standard restraints consisting

of restraints on bond lengths, bond angles, torsion angles, planes, chirality, and non-

bonded repulsion; 2) standard restraints with the addition of secondary-structure

restraints; 3) standard restraints with the addition of Ramachandran plot restraints; and

4) standard restraints with the addition of secondary-structure and Ramachandran plot

restraints.

Quantum-based real- and reciprocal-space refinement was performed using the qr.refine

program of Q|R, using all default settings except for the source of QM restraints

(AQuaRef).

Reciprocal-space refinement in Phenix was performed using phenix.refine[72] with the

exact same four choices of restraints as in real-space refinement.

Software and availability

Phenix software is available at: phenix-online.org. Quantum refinement (Q|R) software is

available at qrefine.com. AQuaRef refinement is available in Phenix starting dev-5395

version. CCTBX-based Python scripts and the data (atomic models, cryo-EM maps, X-

ray diffraction data) used in this study are available at: https://phenix-

online.org/phenix_data/afonine/qr_aimnet2_2024/. Refinement parameters are

documented in README files, as well as in the Python scripts used to run the

refinements. Input data for deposited models were obtained from the Protein Data Bank[73]

and Electron Microscopy Data Bank[74], either by using the Phenix tool phenix.fetch_pdb

or from the CERES server[75].

Graphics software

518 Map and model images were prepared using PyMOL[76]. Routine inspection of maps and

519 models was performed using Coot[77]. Plots were generated using Matplotlib[78].

Acknowledgements

PVA, NWM, and PDA acknowledge funding from the National Institutes of Health (grants R01GM071939, P01GM063210, and R24GM141254), as well as support from the Phenix Industrial Consortium and the US Department of Energy under Contract No. DE-AC02-05CH11231. OI acknowledges support from the US National Science Foundation (NSF CHE-2154447). AER acknowledges support from the US National Science Foundation (NSF CHE-1802831 and OAC-2311632). MB acknowledges support from the COST Action CA21101 "Confined molecular systems: from a new generation of materials to the stars' (COSY)" supported by COST (European Cooperation in Science and Technology) and computer resources provided by Wroclaw Centre for Networking and Supercomputing (http://wcss.pl).

This work used Expanse at SDSC and Bridges-2 at PSC through allocation CHE200122 from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by NSF grants #2138259, #2138286, #2138307, #2137603, and #2138296. This research is part of the Frontera computing project at the Texas Advanced Computing Center. Frontera is made possible by the National Science Foundation award OAC-1818253. This research, in part, was done using resources provided by the Open Science Grid, which is supported by the award 1148698 and the U.S. DOE Office of Science.

Author contributions

Conceptualization: P.V.A., A.E.R., O.I., M.B.; Methodology: R.Z., H.G., K.R.; Software: P.V.A., N.W.M., R.Z., M.P.W, H.K.; Validation: M.B., K.R., H.G., P.D.A.; Formal analysis: M.B.; Data Curation: R.Z., H.G., K.R., P.V.A.; Writing (original draft): P.V.A., M.B., R.Z.; Writing (review & editing): all authors; Visualization: P.V.A., M.B., H.G., R.Z.; Supervision, P.V.A., A.E.R., O.I.; Project administration: P.V.A., A.E.R., O.I., M.B.; Funding Acquisition: P.D.A., P.V.A., A.E.R., O.I.

Competing Interests

The authors declare no competing interests.

References

1. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024).

2. Baek, M. *et al.* Accurate prediction of protein structures and interactions using a three-track neural network. *Science* **373**, 871–876 (2021).

3. Krishna, R. *et al.* Generalized biomolecular modeling and design with RoseTTAFold All-Atom. *Science* **384**, eadl2528 (2024).

4. Terwilliger, T. C. *et al.* AlphaFold predictions are valuable hypotheses and accelerate but do not replace experimental structure determination. *Nat. Methods* **21**, 110–116 (2024).

5. Edich, M., Briggs, D. C., Kippes, O., Gao, Y. & Thorn, A. The impact of AlphaFold2 on experimental structure solution. *Faraday Discuss.* **240**, 184–195 (2022).

6. Urzhumtsev, A. G. & Lunin, V. Y. Introduction to crystallographic refinement of macromolecular atomic models. *Crystallogr. Rev.* **25**, 164–262 (2019).

7. Evans, P. R. An introduction to stereochemical restraints. *Acta Crystallogr. D Biol. Crystallogr.* **63**, 58–61 (2007).

8. Vagin, A. A. *et al.* REFMAC 5 dictionary: organization of prior chemical knowledge and guidelines for its use. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2184–2195 (2004).

9. Engh, R. A. & Huber, R. Accurate bond and angle parameters for X-ray protein

structure refinement. *Acta Crystallogr. A* **47**, 392–400 (1991).

10. Agirre, J. *et al.* The *CCP* 4 suite: integrative software for macromolecular

crystallography. *Acta Crystallogr. Sect. Struct. Biol.* **79**, 449–461 (2023).

11. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons

and electrons: recent developments in *Phenix. Acta Crystallogr. Sect. Struct. Biol.*

75, 861–877 (2019).

12. Afonine, P. V. *et al.* Real-space refinement in *PHENIX* for cryo-EM and

crystallography. *Acta Crystallogr. Sect. Struct. Biol.* **74**, 531–544 (2018).

13. Kovalevskiy, O., Nicholls, R. A. & Murshudov, G. N. Automated refinement of

macromolecular structures at low resolution using prior information. *Acta Crystallogr.*

Sect. Struct. Biol. **72**, 1149–1161 (2016).

14. DeLaBarre, B. & Brunger, A. T. Considerations for the refinement of low-resolution

crystal structures. *Acta Crystallogr. D Biol. Crystallogr.* **62**, 923–932 (2006).

15. Headd, J. J. *et al.* Use of knowledge-based restraints in *phenix.refine* to improve

macromolecular refinement at low resolution. *Acta Crystallogr. D Biol. Crystallogr.*

68, 381–390 (2012).

16. Sobolev, O. V., Afonine, P. V., Adams, P. D. & Urzhumtsev, A. Programming new

geometry restraints: parallelity of atomic groups. *J. Appl. Crystallogr.* **48**, 1130–1141

(2015).

17. De Vries, I. *et al.* New restraints and validation approaches for nucleic acid

structures in *PDB-REDO. Acta Crystallogr. Sect. Struct. Biol.* **77**, 1127–1141 (2021).

18. Afonine, P. V., Sobolev, O. V., Moriarty, N. W., Terwilliger, T. C. & Adams, P. D.

Overall protein structure quality assessment using hydrogen-bonding parameters.

Acta Crystallogr. Sect. Struct. Biol. **79**, 684–693 (2023).

19. Moriarty, N. W., Liebschner, D., Tronrud, D. E. & Adams, P. D. Arginine off-kilter:

guanidinium is not as planar as restraints denote. *Acta Crystallogr. Sect. Struct. Biol.*

76, 1159–1166 (2020).

20. Richardson, J. S. *et al.* Model validation: local diagnosis, correction and when to

quit. *Acta Crystallogr. Sect. Struct. Biol.* **74**, 132–142 (2018).

21. Jiang, Z., Biczysko, M. & Moriarty, N. W. Accurate geometries for "Mountain pass"

regions of the Ramachandran plot using quantum chemical calculations. *Proteins*

Struct. Funct. Bioinforma. **86**, 273–278 (2018).

22. Moriarty, N. W. *et al.* Improved chemistry restraints for crystallographic refinement

by integrating the Amber force field into *Phenix*. *Acta Crystallogr. Sect. Struct. Biol.*

76, 51–62 (2020).

23. Zheng, M., Reimers, J. R., Waller, M. P. & Afonine, P. V. *Q | R*: quantum-based

refinement. *Acta Crystallogr. Sect. Struct. Biol.* **73**, 45–52 (2017).

24. Bergmann, J., Oksanen, E. & Ryde, U. Combining crystallography with quantum

mechanics. *Curr. Opin. Struct. Biol.* **72**, 18–26 (2022).

25. Zheng, M. *et al.* Including crystallographic symmetry in quantum-based refinement:

Q | R #2. *Acta Crystallogr. Sect. Struct. Biol.* **76**, 41–50 (2020).

26. Zheng, M. *et al.* Solving the scalability issue in quantum-based refinement: Q|R#1.

Acta Crystallogr. Sect. Struct. Biol. **73**, 1020–1028 (2017).

27. Wang, L. *et al.* Real-space quantum-based refinement for cryo-EM: *Q | R* #3. *Acta*

Crystallogr. Sect. Struct. Biol. **76**, 1184–1191 (2020).

28. Borbulevych, O. Y., Plumley, J. A., Martin, R. I., Merz, K. M. & Westerhoff, L. M.

Accurate macromolecular crystallographic refinement: incorporation of the linear

scaling, semiempirical quantum-mechanics program *DivCon* into the *PHENIX*

refinement package. *Acta Crystallogr. D Biol. Crystallogr.* **70**, 1233–1247 (2014).

29. Ryde, U. Combined quantum and molecular mechanics calculations on

metalloproteins. *Curr. Opin. Chem. Biol.* **7**, 136–142 (2003).

30. Canfield, P., Dahlbom, M. G., Hush, N. S. & Reimers, J. R. Density-functional

geometry optimization of the 150 000-atom photosystem-I trimer. *J. Chem. Phys.*

124, 024301 (2006).

31. Kulik, H. J., Luehr, N., Ufimtsev, I. S. & Martinez, T. J. Ab Initio Quantum Chemistry

for Protein Structures. *J. Phys. Chem. B* **116**, 12501–12509 (2012).

32. Zheng, M. & Waller, M. P. Adaptive quantum mechanics/molecular mechanics

methods. *WIREs Comput. Mol. Sci.* **6**, 369–385 (2016).

33. Anstine, D., Zubatyuk, R. & Isayev, O. AIMNet2: A Neural Network Potential to Meet

your Neutral, Charged, Organic, and Elemental-Organic Needs. Preprint at

https://doi.org/10.26434/chemrxiv-2023-296ch-v2 (2024).

34. Smith, J. S., Isayev, O. & Roitberg, A. E. ANI-1: an extensible neural network

potential with DFT accuracy at force field computational cost. *Chem. Sci.* **8**, 3192–

3203 (2017).

35. Smith, J. S., Nebgen, B., Lubbers, N., Isayev, O. & Roitberg, A. E. Less is more:

Sampling chemical space with active learning. *J. Chem. Phys.* **148**, 241733 (2018).

36. Devereux, C. *et al.* Extending the Applicability of the ANI Deep Learning Molecular

Potential to Sulfur and Halogens. *J. Chem. Theory Comput.* **16**, 4192–4202 (2020).

641 37. Zubatyuk, R., Smith, J. S., Nebgen, B. T., Tretiak, S. & Isayev, O. Teaching a neural

642   network to attach and detach electrons from molecules. *Nat. Commun.* **12**, 4870

643   (2021).

644 38. Yan, Z., Wei, D., Li, X. & Chung, L. W. Accelerating reliable multiscale quantum

645   refinement of protein–drug systems enabled by machine learning. *Nat. Commun.*

646   **15**, 4181 (2024).

647 39. Brünger, A. T., Karplus, M. & Petsko, G. A. Crystallographic refinement by simulated

648   annealing: application to crambin. *Acta Crystallogr. A* **45**, 50–61 (1989).

649 40. Wang, Y. *et al.* Optimal clustering for quantum refinement of biomolecular

650   structures: Q|R#4. *Theor. Chem. Acc.* **142**, 100 (2023).

651 41. Moriarty, N. W., Tronrud, D. E., Adams, P. D. & Karplus, P. A. Conformation-

652   dependent backbone geometry restraints set a new standard for protein

653   crystallographic refinement. *FEBS J.* **281**, 4061–4071 (2014).

654 42. Engh, R. A. & Huber, R. Structure quality and target parameters. in 474–484 (2012).

655   doi:10.1107/97809553602060000857.

656 43. Afonine, P. V. *et al.* New tools for the analysis and validation of cryo-EM maps and

657   atomic models. *Acta Crystallogr. Sect. Struct. Biol.* **74**, 814–840 (2018).

658 44. Booth, A. D. LXXIV. An expression for following the process of refinement in X-ray

659   structure analysis using fourier series. *Lond. Edinb. Dublin Philos. Mag. J. Sci.* **36**,

660   609–615 (1945).

661 45. Brünger, A. T. Free R value: a novel statistical quantity for assessing the accuracy

662   of crystal structures. *Nature* **355**, 472–475 (1992).

663 46. Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-

atom structure validation. *Protein Sci.* **27**, 293–315 (2018).

47. Lovell, S. C. *et al.* Structure validation by Cα geometry: ϕ,ψ and Cβ deviation.

Proteins Struct. Funct. Bioinforma. **50**, 437–450 (2003).

48. Hintze, B. J., Lewis, S. M., Richardson, J. S. & Richardson, D. C. Molprobity's

ultimate rotamer-library distributions for model validation: MolProbity's Ultimate

Rotamer-Library. *Proteins Struct. Funct. Bioinforma.* **84**, 1177–1189 (2016).

49. Lovell, S. C., Word, J. M., Richardson, J. S. & Richardson, D. C. The penultimate

rotamer library. *Proteins Struct. Funct. Genet.* **40**, 389–408 (2000).

50. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of

Coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).

51. Chen, V. B. *et al. MolProbity* : all-atom structure validation for macromolecular

crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).

52. Sobolev, O. V. *et al.* A Global Ramachandran Score Identifies Protein Structures

with Unlikely Stereochemistry. *Structure* **28**, 1249-1258.e2 (2020).

53. Tickle, I. J., Laskowski, R. A. & Moss, D. S. R_{free} and the R_{free} Ratio. I. Derivation of

Expected Values of Cross-Validation Residuals Used in Macromolecular Least-

Squares Refinement. *Acta Crystallogr. D Biol. Crystallogr.* **54**, 547–557 (1998).

54. Tickle, I. J., Laskowski, R. A. & Moss, D. S. R_{free} and the R_{free} ratio. II. Calculation

of the expected values and variances of cross-validation statistics in

macromolecular least-squares refinement. *Acta Crystallogr. D Biol. Crystallogr.* **56**,

442–450 (2000).

55. Barad, B. A. *et al.* EMRinger: side chain–directed model and map validation for 3D

cryo-electron microscopy. *Nat. Methods* **12**, 943–946 (2015).

56. Moriarty, N. W. *et al.* Improved chemistry restraints for crystallographic refinement by

integrating the Amber force field into Phenix. *Acta Cryst D* **76**, 51–62 (2020).

57. DiMaio, F. *et al.* Improved low-resolution crystallographic refinement with Phenix and

Rosetta. *Nat Methods* **10**, 1102–1104 (2013).

58. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal

structures. *Acta Cryst D* **67**, 355–367 (2011).

59. Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM single-particle

structure refinement and map calculation using Servalcat. *Acta Cryst D* **77**, 1282–1291

(2021).

60. Lin, J., Pozharski, E. & Wilson, M. A. Short Carboxylic Acid–Carboxylate Hydrogen

Bonds Can Have Fully Localized Protons. *Biochemistry* **56**, 391–402 (2017).

61. Sheldrick, G. M. Crystal structure refinement with SHELXL. *Acta Cryst C* **71**, 3–8

(2015).

62. Grosse-Kunstleve, R. W., Sauter, N. K., Moriarty, N. W. & Adams, P. D. The

Computational Crystallography Toolbox : crystallographic algorithms in a reusable

software framework. *J. Appl. Crystallogr.* **35**, 126–136 (2002).

63. Hawkins, P. C. D., Skillman, A. G., Warren, G. L., Ellingson, B. A. & Stahl, M. T.

Conformer Generation with OMEGA: Algorithm and Validation Using High Quality

Structures from the Protein Databank and Cambridge Structural Database. *J. Chem.*

Inf. Model. **50**, 572–584 (2010).

64. Spicher, S. & Grimme, S. Robust Atomistic Modeling of Materials, Organometallic,

and Biochemical Systems. *Angew. Chem. Int. Ed.* **59**, 15665–15673 (2020).

65. Mardirossian, N. & Head-Gordon, M. Mapping the genome of meta-generalized

gradient approximation density functionals: The search for B97M-V. *J. Chem. Phys.*

142, 074111 (2015).

66. Caldeweyher, E. *et al.* A generally applicable atomic-charge dependent London

dispersion correction. *J. Chem. Phys.* **150**, 154122 (2019).

67. Weigend, F. & Ahlrichs, R. Balanced basis sets of split valence, triple zeta valence

and quadruple zeta valence quality for H to Rn: Design and assessment of

accuracy. *Phys. Chem. Chem. Phys.* **7**, 3297 (2005).

68. Weigend, F. Accurate Coulomb-fitting basis sets for H to Rn. *Phys. Chem. Chem.*

Phys. **8**, 1057 (2006).

69. Neese, F., Wennmohs, F., Becker, U. & Riplinger, C. The ORCA quantum chemistry

program package. *J. Chem. Phys.* **152**, 224108 (2020).

70. Barone, V. & Cossi, M. Quantum Calculation of Molecular Energies and Energy

Gradients in Solution by a Conductor Solvent Model. *J. Phys. Chem. A* **102**, 1995–

2001 (1998).

71. Word, J. M., Lovell, S. C., Richardson, J. S. & Richardson, D. C. Asparagine and

glutamine: using hydrogen atom contacts in the choice of side-chain amide

orientation 1 1Edited by J. Thornton. *J. Mol. Biol.* **285**, 1735–1747 (1999).

72. Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with

phenix.refine. *Acta Crystallogr. D Biol. Crystallogr.* **68**, 352–367 (2012).

73. Berman, H. M. The Protein Data Bank. *Nucleic Acids Res.* **28**, 235–242 (2000).

74. The wwPDB Consortium *et al.* EMDB—the Electron Microscopy Data Bank. *Nucleic*

Acids Res. **52**, D456–D465 (2024).

75. Liebschner, D. *et al. CERES* : a cryo-EM re-refinement system for continuous

improvement of deposited models. *Acta Crystallogr. Sect. Struct. Biol.* **77**, 48–61

(2021).

76. Schrödinger, LLC. The PyMOL Molecular Graphics System, Version 1.8. (2015).

77. Emsley, P. & Cowtan, K. *Coot* : model-building tools for molecular graphics. *Acta*

Crystallogr. D Biol. Crystallogr. **60**, 2126–2132 (2004).

78. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95

(2007).

SUPPLEMENTAL METHODS

Case study: Short hydrogen bonds in human DJ-1 and its bacterial homologue
YajL

Atomic models and experimental data for human DJ-1 and the bacterial homologue YajL
were downloaded from the Protein Data Bank (PDB) using accession codes 5SY6 and
5SY4, respectively. Reflection intensities were used in all calculations. Both atomic
models were protonated using the Reduce program from Phenix suite, as the deposited
models lacked hydrogen atoms.

To address uncertainty in refined atomic model parameters due to intrinsic refinement
properties, particularly in the coordinates, interatomic distances are reported as averages
with standard deviations, based on 100 refinement runs. Each refinement run (restrained
and unrestrained with phenix.refine as well as AQuaRef) used identical settings but began
with models where coordinates were randomized with an r.m.s.d. of 0.1 Å.

To avoid biases in assumptions about the protonation states of E/D and D/T residues,
these residues were modeled as unprotonated with idealized geometries in the input
refinement models. For AQuaRef refinements, the proton was positioned exactly between
the Oε2 and Oδ2 atoms for DJ-1 (Fig. 5b) and YajL, as well as between the Oδ2 and Oγ1

atoms for YajL. In phenix.refine, this proton was not explicitly modeled to avoid assuming

its association with a particular residue.

Restrained phenix.refine refinements were performed with 10 macro-cycles to ensure

convergence. Unrestrained refinements used the same settings, except geometric

restraints affecting hydrogen bond interaction moieties in E/D and D/T residues were

eliminated. Excluded restraints were covalent bond and angle restraints for COO moieties

and repulsion restraints between $O\varepsilon 2$, $O\delta 2$, and $O\gamma 1$ atoms. AQuaRef refinements were

performed with all default settings.

The AIMNet2 energy profiles between corresponding oxygen atoms were calculated by

sampling the proton positions along the line connecting the oxygens at 100 equally

spaced points.

The sigma-A weighted Fo-Fc difference map value profiles were interpolated along the

$O\delta 2$-H, $O\varepsilon 2$-H, and $O\gamma 1$-H lines using tri-cubic interpolation and averaged across 100

difference maps from corresponding refinements. The mean bulk-solvent density value

was estimated using the flat bulk-solvent model, as implemented in Phenix.

All Python scripts using Q|R and CCTBX libraries to perform all the above calculations

are available in Supplemental Data.

776